

# More than 18,000 effectors in the *Legionella* genus genome provide multiple, independent combinations for replication in human cells

Laura Gomez-Valero<sup>1</sup>, Christophe Rusniok<sup>2</sup>, Danielle Carson<sup>3</sup>, Sonia Mondino<sup>2</sup>, Ana-Elena Perez-Cobas<sup>2</sup>, Monica Rolando<sup>2</sup>, Shivani Pasricha<sup>4</sup>, Sandra Reuter<sup>5</sup>, Jasmin Demirtas<sup>2</sup>, Johannes Crumbach<sup>2</sup>, Stéphane Descorps-Declere<sup>2</sup>, Elizabeth Hartland<sup>6</sup>, Sophie Jarraud<sup>7</sup>, Gordon Dougan<sup>8</sup>, Gunnar Schroeder<sup>3</sup>, Gad Frankel<sup>3</sup>, Carmen Buchrieser<sup>2</sup>

<sup>1</sup>Institute Pasteur, <sup>2</sup>Institut Pasteur, <sup>3</sup>Imperial College London, <sup>4</sup>CESAR-University of Melbourne, <sup>5</sup>University of Freiburg, <sup>6</sup>Hudson Institute of Medical Research, <sup>7</sup>Centre National de Référence des Legionella, <sup>8</sup>Wellcome Trust Sanger Institute

Submitted to Proceedings of the National Academy of Sciences of the United States of America

The genus *Legionella* comprises 65 species among, which *Legionella pneumophila* is a human pathogen. To understand the evolution of an environmental to an accidental human pathogen, we have functionally analyzed 80 *Legionella* genomes spanning 58 species. Uniquely, an immense repository of 18,000 secreted proteins encoding 137 different eukaryotic-like domains and over 200 eukaryotic-like proteins is paired with a highly conserved T4SS. Specifically, we show that eukaryotic Rho and Rab-GTPase domains are found nearly exclusively in eukaryotes and *Legionella*. Translocation assays for selected Rab-GTPase proteins revealed that they are indeed T4SS secreted substrates. Furthermore, F/U-box and SET domains were present in >70% of all species suggesting that manipulation of host signal transduction, protein turnover and chromatin modification pathways are fundamental intracellular replication strategies for Legionellae. In contrast, the Sec-7 domain was restricted to *L. pneumophila* and seven other species, indicating effector repertoire tailoring within different amoebae. Functional screening of 47 species revealed 60% were competent for intracellular replication in THP-1 cells, but interestingly this phenotype was associated with diverse effector assemblages. These data, combined with evolutionary analysis indicate that the capacity to infect eukaryotic cells has been acquired independently many times within the genus and that a highly conserved T4SS secretes an exceptional number of different proteins shaped by inter-domain gene transfer. Furthermore we revealed the surprising extent to which Legionellae have co-opted genes and thus cellular functions from their eukaryotic hosts and provide a new understanding of how dynamic reshuffling and gene-acquisition has led to the emergence of human pathogens.

Legionella | co-evolution | horizontal gene transfer | protozoa

## Introduction

Legionnaires' disease or legionellosis is an atypical pneumonia caused by bacteria of the genus *Legionella*. Shortly after the discovery of *L. pneumophila* (1) it was reported that this bacterium is pathogenic for freshwater and soil amoebae of the genera *Acanthamoeba* and *Naegleria* (2). This finding led to a new perception in microbiology, whereby bacteria that parasitize protozoa can utilize similar processes to infect human cells. Sequencing and analyses of the *L. pneumophila* genome substantiated this idea, when it revealed the presence of a large number and variety of eukaryotic-like domains within the predicted proteome (3). Many of these proteins, termed effector proteins, were shown to be secreted into the host cell where they facilitate *Legionella* intracellular replication within a specialized compartment termed the *Legionella* containing vacuole (LCV) (3, 4). Overall, the type IV secretion system (T4SS), Dot/Icm, secretes more than 300 different effector proteins into the host cell and is indispensable for virulence of *L. pneumophila* (5-8). The presence of the Dot/Icm T4SS in other *L. pneumophila* strains and in selected *Legionella*

species had also been reported (9-12) but recent genome scale studies of *Legionella* (13-15) indicated that the T4SS system is present in every *Legionella* strain analyzed.

Despite high conservation of the Dot/Icm system among different *Legionella* species, effector repertoires appear to vary greatly. An analysis of putative T4SS effectors of *L. longbeachae*, the second most frequent cause of Legionnaires' disease, revealed that only about 50% of the virulence factors described in *L. pneumophila* were also present in the genome of *L. longbeachae* (16). Recently, Burstein *et al.* (14) analyzed 38 *Legionella* species using a machine learning approach to predict T4SS effectors and Joseph *et al.* (15) examined *Legionella* genome dynamics, both concluding that DNA interchange between different species is rare. However, still little is known about the potential of the different species to cause human disease and about the impact and the specific characteristics of the T4SS effectors on the evolution of new human pathogens within this environmental bacterial genus.


Here we present a comprehensive analysis of the *Legionella* genus genome, covering 80 *Legionella* strains belonging to 58 *Legionella* species and subspecies. We establish a pan-genus pool of putative T4SS effectors and show that this comprises over 18,000 proteins and identify more than 200 new eukaryotic-like proteins and 137 eukaryotic domains, including a unique class

## Significance

*Legionella* comprises 65 species for which aquatic amoebae are the natural reservoirs. Using functional and comparative genomics to deconstruct the entire bacterial genus we reveal the surprising parallel evolutionary trajectories that have led to the emergence of human pathogenic *Legionella*. An unexpectedly large and unique repository of secreted proteins (>18,000) containing eukaryotic-like proteins acquired from all domains of life (plant, animal, fungal, archaea) is contrasting with a highly conserved type 4 secretion system. This study reveals an unprecedented environmental reservoir of bacterial virulence factors, and provides a new understanding of how reshuffling and gene-acquisition from environmental eukaryotic hosts, may allow for the emergence of human pathogens.

## Reserved for Publication Footnotes

137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204



**Fig. 1.** The *Legionella* genomes are diverse in size and gene content. A) Phylogeny of the genomes based on the core genome, genome size, GC content and number of singletons of each species are depicted. Numbers represent bootstrap values. Branches are coloured according to the clade they belong to. Genome size and GC content include plasmids if present in the corresponding species. The number of singletons is based on the results of OrthoMCL (takes into account orthologs and paralogs). Each species has been compared to the others without taking into account strains from the same species to avoid bias due to the number of strains sequenced within a species. B) Occurrence of genes within the 80 analysed *Legionella* genomes. Left end of the x-axis, genes present in a single genome (strain specific genes; 5832 ≈32% of the pangenome); right end of the x-axis, genes present in all 80 genomes (core-genome; 1008 genes ≈6% of the pan-genome) C) Gene accumulation curve for the total number of proteins of the 80 genomes. D) Negative correlation between genome size and GC content indicating high acquisition of foreign genes (Pearson's correlation coefficient equal to -0.46 with a p-value<0.0001)

of putative bacterial Rab GTPases. We confirmed experimentally that a subset of these proteins translocate into the host cell upon infection. We conclude that the T4SS is highly conserved at the sequence level, but the effector proteins secreted are highly diverse.

**Results and discussion**

**The *Legionella* genus genome is dynamic and characterized by frequent genetic exchange.** We sequenced 58 *Legionella* species of which 16 were newly sequenced, and analyzed them in combination with all publicly available genomes (80 genomes in total) (SI Appendix, Table S1). The *Legionella* genomes were extremely diverse, as the genome size varied from 2.37Mb (*L. adalaidensis*) to 4.88Mb (*L. santicrucis*), the GC content from 34.82% (*L. busanensis*) to 50.93% (*L. geestiana*) and the number of clusters of orthologous genes as defined with OrthoMCL was 17,992 of which 5,832 (32%) were strain specific (singletons) (Fig. 1A). Only 1,008 genes (6%) constituted the core genome (Fig. 1B), compared to an earlier analysis of 38 *Legionella* species, which found 16,416


clusters of orthologous and 1,054 core genes (14). The addition of 40 new genomes comprising 16 newly sequenced *Legionella* species in our study increased the number of orthologous gene clusters by over 1,576 and decreased the core genome by 46 genes, underlining the high diversity of the *Legionella* genus. This difference suggested that the *Legionella* genus pan-genome is far from fully described and that sequencing of additional *Legionella* species will increase the genus gene repertoire significantly. This was supported by the rarefaction curve that does not reach a plateau (Fig. 1C).

The highly dynamic nature of these genomes is also seen in the analysis of the strain specific genes and the accessory genome as it highlights the presence of several mobile genetic elements; often associated with genes encoding for transfer regions/conjugative elements such as the type IVA secretion systems (T4ASS). These T4ASSs (classified as T4SSE, G, I and T (17) are present in each strain to varying degrees indicating that they circulate among the different *Legionella* strains (SI Appendix, Table S2) and therefore drive genome dynamics and

205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272

273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340

341  
342  
343  
344  
345  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408



**Fig. 2.** Eukaryotic domains have a diverse distribution within the genus *Legionella* suggesting multiple acquisition events. The number and distribution of the 41 most frequently identified eukaryotic motifs within the genus *Legionella* are shown. Numbers represent the number of proteins containing this eukaryotic motif. Abbreviations used: ANK (ankyrin), F-box, U-box, SET domain, Pkinases (protein kinases), Sec-7 domain, LRR (leucine rich repeats), Miro (Mitochondrial Rho domain), TTL (tubulin-tyrosine ligase), SH2 (The Src homology 2), PAM2 (ataxin-2, C-terminal), PPR (pentatricopeptide repeat), I-set (immunoglobulin I-set), NP (nucleoside phosphatase gda1/cd39), HAD (HAD-superfamily hydrolase), DH (Dbl homology domain), Mit. Substrate (mitochondrial substrate/solute carrier), Rho GTPases-activating protein domain, T-complex (T-complex10/11), PC65 (Peptidase C65 otubain), Ergosterol (Ergosterol biosynthesis), Flavin (flavin monooxygenase-like), Astacin (Peptidase M12A, astacin), Cyt:P450 (Cytochrome P450), Cytokine FAD (Cytokinin dehydrogenase 1, FAD/cytokinin binding domain), PQ loop repeat, Peptidase C2 (calpain, catalytic domain), LR glioma (Leucine-rich glioma-inactivated, ETP repeat), Ovarian (Ovarian tumour, otubain), Papain (Peptidase C1A, papain C-terminal), DOT1 (Histone methylation DOT1), Rab small GTPases, DUF155, C/C (Clathrin/coatomer adaptor, adaptinlike), RCC1 (Regulator of chromosome condensation).

diversification. It has been suggested that the incorporation of foreign DNA via horizontal gene transfer (HGT) is responsible for an increase in the AT content and the increase in genome size (18). Indeed, we found a negative correlation between the genome size and the GC content for the *Legionella* genomes, which also suggests frequent HGT (Fig. 1D) (19). Despite the importance of flagella for transmission to new hosts as shown for *L. pneumophila*, flagella encoding genes were not conserved in all species, but showed a patchy distribution, as 23 of the 80 strains analyzed lacked flagella genes (SI Appendix, Fig. S1). The analyses showed that the *Legionella* genus genome is highly diverse, dynamic and shaped by HGT.

**The genus *Legionella* encodes proteins with 137 different eukaryotic domains.** Interpro scan analysis of all 58 *Legionella* species revealed the presence of 137 different eukaryotic motifs/domains in the genus *Legionella* (SI Appendix, Table S3) according to the definition that an eukaryotic domain is one that is found in >75% of eukaryotic genomes and <25% in prokaryotic

genomes. The most abundant eukaryotic domains identified were ankyrin repeats. Interestingly, *L. santicrucis* and *L. massiliensis* encoded 41 and 39 ankyrin domains, respectively (Fig. 2). Ankyrin motifs were found frequently associated with other eukaryotic motifs and thus constituted modular proteins associated with eukaryotic F-box, U-box, Rab or SET domains. Notably, F-box and U-box domains were present in more than two thirds of the species analyzed (Fig. 2) suggesting manipulation of the host ubiquitin-system is a fundamental virulence strategy of *Legionella* species. Generally, the genomes contained one to three F-box containing proteins with the exception of *L. nautarum* and *L. dronzanskii*, which contained 18 and 10, respectively. The SET domain containing protein RomA of *L. pneumophila* that induces a unique host chromatin modification (20) is present in 46 of the 58 *Legionella* species suggesting the ability of many *Legionella* species to manipulate host chromatin (Fig. 2). Interestingly, the Sec-7 domain present in the effector RalF, a bacterial ARF guanine exchange factor and the first described Dot/Icm effector

**Table 1. Homology of *Legionella* Rab domains-containing proteins against protozoan Rab proteins**

Domain	Protein	First blast hit	Identity	Coverage	e-value
Rab	Lade0491	<i>Entamoeba histolytica</i>	35%	52%	4.E-17
Rab	LgoA0634	<i>Paramecium tetraurelia</i>	33%	51%	2.E-19
Rab	Llo3288	<i>Ichthyophthirius multifiliis</i>	42%	53%	4.E-31
Rab	Lstei0814	<i>Tetrahymena thermophila</i>	34%	86%	3.E-26
Rab	Lstei2185	<i>Stentor coeruleus</i>	38%	55%	6.E-29
Rab	Lbir2252	<i>Entamoeba invadens</i>	32%	55%	5.E-15
Rab	Lges1860	<i>Entamoeba histolytica</i>	34%	55%	7.E-25
Rab+ Fbox	Lwad3214	<i>Paramecium tetraurelia</i>	34%	35%	7.E-14
Rab	Lgra2891	<i>Guillardia theta</i>	36%	56%	2.E-19
Rab	Lgra3435	<i>Entamoeba histolytica</i>	35%	59%	2.E-27
Rab	Lma1540	<i>Paramecium tetraurelia</i>	34%	55%	1.E-17
Rab + ank	LmasA3690	<i>Oxytricha trifallax</i>	34%	19%	2.E-19
Rab	Lqua0234	<i>Dictyostelium fasciculatum</i>	38%	34%	1.E-25
Rab	Lquin3026	<i>Tetrahymena thermophila</i>	34%	57%	1.E-19
Rab	Lspi0161	<i>Naegleria gruberi</i>	34%	24%	7.E-24
Rab	Lwal3261	<i>Paramecium tetraurelia</i>	33%	85%	7.E-18

Each Rab protein listed in the table represents a different orthologous group. Results are based on blastp searches using the non-redundant NCBI database.

of *L. pneumophila* (21) was present in only eight (*L. pneumophila*, *L. longbeachae*, *L. feelei*, *L. sainthelensis*, *L. santacruzis*, *L. shake-speari*, *L. quateirensis* *L. moravica*) of the 58 *Legionella* species analyzed, suggesting that, different effectors may compensate for RalF activity or that LCV biogenesis varies among different species (Fig. 2).

One newly identified motif in *Legionella* was the ergosterol reductase ERG4/ERG24 (IPR001171) domain. Ergosterol is the primary sterol in the cell membranes of filamentous fungi, present in membranes of yeast and mitochondria (22). Importantly, it is also the major sterol of amoebae such as *A. castellanii* and *A. polyphaga*, the natural hosts of *Legionella* (23, 24). We found that 31 *Legionella* species encoded one or two proteins with the ERG4/ERG24 domain (Fig. 2). The *L. longbeachae* protein (L1o1320) containing this domain showed 56% aa identity to that encoded by the amoeba *Naegleria gruberi* and 30% aa identity to that encoded by *A. castellanii* strain Neff. This domain was also present in other amoebae related bacteria such as *Parachlamydia acanthamoebae* and *Protochlamydia naegleriophila*, as well as *Coxiella burnetii*. Phylogenetic analyses suggested that *L. longbeachae* acquired this domain from amoeba (SI Appendix, Fig. S2A).

Phylogenetic analyses of the here identified C-terminal alIinase and Caleosin domains present in *L. beliardensis* and *L. anisa* or the *L. longbeachae* clade (Fig. 2), respectively further supported acquisition of these domains from plants, amoeba or fungi (SI Appendix, Fig. S2B-C). They probably help *Legionella* to fight competitor bacteria or fungi in amoebae or in the environment. Taken together, our analyses highlight key domains preferentially present in protozoa, fungi, plants or animals that have been acquired by different *Legionella* species.

**A unique case in the prokaryotic world: *Legionella* encode small GTPase-like domains** The Ras-related small GTPase superfamily comprises more than 150 members in humans, which function as key regulators of signal transduction in almost all cellular processes(25). These enzymes bind and hydrolyse GTP to GDP and activate downstream effectors when bound to GTP. The first identified member was the p21-Ras protein, an evolutionary conserved small GTPase that controls cell proliferation, survival and migration through its effector binding at RAF/MAPK and PI3K (26). The Ras protein superfamily is subdivided into at least five distinct branches: Ras, Rho, Rab, Arf and Ran (27).


Evolutionarily conserved orthologs are found in *Drosophila*, *C. elegans*, *S. cerevisiae*, *S. pombe*, *Dictyostelium* and plants (28).

The only Rab-like protein in a prokaryotic genome was reported in the *L. longbeachae* genome sequence (16). However, upon analysis of our 80 *Legionella* strains, we identified 184 small GTPases of which 104 could be classified with a very high confidence as Rab, Ras or Rho like proteins (34 Ras, 71 Rab and one Rho domain) (SI Appendix, Table S4 and Fig. S3). Blastp analysis of these proteins in the NCBI database revealed that 149 of the 184 small GTPases of *Legionella* were exclusively present in *Legionella* and eukaryotic organisms (Table 1). The Rab domain was localized to different parts of the effector proteins, and a subset of Rab proteins carried additional domains such as U-box domains, ankyrin motifs or F-box domains (Fig. 3A). Alignment of the different Rab domains identified in the *Legionella* genomes revealed that the structural features of eukaryotic Rab domains were conserved among the *Legionella* proteins (SI Appendix, Fig. S4).

To analyze further the evolutionary history of the Ras-related domains in *Legionella* we undertook phylogenetic analyses of these proteins. For example, the two *L. longbeachae* Rab proteins, Llo1716 and Llo3288, were present in all strains closely related to *L. longbeachae*, suggesting that they and their orthologous share a common origin and evolved from a gene acquired by the ancestor of all these species (SI Appendix, Fig. S5). Further phylogenetic analysis of 16 Rab proteins present in eight different *Legionella* species showed that these Rab domains were acquired by HGT, mainly from protozoa (Fig. 3B and SI Appendix, Fig. S6A-P). Recently a novel isoform of Rab5D was identified in the *Acanthamoeba polyphaga mimivirus* (APMV) and all group I members of the *Mimiviridae* (29). Phylogenetic analyses suggested that the Rab GTPase was acquired by an ancestor of the *Mimiviridae* family and Rabs from *Mimiviridae*, *Plasmodium* and few lower eukaryotes form a separate clade (29). Thus, *Legionella* and APMV that both infect the protozoa *Acanthamoeba* encode Rab proteins most likely to mimic and subvert host cell function. To substantiate that these proteins act in the host cell, we determined whether the Rab containing proteins were bona fide substrates of the Dot/Icm T4SS by creating fusion proteins between the 16 different Rab proteins and the catalytic domain of the TEM-1 beta-lactamase (indicated by a star in SI Appendix, Fig. S5). Translocation assays were performed using wild type *L.*

545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612

613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680



**Fig. 3.** Domain organization of small GTPases in *Legionella* and phylogenetic analyses of the Llo3288 Rab proteins suggests eukaryotic origin. **A)** Domain organization of the different small GTPases proteins identified. **B)** Unrooted tree of Llo3288 and homologues recruited by blastp constructed using likelihood. Local support values are represented with circles on the corresponding branches and size of circles is proportional to the values (only local support of at least 0.7 are shown). **C)** Translocation of selected proteins using the beta-lactamase translocation assay and infection of Raw264.7 cells for 1h with Lp wild type or LpΔdotA expressing BlaM-effector fusions analysed with a microplate reader. Three independent experiments (n=9) were done. Statistical significance was determined by 2-way Anova with multiple comparisons test (\*, P<0.05; \*\*, P<0.01; \*\*\*, P<0.001). **D)** Translocation of selected proteins using the beta-lactamase translocation assay and infection of THP-1 cells at an MOI of 50 during 1h 30min with Lp and Llo strains in before addition of CCF4-AM and analyses by flow cytometry. Histograms show the frequency of BlaM- translocated, blue fluorescence-emitting cells as means ± SD of three independent experiments (n=12). Statistical significance was determined by Wilcoxon matched pairs test (\*\*, P<0.01; \*\*\*, P<0.001). Lp, *L. pneumophila* wild type; Llo, *L. longbeachae* wild type; Lp ΔdotA, *L. pneumophila* ΔdotA; LloΔ, *L. longbeachae* ΔdotA.


*pneumophila* as a surrogate host and compared with an isogenic Dot/Icm mutant ( $\Delta dotA$ ). All 16 Rab motif-containing proteins were translocated by *L. pneumophila* but not by the  $\Delta dotA$  mutant (Fig. 3C-D).

**More than 250 different eukaryotic like proteins are encoded in *Legionella* genomes.** In addition to modular effectors with eukaryotic domains, the *Legionella* genome encodes proteins that are similar to eukaryotic proteins, many of which are proven effectors of the Dot/Icm T4SS. A wider search for eukaryotic like proteins in the *Legionella* genus identified 2196 eukaryotic like proteins representing more than 400 different orthologous groups that matched better to eukaryotes than to prokaryotes from a total of 6809 different orthologous proteins that matched with eukaryotic proteins. Among these, we identified 156 proteins with a eukaryotic domain, and 210 new eukaryotic-like proteins (SI Appendix, Table S5). Furthermore, 152 eukaryotic like proteins detected possess a higher GC content (40%-62%) than the rest of the genome indicating recent HGT. Phylogenetic analysis of

selected, newly identified proteins suggested that these were acquired from eukaryotes. As an example, SI Appendix, Fig. S7 shows the protein LanA0735 from *Legionella anisa*, a species frequently found in artificial water systems. This protein belongs to the pyridine nucleotide-disulfide oxidoreductase family, a sub-family of the FAD dependent oxidoreductase family. LanA0735 showed some similarity to thioredoxin reductase that exists as two major ubiquitous isoenzymes in higher eukaryotic cells, one cytosolic and the other one mitochondrial. The cytosolic form has been implicated in interference with the acidification of the lysosomal compartment in *C. elegans* (30), and thus LanA0735 may help *Legionella* avoid vacuole acidification during infection.

Among the proteins defined as eukaryotic like, two previously described phospholipases of *L. pneumophila*, PlcB (Lpp1411/Lpg1455) and PlcA (Lpp0565/Lpg0502) were identified in our analysis as eukaryotic proteins. The only other bacteria encoding these two enzymes are *Pseudomonas* and amoeba-associated bacteria. The two enzymes have phospholipase activity

681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748



**Fig. 4.** Gain/loss prediction for selected eukaryotic proteins and domain containing proteins. Circles on the branches represent gain events, crosses loss events. The full squares, circles, triangles or stars indicate the presence of the respective protein; the empty squares, circles, triangles or stars indicate that the protein is absent in this species.

(31), but their role in infection is unknown. Here they were predicted as phosphatidylcholine-hydrolyzing phospholipase C. Phosphatidylcholine is a eukaryotic membrane phospholipid that is present in only about 15% of prokaryotic species, in particular bacteria interacting with eukaryotes (32). *L. pneumophila* belongs to the phosphatidylcholine-containing group of bacteria, which includes *Francisella tularensis* or *Brucella abortus* (33). These pathogens use the phosphatidylcholine synthase pathway exclusively for phosphatidylcholine formation and are thought to depend on choline supplied from the host cell (34). Indeed, it has been shown that phosphatidylcholin synthesis is required for *L. pneumophila* virulence (35). Thus, it is tempting to infer that the role of these enzymes may be to help acquire choline from the host cell.

**Evolutionary history of eukaryotic domains and eukaryotic proteins.** It is intriguing that *Legionella* species encode such a diverse repertoire of eukaryotic domains and eukaryotic-like proteins. To understand better this unique feature of the genus we analyzed the evolutionary history of these proteins. After phylogenetic reconstruction of the genus *Legionella* based on the core genome (at least 50% identical) (Fig. 1A), we analyzed the distribution of the eukaryotic motifs and the eukaryotic proteins with respect to the evolution of the genus. For most we found patchy distribution, as the repertoire of these proteins is variable among the different *Legionella* species (Fig. 2). Such a distribution is indicative of gain and loss events during the evolution of the

genus. To analyze further how these proteins may have evolved in *Legionella* we selected 25 eukaryotic motifs representing 2,837 different proteins in over 800 orthologous groups and used the program Gloome to analyze the gain and loss events for these proteins. We found that the number of gain events (1,197/69%) considerably exceeded the number of loss events (549/31%), a bias that was even stronger when using parsimony (1,628 gain events versus 89 loss events) (SI Appendix, Fig. S8). These results were confirmed also when using a more conservative approach by taking a probability cut-off for the stochastic model of 0.8 instead of 0.5, and when analyzing each motif separately.


An exemplary view of this result is shown for four proteins encoding different motifs (U-box and ankyrin repeat, SET domain and ankyrin repeat, astacin domain and allinase domain; Fig. 4). Loss events are indicated by a star and gain events by a dot. The number of gain events exceeds the number of loss events, indicating that in the *Legionella* genus gene acquisition is dominant. Moreover, gene acquisition seems to be an ongoing and frequent process in the genus *Legionella* given the high number of events we observed and the fact that most of them are localized in the terminal branches of the tree (SI Appendix, Fig. S8). To analyse if eukaryotic-like proteins have the same evolutionary history, we took the sphingosine1-phosphate lyse (*LpSpl*) (36, 37) as an example. Indeed, when running the same analyses this gene also appeared to have been gained multiple times during the evolution of the genus (Fig. 4).

Thus, in comparison to most prokaryotic species analysed to date, more gene gain events are evident than loss events during evolution of the *Legionella* genus, which is also corroborated by the fact that the ancestral genomes were probably smaller (Fig. 1A, cluster I). Indeed, as seen in Fig. 1A, in each of the defined phylogenetic clusters only few genomes have a larger size e.g. in cluster II *L. massiliensis* is the only species with a big genome, thus the most parsimonious explanation is that the ancestor of this clade had a small genome and in the branch leading to *L. massiliensis* gene gain occurred. This finding is similar to what was described for the adaptation of louse-borne intracellular pathogens and amoeba associated bacteria. It is well known that the specialization of intracellular bacteria is associated with genome reduction, and extreme genome reduction can be seen in louse-borne human specialists. In contrast, nonspecialized intra-amoebal microorganisms exhibit a genome larger than their relatives due to gene conservation and acquisition (38).

**The Dot/Icm secretion system is a highly conserved machinery secreting thousands of different proteins.** The Dot/Icm T4SS is indispensable for intracellular replication of *L. pneumophila* in both amoeba and macrophages (39). In stark contrast to the high genetic diversity observed in the *Legionella* genomes, the Dot/Icm T4SS is part of the core genome as it is present in all species analyzed and the organization of the constituent proteins is highly conserved, even at the amino acid level. The proteins comprising the secretion machinery show an average amino acid identity of more than 50% and some even more than 90% when compared to the *L. pneumophila* Dot/Icm components (SI Appendix Fig. S9A and Table S6). The most conserved proteins are DotB, a secretion ATPase (86-100% aa identity) and IcmS, a small acidic cytoplasmic protein (74-98% aa identity). This high conservation is even seen with one of the few non-*Legionella* species that encode a Dot/Icm system, *Coxiella burnetii*.

The only gene of the Dot/Icm system that is not present in all *Legionella* species is *icmR*. IcmR interacts with IcmQ as a chaperone preventing IcmQ self-dimerization (40). Although IcmQ is highly conserved, the gene encoding IcmR is frequently replaced by one or two non-homologous genes encoding for proteins that are called FIR because they can functionally replace IcmR (41). When overlapping the occurrence of the different FIR genes with the phylogeny of the species, most phylogenetically closely related

749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816



**Fig. 5.** The replicative capacity of the different *Legionella* species in THP-1 cells correlates with their epidemiological features. Replication of each strain at the time point 72h after infection of THP-1 cells is shown (24h and 48h of infection are shown in SI Appendix, Fig. S14. Intracellular replication was determined by recording the number of colony-forming units (CFU) after plating on BCYE agar. *L. pneumophila* Paris, representative of a replicating strain (blue box); *L. pneumophila*  $\Delta$ dotA, representative of non-replicating strain (red box). The strains are ordered according to the mean replication values. A) *Legionella* species replicating like or significantly better than *L. pneumophila* Paris. B) Species with no or significantly lower replication capacities than *L. pneumophila* Paris.

species share homologous FIR genes (SI Appendix, Fig. S10). Apart from two conserved regions (SI Appendix, Fig. S11), the absence of sequence homology among FIR proteins indicates that *icmR* is an extremely fast evolving gene and therefore probably under positive selection. The reason why this gene is extremely divergent is still unknown but could be also linked to the high variety of Dot/Icm effectors described in this genus. Thus, except for the FIR genes, the Dot/Icm T4SS is highly conserved and encoded in a very dynamic genetic context.

It has been shown previously, that the more than 300 substrates of the *L. pneumophila* Dot/Icm system are not universally present within the genus *Legionella* as among 38 *Legionella* species only seven core effectors had been described (14). Surprisingly, when adding the 40 additional genomes and 16 new *Legionella* species sequenced in this study, we identified 8 core effectors instead of seven. A comparison of the two studies confirmed Lpg0103 (VipF), Lpg0107 (RavC), Lpg2300 (LegA3/AnkH/AnkW), and Lpg2815 (IroT/MavN) as core substrates (14) (SI Appendix, Fig. S9B and Table S7). Three of the previously defined core substrates (Lpg0140, Lpg2832, Lpg3000) were present in two genomes as two consecutive genes instead of one, however, this fragmentation might be a sequencing error, and thus we considered these substrates also as core substrates (SI Appendix Table S7). In our study we identified one additional core effector gene, *lpg1356/lpp1310*. This protein has been reported by Lifshitz and colleagues (42) as secreted protein, but had not been included in the Burstein effector search, which explains the different result (SI Appendix, Fig. S9B and Table S7). Similarly, to most of the other core substrates, their functions are not known, but Lpg1356 encodes eight eukaryotic Sel-1 motifs similar to LpnE, a *L. pneumophila* virulence determinant that influences vacuolar trafficking (43). Furthermore, seven other genes are present in all but one, two or four genomes, thus they might have important functions in host pathogen interactions (SI Appendix Table S7). Interestingly, when the effector repertoire of several strains of one species is compared the conservation of the effectors is very high (between 82 and 97%) (SI Appendix Table S8). However, if more strains than two are available for a species as it is the case for *L. pneumophila* where 11 strains could be compared, the conservation of the effector pool is only 65% (264 of the 408 different effectors identified in the 11 strains) (SI Appendix Table S8). Thus the *L. pneumophila* core effector set is also smaller than previously thought. Taken together, the genus

*Legionella* has 8 core substrates present in all genomes and seven additional ones that are present in nearly all genomes.

Interestingly, whereas the number of core Dot/Icm substrates is extremely small, the number and the diversity of predicted Dot/Icm substrates is extremely high. Indeed, through a machine learning approach, Burstein *et al* predicted that the *Legionella* genus would encode 5,885 effectors (14). Here we extended these analyses and identified 4,767 proteins with eukaryotic motifs that have a high probability to be secreted effectors as shown for the Rab-like proteins. If we consider that the orthologous of these proteins in each species are also effectors then the number raises to 7103 (representing 1145 different orthologous proteins) (SI Appendix Fig. S9C). Moreover, we identified 2,196 eukaryotic like proteins representing 414 different orthologous genes, which form together with the above-mentioned eukaryotic motif carrying proteins 1,400 different putative orthologous substrates of the Dot/Icm T4SS. Finally, when adding to the effectors predicted in this study (based on their similarity to eukaryotic domains and proteins), the effectors previously described in *L. pneumophila* and their orthologues (more than 7000 proteins representing about 300 different orthologous), as well as the effectors predicted by the machine learning approach and their orthologous (more than 10 000 proteins representing about 900 different orthologous) (14) the total number of different effectors rises to almost 18,000 proteins (more than 1,600 orthologous groups) (SI Appendix, Table S9 and Fig. S9C). Therefore, the *Legionella* genus has by far the highest number and widest variety of effectors described for an intracellular bacterium. Furthermore, when calculating the growth accumulation curve for Dot/Icm predicted effectors, this number should still increase with the sequencing of new *Legionella* genomes, as the plateau is not reached yet (SI Appendix, Fig. S9D).

**The ability to infect human cells has been acquired independently several times during the evolution of the genus *Legionella*.** Among the 65 *Legionella* species known, *L. pneumophila* is responsible for over 90% of human disease, followed by *L. longbeachae* (2-7% of cases, except Australia and New Zealand with 30% (44)). Certain *Legionella* species such as *L. micdadei*, *L. dumoffii* or *L. bozemanii* have once or sporadically been associated with human disease (44), and all other species seem to be environmental bacteria only. The reasons for these differences are not known. To explore whether all species are able to replicate in human cells we chose the human macrophage like cell line

THP-1 as model and tested the replication capacity of 47 different *Legionella* species. Infections were carried out in duplicates or triplicates and colony-forming units were recorded at 24h, 48h and 72h post infection. Levels of intracellular replication were compared to wild type *L. pneumophila* strain Paris and an isogenic non-replicating  $\Delta dotA$  mutant as reference strains (Fig. 5 and SI Appendix, Fig. S12 and S13). Results were also compared to data previously reported for different *Legionella* species in THP-1, U937 and A549 cells, Mono Mac 6, mouse and guinea pig derived macrophages, or in guinea pigs (SI Appendix, Table S10). When results at 72 h after infection were analyzed, 28 of the 47 species tested were impaired for intracellular replication whereas nine species replicated similarly to *L. pneumophila* Paris or better (Fig. 5). These nine species were *L. gormanii*, *L. jamestowniensis*, *L. jordanis*, *L. like brunensis*, *L. maceachernii*, *L. micdadei*, *L. nagasakiensis*, *L. parisiensis*, and *L. tucsonensis*. Interestingly, *L. jamestowniensis*, for which one human case has been reported (45), replicated better than *L. pneumophila* Paris. Indeed, *L. jamestowniensis* productively infects human U937-derived phagocytes. The remaining eight species showed variable replication patterns being significantly different from *L. pneumophila* Paris only in one or two of the three analyzed time points (SI Appendix, Fig. S12). Broadly, the species most frequently reported from human disease (*L. pneumophila*, *L. longbeachae*, *L. micdadei*, *L. bozemanii* and *L. dumoffii*) are also those that replicated robustly in THP-1 cells. The only exception was the *L. dumoffii* strains that were impaired for replication in THP-1 cells but which have been shown to replicate in other cell types and guinea pigs. Taken together, there is a convincing correlation between the frequency of isolation from human disease and the ability to grow in macrophage-like cells.

To analyze this further, we overlapped the replication results with the phylogeny of the genus. Apart from the small cluster containing *L. beliardensis*, *L. gresilensis* and *L. busanensis*, which were all unable to grow in THP-1 cells, replicating and non-replicating strains were mixed in the phylogeny (SI Appendix, Fig. S14). This suggests that the capacity to replicate in human cells has been acquired independently several times during evolution of the *Legionella* genus, possibly as a result of recruiting effectors that allow adaptation to particular niches. To understand whether a specific set of effectors is necessary to infect human cells, we further analyzed the combination of effectors present in the strains isolated from human disease and effectors present in strains capable of replicating in THP-1 cells. Surprisingly, no specific set of effectors could be attributed to strains capable of replicating in human cells or isolated from human disease, although among these strains certain conserved motifs always present were identified, such as ankyrin motifs, F-box or SET-domains, suggesting that common pathways need to be subverted to cause human infection. Thus, the capacity to infect human cells has been acquired independently, several times during the evolution of the genus *Legionella*.

In conclusion, the analysis of 80 *Legionella* strains representing 58 different *Legionella* species has revealed a contrasting picture of the *Legionella* genus. It encodes a highly conserved T4SS predicted to secrete more than 18,000 proteins, of which only 8 are conserved throughout the genus. Together the genomes portray an extremely diverse genus shaped by massive inter-domain horizontal gene transfer, circulating mobile genetic elements and eukaryotic like proteins. Our in-depth analyses of eukaryotic features of the *Legionella* genomes identified 137 different eukaryotic domains of which Rab or Ras domain-containing proteins were quasi unique to the genus *Legionella*. The secretion assays

undertaken for 16 of these Rab or Ras domain-containing proteins confirmed that these were translocated Dot/Icm effectors. In addition to the eukaryotic domains, we identified 210 orthologous groups of eukaryotic like proteins. If all these proteins in the different species and their orthologues are taken into account, we found more than 8,000 proteins that have been shaped by inter-domain horizontal gene transfer in the genus *Legionella*. Thus, to our knowledge the genus *Legionella* contains the widest variety and highest number of eukaryotic proteins and domains of any prokaryotic genus genome analyzed to date. Analyzing more strains per species will probably discover new unknown effectors increasing our knowledge of the set of tools used by *Legionella* to infect eukaryotic cells. Although eukaryotic proteins and domains were a universal feature of the genus *Legionella*, the repertoire of these proteins for each species was different. Surprisingly, even when the same motif was present in different species, these were often present in different proteins with no orthology. In accordance with this finding, our evolutionary analysis of the presence/absence of these domains and proteins suggested that these proteins were mostly acquired through gene gain events.

When exploring the replication capacity of 47 different *Legionella* species in human macrophage-like cell line THP-1, we found that the 23 species were capable of replicating in THP-1 cells. However, these did not cluster in the phylogeny, indicating that the capacity to replicate in macrophages can be achieved by different combinations of effectors, and this capacity has been acquired several times during the evolution of the *Legionella* genus. As humans are an accidental host for *Legionella*, the capacity to replicate in macrophages may also have been obtained by a coincidental acquisition of different virulence properties initially needed to adapt to a specific natural host, such as amoebae. Indeed, due to the high conservation of key signaling pathways in professional phagocytes such as amoebae and human macrophages, different combinations of effectors may allow *Legionella* species to infect higher eukaryotic cells by chance.

Here we show that all *Legionella* species have acquired eukaryotic proteins that likely modulate specific host functions to allow intracellular survival and replication in eukaryotic host cells. At a certain point, the evolution of a combination of effector proteins that allow replication in human cells may inadvertently lead to the emergence of new human pathogens from environmental bacteria.

#### Material and Methods

The materials and methods are described at length in SI Appendix. This includes: Sequencing and assembly, sequence processing and annotation, pan/core genome, ortholog and singleton definition, phylogenetic reconstruction and evolutionary analysis, phylogenetic analyses of Rab and eukaryotic-like proteins, infection assays, statistical analysis, and translocation assays. The raw sequence reads were deposited in the European Nucleotide Archive (study accession number PRJEB24896). The sequences and annotations can be accessed through: [https://github.com/bbi-ip/Legionella\\_genus.proteins.git](https://github.com/bbi-ip/Legionella_genus.proteins.git)

#### Acknowledgements

We would like to thank Tim P. Stinear for critical reading of the manuscript and helpful comments and we acknowledge the receipt of 53 different *Legionella* strains from the Collection of the Institut Pasteur (CIP). Work in the CB laboratory is financed by the Institut Pasteur, the grant n°ANR-10-LABX-62-IBEID and the Fondation pour la Recherche Médicale (FRM) grant N° DEQ20120323697. **Author contributions** SJ and LGV, ELH contributed to sample collection and strain analyses DNA extraction and sequencing; CR, DC, SM, AEPC, MR, SP, SR, JD, JC, SDD, GNS to functional experiments, data analyses and interpretation. The manuscript was written by LGV and CB with input from co-authors. The project was conceived, planned and supervised by LGV, GD, GF and CB.

freshwater and soil amoebae. *J Clin Pathol* 33(12):1179-1183.

3. Cazalet C, et al. (2004) Evidence in the *Legionella pneumophila* genome for exploitation of host cell functions and high genome plasticity. *Nat Genet* 36(11):1165-1173.

1. Fraser DW, et al. (1977) Legionnaires' disease: description of an epidemic of pneumonia. *N Engl J Med* 297(22):1189-1197.
2. Rowbotham TJ (1980) Preliminary report on the pathogenicity of *Legionella pneumophila* for



1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156

4. Bruggemann H, Cazalet C, & Buchrieser C (2006) Adaptation of *Legionella pneumophila* to the host environment: role of protein secretion, effectors and eukaryotic-like proteins. *Curr Opin Microbiol* 9(1):86-94.
5. Komano T, Yoshida T, Narahara K, & Furuya N (2000) The transfer region of Inc11 plasmid R64: similarities between R64 tra and *Legionella icm/dot* genes. *Mol Microbiol* 35(6):1348-1359.
6. Escoll P, Mondino S, Rolando M, & Buchrieser C (2016) Targeting of host organelles by pathogenic bacteria: a sophisticated subversion strategy. *Nat Rev Microbiol* 14(1):5-19.
7. Finsel I & Hilbi H (2015) Formation of a pathogen vacuole according to *Legionella pneumophila*: how to kill one bird with many stones. *Cell Microbiol* 17(7):935-950.
8. Nora T, Lomma M, Gomez-Valero L, & Buchrieser C (2009) Molecular mimicry: an important virulence strategy employed by *Legionella pneumophila* to subvert host functions. *Future Microbiol* 4:691-701.
9. Burstein D, et al. (2009) Genome-scale identification of *Legionella pneumophila* effectors using a machine learning approach. *PLoS Pathog* 5(7):e1000508.
10. Gomez-Valero L, et al. (2011) Extensive recombination events and horizontal gene transfer shaped the *Legionella pneumophila* genomes. *BMC Genomics* 12:536.
11. Gomez-Valero L, et al. (2014) Comparative analyses of *Legionella* species identifies genetic features of strains causing Legionnaires' disease. *Genome Biol* 15(11):505.
12. Morozova I, et al. (2004) Comparative sequence analysis of the *icm/dot* genes in *Legionella*. *Plasmid* 51(2):127-147.
13. Sanchez-Buso L, Comas I, Jorques G, & Gonzalez-Candelas F (2014) Recombination drives genome evolution in outbreak-related *Legionella pneumophila* isolates. *Nat Genet* 46(11):1205-1211.
14. Burstein D, et al. (2016) Genomic analysis of 38 *Legionella* species identifies large and diverse effector repertoires. *Nat Genet* 48(2):167-175.
15. Joseph SJ, et al. (2016) Dynamics of genome change among *Legionella* species. *Sci Rep* 6:33442.
16. Cazalet C, et al. (2010) Analysis of the *Legionella longbeachae* genome and transcriptome uncovers unique strategies to cause Legionnaires' disease. *PLoS Genet* 6(2):e1000851.
17. Guglielmini J, de la Cruz F, & Rocha EP (2013) Evolution of conjugation and type IV secretion systems. *Mol Biol Evol* 30(2):315-331.
18. Bohlin J, Brynildsrud OB, Sekse C, & Snipen L (2014) An evolutionary analysis of genome expansion and pathogenicity in *Escherichia coli*. *BMC Genomics* 15:882.
19. Bohlin J, Sekse C, Skjerve E, & Brynildsrud O (2014) Positive correlations between genomic %AT and genome size within strains of bacterial species. *Environ Microbiol Rep* 6(3):278-286.
20. Rolando M, et al. (2013) *Legionella pneumophila* effector RomA uniquely modifies host chromatin to repress gene expression and promote intracellular bacterial replication. *Cell Host Microbe* 13(4):395-405.
21. Nagai H, Kagan JC, Zhu X, Kahn RA, & Roy CR (2002) A bacterial guanine nucleotide exchange factor activates ARF on *Legionella* phagosomes. *Science* 295(5555):679-682.
22. Pasanen AL, Yli-Pietila K, Pasanen P, Kalliokoski P, & Tarhanen J (1999) Ergosterol content in various fungal species and biocontaminated building materials. *Appl Environ Microbiol* 65(1):138-142.
23. Smith FR & Korn ED (1968) 7-Dehydrostigmaterol and ergosterol: the major sterols of an amoeba. *J Lipid Res* 9(4):405-408.
24. Thomson S, et al. (2017) Characterisation of sterol biosynthesis and validation of 14alpha-demethylase as a drug target in *Acanthamoeba*. *Sci Rep* 7(1):8247.
25. Wennerberg K, Rossman KL, & Der CJ (2005) The Ras superfamily at a glance. *J Cell Sci* 118(Pt 5):843-846.
26. Simanshu DK, Nissley DV, & McCormick F (2017) RAS Proteins and Their Regulators in Human Disease. *Cell* 170(1):17-33.
27. Rojas AM, Fuentes G, Rausell A, & Valencia A (2012) The Ras protein superfamily: evolutionary tree and role of conserved amino acids. *J Cell Biol* 196(2):189-201.
28. Colicelli J (2004) Human RAS superfamily proteins and related GTPases. *Sci STKE* 2004(250):RE13.
29. Zade A, Sengupta M, & Kondabagil K (2015) Extensive *in silico* analysis of Mimivirus coded Rab GTPase homolog suggests a possible role in virion membrane biogenesis. *Front Microbiol* 6:929.
30. Li W, et al. (2012) Two thioredoxin reductases, trxr-1 and trxr-2, have differential physiological roles in *Caenorhabditis elegans*. *Mol Cells* 34(2):209-218.
31. Hiller M, Lang C, Michel W, & Flieger A (2017) Secreted phospholipases of the lung pathogen *Legionella pneumophila*. *Int J Med Microbiol*.
32. Aktas M, et al. (2010) Phosphatidylcholine biosynthesis and its significance in bacteria interacting with eukaryotic cells. *Eur J Cell Biol* 89(12):888-894.
33. Geiger O, Lopez-Lara IM, & Sohlenkamp C (2013) Phosphatidylcholine biosynthesis and function in bacteria. *Biochim Biophys Acta* 1831(3):503-513.
34. Comerici DJ, Altabe S, de Mendoza D, & Ugalde RA (2006) *Brucella abortus* synthesizes phosphatidylcholine from choline provided by the host. *J Bacteriol* 188(5):1929-1934.
35. Conover GM, et al. (2008) Phosphatidylcholine synthesis is required for optimal function of *Legionella pneumophila* virulence determinants. *Cell Microbiol* 10(2):514-528.
36. Degtyar E, Zusman T, Ehrlich M, & Segal G (2009) A *Legionella* effector acquired from protozoa is involved in sphingolipids metabolism and is targeted to the host cell mitochondria. *Cell Microbiol* 11(8):1219-1235.
37. Rolando M, et al. (2016) *Legionella pneumophila* S1P-lyase targets host sphingolipid metabolism and restrains autophagy. *Proc Natl Acad Sci U S A* 113(7):1901-1906.
38. Moliner C, Fournier PE, & Raoult D (2010) Genome analysis of microorganisms living in amoebae reveals a melting pot of evolution. *FEMS Microbiol Rev* 34(3):281-294.
39. Segal G & Shuman HA (1999) *Legionella pneumophila* utilizes the same genes to multiply within *Acanthamoeba castellanii* and human macrophages. *Infect Immun* 67(5):2117-2124.
40. Dumenil G & Isberg RR (2001) The *Legionella pneumophila* IcmR protein exhibits chaperone activity for IcmQ by preventing its participation in high-molecular-weight complexes. *Mol Microbiol* 40(5):1113-1127.
41. Feldman M, Zusman T, Hagag S, & Segal G (2005) Coevolution between nonhomologous but functionally similar proteins and their conserved partners in the *Legionella* pathogenesis system. *Proc Natl Acad Sci U S A* 102(34):12206-12211.
42. Lifshitz Z, et al. (2013) Computational modeling and experimental validation of the *Legionella* and *Coxiella* virulence-related type-IVB secretion signal. *Proc Natl Acad Sci U S A* 110(8):E707-715.
43. Newton HJ, et al. (2007) Sell repeat protein LpnE is a *Legionella pneumophila* virulence determinant that influences vacuolar trafficking. *Infect Immun* 75(12):5575-5585.
44. Yu VL, et al. (2002) Distribution of *Legionella* species and serogroups isolated by culture in patients with sporadic community-acquired legionellosis: an international collaborative survey. *J Infect Dis* 186(1):127-128.
45. Prochazka B, et al. (2016) Draft Genome Sequence of *Legionella jamestowniensis* Isolated from a Patient with Chronic Respiratory Disease. *Genome announcements* 4(5).

1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224