

MORITA EQUIVALENCE FOR RINGS WITHOUT IDENTITY

By

P. N. ÁNH and L. MÁRKI

In the paper [1] Abrams made a first step in extending the theory of Morita equivalence to rings without identity. He considered rings in which a set of commuting idempotents is given such that every element of the ring admits one of these idempotents as a two-sided unit, and the categories of all left modules over these rings which are unitary in a natural sense. He proved that two such module categories over the rings R and S , say, are equivalent if and only if there exists a unitary left R -module P which is a generator, the direct limit of a given kind of system of finitely generated projective modules, and such that S is isomorphic to the ring of certain endomorphisms of P .

The aim of the present paper is to extend this theory in two ways: to cover a wider range of rings, and to transfer more of the classical Morita theory. Firstly, one can weaken the condition of commutativity of the idempotents in question: it suffices to require that any two of them have a common upper bound under the natural partial order (i. e., any two elements of the ring admit a common two-sided identity), a condition which is fulfilled by all regular rings (regular in the sense of Neumann). Whenever one has such a system of idempotents, then any larger system, in particular, the set of all idempotents, is also such, which is not the case for the systems of Abrams. Secondly, by a suitable modification of some homological lemmas we obtain also the two-sided characterizations of Morita equivalence, arriving thus at a complete analogy to the classical case of rings with identity. Our presentation is a combination of those in Anderson-Fuller [2], §§ 21-22, and Bass [5] (see also [6], Chapter II). This machinery allows us to avoid the elaborate construction of Abrams. As examples we describe, among others, those rings with local units which are Morita equivalent to division rings and primary rings, respectively. The Rees matrix rings studied in [4] turn out to have a natural place in this theory.

The theory we present here is a counterpart of the theory of Morita duality developed by Yamagata [10]. On the one hand, we shall use the same modified Hom-functors but for projective and not injective modules, and on the

other hand, it turns out that every Morita equivalence class of rings with local units contains rings with enough idempotents, i. e., rings considered by Yamagata.

Notice also that the module categories we consider are full subcategories of the categories of modules over unital overrings of the respective rings with local units. Nevertheless, Sato's [8] theory of equivalence does not apply because he considers the usual Hom-functors, which does not work in our case.

1. Preparations.

DEFINITION 1. R is a *ring with local units* if every finite subset of R is contained in a subring of the form eRe where $e=e^2 \in R$.

We call a left module M over R *unitary* if $RM=M$, i. e., for each $m \in M$ there are $r_1, \dots, r_n \in R$ and $m_1, \dots, m_n \in M$ such that $r_1m_1 + \dots + r_nm_n = m$. If R is a ring with local units then this implies that for every finite subset $M' \subset M$ there is an idempotent $e \in R$ such that $em=m$ for all $m \in M'$. By $R \text{ Mod}$ we denote the category of unitary left R -modules together with the usual R -homomorphisms. Dually, $\text{Mod } R$ denotes the category of unitary right R -modules. Similarly to the case considered in Abrams [1], $R \text{ Mod}$ (or $\text{Mod } R$) is a complete and cocomplete additive category. We call a bimodule unitary if it is unitary on both sides.

In what follows, R denotes a ring with local units. The most important thing for us is to find those modules in $R \text{ Mod}$ which play the role of the progenerators in the case of rings with identity. Of course, projective generators make sense in $R \text{ Mod}$ for this a categorical notion; however, ${}_R R$ is neither finitely generated nor projective if R has no identity, and the notion we need ought to include ${}_R R$, too. Therefore we define:

DEFINITION 2. $P \in R \text{ Mod}$ is a *locally projective module* if there is a direct system $(P_i)_{i \in I}$ of finitely generated projective direct summands of P together with projections $\phi_i: P \rightarrow P_i$ such that ϕ_i factors through ϕ_j whenever $i \leq j$, and such that $\varinjlim P_i = P$. Notice that ${}_R R$ is locally projective if R has local units, as Re is a projective direct summand of R for every idempotent $e \in R$, and the multiplication maps $\phi_e: R \rightarrow Re$ satisfy the condition on ϕ_i if we define $e \leq f \Leftrightarrow ef = fe = e$.

The role of progenerators will be played by the locally projective generators in $R \text{ Mod}$. But before turning to them, we shall establish homological properties of locally projective modules. In doing so, we shall need a more restrictive notion instead of the Hom-sets. For the sake of convenience, homomorphisms of modules will be written opposite the scalars.

Notice that the usual definition of tensor product makes no use of the identity in the ring, hence it makes sense in our case, too.

The following Propositions 1.1-1.5 and 1.7 can be proved along the same lines as in the case of rings with identity (see e. g. Anderson-Fuller [2], §20), therefore we present them without proof. Before stating Proposition 1.1, observe the following. If R and S are rings with local units and ${}_S N$ and ${}_S U_R$ are unitary then $\text{Hom}_S(U, N)$ is a left R -module by putting, for $\phi \in \text{Hom}_S(U, N)$ and $r \in R$, $r\phi : u \in U \mapsto (ur)\phi \in N$. The submodule $R \text{Hom}_S(U, N)$ is the largest unitary R -submodule of $\text{Hom}_S(U, N)$. By $R \text{Hom}_S(U, -)$ we denote the functor induced by the mapping $N \mapsto R \text{Hom}_S(U, N)$.

PROPOSITION 1.1. For all $M, M' \in R \text{Mod}$, $m \in M$, $\phi \in \text{Hom}_R(M, M')$, put

$$m\rho_M : r \mapsto mr \quad (r \in R) \text{ (thus } \rho : M \rightarrow R \text{Hom}_R(R, M))$$

and

$$\rho_\phi : \gamma \mapsto \gamma \circ \phi \quad (\gamma \in R \text{Hom}_R(R, M)).$$

Then $\rho : {}_R \text{Mod} \rightarrow R \text{Hom}_R(R, -)$ is a natural isomorphism.

PROPOSITION 1.2. For all $M, M' \in R \text{Mod}$ and $\phi \in \text{Hom}_R(M, M')$, put

$$(r \otimes m)\mu_M = rm \quad (r \in R, m \in M) \text{ (thus } \mu_M : R \otimes M \rightarrow M)$$

and

$$\mu_M : R \otimes \phi \mapsto \phi.$$

Then $\mu : R \otimes_R - \rightarrow {}_R \text{Mod}$ is a natural isomorphism.

COROLLARY 1.3. For all $e^2 = e \in R$ and $M \in R \text{Mod}$, $eR \otimes M \cong eM$.

PROPOSITION 1.4. Let $\theta : {}_R U_S \rightarrow {}_R V_S$ be a bimodule homomorphism between unitary bimodules ${}_R U_S$ and ${}_R V_S$ where R and S are rings with local units. Put, for all $M, M' \in R \text{Mod}$ and $\phi \in \text{Hom}_R(M, M')$,

$$\eta_M : \gamma \mapsto \theta \circ \gamma \quad (\gamma \in S \text{Hom}_R(V, M)),$$

$$\eta_\phi : \delta \mapsto \theta \circ \delta \quad (\delta \in S \text{Hom}_R(V, \phi)).$$

Then $\eta : S \text{Hom}_R(V, -) \rightarrow S \text{Hom}_R(U, -)$ is a natural transformation between two functors from $R \text{Mod}$ to $S \text{Mod}$. Moreover, if θ is an isomorphism then η is a natural isomorphism.

Before stating the next proposition, observe the following. If N_S and ${}_R U_S$ are unitary, then $\text{Hom}_S(N, U)$ is a left R -module if we put, for all $\phi \in \text{Hom}_S(N, U)$ and $r \in R$, $r\phi : n \in N \mapsto r(\phi n)$. Then $R \text{Hom}_S(N, U)$ is a unitary

left R -module. Similarly, if ${}_R M$ is any unitary left R -module then $\text{Hom}_R(M, U)$ is naturally a right S -module and hence $\text{Hom}_R(M, U)S$ is a unitary right S -module. Further, notice that ${}_S(K \otimes_R M) \in S \text{ Mod}$ whenever ${}_S K_R$ and ${}_R M$ are unitary modules.

PROPOSITION 1.5. *For every triple $({}_R P, {}_R U_S, {}_S M)$ such that ${}_R P$ is a finitely generated projective module, there is an isomorphism of abelian groups*

$$\eta : \text{Hom}_R(P, U) \otimes_S M \rightarrow \text{Hom}_R(P, U \otimes_S M)$$

defined via

$$\eta(\gamma \otimes m) : p \mapsto p\gamma \otimes m$$

that is natural in each of the three variables P, U, M .

COROLLARY 1.6. *For every triple of unitary modules $({}_R P_S, {}_R U_S, {}_S M)$ such that ${}_R P$ is locally projective and Pf is a finitely generated left R -module for all $f^2 = f \in S$, there is an isomorphism of left S -modules*

$$\eta : S \text{ Hom}_R(P, U) S \otimes_S M \rightarrow S \text{ Hom}_R(P, U \otimes_S M)$$

defined via

$$(\gamma \otimes m)\eta : p \mapsto p\gamma \otimes m$$

that is natural in each of the three variables P, U, M .

PROOF. It is routine to verify that η is a homomorphism which is natural in each variable. Next we show that η is injective. In fact, assume $(\sum \gamma_i \otimes m_i)\eta = 0$. Since $\gamma_i \in S \text{ Hom}_R(P, U)S$, there is an idempotent $f^2 = f \in S$ with $f\gamma_i f = \gamma_i$ for all i . By assumption the left R -submodule Pf is finitely generated, hence it is contained in a finitely generated projective direct summand P' of P . By $P' = P'f \oplus P'(1-f) = Pf \oplus P'(1-f)$, where $P'(1-f) = \{p \in P' \mid P'f = 0\}$, we obtain that Pf is also projective. By $(\sum \gamma_i \otimes m_i)\eta = 0$, the homomorphism $\phi' : Pf \rightarrow U \otimes_S M : pf \mapsto \sum p\gamma_i \otimes m_i$ is trivial. Therefore by Proposition 1.5 we have $\sum \gamma_i \otimes m_i = 0$ in $\text{Hom}_R(Pf, U) \otimes_S M$, but $\text{Hom}_R(Pf, U) = f \text{ Hom}_R(P, U)S$, hence $\sum \gamma_i \otimes m_i$ must be zero in $S \text{ Hom}_R(P, U)S \otimes_S M$. For proving the surjectivity of η , if ϕ is any element in $S \text{ Hom}_R(P, U \otimes_S M)$, then there is an idempotent $f^2 = f \in S$ with $f\phi = \phi$. Consider the restriction ϕ' of ϕ to Pf which is a finitely generated projective direct summand of ${}_R P$. By Proposition 1.5, there is an element $\sum \gamma'_i \otimes m_i$ of $\text{Hom}_R(Pf, U) \otimes_S M$ which corresponds to ϕ' . Extend γ'_i to a γ_i defined on P by putting $(P(1-f))\gamma_i = 0$. Now it is clear that $(\sum \gamma_i \otimes m_i)\eta = f\phi = \phi$, and we are done.

PROPOSITION 1.7. *Let P_S , ${}_R U_S$ and ${}_R M$ be unitary modules such that P_S is finitely generated and projective. Then there is an isomorphism of abelian groups*

$$\eta : P \otimes_S \text{Hom}_R(U, M) \rightarrow \text{Hom}_R(\text{Hom}_S(P, U), M)$$

defined via

$$\eta(p \otimes \gamma) : \delta \in \text{Hom}_S(P, U) \mapsto (\delta p) \gamma$$

that is natural in each of the three variables P, U, M .

COROLLARY 1.8. *Let ${}_R P_S$, ${}_R U_S$ and ${}_R M$ be unitary modules such that P_S is locally projective and eP is a finitely generated right S -module for all idempotents $e^2 = e \in R$. Then there is an isomorphism of left R -modules*

$$\eta : P \otimes_S \text{Hom}_R(U, M) \rightarrow R \text{Hom}_R(R \text{Hom}_S(P, U)R, M)$$

defined via

$$(p \otimes \gamma) \eta : \delta \in R \text{Hom}_S(P, U)R \mapsto (\delta p) \gamma$$

that is natural in each of the three variables P, U, M .

PROOF. It is routine to verify that η is a homomorphism which is natural in P, U and M . Assume now $(\sum p_i \otimes \gamma_i) \eta = 0$. Since ${}_R P$ is unitary, there is an idempotent $e \in R$ with $e p_i = p_i$ for all i . From our assumption it follows that eP is a finitely generated projective direct summand of P_S . Since $(\sum p_i \otimes \gamma_i) \eta = 0$, the element ϕ of $\text{Hom}_R(\text{Hom}_S(eP, U), M)$ defined by $\delta \phi = \sum (\delta p_i) \gamma_i$, $\delta \in \text{Hom}_S(eP, U)$, is zero. Hence we can apply Proposition 1.7 and obtain that the element $\sum p_i \otimes \gamma_i$ is zero in $eP \otimes_S \text{Hom}_R(U, M)$ and therefore it must be zero in $P \otimes_S \text{Hom}_R(U, M)$, too. The surjectivity of η is seen as in the proof of Corollary 1.6.

COROLLARY 1.9. *Let ${}_R P_S$ and ${}_S N$ be unitary modules such that P_S is locally projective and eP is a finitely generated right S -module for every idempotent $e \in R$. Then there is an isomorphism of left R -modules*

$$\eta : P \otimes_S N \rightarrow R \text{Hom}_S(\text{Hom}_S(P, S)R, N)$$

defined via

$$(p \otimes n) \eta : \delta \in \text{Hom}_S(P, S)R \mapsto (\delta p) n$$

that is natural in P and N .

PROOF. Putting $R=S$, $U=S$ and $N=M$ in Proposition 1.7, we obtain that $\eta : P \otimes_S N \rightarrow \text{Hom}_S(\text{Hom}_S(P, S), N)$ is an isomorphism of abelian groups which is

natural in P and N , provided that P_S is finitely generated and projective. In the general case, it is straightforward to check that η is a homomorphism of left R -modules which is natural in P and N . Next we show that η is injective. In fact, assume $(\sum p_i \otimes n_i)\eta = 0$. Since ${}_R P$ is unitary, there is an idempotent $e \in R$ such that $ep_i = p_i$ for all i , and eP is a projective right S -module for P_S is locally projective. Now the element ϕ of $\text{Hom}_S(\text{Hom}_S(eP, S), N)$ defined by $\delta\phi = \sum(\delta p_i)n_i$ is zero since $(\sum p_i \otimes n_i)\eta = 0$, and by the remark made at the beginning of the proof we have that $\sum p_i \otimes n_i$ is the zero element in $eP \otimes_S N$, hence it is the zero in $P \otimes_S N$, too. The surjectivity of η is seen as in the proof of Corollary 1.6.

LEMMA 1.10. *Let ${}_R P$ be a locally projective generator, and let S be a subring of $\text{End}_R P$ having local units such that $P \in \text{Mod } S$, $S \text{End}_R P = S$, and Pf is a finitely generated submodule of ${}_R P$ for every idempotent $f \in S$. Then*

- 1) P_S is a locally projective generator,
- 2) the mapping $\lambda: R \rightarrow \text{End}_S P: r \mapsto \lambda_r$, where $\lambda_r: p \mapsto rp$, is an embedding of R into $\text{End}_S P$ such that $(\text{End}_S P)(\lambda(R)) = \lambda(R)$.

PROOF. By the assumption, P can be considered as a unitary R - S -bimodule. Since ${}_R P$ is a generator, it generates Re for any idempotent $e \in R$, i.e., there are a natural number n and a unitary left R -module P' such that $P^n \cong Re \oplus P'$. Then it follows

$$(\text{End}_R P)^n \cong \text{Hom}_R(P^n, P) \cong \text{Hom}_R(Re, P) \oplus \text{Hom}_R(P', P) \cong eP \oplus \text{Hom}_R(P', P).$$

This fact implies, since $\text{End}_R P$ is a ring with identity, that eP is a finitely generated right $\text{End}_R P$ -module, i.e., there are finitely many elements $p_1, \dots, p_k \in eP$ such that every element $p \in eP$ can be expressed as $p = p_1\phi_1 + \dots + p_k\phi_k$ where $\phi_1, \dots, \phi_k \in \text{End}_R P$. On the other hand, we know that $P \in \text{Mod } S$, hence there is an idempotent $f \in S$ with $p_i f = p_i$ for all i and then $p = p_1(f\phi_1) + \dots + p_k(f\phi_k)$. Since $f\phi_1, \dots, f\phi_k$ are contained in S by the assumption, we see that eP is a finitely generated right S -module. Since eP is a projective right $\text{End}_R P$ -module and every right S -module can be considered as a right $\text{End}_R P$ -module, we deduce immediately that eP is a projective right S -module, too. Furthermore, if $e_1, e_2 \in R$ are idempotents such that $e_1 \leq e_2$, then the map $\phi_{e_1}: P \rightarrow e_1 P: p \mapsto e_1 p$ factors through the corresponding map ϕ_{e_2} . All this shows that P_S is locally projective.

For any idempotent $f \in S$, Pf is a finitely generated projective left R -module by the assumptions, hence there is an idempotent $e \in R$ with $Pf \oplus T \cong (Re)^n$ for a natural number n and a unitary left R -module T . Therefore

$$\begin{aligned} (eP)^n &\cong [\text{Hom}_R(Re, p)]^n \cong \text{Hom}_R((Re)^n, P) \cong \text{Hom}_R(Pf, P) \oplus \text{Hom}_R(T, P) \\ &\cong fS \oplus \text{Hom}_R(T, P) \end{aligned}$$

which implies, since S has local units, that P_S is a generator.

Since ${}_R P$ is a generator, ${}_R R$ is a sum of homomorphic images of ${}_R P$, but R as a ring has local units, and so $\text{ann}_R(P)=0$ must hold. This implies that the mapping λ , which is clearly a homomorphism, is an embedding. In what follows we shall identify R with the subring $\lambda(R)$ of $\text{End}_S P$.

In order to see $(\text{End}_S P)R=R$, take any $\rho \in (\text{End}_S P)R$. Since R has local units, there is an idempotent $e \in R$ such that $\rho e = \rho$. As eP is a finitely generated right S -module, we have $eP = p_1 S + \dots + p_n S$. Let K denote the submodule of ${}_R P^n$ generated by (p_1, \dots, p_n) . Since ${}_R P$ is a generator, K is a sum of homomorphic images of ${}_R P^n$, i.e., $(p_1, \dots, p_n) = x_1 \phi_1 + \dots + x_k \phi_k$ where $x_1, \dots, x_k \in P^n$ and $\phi_1, \dots, \phi_k : P^n \rightarrow K$. As $P \in \text{Mod } S$ and S has local units, each x_i is contained in a $(Pf)^n$, $f^2 = f \in S$, so we can replace each ϕ_i by $f\phi_i$. Now $f\phi_i : P^n \rightarrow K$ can be considered as an $n \times n$ matrix with entries from $f \text{End}_R P$, by one of our assumptions we have $S \text{End}_R P = S$, hence each $f\phi_i$ can be considered as an element of S_n , the ring of $n \times n$ matrices over S . All this shows that $\rho = \rho e$ can be considered as an endomorphism of $(P^n)_{S_n}$ and therefore we have

$$\rho(p_1, \dots, p_n) = \rho(x_1(f\phi_1) + \dots + x_k(f\phi_k)) = (\rho x_1)(f\phi_1) + \dots + (\rho x_k)(f\phi_k),$$

and here $(\rho x_i)(f\phi_i) \in P^n f\phi_i \subseteq K$ for $i=1, \dots, k$. Hence $\rho(p_1, \dots, p_n) \in K = R(p_1, \dots, p_n)$, thus we have that $\rho = \rho e = r e$ for some $r \in R$.

2. The Morita equivalence.

THEOREM 2.1. *Let R and S be equivalent rings with local units via inverse equivalences $G : R \text{ Mod} \rightarrow S \text{ Mod}$ and $H : S \text{ Mod} \rightarrow R \text{ Mod}$. Set*

$$P = H({}_S S) \quad \text{and} \quad Q = G({}_R R).$$

Then P and Q are naturally unitary bimodules ${}_R P_S$ and ${}_S Q_R$ such that

- 1) ${}_R P, P_S, {}_S Q, Q_R$ are locally projective generators and
 $S \text{End}_R P = S = (\text{End}_R Q)S, (\text{End}_S P)R = R = R \text{End}_S Q$;
- 2) ${}_R P_S \cong \text{Hom}_R(Q, R)S \cong R \text{Hom}_S(Q, S), {}_S Q_R \cong \text{Hom}_S(P, S)R \cong S \text{Hom}_R(P, R)$;
- 3) $G \cong S \text{Hom}_R(P, -), H \cong R \text{Hom}_S(Q, -)$;
- 4) $G \cong Q \otimes_R -, H \cong P \otimes_S -$;
- 5) *identifying ${}_S Q_R$ with $S \text{Hom}_R(P, R)$ and S with $S \text{Hom}_R(P, P)$ (see 2 and 1 above), consider the bilinear products*

$$(-, -): P \times Q \rightarrow R : (p, q) = pq \in R,$$

$$\langle -, - \rangle: Q \times P \rightarrow S : \langle q, p \rangle = (-, q)p \in S;$$

then $P \otimes_S Q$ and $Q \otimes_R P$ become rings if we put $(p_1 \otimes q_1)(p_2 \otimes q_2) = p_1 \otimes \langle q_1, p_2 \rangle q_2$ and $(q_1 \otimes p_1)(q_2 \otimes p_2) = q_1 \otimes (p_1, q_2)p_2$, and we have $R \cong P \otimes_S Q$ and $S \cong Q \otimes_R P$.

PROOF. By Proposition 1.1, $R \cong R \text{Hom}_R(R, R)$, moreover, this is also an isomorphism of rings; furthermore, G yields the isomorphism of rings $\text{End}_{R(R)} \cong \text{End}_{S(S)}Q$, and therefore Q can be considered as a right R -module. In order to show that Q_R is unitary, take an arbitrary element $q \in Q$. Since $Q = \cup \{G(Re) : e^2 = e \in R\}$, there is an idempotent $e \in R$ such that $q \in G(Re)$. Now the right translation $\rho_e \in \text{End}_S Q$ induced by e acts as an identity on $G(Re)$, hence $qe = q$. Similarly, P is a unitary right S -module. It is clear that $Qe = G(Re)$ is a finitely generated left S -module.

Since ${}_R R$ is a locally projective generator, the same holds for ${}_S Q$, too. In the same way, ${}_R P$ is a locally projective generator. Now we can apply Lemma 1.10 and obtain that P_S and Q_R are also locally projective generators and it holds $R \text{End}_S Q = R = (\text{End}_S P)R$ and $S = S \text{End}_R P = (\text{End}_R Q)S$, and our first claim is proven.

Next we turn to the proof claim 3). Since G and H are equivalences, for every $M \in R \text{Mod}$ we have the left S -isomorphism

$$\text{Hom}_S(S, G(M)) \cong \text{Hom}_R(H(S), M) = \text{Hom}_R(P, M).$$

Furthermore, by Proposition 1.1, $G(M) \cong S \text{Hom}_S(S, G(M))$ is a natural isomorphism in M . All this shows that $G \cong S \text{Hom}_R(P, -)$ and similarly $H \cong R \text{Hom}_S(Q, -)$, and claim 3) is proven.

Now we have

$${}_S Q_R = {}_S G(R)_R \cong S \text{Hom}_R(P, R) \cong S \text{Hom}_R(P, \text{Hom}_S(P, P)R),$$

and there is also an S - R -bimodule isomorphism η between $S \text{Hom}_R(P, \text{Hom}_S(P, P)R)$ and $\text{Hom}_S(P, S \text{Hom}_R(P, P))R \cong \text{Hom}_S(P, S)R$ defined by

$$\eta(\gamma) \in \text{Hom}_S(P, S \text{Hom}_R(P, P))R : a \mapsto \eta(\gamma)a \in S \text{Hom}_R(P, P) : b \in P \mapsto (a\gamma)b$$

for every element $\gamma \in S \text{Hom}_R(P, \text{Hom}_S(P, P)R)$. (For proving that η is an isomorphism, notice that its inverse is $b\eta^{-1}(\alpha) : a \in P \mapsto a(\alpha b)$.) Hence we get ${}_S Q_R \cong \text{Hom}_S(P, S)R$. Similarly we have ${}_R P_S \cong \text{Hom}_R(Q, R)S \cong R \text{Hom}_S(Q, S)$.

Now Proposition 1.4 and Corollary 1.9 together with claims 2) and 3) proven just above yield

$$H \cong R \operatorname{Hom}_S(Q, -) \cong R \operatorname{Hom}_S(\operatorname{Hom}_S(P, S)R, -) \cong P \otimes_S -$$

and similarly $G \cong Q \otimes_R -$.

To prove 5), consider the mapping

$$\lambda: P \otimes_S Q \rightarrow R: \sum p_i \otimes q_i \mapsto \sum (p_i, q_i).$$

It is clear that λ is a homomorphism of abelian groups. Next,

$$\begin{aligned} \lambda[(p_1 \otimes q_1)(p_2 \otimes q_2)] &= \lambda[p_1 \otimes \langle q_1, p_2 \rangle q_2] = (p_1, \langle q_1, p_2 \rangle q_2) = (p_1 \langle q_1, p_2 \rangle, q_2) \\ &= ((p_1, q_1)p_2, q_2) = (p_1, q_1)(p_2, q_2) = \lambda(p_1 \otimes q_1)\lambda(p_2 \otimes q_2), \end{aligned}$$

hence λ is a ring homomorphism. Since ${}_R P$ is a generator, ${}_R R$ is a sum of homomorphic images of P , so every $r \in R$ can be written as a finite sum $r = \sum p_i \phi_i$, $p_i \in P$, $\phi_i \in \operatorname{Hom}_R(P, R)$. Now P_S is unitary and S has local units, hence there is an idempotent $f \in S$ such that $p_i = p_i f$ for all i . Therefore we can replace ϕ_i by $f \phi_i = q_i \in Q$, and then $r = \sum p_i q_i = \lambda(\sum p_i \otimes q_i)$. Thus the mapping λ is surjective. Finally, suppose that $\sum (p_i, q_i) = 0$. Since Q_R is unitary, there is an $e \in R$ such that $q_i e = q_i$ for all i , and by the surjectivity of λ , e can be written as $\sum (p'_j, q'_j)$. Now we have $\sum p_i \otimes q_i = \sum p_i \otimes q_i (\sum p'_j, q'_j) = \sum_{i,j} p_i \otimes q_i (p'_j, q'_j)$. At this point, notice that for any $p' \in P$ and $q, q' \in Q$, $q(p', q') \in Q \cong \operatorname{Hom}(P, R)$ and $\langle q, p' \rangle q' \in Q \cong \operatorname{Hom}(P, R)$, and for all $p \in P$ it holds $p(q(p', q')) = (p, q)(p', q') = ((p, q)p', q') = (p \langle q, p' \rangle, q') = (p, \langle q, p' \rangle q')$, hence $q(p', q') = \langle q, p' \rangle q'$. Therefore we can continue:

$$\begin{aligned} \sum p_i \otimes q_i &= \sum_{i,j} p_i \otimes \langle q_i, p'_j \rangle q'_j = \sum_{i,j} p_i \langle q_i, p'_j \rangle \otimes q'_j = \sum_{i,j} (p_i, q_i) p'_j \otimes q'_j \\ &= \sum_j (\sum_i (p_i, q_i) p'_j) \otimes q'_j = 0, \end{aligned}$$

which proves that λ is injective.

Finally we consider $\rho: Q \otimes_R P \rightarrow S: \sum q_i \otimes p_i \mapsto \sum \langle q_i, p_i \rangle$. It is clear that ρ is a homomorphism of abelian groups. The fact that ρ is a ring homomorphism and the injectivity of ρ are proven in the same way as was done for λ above. To prove the surjectivity, consider first an idempotent $f \in S$. Since Pf is a finitely generated projective left R -module, there are a $P' \in R \operatorname{Mod}$, an idempotent $e \in R$, and a natural number n such that $Pf \oplus P' \cong (Re)^n$. Denote by p_1, \dots, p_n the canonical image of the basis $(e)_1, \dots, (e)_n$ of $(Re)^n$. Then every element $p \in Pf$ admits a unique decomposition $p = (r_1 e)p_1 + \dots + (r_n e)p_n$. Denote by q_i ($i=1, \dots, n$) the mapping which assigns to each $p \in Pf$ the corresponding element $r_i e$. Clearly, this q_i is a homomorphism from Pf to $Re \subseteq R$. We extend q_i to the whole of P by putting $(P(1-f))q_i = 0$ (here $1-f$ makes sense for f is an endomorphism of ${}_R P$) and denote this extended mapping also by q_i .

By the definition of q_i we have, for every $p \in P$, $(p, q_i) = (pf, q_i)$ and $pf = \sum (pf, q_i) p_i$. Therefore $p \sum \langle q_i, p_i \rangle = \sum (p, q_i) p_i = pf$ for all $p \in P$, hence $f = \sum \langle q_i, p_i \rangle$. Finally, if $s \in S$ and $p \in P$ are arbitrary, then $s = fs$ for an idempotent $f = \sum \langle q_i, p_i \rangle \in S$, and then $ps = pfs = p(\sum \langle q_i, p_i \rangle s) = \sum (p, q_i) (p_i s) = p \sum \langle q_i, p_i s \rangle$, i. e., $s = \sum \langle q_i, p_i s \rangle$, and we are done.

The usual definition of a Morita context makes no use of the identities of the rings, hence it makes sense in our case. Now we have:

THEOREM 2.2. *Let $R, S, {}_R P_S, {}_S Q_R, (,): P \times Q \rightarrow R, \langle, \rangle: Q \times P \rightarrow S$ be a Morita context where R, S are rings with local units and P, Q are unitary bimodules. Then $P \otimes_S -: S \text{ Mod} \rightarrow R \text{ Mod}$ and $Q \otimes_R -: R \text{ Mod} \rightarrow S \text{ Mod}$ are equivalences inverse to each other if and only if both $(,)$ and \langle, \rangle are surjective.*

PROOF. If $P \otimes_S -$ and $Q \otimes_R -$ are inverse equivalences then the surjectivity of $(,)$ and \langle, \rangle follows from 5) in Theorem 2.1. Conversely, if these mappings are surjective then they induce surjective bimodule homomorphisms from ${}_R(P \otimes_S Q)_R$ to R and from ${}_S(Q \otimes_R P)_S$ to S . Next we see that these homomorphisms are also injective. Indeed, let $\sum (p_i, q_i) = 0$. Since Q_R is unitary, there is an $e \in R$ such that $q_i e = q_i$ for all i , and by the surjectivity of $(,)$, e can be written as $\sum (p'_j, q'_j)$. Now we have $\sum p_i \otimes q_i = \sum p_i \otimes q_i (\sum p'_j, q'_j) = \sum_{i,j} p_i \otimes q_i (p'_j, q'_j) = \sum_{i,j} p_i \otimes \langle q_i, p'_j \rangle q'_j = \sum_{i,j} p_i \langle q_i, p'_j \rangle \otimes q'_j = \sum_{i,j} (p_i, q_i) p'_j \otimes q'_j = \sum_j (\sum_i (p_i, q_i)) p'_j \otimes q'_j = 0$. The injectivity of \langle, \rangle is proved dually. Now we obtain, for every $M \in R \text{ Mod}$ and $N \in S \text{ Mod}$, $P \otimes_S (Q \otimes_R M) \cong (P \otimes_S Q) \otimes_R M \cong R \otimes_R M \cong M$ and similarly $Q \otimes_R (P \otimes_S N) \cong N$.

REMARK. In Taylor [9] Morita contexts with surjective mappings are shown to yield Morita equivalence, and vice versa, for central separable algebras over a commutative ring with identity. However, central separable algebras need not have local units and the converse implication does not hold either.

COROLLARY 2.3. *For any rings R, S with local units, $R \text{ Mod}$ and $S \text{ Mod}$ are equivalent if and only if $\text{Mod } R$ and $\text{Mod } S$ are equivalent.*

Next we proceed to characterize Morita equivalence in a way similar to the case of rings with identity. Conform to that terminology, call a unitary bimodule ${}_R M_S$ *balanced* if the canonical homomorphisms $S \rightarrow \text{End}_R M$ and $R \rightarrow \text{End}_S M$ are injective and, identifying R and S with the corresponding subrings of endomorphisms of M , it holds $S \text{End}_R M = S$ and $(\text{End}_S M)R = R$.

THEOREM 2.4. *Let R, S be rings with local units and $G: R \text{ Mod} \rightarrow S \text{ Mod}$, $H: S \text{ Mod} \rightarrow R \text{ Mod}$ be additive functors. Then G and H are equivalences inverse*

to each other if and only if there exists a unitary bimodule ${}_R P_S$ such that

- 1) both ${}_R P$, P_S are locally projective generators,
- 2) ${}_R P_S$ is balanced,
- 3) $G \cong S \text{Hom}_R(P, -)$ and $H \cong P \otimes_S -$.

Moreover, if P satisfies these conditions then, putting $Q = S \text{Hom}_R(P, R)$, ${}_S Q_R$ is a balanced bimodule, both ${}_S Q$ and Q_R are locally projective generators, $H \cong R \text{Hom}_S(Q, -)$ and $G \cong Q \otimes_R -$.

PROOF. The necessity of the conditions as well as the final assertion follow from Theorem 2.1. To prove the sufficiency, let $M \in R \text{Mod}$ be arbitrary. Then we have

$$\begin{aligned} HG(M) &\cong P \otimes_S S \text{Hom}_R(P, M) \stackrel{\text{Cor. 1.8}}{\cong} R \text{Hom}_R(R \text{Hom}_S(P, P)R, M) \stackrel{\text{Prop. 1.3}}{\cong} \\ &\cong R \text{Hom}_R(R, M) \stackrel{\text{Prop. 1.1}}{\cong} M. \end{aligned}$$

On the other hand, for any $N \in S \text{Mod}$,

$$GH(N) \cong S \text{Hom}_R(P, P \otimes_S N) \stackrel{\text{Cor. 1.6}}{\cong} S \text{Hom}_R(P, P) S \otimes_S N \cong S \otimes_S N \stackrel{\text{Prop. 1.2}}{\cong} N.$$

Following Abrams [1], now we present a concrete way to construct rings with local units Morita equivalent to a given ring R of this kind.

For a locally projective module P , the endomorphisms of each P_i extend to endomorphisms of P when composed by ϕ_i , and in this way the endomorphism rings of the components P_i form a direct system of subrings of $\text{End}_R P$. Their limit $S = \varinjlim \text{End}_R P_i$ consists exactly of those endomorphisms of P which factor through one of the projections ϕ_i . The ring S has local units because if the endomorphism $s \in S$ factors through ϕ_i then, choosing a P_j which contains P_i and the image of s (notice that the latter is finitely generated hence such a P_j exists), the projection ϕ_j is a unit to s . Now it is clear that $P \in \text{Mod } S$ and $S \text{End}_R P = S$. If, in addition, ${}_R P$ is a generator then by Lemma 1.10 we obtain that P_S is also a locally projective generator and ${}_R P_S$ is balanced. Then Theorem 2.4 says that the functors $S \text{Hom}_R(P, -)$ and $P \otimes_S -$ are inverse equivalences between $R \text{Mod}$ and $S \text{Mod}$. Furthermore, it is also clear from the above that, for any $M \in R \text{Mod}$, $S \text{Hom}_R(P, M)$ consists exactly of those R -homomorphisms from P to M which factor through one of the ϕ_i . Therefore $S \text{Hom}_R(P, M)$ is, as an abelian group, just the direct limit of the $\text{Hom}_R(P_i, M)$. Thus we have:

THEOREM 2.5 (cf. Abrams [1]). *Two rings R, S with local units are Morita equivalent if and only if there exists a locally projective generator ${}_R P$ such that, using the notation above, $S \cong \varinjlim \text{End}_R P_i$.*

REMARK. Let R be an arbitrary ring with local units, and consider the module ${}_R P = \bigoplus_{e^2=e \in R} Re$. (Notice that if e, f are idempotents with $Re = Rf$ then this left ideal appears (at least) twice in the decomposition of P .) Clearly, ${}_R P$ is a locally projective module. By Theorem 2.5, the ring $S = \varinjlim \text{End}_R Re$ is Morita equivalent to R . To every idempotent $e \in R$ we can assign the endomorphism θ_e of P defined to act identically on the direct component Re and as a zero on all other components. Clearly, the θ_e are orthogonal idempotents in S , and by the definition of S we have

$$S = \bigoplus_{e^2=e \in R} \theta_e S = \bigoplus_{e^2=e \in R} S \theta_e.$$

This shows that S is a ring with enough idempotents in the sense of Fuller [7]. Thus every Morita equivalence class of rings with local units contains rings with enough idempotents (which are even more special than the rings considered in Abrams [1]). A theory of Morita duality for rings with enough idempotents is presented in Yamagata [10].

3. Examples and applications

Of course, all the examples given in Abrams [1] are examples for our theory, too; we are not going to list them again.

EXAMPLE 1. Every regular ring is a ring with local units (but not necessarily in the sense of Abrams [1]). Indeed, let a_1, \dots, a_n be arbitrary elements of a regular ring R . Then there is a $g = g^2 \in R$ such that $a_i g = a_i$, $i = 1, \dots, n$, further there is an $f = f^2 \in R$ such that $f a_i = a_i$, $i = 1, \dots, n$, and $f g = g$. Putting $e = g + f - g f$, it is straightforward to check that $e^2 = e$ and $a_i e = a_i = e a_i$, $i = 1, \dots, n$.

PROPOSITION 3.1. *If R and S are Morita equivalent rings with local units and R is regular then S is also regular.*

PROOF. By Theorem 2.5, S is a direct limit of endomorphism rings of finitely generated projective R -modules. Since R is regular, all these rings are regular, too, and the same holds for S , being the union of these endomorphism rings.

EXAMPLE 2. Let R be a ring with identity, S be a Rees matrix ring over R with canonical decomposition $S \cong Se \underset{eSe}{\otimes} eS$, $e^2 = e \in S$, $eSe \cong R$ (for the definitions of the notions occurring in this example, see Anh-Márki [4]). If S is finitely orthogonal with respect to e , then S is obviously a ring with local units. Now Se is a finitely generated projective left S -module. For any $M \in S \text{ Mod}$ and $m \in M$, consider the mapping $\rho_m : Se \rightarrow M : se \rightarrow sem$. These ρ_m 's together define a homomorphism from $(Se)^{(M)}$ to M whose image is $SeM = Se(SM) = (SeS)M = SM = M$. This proves that Se is a generator for $S \text{ Mod}$. By Theorem 2.5, S is then Morita equivalent to $\text{End}_S(Se) \cong eSe \cong R$.

In what follows, a ring S as in Example 2 will be called a *finitely orthogonal Rees matrix ring*.

Next, observe that § 21 in Anderson-Fuller [2] makes no use of the identity in the given rings, all the results (and proofs) presented there are valid for our module categories, too. Thus we have:

PROPOSITION 3.2 (cf. [2], Corollary 21.9). *Let R and S be equivalent rings with local units. Then R is primitive or a ring with zero Jacobson radical if and only if S is such.*

PROPOSITION 3.3 (cf. [2], Proposition 21.11). *Equivalent rings with local units have isomorphic lattices of ideals; in particular, one of them is simple if and only if so is the other.*

We can also prove the following.

PROPOSITION 3.4. *Let R and S be equivalent rings with local units. If both R and S are commutative then they are isomorphic.*

PROOF. Consider the unitary bimodule ${}_R P_S$ given in Theorem 2.1. Since R is commutative, for any idempotent $e \in R$, R is the direct sum of the rings eR and $(1-e)R$, and we have $P \cong {}_R eP \oplus {}_R (1-e)P = {}_R eP \oplus {}_R (1-e)P$. Now S , being a ring of certain endomorphisms of ${}_R P$, also decomposes into a direct sum $S_1 \oplus S_2$, and again by Theorem 2.1, S_1 and S_2 are equivalent to eR and $(1-e)R$, respectively. By the construction of P , eP is finitely generated as an S -module, hence also as an S_1 -module. Then $eR \cong \text{End } eP_{S_1}$, and since S_1 is commutative, we obtain an embedding of S_1 into eR , but then eP is finitely generated as an eR -module, hence also as an R -module. Herefrom we conclude that rP is a finitely generated R -module for every $r \in R$, hence by the commutativity of R , r can be considered as an element of S , and similarly, every element of S can be considered as an element of R , whence the assertion

follows.

Contrary to the case of unital rings, it is not true that if R and S are equivalent rings with local units then their centres must be isomorphic. In fact, if R is any ring with local units, N is a countably infinite set and R_N^f denotes the ring of $N \times N$ matrices over R with finitely many non-zero entries, the R_N^f is a finitely orthogonal Rees matrix ring over R (see [4]), hence it is Morita equivalent to R by Example 2, and R_N^f is centreless. (We thank Dr. G. Abrams for calling our attention to this simple example.)

Now we characterize rings which are Morita equivalent to rings of certain 'nice' kinds. The first result is essentially Corollary 4.3 in Abrams [1].

PROPOSITION 3.5. *A ring R with local units is Morita equivalent to a ring with identity if and only if there exists an idempotent $e \in R$ with $R = ReR$. If this is the case then R is Morita equivalent to eRe .*

The proof is the same as that of Corollary 4.3 in [1], therefore it is omitted here.

PROPOSITION 3.6. *A ring with local units is Morita equivalent to a division ring if and only if it is a simple ring with minimal one-sided ideals.*

PROOF. Let R be a ring with local units which is Morita equivalent to a division ring D . By Proposition 3.5 there is an idempotent $e \in R$ such that eRe and R are Morita equivalent. Given any finite subset X of R , $X \cup \{e\}$ has a local unit f ; then $X \subseteq fRf$, $R \supseteq RfR \supseteq RefR = ReR = R$, so $R = RfR$, and by Proposition 3.5 fRf is Morita equivalent to R , hence also to D . Now fRf , being a ring with identity Morita equivalent to the division ring D , must be isomorphic to a full matrix ring over D . Theorem 1 in Anh [3] tells us now that R is a simple ring with minimal one-sided ideals.

Conversely, if R is a simple ring with minimal one-sided ideals then it is regular, hence a ring with local units. On the other hand, for any primitive idempotent $e \in R$, eRe is a division ring and $ReR = R$. By Proposition 3.5, R is then Morita equivalent to eRe .

REMARK. Notice that, by a result of E. Hotzel (see Corollary 3.5 in [4]), simple rings with minimal one-sided ideals are just the finitely orthogonal Rees matrix rings over division rings.

By a *primary ring* A we mean a ring with identity whose factor $A/J(A)$ by its Jacobson radical is a simple artinian ring such that idempotents can be lifted. If, moreover, $A/J(A)$ is a division ring then A is said to be a *local*

ring. A ring R is said to be a *strongly locally matrix ring* over a (unital) ring S if for every finite subset $U \subseteq R$ there is an idempotent $e \in R$ such that $U \subseteq eRe$ and eRe is isomorphic to the matrix ring S_n for some n .

PROPOSITION 3.7. *A ring R with local units is Morita equivalent to a primary ring if and only if R is isomorphic to a strongly locally matrix ring over a local ring. If, in addition, ${}_R R$ and R_R are projective modules then R is isomorphic to a finitely orthogonal Rees matrix ring over a local ring.*

PROOF. Let R be Morita equivalent to a primary ring. By Proposition 3.5 there is an idempotent $e \in R$ such that eRe and R are Morita equivalent, hence eRe is Morita equivalent to a primary ring S . Now both eRe and S are rings with identity, hence eRe isomorphic to $fS_n f$ for an idempotent f in a full matrix ring S_n over S . Here $J(fS_n f) = fJ(S_n)f$, so $fS_n f / J(fS_n f) = fS_n f / fJ(S_n)f \cong \bar{f}(S_n / J(S_n))\bar{f}$ where \bar{f} denotes the image of f under the canonical homomorphism of S_n corresponding to $J(S_n)$, the last ring is obviously simple and artinian, and it is also clear that the idempotents can be lifted. Therefore eRe is itself a primary ring, hence there is an idempotent $g \in eRe$ such that gRg is a local ring. Now we have $(eRe)g(eRe) = eRe$ and, by $R = ReR$, also $R = ReR = ReReR = ReRegeReR = ReRgReR = RgR$. Hence, by Proposition 3.5, the bimodule ${}_R Rg {}_{gRg}$ induces Morita equivalence between R and gRg . Furthermore, similarly to the case treated in Example 2, $Rg \in R \text{ Mod}$ is a finitely generated projective generator. Then by Lemma 1.10, $Rg {}_{gRg}$ is a locally projective generator. The canonical components of $Rg {}_{gRg}$ are free modules, being finitely generated projectives over a local ring. Therefore the endomorphism ring of each of them is a finite matrix ring over gRg , and the assertion follows from Theorem 2.5.— The converse is obvious by Proposition 3.5.

Suppose now that, in addition, ${}_R R$ and R_R are projective. Then in the proof above, $Rg {}_{gRg}$ and ${}_{gRg} gR$ are also projective, hence they are free modules, for gRg is a local ring. Now [4], Theorem 3.1 says that $R = RgR$ is a Rees matrix ring over gRg , and it is finitely orthogonal for R has local units.

PROPOSITION 3.8. *A ring S with local units is Morita equivalent to a two-sided perfect local ring R if and only if S is an orthogonal Rees matrix ring over R .*

PROOF. Suppose that R and S are Morita equivalent. By Theorem 2.1, there exist locally projective bimodules ${}_R P_S$ and ${}_S Q_R$ such that $S \cong Q \otimes_R P$. Since R is perfect, ${}_R P$ and Q_R are projective, and since R is local, projectives are

free. Now the assertion follows from [4], Theorem 3.1. The converse is obvious by Example 2.

References

- [1] G.D. Abrams, Morita equivalence for rings with local units, *Commun. Algebra* 11 (1983), 801-837.
- [2] F.W. Anderson and K.R. Fuller, *Rings and Categories of Modules*, Graduate Texts in Mathematics, 13, Springer, Berlin, 1974.
- [3] P.N. Ánh, On Litoff's theorem, *Studia Sci. Math. Hungar.* 16 (1981), 255-259.
- [4] P.N. Ánh and L. Márki, Rees matrix rings, *J. Algebra* 81 (1983), 340-369.
- [5] H. Bass, *The Morita theorems*, Lecture Notes, University of Oregon, 1962.
- [6] H. Bass, *Algebraic K-theory*, W. A. Benjamin, New York, 1968.
- [7] K.R. Fuller, On rings whose left modules are direct sums of finitely generated modules, *Proc. Amer. Math. Soc.* 54 (1976), 39-44.
- [8] M. Sato, On equivalences between module subcategories, *J. Algebra* 59 (1979), 412-420.
- [9] J.L. Taylor, A bigger Brauer group, *Pacific J. Math.* 103 (1982), 163-203.
- [10] K. Yamagata, On Morita duality for additive group valued functors, *Commun. Algebra* 7 (1979), 367-392.

Mathematical Institute
Hungarian Academy of Sciences
H-1364 Budapest, Pf. 127
Hungary