



Morphological and molecular studies of *Neosynechococcus sphagnicola*, gen. et sp. nov. (Cyanobacteria, Synechococcales)

PETR DVOŘÁK¹*, FRANTIŠEK HINDÁK², PETR HAŠLER¹, ALICA HINDÁKOVÁ² & ALOISIE POULÍČKOVÁ¹

¹ Department of Botany, Faculty of Sciences, Palacký University, Šlechtitelů 11, CZ-78371 Olomouc, Czech Republic,

² Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 9, SK-84523 Bratislava, Slovakia

* Corresponding author (email: p.dvorak@upol.cz)

Abstract

The genus *Synechococcus* represents an enigmatic group of cyanobacteria with very simple unicellular morphology and polyphyletic evolutionary origin. Here, we describe a new genus based on strain of *Synechococcus*-like cyanobacterium. The strain was isolated from the peat bog Klin (Slovakia), where it occupies different niches such as hyaline cells of *Sphagnum*, sheaths of cyanobacteria, dead cells of desmids, carapaces of dead crustaceans, and solitary in detritus. We describe this new genus using a combination of molecular, morphological and ecological features. A phylogeny of the 16S rRNA gene, 16S-23S ITS and *rbcL* loci showed a separate position of the investigated strain and its close proximity to filamentous cyanobacteria. Therefore, it is a novel lineage of *Synechococcus*-like cyanobacteria illustrating the polyphyletic nature of the genus *Synechococcus*. Moreover, the strain exhibits unique morphological and ecological features, which allow us to erect the new monospecific genus *Neosynechococcus*.

Key words: new genus, new species, peat bog, Slovakia, 16S rRNA, 16S-23S ITS, *rbcL*

Introduction

The genus *Synechococcus* Nägeli (1849: 56) is a widely distributed group of cyanobacteria exhibiting very simple morphology, which is characterized by unicellular or pseudofilamentous thalli, asymmetric binary fission, and small elongated to cylindrical or rod-like cells (Komárek & Anagnostidis 1998). Its distribution has been extensively investigated in marine environments where, together with *Prochlorococcus* Chisholm *et al.* (1992: 299), it represents the most abundant autotrophs (e.g. Schmidt *et al.* 1991, Li 1994). However, there is important species diversity in other environments, e.g. freshwater planktic, benthic or soil biotopes (see Komárek & Anagnostidis (1998) and John *et al.* (2011) for review). Other taxa include endogloecic species inhabiting mucilage of other cyanobacteria or epiphytic taxa colonizing the surface of colonies. Hindák (1996) described *S. epigloecicus* Hindák (1996: 77) and *S. endogloecicus* Hindák (1996: 78) inhabiting a surface and an interior of mucilage of the cyanobacterium *Microcystis aeruginosa* Kützinger (1846: 6) and *Woronichinia naegeliana* (Unger 1854: 195) Elenkin (1933: 30). Another endogloecic species, *S. mucicolus* Joosten (2006: 76), occurs in the mucilage of *Microcystis* spp. and was described by Joosten (2006).

Besides endogloecic species, cyanobacteria often live in association with other organisms including fungi (lichens), corals or ferns. Moreover, cyanobacteria may be partners in symbiosis with mosses (see Solheim & Zielke (2002) and Adams & Duggan (2008) for a review), because of their ability to fix atmospheric nitrogen. Nitrogenase activity, which is indicative of fixation of atmospheric nitrogen, has been documented in the *Sphagnum* Linnaeus (1753: 1106) mosses (Granhall & Hofsten 1976), where diazotrophic cyanobacteria often inhabit *Sphagnum* hyaline cells. This consortium provides nitrogen to the *Sphagnum* while the cyanobacteria benefit from a stable and more favourable environment than acidified peat bog (Solheim & Zielke 2002).

Evolutionary relationships within the cyanobacteria are the most frequently reconstructed using 16S rRNA gene phylogeny. While it is an effective molecular marker for resolution at the genus level (e.g. Komárek 2010), better taxonomical resolution (lower than genus) is achieved using additional molecular markers. For instance, 16S-

Acknowledgements

The research was supported by IGA UP Prf-2014001 and Post-UP II CZ.1.07/2.3.00/30.0041 (Dvořák, Hašler, Pouličková), and by VEGA projects No 2/0073/13 and No 2/0113/11 (Hindák, Hindáková). We would like to express our gratitude to the referees for helpful comments. Moreover, we are grateful to Dale A. Casamatta from University of North Florida, who edited English throughout the text.

References

- Anagnostidis, K. (2001) Nomenclatural changes in cyanoprokaryotic order Oscillatoriales. *Preslia* 73: 359–375.
- Adams, D.G. & Duggan, P.S. (2008) Cyanobacteria-bryophytes symbioses. *Journal of Experimental Botany* 59: 1047–1058.
<http://dx.doi.org/10.1093/jxb/ern005>
- Bornet, E. & Flahault, C. (1888) Revision des Nostocacées Hétérocystées. *Annales des Sciences Naturelles; Botanique, Série 7*: 177–262.
- Boyer, S.L., Fletchner, V. & Johansen, J.R. (2001) Is the 16S-23S rRNA internal transcribed spacer (ITS) region a good tool for use in molecular systematics and population genetics? A case study in cyanobacteria. *Molecular Biology and Evolution* 18: 1057–1069.
<http://dx.doi.org/10.1093/oxfordjournals.molbev.a003877>
- Boyer, S.L., Johansen, J.R. & Howard, G.L. (2002) Phylogeny and genetic variance in terrestrial *Microcoleus* (Cyanophyceae) species based on sequence analysis of the 16S rRNA gene and associated 16S-23S ITS region. *Journal of Phycology* 38: 1222–1225.
<http://dx.doi.org/10.1046/j.1529-8817.2002.01168.x>
- Casamatta, D.A., Johansen J.R., Vis, M.L. & Broadwater S.T. (2005) Molecular and morphological characterization of ten polar and near-polar strains within the Oscillatoriales (Cyanobacteria). *Journal of Phycology* 41: 421–438.
<http://dx.doi.org/10.1111/j.1529-8817.2005.04062.x>
- Castresana, J. (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17: 540–552. <http://dx.doi.org/10.1093/oxfordjournals.molbev.a026334>
- Chisholm, S.W., Frankel, S.L., Goericke, R., Olson, R.J., Palenik, B., Waterbury, J.B., West-Johnsrud, L. & Zettler, E.R. (1992) *Prochlorococcus marinus* nov. gen. nov. sp.: an oxyphototrophic marine prokaryote containing divinyl chlorophyll a and b. *Archives of Microbiology* 157: 297–300.
<http://dx.doi.org/10.1007/bf00245165>
- Dvořák, P., Hašler, P. & Pouličková, A. (2012) Phylogeography of the *Microcoleus vaginatus* (cyanobacteria) from three continents – a spatial and temporal characterization. *PLoS ONE* 7(6): e40153.
<http://dx.doi.org/10.1371/journal.pone.0040153>
- Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792–1797.
<http://dx.doi.org/10.1093/nar/gkh340>
- Elenkin, A.A. (1933) O novych semejstvach sinezelenych vodoroslej grupy Stereometrae Elenk. (podgrupy Excavatae Elenk.) porjadka Chrococcales Geitler (1925). *Trudy Botanicheskogo Instituta Akademii Nauk SSSR Serie 2* (1–2): 23–34.
- Fox, G.E., Wisotzkey, J.D. & Jurtshuk Jr., P. (1992) How close is close: 16S rRNA sequence identity may not be sufficient to guarantee species identity. *International Journal of Systematic Bacteriology* 42: 166–170.
<http://dx.doi.org/10.1099/00207713-42-1-166>
- Fredslund, J., Schauser, L., Madsen, L.H., Sandal, N. & Stougaard, J. (2005) PriFi: using a multiple alignment of related sequences to find primers for amplification of homologs. *Nucleic Acids Research* 33.
<http://dx.doi.org/doi:10.1093/nar/gki425>
- Gomont, M. (1899). Sur quelques Oscillariées nouvelles. *Bulletin de la Société Botanique de France* 46: 25–41.
- Granhall, U. & v. Hofsten, A. (1976) Nitrogenase activity in relation to intracellular organisms in *Sphagnum* mosses. *Physiologia Plantarum* 36: 88–94.
<http://dx.doi.org/10.1111/j.1399-3054.1976.tb05033.x>
- Hansgirg, A. (1892) Prodrum der Algenflora von Böhmen. Zweiter Theil welcher die blaugrünen Algae (Myxophyceen, Cyanophyceen), nebst Nachträgen zum ersten Theile und einer systematischen Bearbeitung der in Böhmen verbreiten saprophytischen Bacterien und Euglenen enthält. Mit den opiz-preise Gekrönte Arbeit auf Kosten des Opizfondes. *Archiv für die naturwissenschaftliche Landesdurchforschung von Böhmen* 8(4): 1–268.
- Hašler, P., Dvořák, P., Johansen, J.R., Kitner, M., Ondřej, V. & Pouličková, A. (2012) Morphological and molecular study of epipellic filamentous genera *Phormidium*, *Microcoleus* and *Geitlerinema* (Oscillatoriales, Cyanophyta/Cyanobacteria). *Fottea* 12: 341–358.

- Haverkamp, T.H., Schouten, D., Doeleman, M., Wollenzien, U., Huisman, J. & Stal, L.J. (2009) Colorful microdiversity of *Synechococcus* strains (picocyanobacteria) isolated from Baltic Sea. *The ISME Journal* 3: 397–408.
<http://dx.doi.org/10.1038/ismej.2008.118>
- Hindák, F. (1996) Cyanophytes colonizing mucilage of chroococcal water blooms. *Nova Hedwigia* 112: 69–82.
- Hindák, F. & Hindáková, A. (2012) *Chalarodora azurea* Pascher 1929 – a rare glaucophyte found in the peat–bog Klin (Orava, northern Slovakia). In: Wołowski, K., Kaczmarska, I., Ehrman, J.M. & Wojtal, A.Z. (eds.) *Current advances in algal taxonomy and its applications: phylogenetic, ecological and applied perspective*. Institute of Botany, Polish Academy of Sciences, Kraków, pp. 53–60.
- Hoffmann, L., Komárek, J. & Kaštovský, J. (2005) System of cyanoprokaryotes (cyanobacteria) – state in 2004. *Algological Studies* 117: 95–115.
<http://dx.doi.org/10.1127/1864-1318/2005/0117-0095>
- Honda, D., Yokota, A. & Sugiyama, J. (1999) Detection of seven major evolutionary lineages in cyanobacteria based on the 16S rRNA gene sequence analysis with new sequences of five marine *Synechococcus* strains. *Journal of Molecular Evolution* 48: 723–739
<http://dx.doi.org/10.1007/PL00006517>
- John, D.M., Whitton, B.A., Brook A.J. (2011) *The freshwater algal flora of the British Isles*. University Press, Cambridge, 878 pp.
- Joosten, A.M.T. (2006) *Flora of the blue-green algae of the Netherlands. I. the non-filamentous species of inland waters*. KNNV Publishing, Utrecht, 239 pp.
- Katoh, H., Itoh, S., Shen, J.R. & Ikeuchi, M. (2001) Functional analysis of psbV and a novel c-type cytochrome gene psbV2 of the thermophilic cyanobacterium *Thermosynechococcus elongatus* strain BP-1. *Plant Cell Physiology* 42: 599–607.
<http://dx.doi.org/10.1093/pcp/pce074>
- Kimura, M. (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16: 111–120. <http://dx.doi.org/10.1007/BF01731581>
- Komárek, J. (2010) Recent changes (2008) in cyanobacteria taxonomy based on a combination of molecular background with phenotype and ecological consequences (genus and species concept). *Hydrobiologia* 639: 245–259.
<http://dx.doi.org/10.1007/s10750-009-0031-3>
- Komárek, J. & Anagnostidis, K. (1986) Modern approach to the classification system of cyanophytes 2 – Chroococcales. *Algological Studies* 43: 157–226.
- Komárek, J. & Anagnostidis, K. (1988) Modern approach to the classification system of cyanophytes 3 – Oscillatoriales. *Algological Studies* 50–53: 327–472.
- Komárek, J. & Anagnostidis, K. (1998) Cyanoprokaryota 1. Teil: Chroococcales. In: Ettl, H., Gärtner, G., Heynig, H. & Mollenhauer, D. (eds.): *Süßwasserflora von Mitteleuropa 19/1*. Gustav Fischer, Jena-Stuttgart-Lübeck-Ulm, 548 pp.
- Kützing, F.T. (1846) *Tabulae phycologicae; oder, Abbildungen der Tange. Vol. I*. Nordhausen, 160 pp.
<http://dx.doi.org/10.5962/bhl.title.4971>
- Latysheva, N., Junker, V.L., Palmer, W.J., Codd, G.A & Baker, D. (2012) The evolution of nitrogen fixation in cyanobacteria. *Bioinformatics*, 28: 603–606.
<http://dx.doi.org/10.1093/bioinformatics/bts008>
- Li, W.K.W. (1994) Primary production of prochlorophytes, cyanobacteria, and eucaryotic ultraphytoplankton: measurements from flow cytometric sorting. *Limnology and Oceanography* 39: 169–175.
<http://dx.doi.org/10.4319/lo.1994.39.1.0169>
- Linnaeus, C. (1753) *Species plantarum, ed 1*. Holmiae, Stockholm, 1200 pp.
- Miyake, M., Erata, M. & Asada, Y. (1996) A thermophilic cyanobacterium, *Synechococcus* sp. MA19 capable of accumulating poly-β-hydroxybutyrate. *Journal of Fermentation and Bioengineering* 82: 512–514.
[http://dx.doi.org/10.1016/s0922-338x\(97\)86995-4](http://dx.doi.org/10.1016/s0922-338x(97)86995-4)
- Mrva, M. & Hindák, F. (2010) Opätovný nález schránkatej meňavky *Lesquereusia spiralis* (Ehrenberg, 1840) (Testacealobosia: Lesquereusiidae) v Klinskem rašelinisku (Horná Orava, Slovensko). *Limnologický spravodajca* 4: 61–64.
- Nägeli, C. (1849) Gattungen einzelliger Algen. *Neue Denkschriften der Allgemeinen Schweizerischen Gesellschaft für die Gesamten Naturwissenschaften* 10: 1–139.
- Perkerson, R.B., Johansen, J.R., Kováčik, L., Brand, J., Kaštovský, J. & Casamatta, D.A. (2011) A unique pseudanabaenalean (cyanobacteria) genus *Nodosilinea* gen. nov. based on morphological and molecular data. *Journal of Phycology* 47: 1397–1412.
<http://dx.doi.org/10.1111/j.1529-8817.2011.01077.x>
- Robertson, B.R., Tezuka, N. & Watanabe, M. (2001) Phylogenetic analyses of *Synechococcus* strains (cyanobacteria) using sequences of 16S rDNA and part of the phycocyanin operon reveal multiple evolutionary lines and reflect phycobilin content. *International Journal of Systematic and Evolutionary Microbiology* 51: 861–871.
<http://dx.doi.org/10.1099/00207713-51-3-861>
- Schmidt, T.M., DeLong, E.F. & Pace, N.R. (1991) Analysis of a marine picoplankton community by 16S rRNA gene cloning and sequencing. *Journal of Bacteriology* 173: 4371–4378.

- Shirrmeyer, B.E., Antonelli, A. & Bagheri, H.C. (2011) The origin of multicellularity in cyanobacteria. *BMC Evolutionary Biology* 11: 45.
<http://dx.doi.org/10.1186/1471-2148-11-45>
- Siegesmund, M.A., Johansen, J.R., Karsten, U. & Friedl, T. (2008) *Coleofasciculus* gen. nov. (cyanobacteria): morphological and molecular criteria for revision of the genus *Microcoleus* Gomont. *Journal of Phycology* 44: 1572–1585.
<http://doi.wiley.com/10.1111/j.1529-8817.2008.00604.x>
- Solheim, B. & Zielke, M. (2002) Associations between cyanobacteria and mosses. In: Rai, A.N., Bergman, B. & Rasmussen, U. (eds.) *Cyanobacteria in symbiosis*. Kluwer Academic Publishers, Dordrech, pp. 137–152.
- Spurr, A.R. (1969) A low-viscosity resin embedding medium for electron microscopy. *Journal of Ultrastructure Research* 26: 31–43.
[http://dx.doi.org/10.1016/S0022-5320\(69\)90033-1](http://dx.doi.org/10.1016/S0022-5320(69)90033-1)
- Stal, L.J. (1992) Poly(hydroxyalkanoates) in cyanobacteria: an overview. *FEMS Microbiology Letter* 103: 169–180.
<http://dx.doi.org/10.1111/j.1574-6968.1992.tb05835.x>
- Stackerbrandt, E. & Goebel, B.M. (1994) Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *International Journal of Systematic Bacteriology* 44: 846–849.
- Staub, R. (1961) Research on physiology of nutrients of the planktonic cyanobacterium *Oscillatoria rubescens*. *Schweizerische Zeitschrift Für Hydrologie* 23: 83–198.
- Swofford, D.L. (2002) *PAUP. Phylogenetic Analysis of Parsimony (and Other Methods) ver. 4.02*. Sinauer Associates, Sunderland, Massachusetts.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28: 2731–2739.
<http://dx.doi.org/10.1093/molbev/msr121>
- Taton, A., Grubisic, S., Ertz, D., Hodgson, D.A., Piccardi, R., Biondi, N., Tredici, M.R., Mainini, M., Losi, D., Marinelli, F. & Wilmotte, A. (2006) Polyphasic study of Antarctic cyanobacterial strains. *Journal of Phycology* 42: 1257–1270.
<http://dx.doi.org/10.1111/j.1529-8817.2006.00278.x>
- Tomitani, A., Knoll, A.H., Cavanaugh, C.M. & Ohno, T. (2006) The evolutionary diversification of cyanobacteria: molecular-phylogenetic and paleontological perspectives. *Proceedings of the National Academy of Sciences of the United States of America* 103: 5442–5447.
<http://dx.doi.org/10.1073/pnas.0600999103>
- Unger, F. (1854) Beiträge zur Kenntnis der niedersten Algenformen, nebst Versuchen ihre Entstehung betreffend. *Denkschriften der Kaiserlichen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Classe* 7(1): 185–196.
- Zammit, G., Billi, D. & Albertano, P. (2012) The subaerophytic cyanobacterium *Oculatella subterranea* (Oscillatoriales, Cyanophyceae) gen. et sp. nov.: a cytomorphological and molecular description. *European Journal of Phycology* 47: 341–354.
<http://dx.doi.org/10.1080/09670262.2012.717106>