

MOSAIC RING CHROMOSOME 18, RING CHROMOSOME 18 DUPLICATION/DELETION AND DISOMY 18: PERINATAL FINDINGS AND MOLECULAR CYTOGENETIC CHARACTERIZATION BY FLUORESCENCE *IN SITU* HYBRIDIZATION AND ARRAY COMPARATIVE GENOMIC HYBRIDIZATION

Chih-Ping Chen^{1,2,3,4,5,6†*}, Yung-Ting Kuo^{7†}, Shuan-Pei Lin^{2,8}, Yi-Ning Su⁹, Yann-Jang Chen¹⁰, Rui-Yuan Hsueh¹¹, Yi-Hui Lin¹², Pei-Chen Wu¹, Chen-Chi Lee¹, Yu-Ting Chen², Wayseen Wang^{2,13}

*Departments of*¹*Obstetrics and Gynecology,* ²*Medical Research, and* ⁸*Pediatrics, Mackay Memorial Hospital,* ⁵*Institute of Clinical and Community Health Nursing,* ⁶*Department of Obstetrics and Gynecology, School of Medicine, and* ¹⁰*Faculty of Life Sciences and Institute of Genome Sciences, National Yang-Ming University, Departments of* ⁷*Pediatrics and* ¹¹*Obstetrics and Gynecology, Taipei Medical University-Shuang Ho Hospital,* ⁹*Department of Medical Genetics, National Taiwan University Hospital,* ¹²*Department of Obstetrics and Gynecology, Taipei Medical University-Wan Fang Hospital, and* ¹³*Department of Bioengineering, Tatung University, Taipei;* ³*Department of Biotechnology, Asia University, and* ⁴*School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan.*

SUMMARY

Objective: To present the perinatal findings and molecular cytogenetic analysis of a rare chromosomal abnormality involving structural and numerical abnormalities of chromosome 18.

Materials, Methods and Results: A 36-year-old woman, gravida 5, para 3, underwent amniocentesis because of her advanced maternal age. Amniocentesis revealed a karyotype of 46,XY,r(18) [27]/45,XY,-18[5]/46,XY[5]. The parents decided to continue the pregnancy. Level II ultrasound revealed ventriculomegaly. At 38 weeks of gestation, a 3,725 g male fetus was delivered. The fetus had microcephaly, hypertelorism, epicanthal folds, cleft palate, a broad flat nose, simian creases, broad hands, tapered fingers, clubfeet, micropenis, a sacral dimple, hypotonia, ventriculomegaly, and a ventricular septal defect. The peripheral blood lymphocytes revealed a karyotype of 46,XY,r(18)[81]/45,XY,-18[3]/46,XY,idic r(18)[3]/46,XY[13]. Fluorescence *in situ* hybridization using chromosome 18 centromeric probe (cep18) and subtelomeric (18pter, 18qter) identified four types of cells, r(18), idic r(18), monosomy 18, and disomy 18. Array comparative genomic hybridization analysis of the blood demonstrated a 14.9-Mb deletion at chromosome 18p [arr cgh 18p11.32p11.21 (0-14,941,330) × 1] and a 29.6-Mb deletion at chromosome 18q [arr cgh 18q21.2q23 (46,533,430-76,117,153) × 1]. The proband's karyotype was 46,XY,r(18)(p11.21q21.2)[81]/45,XY,-18[3]/46,XY,idic r(18)(p11.21q21.2;p11.21q21.2)[3]/46,XY[13].



ELSEVIER

*Correspondence to: Dr Chih-Ping Chen, Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.
E-mail: cpc_mmh@yahoo.com

Accepted: June 8, 2010

†Chih-Ping Chen and Yung-Ting Kuo contributed equally to this work.

Conclusion: Array comparative genomic hybridization is useful to determine the breakpoints of a ring chromosome, particularly in cases where the ring chromosome comprises the majority of the mosaicism. [*Taiwan J Obstet Gynecol* 2010;49(3):327–332]

Key Words: array comparative genomic hybridization, fluorescence *in situ* hybridization, mosaic ring chromosome, prenatal diagnosis, ring chromosome 18 duplication/deletion

Introduction

The 18q deletion syndrome (OMIM 601808) is caused by variable deletions ranging from 18q21.2, 18q21.3, or 18q22.2 to 18qter [1–3]. The diverse phenotypic features of the 18q deletion syndrome include low birth weight, short stature, microcephaly, midfacial hypoplasia, prognathism, a carp-shaped mouth, a protuberant lower lip, dysplastic ears with prominent antihelix and antitragus, abnormal skull, vertebrae and ribs, atretic ear canals, clubfeet, vertical tali, tapered fingers, dimples over limb joints, hypoplasia of labia or scrotum, micropenis, cryptorchidism, hypospadias, nystagmus, strabismus, glaucoma, tapetoretinal degeneration, bilateral optic atrophy, hypotonia, seizures, deafness, enlarged ventricles, hydrocephalus, porencephaly, holoprosencephaly (HPE), cerebellar hypoplasia, decreased white matter, impaired or delayed myelination, and congenital heart defects [3–12].

The clinical phenotype of the 18p deletion syndrome (OMIM 146390) usually includes growth and mental retardation, hypotonia, epicanthic folds, ptosis, a low nasal bridge, a rounded face, micrognathia, a short neck, abnormal ears, small hands and feet, clinodactyly of the fifth finger, cardiac defects, abnormal genitalia, and cerebral malformations [13–15].

A ring chromosome 18, or r(18) exhibits breakage and reunion at the breakpoints on the long and short arms of chromosome 18, with deletions of the chromosomal segments distal to the breakpoints. The r(18) phenotype is associated with the anomalies of 18p deletion and 18q deletion, and can be associated with the features of both 18p deletion and 18q deletion syndromes [16]. Here, we present the perinatal findings and array comparative genomic hybridization (aCGH) characterization of a rare chromosomal abnormality associated with different cell lines involving structural and numerical abnormalities of chromosome 18.

Materials, Methods and Results

A 36-year-old woman, gravida 5, para 3, underwent amniocentesis at a community obstetric clinic because

of her advanced maternal age. Her husband was 40 years old. She and her husband were both healthy and unrelated, and there was no family history of congenital malformations. The parents had two healthy daughters, one was 12 years old and the other was 9 years old. Amniocentesis at 16 weeks of gestation revealed a male fetus with mosaic ring chromosome 18 [r(18)], monosomy 18 and disomy 18, or 46,XY,r(18)[27]/45,XY,-18[5]/46,XY[5]. The parents decided to continue the pregnancy. Level II ultrasound was unremarkable, except for ventriculomegaly. A 3,725 g male fetus was delivered uneventfully at 38 weeks of gestation. The baby had a head circumference of 33 cm (15th centile) and a body length of 51 cm (60th centile). On examination, the baby manifested microcephaly, hypertelorism, epicanthal folds, cleft palate, a broad flat nose, simian creases, broad hands, tapered fingers, clubfeet, micropenis, a sacral dimple, and hypotonia. Brain ultrasound revealed ventriculomegaly, and echocardiography revealed a ventricular septal defect (VSD). Conventional cytogenetic analysis of peripheral blood lymphocytes revealed a karyotype of 46,XY,r(18)[81]/45,XY,-18[3]/46,XY,idic r(18)[3]/46,XY[13] (Figures 1–4). The deletion of distal 18p and distal 18q on r(18) and idic r(18), and the duplication of the centromere of chromosome 18 were demonstrated by fluorescence *in situ* hybridization (FISH) using an 18p terminal probe (18pter, RP11-324G2)




Figure 1. A karyotype of 46,XY,r(18)(p11.2;q21.2).




Figure 2. A karyotype of 46,XY,idic r(18)(p11.21q21.2;p11.21q21.2).




Figure 3. A karyotype of 45,XY,-18.



Figure 4. A karyotype of 46,XY.

(spectrum green) at 18p11.32, an 18q terminal probe (18qter, RP11-89N1) (spectrum yellow) at 18q23, and an 18 centromeric probe (cep18) (spectrum red) (TelVysion, Downers Grove, IL, USA) (Figure 5). A normal

chromosome 18 contained one cep18 signal (red), one 18pter signal (green), and one 18qter (yellow). The r(18) contained only one cep18 signal (red) and lacked the 18pter signal (green) and the 18qter signal (yellow). The idic r(18) contained two cep18 signals (red) and lacked the 18pter signal (green) and the 18qter signal (yellow). A monosomy 18 cell contained only one normal chromosome 18. A disomy 18 cell contained only two normal chromosomes 18. Oligonucleotide-based aCGH of the blood using Oligo HD Scan (CMDX, Irvine, CA, USA) revealed a 14.9-Mb deletion at chromosome 18p [arr cgh 18p11.32p11.21 (0-14,941,330) × 1] and a 29.6-Mb deletion at chromosome 18q [arr cgh 18q21.2q23 (46,533,430-76,117,153) × 1] (Figure 5). Therefore, the proband's karyotype was 46,XY,r(18)(p11.21q21.2)[81]/45,XY,-18[3]/46,XY,idic r(18)(p11.21q21.2;p11.21q21.2)[3]/46,XY[13].

Discussion

We previously demonstrated the usefulness of aCGH for prenatal detection of microdeletions and unbalanced translocations [17,18]. In this report, we also demonstrate the use of aCGH to determine the breakpoints of a ring chromosome in a case in which the ring chromosome comprises the majority of the mosaicism. The present case had high-level mosaicism for r(18) and low-level mosaicism for disomy 18, r(18) duplication and r(18) deletion. The aCGH findings were consistent with a deletion of 18p11.21→pter and a deletion of 18q21.2→qter in the r(18). aCGH can detect DNA dosage imbalances, including deletions and duplications, but shows limitations for the detection of low-level mosaicism, balanced translocations, inversions and polyploidy. Recent studies have suggested that aCGH can detect as little as 20% mosaicism in peripheral blood cells [19,20]. In this study, aCGH was unable to detect low-level mosaicism.

The present case had a 29.6-Mb 18q deletion, a 14.9-Mb 18p deletion, and phenotypic abnormalities included microcephaly, ventriculomegaly, clubfeet, abnormal external genitalia, ventricular septal defect, cleft palate, hypotonia, and facial dysmorphism. The 18q 21.2→qter deletion in this case encompassed the critical regions for orofacial cleft, microcephaly, and the typical 18q deletion phenotype. Dostal et al [21] suggested that the potential critical region for orofacial cleft is at 18q22.3, between markers D18S879 and D18S1141, and contains the orofacial cleft candidate genes, *SALL3* and *TSHZ1*. Feenstra et al [12] suggested the critical region for the typical phenotype of 18q deletion syndrome is 4.3-Mb region located within 18q22.3-q23,




Figure 5. Fluorescence *in situ* hybridization studies of four types of cells with an 18p terminal probe RP11-324G2 (spectrum green), an 18q terminal probe RP11-89N1 (spectrum yellow) and an 18 centromeric probe (spectrum red). (A) A metaphase cell with monosomy 18 and one chromosome 18 with three signals (green, red and yellow). (B) A metaphase cell with disomy 18 and two chromosomes 18. (C) A metaphase cell with r(18) and one chromosome 18 and one r(18), which has only one red signal. (D) A metaphase cell with idic r(18) and one chromosome 18 and one idic r(18), which has two red signals. r(18)=ring chromosome 18; idic r(18)=isodicentric ring chromosome 18; 18=chromosome 18.

and the critical region for microcephaly is at 18q21.33. The 18p11.21→pter deletion in this case encompassed *HPE4*. However, the present case lacked the HPE phenotype. An HPE critical region on 18p11.3 has been defined as *HPE4* (OMIM 142946) [22] and the responsible gene is *TGIF* (OMIM 602630) [23]. Only ~10% of cases with 18p deletion have HPE [13]. The low incidence of HPE in patients with 18p deletion and *TGIF* haploinsufficiency is because of the autosomal-dominant inheritance pattern of *TGIF* with low penetrance [23], and the requirement for other genetic or environmental factors [24]. Nanni et al [25] suggested that either maternal retinoic acid levels or altered activity in another protein can modify the effects of *TGIF*, and thus lead to phenotypic variability among patients with a *TGIF* deletion.

Prenatal diagnosis of mosaic r(18) with r(18) deletion/duplication is rare [26–29]. Eiben et al [26] reported prenatal diagnosis of 45,XX,-18/46,XX,r(18) (p11q12) by amniocentesis in a fetus with cebocephaly and HPE. The skin fibroblasts had 26.7% (8/30) of

monosomy 18 and 73.3% (22/30) of r(18). Fischer et al [27] reported prenatal diagnosis of 46,XY,r(18)(p11q23)[103]/45,XY,-18[13] by amniocentesis in a fetus with amniotic band syndrome. The cord blood had a karyotype of 46,XY,r(18). Carreira et al [28] reported prenatal diagnosis of mosaic r(18) and r(18) deletion/duplication in two cases, and used multicolor banding to determine the breakpoints. In the first case, a fetus with HPE, 46,XY,r(18)(p11.1q22)[36]/45,XY,-18[7]/47,XY,-18,+r(18)(p11.1q22)×2[1]/46,XY,dupr(18)(p11.1q22)[1] was diagnosed by amniocentesis. Postnatal cytogenetic analysis of fetal tissue revealed a karyotype of 46,XY,r(18)[89]/45,XY,-18[22]/46,XY,dupr(18)[2]/46,XY,mar(18)[4]. In the second case, a fetus with congenital heart defects and obstructive uropathy, 46,XY,r(18)(p11.22q21.2)[12]/46,XY,r(18;18)(p11.22q21.2;p11.22q21.2)[4]/47,XY,-18,+r(18)(p11.22q21.2)×2[1]/del(18)(:p11.21→q11.2:)[1]/ace(18)(:q11.2→q12.2:)[1] was diagnosed by amniocentesis. Mello et al [29] reported prenatal diagnosis of 46,XY,t(3;6)(p26.2;q15),r(18)[22]/45,XY,t(3;6)(p26.2;




Figure 6. Oligonucleotide-based array comparative genomic hybridization shows a 14.9-Mb deletion (0–14,941,330 bp) of the short arm of chromosome 18 and a 29.6-Mb deletion (46,533,430–76,117,153 bp) of the long arm of chromosome 18.

q15),-18[6] by amniocentesis in a fetus with a single umbilical artery, an increased nuchal fold thickness and tricuspid regurgitation. aCGH revealed a 2.8-Mb of 18p11.31p11.32 deletion and a 22.5-Mb 18q21.3 q23 deletion. The fetus was carried to term and had

microcephaly, micropenis, a short stature, congenital heart defects, a high-arched palate, abnormal hands, and facial dysmorphism.

To our knowledge, this is the first report of mosaic r(18), r(18) deletion/duplication and disomy 18 in which

the baby was carried to term. Perinatally, this case represents a difficulty for genetic counseling because of the complexity of the karyotype and the molecular evidence of large deletions of 18p and 18q. The parents elected to continue the pregnancy based on a desire for a son. Although the proband has many features of both 18p and 18q deletion syndromes, he is making great progress and surviving the neonatal period.

Acknowledgments

This work was supported by research grants NSC-96-2314-B-195-008-MY3 and NSC-97-2314-B-195-006-MY3 from the National Science Council, and MMH-E-99004 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- Kline AD, White ME, Wapner R, et al. Molecular analysis of the 18q- syndrome and correlation with phenotype. *Am J Hum Genet* 1993;52:895-906.
- Silverman GA, Schneider SS, Massa HF, et al. The 18q- syndrome: analysis of chromosomes by bivariate flow karyotyping and the PCR reveals a successive set of deletion breakpoints within 18q21.2-q22.2. *Am J Hum Genet* 1995; 56:926-37.
- Strathdee G, Zackai EH, Shapiro R, Kamholz J, Overhauser J. Analysis of clinical variation seen in patients with 18q terminal deletions. *Am J Med Genet* 1995;59:476-83.
- de Grouchy J, Royer P, Salmon C, Lamy M. Délétion partielle du bras long du chromosome 18. *Pathol Biol* 1964;12:579-82.
- Greenberg F. Chromosome 18, monosomy 18q. In: ML Buyse, ed. *Birth Defect Encyclopedia*. Cambridge: Blackwell Scientific Publications, 1990:382-3.
- Chen CP, Chern SR, Liu FF, Jan SW, Lee CC, Chang YC, Yue CT. Prenatal diagnosis of a deletion of 18q in a fetus associated with multiple-marker screen positive results. *Prenat Diagn* 1997;17:571-6.
- Chen CP, Tzen CY, Chang TY, et al. Prenatal diagnosis of *de novo* mosaic distal 18q deletion associated with congenital anomalies. *Ultrasound Obstet Gynecol* 2003;21:202-4.
- Chen CP, Chern SR, Hung FY, et al. Prenatal diagnosis of pure distal 18q deletion. *Prenat Diagn* 2006;26:184-5.
- Chen CP, Lin SP, Chern SR, Lee CC, Huang JK, Wang W. Direct transmission of the 18q- syndrome from mother to daughter. *Genet Counsel* 2006;17:185-9.
- Cody JD, Pierce JF, Brkanac Z, Plaetke R, Ghidoni PD, Kaye CI, Leach RJ. Preferential loss of the paternal alleles in the 18q- syndrome. *Am J Med Genet* 1997;69:280-6.
- Cody JD, Ghidoni PD, Dupont BR, et al. Congenital anomalies and anthropometry of 42 individuals with deletions of chromosome 18q. *Am J Med Genet* 1999;85:455-62.
- Feenstra I, Vissers LELM, Orsel M, et al. Genotype-phenotype mapping of chromosome 18q deletions by high-resolution array CGH: an update of the phenotypic map. *Am J Med Genet* 2007;143A:1858-67.
- Schinzel A. *Catalogue of Unbalanced Chromosome Aberrations in Man*, 2nd edn. Schinzel A, ed. Berlin: Walter de Gruyter, GmbH & Co. 2001;717-22.
- Chen CP, Chern SR, Wang W, et al. Prenatal diagnosis of partial monosomy 18p (18p11.2→pter) and trisomy 21q (21q22.3→qter) with alobar holoprosencephaly and premaxillary agenesis. *Prenat Diagn* 2001;21:346-50.
- Wester U, Bondeson ML, Edeby C, Annerén G. Clinical and molecular characterization of individuals with 18p deletion: a genotype-phenotype correlation. *Am J Med Genet* 2006; 140A:1164-71.
- Stankiewicz P, Brozek I, Hélias-Rodzewicz Z, et al. Clinical and molecular-cytogenetic studies in seven patients with ring chromosome 18. *Am J Med Genet* 2001;101:226-39.
- Chen CP, Su YN, Chang TY, Chern SR, Tsai FJ, Hwang JK, Wang W. 22q11.2 microdeletion in a fetus with double-outlet right ventricle, pulmonary stenosis and a ventricular septal defect: prenatal diagnosis by array comparative genomic hybridization. *Taiwan J Obstet Gynecol* 2009;48:437-40.
- Chen C-P, Su Y-N, Tsai F-J, et al. Terminal 2q deletion and distal 15q duplication: prenatal diagnosis by array comparative genomic hybridization using uncultured amniocytes. *Taiwan J Obstet Gynecol* 2009;48:441-5.
- Ballif BC, Rorem EA, Sundin K, et al. Detection of low-level mosaicism by array CGH in routine diagnostic specimens. *Am J Med Genet* 2006;140A:2757-67.
- Shaffer LG, Kashork CD, Saleki R, Rorem E, Sundin K, Ballif BC, Bejjani BA. Targeted genomic microarray analysis for identification of chromosome abnormalities in 1500 consecutive clinical cases. *J Pediatr* 2006;149:98-102.
- Dostal A, Nemeckova J, Gaillyova R. The 18q deletion syndrome and analysis of the critical region for orofacial cleft at 18q22.3. *J Craniomaxillofac Surg* 2009;37:272-5.
- Overhauser J, Mitchell HF, Zackai EH, Tick DB, Rojas K, Muenke M. Physical mapping of the holoprosencephaly critical region in 18p11.3. *Am J Hum Genet* 1995;57:1080-5.
- Gripp KW, Edwards MC, Mowat D, et al. Mutations in the transcription factor TGIF in holoprosencephaly. *Am J Hum Genet* 1998;63:A32.
- Portnoi MF, Gruchy N, Marlin S, et al. Midline defects in deletion 18p syndrome: clinical and molecular characterization of three patients. *Clin Dysmorphol* 2007;16:247-52.
- Nanni L, Ming JE, Bocian M, et al. The mutational spectrum of the *Sonic Hedgehog* gene in holoprosencephaly: *SHH* mutations cause a significant portion of autosomal dominant holoprosencephaly. *Hum Mol Genet* 1999;8:2479-88.
- Eiben B, Unger M, Stoltenberg G, et al. Prenatal diagnosis of monosomy 18 and ring chromosome 18 mosaicism. *Prenat Diagn* 1992;12:945-50.
- Fischer W, Dermitzel A, Osmers R, Pruggmayer M. Complete karyotype discrepancy between placental and fetal cells in a case of ring chromosome 18. *Prenat Diagn* 2001;21:481-3.
- Carreira IM, Mascarenhas A, Matoso E, et al. Three unusual but cytogenetically similar cases with up to five different cell lines involving structural and numerical abnormalities of chromosome 18. *J Histochem Cytochem* 2007;55:1123-8.
- Mello AL, Crotwell PL, Flanagan JD, et al. Clinical course of a 20-month-old child diagnosed prenatally with mosaic ring chromosome 18 and monosomy 18. *SD Med* 2008;61: 327-9, 331.