

NBER WORKING PAPER SERIES

MULTI-GENERATIONAL IMPACTS OF CHILDHOOD ACCESS TO THE SAFETY NET:
EARLY LIFE EXPOSURE TO MEDICAID AND THE NEXT GENERATION'S HEALTH

Chloe N. East
Sarah Miller
Marianne Page
Laura R. Wherry

Working Paper 23810
<http://www.nber.org/papers/w23810>

NATIONAL BUREAU OF ECONOMIC RESEARCH

1050 Massachusetts Avenue

Cambridge, MA 02138

September 2017, Revised January 2022

First version: September 2017. This version: January 23, 2022. We thank Henrik Kleven, Thomas Lemieux, Liran Einav, Andrew Goodman-Bacon, Michelle Marcus, Adam Schickedanz, Barton Willage, four anonymous referees, and seminar participants at Duke University, Indiana University O'Neill School, Institute for Poverty Research, National Bureau of Economic Research, New York University, Norwegian School of Economics, Purdue University, San Diego State University, Texas A&M, UCLA, UC Davis, University of Connecticut, University of Mannheim, University of Notre Dame, University of Pennsylvania, University of Southern California, Uppsala University, and University of Virginia. We also thank session participants at the All-California Labor Economics, American Economic Association, American Society of Health Economists, Association of Public Policy and Management, Federal Statistical Research Data Center, Population Association of America, and Western Economic Association annual meetings, for their helpful comments. We also thank Pat Barnes, Amy Branum, Clint Carter, and John Sullivan for their assistance in accessing the restricted data used in this project at the California and Michigan Census Research Data Centers. This project was supported by funding from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD, R01 HD093898), National Science Foundation (SMA1327768), the Robert Wood Johnson Foundation (74926-0), and a University of California Davis Interdisciplinary Frontiers in the Humanities and Arts seed grant. Wherry benefited from facilities and resources provided by the California Center for Population Research at UCLA (CCPR), which receives core support (R24-HD041022) from the NICHD. The views expressed herein are those of the authors and do not necessarily reflect the views of the National Bureau of Economic Research.

NBER working papers are circulated for discussion and comment purposes. They have not been peer-reviewed or been subject to the review by the NBER Board of Directors that accompanies official NBER publications.

© 2017 by Chloe N. East, Sarah Miller, Marianne Page, and Laura R. Wherry. All rights reserved. Short sections of text, not to exceed two paragraphs, may be quoted without explicit permission provided that full credit, including © notice, is given to the source.

Multi-generational Impacts of Childhood Access to the Safety Net: Early Life Exposure to Medicaid and the Next Generation's Health

Chloe N. East, Sarah Miller, Marianne Page, and Laura R. Wherry

NBER Working Paper No. 23810

September 2017, Revised January 2022

JEL No. I1,I13,I14,I18

ABSTRACT

We examine multi-generational impacts of positive in utero health interventions using a new research design that exploits sharp increases in prenatal Medicaid eligibility that occurred in some states. Our analyses are based on U.S. Vital Statistics Natality files, which enables linkages between individuals' early life Medicaid exposure and the next generation's health at birth. We find evidence that the health benefits associated with treated generations' early life program exposure extend to later offspring. Our results suggest that the returns on early life health investments may be substantively underestimated.

Chloe N. East
University of Colorado Denver
chloe.east@ucdenver.edu

Sarah Miller
Ross School of Business
University of Michigan
701 Tappan Street
Ann Arbor, MI 48109
and NBER
mille@umich.edu

Marianne Page
Department of Economics
University of California, Davis
Davis, CA 95616-8578
and NBER
mepage@ucdavis.edu

Laura R. Wherry
Wagner Graduate School of Public Service
New York University
295 Lafayette Street
New York, NY 10012
and NBER
laura.wherry@nyu.edu

1 Introduction

There is substantial evidence that health and socioeconomic inequalities persist across generations. A growing number of studies suggest that differences in early life health environments may causally contribute to these disparities. Negative shocks to the *in utero* environment, in particular, have been found to be harmful to individuals' later life health and earnings. A handful of studies also examine positive interventions and find that policies intended to improve early life experiences generate better adult outcomes (Almond and Currie, 2011; Almond et al., 2018). By extension, literatures in economics, epidemiology and child development predict that the causal impacts of these interventions should echo beyond the exposed generation, onto later offspring. Little is known, however, about the extent to which the early life environment affects future generations, or the potential for public policy to alter such linkages.

We consider whether public health interventions experienced *in utero* affect the next generation's health. We focus on the impact of the largest source of health-related services for low-income individuals in the United States: the Medicaid program. Changes in program rules during the 1980s expanded access to low-income pregnant women who were not already connected to Medicaid through the welfare system, and led to a dramatic increase in prenatal coverage. There was considerable variation in the timing and magnitude of these expansions across states, which prior empirical research has harnessed to document the program's beneficial effects on cohorts who gained *in utero* access. We build on this "first generation" research to investigate whether positive policy interventions in one generation transmit to the next generation.

Our study makes three contributions. First, within the active "early origins" literature, the vast majority of causal studies confine their analyses to treated cohorts. While an ever-expanding number of animal experiments provide compelling evidence that the effects of early life environments can be transmitted to later generations,¹ human studies are more rare. We contribute to this literature by using a quasi-experimental design to document multi-generational effects in humans.

Second, we focus on documenting generationally persistent effects of a widespread policy-driven intervention. Most of what we do know about how individuals' early life health experiences affect their later offspring comes from studies of exposure to disasters such as famine and disease outbreaks, which may not compare well to the effects of more common (and malleable) health experiences. This is an important gap – particularly in light of current political debates about the cost of publicly provided health insurance – as substantive multiplier effects to future generations would suggest that existing benefit-cost calculations underestimate the true value of government investments in children's health.

Third, we advance knowledge of the impacts of the 1980s Medicaid expansions by putting forward a new research design that explicitly addresses ongoing debates about policy endogeneity and the validity of using state-level variation in the expansions' timing and magnitude as a natural experiment. Following the pioneering work of Currie and Gruber (1996a,b), most investigations employ an instrumental variables model to isolate policy effects from other potentially correlated changes in state characteristics. However, lack of information on the pre-expansion period (before 1979) has left open the possibility that estimates of the program's benefits are contaminated by unobserved state pre-trends

¹Useful reviews of this literature include Daxinger and Whitelaw (2010, 2012); Heard and Martienssen (2014); Hochberg et al. (2011); Nadeau (2009).

that were correlated with the expansions' timing and magnitude. We have collected four additional years of information on state Medicaid rules, which allows us to measure state eligibility going back to 1975. With these measures, we estimate an event study model that allows flexible estimation of the dynamic effects of changes in Medicaid, and a direct assessment of the extent to which both Currie and Gruber's estimates, and our own estimates of the program's effects on the next generation's health, reflect other underlying factors. To implement this, we classify states as either treated states or control states based on their expansion patterns across the 1975-1988 period, during which time prenatal Medicaid was made available to the lowest income families. Treated states are those that experienced sharp increases in Medicaid eligibility, while control states are those that trended more incrementally over time. We also use restricted data on Medicaid coverage among labor and delivery hospital discharges to validate the event study design, and provide important new evidence on the impacts of the expansions on actual take-up.

Our main analyses are based on information that is available in restricted-use versions of the U.S. Vital Statistics birth records. With these data, we analyze infant health outcomes among two generations. First, we consider those who were directly exposed to the expansions: infants born between 1975 and 1988. We call these cohorts the "first generation." Most previous research has focused on a subset of these cohorts, who were born after 1979. Then, we move on to investigate whether the benefits of *in utero* Medicaid exposure affected the next generation by looking at birth outcomes among the first generation's offspring. We call these infants the "second generation." For the first generation, we use state and year of birth to assign policy generosity, and, for the second generation, we use *mother's* state and year of birth to link to measures of mother's early life Medicaid exposure.

We estimate Medicaid's impacts on standard measures of infant health, including birthweight and low birthweight. These outcomes are strong candidates for a second generation exploration because previous research has shown that birthweight is tied to measures of maternal health such as obesity and diabetes, for which there is already evidence of long-term program effects (Miller and Wherry, 2019). Moreover, birthweight is highly predictive of later life health and economic outcomes, making it an outcome of particular interest. We also investigate other measures of second generation health, including variants of gestational length and weight-for-gestational age. Finally, we explore potential transmission mechanisms. In addition to confirming Medicaid's impacts on the first generation's health at birth as a potential pathway, we consider changes in first generation fertility patterns that might be indicative of selection, and changes in first generation health and health behaviors during pregnancy.

We find that the expansions increased the first generation's likelihood of being enrolled in Medicaid at birth, and we validate the past research finding that the expansions reduced the percent of the first generation who were born low birthweight. Then we go on to show the benefits of expanding early life access for the next generation: the first generation's *in utero* exposure to Medicaid leads to a statistically significant increase in the second generation's birthweight and reduces the incidence of very low birthweight and small for gestational age. We also observe less precisely estimated, but suggestive, declines in low birthweight and very preterm births.

Analyzing the effect of a policy across multiple generations is complicated because of the time that passes between the first generation's treatment and the second generation's outcomes. Our identification assumption requires that among cohorts born in treated states around the time of the expansions

there are no other cohort-state varying economic or health shocks that also affect the health of the next generation. As with all such research designs, we cannot be 100% sure that we have eliminated all possible confounders; nevertheless, our event study approach offers a transparent way to assess the validity of our design and our results. We also show that our results are robust to a variety of specification checks, including alternative state and year control variables, changes in the sample definition, alternative measures of eligibility, omitting the control states entirely, and versions of the model that use a synthetic control approach to construct the comparison group. They are also robust to new methods proposed by [Callaway and Sant’Anna \(2020\)](#) that account for biases that are sometimes found in traditional two-way fixed effects and event study models. In some cases, these alternative specifications provide wider confidence intervals than our baseline results or moderately change the size of the effect. When taken together, however, the pattern of results paints a compelling picture of improvements in second generation infant health.

Documenting the presence of multi-generational spillovers is an important contribution in its own right. Moreover, back-of-the envelope calculations suggest that the magnitude of the spillovers is economically important. Our point estimates of the effects of *in utero* Medicaid access on the second generation’s birthweight suggest medical cost savings in the first year of life that are about 60 percent of the costs of providing the first generation with *in utero* coverage. Even the lower bound of the 95 percent confidence interval suggests cost savings of around 16 percent of initial investment costs. Importantly, this calculation does not include any other benefits that were likely accrued to the second generation, such as later life savings in medical costs or social supports. Nor does it include benefits associated with previously documented improvements in the first generation’s health and economic outcomes. If these benefits were incorporated, the costs savings would be substantially larger.

We find no evidence that changes in overall fertility can explain the effects on infant health, but we do find a small shift in the racial composition of women giving birth: first-generation Medicaid exposure increases the percent of second-generation births that are to white women, and decreases the percent to Black women. Controlling for mothers’ demographic characteristics does not change the baseline results, however, and results are similar when we focus on white births, a sub-group for whom we would not expect to see effects if the main estimates are driven by changes in racial composition. Therefore, we do not believe that selection is driving the results.

Our results indicate that public investments in prenatal health have persistent impacts beyond the treated generation. By quantifying these effects, we establish that benefit/cost ratios based only on outcomes directly experienced by cohorts who were immediately affected by the Medicaid expansions underestimate the program’s overall efficacy. More broadly, our analyses suggest that even “long-run” studies of early-life interventions may fail to capture the full extent of benefits conferred.

The remainder of our paper proceeds in the following way: Section 2 provides further information about the existing literature on “early life” health and multi-generational processes. In Section 3, we describe the Medicaid program and the nature of the 1980s expansions. Sections 4 and 5 describe our empirical strategy and data. We present our results in Section 6 and conclude with a discussion in Section 7.

2 Background

More than 40 years ago, [Forsdahl \(1977\)](#) put forward a provocative hypothesis that the period of gestation has significant impacts on individual health that reach well into adulthood. This theory gained further traction following [Barker \(1990\)](#), and in recent years there has been growing scientific agreement that the time both before and immediately after birth are critical periods when the developing body takes adaptive cues from its surrounding environment (often called the “Barker Hypothesis”). A key feature of the “fetal origins” hypothesis is that the health effects of the *in utero* environment can remain latent for many years. We have yet to achieve a full understanding of the processes underlying these phenomena, but a leading theory is that the fetus’s surrounding environment alters genetic programming through the “switching on” of specific genes.

Numerous economists and epidemiologists have used quasi-experimental designs to test the fetal origins hypothesis, and have found that *in utero* and early life health experiences can have important effects on later life outcomes. Although the vast majority of studies have identified these effects using short-term, extreme events such as disease outbreaks and famines,² a few have recently emerged demonstrating the long term efficacy of more wide-spread positive health interventions.³ Most of these studies focus on historical events, or settings outside the U.S. Three exceptions, [Bailey et al. \(2020\)](#); [Bitler and Figinski \(2019\)](#); [Hoynes et al. \(2016\)](#), investigate the 1960’s roll out of the U.S. Food Stamp program and find that access to the program in early childhood generated improvements in later life economic and health outcomes, including conditions related to cardiovascular disease such as obesity, high blood pressure, and diabetes.⁴

In this vein, recent work finds that changes to Medicaid rules that expanded coverage to low-income pregnant women had long-term benefits on their first generation offspring. Focusing on variation generated by the 1980s prenatal expansions, [Miller and Wherry \(2019\)](#) find that, like the Food Stamp program, *in utero* exposure to Medicaid reduced the likelihood of having metabolic-syndrome and circulatory-system linked chronic illnesses in adulthood. Importantly for our study, when these later life diseases are experienced during pregnancy, women and their children are put at risk for a variety of health problems, including an increased risk of gestational diabetes, complications related to high blood pressure, and preterm birth ([Catalano and Ehrenberg, 2006](#)). Moving beyond health, Miller and Wherry also document that *in utero* exposure to the program is associated with increases in educational attainment.⁵ Similar results are found in a large number of studies evaluating the long-term effects of the 1980s and 1990s expansions to broader age groups ([Brown et al., 2020](#); [Cohodes et al., 2016](#); [Currie et al., 2008](#); [Thompson, 2017](#); [Wherry and Meyer, 2016](#); [Wherry et al., 2017](#)),⁶ although these studies

²Examples from this large literature include [Almond \(2006\)](#); [Almond and Mazumder \(2005\)](#); [Barreca \(2010\)](#); [Mazumder et al. \(2010\)](#); [Neelsen and Stratmann \(2012\)](#) [disease], [Almond et al. \(2010\)](#); [Almond and Mazumder \(2011\)](#); [Chen and Zhou \(2007\)](#); [Painter et al. \(2005\)](#); [Ravelli et al. \(1976\)](#); [Roseboom et al. \(2001\)](#); [Scholte et al. \(2015\)](#); [Stein et al. \(1975\)](#); [Susser \(1992\)](#); [van Ewijk \(2011\)](#) [nutrition], [Almond et al. \(2009\)](#) [pollution]. Quasi-experimental studies of stress ([Persson and Rossin-Slater, 2018](#)) and pollution ([Sanders, 2012](#)) also find detrimental effects.

³E.g. [Bhalotra and Venkataramani \(2015\)](#); [Bharadwaj et al. \(2013\)](#); [Bütikofer et al. \(2019\)](#); [Fitzsimmons and Vera-Hernández \(2013\)](#); [Glied and Neidell \(2010\)](#).

⁴Related literatures examine the long-term effects of education interventions such as Head Start, which includes a health component (e.g. [Carneiro and Ginja, 2014](#); [Deming, 2009](#); [Garces et al., 2002](#); [Ludwig and Miller, 2007](#); [Conti et al., 2016](#)), and policies that reduce pollution exposure ([Isen et al., 2017](#); [Nilsson, 2009](#)).

⁵This finding is consistent with earlier work by [Levine and Schanzenbach \(2009\)](#) linking Medicaid-induced improvements in infant health to higher test scores in 4th and 8th grade.

⁶Three other studies document how the introduction of Medicaid between 1966 and 1970 improved later life health.

do not separate the effect of *in utero* exposure from later childhood exposure. Miller and Wherry find that this distinction is important, with the prenatal exposures generating substantially bigger impacts.

Taken as a whole, the existing literature has generated two broad conclusions. First, early life health shocks have long-term impacts on the health and economic outcomes of those who experience them. Second, many widespread public health interventions targeted at children have substantive positive benefits that last well into adulthood. A natural question is whether these effects endure to the next generation. Economists have previously documented that health and economic status persist across multiple generations (Clark, 2014; Halliday et al., 2021; Solon, 2018), but quasi-experimental investigations are rare. We know little about what drives the correlations, or the potential for policy based treatments to alter them. The dearth of work among social scientists likely results from the multiple challenges of identifying exogenous variation in early life health environments and linking that variation to data that provides relevant information on multiple generations.

These challenges can be overcome in biological studies, where an accumulation of evidence based on animal experiments finds that prenatal health shocks have persistent generational effects. As an example, studies have documented that rats that are malnourished before or during pregnancy produce offspring with smaller brains and reduced cognition, even if the offspring receive sufficient nutrition after birth. Importantly, these effects are not only observed in the immediate offspring, but are present in later generations as well.⁷ Similar multi-generational patterns have been found with *in-utero* exposure to disease, stress, and smoke.⁸ One explanation for this pattern is that the biological predecessors of the ovaries and sperm cells, which produce the next generation, are already present at the fetal stage and are therefore exposed to any insult experienced by the fetus.

In spite of the methodological challenges, a few epidemiologists and economists have been able to shed light on this question by exploiting historical shocks. Painter et al. (2008) investigate the multi-generational impacts of the Dutch Hunger Winter of 1944-1945, which reduced the food consumption of a previously well-nourished population by more than 75 percent. They find that the offspring of those who were exposed to the famine *in utero* had worse health in later life. Similarly, Almond et al. (2010) find that fetal exposure to malnutrition resulting from the 1959-1961 Chinese famine increased low birthweight incidence in the next generation. Looking beyond the effects of extreme nutritional deprivation, Richter and Robling (2016) find that the children of those who were exposed *in utero* to the 1918-1919 influenza pandemic grew up to have lower levels of educational attainment. Similarly, Black et al. (2019) find that Norwegian cohorts exposed to radioactive fallout during the *in utero* period had children with lower cognitive ability.⁹

Using geographic variation in program roll-out to identify the effects of exposure to Medicaid under age 6, Boudreaux et al. (2016) find that Medicaid reduced the probability of having a chronic health condition in adulthood. Using a similar strategy, Sohn (2017) finds that Medicaid's initial roll-out was associated with lower adult mortality. Goodman-Bacon (2021) uses variation in pre-existing welfare eligibility levels (since Medicaid was originally linked to welfare receipt) and finds that the introduction of Medicaid reduced later life mortality and disability, increased employment, and reduced disability benefits for cohorts who were exposed to the program early in life.

⁷A few examples include Aerts and Van Assche (2006); Cowley and Griesel (1966); Dunn and Bale (2009); Jimenez-Chillaron et al. (2009); Martin et al. (2008); Martínez et al. (2014); Masuyama et al. (2015); Radford et al. (2014); Schöpfer et al. (2012); Zamenhof et al. (1971).

⁸Examples include Francis et al. (1999); Grundwald and Brunton (2015); Maritz and Mutemwa (2014); Matthews and Phillips (2012); Morgan and Bale (2011); Rehan et al. (2012).

⁹A few studies also investigate multi-generational effects of health events that occur at later ages. van den Berg and Pinger (2016) investigate multi-generational effects of pre-pubertorial exposure to the German famine of 1916-1918 and

A small number of studies examine intergenerational health effects in the more modern U.S. context. Two studies relate local variation in infant mortality rates at the time of the mother’s birth to her later offspring’s health. Here, the infant mortality rate proxies for broad disease exposure, which can be driven by many factors, including access to medical care. [Almond et al. \(2012\)](#) find that higher exposure to disease is associated with worse long-term outcomes and an increase in the probability that future offspring are below the low birthweight threshold. [Almond and Chay \(2006\)](#) focus on the dramatic improvements in Black infant mortality rates that coincided with the Civil Rights Act, and find that Black women born during the late 1960s had a reduced likelihood of giving birth to a low birthweight infant. A third study, [Colmer and Voorheis \(2020\)](#), documents improved educational outcomes among the grandchildren of cohorts who benefited from reductions in pollution exposure driven by the 1970 Clean Air Act Amendments.¹⁰

We build on this small number of studies by harnessing a policy driven increase in access to a widespread public health program that is a critical component of the U.S. safety net. This allows us to establish multi-generational linkages associated with more common and contemporaneous variation in early life health experiences, while simultaneously quantifying long-term benefits of the Medicaid program that have not been previously considered. Medicaid may alter biological associations across generations by increasing the use of prenatal care, which provides nutrition and drug counseling, immunizations, early diagnoses and direct interventions.¹¹ Along these lines, [Currie and Gruber \(1996b\)](#) link changes in prenatal Medicaid access to reductions in treated cohorts’ probability of being low birthweight, and maternal birthweight is predictive of later offspring’s birthweight (e.g. [Currie and Moretti, 2007](#); [Black et al., 2007](#); [Royer, 2009](#)). Access to Medicaid may also reduce maternal stress: in an analysis of the Oregon Health Insurance Experiment, [Finkelstein et al. \(2012\)](#) find that those who gained health insurance through the experiment experienced substantive improvements in mental health. Several studies have linked parental and *in utero* stress to children’s well-being ([Black et al., 2016](#); [Camacho, 2008](#); [Mansour and Rees, 2012](#); [Persson and Rossin-Slater, 2018](#); [Valente, 2011](#)), with possible ramifications for the next generation’s health.

Besides biological pathways, Medicaid may of course affect the next generation’s health through its documented impacts on the treated generation’s human capital and earnings. As described earlier, numerous studies find that the Medicaid expansions had positive effects on treated cohorts’ educational and economic outcomes in adulthood, and it is well known that children living in high income families are healthier than children living in low income families ([Currie and Almond, 2011](#); [Case et al., 2002, 2005](#)). This is intuitive, as parents with more income have more resources to invest in their children, experience lower stress levels ([Aizer et al., 2016](#); [Evans and Garthwaite, 2014](#)) and engage in healthier

find evidence of mental health effects on later generations, which they attribute to biological rather than social processes. [Bütikofer and Salvanes \(2020\)](#) find that in Norway, generational persistence in educational attainment was mitigated by a 1940s tuberculosis control program.

¹⁰Two additional studies examine the multi-generational effects of post-natal early life educational interventions that include a health component. [Barr and Gibbs \(2019\)](#) and [Rossin-Slater and Wüst \(2020\)](#) examine the effects of exposure to preschool in the U.S. and Denmark, respectively, on educational outcomes for the next generation and find positive effects. In addition, Barr and Gibbs find evidence of reduced teen pregnancy and criminal activity in the second generation.

¹¹Several studies of Medicaid’s prenatal expansions document increased use or improved timing and adequacy of prenatal care ([Currie and Gruber, 1996b, 2001](#); [Dave et al., 2008](#); [Dubay et al., 2001](#); [Howell, 2001](#)). The medical literature also hints that prenatal care may be a pathway for improving the intergenerational transmission of health ([Lu et al., 2003](#)).

behaviors (Hoynes et al., 2015).

3 Medicaid and the 1980s Expansions

Medicaid currently provides health insurance coverage for nearly half of all births in the U.S. (Markus et al., 2013). Eligibility criteria for pregnant women were relatively restrictive until the 1980s, when a series of state and federal policy changes greatly expanded access. We describe the inception of the Medicaid program, including which groups were initially eligible, and the later policy changes that expanded eligibility, below.

Created in 1965 as part of the Social Security Amendments, Medicaid was initially available to low-income, non-disabled women of reproductive age who received cash assistance through the Aid to Families with Dependent Children (AFDC) program.¹² Eligibility for AFDC was restricted to single women with at least one dependent child. Low-income women with first time pregnancies and those with marital partners did not qualify for coverage. Moreover, AFDC income eligibility thresholds were generally much lower than the federal poverty line. The average threshold was 61% of the federal poverty line (FPL) in 1979, and ranged from 24% to 99% across states.¹³

Because Medicaid is a joint federal-state program, the federal government sets mandatory eligibility requirements, but states have some flexibility to extend eligibility to other especially needy population groups. Dating back to 1966, a number of states exercised different options to extend eligibility to certain pregnant women not eligible for AFDC. The options offered coverage to specific groups of pregnant women, such as first-time pregnant women who would later qualify for AFDC, or pregnant women in two-parent families where the principal earner was unemployed and who met the income and resource requirements for AFDC.¹⁴ Some options extended eligibility to all pregnant women who were financially eligible for AFDC, but who did not meet the program’s family structure requirements (i.e. women who were *not* single mothers with dependent children). Adoption of these options was not uniform across states, and, prior to the 1980s, there were substantive differences in eligibility criteria that were applied to pregnant women.

Beginning in the early 1980s, there was emerging national consensus on the importance of prenatal care for pregnant women, which led to major changes to the Medicaid program (Howell, 2001). More states began to exercise options to expand Medicaid eligibility to pregnant women who did not meet the family structure requirements for AFDC, but who did meet the AFDC financial eligibility criteria. This wave of new optional state expansions was followed by two federal mandates requiring that *all* states expand eligibility to these women. These changes caused abrupt expansions in states with historically less generous eligibility rules and lower eligibility levels.

Figure 1 Panel (a), which is based on information we have gathered on state policy rules back to 1975, documents that these changes had a large effect: between 1975 and 1988 the share of women who qualified for pregnancy-related Medicaid climbed from about 12 to 20 percent.¹⁵ Here, and throughout

¹²Since Medicaid was initially tied directly to AFDC, many women received benefits from both programs. However, as we describe in detail in Appendix Section A.1, the policy variation we exploit is driven by expansions to Medicaid and not by changes in AFDC benefits.

¹³Authors’ calculation based on payment standard for a family of 3 in 1979.

¹⁴Appendix Section A provides a detailed discussion of each of the different state options described in this section.

¹⁵Additional information about the calculator and data used to calculate eligibility in this figure is provided in Section 4 and Appendix Section A.

the rest of the paper, we multiply all proportions by 100 for ease of presentation. Panel (b), which uses data on hospital discharges for labor and delivery between 1979-1988, shows that this increase corresponded to a steep increase in the share of mothers who were enrolled in the Medicaid program at the time of their child’s birth.¹⁶ To our knowledge, this is the first time that administrative data have been used to document that take-up patterns among pregnant women mirrored these eligibility expansions.¹⁷

Following the original terminology used by Currie and Gruber (1996b), we refer to these changes in eligibility as the “targeted” expansions. They differ from “broad” expansions that occurred later because they were aimed at very low income pregnant women (i.e. those with income levels below AFDC thresholds). Beginning in the late 1980s, the broad expansions allowed states to further expand coverage to pregnant women and children with higher incomes. We cannot make use of variation generated by broad changes in our analyses because most of the affected cohorts are still too young to allow for multi-generational analyses. We note, however, that previous studies have documented that the targeted expansions had stronger effects on treated cohorts’ outcomes (Currie and Gruber, 1996b; Miller and Wherry, 2019), making them a preferred candidate to test for multi-generational effects. We also note that pregnant women who enrolled in Medicaid received coverage for prenatal care and services, hospital and postpartum care, and one year of Medicaid eligibility for their newborns (Congressional Research Service, 1988).

With this background in mind, we investigate multi-generational effects of *in utero* Medicaid access by focusing on the offspring of cohorts who were born between 1975 and 1988. We refer to the mothers in these cohorts as the “first” (exposed) generation, and to their later infants as the “second” generation.

4 Empirical Strategy

4.1 Background

Currie and Gruber examine the impact of the prenatal Medicaid expansions on first generation infant health by estimating a model that exploits variation in the magnitude of the expansions across states and over time. Specifically, they estimate:

$$y_{nb} = \alpha + \phi Elig_{nb} + \mu_n + \lambda_b + \gamma X_{nb} + \epsilon_{nb} \tag{1}$$

where y_{nb} is the fraction of infants born in state n and year b who were below the low birthweight threshold, μ_n and λ_b are state of birth and year of birth fixed effects, respectively, and X_{nb} is a vector of state-year control variables that reflect the demographic, economic, and policy environment. $Elig_{nb}$ is the percent of women of child-bearing age in the state-year who were eligible for Medicaid in the event of a pregnancy. Currie and Gruber’s analyses, as well as later studies that build upon this research design, begin in 1979, on the eve of the Medicaid program’s rapid expansion.

Acknowledging that changes in states’ demographic and economic conditions could cause changes

¹⁶While the changes track quite well across both of these measures, there are a few years in which coverage actually exceeds eligibility. We believe this is due to the fact that our eligibility estimates are based on women of child-bearing age, whereas our estimates of coverage are derived from a sample of women actually giving birth. These groups may differ on some important dimensions such as age and income.

¹⁷Dave et al. (2010) use this source of hospital discharge data to examine the effects of later Medicaid expansions (1985-1996) on the health insurance coverage of pregnant women at the time of child birth.

in the percent eligible for Medicaid even without a change in policy, Currie and Gruber pioneered a “simulated instrument” approach in which they instrument actual eligibility with a simulated measure that isolates changes driven only by changes in Medicaid rules and is independent of changes in states’ demographic and economic characteristics. The simulated measure is constructed by applying Medicaid eligibility rules in each state and year to a *national* sample of potential mothers (women ages 15-44) drawn from each year of the Annual Social and Economic Supplement of the Current Population Survey. Our analyses also rely on simulated eligibility measures to capture changes in Medicaid policy, following the convention that has been adopted by much of the literature.¹⁸ Appendix Figure A1 shows that national trends in actual and simulated eligibility measures are very closely related.

Using this model, Currie and Gruber find that a 10 percentage point increase in prenatal Medicaid eligibility under the targeted expansions reduced the first generation’s incidence of low birthweight by 2.6 percent.¹⁹ Subsequent papers have extended their approach to look at outcomes associated with later childhood expansions by replacing $Elig_{nb}$ with a measure of cumulative or average childhood eligibility from birth to age 18 (e.g. Brown et al., 2020; Cohodes et al., 2016; Thompson, 2017) and using a comparable simulated eligibility instrumental variables approach. Noting that the prenatal period is a particularly receptive stage of development, with the potential to yield large returns on investment, Miller and Wherry (2019) further extend this design by including separate measures of prenatal and later childhood eligibility. They find that this distinction is important, with most of the long-run health improvements driven by the expansions in prenatal access.

It would be natural to extend this framework to examine the expansions’ impact on second generation outcomes.²⁰ As Currie and Gruber note, however, identification in this model rests on the assumption that state Medicaid policy was exogenous to treated cohorts’ birth outcomes. Although previous analyses have included a large number of state and year varying controls, some researchers have argued that the changes are not “randomly occurring natural experiments,” and they speculate that estimates of the expansions’ impacts reflect other underlying state-level changes (Dave et al., 2008). The traditional research design has previously been estimated with very limited years of data from the pre-expansion era, which restricts researchers’ ability to test for differential changes across states prior to the expansions. Therefore, we have collected additional years of information on state program rules, and use these data to document changes in state-level prenatal Medicaid eligibility back

¹⁸To create the simulated *in utero* measure, we use a national random sample of 3,000 women from each year of the Annual Social and Economic Supplement of the Current Population Survey (CPS). Since childhood eligibility was also changing for these cohorts, we also examine whether the changes in prenatal eligibility were correlated with changes in childhood eligibility, and whether including controls for childhood eligibility affects the estimates. To construct measures of simulated childhood eligibility, we use national random samples of 1,000 children at each age between 1 and 18 in each survey year and calculate the percent of children in each state, year, and age that would be eligible for Medicaid based on state eligibility rules. We then add the estimates across ages for each cohort, to create a measure of cumulative simulated eligibility throughout childhood for each cohort and state and use this as a measure of generosity during childhood. We describe these calculations in more detail in Appendix Section A.3.

¹⁹We replicate this result using their empirical approach in Appendix Table A1. Note that, for the purpose of this replication, we follow Currie and Gruber and express the outcome as the number of low birthweight births per 1000 infants. The first column displays the results reprinted from their paper and the second column is our replication exercise. The remaining columns update the analysis to cluster standard errors by state (column 3), add the state by year controls we implement in our model, described in detail below (column 4), and weight by the number of births in each cell (column 5).

²⁰An earlier version of this paper used this approach. That version is available upon request.

to 1975.²¹ Then we implement a more transparent event study design that allows direct examination of the extent to which state-level pre-period outcomes varied with state-level Medicaid expansions, and addresses concerns about potential contaminants.

Our analyses start by documenting the evolution of state-level Medicaid eligibility. Appendix Figure A2 shows the time path of eligibility from 1975-1988 for each of the fifty states. It is immediately clear that states exhibit different expansion patterns, and that these patterns fall into two dominant types: many states experienced small, gradual increases in eligibility throughout the period (e.g. California and Rhode Island), while others exhibit flat, low levels of prenatal coverage during the 1970s that are later punctuated by a large, abrupt, increase (e.g. South Carolina and Maine).

This is exactly what the discussion in Section 3 should lead us to expect. Specifically, in early years, states that initially embraced strict eligibility criteria and did not avail themselves of the state options, exhibit low levels of eligibility. In later years, we observe discrete jumps in these states that correspond to a large, voluntary state-level expansion or one of the later federally mandated expansions. In contrast, states that took early advantage of the existing state options to cover low-income pregnant women have relatively high levels of eligibility from the beginning of the study period. As a result, these states exhibit smaller, more gradual increases that are less pronounced in the wake of the subsequent state and federal changes. These two different eligibility trajectories will be demonstrated visually in the next subsection.

4.2 Classification of Treatment and Control States

Using Appendix Figure A2 as a guide, we organize states into “treated” states (those that experienced a sharp jump in eligibility) and “control” states (those for whom eligibility trended smoothly). Our event study is based on differences in the evolution of prenatal eligibility across these two groups. The treatment group consists of the 28 states in which eligibility was stagnant for a minimum of the five years between 1975 and 1979, and that later experienced a large positive shock. We focus on shocks that took place between 1980 and 1985 to ensure that we have at least four “post-event” years over which to observe the expansions’ effects. We include the 22 remaining states as “control” states to help identify secular changes that were separate from the Medicaid expansions.²²

Figure 2 shows which states we classify as control states, which states we classify as treatment states, and the timing of the expansion for the treatment states. Given that “treatment” requires a large, abrupt increase in eligibility, it is unsurprising that treated states are disproportionately located in the South and Midwest; Medicaid was less generous in the pre-expansion period in these states, and transfer policies have historically been less generous in these parts of the country as well.²³

²¹Information on eligibility prior to 1975 is very spotty, therefore we begin our sample period in 1975.

²²We did not have a specific algorithm to define treatment and control states or event time. Rather, each of the four authors individually classified states based on Appendix Figure A2, and then we decided on a consistent classification as a group. We relied on measures of simulated eligibility, rather than actual eligibility, to ensure that changes in eligibility are driven by policy changes and not by demographic or economic shifts, and there were few classification discrepancies across the authors. States with large changes outside the 1980-1985 period are included in the control group, but the few states that fall into this category do not affect our results: we show that estimates are similar when we drop the control states from the analysis.

²³Prior work has found that racism played a role in generating a less generous safety net (Lee and Roemer, 2006), and, Appendix Table A2 shows that treated states have slightly higher Black populations than control states. We multiply all proportions by 100 for ease of presentation. There are a few small differences in other demographic and economic characteristics across treatment and control states. Our analyses include state fixed effects to account for any time-invariant

Figure 3 Panel (a) provides further insights into the sources of our identifying variation by combining all of the treated states together, and documenting how the changes in eligibility break down by source. The sources are shown additively from most to least restrictive (shown from lowest to highest lines on the graph), and include changes in eligibility driven by: changes in AFDC rules (solid blue line), targeted expansions occurring via state options (long dashed red line), targeted expansions occurring under federal mandates (short dashed green line), and broader income-based expansions (dash dot yellow line). The last category is concentrated at the end of the period, and reflects optional state expansions that included pregnant women with family incomes up to the poverty line.²⁴

It is easy to see that prior to the 1980s, Medicaid eligibility in treated states was largely determined by state AFDC rules, and that these states were less likely to take up state options than the control states. Beginning in the early 1980s, a few treatment states increased eligibility through the state options. This was followed by a sizeable bump under the federal mandates. Between 1975 and 1988, prenatal eligibility roughly doubled in treatment states, from just under 9 percent to 18 percent. In contrast, Figure 3 Panel (b) shows that control states started off with higher eligibility, because they had already implemented the state options, and that eligibility gradually increased over time as more options were adopted. As a result, the federal mandates had smaller effects on eligibility.

Not only is the change in eligibility greater for treatment states than control states, but the women affected are more disadvantaged due to the more stringent baseline eligibility rules for these states. To demonstrate this, we compare the characteristics of women who gained eligibility over this period in both groups of states in Appendix Table A3.²⁵ The women in treated states are noticeably more disadvantaged in terms of their family income and poverty rate than women in control states. Notice that nearly half of the women gaining eligibility in treatment states have family incomes below 50% of the FPL, versus only 13 percent in the control states. Women in the treatment states are also more likely to be single and non-white.

Finally, Figure 4 demonstrates that among treated states there is also variation in when the large jumps occur. States are grouped together by the initial year of the abrupt increase (the “expansion year”), and eligibility is plotted relative to the year prior to expansion, with the number of states expanding in each year shown in parentheses. While the most common expansion years are 1982 (11 states) and 1985 (7 states), when federal policy provided states with new options to expand coverage (1982), and mandated expansion of coverage to more women (1985), there are also states that experience large expansions in every other year between 1980 and 1985.²⁶ In our main event study framework, we rely on the differential timing of expansions across states, as well as differences between treated and control states, to identify the effects of Medicaid separately from time trends.

differences, and we show that our estimates are robust to including a large number of state-year varying economic and policy controls, and region-year fixed effects that control for common shocks to outcomes within regions over time.

²⁴Under the Omnibus Budget Reconciliation Act of 1986, states were allowed to expand Medicaid to pregnant women with family incomes below 100% FPL starting in April 1987. See additional discussion in Appendix Section A.1, along with more detailed breakouts of the different eligibility paths in Appendix Figures A3 and A4.

²⁵Using the CPS, we calculate mean characteristics for the women in treatment states who are eligible during the post-period (defined using each state’s specific treatment timing), but who were not eligible under the rules that were in place during the last pre-treatment year. We calculate mean characteristics for women in control states who are eligible during 1982-1988 but were not eligible under the rules in place in 1981.

²⁶See Appendix A.1 for more details on these policy changes.

4.3 Event Study

The event study takes the following form:

$$y_{nb} = \alpha + \sum_{t=-5, t \neq -1}^3 \beta_t 1\{b - e_n^* = t\} + \mu_n + \lambda_b + \gamma X_{nb} + \epsilon_{nb} \quad (2)$$

where y_{nb} is an outcome for individuals born in state n and year b . Initially, we estimate the expansions' effects on the first generation's *in utero* Medicaid eligibility and coverage at birth. Then, in line with the existing literature, we examine the first generation's health outcomes at birth. The primary goal of our study, however, is to understand whether the benefits associated with the first generation's prenatal access to Medicaid had spillover effects onto their offspring. To do so, we replace first generation outcomes with the outcomes of the offspring of women who were born in state n and year b .

The key regressors are the series of dummy variables $1\{b - e_n^* = t\}$ that take on a value of one for each event time year, where event time is defined for each treated state relative to the year in which it first experienced a discrete jump in eligibility (e_n^*). We omit the year before each state's large expansion $t = -1$, so the estimated β_t s are relative to the year before the expansion occurred. For example, β_1 is the effect one year after the discrete change in eligibility, relative to one year before the jump.²⁷ We do not define event time for the control states since they do not experience a large discrete expansion. The control states help estimate secular trends across cohorts and the effects of the control variables.²⁸

As in previous studies, we include first generation state of birth fixed effects, μ_n , to account for fixed differences in the outcomes of mothers and their children across states, and first generation year of birth fixed effects, λ_b , to account for national changes over time. Our baseline model also follows the literature by incorporating a large number of first generation state and year of birth control variables including the unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws, state Medicaid restrictions for abortion, and population demographic controls for each state and year. Sensitivity analyses include versions of the model that eliminate the control variables, or add additional controls, such as region-year fixed effects, and state-year controls at the time of the *second* generation's birth, which we discuss in more detail below. Our results are not sensitive to their inclusion. Our estimates are weighted by the size of the second generation birth cohort.²⁹ We cluster our standard errors by mothers' state of birth.

Appendix Figure A5 plots second generation health outcomes by event time, where each treatment timing group is shown in a different color. Control states are shown in pink, with event time centered at 1982. It is easy to see that, consistent with the decades long national trend in infant health ([United](#)

²⁷By choosing to classify treated states as those with a large, abrupt eligibility change in the middle of the 1975-1988 period, we are able to observe at least 5 years before the event and 4 years after the event. We bin event time observations that are more than 5 years before the event and more than 4 years after the event. We estimate, but do not report these coefficients, because they are based on an unbalanced sample. Binning allows us to separately identify treatment effects from secular time trends even when we do not include the control states in the model ([Schmidheiny and Siegloch, 2019](#)).

²⁸A potential concern is that using partially treated states (those with smaller and less abrupt changes in eligibility) as a control group will lead to biased estimates ([Schmidheiny and Siegloch, 2019](#)). Figure 9 documents that the pattern of estimates is similar (albeit with larger standard errors) when the control states are not included in the analyses. We also note that there is little difference in the magnitude of the eligibility changes across treated states (Figure 4). Therefore, we do not use variation in the magnitude of the expansions as an additional source of identifying variation in our model.

²⁹The results are robust to weighting by the size of the first generation cohort instead.

Health Foundation, 2020; Wang, 2010), and pre-dating the expansions, second generation health was declining. Linear trend estimates based on *pre-expansion* data are shown in solid lines, and clearly vary across state groups, possibly reflecting differing trends in the demographics of women giving birth, adoption of health technologies, or changes in state policies (including expansions of prenatal Medicaid eligibility that occurred before those that are the focus of this study). Appendix Figure A6 documents that first generation health also trended differently across state groups. To ensure that our estimates are not contaminated by state differences in pre-expansion trajectories, we remove linear pre-trends following the two-step de-trending procedure implemented in Goodman-Bacon (2020) and described in more detail in Section 5.

This research design is a departure from the conventional approach, which exploits state and cohort variation in the timing and magnitude of the expansions but does not address potential differences in states' pre-period trajectories. Another advantage of our research design is that it provides non-parametric estimates of the expansions' dynamic effects. Assuming that treatment effects are homogeneous across treatment groups, estimates generated by the event study will be unbiased, even in the case of staggered treatment timing. As described below, we examine the importance of this assumption using the method proposed by Callaway and Sant'Anna (2020) and obtain similar results.

5 Data

Our measures of Medicaid eligibility are based on data in the Annual Social and Economic Supplement of the Current Population Survey and are described in more detail in Section 4 and Appendix Section A.3. We also analyze Medicaid take-up among pregnant mothers using restricted-use data from the National Hospital Discharge Survey (NHDS). The NHDS data provide discharge-level information for a nationally representative sample of non-Federal, short-stay hospitals, and include the expected payer for the hospital visit. This allows us to identify births paid for by Medicaid. Because the NHDS data are based on administrative hospital records, they are not subject to misreporting issues that are common in survey data (see Davern et al., 2009; Klerman et al., 2005), but unlike our other data series, they are not available before 1979.³⁰ In our event study analyses, we estimate the pre-period coefficients using all available data. We also apply the NHDS survey weights. Using diagnosis and procedure codes based on the International Classification of Diseases coding system,³¹ we identify 187,488 labor and delivery hospitalizations between 1979 and 1988 that include information on the expected payer.

To conduct our main birth outcome analyses, we use the 1975-2017 U.S. Vital Statistics Natality Data Files.³² These files contain individual birth records for the full census of U.S. births. The data include information on infants' health, gender, parity, and year and state of birth. They also include

³⁰We acknowledge that if some pregnant women did not enroll in Medicaid until delivery then the NHDS data could produce an upper bound estimate of the percent of infants who gained *in utero* coverage. Using Medicaid claims data from four states, Ellwood and Kenney (1995) document that the share of women with Medicaid deliveries who enrolled during the *first trimester* were 50 percent (California), 52 percent (Georgia), 69 percent (Michigan), and 49 percent (Tennessee) in 1987. The authors also document higher first trimester enrollment among women receiving cash assistance, ranging from 66-79 percent. These take up rates were almost certainly higher at later stages of pregnancy, but before delivery, suggesting that the vast majority of women enrolled prior to delivery. Related to this, we remind the reader that all children who were covered at the time of delivery were eligible for coverage during their first year of life.

³¹Details are found in Appendix Section D.

³²Beginning in 2005, mother's state of birth information is only available in the restricted access data, so we use the restricted data from 2005-2017.

detailed demographic information about each infant’s mother, including her state of birth and age, which allows us to approximate her year of birth.³³ The latter variables are critical to our second generation analyses, as they allow measures of *in utero* Medicaid eligibility to be matched to each mother in the first generation. In the second generation analyses we exclude infants whose mothers were born outside of the United States, as well as mothers born in Arizona, which did not adopt a state Medicaid program until 1982.

We begin by using the event study framework to reassess Currie and Gruber’s first-generation results. Currie and Gruber focus on the incidence of low birthweight (less than 2500 grams) among infants born between 1979 and 1992. Appendix Table A1 shows that we are able to replicate their results when we apply their research design to the same cohorts. Moving forward, our event study analysis changes the included cohorts in two ways. First, because a convincing event study requires pre-period data, we add the four cohorts born between 1975 and 1978. Second, the second generation analyses would ideally include all births to women born between 1975 and 1992, but many women born in the 1990s have not given birth by 2017, so we focus on cohorts born between 1975 and 1988. This ensures that all first generation cohorts that were affected by the targeted prenatal expansions are included, while allowing us to observe each cohort’s fertility through age 28. The analyses of second generation outcomes is restricted to infants whose mothers meet this age criteria to ensure that each maternal cohort contributes equally to the identifying variation, and that cohort-level comparisons are across women who are giving birth at the same age. We further restrict the second generation analyses to first births only, as this generates a more representative sample of births across cohorts. During our time frame, 81% of first births, and 62% of all births, were to women aged 28 or younger.³⁴ We test the robustness of our results to both of these restrictions below.

The second generation analyses include infants born between 1990 and 2017.³⁵ We examine the impacts of the prenatal expansions on the next generation’s average birthweight and incidence of low birthweight. These outcomes are standard measures of infant health and are highly predictive of later life health, cognitive and economic outcomes (e.g. Black et al., 2007; Figlio et al., 2014). We also explore other outcomes available in the national natality data, including very low birthweight (weighing less than 1500 grams), gestational length (in weeks), preterm (born before 37 weeks gestation), and very preterm (born before 28 weeks gestation). As described in Kramer (1987a,b), birthweight is determined in part by gestational length. For this reason, we also consider a common measure of intrauterine growth: whether the infant is below the 10th percentile of birthweight for gestational age (“small for gestational age”).³⁶

We collapse the data into cells based on the first generation’s state of birth and year of birth. We then merge each cell with corresponding measures of actual and simulated Medicaid eligibility, and with information on states’ economic conditions (state unemployment rate and per capita income), demographic composition (age distribution, marital status, educational attainment and race), safety

³³Miller and Wherry (2016) show that using age to impute birth year leads to nearly identical simulated eligibility assignments as using actual birth year.

³⁴Authors’ calculations from the Vital Statistics Natality Files.

³⁵We restrict births to women ages 15 or older. The second generation sample, therefore begins in 1990, since this is year that the 1975 cohort turns 15.

³⁶This variable is constructed using two potentially noisy variables, increasing the likelihood that the variable is measured with error.

net generosity, and abortion policies. Additional details about these control variables and sources are provided in Appendix Section D.

As discussed in Section 4, second generation outcomes were trending differently in treatment and control states in the pre-expansion era. To ensure that our estimates are not contaminated by differential linear pre-trends, we directly remove them using the two-step de-trending method implemented in Goodman-Bacon (2020). For each treated state and outcome,³⁷ we estimate a linear trend using data only from years prior to the expansion, then extrapolate this estimated trend through all years of data, and subtract the predicted values of each outcome from the observed values. Since control states do not have a clear expansion year, the linear “pre-trend” is estimated using data from 1975-1981, as 1982 is the year of the national trend break in eligibility (Figure 1). The underlying assumption in the event study model is therefore that outcomes would have continued to follow their linear pre-trends in the absence of the expansions. Since linear pre-trends are removed from our analysis, the event study provides information on any remaining non-linear differences across states in the pre-period.

6 Results

6.1 Eligibility Estimates

Figure 5 provides “first stage” event study estimates based on equation (2), where Panel (a) shows the estimated change in prenatal eligibility and Panel (b) shows the estimated change in simulated prenatal eligibility, both of which are calculated using the CPS, as described in Appendix Section A.3. Recall that both of these variables capture the percent of women who are *eligible* for Medicaid, rather than the percent who *received* Medicaid. The horizontal axis denotes the number of years before and after the expansion. Event time zero is the first year of the expansion. We omit event time -1, so all estimates are relative to the year before the expansion. The estimates are plotted along with their 95% confidence intervals.

It is immediately clear that after accounting for differences in linear pre-period trends there are no remaining pre-expansion differences between treatment and control states, and that in treated states, the period following the expansion is associated with an abrupt increase in both actual and simulated eligibility. This is expected because of the way in which we assign treatment, and is evidence of a strong first stage. Our estimates indicate that four years after the initial expansion, simulated eligibility had increased by 6.6 percentage points. Coefficient and standard error estimates are reported in Table 1.

Appendix Figure A7 shows the results of a variety of additional analyses, with the baseline estimates shown as solid black circles. Alternative specifications include models that eliminate the state-year controls (solid blue triangles), and add region-year fixed effects (solid green square). The estimates barely change across these specifications indicating that the eligibility expansions used for identification were not strongly related to other state-year policy, demographic, or economic changes, or to other changes within regions over time. The results are very similar when we do not de-trend (hollow gray circle), weight by the number of first generation (female) births rather than the number of second generation births (hollow purple square), and when we estimate the model without including the control states (hollow red triangle).³⁸

³⁷An exception is that we do not de-trend the NHDS payment data, given the limited number of pre-expansion years.

³⁸In Appendix Figure A8, we also document that our results are robust to an alternative measure of simulated eligibility

6.2 Medicaid Coverage at Birth

Although Figure 5 confirms that eligibility increased in the wake of the expansions, we would not expect broader eligibility to translate into better infant health outcomes without a corresponding increase in program take-up. To investigate Medicaid take-up, we estimate the event study model using NHDS data on Medicaid coverage at the time of birth. Figure 6 shows estimates based on equation (2), where the dependent variable is the percent of hospitalized births that were covered by Medicaid. As in Figure 5, the pre-expansion estimates are close to zero, and there is a clear increase in coverage following the initial expansion.³⁹ Importantly, within four years of the initial expansion, the percent of births covered by Medicaid had increased by 4.6 percentage points, which, when compared to the 6.6 percentage point increase in simulated eligibility, implies a 70% take-up rate among newly eligible mothers. In contrast to eligibility in Figure 5, however, the increase in coverage phases in more slowly, suggesting that it took time for program take-up to fully respond. Given this, we anticipate a similar ramp-up pattern in the health estimates.

6.3 First Generation Estimates

Having established the event study’s validity, we use the same model to re-examine Currie and Gruber’s first generation results. Independent of our upcoming second generation analyses, here we make an important contribution to the literature by shedding light on the extent to which previous estimates of the expansions’ effects may have been confounded by the presence of other changes that were occurring within states.

Figure 7 and Table 1 present our event study estimates of the expansions’ effects on the percent of first generation newborns who were low birthweight. The estimates strongly support Currie and Gruber’s original findings. Our pre-period estimates are close to zero and not distinguishable from each other, and there is an abrupt decrease in low birthweight in the treated states, compared to the control states, immediately following treated states’ initial expansion. The magnitude of the point estimates grows over the post-expansion period, so that, four years post treatment, the incidence of low birthweight has declined by 0.25 percentage points ($p < 0.001$), or 3.3 percent of the treated states’ pre-period mean (Table 1).⁴⁰ Our confidence intervals include effects as small as a 0.09 percentage point reduction (1.2% when compared to the treated states’ pre-expansion mean) and as large as a 0.40 percentage point reduction (5.4%). When coupled with our estimated 6.6 percentage point increase in prenatal eligibility, the results suggest that a 10 percentage point increase in eligibility reduces the incidence of low birthweight in the first generation by 0.37 percentage points, or about 5 percent.⁴¹ As

that is based on a fixed national sample where the data are pooled across all of our sample years (1975-1988). This version of simulated eligibility ensures that any changes in national population demographics over time are not driving variation in the simulated measure. Appendix Figure A9 shows that our estimates of the expansions’ effects on eligibility are robust to the inclusion of additional control variables that account for factors that might change between the time the mother (first generation) was born and the child (second generation) was born. We discuss these robustness checks in more detail when we examine second generation health outcomes in Section 6.4.

³⁹Note, however, that as discussed in Section 5, the pre-expansion coefficients in this analysis are estimated using all available data, but necessarily include different treatment states at each event time (i.e. are “unbalanced”) due to the limited years of pre-data available for the NHDS. This is not the case for any post-expansion coefficients.

⁴⁰Recall that for ease of presentation, we multiply all variables with values between 0 and 1 by 100.

⁴¹Note that the baseline mean among women who gained eligibility may be substantially higher than the population average because those who gained eligibility were extremely disadvantaged. For example, among children living in the lowest income households in the 1981 National Health Interview Survey (under \$3000 in annual income), the incidence of

a point of comparison, Currie and Gruber estimate that a 10 percentage point increase in eligibility led to a 2.6 percent decline in low birthweight, which falls squarely within our 95% confidence interval.

To interpret these effects as causal, it must be the case that the timing of the expansions was unrelated to other factors that affect infant health. For example, one might be concerned that a state's decision to expand was influenced by the state of its economy. To check for this, we examine how our estimates change under the same set of alternative specifications described in Section 6.1, which include models that eliminate the state-year controls or add region-year fixed effects, a version of the baseline model that does not remove linear pre-trends, and a version of the model that does not include the control states. Appendix Figure A10 Panel (a) shows that estimates are similar across the specifications. The coefficient estimates during the post-period are smaller when we do not account for pre-existing linear trends, supporting the possibility that control states' early adoption of the expansion options may have already altered their infant health trajectories prior to the 1980s. Unsurprisingly, given the reduced sample size, confidence intervals are larger when we estimate the model without including the control states. Even with the loss of precision that accompanies some of these checks, however, the pattern of results remains very similar.

Taken as a whole, our event study analysis bolsters the credibility of Currie and Gruber's landmark findings. The new research design also provides insights into the time path of effects, specifically, that the expansions' impact on the first generation's health took time to fully ramp up. A likely explanation is that it took time for pregnant women to learn about their eligibility and enroll in the program. Moreover, although improvements in the first generation's birthweight are not a necessary pre-condition for the presence of second generation effects, the results presented in Figure 7 allude to mothers' health at birth as a potential conduit, as previous studies have documented intergenerational linkages in low birthweight (e.g. Currie and Moretti, 2007).⁴²

low birth weight was 17.4 percent, which is more than twice as large as the baseline incidence of low birthweight in the treated states (7.44 percent, see Table 1). Even so, the effect sizes contained within the 95% confidence interval around the low birthweight estimate are sufficiently large to suggest that they may not arise from increases in prenatal care alone. A combination of pathways may contribute to the effects. For example, prenatal care may put mothers in touch with other social services: three quarters of women receiving Medicaid funded prenatal care in 1988 reported that they received WIC during their pregnancy (see Table 1 in Miller and Wherry, 2019), and forty percent reported that they learned about WIC through a doctor, nurse or health provider. Related to this, it is possible that prenatal counseling changes nutrition or substance use behaviors sufficiently to affect birth outcomes. Again, as discussed in Miller and Wherry (2019), 90% of women who obtained prenatal care through Medicaid received nutritional counseling, 95% were counseled to take vitamin supplements, 69% received counseling with respect to smoking, and 61% received counseling with respect to alcohol. These types of instructions may be more common among low-income women, or they may have a stronger impact on low-income women relative to the average woman who gets a prenatal care visit. Another potential pathway is through an increase in families' financial resources. Based on the following website, <https://www.cryo-cell.com/blog/april-2017/when-childbirth-cost-100-dollars>, out-of-pocket expenditures for an uninsured birth during this time period were about \$1500 (\$6500 in 2012\$s). Multiple studies document that cash and near-cash transfers improve birth outcomes among infants born to low-income women (e.g. Kehrer and Wolin, 1979; Hoynes et al., 2015; Almond et al., 2011; East, 2020). Another possible pathway is that insurance may reduce maternal stress. A significant body of research links prenatal maternal stress to adverse birth outcomes, including low birthweight (Persson and Rossin-Slater, 2018). Moreover, poor urban women from minority backgrounds are twice as likely as middle-class women to meet diagnostic criteria for major and minor depression during pregnancy and the postpartum period (Grote et al., 2010). Other potential mechanisms include changes in maternal employment, or spillover effects onto non-participants.

⁴²We also consider the expansions' effects on average birthweight and the incidence of very low birthweight, which are not examined in the original Currie and Gruber paper. Event study estimates and robustness analyses are provided in Panels (b) and (c) of Appendix Figure A10 and Appendix Table A4. The estimated effects are suggestive of expansion related declines in very low birthweight. We do not examine changes in outcomes related to gestational length because prior to 1981 the birth certificate records have a high rate of missing values for this variable.

6.4 Second Generation Estimates

Figure 8 shows the relationship between mothers' *in utero* Medicaid access and second generation outcomes. Focusing first on birthweight (Panel (a)), we see evidence of an increase following the expansions. The point estimates grow over time, reaching 4.7 grams four years after expansion (see Table 1). The 95% confidence interval allows us to rule out effects of more than 8.2 grams and less than 1.2 grams. Although the impacts on some of the other infant health outcomes are less precisely estimated, we see largely similar patterns, particularly at the lower end of the birthweight distribution (low birthweight and very low birthweight).⁴³ We also see evidence of a decrease in very preterm births. As discussed in Section 5, low birthweight and prematurity are of particular interest because they are closely linked to other early and later life health and cognitive outcomes. Also, low birthweight and preterm births are associated with maternal health characteristics that are known to have been improved by the prenatal expansions (Institute of Medicine, 2017; Miller and Wherry, 2019).⁴⁴ We also see some evidence of improvements in small for gestational age, which is indicative of gains in intrauterine growth.⁴⁵

As with the first generation estimates, Figure 9 shows that the second generation results are, for the most part, similar when we employ the specification checks described in Section 6.1. Our second generation estimates often have large confidence intervals associated with them, but across all of these checks the patterns remain very stable, even if not always statistically different from zero at conventional significance levels. Excluding the state-year covariates, or adding region-year fixed effects does not change the results, which again demonstrates that these large expansions were not correlated in a meaningful way with state-level changes in demographics, policies, or economic conditions, or with time-varying changes within regions. The figure also documents similar patterns when we use different weights (mother rather than child cohort size). We see larger deviations from the baseline estimates when we do not account for pre-existing differences in state trends (gray circles in the figure). When we remove control states from our sample, the estimated intergenerational effects often become larger but more imprecise.⁴⁶

Appendix Figure A12 shows the sensitivity of the estimates to additional sets of state-year controls. During the 1980s and 1990s, Medicaid coverage was also extended to older and higher income children. If the prenatal expansions we focus on are correlated with these other childhood expansions, then we might be erroneously attributing the observed health improvements among second generation infants to their mothers' *in utero* coverage when, in fact, the improvements result from mothers' increased Medicaid access in later childhood. Most studies estimate the impact of expanding children's access

⁴³The 95% confidence intervals around the year four point estimates for low birthweight (very low birthweight) allow us to rule out declines of more than 0.27 (0.14) percentage points and gains of more than 0.03 (declines of less than 0.02) compared to baseline means of 7.52 and 1.45.

⁴⁴For example, low birthweight has been linked to chronic hypertension, pre-pregnancy diabetes, and maternal obesity (Institute of Medicine (US) Committee on Understanding Premature Birth and Assuring Healthy Outcomes, 2007).

⁴⁵The differences in outcomes between treated and control states are evident even in the raw means. In Appendix Figure A11 we plot mean outcomes by event time, across treatment and control groups (solid black and orange lines). The trends are very similar prior to the expansions, and then deviate for the outcomes for which we find (at least suggestive) evidence of effects – birthweight, low birthweight, very low birthweight, very preterm, and small for gestational age. These same patterns persist after we residualize with the state-year controls in our baseline models – shown in the gray and orange dashed lines.

⁴⁶We report the coefficient and standard error estimates from all of these robustness checks and for each outcome in Appendix Tables A5-A11.

to Medicaid using a single index that aggregates eligibility across the length of childhood, without evaluating linkages between the *in utero* and later childhood expansions. In Appendix Figure A13, we use the same event study framework to directly estimate the relationship between the prenatal expansions and the childhood expansions, and find no evidence that they are correlated.⁴⁷ Therefore, it is unsurprising that our second generation estimates are very similar when we add measures of later childhood eligibility to our baseline model (shown in the solid purple triangles in Appendix Figure A12).

Appendix Figure A12 also documents that the results are similar when we control for mother’s own eligibility during adulthood (open green circles), and when we control for the second generation’s own prenatal eligibility (open pink triangles). They are also unaffected when we include the same set of state-year controls listed in Section 4 augmented with additional controls capturing the generosity of welfare and access to family planning services, but measured in the *child’s* year of birth rather than the mother’s year of birth (shown in the solid blue squares).⁴⁸

A remaining concern with event study estimates is that they may be biased in the presence of heterogeneous treatment effects across expansion timing groups (Sun and Abraham, 2020). To alleviate this concern, we implement a new estimation method proposed by Callaway and Sant’Anna (2020) that is robust to heterogeneous treatment effects across groups. Essentially, this approach avoids using earlier treated units as controls for later treated units (which can lead to biased estimates). The results are shown in Appendix Figure A14.⁴⁹ Across all outcomes, the results are very similar to our baseline estimates.⁵⁰ This reassures us that the baseline event study results are robust to possible treatment effect heterogeneity.

As an alternative to our main event study model, we also evaluate the impact of the eligibility expansions using a synthetic control approach.⁵¹ This method constructs a weighted average of untreated states to serve as a synthetic control for each treated state, with weights selected to minimize the pre-treatment differences in the outcome and control variables across the treated state and synthetic control units, as described in Abadie et al. (2010). We select weights following the Abadie et al. (2010) method to obtain a synthetic control unit for each treated state, then stack the observations for each treated state and its synthetic control, and estimate an event study model, following the methods described in Kleven (2021). The results of this exercise are reported in Appendix Figure A15. While in some cases the synthetic control approach produces smaller point estimates, the overall patterns remain very similar.

⁴⁷We cannot rule out that changes in prenatal eligibility affected childhood *take-up* of Medicaid, however.

⁴⁸See the discussion in Appendix Sections A and B for detailed information about how we construct these other eligibility variables. Additional details on the other control variables are found in Appendix Section D.

⁴⁹We implement the Callaway and Sant’Anna method using R code they provided. We use the same weighting as in our main analysis (by size of birth cohort) and cluster by state. To simplify comparisons across estimators, we do not include control variables in this model. Note that our application of this method continues to use the control states as a comparison group and that some of these states may have experienced similar expansions in eligibility (i.e. “treatment”) prior to our study period.

⁵⁰Specifically, with this method, we find that four years after the expansion average birthweight had increased by 4.2 grams, low birthweight had decreased by 0.09 percentage points, very low birthweight decreased by 0.08 percentage points, very preterm decreased by 0.05 percentage points, and small-for-gestation-age decreased by 0.14 percentage points.

⁵¹This method offers an alternative way to account for pre-trends; when using this method we do not de-trend the outcome variables and instead use the matching procedure to find control states that match the treated states’ pre-trends. Appendix Section D.4 contains further details on our implementation of this method.

Next, we verify that our results are not sensitive to our sample selection criteria. Appendix Figure A16 shows estimates based on a second generation sample of higher parity births. This sample generates similar birthweight and low birthweight estimates as the main sample, but we no longer observe improvements in the incidence of very preterm or small for gestational-age. Tabulations we ran using pre-period data indicate that these outcomes are less common among higher order births, which may explain the difference. Appendix Figure A17 shows what happens when we relax our baseline maternal age restriction from ages 15-28 to include births to older mothers (recall that this produces a sample that is unbalanced in maternal age). Again, the estimates are very similar to those produced by our main sample.

Finally, we repeat our analysis including only children of foreign-born mothers, who are excluded in the main analysis, and who we assume were unaffected by the first generation prenatal expansions, since the first generation was not born in the U.S. This placebo check is another way of addressing concerns about unexplained cohort by state specific changes in infant health that are correlated with, but separate from, exposure to the prenatal expansions. We use the state of residence at the time of the child’s birth to assign policy and control variables. The results from this analysis are found in Appendix Figure A18.⁵² As expected, we do not detect any changes in infant health that correspond with the expansions’ timing.

6.4.1 Magnitudes

Our results indicate that the benefits of prenatal Medicaid spill over onto later offspring. Taken at face value, the point estimate for average birthweight suggests that the 6.6 percentage point increase in *in utero* eligibility generated by the 1980s expansions increased the second generation’s average birthweight by a statistically significant 4.7 grams, which corresponds to an increase per newly eligible woman of about 71 grams (2 percent relative to the mean).⁵³ This is a bit less than half of the observed gap in average birthweight between mothers with vs. without a high school degree at the beginning of our sample period. Comparable estimates for low birthweight and very low birthweight are 1.8 percentage points and 1.2 percentage points. Of course, the confidence intervals around these estimates include both much smaller and much larger effect sizes.

Not everyone in the first generation who became eligible for Medicaid under the expansions actually received coverage. We therefore interpret the estimated effects per newly eligible woman as “intent to treat” estimates, where treatment is defined as enrollment in the program. Assuming that the benefits of Medicaid eligibility accrue only to those who were actually enrolled *in utero*, and that there were no effects on those who were eligible but did not enroll, we can obtain the treatment effect of Medicaid enrollment by dividing the estimated expansion effects by the estimated increase in Medicaid coverage

⁵²We note that the sample of births to foreign-born women is much smaller than the sample of births to U.S.-born women. For example, in 1980 there are an average of 19,000 births per state to US-born women, compared to 4,500 births to foreign-born women. Not surprisingly, the estimates for the sample of births to foreign-born women are accompanied by larger confident intervals.

⁵³Note that this calculation relies on the assumption that there are no spillover effects of the expansions onto women who did not gain eligibility. This assumption may be violated if, for example, those who gained prenatal eligibility shared health information with friends or neighbors who did not gain eligibility, or if the expansions induced changes in providers’ behavior or policies that affected communities more broadly. If such spillovers exist, these treatment effect calculations may be biased upwards.

in Table 1 (4.6 percentage points in the fourth year of expansion). Using the point estimates above, this calculation implies that among the offspring of women whose pregnant mothers enrolled in Medicaid, average birthweight increased by a little over 100 grams.⁵⁴ Similarly, the treatment effect estimates for low birthweight and very low birthweight are 2.6 and 1.8 percentage points, respectively. While some of these treatment effect estimates are large in comparison to our baseline means, it is important to recall that the early expansions targeted very poor pregnant women for whom the incidence of poor birth outcomes was substantially higher than in the full population.⁵⁵

We can also compare our estimates of the expansions' effects across generations. We note that the first generation receives a direct and clearly defined treatment, which is access to Medicaid coverage during the *in utero* period. In contrast, the second generation's treatment is the bundle of biological and economic outcomes experienced by the mother as a result of her access to Medicaid, and subsequently passed on to her infant. It is therefore not clear *ex ante* which generation should experience larger effects, or whether effects must be present at birth in the first generation in order for the second generation to be plausibly affected. Comparing the point estimates in columns 4 and 6 in Table 1 we see that the decline in low birthweight experienced by the second generation is about 48% as large as the effect in the first generation. This is consistent with Currie and Moretti (2007), who find that the probability of being low birthweight is nearly 50 percent higher among children whose mothers were themselves born below the low birthweight threshold.⁵⁶

Finally, keeping in mind that there are wide confidence intervals around the estimates, we consider possible implications for the program's return on investment. Currie and Gruber (1996b) report that the targeted expansions increased Medicaid spending per eligible woman by \$450 (this cost is estimated in 1981 and we inflate to 2011 dollars). We compare this with the medical cost associated with lower birthweight averted in the second generation. Based on estimates provided in Almond et al. (2005), we calculate that each additional gram of birthweight reduces hospital costs by \$8.29.⁵⁷ Our estimates suggest that first generation eligibility increases average second generation birthweight by 71 grams, resulting in about \$589 in savings per first generation woman made eligible. Given the amount of time that has passed, we discount these cost savings back to the period of the initial Medicaid outlays (1981), using the discount rate recommended by the Department of Commerce for life-cycle studies (3%, see Lavappa and Kneifel, 2016). For this calculation, we assume cost savings accrue equally across all second generation birth years.⁵⁸ This calculation suggests that the expansions generated average cost savings of \$287 per newly eligible woman, or more than 60% of the cost of the initial investment.⁵⁹ Even

⁵⁴We obtain this estimate by dividing 4.8 grams by the 4.6 percentage point increase in Medicaid coverage.

⁵⁵As noted earlier, our calculations indicate that the incidence of low birthweight among children living in the lowest income households in the 1981 NHIS was more than twice that of the full population.

⁵⁶Other studies have estimated smaller intergenerational birthweight correlations (Black et al., 2007; Royer, 2009) but importantly, Currie and Moretti (2007) find that poverty increases the transmission of low birthweight from mother to child.

⁵⁷Almond et al. (2005) estimate that an additional gram of birthweight reduces hospital charges by \$22 for infants in the 2000-2100 gram range in 2000 dollars, or \$28.60 in 2011 dollars. Since hospital charges do not accurately reflect hospital resource costs (due to markups), we deflate this estimate using national charge-to-cost ratios (Bai and Anderson, 2015) to arrive at \$8.29 per gram.

⁵⁸We estimate cost savings for each second generation birth year from 1995 to 2017 since the first expansions occurred in 1980 and first generation mothers born in that year reached age 15 in 1995.

⁵⁹Using the discount rate recommended by the Office of Management and Budget of 0.5% instead, the discounted value of the benefits is \$520 (U.S. Office of Management and Budget, 2016). Details of these calculations are in the Appendix D.5.

if we use a more conservative birthweight estimate from the bottom of the 95% confidence interval, we calculate an average cost savings of more than 16% of the cost of the initial investment. Notably, this calculation is focused solely on benefits that translate to the second generation, and ignores any improvements in the first generation’s health and human capital. The calculation also ignores medical cost savings that result from any second generation health improvements in later life that are tied to better health at birth, as well as improvements in later life earnings (and tax revenues), which have also been tied to birthweight (e.g. [Black et al., 2007](#); [Bharadwaj et al., 2018](#)).

It may also be reasonable to expect that health benefits accruing to the first two generations will continue to be passed on in the future, albeit incompletely. We could therefore model the intergenerational benefits of the Medicaid program as an annuity that pays off once each generation. We assume that half of the benefit to the mother’s generation is transmitted to the child (as we observe for low birthweight), and then only half of that received benefit to the child’s offspring, and so forth, such that the benefit of an investment decays exponentially across generations. Applying a 3% discount rate, we calculate that accounting for the intergenerational aspects of the program’s effects in this way results in estimated benefits that are more than 30% higher than what would be observed if the analysis focused only on the first generation.⁶⁰

6.4.2 Additional Outcomes

We extend our analysis of the intergenerational effects of the Medicaid expansions by looking at additional health indicators, including the presence of congenital anomalies, whether the birth was associated with any abnormal conditions, and the sex ratio at birth.⁶¹ Congenital anomalies are more common among low birthweight and preterm infants, and have been linked to environmental factors that are often associated with low income, including poorer access to nutritious foods, health care and screening ([World Health Organization, 2012](#)). Similarly, some abnormal conditions such as NICU admission or surfactant replacement therapy could be indicative of underlying health conditions. Our examination of the sex ratio is motivated by the [Trivers and Willard \(1973\)](#) hypothesis, which suggests that, because male fetuses are more sensitive to negative health environments than female fetuses, improvements in maternal health might disproportionately reduce the number of *in utero* losses that are male. The results of these analyses are provided in Appendix Figure [A19](#) and lend further support to the hypothesis that the expansions’ health benefits extended beyond the treated generation. Although there is no evidence that the expansions affected abnormal conditions at birth, the patterns of estimates for congenital anomalies and the sex ratio are similar to our main results.

⁶⁰Specifically, we assume the Medicaid investment generates a payoff that falls by 50% in value and is paid out every 25 years. The present value of such a payoff of size X would be:

$$\$X \times \sum_{n=0}^{\infty} \left(\frac{0.5}{(1+r)^{25}} \right)^n.$$

Since $\frac{0.5}{(1+r)^{25}} < 1$ we can apply the rules for geometric series. Plugging in $r = 0.03$ for a 3% discount rate, this simplifies to $\$X \times 1.31$.

⁶¹Examples of abnormal conditions include the provision of assisted ventilation, NICU admission, surfactant replacement therapy, antibiotics, seizures, or significant birth injury.

6.5 Mechanisms

6.5.1 Changes in Fertility or Maternal Characteristics

What are the mechanisms generating these intergenerational spillovers? We investigate potential pathways using additional information provided in the natality files. First, we consider changes in the first generation’s fertility. The same (or related) biological processes that lead to improvements in the first generation’s health may have also affected the first generation’s fecundity.⁶² We also consider whether the Medicaid expansions led to changes in the composition of women giving birth: if children are a normal good, then Medicaid induced increases in the first generation’s earnings might also lead to increases in the desired number of children. On the other hand, improved economic opportunities might also lead to delays in childbearing (Brown et al., 2020).⁶³

We explore these potential mechanisms in Figure 10. We estimate regressions similar to equation (2), replacing the dependent variable with measures of fertility or maternal characteristics (age, educational attainment, marital status and race).⁶⁴ We find no evidence that the Medicaid expansions led to changes in fertility behavior; neither the likelihood of having a first birth, nor the age at first birth, are affected by the expansions. In addition to their statistical insignificance, the point estimates are very small (four years after the expansions, there is a 0.3% decrease in the first birth rate and a 0.1% increase in age at first birth, relative to the baseline means).

We do see some evidence of changes in the characteristics of mothers, particularly with respect to race. The expansions are associated with an increase in the percent of births to white mothers (0.7% relative to the pre-treatment mean) and a decrease in the percent of births to Black mothers (3% relative to the pre-treatment mean). Since white infants tend to be healthier on average than Black infants, we consider whether the shift towards white births explains the expansions’ apparent effects on the second generation’s health. To do this, we use the estimated effects four years after the primary expansion. Table 2 shows that by year four the percent of births to white mothers had increased by 0.54 percentage points, with a roughly similar decrease in the percent of births to Black mothers.⁶⁵ A back-of-the-envelope calculation based on this compositional shift suggests that selection can explain, at most, about 30 percent of the overall effect of the Medicaid expansion on average birthweight.⁶⁶ In

⁶²We are not able to measure total fertility because we cannot observe most of our first generation cohorts throughout their childbearing years.

⁶³Brown et al. (2020) estimate that each additional year of Medicaid eligibility from birth to age 18 is associated with a 1.2 percentage point reduction in the probability that a woman has her first child by age 28, but the authors do not examine the effect of *in utero* eligibility on fertility.

⁶⁴Three of the outcomes analyzed in this section and the next (Section 6.5.2) – mother’s educational attainment, prenatal care utilization, and race – were affected by the introduction of the 2003 revision of the U.S. Standard Certificate of Live Birth. This revised version replaced the 1989 version that was in use during the remainder of the period covered by our analyses. See Appendix Section D for more details on how we account for this change in our analyses.

⁶⁵We appreciate that by year four the estimated effects on racial composition are no longer statistically significant, but we use these estimates, which are very similar to the estimates in year 3, so as to be consistent with the focal event year used in the rest of the paper.

⁶⁶If we apply these estimates to the race specific averages for birthweight (3329.6 for children of white mothers and 3080.5 grams for children of Black mothers), we predict an increase in average birthweight of 1.24 grams due solely to the expansion’s effect on the racial composition of births (i.e. $3329.6 \cdot 0.005 - 3080.5 \cdot 0.005$). This estimate is 26% of the program’s estimated increase in the second generation’s average birthweight of 4.7 grams in year 4 of the expansion. We also explore whether the observed changes in infant health persist when we restrict the sample of second generation births to white mothers (78% of the sample). The results, shown in Appendix Figure A20, are very similar to those produced by the full baseline sample, indicating that selection based solely on race cannot explain all of our findings. Unfortunately,

addition, we re-estimate our main model directly controlling for mother’s education, marital status, and race. The results, shown in Appendix Figure A21, are very similar to our baseline estimates. This is a strong indication that the observed health improvements are not driven by changes in the composition of mothers giving birth.⁶⁷

6.5.2 Maternal Health and Behaviors

Finally, we consider the role of maternal health and maternal health behaviors. The results of these analyses should be interpreted cautiously, as health conditions reported on birth certificates are relatively limited and known to be under-reported (Lain et al., 2012). Nevertheless, Appendix Figure A22 hints at the presence of biological pathways. We see suggestive evidence that the prenatal expansions reduced the incidence of medical risk factors, which include diabetes, chronic hypertension, pregnancy-related hypertension, and eclampsia, among first generation mothers. Consistent with first generation studies that document a positive link between early-life Medicaid and later-life metabolic health (e.g. Boudreaux et al., 2016; Thompson, 2017; Miller and Wherry, 2019), the decline in maternal medical risk is driven by a reduction in pregnancy-related hypertension. As described in Section 2, an improvement in this outcome would be expected to reduce the likelihood of giving birth to a low birthweight or preterm infant.⁶⁸

7 Conclusion

Social scientists have long known that children’s outcomes are predicted by their parents’ socioeconomic status (e.g. Mazumder, 2005; Solon, 1992), and recent research suggests that health plays an important role in the intergenerational transmission process (Halliday et al., 2020, 2021). The extent to which the impacts of policy-driven health investments spill over onto later generations has been largely unexplored, however. Understanding such effects is important, both because it sheds light on how economic and health status are conferred across generations, and because it provides insights on the extent to which generationally persistent disadvantage can be ameliorated by policy choices. Furthermore, if safety net programs’ benefits extend beyond treated generations, then those benefits need to be accounted for in order to accurately assess both net-costs to the government and their total value to recipients (direct and indirect). Neglecting policies’ multi-generational effects could result in dramatic undervaluation.

Any exercise that seeks to quantify a policy’s multigenerational effects will inherently face substantial uncertainty: multigenerational outcomes are measured generations away from the initial treatment, and the individuals who were directly affected by a policy early in life have been subject to a multitude of economic, social, educational and cultural exposures before having children themselves. Furthermore, the factors that generate maternal health accrue over a lifetime, and the decision whether and

small sample sizes impede further subgroup analyses.

⁶⁷It is also possible that marital matching changed. Unfortunately, there is limited information about fathers’ characteristics provided on birth certificates (age, race, and Hispanic origin), which makes a direct examination of this hypothesis infeasible.

⁶⁸Surprisingly, we see some evidence that Medicaid eligibility is associated with reductions in the use of prenatal care during pregnancy. If anything, this would lead us to expect worse infant health outcomes, and suggests that the effects would be larger in the absence of this association. We do not examine changes in alcohol or tobacco use due to limited availability on the birth certificate records for all states and years. Alcohol use is only available through 2006. Information on tobacco use is available through 2008. This would only allow us to observe women through age 18 and age 20, respectively, if we were to balance the sample on mothers’ age at birth.

when to have children is multifaceted and varied. Given this, disentangling the exact mechanism(s) is practically very difficult. Nevertheless, such explorations are critical to the accurate assessment of interventions' benefits, and our results are an important contribution to this effort.

In this paper, we provide new evidence that expanding health related services to low-income pregnant women has persistent impacts on later generations' health. We introduce a new event study approach to analyze the effects of the 1980s Medicaid expansions that exploits large, discrete jumps in state eligibility resulting from a combination of state policy choices and federal mandates. Using this new approach, we confirm previous studies' findings that the targeted expansions generated improvements in the first generation's birth outcomes. Then we document that this treated generation went on to give birth to healthier offspring. Despite the large confidence intervals accompanying some of our estimates, they all point to a consistent pattern of improvement in second generation health. Moreover, these results do not appear to be driven by changes in the first generation's fertility behavior or selection into childbearing. Instead, the positive spillovers likely reflect improvements in the first generation's health and economic outcomes prior to giving birth. In particular, our analyses suggest that Medicaid-induced health gains among first generation mothers may be an important underlying mechanism.

Our study offers a new perspective on health inequalities and the potential role for government intervention. Generational persistence in the impacts of early life environments suggest that historical differences in fetal health conditions between advantaged and disadvantaged groups may undermine contemporaneous efforts to close health and economic gaps. At the same time, our results indicate that early life health investments have payoffs that extend well beyond those that social policymakers usually consider. It is notable that Medicaid's second generation effects are observed among cohorts who were born during roughly the same time frame for which recent studies by [Aizer and Currie \(2014\)](#) and [Currie and Schwandt \(2016a,b\)](#) document large improvements in child health and declining health inequality. Investigating a more complete range of program benefits to later generations is an important goal of future work, and is critical in light of increasing debates about the efficacy of the U.S. safety net.

References

- Abadie, A., A. Diamond, and J. Hainmueller (2010, June). Synthetic Control Methods for Comparative Case Studies: Estimating the Effect of California’s Tobacco Control Program. *Journal of the American Statistical Association* 105(490), 493–505.
- Aerts, L. and F. A. Van Assche (2006, May). Animal evidence for the transgenerational development of diabetes mellitus. *The International Journal of Biochemistry & Cell Biology* 38(5), 894–903.
- Aizer, A. and J. Currie (2014, May). The intergenerational transmission of inequality: Maternal disadvantage and health at birth. *Science* 344(6186), 856–861.
- Aizer, A., L. Stroud, and S. Buka (2016, August). Maternal Stress and Child Outcomes: Evidence from Siblings. *Journal of Human Resources* 51(3), 523–555.
- Almond, D. (2006). Is the 1918 Influenza Pandemic Over? Long-Term Effects of In Utero Influenza Exposure in the Post-1940 U.S. Population. *Journal of Political Economy* 114(4), 672–712.
- Almond, D. and K. Y. Chay (2006, February). The Long-Run and Intergenerational Impact of Poor Infant Health: Evidence from Cohorts Born During the Civil Rights Era. Working Paper.
- Almond, D., K. Y. Chay, and D. S. Lee (2005, August). The Costs of Low Birth Weight. *The Quarterly Journal of Economics* 120(3), 1031–1083.
- Almond, D. and J. Currie (2011, September). Killing Me Softly: The Fetal Origins Hypothesis. *Journal of Economic Perspectives* 25(3), 153–172.
- Almond, D., J. Currie, and V. Duque (2018, December). Childhood Circumstances and Adult Outcomes: Act II. *Journal of Economic Literature* 56(4), 1360–1446.
- Almond, D., J. Currie, and M. Herrmann (2012, August). From infant to mother: Early disease environment and future maternal health. *Labour Economics* 19(4), 475–483.
- Almond, D., L. Edlund, H. Li, and J. Zhang (2010). Long-Term Effects of Early-Life Development: Evidence from the 1959 to 1961 China Famine. In *The Economic Consequences of Demographic Change in East Asia*, pp. 321–345. Chicago: University of Chicago Press.
- Almond, D., L. Edlund, and R. Palme (2009, November). Chernobyl’s Subclinical Legacy: Prenatal Exposure to Radioactive Fallout and School Outcomes in Sweden. *Quarterly Journal of Economics* 124(4), 1729–1772.
- Almond, D., H. W. Hoynes, and D. W. Schanzenbach (2011). Inside the War on Poverty: The Impact of Food Stamps on Birth Outcomes. *The Review of Economics and Statistics* 93(2), 387–403. Publisher: The MIT Press.
- Almond, D. and B. Mazumder (2005, May). The 1918 Influenza Pandemic and Subsequent Health Outcomes: An Analysis of SIPP Data. *American Economic Review* 95(2), 258–262.

- Almond, D. and B. Mazumder (2011, October). Health Capital and the Prenatal Environment: The Effect of Ramadan Observance during Pregnancy. *American Economic Journal: Applied Economics* 3(4), 56–85.
- Bai, G. and G. F. Anderson (2015, June). Extreme Markup: The Fifty US Hospitals With The Highest Charge-To-Cost Ratios. *Health Affairs* 34(6), 922–928.
- Bailey, M., H. Hoynes, M. Rossin-Slater, and R. Walker (2020, April). Is the Social Safety Net a Long-Term Investment? Large-Scale Evidence from the Food Stamps Program. Working Paper w26942, National Bureau of Economic Research, Cambridge, MA.
- Barker, D. J. (1990, November). The fetal and infant origins of adult disease. *BMJ : British Medical Journal* 301(6761), 1111.
- Barr, A. and C. R. Gibbs (2019, October). Breaking the Cycle? Intergenerational Effects of an Anti-Poverty Program in Early Childhood. EdWorking Paper No. 19-141, Brown University.
- Barreca, A. I. (2010). The Long-Term Economic Impact of In Utero and Postnatal Exposure to Malaria. *The Journal of Human Resources* 45(4), 865–892.
- Bhalotra, S. R. and A. Venkataramani (2015, August). Shadows of the Captain of the Men of Death: Early Life Health Interventions, Human Capital Investments, and Institutions. SSRN Scholarly Paper ID 1940725, Social Science Research Network, Rochester, NY.
- Bharadwaj, P., K. V. Løken, and C. Neilson (2013, August). Early Life Health Interventions and Academic Achievement. *American Economic Review* 103(5), 1862–1891.
- Bharadwaj, P., P. Lundborg, and D.-O. Rooth (2018, January). Birth Weight in the Long Run. *Journal of Human Resources* 53(1), 189–231.
- Bitler, M. P. and T. F. Figinski (2019). Long-Run Effects of Food Assistance: Evidence from the Food Stamp Program. Working Paper, Economic Self-Sufficiency Policy Research Institute.
- Black, S. E., A. Bütikofer, P. J. Devereux, and K. G. Salvanes (2019, July). This Is Only a Test? Long-Run and Intergenerational Impacts of Prenatal Exposure to Radioactive Fallout. *The Review of Economics and Statistics* 101(3), 531–546.
- Black, S. E., P. J. Devereux, and K. G. Salvanes (2007, February). From the Cradle to the Labor Market? The Effect of Birth Weight on Adult Outcomes. *The Quarterly Journal of Economics* 122(1), 409–439.
- Black, S. E., P. J. Devereux, and K. G. Salvanes (2016, January). Does Grief Transfer across Generations? Bereavements during Pregnancy and Child Outcomes. *American Economic Journal: Applied Economics* 8(1), 193–223.
- Boudreaux, M. H., E. Golberstein, and D. D. McAlpine (2016, January). The Long-Term Impacts of Medicaid Exposure in Early Childhood: Evidence from the Program’s Origin. *Journal of Health Economics* 45, 161–175.

- Brown, D. W., A. E. Kowalski, and I. Z. Lurie (2020, March). Long-Term Impacts of Childhood Medicaid Expansions on Outcomes in Adulthood. *The Review of Economic Studies* 87(2), 792–821.
- Bütikofer, A., K. Loken, and K. Salvanes (2019, May). Infant Health Care and Long-Term Outcomes. *Review of Economics and Statistics* 101(2), 341–354.
- Bütikofer, A. and K. G. Salvanes (2020, October). Disease Control and Inequality Reduction: Evidence from a Tuberculosis Testing and Vaccination Campaign. *The Review of Economic Studies* 87(5), 2087–2125.
- Callaway, B. and P. H. Sant’Anna (2020, December). Difference-in-Differences with multiple time periods. *Journal of Econometrics* (Forthcoming).
- Camacho, A. (2008). Stress and Birth Weight: Evidence from Terrorist Attacks. *The American Economic Review* 98(2), 511–515.
- Carneiro, P. and R. Ginja (2014). Long-Term Impacts of Compensatory Preschool on Health and Behavior: Evidence from Head Start. *American Economic Journal: Economic Policy* 6(4), 135–173.
- Case, A., A. Fertig, and C. Paxson (2005). The Lasting Impact of Childhood Health and Circumstance. *Journal of Health Economics* 24, 365–389.
- Case, A., D. Lubotsky, and C. Paxson (2002, December). Economic Status and Health in Childhood: The Origins of the Gradient. *American Economic Review* 92(5), 1308–1334.
- Catalano, P. M. and H. M. Ehrenberg (2006). Review article: The short- and long-term implications of maternal obesity on the mother and her offspring. *BJOG: An International Journal of Obstetrics & Gynaecology* 113(10), 1126–1133.
- Centers for Disease Control and Prevention (2011). User Guide to the 2011 Natality Public Use File. Technical report.
- Chen, Y. and L.-A. Zhou (2007, July). The long-term health and economic consequences of the 1959–1961 famine in China. *Journal of Health Economics* 26(4), 659–681.
- Clark, G. (2014). *The Son Also Rises: Surnames and the History of Social Mobility*. Princeton University Press.
- Cohodes, S. R., D. S. Grossman, S. A. Kleiner, and M. F. Lovenheim (2016, August). The Effect of Child Health Insurance Access on Schooling: Evidence from Public Insurance Expansions. *Journal of Human Resources* 51(3), 727–759.
- Colmer, J. and J. Voorheis (2020, November). The Grandkids Aren’t Alright: The Intergenerational Effects of Prenatal Pollution Exposure. Working Paper.
- Congressional Research Service (1988). *Medicaid Source Book: Background Data and Analysis: A Report*. Washington D.C.: U.S. Government Printing Office.

- Conti, G., J. J. Heckman, and R. Pinto (2016). The Effects of Two Influential Early Childhood Interventions on Health and Healthy Behaviour. *The Economic Journal* 126(596), F28–F65.
- Cowley, J. and R. Griesel (1966). The Effect on Growth and Behavior of Rehabilitating First and Second Generation Low Protein Rats. *Animal Behavior* 14, 506–517.
- Currie, J. and D. Almond (2011, January). Human Capital Development Before Age Five. In D. Card and O. Ashenfelter (Eds.), *Handbook of Labor Economics*, Volume 4, pp. 1315–1486. Elsevier.
- Currie, J., S. Decker, and W. Lin (2008). Has Public Health Insurance for Older Children Reduced Disparities in Access to Care and Health Outcomes? *Journal of Health Economics* 27, 1567–1581.
- Currie, J. and J. Gruber (1994, February). Saving Babies: The Efficacy and Cost of Recent Expansions of Medicaid Eligibility for Pregnant Women. Working Paper No. w4644, National Bureau of Economic Research.
- Currie, J. and J. Gruber (1996a, May). Health Insurance Eligibility, Utilization of Medical Care, and Child Health. *The Quarterly Journal of Economics* 111(2), 431–466.
- Currie, J. and J. Gruber (1996b). Saving Babies: The Efficacy and Cost of Recent Changes in the Medicaid Eligibility of Pregnant Women. *Journal of Political Economy* 104(6), 1263–1296.
- Currie, J. and J. Gruber (2001, October). Public health insurance and medical treatment: the equalizing impact of the Medicaid expansions. *Journal of Public Economics* 82(1), 63–89.
- Currie, J. and E. Moretti (2007, April). Biology as Destiny? Short- and Long-Run Determinants of Intergenerational Transmission of Birth Weight. *Journal of Labor Economics* 25(2), 231–264.
- Currie, J. and H. Schwandt (2016a, May). Inequality in mortality decreased among the young while increasing for older adults, 1990–2010. *Science* 352(6286), 708–712.
- Currie, J. and H. Schwandt (2016b, May). Mortality Inequality: The Good News from a County-Level Approach. *Journal of Economic Perspectives* 30(2), 29–52.
- Dave, D., S. Decker, R. Kaestner, and K. Simon (2008, December). Re-examining the Effects of Medicaid Expansions for Pregnant Women. Working Paper No. w14591, National Bureau of Economic Research, Cambridge, MA.
- Dave, D. M., S. L. Decker, R. Kaestner, and K. I. Simon (2010, November). The Effect of Medicaid Expansions on the Health Insurance Coverage of Pregnant Women: An Analysis Using Deliveries. *INQUIRY: The Journal of Health Care Organization, Provision, and Financing* 47(4), 315–330.
- Davern, M., J. A. Klerman, D. K. Baugh, K. T. Call, and G. D. Greenberg (2009, June). An Examination of the Medicaid Undercount in the Current Population Survey: Preliminary Results from Record Linking. *Health Services Research* 44(3), 965–987.
- Daxinger, L. and E. Whitelaw (2010, December). Transgenerational Epigenetic Inheritance: More Questions than Answers. *Genome Research* 20(12), 1623–1628.

- Daxinger, L. and E. Whitelaw (2012, March). Understanding Transgenerational Epigenetic Inheritance via the Gametes in Mammals. *Nature Reviews Genetics* 13(3), 153–162.
- Deming, D. (2009, July). Early Childhood Intervention and Life-Cycle Skill Development: Evidence from Head Start. *American Economic Journal: Applied Economics* 1(3), 111–134.
- Dubay, L., T. Joyce, R. Kaestner, and G. M. Kenney (2001, June). Changes in Prenatal Care Timing and Low Birth Weight by Race and Socioeconomic Status: Implications for the Medicaid Expansions for Pregnant Women. *Health Services Research* 36(2), 373–398.
- Dunn, G. A. and T. L. Bale (2009, November). Maternal High-Fat Diet Promotes Body Length Increases and Insulin Insensitivity in Second-Generation Mice. *Endocrinology* 150(11), 4999–5009.
- East, C. N. (2020). The Effect of Food Stamps on Children’s Health: Evidence from Immigrants’ Changing Eligibility. *Journal of Human Resources* 55(2), 387–427. Publisher: University of Wisconsin Press.
- Ellwood, M. R. and G. Kenney (1995). Medicaid and Pregnant Women: Who is Being Enrolled and When. *Health Care Financing Review* 17(2), 7–28.
- Evans, W. N. and C. L. Garthwaite (2014, May). Giving Mom a Break: The Impact of Higher EITC Payments on Maternal Health. *American Economic Journal: Economic Policy* 6(2), 258–290.
- Figlio, D., J. Guryan, K. Karbownik, and J. Roth (2014, December). The Effects of Poor Neonatal Health on Children’s Cognitive Development. *American Economic Review* 104(12), 3921–3955.
- Finkelstein, A., S. Taubman, B. Wright, M. Bernstein, J. Gruber, J. P. Newhouse, H. Allen, and K. Baicker (2012, August). The Oregon Health Insurance Experiment: Evidence from the First Year. *The Quarterly Journal of Economics* 127(3), 1057–1106.
- Fitzsimmons, E. and M. Vera-Hernández (2013, December). Food for Thought? Breastfeeding and Child Development. Working Paper, Institute for Fiscal Studies.
- Forsdahl, A. (1977, June). Are Poor Living Conditions in Childhood and Adolescence an Important Risk Factor for Arteriosclerotic Heart Disease? *British Journal of Preventive & Social Medicine* 31(2), 91–95.
- Francis, D., J. Diorio, D. Liu, and M. J. Meaney (1999, November). Nongenomic Transmission Across Generations of Maternal Behavior and Stress Responses in the Rat. *Science* 286(5442), 1155–1158.
- Garces, E., D. Thomas, and J. Currie (2002, September). Longer-Term Effects of Head Start. *American Economic Review* 92(4), 999–1012.
- Glied, S. and M. Neidell (2010, March). The Economic Value of Teeth. *Journal of Human Resources* 45(2), 468–496.
- Goodman-Bacon, A. (2020, August). Difference-in-Differences with Variation in Treatment Timing. *Journal of Econometrics* (Forthcoming).

- Goodman-Bacon, A. (2021). The Long-Run Effects of Childhood Insurance Coverage: Medicaid Implementation, Adult Health, and Labor Market Outcomes. *American Economic Review* (Forthcoming).
- Grote, N. K., J. A. Bridge, A. R. Gavin, J. L. Melville, S. Iyengar, and W. J. Katon (2010, October). A Meta-analysis of Depression During Pregnancy and the Risk of Preterm Birth, Low Birth Weight, and Intrauterine Growth Restriction. *Archives of General Psychiatry* 67(10), 1012.
- Grundwald, N. J. and P. J. Brunton (2015, December). Prenatal Stress Programs Neuroendocrine Stress Responses and Affective Behaviors in Second Generation Rats in a Sex-Dependent Manner. *Psychoneuroendocrinology* 62, 204–216.
- Halliday, T., B. Mazumder, and A. Wong (2021, January). Intergenerational Mobility in Self-Reported Health Status in the US. *Journal of Public Economics* 193(104307), 1–20.
- Halliday, T. J., B. Mazumder, and A. Wong (2020, March). The Intergenerational Transmission of Health in the United States: A Latent Variables Analysis. *Health Economics* 29(3), 367–381.
- Heard, E. and R. A. Martienssen (2014, March). Transgenerational Epigenetic Inheritance: Myths and Mechanisms. *Cell* 157(1), 95–109.
- Hill, I. T. (1987, February). Broadening Medicaid Coverage of Pregnant Women and Children: State Policy Responses. Technical report, National Governors' Association, Washington, DC.
- Hochberg, Z., R. Feil, M. Constancia, M. Fraga, C. Junien, J.-C. Carel, P. Boileau, Y. Le Bouc, C. L. Deal, K. Lillycrop, R. Scharfmann, A. Sheppard, M. Skinner, M. Szyf, R. A. Waterland, D. J. Waxman, E. Whitelaw, K. Ong, and K. Albertsson-Wikland (2011, April). Child Health, Developmental Plasticity, and Epigenetic Programming. *Endocrine Reviews* 32(2), 159–224.
- Howell, E. M. (2001, March). The Impact of the Medicaid Expansions for Pregnant Women: A Synthesis of the Evidence. *Medical Care Research and Review* 58(1), 3–30.
- Hoynes, H., D. Miller, and D. Simon (2015, February). Income, the Earned Income Tax Credit, and Infant Health. *American Economic Journal: Economic Policy* 7(1), 172–211.
- Hoynes, H., D. W. Schanzenbach, and D. Almond (2016, April). Long-Run Impacts of Childhood Access to the Safety Net. *American Economic Review* 106(4), 903–934.
- Institute of Medicine (2017). *Preterm Birth: Causes, Consequences, and Prevention*. Washington, DC: The National Academies Press.
- Institute of Medicine (US) Committee on Understanding Premature Birth and Assuring Healthy Outcomes (2007). Medical and Pregnancy Conditions Associated with Preterm Birth. In R. E. Behrman and A. S. Butler (Eds.), *Preterm Birth: Causes, Consequences, and Prevention*. National Academies Press (US).
- Isen, A., M. Rossin-Slater, and W. R. Walker (2017, May). Every Breath You Take—Every Dollar You'll Make: The Long-Term Consequences of the Clean Air Act of 1970. *Journal of Political Economy* 125(3), 848–902.

- Jimenez-Chillaron, J. C., E. Isganaitis, M. Charalambous, S. Gesta, T. Pentinat-Pelegrin, R. R. Faucette, J. P. Otis, A. Chow, R. Diaz, A. Ferguson-Smith, and M.-E. Patti (2009, February). Intergenerational Transmission of Glucose Intolerance and Obesity by In Utero Undernutrition in Mice. *Diabetes* 58(2), 460–468.
- Kehrer, B. H. and C. M. Wolin (1979). Impact of Income Maintenance on Low Birth Weight: Evidence from the Gary Experiment. *The Journal of Human Resources* 14(4), 434.
- Klerman, J. A., J. S. Ringel, and E. Roth (2005, May). Under-Reporting of Medicaid and Welfare in the Current Population Survey. *RAND Working Paper* (WR-169-3). Publisher: RAND Corporation.
- Kleven, H. (2021, November). The EITC and the Extensive Margin: A Reappraisal. *NBER Working Paper Series* (No 26405), 201.
- Kramer, M. S. (1987a). Determinants of low birth weight: methodological assessment and meta-analysis. *Bulletin of the World Health Organization* 65(5), 663–737.
- Kramer, M. S. (1987b). Intrauterine Growth and Gestational Duration Determinants. *Pediatrics* 80(4), 502–511.
- Kuklina, E. V., M. K. Whiteman, S. D. Hillis, D. J. Jamieson, S. F. Meikle, S. F. Posner, and P. A. Marchbanks (2008, July). An Enhanced Method for Identifying Obstetric Deliveries: Implications for Estimating Maternal Morbidity. *Maternal and Child Health Journal* 12(4), 469–477.
- Lain, S. J., R. M. Hadfield, C. H. Raynes-Greenow, J. B. Ford, N. M. Mealing, C. S. Algert, and C. L. Roberts (2012, April). Quality of Data in Perinatal Population Health Databases: A Systematic Review. *Medical Care* 50(4), e7.
- Lavappa, P. D. and J. D. Kneifel (2016). Energy Price Indices and Discount Factors for Life-Cycle Cost Analysis – 2019 Annual Supplement to NIST Handbook 135. NIST Interagency/Internal Report 85-3273-32.
- Lee, W. and J. E. Roemer (2006, August). Racism and redistribution in the United States: A solution to the problem of American exceptionalism. *Journal of Public Economics* 90(6-7), 1027–1052.
- Levine, P. B. and D. Schanzenbach (2009). The Impact of Children’s Pubic Health Insurance Expansions on Educational Outcomes. *Forum for Health Economics and Policy* 12(1), 1–26.
- Lu, M. C., V. Tache, G. R. Alexander, M. Kotelchuck, and N. Halfon (2003, January). Preventing low birth weight: is prenatal care the answer? *Journal of Maternal-Fetal and Neonatal Medicine* 13(6), 362–380.
- Ludwig, J. and D. L. Miller (2007, February). Does Head Start Improve Children’s Life Chances? Evidence from a Regression Discontinuity Design. *The Quarterly Journal of Economics* 122(1), 159–208.
- Mansour, H. and D. I. Rees (2012, September). Armed Conflict and Birth Weight: Evidence from the al-Aqsa Intifada. *Journal of Development Economics* 99(1), 190–199.


- Maritz, G. S. and M. Mutemwa (2014). The Effect of Grand Maternal Nicotine Exposure During Gestation and Lactation on Lung Integrity of the F2 Generation. *Pediatric Pulmonology* 49(1), 67–75.
- Markus, A. R., E. Andres, K. D. West, N. Garro, and C. Pellegrini (2013, September). Medicaid Covered Births, 2008 Through 2010, in the Context of the Implementation of Health Reform. *Women’s Health Issues* 23(5), e273–e280.
- Martin, D. I., J. E. Cropley, and C. M. Suter (2008, August). Environmental Influence on Epigenetic Inheritance at the Avy Allele. *Nutrition Reviews* 66(suppl.1), S12–S14.
- Martínez, D., T. Pentinat, S. Ribó, C. Daviaud, V. W. Bloks, J. Cebrià, N. Villalmanzo, S. G. Kalko, M. Ramón-Krauel, R. Díaz, T. Plösch, J. Tost, and J. C. Jiménez-Chillarón (2014, June). In Utero Undernutrition in Male Mice Programs Liver Lipid Metabolism in the Second-Generation Offspring Involving Altered Lxra DNA Methylation. *Cell Metabolism* 19(6), 941–951.
- Masuyama, H., T. Mitsui, E. Nobumoto, and Y. Hiramatsu (2015, July). The Effects of High-Fat Diet Exposure In Utero on the Obesogenic and Diabetogenic Traits Through Epigenetic Changes in Adiponectin and Leptin Gene Expression for Multiple Generations in Female Mice. *Endocrinology* 156(7), 2482–2491.
- Matthews, S. G. and D. I. Phillips (2012, January). Transgenerational inheritance of stress pathology. *Experimental Neurology* 233(1), 95–101.
- Mazumder, B. (2005, May). Fortunate Sons: New Estimates of Intergenerational Mobility in the United States Using Social Security Earnings Data. *Review of Economics and Statistics* 87(2), 235–255.
- Mazumder, B., D. Almond, K. Park, E. M. Crimmins, and C. E. Finch (2010, February). Lingering Prenatal Effects of the 1918 Influenza Pandemic on Cardiovascular Disease. *Journal of Developmental Origins of Health and Disease* 1(1), 26–34.
- Meyer, B. D. and D. T. Rosenbaum (2001, August). Welfare, the Earned Income Tax Credit, and the Labor Supply of Single Mothers. *The Quarterly Journal of Economics* 116(3), 1063–1114.
- Miller, S. and L. Wherry (2016, August). The Long-Term Effects of Early Life Medicaid Coverage. SSRN Scholarly Paper ID 2466691, Social Science Research Network, Rochester, NY.
- Miller, S. and L. R. Wherry (2019). The Long-Term Effects of Early Life Medicaid Coverage. *Journal of Human Resources* 54(3), 785–824.
- Morgan, C. P. and T. L. Bale (2011, August). Early Prenatal Stress Epigenetically Programs Dysmasculinization in Second-Generation Offspring via the Paternal Lineage. *Journal of Neuroscience* 31(33), 11748–11755.
- Nadeau, J. H. (2009, October). Transgenerational Genetic Effects on Phenotypic Variation and Disease Risk. *Human Molecular Genetics* 18(R2), R202–R210.

- Neelsen, S. and T. Stratmann (2012, January). Long-run Effects of Fetal Influenza Exposure: Evidence from Switzerland. *Social Science & Medicine* 74(1), 58–66.
- Nilsson, J. P. (2009, January). The Long Term Effects of Early Childhood Lead Exposure: Evidence from the Phase-out of Leaded Gasoline. Working Paper.
- Painter, R. C., C. Osmond, P. Gluckman, M. Hanson, D. I. W. Phillips, and T. J. Roseboom (2008). Transgenerational Effects of Prenatal Exposure to the Dutch Famine on Neonatal Adiposity and Health in Later Life. *BJOG: An International Journal of Obstetrics & Gynaecology* 115(10), 1243–1249.
- Painter, R. C., T. J. Roseboom, and O. P. Bleker (2005, September). Prenatal Exposure to the Dutch Famine and Disease in Later Life: An Overview. *Reproductive Toxicology* 20(3), 345–352.
- Persson, P. and M. Rossin-Slater (2018, April). Family Ruptures, Stress, and the Mental Health of the Next Generation. *American Economic Review* 108(4-5), 1214–1252.
- Radford, E. J., M. Ito, H. Shi, J. A. Corish, K. Yamazawa, E. Isganaitis, S. Seisenberger, T. A. Hore, W. Reik, S. Erkek, A. H. F. M. Peters, M.-E. Patti, and A. C. Ferguson-Smith (2014, August). In Utero Undernourishment Perturbs the Adult Sperm Methyloome and Intergenerational Metabolism. *Science* 345(6198).
- Ravelli, G.-P., Z. A. Stein, and M. W. Susser (1976, August). Obesity in Young Men after Famine Exposure in Utero and Early Infancy. *New England Journal of Medicine* 295(7), 349–353.
- Rehan, V. K., J. Liu, E. Naeem, J. Tian, R. Sakurai, K. Kwong, O. Akbari, and J. S. Torday (2012, October). Perinatal Nicotine Exposure Induces Asthma in Second Generation Offspring. *BMC Medicine* 10(1), 129.
- Richter, A. and P. O. Robling (2016, September). Multigenerational Effects of the 1918-19 Influenza Pandemic on Educational Attainment: Evidence from Sweden. Working Paper.
- Roseboom, T. J., J. H. P. van der Meulen, A. C. J. Ravelli, C. Osmond, D. J. P. Barker, and O. P. Bleker (2001, December). Effects of Prenatal Exposure to the Dutch Famine on Adult Disease in Later Life: An Overview. *Molecular and Cellular Endocrinology* 185(1), 93–98.
- Rossin-Slater, M. and M. Wüst (2020, July). What Is the Added Value of Preschool for Poor Children? Long-Term and Intergenerational Impacts and Interactions with an Infant Health Intervention. *American Economic Journal: Applied Economics* 12(3), 255–286.
- Royer, H. (2009, January). Separated at Girth: US Twin Estimates of the Effects of Birth Weight. *American Economic Journal: Applied Economics* 1(1), 49–85.
- Sanders, N. J. (2012, July). What Doesn't Kill You Makes You Weaker Prenatal Pollution Exposure and Educational Outcomes. *Journal of Human Resources* 47(3), 826–850.
- Schmidheiny, K. and S. Siegloch (2019). On Event Study Designs and Distributed-Lag Models: Equivalence, Generalization and Practical Implications. *CESifo Working Paper no. 7481*.


- Scholte, R. S., G. J. van den Berg, and M. Lindeboom (2015, January). Long-run Effects of Gestation During the Dutch Hunger Winter Famine on Labor Market and Hospitalization Outcomes. *Journal of Health Economics* 39, 17–30.
- Schöpfer, H., R. Palme, T. Ruf, and S. Huber (2012, March). Effects of Prenatal Stress on Hypothalamic–pituitary–Adrenal (HPA) Axis Function Over Two Generations of Guinea Pigs (*Cavia aperea f. porcellus*). *General and Comparative Endocrinology* 176(1), 18–27.
- Sohn, H. (2017, March). Medicaid’s Lasting Impressions: Population Health and Insurance at Birth. *Social Science & Medicine* 177, 205–212.
- Solon, G. (1992). Intergenerational Income Mobility in the United States. *The American Economic Review* 82(3), 393–408.
- Solon, G. (2018). What Do We Know So Far about Multigenerational Mobility? *The Economic Journal* 128(612), F340–F352.
- Stein, Z., M. Susser, G. Saenger, and F. Marolla (1975). *Famine and Human Development: The Dutch Hunger Winter of 1944-1945*. New York: Oxford University Press.
- Sun, L. and S. Abraham (2020, August). Estimating Dynamic Treatment Effects in Event Studies With Heterogeneous Treatment Effects. *Journal of Econometrics* (Forthcoming).
- Susser, E. S. (1992). Schizophrenia After Prenatal Exposure to the Dutch Hunger Winter of 1944-1945. *Arch Gen Psychiatry* 49, 983–988.
- Thompson, O. (2017, January). The Long-Term Health Impacts of Medicaid and CHIP. *Journal of Health Economics* 51, 26–40.
- Trivers, R. L. and D. E. Willard (1973, January). Natural Selection of Parental Ability to Vary the Sex Ratio of Offspring. *Science* 179(4068), 90–92.
- United Health Foundation (2020). Low Birthweight in the United States, 1990-2018.
- U.S. Office of Management and Budget (2016, November). OMB Circular A-94 Appendix C. Technical report.
- Valente, C. (2011, July). Children of the Revolution: Fetal and Child Health amidst Violent Civil Conflict. Technical Report 11/12, HEDG, c/o Department of Economics, University of York.
- van den Berg, G. J. and P. R. Pinger (2016, December). Transgenerational Effects of Childhood Conditions on Third Generation Health and Education Outcomes. *Economics & Human Biology* 23, 103–120.
- van Ewijk, R. (2011, December). Long-term Health Effects on the Next Generation of Ramadan Fasting During Pregnancy. *Journal of Health Economics* 30(6), 1246–1260.
- Wang, S. S. (2010, January). Birth Weights Fell From 1990 to 2005. *Wall Street Journal*.

- Wherry, L. R. and B. D. Meyer (2016, August). Saving Teens: Using a Policy Discontinuity to Estimate the Effects of Medicaid Eligibility. *Journal of Human Resources* 51(3), 556–588.
- Wherry, L. R., S. Miller, R. Kaestner, and B. D. Meyer (2017, April). Childhood Medicaid Coverage and Later-Life Health Care Utilization. *The Review of Economics and Statistics* 100(2), 287–302.
- World Health Organization (2012, October). Congenital Anomalies. Fact Sheet No. 370.
- Zamenhof, S., E. v. Marthens, and L. Grauel (1971, May). DNA (Cell Number) in Neonatal Brain: Second Generation (F2) Alteration by Maternal (F0) Dietary Protein Restriction. *Science* 172(3985), 850–851.

Figure 1: Trends in Prenatal Medicaid Eligibility and Medicaid Coverage at Birth, 1975 to 1988




(a) Eligibility




(b) Coverage

Notes: Authors' calculation from the Current Population Survey and Medicaid eligibility rules and from the National Hospital Discharge Survey. For eligibility estimates, state averages are weighted using the number of births in each state-year cohort. For coverage estimates, sample weights are applied. All estimates are reported in percents. See text for further details.

Figure 3: Prenatal Eligibility by Source and Treatment Status, 1975-1988




(a) Treated States



(b) Control States


Notes: Authors' calculation from the Current Population Survey and Medicaid eligibility rules. Each line represents Medicaid eligibility through each of the state options and federal mandates. Specifically, we construct this figure by calculating the percent of women who would be eligible under each pathway in an additive fashion that reflects the order of eligibility pathways from lowest to highest pathway on the figure. These pathways are discussed in more detail in Appendix Section A. All estimates are reported in percents.

Figure 4: Trends in Simulated Prenatal Medicaid Eligibility Relative to Year Prior to Large Expansion




Notes: Authors' calculation from the Current Population Survey and Medicaid eligibility rules. Difference in simulated eligibility from the year prior to the large expansion in each group of treated states is depicted. The number of treated states in each group is listed in parenthesis. All estimates are reported in percents. See text for further details.

Figure 5: Event Study Coefficients for Prenatal Eligibility




(a) Actual Eligibility



(b) Simulated Eligibility


Notes: Coefficient estimates are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth.

Figure 6: Event Study for Medicaid Coverage Among Labor and Delivery Hospital Discharges




Notes: Coefficient estimates are reported in percentage points. Estimated for hospital discharges for labor and delivery between 1979-1988. Regressions are weighted by NHDS sample weights and include state of birth and year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by state.

Figure 7: Event Study for First-Generation Outcome: Low Birth Weight




Notes: Coefficient estimates are reported in percentage points. Estimated for infants born in 1975-1988. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size and include state of birth and year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by infant's state of birth.

Figure 8: Event Study for Second Generation Outcomes




Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth.

Figure 9: Event Study for Second Generation Outcomes, Robustness to Alternative Controls and Specifications



Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects. Unless indicated otherwise, regressions also include state-year control variables. Standard errors are clustered by mother's state of birth.

Figure 10: Event Study for Fertility Outcomes and Maternal Characteristics



Notes: Coefficient estimates for (a), (c), (d), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth. Additional controls for the share of births with revised birth certificate records are included for education and race outcomes, and the share of births with birth certificate records allowing for the report of multiple race categories are included for the race outcomes. Standard errors are clustered by mother's state of birth.

Table 1: Event Study Estimates on Eligibility, Coverage, and Health Outcomes

	<i>First Stage</i>			<i>First Gen.</i>			<i>Second Generation</i>				
	Actual Eligibility	Simulated Eligibility	Coverage At Birth	Low Birth Weight	Birth Weight	Low Birth Weight	Very Low Birth Weight	Gestation Length	Preterm	Very Preterm	Small for Gest Age
Event Time 3	6.013*** (1.171)	6.578*** (1.094)	4.6*** (1.4)	-0.247*** (0.077)	4.678** (1.780)	-0.118 (0.075)	-0.081*** (0.029)	0.002 (0.009)	0.075 (0.115)	-0.046 (0.028)	-0.195** (0.084)
Event Time 2	5.735*** (0.952)	6.047*** (0.859)	5.7*** (1.1)	-0.236*** (0.055)	3.592** (1.755)	-0.088 (0.094)	-0.036 (0.027)	-0.003 (0.008)	0.078 (0.087)	-0.044** (0.019)	-0.165* (0.097)
Event Time 1	5.789*** (0.906)	5.752*** (0.796)	3.7*** (0.9)	-0.134*** (0.042)	1.878* (0.969)	-0.033 (0.065)	-0.053* (0.028)	-0.000 (0.007)	0.074 (0.082)	-0.048** (0.023)	-0.066 (0.069)
Event Time 0	4.856*** (0.551)	4.677*** (0.528)	2.6*** (0.7)	-0.100** (0.039)	-0.071 (1.470)	0.015 (0.079)	-0.045* (0.024)	0.005 (0.009)	0.006 (0.116)	-0.063*** (0.019)	0.017 (0.080)
Event Time -1	Omitted	Omitted	Omitted	Omitted	Omitted	Omitted	Omitted	Omitted	Omitted	Omitted	Omitted
Event Time -2	-0.342 (0.491)	-0.524 (0.544)	-0.7 (0.8)	-0.034 (0.039)	-0.518 (1.080)	-0.004 (0.046)	-0.004 (0.026)	-0.002 (0.006)	-0.011 (0.074)	-0.004 (0.021)	0.087 (0.074)
Event Time -3	-0.310 (0.591)	-0.356 (0.573)	-0.9 (0.9)	0.000 (0.039)	0.806 (1.672)	-0.035 (0.055)	0.027 (0.023)	0.011* (0.006)	-0.093 (0.090)	-0.008 (0.022)	0.001 (0.096)
Event Time -4	-0.193 (0.640)	0.035 (0.572)	-0.5 (1.2)	0.043 (0.049)	-0.473 (1.234)	0.067 (0.054)	0.053** (0.024)	-0.002 (0.006)	0.016 (0.093)	0.024 (0.021)	0.000 (0.068)
Event Time -5	-0.537 (0.774)	0.265 (0.612)	-1.7 (1.6)	0.034 (0.057)	-0.088 (1.208)	0.051 (0.052)	-0.001 (0.027)	0.008 (0.006)	0.013 (0.111)	-0.010 (0.023)	0.018 (0.072)
Mean Y – Overall Sample	14	15	16.8	6.92	3257.81	7.61	1.44	38.93	10.68	0.77	11.51
Mean Y – Treated States in 1975	8.46	9.36	14.3	7.44	3276.99	7.52	1.45	39.02	11.17	0.82	11.25
N	700	700	187,488	700	700	700	700	700	700	700	700

Notes: Coefficient estimates for all outcomes other than birth weight and gestation length are reported in percentage points. Models of eligibility and first generation low birthweight estimated for all infants born 1975-1988. Model of coverage uses hospital discharges related to labor and delivery for 1979-1988. Models of second generation health outcomes estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation for all outcomes except coverage. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size or survey weights where applicable and include state and year of birth (first generation) or mother’s state of birth and mother’s year of birth fixed effects (second generation). All models include controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by state of birth (first generation) or mother’s state of birth (second generation). We report the mean of the outcome variable without de-trending for the full sample and for treated states in 1975 (for coverage we use the first year of data available: 1979). Significance levels: * p<0.10, ** p<0.05, *** p<0.01.

Table 2: Event Study Estimates on Fertility Outcomes and Maternal Characteristics

	First Birth Rate	Age at First Birth	High School Education	Married	White	Black	Other
Event Time 3	-0.190 (0.515)	0.028 (0.032)	-0.524 (0.484)	0.390 (0.308)	0.539 (0.386)	-0.549 (0.388)	0.010 (0.104)
Event Time 2	0.172 (0.416)	0.011 (0.023)	-0.524 (0.383)	0.280 (0.229)	0.491* (0.280)	-0.551* (0.281)	0.059 (0.094)
Event Time 1	0.098 (0.405)	-0.008 (0.021)	-0.362 (0.276)	0.155 (0.196)	0.354** (0.173)	-0.388** (0.185)	0.034 (0.068)
Event Time 0	0.084 (0.318)	-0.008 (0.009)	-0.228 (0.195)	0.076 (0.182)	0.111 (0.166)	-0.150 (0.158)	0.039 (0.047)
Event Time -1	Omitted	Omitted	Omitted	Omitted	Omitted	Omitted	Omitted
Event Time -2	-0.071 (0.158)	0.024* (0.013)	0.264 (0.170)	0.125 (0.134)	-0.123 (0.131)	0.088 (0.124)	0.035 (0.078)
Event Time -3	-0.129 (0.249)	0.024 (0.018)	0.144 (0.263)	0.137 (0.224)	0.015 (0.165)	-0.004 (0.180)	-0.011 (0.072)
Event Time -4	0.094 (0.255)	0.032* (0.016)	0.241 (0.292)	0.219 (0.213)	0.008 (0.179)	0.019 (0.195)	-0.027 (0.081)
Event Time -5	-0.131 (0.342)	0.020 (0.019)	-0.038 (0.403)	-0.096 (0.312)	-0.146 (0.237)	0.074 (0.244)	0.072 (0.109)
Mean Y – Overall Sample	54.69	21.67	74.87	42.50	77.76	19.70	2.54
Mean Y – Treated States in 1975	62.28	21.38	73.13	51.68	77.15	21.27	1.59
N	700	700	700	700	700	700	700

Notes: Coefficient estimates for all outcomes other than age at first birth are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Additional controls for the share of births with revised birth certificate records are included for education and race outcomes, and the share of births with birth certificate records allowing for the report of multiple race categories are included for the race outcomes. Standard errors are clustered by mother's state of birth. We report the mean of the outcome variable without de-trending. Significance levels: * $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$.

Multi-Generational Impacts of Childhood Access to the Safety Net: Early Life Exposure to Medicaid and the Next Generation’s Health

Appendix

Chloe N. East Sarah Miller Marianne Page Laura R. Wherry

A Mother’s Eligibility at the Time of Birth and During Childhood

A.1 Additional Details on Prenatal Eligibility and the 1980s Expansions

In this section, we describe in more detail the pathways for prenatal Medicaid eligibility in our study period. At the beginning of this period, the primary pathway for Medicaid eligibility for pregnant women was through the AFDC program. To qualify for AFDC and its Medicaid benefit, women needed to be single parents, and therefore already have at least one dependent child, as well as meet the AFDC financial eligibility criteria. The solid red line in Appendix Figure A3 tracks the percent of women who were eligible for AFDC, and, therefore, qualified for prenatal Medicaid coverage under this eligibility pathway in the event of a pregnancy.⁶⁹

In addition, optional state AFDC and/or Medicaid eligibility programs expanded coverage to certain groups that, while not specifically targeting pregnant women, sometimes included pregnant women. Some states had AFDC-Unemployed Parent (AFDC-UP) programs, which allowed two-parent families in which the principal earner was unemployed to qualify for AFDC and Medicaid. In addition, optional state Ribicoff children programs allowed minors who met the financial standards for AFDC, but did not qualify due to family structure, to receive Medicaid coverage. Finally, some states also had “Medically Needy” programs that provided coverage for individuals with incomes higher than AFDC levels but with large medical expenses. These three eligibility pathways are depicted in the next three lines on Appendix Figure A3 – the red dashed line, the yellow dashed line, and green dashed line, respectively. As may be seen here, these options predated the study period and there was little to no change in prenatal eligibility under these pathways by 1988.

The focus of our paper is expansions in prenatal Medicaid eligibility that occurred starting in the 1980s under both state options and federal mandates. The 1980s saw a larger number of states exercising existing and new options to extend Medicaid eligibility, followed by two federal law changes that mandated Medicaid eligibility for pregnant women meeting the AFDC financial requirements but not other AFDC eligibility criteria. We describe these changes that specifically impacted prenatal eligibility next.

First, a number of states offered AFDC benefits, and Medicaid coverage, to first-time pregnant women, provided that they met the financial criteria for AFDC under an “AFDC unborn” option. However, the Omnibus Reconciliation Act of 1981 (OBRA81) restricted participation for these women

⁶⁹We construct this figure by calculating the percent of women who would be eligible under each pathway in an additive fashion that reflects the order of eligibility pathways from the most to least restrictive on the figure (shown from lowest to highest lines on the graph). So, for example, we calculate the percent of women eligible under AFDC rules only to plot the solid red line. Then we calculate how many more women would be eligible under the AFDC-UP rules to plot the dashed red line, and so on.

until the sixth month of pregnancy. Following [Currie and Gruber \(1994\)](#), we considered this a source of prenatal eligibility if the state covered women starting during the first trimester of pregnancy. The role of the AFDC unborn pathway on prenatal eligibility may be seen in Appendix Figure [A3](#) in the light blue dashed line.

Second, despite the OBRA81 restriction on AFDC eligibility for unborn children, states exercised alternative options under the Medicaid program (rather than AFDC) to cover pregnant women. A new “Ribicoff unborn” option introduced in 1982 allowed states to provide Medicaid coverage to unborn children, which meant coverage of pregnant women, whose family income qualified them for AFDC. This is shown in the dark blue dashed line.

Third, dating as far back as 1966, some states exercised additional options to cover women with a first-time pregnancy, and pregnant women in two-parent families where the principal earner was unemployed, as long as the families met the financial requirements for AFDC. We call these options “pre-DEFRA state options” since they effectively cover the same groups as a federal mandate to cover these groups under the Deficit Reduction Act (DEFRA, effective October 1984). These state options are shown in the purple dashed line, and the DEFRA mandated changes are shown in the solid purple line.

Fourth, some states also had optional rules in place to provide Medicaid coverage to all AFDC financially eligible pregnant women, regardless of family structure or employment status. We refer to this as a “pre-COBRA state option”, since the Consolidated Omnibus Budget and Reconciliation Act (COBRA, effective July 1986) required states to extend Medicaid eligibility to all pregnant women meeting the AFDC financial requirements. This state option is in the pink dashed line and the federal mandate is in the solid pink line.

Finally, the Omnibus Reconciliation Act of 1986 (OBRA86) authorized states to further expand eligibility for pregnant women with incomes up to the poverty line starting in April 1987 (i.e. “income expansions”). States were only beginning to take up this option at the very end of our study period. As may be seen in Appendix Figure [A3](#) in the dashed red line, broad income expansions came in late during the period and were only adopted on a very small scale.

Appendix Figure [A4](#) displays the role of the different eligibility pathways separately for our treated and control states. The treated states experienced clear increases in the use of pre-DEFRA state options starting in 1980, followed by pre-COBRA state options and the introduction of the federal mandates. The control states were already exercising both pre-DEFRA and pre-COBRA state options throughout the 1970s, and only experienced very gradual increases in prenatal eligibility during the 1980s.

A potentially important point for interpreting our results is that Medicaid eligibility sometimes comes with AFDC eligibility, so we may be estimating the impact of both AFDC and Medicaid. However, the policy changes we exploit in the treated states – pre-DEFRA and pre-COBRA options, DEFRA, and COBRA – only expanded eligibility for Medicaid and not AFDC. Additionally, while control states experience a decline in AFDC Unborn between 1980 and 1982, which also means a reduction in AFDC, we do not believe our results are generated by this change, due to the fact that we get very similar results when we omit all the control states in Figure [9](#).

Each option and the sources for the relevant eligibility rules are listed below. We relied heavily on the Appendix of [Currie and Gruber \(1994\)](#) for information about the different options for Medicaid

eligibility during this period, as well as the sources used by these authors. We supplemented this information with additional sources to compile rules for earlier years.

1. AFDC and AFDC-UP: AFDC program parameters and information on the presence of state AFDC-UP programs for the 1975-1996 years were taken from the Urban Institute's [Transfer Income Model](#), version 3 (TRIM3). Using these parameters, we were able to calculate whether a family was eligible for either program based on state rules, monthly total family income, and family size. For the AFDC-UP program, we assume that a two-parent family is eligible if the state had a program and their maximum hours worked were less than 1200 in the prior year. Following [Currie and Gruber \(1994\)](#), we adjust family income for minors residing with their parents by subtracting the needs standard for a family of that size. For adults ages 19 and older residing with their parents, we do not include parents' income in the eligibility determination.
2. Ribicoff children and unborn options: Information on Ribicoff children programs for 1988 forward were drawn from materials provided by Bruce Meyer and used in [Meyer and Rosenbaum \(2001\)](#). Rules for the 1985-1986 years were drawn from the TRIM3 database and for 1983 from the 1983 Health Care Financing Administration (HCFA)'s *Analysis of State Medicaid Program Characteristics* report. State rules regarding coverage of unborn children under Ribicoff programs, which meant coverage of pregnant women whose income qualified them for AFDC, were taken from the 1983 HCFA report as well. We were unable to locate earlier records of these state Ribicoff programs. We therefore assume that the same Ribicoff children provisions were in place during earlier years, with the exception of the unborn children, which was established in 1982.⁷⁰
3. Medically Needy: State Medically Needy thresholds as a percent of poverty were drawn from TRIM3 (for years 1985 forward), the 1981, 1983, 1984, and 1986 *Medicare and Medicaid Data Books* issued by the Health Care Financing Administration, and the 1977 and 1979 *Data on the Medicaid Program* reports. For the in between years when we were unable to locate Medically Needy rules, we impute the values as the average of the thresholds for the years book-ending the given year. For 1975, we assume the rules were the same as those in place in 1976. Following the description of state Medically Needy programs in these sources, we include all pregnant women who were categorically eligible for state Medicaid programs and compare their net income to the Medically Needy income thresholds for their family size.
4. AFDC-unborn: Prior to OBRA 1981, many states offered AFDC coverage for first-time pregnant women. Following [Currie and Gruber \(1996b\)](#), we consider this as a pathway to Medicaid if the state covered a woman starting in her first trimester. Information on the presence of these programs and whether states included the unborn child in the benefit calculation were taken from the 1974, 1976, 1978, 1980, 1981 *Characteristics of State Plans for Aid to Families with Dependent Children* reports published by the Department of Health and Human Services. For

⁷⁰In 1982, a new Medicaid eligibility category was established for pregnant women by Congress. This is described by Sara Rosenbaum in her 1983 report "The Prevention of Infant Mortality: The Unfulfilled Promise of Federal Health Programs for the Poor." Prior to this new category, it appears that states were able to cover unborn children under the regular Ribicoff children program, but we were unable to find any additional information about states doing so.

any policy changes that occurred between reports, we use the midpoint of the dates reported as the implementation date.

5. Pre-DEFRA state option 1: Separately, a number of states offered Medicaid coverage to first-time pregnant women from the point of medical verification, provided that they met the financial criteria for AFDC. Information on this option for all states and the implementation dates are taken from [Hill \(1987\)](#).
6. Pre-DEFRA state option 2: Some states offered Medicaid coverage to pregnant women in two-parent families if the principal earner was unemployed and the family met the AFDC financial eligibility criteria. Information on this option for all states and the implementation dates are also from [Hill \(1987\)](#).
7. Pre-COBRA state option: Some states offered Medicaid coverage to all pregnant women meeting the financial eligibility criteria for AFDC. Information on this option for all states and the implementation dates are also from [Hill \(1987\)](#).

A.2 Mother’s Childhood Eligibility

In addition to changes in prenatal eligibility, eligibility for children who were not traditionally eligible for AFDC and with family income levels exceeding AFDC cutoffs was also expanded starting in the 1980s. These eligibility changes were first introduced as state options and later by federal mandate. The initial eligibility changes were more “targeted” (following the terminology first used by [Currie and Gruber, 1996b](#)) to the lowest income children – those whose families met the income and resource eligibility criteria for AFDC but who did not otherwise meet the family structure requirements for the AFDC program. The later “broad” eligibility expansions extended eligibility to children with incomes that exceeded the AFDC eligibility levels, and are often referred to as poverty-related expansions. Eligibility levels for children continued to grow during the 1990s as Medicaid eligibility changes continued to be phased in, and later through the 2000s under optional state expansions to higher income children under the Children’s Health Insurance Program (CHIP).

Given these concurrent changes in childhood eligibility, we examine the correlation between the mother’s *in utero* and childhood eligibility in the paper. We also control for her simulated childhood eligibility as a robustness check. Each eligibility pathway and the sources for the relevant eligibility rules are listed below.

For the years 1975 to 1996, Medicaid eligibility is calculated under the eligibility rules for the AFDC and the AFDC-Unemployed Parents (AFDC-UP) programs, optional state programs (e.g. Ribicoff children, Medically Needy described above), and both targeted and poverty-related expansions for children. For the years 1997 to 2006, public eligibility under Medicaid *and* state Children’s Health Insurance Programs (CHIP) is calculated under the rules for poverty-related Medicaid expansions and additional Medicaid expansions or new state programs under CHIP.

Sources for eligibility rules under pathways related to AFDC, AFDC-UP, Ribicoff children, and Medically Needy programs are listed in Section [A.1](#). Information on the poverty-related Medicaid expansions, and later CHIP-related expansions in eligibility by state, including the population targeted, implementation date, and income cutoffs were compiled from the sources below. Income disregard rules

by state and year were downloaded from the Urban Institute’s TRIM3 database.

- [Maternal and Child Update](#), National Governors Association: 9/97, 9/98, 2/99, 1/00, 2/01, 2/02, 2/03
- Enrollment Increases in State CHIP Programs: December 1998 to June 1999, prepared by Vernon K. Smith at Health Management Associates for the Kaiser Commission on Medicaid and the Uninsured, July 30, 1999
- *Implementation of the State Children’s Health Insurance Program: Momentum is Increasing After a Modest Start: First Annual Report*, January 2001 report prepared by Mathematica Policy Research, Inc. by Rosenbach et al.
- [Kaiser Commission on Medicaid and the Uninsured](#) (mostly) annual surveys of state Medicaid/CHIP programs beginning in 2000: available for years 2000, 2002, 2003-2005, and 2007

A.3 Construction of Eligibility Measures

To construct measures of the mother’s eligibility at the time of birth and during childhood (ages 1-18), we used detailed eligibility rules compiled for each state during the period 1975 to 2006. Eligibility was estimated using the year of the eligibility determination and family characteristics, including family structure, income, and information on parental employment. For measures of prenatal eligibility, we calculate eligibility during each month of a given year and use the average monthly eligibility level for that year.

We used the 1977-1989 Annual Social and Economic Supplements (ASEC) of the Current Population Survey (CPS) to estimate mother’s eligibility for public health insurance at the time of birth which is our measure of prenatal eligibility. This measure was estimated using women ages 15-44 and determining their eligibility in event of a pregnancy by state and year during the period 1975-1988. To construct a simulated eligibility measure, we drew a national sample of 3,000 women ages 15-44 for each year and estimated eligibility for this sample using state-specific eligibility rules during that year. Not all states are identifiable in the 1976 CPS, so, to estimate eligibility for the 1975 year, we relied on the CPS for next year (i.e. 1977 year of data) and deflate dollar values using the CPI-U.⁷¹

We used the 1977-2007 ASEC to estimate eligibility for childhood Medicaid coverage by single year of age for cohorts born between 1975-1988. We assumed that birth year was equal to calendar year minus age in order to estimate eligibility by birth year x age x state. These estimates were then added across ages 1-18 in order to create a measure of cumulative childhood eligibility for each birth year by state. To construct a simulated eligibility measure, we used a national sample of 1,000 children of each age for each year and estimated eligibility for this sample using state-specific eligibility rules during that year.

B Mother’s Adult Eligibility

When examining public health insurance eligibility for the first generation during adulthood, we consider eligibility for low-income parents under Medicaid Section 1931 criteria in each state, as well as

⁷¹In the 1976 CPS, 38 states cannot be separately identified but share state codes with other states.

expanded eligibility for health care coverage for parents and childless adults under both waiver and state-funded programs. We also consider expansions under the Affordable Care Act. Information on state eligibility thresholds for coverage for adults for the years 1998-2017 were compiled from the sources listed below.

- [Maternal and Child Update](#), National Governors Association: 2002 through 2010 reports
- [Kaiser Commission on Medicaid and the Uninsured](#) annual surveys of state Medicaid/CHIP programs: 2002-2005, 2007-2009, 2011-2013, and 2015-2018 reports
- Broaddus M, Blaney S, Dude A, Guyer J, Ku L, Peterson J. *Expanding Family Coverage: States' Medicaid Eligibility Policies for Working Families in the Year 2000*. Washington, DC: Center on Budget and Policy Priorities; 2001.
- Busch SH, Duchovny N. Family coverage expansions: Impact on insurance coverage and health care utilization of parents. *Journal of Health Economics*. 2005;24(5):876-890.
- Hearne J. *Medicaid Eligibility for Adults and Children*. Washington, DC: Congressional Research Service, The Library of Congress; 2005.
- Indiana Legislative Services Agency. *The Healthy Indiana Plan and Health Coverage of Childless Adults Across the States*. Indianapolis, IN: Health Finance Committee, Indiana Legislative Services Agency; 2011.
- National Conference of State Legislatures. [State Health Programs to Covered the Uninsured](#), 2009-10. 2010. Accessed May 19, 2014.
- [National Conference of State Legislatures](#). Using Medicaid Dollars to Cover the Uninsured: States Use of Medicaid Dollars to Cover the Uninsured. 2009. Accessed May 19, 2014.
- Somers SA, Hamblin A, Verdier JM, Byrd VL. *Covering Low-Income Childless Adults in Medicaid: Experiences from Selected States*. Center for Health Care Strategies, Inc.; 2010.

Federal law for family coverage under Section 1931 requires that states disregard at least \$90 of earned income per month when assessing Medicaid eligibility (Birnbaum 2000). In 2000, most states were using this minimum earnings disregard in eligibility determinations (Broaddus et al. 2001). Therefore, we chose to apply this rule for all states for the years 1998-2013. For 2014-2017, following the implementation of the Affordable Care Act Medicaid expansions, a standard disregard of five percentage points of the federal poverty level is built into the eligibility thresholds.

We construct a measure of average cumulative adult Medicaid simulated eligibility from age 19 to the age we observe women at child birth by state and birth year cohort. This measure is constructed using a sample of adults ages 19-28 from the Annual Social and Economic Supplements (ASEC) of the Current Population Survey (CPS). We use a random national sample of 1,000 adults per year of age and survey year and estimated eligibility for this sample using state-specific eligibility rules during that year.

C Second Generation’s Own *In Utero* Eligibility at the Time of Birth

To calculate the second generation child’s own *in utero* eligibility (which is the same as the first generation mother’s eligibility at the time of second generation child’s birth), we use the eligibility rules under Medicaid Section 1931 eligibility, poverty-related Medicaid expansions for pregnant women, expanded Medicaid rules authorized under the Balanced Budget Act of 1997, and separate state programs created under the State Children’s Health Insurance Program over the period 1989-2017. Income eligibility cutoffs by state and year were compiled from the sources listed under Appendix Sections A and B.

We use the 1990-2018 Annual Social and Economic Supplements (ASEC) of the Current Population Survey (CPS) to estimate mother’s simulated eligibility for public health insurance at the time of infant’s birth. This measure of *in utero* eligibility was estimated using women ages 15-44. We drew national sample of 3,000 women ages 15-44 for each year between 1990-2017 and estimated eligibility in event of a pregnancy for this sample using state-specific eligibility rules during that year.

D Additional Details on Data Construction

D.1 State-Year Control Variables

In our main analyses we include controls for state economic conditions, demographics, safety net policy, and abortion policy based on the state and year of mother’s birth (first generation). These controls are described in detail below. In additional robustness checks, we include these same variables, as well as several additional variables also described below, at the time of the second generation’s birth.

We include the following demographic controls: the fraction of the state population between the ages of 0-4, 5-17, 18-24, 25-44, 44-64; the fraction of the state population that identifies as Black and identifies as a race other than white or Black; the fraction of the state population with a high school degree, some college, college or more. These were constructed by the authors using the ASEC.

We include the following economic controls: state median household income (from the U.S. Bureau of Labor Statistics) and unemployment rate (from the U.S. Census Bureau). We include maximum welfare benefits at the time of the mother’s birth. In addition, in certain specifications, we include the following measures of welfare generosity at the time of the second generation’s birth: state welfare family cap; whether the state had an EITC program, whether the state had implemented TANF. The sources are:

- Crouse, Gil. 1999. “State Implementation of Major Changes to Welfare Policies, 1992-1998.” Office of Human Services Policy, ASPE, U.S. Department of Health and Human Services.
- Urban Institute TRIM3 Program Rules for 1990-1995
- Urban Institute Welfare Rules Database for 1996-2017
- [NBER TAXSIM](#)
- [Tax Credits for Working Families](#)
- Urban Institute Tax Policy Center

- [University of Kentucky Center for Poverty Research National Welfare Data, 1980-2017](#)
- [Robert Moffitt's Welfare Benefits Data Base](#)

We include the following measures of family planning coverage at the time of the mother's birth: state parental consent and notification laws for abortion and state Medicaid restrictions for abortion. In addition, in certain specifications, we also include the following at the time of the second generation's birth: mandatory delay for abortion laws; income based and duration based Medicaid family planning waivers; state mandate for private health insurance coverage of contraceptives; an indicator that emergency contraceptives can be provided over-the-counter; an indicator that minor may consent to contraceptive services in all or limited circumstances. The sources are:

- *Our Daughters' Decision: The Conflict in State Law on Abortion and Other Issues* by Patricia Donovan, The Alan Guttmacher Institute 1992.
- "Minors and the Right to Consent to Health Care" by Heather Boonstra and Elizabeth Nash, *The Guttmacher Report on Public Policy*, August 2000.
- State Policies in Brief from the Guttmacher Institute on Medicaid Family Planning Eligibility Expansions, Minors' Access to Contraception, State Funding of Abortion Under Medicaid, Mandatory Waiving Periods for Abortion and Parental Involvement in Minors' Abortions
- Kearney, Melissa S. and Phillip B. Levine. 2009. "Subsidized Contraception, Fertility, and Sexual Behavior." *Review of Economics and Statistics* 91(1): 137-151.
- Insurance Coverage for Contraception Laws by the National Conference of State Legislatures
- Oza, Anjali D. The Economics of Emergency Contraception. 2010. University of Chicago PhD Dissertation.
- Levine, Phillip. 2004. Sex and Consequences: Abortion, Public Policy, and the Economics of Fertility.

D.2 Additional Details on Birth Certificate Revision

The variables used to examine mother's (i.e. first generation's) educational attainment, prenatal care utilization, and race were affected by the introduction of the 2003 revision of the U.S. Standard Certificate of Live Birth, which replaced the 1989 revision that was in use during the remainder of the period covered by our analyses. State adoption of the revision is staggered over the period. By January 2011, 36 states and the District of Columbia had implemented the revised birth certificate. These states represent 83 percent of births to U.S. residents ([Centers for Disease Control and Prevention, 2011](#)). Starting in 2011, the CDC no longer made available certain data items from the unrevised birth certificate, including maternal education and prenatal care utilization. As a result, information on these variables is incomplete, and only available for states that had fully implemented the 2003 revision. Fourteen states in 2011, 12 states in 2012, 9 states in 2013, 3 states in 2014, and 2 states in 2015 have incomplete information for these data fields. By 2016 all states had implemented the revised birth certificate.

In addition, even when the data fields are available, these measures are not considered comparable before and after the 2003 revision. Prior to the revision, mother’s education was classified into years of education: no formal education, 1-8 years of elementary school, 1-4 years of high school, 1-4 years of college, and 5 or more years of college. The 2003 revision classified mother’s education into the following categories: 8th grade or less; 9th through 12th grade with no diploma; high school graduate or GED completed; some college credit, but not a degree; associate degree; Bachelor’s degree; Master’s degree; and, doctorate or professional degree. In our analyses, we code high school or less as having at least 4 years of high school under the 1989 revision, and being a high school graduate or having a GED completed under the 2003 revision. In addition, changes occurred in information collected on mother’s race with the 2003 revision including more detailed race categories. Also, beginning in 2003, states had the option of allowing the report of multiple race categories. These multiple race combinations are bridged to a single race category for comparability to other reporting areas and years.

We address the incomparability of these measures after the birth certificate revision by including in regressions for which maternal education, race, or prenatal care utilization are dependent variables a measure of the fraction of birth records in that cell (mother’s birth year x mother’s state of birth) with a revised birth certificate.

D.3 Identifying Labor and Delivery Hospitalizations in the National Hospital Discharge Survey

We identify hospitalizations related to labor and delivery in the National Hospital Discharge survey using ICD-9 diagnosis and procedure codes and following the method validated in [Kuklina et al. \(2008\)](#). First, we code all visits with a diagnosis code starting with “V27” (corresponding to live or still births) or “650” (normal delivery) as discharges related to labor and delivery. Second, if the first three or four digits of the procedure code falls in the subsequent list, we classify the hospitalization as being a labor or delivery: “7251”, “7252”, “7253”, “7254”, “7271”, “7279”, “728”, “720”, “721”, “7221”, “7229”, “7231”, “7239”, “7322”, “7359”, “736”, “740”, “741”, “742”, “744”, “7499”, “724”, “726” or “729.” These codes are associated with different obstetrics procedures. Finally, we exclude discharges with a diagnosis code that indicates an ectopic pregnancy or abortive outcome.

D.4 Additional Details on Implementation of Synthetic Control Event Study Approach

We assess the robustness of our results to using the synthetic control methods pioneered by [Abadie et al. \(2010\)](#) and following the implementation procedure described in [Kleven \(2021\)](#). Specifically, we do the following. First, for each outcome, we match each treated state to a weighted average of the untreated states using pre-expansion values of the outcome and control variables. This results in two observations for each outcome per year for each treated state: the outcome in the treated state, and the outcome in its synthetic control. Second, we stack each set of observations into a single dataset and estimate the following model, using robust standard errors clustered at the state level, with the synthetic control unit treated as a separate state:


$$y_{nt} = \alpha + \sum_{t=-5, t \neq -1}^3 \kappa_t 1\{b - e_n^* = t\} \times Treated_n + \delta_t 1\{b - e_n^* = t\} + \mu_n + \lambda_t + \epsilon_{nt} \quad (3)$$

Note that this equation varies from our baseline model described in equation (2) in a few ways. First, because event times are now defined both for the treated state and the synthetic controls state, we include event time indicators (λ_t) rather than calendar time indicators (λ_b) as in our main specification. Second, in the synthetic control approach we match untreated states to treated states based on the state-level controls in the pre-expansion period rather than control for them directly in the model. Third, we include separate fixed effects for the treated state and its synthetic control counterpart, analogous to the state fixed effects in our baseline model. Finally, we weight these regressions using the average number of births in the treated state applied to both the treated state and its synthetic control counterpart. Our estimates of κ_t are reported in Appendix Figure A15.

D.5 Additional Details on Discounting of Long-Run Benefits


To calculate the discounted value of the second generation benefits, we estimate that in 2011 \$s the value of the benefits from increasing birthweight by 71 grams (our estimated change in birthweight for each newly eligible woman in the first generation) is \$589. We then apply this value of the benefit to each second generation birth cohort from 1995-2017, and calculate the discounted value of this benefit between 1981 and each birth cohort. We have chosen 1981 as the reference point because this is the year for which we have estimated Medicaid costs (Currie and Gruber, 1996b). We then take the average discounted value across second generation birth cohorts to generate a summary measure. We do this for two different discount rates: 1) 0.5%, which is the discount rate recommended for 20-year studies by the Office of Management and Budget (U.S. Office of Management and Budget, 2016), and 2) 3%, which is the discount rate recommended for life-cycle studies by the Department of Commerce (Lavappa and Kneifel, 2016).



















Figure A1: Trends in Actual and Simulated Prenatal Medicaid Eligibility, 1975 to 1988




Notes: Authors' calculation from the Current Population Survey and Medicaid eligibility rules. All estimates are reported in percents. See text for further details.

Figure A2: Trends in Simulated Prenatal Medicaid Eligibility by State, 1975 to 1988









Notes: Authors' calculation from the Current Population Survey and Medicaid eligibility rules. All estimates are reported in percents. See text for further details.

Figure A3: Prenatal Eligibility by Source, 1975-1988




Notes: All estimates are reported in percents. Authors' calculation from the Current Population Survey and Medicaid eligibility rules. Each line represents Medicaid eligibility through each of the state options and federal mandates. Specifically, we construct this figure by calculating the percent of women who would be eligible under each pathway in an additive fashion that reflects the order of eligibility pathways from the most to least restrictive on the figure. These pathways discussed in more detail in Appendix Section A.1.

Figure A4: Prenatal Eligibility by Source and Treatment Status, 1975-1988




Notes: All estimates are reported in percents. Authors' calculation from the Current Population Survey and Medicaid eligibility rules. Each line represents Medicaid eligibility through each of the state options and federal mandates. Specifically, we construct this figure by calculating the percent of women who would be eligible under each pathway in an additive fashion that reflects the order of eligibility pathways from the most to least restrictive on the figure. These pathways discussed in more detail in Appendix Section A.1.

Figure A5: Trends in Second Generation Outcomes by Expansion Cohort




Notes: Estimates for all outcomes other than birthweight and gestation length are reported in percents. Figure plots average values by event time for states that expanded Medicaid eligibility at different years, and the control states, as denoted by the legend. These averages are produced using first-born infants of mothers born in 1975-1988 and ages 15-28. For treated states, the linear trend is estimated using all pre-period years for each treatment cohort. For control states, we use the period 1975-1981 to estimate this trend.

Figure A6: Trends in First Generation Outcomes by Expansion Cohort




Notes: Estimates for low birthweight and very low birthweight are reported in percents. Figure plots average values by event time for states that expanded Medicaid eligibility at different years, and the control states, as denoted by the legend. These averages produced using infants born in 1975-1988. For treated states, the linear trend is estimated using all pre-period years for each treatment cohort. For control states, we use the period 1975-1981 to estimate this trend.

Figure A7: Event Study for Prenatal Eligibility, Robustness to Alternative Controls and Specifications




(a) Actual Eligibility



(b) Simulated Eligibility


Notes: Coefficient estimates are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year) except where otherwise noted. Standard errors are clustered by mother's state of birth.

Figure A8: Event Study for Prenatal Eligibility, Robustness to using a Pooled National Sample to Calculate Eligibility Measure




Notes: Coefficient estimates are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth.

Figure A9: Event Study for Prenatal Eligibility, Robustness to Controls for Mother’s Eligibility at Other Ages, Second Generation’s Own Prenatal Eligibility, and State-Year Controls in Second Generation




(a) Actual Eligibility




(b) Simulated Eligibility

Notes: Coefficient estimates are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother’s state of birth and mother’s year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother’s state of birth.


Figure A10: Event Study for First Generation Outcomes, Robustness to Alternative Controls and Specifications



(a) Low birthweight




(b) Birthweight



(c) Very low birthweight


Notes: Coefficient estimates for low birthweight and very low birthweight are reported in percentage points. Estimated for infants born in 1975-1988. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation except where otherwise noted. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size and include state of birth and year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year) except where otherwise noted. Standard errors are clustered by infant's state of birth.

Figure A11: Trends in Second Generation Outcomes across Treated and Control States, Detrended and Residualized




Notes: Estimates for all outcomes other than birthweight and gestation length are reported in percents. Figure plots average values by event time for treated and control states, as denoted by the legend. These averages are produced using first-born infants of mothers born in 1975-1988 and ages 15-28. For treated states, the linear trend is estimated using all pre-period years for each treatment cohort. For control states, we use the period 1975-1981 to estimate this trend. To residualize, we regress each de-trended outcome on the state by year controls in our baseline model, take the residual, and add back the state-specific pre-period mean.

Figure A12: Event Study for Second Generation Outcomes, Robustness to Controls for Mother’s Eligibility at Other Ages, Second Generation’s Own Prenatal Eligibility, and State-Year Controls in Second Generation




Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother’s state of birth and mother’s year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother’s state of birth.

Figure A13: Event Study for Childhood Eligibility




(a) Actual Eligibility



(b) Simulated Eligibility


Notes: Coefficient estimates are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth.

Figure A14: Event Study for Second Generation Outcomes, Robustness to Callaway and Sant'Anna




Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size. Standard errors are clustered by mother's state of birth.

Figure A15: “Stacked” Synthetic Control Event Studies




Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. We generate a “synthetic control” for each treated state as a weighted average of the untreated states based on pre-treatment values of the outcome variable and the socio-economic and demographic control variables. We then stack the treated units and their synthetic controls and estimate an event study model that includes indicators for state, event time, and event time interacted with an indicator that the state is the “treated” rather than the “synthetic control” unit. Regressions are weighted by second generation birth cohort size. Standard errors are clustered by mother’s state of birth.

Figure A16: Event Study for Second Generation Outcomes, Parity=2+




Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state and parity group prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth.

Figure A17: Event Study for Second Generation Outcomes, All Mom Ages




Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth.

Figure A18: Event Study Coefficients for Second Generation Outcomes, Foreign-Born Mothers Only




Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born outside of the U.S. in 1975-1988 and ages 15-28. We use the state of residence at child's birth rather than mother's state of birth to assign policy and control variable information. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of residence and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of residence.

Figure A19: Event Study for Second Generation Outcomes, Other Outcomes



(a) Any Congenital Anomalies


(b) Any Abnormal Conditions



(c) Female


Notes: Coefficient estimates are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). An additional control for the share of births with missing information for each type of outcome is included. Standard errors are clustered by mother's state of birth.

Figure A20: Event Study for Second Generation Outcomes, White Births Only




Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth.

Figure A21: Event Study for Second Generation Outcomes, Robustness to Controlling for Mother's Demographics



Notes: Coefficient estimates for (b), (c), (e), (f), and (g) are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). We also include controls for mother's education, marital status, and race. Standard errors are clustered by mother's state of birth.

Figure A22: Event Study for Second Generation Outcomes, Mother's Health Behaviors and Health Outcomes



Notes: Coefficient estimates for all outcomes other than number of prenatal visits are reported in percentage points. Estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by second generation birth cohort size and include mother's state of birth and mother's year of birth fixed effects and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). An additional control for the share of births with revised birth certificate records is included for the outcomes related to prenatal care utilization. Standard errors are clustered by mother's state of birth.

Table A1: Replication of Currie and Gruber (1996) Low Birth Weight Result Using Simulated Eligibility IV Approach

	Currie and Gruber (1996) Table 3	Replication	Cluster by State	Add State Controls	Weight by # Births
<i>All Expansions</i>					
Prenatal eligibility	-4.347* (2.601)	-5.876** (2.927)	-5.876 (5.013)	-5.207 (4.927)	-6.956* (3.950)
Mean Y	68.12	68.09	68.09	68.09	68.96
N	700	700	700	700	700
<i>Targeted Expansions</i>					
Prenatal eligibility, targeted	-17.81*** (4.294)	-16.202*** (3.819)	-16.202** (6.568)	-15.499** (7.013)	-10.836** (5.194)
Mean Y	68.12	68.09	68.09	68.09	68.96
P-Value	0.00	0.00	0.01	0.03	0.04
N	700	700	700	700	700

Notes: The dependent variable is the number of low birth weight births per 1000 births. Models are estimated for all births born between 1979-1992. All models include include state of birth and year of birth fixed effects. Where noted, regressions are weighted by the birth cohort size and controls for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by infant's state of birth, where noted. Significance levels: * p<0.10, ** p<0.05, *** p<0.01.

Table A2: Infant Health, Fertility, Maternal Characteristics, Eligibility, and Controls in Base Year

	Full Sample	Treated States	Control States
<i>First Generation Infant Health Outcomes</i>			
Low Birthweight	7.28	7.44	7.08
<i>Second Generation Infant Health Outcomes</i>			
Birthweight	3286.69	3276.99	3296.67
Low Birthweight	7.22	7.52	6.92
Very Low Birthweight	1.39	1.45	1.32
Gestational Length	39.08	39.02	39.14
Preterm	10.62	11.17	10.05
Very Preterm	0.76	0.82	0.70
Small for Gestational Age	11.14	11.25	11.03
<i>Second Generation Fertility Outcomes</i>			
First Birth Rate	59.36	62.28	56.34
Age at First Birth	21.44	21.38	21.50
<i>Second Generation Maternal Characteristics</i>			
High School Education	73.84	73.13	74.57
Married	49.69	51.68	47.65
White	78.71	77.15	80.32
Black	19.31	21.27	17.28
Other	1.98	1.59	2.40
Any Prenatal Care	98.99	98.99	98.98
Num Prenatal Visits	11.57	11.57	11.55
1st Trimester Prenatal	79.46	79.39	79.54
Any Medical Risk Factor	7.93	8.43	7.41
Diabetes	1.81	1.82	1.79
Chronic Hypertension	0.58	0.62	0.53
Pregnancy-Related Hypertension	5.45	5.90	4.98
Eclampsia	0.56	0.58	0.54
<i>State Demographics</i>			
Married	45.26	45.52	44.98
Black	12.23	14.51	9.87
Other	1.75	1.06	2.45
High School Drop Out	33.27	34.86	31.62
High School Education	37.59	36.68	38.52
Some College	23.88	23.26	24.51
Population Age 0-4	19.56	19.89	19.23
Population Age 5-17	23.14	23.61	22.66
Population Age 18-24	12.97	13.14	12.79
Population Age 25-44	26.04	26.13	25.94
Population Age 45-64	20.20	19.69	20.73
<i>State Economic and Policy Variables</i>			
Unemployment Rate	8.33	7.30	9.39
Income per Capita	27.55	26.32	28.83
Max AFDC Benefits for Fam of 4	1235.74	1038.43	1438.86
Whether Consent/Notification for Abortion	0	0	0
Whether Medicaid covers Abortion	0	0	0
<i>Medicaid Generosity</i>			
Actual Prenatal Eligibility	11.63	8.46	14.90
Simulated Prenatal Eligibility	12.64	9.36	16.02
Number of States	50	28	22

Notes: All proportions are multiplied by 100. First generation outcomes means estimated using a sample of first-born infants of mothers born in 1975 and ages 15-28 at child birth and weighted by size of maternal birth cohort. Means of second generation outcomes, state demographics, state economic and policy variables, and Medicaid generosity estimated using a sample of first-born infants of mothers born in 1975 and ages 15-28 at child birth and weighted by the second generation birth cohort size.

Table A3: Characteristics of Women Gaining Eligibility In Treatment vs. Control States

	Treatment States	Control States
Family Income, 1981 \$ (SD)	3,587.87 (3,138.04)	5,858.92 (3,014.45)
Less than 50% FPL, %	46.6	12.9
Less than 100% FPL, %	85.1	60.0
Married, %	28.1	34.8
Number of kids, N (SD)	0.81 (1.33)	1.11 (1.39)
White race, %	77.1	81.2
Black race, %	18.8	12.7
Other race, %	5.5	4.2
Employed, %	48.8	54.1

Notes: Data are weighted means from the 1981-1989 ASEC IPUMS-CPS. Means for treatment states are calculated for women who are eligible during post-period but were not eligible under the rules in place in the last pre-treatment year. Means for control states are calculated for women who are eligible during 1982-1988 but were not eligible under the rules in place in 1981.

Table A4: Event Study Estimates on First Generation Health

	Birthweight	Low Birthweight	Very Low Birthweight
Event Time 3	1.200 (2.764)	-0.247*** (0.077)	-0.065** (0.029)
Event Time 2	1.437 (2.233)	-0.236*** (0.055)	-0.052** (0.023)
Event Time 1	0.440 (1.830)	-0.134*** (0.042)	-0.014 (0.017)
Event Time 0	-0.842 (1.583)	-0.100** (0.039)	-0.018 (0.016)
Event Time -1	Omitted	Omitted	Omitted
Event Time -2	0.163 (1.312)	-0.034 (0.039)	-0.004 (0.018)
Event Time -3	-1.185 (1.705)	0.000 (0.039)	0.003 (0.021)
Event Time -4	-1.284 (1.870)	0.043 (0.049)	0.032 (0.025)
Event Time -5	-2.098 (2.446)	0.034 (0.057)	0.003 (0.021)
Mean Y - Overall Sample	3338.34	6.92	1.18
Mean Y - Treated States in 1975	3302.71	7.44	1.19
N	700	700	700

Notes: Coefficient estimates for low birthweight and very low birthweight are reported in percentage points. Models are estimated for all infants born 1975-1988. The dependent variables are the average birthweight, the percent low birthweight, and the percent very low birthweight in each state by year of birth cell. Pre-period trend is estimated and removed from all observations for each state prior to the event study estimation. For treated states, this is estimated in using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. All models include state of birth and year of birth fixed effects. Regressions are weighted by the birth cohort size and control for state-year variables (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by infant's state of birth. We report the mean of the outcome variable without de-trending for the full sample and for treated states in 1975 (for coverage we use the first year of data available: 1979). Significance levels: * $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$.

Table A5: Event Study Estimates on Second Generation Birthweight, Robustness Checks

	Baseline	No State*Year Controls	Control Reg*Year FE	Don't Detrend	Mother Cohort Weights	No Control States
3	4.678** (1.780)	4.372** (1.991)	2.735 (1.985)	2.169 (1.387)	4.919*** (1.779)	7.928*** (2.018)
2	3.592** (1.755)	3.477* (1.907)	2.082 (1.810)	2.440* (1.347)	3.538** (1.732)	6.254*** (2.100)
1	1.878* (0.969)	1.485 (1.055)	0.358 (1.261)	0.536 (1.007)	1.883* (0.972)	4.041** (1.462)
0	-0.071 (1.470)	-0.580 (1.428)	-1.009 (1.634)	-1.085 (1.470)	0.037 (1.435)	1.470 (1.714)
-1	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)
-2	-0.518 (1.080)	-0.657 (1.137)	-0.553 (1.323)	-0.373 (1.121)	-0.351 (1.078)	-0.908 (1.299)
-3	0.806 (1.672)	-0.121 (1.694)	0.837 (1.913)	-0.600 (1.413)	1.057 (1.605)	0.039 (2.339)
-4	-0.473 (1.234)	-1.257 (1.296)	-0.487 (1.690)	-1.545 (1.507)	-0.440 (1.212)	-1.513 (2.930)
-5	-0.088 (1.208)	-1.058 (1.264)	0.154 (1.657)	-1.232 (1.502)	0.024 (1.223)	-0.420 (2.937)
N	700	700	700	700	700	392

Notes: Models of second generation health outcomes estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. The pre-period trend is estimated and removed from all observations for each state prior to the event study estimation for all outcomes except where noted. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size except where noted and include mother's state of birth and mother's year of birth fixed effects (second generation). All models include controls for state-year variables except where noted (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth. Significance levels: * p<0.10, ** p<0.05, *** p<0.01.

Table A6: Event Study Estimates on Second Generation Low Birthweight, Robustness Checks

	Baseline	No State*Year Controls	Control Reg*Year FE	Don't Detrend	Mother Cohort Weights	No Control States
3	-0.118 (0.075)	-0.090 (0.069)	-0.085 (0.079)	-0.062 (0.053)	-0.120 (0.075)	-0.233** (0.090)
2	-0.088 (0.094)	-0.064 (0.090)	-0.084 (0.092)	-0.062 (0.070)	-0.085 (0.094)	-0.201* (0.098)
1	-0.033 (0.065)	0.000 (0.056)	0.013 (0.065)	-0.001 (0.048)	-0.031 (0.065)	-0.112 (0.076)
0	0.015 (0.079)	0.039 (0.072)	0.037 (0.081)	0.042 (0.071)	0.021 (0.076)	-0.054 (0.087)
-1	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)
-2	-0.004 (0.046)	0.013 (0.045)	0.021 (0.052)	0.002 (0.047)	-0.008 (0.045)	0.006 (0.064)
-3	-0.035 (0.055)	-0.004 (0.053)	-0.016 (0.064)	0.011 (0.058)	-0.041 (0.055)	-0.030 (0.083)
-4	0.067 (0.054)	0.091* (0.048)	0.094 (0.063)	0.102* (0.055)	0.062 (0.053)	0.106 (0.074)
-5	0.051 (0.052)	0.080 (0.048)	0.098 (0.059)	0.096* (0.050)	0.046 (0.054)	0.082 (0.088)
N	700	700	700	700	700	392

Notes: Coefficient estimates are reported in percentage points. Models of second generation health outcomes estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. The pre-period trend is estimated and removed from all observations for each state prior to the event study estimation for all outcomes except where noted. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size except where noted and include mother's state of birth and mother's year of birth fixed effects (second generation). All models include controls for state-year variables except where noted (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth. Significance levels: * p<0.10, ** p<0.05, *** p<0.01.

Table A7: Event Study Estimates on Second Generation Very Low Birthweight, Robustness Checks

	Baseline	No State*Year Controls	Control Reg*Year FE	Don't Detrend	Mother Cohort Weights	No Control States
3	-0.081*** (0.029)	-0.080*** (0.027)	-0.081** (0.032)	-0.042* (0.023)	-0.086*** (0.030)	-0.065 (0.046)
2	-0.036 (0.027)	-0.036 (0.027)	-0.034 (0.029)	-0.007 (0.023)	-0.040 (0.029)	-0.033 (0.036)
1	-0.053* (0.028)	-0.053** (0.025)	-0.052* (0.030)	-0.033 (0.023)	-0.059** (0.028)	-0.052 (0.034)
0	-0.045* (0.024)	-0.035* (0.020)	-0.025 (0.025)	-0.028 (0.022)	-0.050** (0.023)	-0.052* (0.027)
-1	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)
-2	-0.004 (0.026)	-0.007 (0.025)	-0.006 (0.029)	-0.013 (0.026)	-0.007 (0.026)	-0.011 (0.031)
-3	0.027 (0.023)	0.027 (0.023)	0.030 (0.026)	0.019 (0.023)	0.023 (0.023)	0.003 (0.031)
-4	0.053** (0.024)	0.049** (0.024)	0.068** (0.028)	0.033 (0.023)	0.052** (0.024)	0.020 (0.035)
-5	-0.001 (0.027)	0.001 (0.022)	0.001 (0.030)	-0.017 (0.025)	-0.002 (0.027)	-0.054 (0.033)
N	700	700	700	700	700	392

Notes: Coefficient estimates are reported in percentage points. Models of second generation health outcomes estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. The pre-period trend is estimated and removed from all observations for each state prior to the event study estimation for all outcomes except where noted. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size except where noted and include mother's state of birth and mother's year of birth fixed effects (second generation). All models include controls for state-year variables except where noted (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth. Significance levels: * p<0.10, ** p<0.05, *** p<0.01.

Table A8: Event Study Estimates on Second Generation Gestational Length, Robustness Checks

	No State*Year		Control	Don't	Mother Cohort	No Control
	Baseline	Controls	Reg*Year FE	Detrend	Weights	States
3	0.002 (0.009)	0.006 (0.010)	0.001 (0.010)	0.014* (0.008)	0.003 (0.009)	0.030** (0.014)
2	-0.003 (0.008)	0.001 (0.008)	-0.004 (0.008)	0.012* (0.006)	-0.001 (0.008)	0.018 (0.013)
1	-0.000 (0.007)	0.001 (0.008)	-0.002 (0.008)	0.008 (0.008)	0.000 (0.007)	0.017 (0.011)
0	0.005 (0.009)	0.004 (0.008)	0.002 (0.009)	0.007 (0.008)	0.006 (0.009)	0.015 (0.011)
-1	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)
-2	-0.002 (0.006)	-0.001 (0.007)	-0.004 (0.007)	-0.007 (0.007)	-0.001 (0.006)	-0.010 (0.007)
-3	0.011* (0.006)	0.009 (0.006)	0.010 (0.006)	-0.003 (0.008)	0.012** (0.006)	-0.001 (0.009)
-4	-0.002 (0.006)	-0.004 (0.007)	-0.001 (0.006)	-0.016* (0.009)	-0.001 (0.005)	-0.017 (0.010)
-5	0.008 (0.006)	0.008 (0.006)	0.009* (0.005)	-0.010 (0.009)	0.008 (0.006)	-0.007 (0.011)
N	700	700	700	700	700	392

Notes: Models of second generation health outcomes estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. The pre-period trend is estimated and removed from all observations for each state prior to the event study estimation for all outcomes except where noted. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size except where noted and include mother's state of birth and mother's year of birth fixed effects (second generation). All models include controls for state-year variables except where noted (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth. Significance levels: * $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$.

Table A9: Event Study Estimates on Second Generation Preterm Birth, Robustness Checks

	Baseline	No State*Year Controls	Control Reg*Year FE	Don't Detrend	Mother Cohort Weights	No Control States
3	0.075 (0.115)	0.090 (0.111)	0.024 (0.112)	-0.066 (0.058)	0.060 (0.115)	-0.016 (0.179)
2	0.078 (0.087)	0.062 (0.090)	0.027 (0.087)	-0.059 (0.056)	0.067 (0.085)	0.018 (0.137)
1	0.074 (0.082)	0.084 (0.080)	0.107 (0.082)	0.001 (0.059)	0.063 (0.078)	-0.000 (0.130)
0	0.006 (0.116)	0.023 (0.110)	0.015 (0.106)	-0.018 (0.104)	-0.008 (0.113)	-0.030 (0.145)
-1	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)
-2	-0.011 (0.074)	-0.056 (0.056)	0.012 (0.067)	0.006 (0.066)	-0.020 (0.076)	0.022 (0.072)
-3	-0.093 (0.090)	-0.149** (0.072)	-0.076 (0.089)	-0.029 (0.083)	-0.104 (0.091)	-0.002 (0.118)
-4	0.016 (0.093)	-0.047 (0.076)	0.034 (0.085)	0.090 (0.086)	-0.002 (0.094)	0.101 (0.116)
-5	0.013 (0.111)	-0.080 (0.079)	0.048 (0.088)	0.103 (0.092)	0.007 (0.112)	0.087 (0.147)
N	700	700	700	700	700	392

Notes: Coefficient estimates are reported in percentage points. Models of second generation health outcomes estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. The pre-period trend is estimated and removed from all observations for each state prior to the event study estimation for all outcomes except where noted. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size except where noted and include mother's state of birth and mother's year of birth fixed effects (second generation). All models include controls for state-year variables except where noted (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth. Significance levels: * p<0.10, ** p<0.05, *** p<0.01.

Table A10: Event Study Estimates on Second Generation Very Preterm Birth, Robustness Checks

	Baseline	No State*Year Controls	Control Reg*Year FE	Don't Detrend	Mother Cohort Weights	No Control States
3	-0.046 (0.028)	-0.047* (0.026)	-0.047 (0.028)	-0.034 (0.023)	-0.055* (0.029)	-0.015 (0.031)
2	-0.044** (0.019)	-0.046** (0.018)	-0.051** (0.022)	-0.037** (0.015)	-0.052*** (0.019)	-0.016 (0.028)
1	-0.048** (0.023)	-0.047** (0.021)	-0.042* (0.024)	-0.043** (0.019)	-0.056** (0.023)	-0.035 (0.033)
0	-0.063*** (0.019)	-0.055*** (0.018)	-0.051** (0.020)	-0.053*** (0.018)	-0.068*** (0.020)	-0.055** (0.022)
-1	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)
-2	-0.004 (0.021)	-0.002 (0.022)	-0.003 (0.022)	-0.005 (0.022)	-0.008 (0.023)	-0.000 (0.025)
-3	-0.008 (0.022)	0.000 (0.020)	-0.009 (0.022)	-0.009 (0.019)	-0.011 (0.021)	-0.008 (0.024)
-4	0.024 (0.021)	0.027 (0.023)	0.022 (0.023)	0.014 (0.020)	0.024 (0.021)	0.002 (0.025)
-5	-0.010 (0.023)	-0.012 (0.019)	-0.014 (0.025)	-0.022 (0.019)	-0.012 (0.023)	-0.039* (0.021)
N	700	700	700	700	700	392

Notes: Coefficient estimates are reported in percentage points. Models of second generation health outcomes estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. The pre-period trend is estimated and removed from all observations for each state prior to the event study estimation for all outcomes except where noted. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size except where noted and include mother's state of birth and mother's year of birth fixed effects (second generation). All models include controls for state-year variables except where noted (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth. Significance levels: * p<0.10, ** p<0.05, *** p<0.01.

Table A11: Event Study Estimates on Second Generation Small for Gestational Age, Robustness Checks

	Baseline	No State*Year Controls	Control Reg*Year FE	Don't Detrend	Mother Cohort Weights	No Control States
3	-0.195** (0.084)	-0.158* (0.091)	-0.144 (0.098)	-0.014 (0.089)	-0.210** (0.082)	-0.261 (0.167)
2	-0.165* (0.097)	-0.137 (0.096)	-0.137 (0.100)	-0.053 (0.080)	-0.162 (0.098)	-0.218* (0.126)
1	-0.066 (0.069)	-0.028 (0.064)	-0.033 (0.073)	0.036 (0.060)	-0.070 (0.073)	-0.093 (0.098)
0	0.017 (0.080)	0.047 (0.074)	0.022 (0.087)	0.065 (0.072)	0.016 (0.080)	-0.028 (0.089)
-1	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)	0.000 (.)
-2	0.087 (0.074)	0.119* (0.068)	0.050 (0.084)	0.049 (0.083)	0.084 (0.071)	0.067 (0.098)
-3	0.001 (0.096)	0.064 (0.097)	-0.031 (0.099)	-0.016 (0.089)	-0.017 (0.090)	-0.032 (0.114)
-4	0.000 (0.068)	0.058 (0.068)	-0.016 (0.092)	-0.016 (0.070)	0.001 (0.064)	-0.003 (0.128)
-5	0.018 (0.072)	0.103* (0.060)	0.006 (0.094)	-0.012 (0.077)	0.011 (0.070)	-0.047 (0.158)
N	700	700	700	700	700	392

Notes: Coefficient estimates are reported in percentage points. Models of second generation health outcomes estimated for first-born infants of mothers born in 1975-1988 and ages 15-28. The pre-period trend is estimated and removed from all observations for each state prior to the event study estimation for all outcomes except where noted. For treated states, this is estimated using all pre-period years for each state. For control states, we use the period 1975-1981 to estimate this trend. Regressions are weighted by birth cohort size except where noted and include mother's state of birth and mother's year of birth fixed effects (second generation). All models include controls for state-year variables except where noted (unemployment rate, personal income per capita, maximum welfare benefit for a family of 4, indicators for state parental consent and notification laws and state Medicaid restrictions for abortion, and demographic controls for each state and year). Standard errors are clustered by mother's state of birth. Significance levels: * p<0.10, ** p<0.05, *** p<0.01.