

 Open access • Book Chapter • DOI:10.1002/9781119961154.CH19

Multilevel Models for Ordinal Data — [Source link](#)

Leonardo Grilli, Carla Rampichini

Institutions: University of Florence

Published on: 09 Nov 2011

Topics: Ordinal data, Ordinal regression, Multilevel model, Covariate and Regression analysis

Related papers:

- [Analysis of ordinal categorical data](#)
- [mvord: An R Package for Fitting Multivariate Ordinal Regression Models](#)
- [Estimation of nonparametric ordinal logistic regression model using local maximum likelihood estimation](#)
- [Binary Dynamic Logit for Correlated Ordinal: estimation, application and simulation](#)
- [Higher Order Markov Structure-Based Logistic Model and Likelihood Inference for Ordinal Data](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/multilevel-models-for-ordinal-data-1xytsw1y>

Index

- ABC Company, 4, 19
ABC survey, 155, 160, 164, 193, 224, 343, 420, 430
ACSS questionnaire, 6, 15, 20, 108, 119–120, 122
adjustment classes, 47–48
alternating least squares (ALS), 309, 335
American Customer Satisfaction Index (ACSI), 15, 116, 122, 232
annual customer satisfaction survey (ACSS), 6, 15, 20, 29, 145–146
assignment mechanism, 174–175, 179–180
association test, 297–300
- Bayesian control chart, 432–433
Bayesian inference, 133, 145, 150, 185, 211
Bayesian network (BN), 4, 79, 193–195, 197–199
BOT1+2, 14, 24–26
- calibration, 48–49
causal estimands, 177–179, 185, 189
cause and effect, 195, 201, 211, 213
census, 5, 9, 15, 38–39, 41, 43
chi-square automatic interaction detection (CHAID), 284
classical MDS, 358, 362, 364, 368
classification and regression trees (CART), 284, 286
clustered data, 16, 391
combined uniform and shifted binomial (CUB) model, 250
computer assisted paper interview (CAPI), 89
computer assisted telephone interview (CATI), 45
computer assisted web interview (CAWI), 45, 90–91
conceptual construct, 12–13
conditional probability (CP), 399
contextual effects, 231, 399
contingency tables, 16, 211, 222–223, 228
control chart, 16, 413, 415–416
correlated responses, 395
covariates, 131, 133–134, 178–179, 182, 184
cumulative model, 394, 396
cumulative sum (CUSUM), 426
customer loyalty, 4–5, 15, 24, 37, 72, 85, 108, 112
customer loyalty index, 24, 488
customer satisfaction, 4–5
customer satisfaction surveys (CSS), 4–7
- data sources, 73
decision drivers, 27, 120
descriptive statistics, 4
design of observational studies, 185
directed acyclic graph (DAG), 194
- e-Research, 89
Eurobarometer (EB), 16, 193, 201, 203, 213
- fixed cost, 92–93
forward search, 157–158, 382–383
fuzzy numbers, 439–442
- gap model, 12–13, 110
generalized linear models, 162, 394
Gower's general coefficient of similarity, 361
Guttman scale, 261
- Heckman two-step procedure, 50–51
hierarchical Bayesian approach, 50–51
- ignorability, 132–134, 137
impurity measure, 288–289
imputation methods, 130
information quality (InfoQ), 5, 8, 11, 72
inner model, 310–312
insights, 3, 6–7, 87, 108
integrated analysis, 71–87
internet survey, 97
ISO 10004, 16, 413–415
ISO 7870, 16, 413
item non response, 42, 44, 46, 49–50
item response theory (IRT), 263
- joint probability distribution (JPD), 194–195

500 INDEX

- latent variable (LV), 14, 16
- likelihood function, 133, 144–145, 196, 228, 236, 237, 239, 242, 244, 246, 263, 401
- linear structural models with latent variables (LISREL), 14–15
- linguistic variables, 439
- logistic model, 14, 261, 264, 428
- log-linear model, 16, 141, 213, 218–219
- marketing integrated system (MIT), 15, 71
- masking effect, 168
- maximum likelihood, 129, 145, 198, 217, 219–222, 237, 260–261, 400, 422
- maximum likelihood methods, 217
- measurement scales, 12, 15, 55
- measurement theory, 56
- membership function, 439–441
- Minkowski family, 360–361, 368
- missing at random (MAR), 132, 400
- missing completely at random (MCAR), 47, 132
- missing data, 10, 15
- missing not at random (MNAR), 10, 15–16, 129, 131
- mixed model, 278, 403
- mixed-type data, 360–361, 368, 373
- mixture models, 144
- M-test, 97, 413, 433–434
- multidimensional scaling (MDS), 16, 357
- multilevel models, 213, 250, 404–408
- multiple imputation (MI), 138–140
- multivariate outliers, 160, 374
- nominal data, 16
- non sampling errors, 41–42
- nonlinear principal component analysis (NPCA), 336–338
- nonmetric MDS, 357
- nonresponse in surveys, 130
- observational studies, 175, 178
- observed variable, 56–57, 59, 133, 136
- online survey, 97–98
- optimal scaling, 335, 355
- ordinal data, 15, 51
- ordinal outliers, 160–161
- ordinal regression, 155, 161
- ordinary least squares (OLS), 184, 309
- outer model, 310, 312
- overall satisfaction, 22–24
- panel, 88
- paper and pencil interview (PAPI), 89
- partial credit model (PCM), 261
- partial least squares (PLS), 268
- perceived service quality, 108–109
- Poisson counts, 426, 433
- potential outcomes, 174–175
- principal component analysis (PCA), 272, 333–334
- principal stratification, 189
- Procrustes analysis, 358, 371, 374, 380
- propensity score, 49–50
- proportional odds model, 251
- proximity, 357, 358
- pruning, 291–292
- R code, 423, 459
- random effects, 16, 250
- randomized studies, 16, 132
- Rasch analysis, 259
- Rasch model, 14, 16, 228, 259, 263
- rating scale model (RSM), 261
- recursive partitioning, 285–286
- recursive partitioning (PARTY), 291–297
- representational measurement theory, 56
- return rates, 85, 435
- R-indicators, 46–47
- robust methods, 157
- root cause analysis (RCA), 71, 78–79
- Rubin causal model, 173, 179, 185
- sample surveys, 37, 51
- sampling frame, 9, 40, 94, 241, 243
- satisfaction indicators, 12, 38, 340
- scale transformations, 60–61
- scaling by majorizing a complicated function (SMACOF), 367
- scenario analysis, 200–201, 211, 213
- sensitivity analysis, 212
- sequential probability ratio test (SPRT), 428–429
- service level agreement (SLA), 6, 8
- service quality, 4, 12, 15, 85
- services of general interest (SGI), 201
- SERVPERF, 13, 111
- SERVQUAL, 13, 110
- splitting rule, 284–285
- stable unit treatment assumption (SUTVA), 176–177
- statistical process control (SPC), 260–417
- strengths, 27, 112
- structural equation model (SEM), 310
- structural equation model with latent variable (SEM-LV), 65
- survey items, 86
- survey respondents, 9, 96, 421, 436
- survey response rates, 9
- survey topics, 20, 24–25, 27
- TOP5, 14, 24–26, 425
- tree graphs, 284–285
- unobserved heterogeneity, 397, 405
- unobserved variable, 3, 341
- variable cost, 92
- weaknesses, 27
- web survey (WS), 4, 15, 45, 50, 94
- weighted classical MDS, 368
- weighting criteria, 440, 445–446, 448, 451