

Multilocus sequence typing suggests the chytrid pathogen of amphibians is a recently emerged clone

ERICA A. MOREHOUSE,*TIMOTHY Y. JAMES,*AUSTEN R. D. GANLEY,*RYTAS VILGALYS,*
LEE BERGER,†‡PETER J. MURPHY§ and JOYCE E. LONGCORE¶

*Department of Biology, Biological Sciences Building, Room 139, Duke University, Durham, NC 27708, USA, †Australian Animal Health Laboratory, Commonwealth Scientific Industrial Research Organization, Geelong 3200, Victoria, Australia, ‡School of Public Health and Tropical Medicine, James Cook University, Townsville 4811, Queensland, Australia, §Department of Zoology, Southern Illinois University, Carbondale, IL 62901 USA, ¶Department Biological Sciences, University of Maine, Orono, ME 04469, USA

Abstract

Chytridiomycosis is a recently identified fungal disease associated with global population declines of frogs. Although the fungus, *Batrachochytrium dendrobatidis*, is considered an emerging pathogen, little is known about its population genetics, including the origin of the current epidemic and how this relates to the dispersal ability of the fungus. In this study, we use multilocus sequence typing to examine genetic diversity and relationships among 35 fungal strains from North America, Africa and Australia. Only five variable nucleotide positions were detected among 10 loci (5918 bp). This low level of genetic variation is consistent with the description of *B. dendrobatidis* as a recently emerged disease agent. Fixed (i.e. 100%) or nearly fixed frequencies of heterozygous genotypes at two loci suggested that *B. dendrobatidis* is diploid and primarily reproduces clonally. In contrast to the lack of nucleotide polymorphism, electrophoretic karyotyping of multiple strains demonstrated a number of chromosome length polymorphisms.

Keywords: amphibian declines, *Batrachochytrium dendrobatidis*, chromosome length polymorphism, clonality, emerging infectious disease, resting spore

Received 12 August 2002; revision received 14 October 2002; accepted 14 October 2002

Introduction

Emerging infectious diseases of wildlife are newly recognized or expanding diseases that threaten biodiversity and ecosystem function. Most emerging wildlife diseases are believed to result from anthropogenic disturbances such as 'spill-over' of infectious agents from humans or domestic animals, human-mediated host or parasite translocations, or climate change (Daszak *et al.* 2000). If their origins are related to disturbance, emerging pathogens may display nonequilibrium population genetics because their geographical or host range has expanded. For example, recent expansion has been invoked to explain low levels of detectable genetic variation in emerging diseases of humans (Sreevatsan *et al.* 1997; Achtman *et al.* 1999).

Chytridiomycosis, a disease of amphibians caused by the chytrid fungus *Batrachochytrium dendrobatidis*, has been

implicated as the cause of rapid declines of amphibian populations in pristine and disturbed habitats in Central America, the western United States, Spain and Australia (Berger *et al.* 1998; Daszak *et al.* 1999; Lips 1999; Bosch *et al.* 2001; Bradley *et al.* 2002). The fungus, which infects keratinized epidermal cells, can cause irregular epidermal hyperplasia and hyperkeratosis, but reasons for death are uncertain (Berger *et al.* 1998; Pessier *et al.* 1999). The pathogen has been found in museum specimens collected as early as 1974 in the United States (Carey *et al.* 1999) and 1978 in Australia (Speare *et al.* 2001). The recent discovery of the disease (Berger *et al.* 1998; Pessier *et al.* 1999) and increasing recognition of its widespread occurrence qualifies chytridiomycosis as an emerging infectious disease (Daszak *et al.* 1999).

Batrachochytrium is a member of the fungal phylum Chytridiomycota in the order Chytridiales. Chytrids produce motile, aquatically distributed zoospores that typically possess a single posterior flagellum. Axenic culture of *Batrachochytrium* has made developmental (Longcore

Correspondence: Tim James. Fax: 919-660-7293; E-mail: tyj2@duke.edu

et al. 1999) and experimental studies possible, including fulfilling Koch's postulates (Nichols *et al.* 2001). Sexual reproduction in chytrid fungi is associated with the production of a resistant, thick-walled resting spore in 'all well-authenticated occurrences' (Sparrow 1960). For *B. dendrobatidis*, no sexual or asexual resting structures have been observed on the amphibian host or in culture. The possession of a resting stage could confer the ability to persist in the absence of amphibians and to be transported by wind, perhaps explaining the widespread distribution of the disease in relatively pristine areas (Berger *et al.* 1998; Daszak *et al.* 1999).

One hypothesis for the emergence of chytridiomycosis is recent human introduction of the fungus into naïve populations (Berger *et al.* 1999; Daszak *et al.* 1999). Alternatively, or in concert, climate change or pollutants may have altered a pre-existing host-parasite relationship (Daszak *et al.* 2000; Blaustein & Kiesecker 2002). In this study, we investigate the genetic diversity of this pathogen using multilocus sequence typing (MLST) to test these two hypotheses. The prediction for a disease that has emerged because of recent introduction or translocation is that of lower geographical population structure and lower overall genetic diversity than would be found if the disease has emerged because of extrinsic factors such as environmental change. We also used the MLST data to test whether *B. dendrobatidis* reproduces clonally or sexually.

Materials and methods

Strains and cultivation techniques

Our sample consisted of 35 strains isolated from 19 different species (Table 1). The majority of the amphibian hosts sampled displayed clinical signs of chytridiomycosis. Strains were isolated (Longcore *et al.* 1999) primarily from amphibians from North America, but include strains from Africa, Australia and Panama. Due to the difficulty in obtaining and isolating the fungus, a more robust population sample was unavailable. However, our sample included strains representing all areas from which the fungus has been cultured. Strains were maintained in glass culture tubes in tryptone-glucose (TG) broth (Longcore *et al.* 1999). Stock cultures were stored at 5–6 °C and transferred at 4–5-month intervals. Additional information about the strains is available from the authors upon request.

DNA extraction

For DNA harvest, Petri-plates of TGhL agar medium (Longcore *et al.* 1999) were inoculated with 0.5–1 mL of stock culture in broth, allowed to dry in a laminar flow hood, sealed and incubated at 23 °C for ~2 weeks. Sporangia and zoospores were harvested by scraping from the plate

surface. Tissues were dehydrated, and DNA was extracted from ~20 mg dry weight, following a CTAB miniprep procedure (Zolan & Pukkila 1986).

Library construction and sequencing

An incomplete genomic library of strain JEL197 (type strain) was created for the generation of molecular markers from random DNA sequences. DNA was partially digested with *Sau3A* and 500–1500 base pairs (bp) restriction fragments were isolated from an agarose gel and ligated into plasmid pZERO (Invitrogen). Plasmids were propagated in *Escherichia coli* strains XLI-Blue (Stratagene) and INVa1F (Invitrogen). Plasmid templates for DNA sequencing were prepared with the QIAprep Spin Miniprep Kit (Qiagen) and sequenced on both strands using universal forward and reverse M13 primers. Sequencing reactions were accomplished with the BigDye sequencing kit (Applied Biosystems) and analysed on an ABI3700 DNA sequencer.

Primer design

We sequenced 33 clones from the genomic library (average insert size ~950 bp). Polymerase chain reaction (PCR) primer pairs were designed to amplify eight of the cloned DNA regions using the software package Primer3 (Rozen & Skaletsky 1997). We focused primarily on clones that had significant matches to GenBank, under the assumption that coding regions were more likely to amplify single, homologous gene regions in contrast to noncoding regions that might be redundant or repetitive. Putative gene regions and *P*-values from BLAST searches were: cysteinyl tRNA synthase (*ctsyn1*; $P = 3 \times 10^{-13}$), anthranilate phosphoribosyltransferase (*aprt13*; $P = 5 \times 10^{-6}$), and 60S ribosomal protein (*r6046*; $P = 6 \times 10^{-13}$). Other targeted gene regions matched unidentified genes (*bdc42*; $P = 10^{-12}$, *uorf48*; $P = 5 \times 10^{-5}$) or did not show significant matches (*bdc3*, *bdc33*, *rnap50*). In addition, we examined sequences in GenBank to design primers to amplify translation elongation factor 1 α (*tef1*) and the nuclear subunit ribosomal RNA gene (*lsu35*). The sequences for the 33 clones have been deposited in GenBank (accession nos BH001009–BH001047).

PCR conditions and DNA sequencing

We amplified gene regions with reaction conditions (Vilgalys & Hester 1990) that consisted of: ~1 ng template DNA, 1.5 mM MgCl₂, 0.2 mM each dNTP, 0.5 μ M each primer and 0.625 U *Taq* polymerase (Applied Biosystems) in a 25- μ L reaction. Thermal cycling parameters were an initial denaturation step at 94 °C for 3 min followed by 35 cycles of denaturing at 94 °C for 1 min, annealing at 50 °C for 30 s, extension at 72 °C for 1 min, and a 7-min final extension at 72 °C. For locus *aprt13*, the annealing temperature was raised to 55 °C. PCR

Table 1 Strain origins, hosts and genotypes at four variable loci. Strain MLST is the multilocus sequence type

Strain	Origin	Host	<i>ctsyn1</i>	<i>aprt13</i>	<i>lsu35</i>	<i>r6046</i>	MLST
JEL197	National Zoological Park, DC, USA	<i>Dendrobates azureus</i>	(A/G)*	(A/G)†	(G)‡	(CA/-)§	1
JEL198	National Zoological Park, DC, USA	<i>Dendrobates auratus</i>	A/G	A/G	G	CA/-	1
JEL213	Mono Co., California, USA	<i>Rana muscosa</i>	A/G	A/G	G	CA/-	1
JEL214	Mono Co., California, USA	<i>Rana muscosa</i>	A/G	A/G	G	CA/-	1
JEL215	Mono Co., California, USA	<i>Rana muscosa</i>	A/G	A/G	G	CA/-	1
JEL226	Yavapai Co., Arizona, USA	<i>Rana yavapaiensis</i>	A/G	A/G	G	CA/-	1
JEL229	Montrose Canyon, Arizona, USA	<i>Hyla arenicolor</i>	A/G	A/G	G	CA/-	1
JEL230	Montrose Canyon, Arizona, USA	<i>Rana yavapaiensis</i>	A/G	A/G	G	CA/-	1
JEL260	Quebec, Canada	<i>Rana catesbeiana</i>	A/G	A/G	G	CA/-	1
JEL262	Quebec, Canada	<i>Rana catesbeiana</i>	A/G	A/G	G	CA/-	1
JEL264	Quebec, Canada	<i>Rana catesbeiana</i>	A/G	A/G	G	CA/-	1
JEL277	Arizona, USA	<i>Ambystoma tigrinum</i>	A/G	A/G	G	CA/-	1
JEL284	Wisconsin, USA (captive)	<i>Rana pipiens</i>	A/G	A/G	G	CA	2
JEL289	Milford, Maine, USA	<i>Rana pipiens</i>	A/G	A/G	G	CA	2
JEL231	Mesquite Wash, Arizona, USA	<i>Rana yavapaiensis</i>	A/G	A/G	A/G	CA	3
JEL254	Orono, Maine, USA	<i>Rana pipiens</i>	A/G	A/G	A/G	CA/-	4
JEL258	Orono, Maine, USA	<i>Rana sylvatica</i>	A/G	A/G	A/G	CA/-	4
JEL203	Bronx Zoo, New York, USA	<i>Dyscophus guineti</i>	A/G	A/G	A	-	5
JEL270	Point Reyes, California, USA	<i>Rana catesbeiana</i>	A/G	A/G	A	-	5
JEL271	Point Reyes, California, USA	<i>Rana catesbeiana</i>	A/G	A/G	A	-	5
JEL282	Toledo Zoo, Ohio, USA	<i>Bufo americana</i>	A/G	A/G	A	-	5
00 545	Melbourne, Victoria, Australia	<i>Litoria lesueuri</i>	A/G	A/G	A/G	-	6
98 1810 3	Tully, Queensland, Australia	<i>Nyctimystes dayi</i>	A/G	A/G	A/G	-	6
99 1385 12	Rockhampton, Queensland, Australia	<i>Litoria caerulea</i>	A/G	A/G	A/G	-	6
PM5	Panama	<i>Smilisca phaeota</i>	A/G	A/G	A/G	-	6
PM7	Panama	<i>Smilisca phaeota</i>	A/G	A/G	A/G	-	6
JEL253	Melbourne, Victoria, Australia	<i>Limnodynastes dumerilii</i>	G	A/G	A/G	-	7
JEL225	Wisconsin, USA (captive)	<i>Silurana (Xenopus) tropicalis</i>	G	A/G	G	-	8
JEL239	Imported from Ghana	<i>Silurana (Xenopus) tropicalis</i>	A/G	A/G	G	-	9
JEL240	Imported from Ghana	<i>Silurana (Xenopus) tropicalis</i>	A/G	A/G	G	-	9
JEL245	Imported from Ghana	<i>Silurana (Xenopus) tropicalis</i>	A/G	A/G	G	-	9
JEL273	Clear Creek Co., Colorado, USA	<i>Bufo boreas</i>	A/G	A/G	G	-	9
JEL275	Clear Creek Co., Colorado, USA	<i>Bufo boreas</i>	A/G	A/G	G	-	9
PM1	Panama	<i>Eleuthodactylus caryophyllaceum</i>	A/G	A/G	G	-	9
JEL274	Clear Creek Co., Colorado, USA	<i>Bufo boreas</i>	A	A/G	G	-	10

*Genotype at base 400; A/G (adenine and guanine) indicates heterozygote at this position; GenBank accession no. = BH001044.

†Genotype at base 679; GenBank accession no. = BH001045.

‡Genotype at base 315; GenBank accession no. = BH001046.

§Genotype between bases 499–500; CA = two basepair insertion, - = strains that lack the insertion, CA/- = heterozygote at this position; GenBank accession no. = BH001047.

amplicons were purified with ULTRAFree-MC centrifugal columns (Millipore Corp.), except for *aprt13*, which was purified from an agarose gel with a QIAquick Gel Extraction Kit (Qiagen). Amplicons were labelled with the BigDye kit and electrophoresed on an ABI3700. The following primers were used for PCR amplification (convention used is locus name followed by F and R designations for forward and reverse primers): *ctsyn1F* (5'-ACCAACTATAACATCATCAAG-3'), *ctsyn1R* (5'-CGAATATCAGTCAACGCAAGC-3'), *aprt13F* (5'-GTCAGGGTTGGCTATTGTTCT-3'), *aprt13R* (5'-TGCTACTATTGCTGCTGTTGC-3'), *lsu35F* (5'-ATC-CCTGTGGTAACTTTTCTG-3'), *lsu35R* (5'-ACGGACAT-

GGGGAATCTGACT-3'), *r6046F* (5'-CTATCTGCGCTCCC-GTGTCAA-3'), *r6046R* (5'-AGGGCTGCAACAACCTGGA-TTT-3'), *uorf48F* (5'-TCGAGGTGCAGACAAAACCTC-3'), *uorf48R* (5'-CAAACCTGAGCCACAATAATGC-3'), *rnaf50F* (5'-AATCCTATCCACCAGTTTCAG-3'), *rnaf50R* (5'-TAA-CGATGAACGCCTTGTAGA-3'), *bdc3F* (5'-TTCTGCTGC-AAGAATCATCG-3'), *bdc3R* (5'-AGTAGAAGCGGGTCCG-TTGAA-3'), *bdc33F* (5'-ATAGACCTTCGGGCTCTGGT-3'), *bdc33R* (5'-TTTCGTGTTAACCCAAAGGC-3'), *bdc42F* (5'-GGCCAACCTGTTGGATTTGT-3'), *bdc42R* (5'-TTGGAGC-TCTGGTTCGACTT-3'), *tef1F* (5'-TACAARTGYGGTGGTA-TYGACA-3'), *tef1R* (5'-ACNGACTTGACYTCAGTRGT-3').

Southern hybridization

Southern capillary transfers were carried out according to Ausubel *et al.* (1998). Probes were labelled with digoxigenin by PCR amplification (Roche). DNA hybridizations were performed in DIG Easy Hyb buffer (Roche) at 42 °C, with post-hybridization washes (65 °C) and chemiluminescent detection as per the manufacturer's instructions (Roche).

Pulsed field gel electrophoresis

Chromosomal DNA was prepared by a modification of the method of Iadonato & Gnirke (1996). Briefly, zoospores were harvested from plate cultures by flooding with water, centrifuged, washed in 50 mM EDTA (pH 8.0) and mixed with an equal volume of 2% LMP agarose to give 10⁹ zoospores/mL. Plugs were cast and incubated in spheroplasting solution containing 1 M sorbitol, 20 mM EDTA (pH 8.0), 10 mM Tris-HCl (pH 7.5), 14 mM β-mercaptoethanol, 2 mg/mL Zymolase (Seikogaku) and 5 mg/mL lysing enzyme (Sigma) at 37 °C for 5 h. Plugs were incubated in LDS buffer (100 mM EDTA, 10 mM Tris-HCl, 1% sodium dodecyl chloride, pH 8.0) at 37 °C overnight. Three washes for 30 min each at RT in NDS buffer (0.1 M EDTA, 2 mM Tris-HCl, 0.2% Sarkosyl, pH 8.0) were followed by five 30-min washes in TE buffer. Chromosomes were separated with the CHEF DR-II pulsed field gel system (Bio-Rad) on a 1% agarose gel with the following parameters: 5.0 V/cm with a 100–200-s ramped switch time for 24 h followed by 2.9 V/cm with a 190–300-s ramped switch time for 24 h, all in 0.5 × TBE buffer at 10 °C.

Data analysis

A tree-like network connecting multilocus genotypes was calculated with statistical parsimony using the software package *tcs* v1.13 (Clement *et al.* 2000). The pairwise distance matrix between isolates was calculated by considering each locus as a separate character and the three possible genotypes at each locus as unique character states. The distance between the two homozygous genotypes at a locus was assumed to be two steps, and heterozygous genotypes were assumed to be one step from each homozygous type.

Results

Nucleotide variation

We screened for sequence variation in *B. dendrobatidis* by PCR/DNA sequencing of 10 gene regions (eight anonymous loci *lsu35* and *tef1*). Direct sequencing of PCR amplicons from 35 strains revealed few polymorphic sites among the 10 loci (5 variable sites in 5918 nucleotides). Six of the 10 loci

showed zero nucleotide substitutions among all 35 samples (*bdc3*, *bdc33*, *bdc42*, *uorf48*, *rnap50*, *tef1*; 3551 nucleotides). The four other loci contained one or two variable sites. More than one allele was present in many strains. A site with a double-peak in the sequence chromatogram indicated this heterogeneity within a PCR product. These double-peaks are referred to hereafter as 'heterozygous sites'; the justification for this terminology is given below. We found five heterozygous sites: two at *ctsyn1*, and one each at *aprt13*, *lsu35* and *r6046*. Four of the five nucleotide heterozygosities are A ↔ G transitions, none of which are predicted to cause a change in amino acid sequence in the most probable translations. The fifth heterozygosity, at locus *r6046*, is a 2-bp indel in a noncoding region. The five variable positions displayed both heterozygous and homozygous genotypes, and the combination of genotypes over the four variable loci formed 10 multilocus sequence types (MLSTs), shown in Table 1.

The frequency of heterozygous genotypes in the population sample varied but was as high as 100% (*aprt13*). At locus *ctsyn1*, 32 of the 35 strains were heterozygous (~91%). Two individuals of one homozygous type and one individual of the other homozygous type were found (Fig. 1). At loci *r6046* and *lsu35*, both types of homozygotes were detected and heterozygous genotypes were in the minority, 40% for *r6046* and ~26% for *lsu35*.

We confirmed that the two alleles at the putatively heterozygous loci were present in the amplified PCR products rather than being artefacts of the fluorescent DNA sequencing procedure by subcloning and sequencing PCR products from one or two strains for each of the four loci. Each subclone (*n* = 8–12) possessed separately one of the two possible alleles at each polymorphic site, and each PCR product showed segregation of the polymorphism among subclones, suggesting that the *B. dendrobatidis* strains possess two different alleles in their genomes. Additionally,




Fig. 1 Digestion of *ctsyn1* PCR amplicons with the restriction enzyme *Fnu4HI* reveals heterozygosity for most individuals at this locus. *Fnu4HI* cuts the 551-bp amplicon of one allele (homozygote in lane 1) into 227-bp, 171-bp, 115-bp and 38-bp fragments. Substitution of A for G in the other allele creates an additional *Fnu4HI* site in the 171-bp fragment, generating 133-bp and 38-bp fragments (homozygotes in lanes 8 and 9). Individuals in lanes 2–7 are heterozygous at this locus for the *Fnu4HI* RFLP. Lanes 1–9 are: JEL274, JEL197, JEL213, JEL230, JEL264, JEL284, JEL258, JEL225, JEL253. Reverse image of a 3% agarose gel stained with ethidium bromide is shown.




Fig. 2 Southern hybridization of genomic DNA digests of three *B. dendrobatidis* strains to a probe for locus *r6046*. In each lane, a single hybridization signal is observed. Numbers above lane refer to strain number, and the restriction endonucleases used are indicated.

restriction digests of some PCR amplicons demonstrated the presence of two alleles at the polymorphic loci. For example, locus *ctsyn1* displayed a restriction fragment length polymorphism (RFLP) and digestion of these amplicons with the enzyme *Fnu4HI* showed that whereas most strains possessed two alleles, strains with only one of the two alleles were also present, confirming genotype assignments based on DNA sequencing (Fig. 1).

We confirmed that these loci are present in a single copy in the genome by hybridizing probes of the polymorphic loci to DNA digested with three different restriction endonucleases. With the exception of locus *Isu35*, only a single band hybridized to these probes under stringent conditions (Fig. 2). For locus *Isu35*, the large subunit rDNA gene, additional, weaker bands were observed with most endonucleases. These data suggest that the presence of

two alleles within a strain is probably a result of heterozygosity at a single locus and, therefore, *B. dendrobatidis* is likely of diploid or higher ploidy.

Population structure

We analysed genetic relationships among strains with statistical parsimony to determine whether the MLST data reveal any evidence for population substructure (Fig. 3). This network groups strains with related MLSTs such that each branch represents a change from a heterozygous state to a homozygous one at a single locus. In general, strains collected from the same locale displayed the same MLST (Table 1 and Fig. 3). For example, strains isolated from *Rana muscosa* from the Sierras in California are in MLST no.1 (JEL 213, 214 and 215). The other two California strains, which were isolated from *R. catesbeiana* collected from Point Reyes, are both in MLST no. 5. The strains from Africa, Australia, Panama and Colorado also demonstrate geographical relatedness. In each example, the strains within each geographical region are identical or differ from each other by the loss of heterozygosity at only a single locus (i.e. are connected by single branches in Fig. 3). Interestingly, MLST no. 6 is comprised of two Panamanian strains and three Australian strains, showing high genetic relatedness of certain strains from two areas for which chytridiomycosis has been implicated in amphibian declines (Berger *et al.* 1998). In general, the MLSTs show no clear global pattern of genotype distribution. However, the most frequently collected genotype in North America (MLST no. 1) was not recovered outside the continent. Moreover, the allele at locus *r6046* containing the 2-bp insertion ('CA' in Table 1) has never been recorded outside




Fig. 3 Depiction of MLSTs as a network of related genotypes. Isolates with the same MLST are grouped into abutting boxes and labelled according to MLST numbers in Table 1. Each branch interconnecting groups represents a change in genotype at a single locus. All changes can be regarded as the change from a heterozygous to a homozygous state. *Strains that are heterozygous at all four loci. Network estimated using the statistical parsimony algorithm as implemented in the TCS software package (Clement *et al.* 2000).




Fig. 4 Separation of *B. dendrobatidis* chromosomes with CHEF gel electrophoresis shows multiple chromosome length polymorphisms. Lane 1 = *Saccharomyces cerevisiae* standard; lane 2 = JEL197; lane 3 = JEL245; lane 4 = JEL253; lane 5 = JEL289. Sizes of the largest *S. cerevisiae* chromosomes are indicated to the left of the photograph.

North America, wherein its frequency is 58%. Genotype frequencies at *r6046* are significantly different ($P < 0.001$; χ^2 test) between North America isolates vs. isolates collected from outside North America. This result is also observed at *lsu35* ($P < 0.05$).

Chromosome polymorphism

We karyotyped several *B. dendrobatidis* strains with CHEF gel electrophoresis. Chromosomes ranged in size from ~0.7–6.0 Mb, and numerous chromosome length polymorphisms existed among strains (Fig. 4). The estimated genome size is 35–40 Mb, although this estimate is difficult as the chromosomal bands stained at different intensities. If *B. dendrobatidis* is diploid, the intensely staining chromosomal bands may reflect comigrating homologous chromosomes, and length polymorphisms may reflect homologous chromosomes that vary in size, such as reported for *Candida albicans* (Iwaguchi *et al.* 1990). In this case, the genome size estimate would be between the true haploid and diploid genome size.

Discussion

This study demonstrated a low level of genetic variation within an intercontinental sample of *B. dendrobatidis* strains. Among the 10 loci surveyed only five variable sites were detected. For each of the loci that demonstrated variation heterozygous isolates were detected, and at two of the loci the majority of strains were heterozygous. From these data we conclude: our strains of *B. dendrobatidis* have the population genetic signature of a newly emerged

pathogen, the fungus is probably diploid and reproduction is primarily clonal.

The low level of genetic variation among strains of *B. dendrobatidis* from North America, Central America, Africa and Australia implies that the coalescence time of the entire sample is relatively recent and that the fungus has become dispersed geographically over a time scale during which few mutations have occurred. Comparable data from other fungal pathogens are not extensive. Similar to *B. dendrobatidis*, other pathogenic fungal species such as *Trichophyton rubrum* (Gräser *et al.* 1999) and *Fusarium oxysporum* f. sp. *ciceris* (Jiménez-Gasco *et al.* 2002) have demonstrated an absence of nucleotide polymorphism among several nuclear loci. Nonetheless, population studies with other fungi generally reveal much greater genetic variation than that observed here (e.g. Forche *et al.* 1999; O'Donnell *et al.* 2000; Xu *et al.* 2000).

Low levels of polymorphism suggest at least four possible, but not mutually exclusive, explanations: (1) the global effective population size in *B. dendrobatidis* is small; (2) mutation rates in *B. dendrobatidis* are low; (3) there has been a recent bottleneck in population size; or (4) the species has recently emerged from a genotype(s) favoured by natural selection. In dismissal of point one, the generally broad distribution and host range of the fungus do not suggest a small effective population size. Additionally, with no a priori evidence to suggest that mutation rates in *B. dendrobatidis* are extraordinarily low (point 2), it seems reasonable to conclude that either a recent bottleneck occurred in the history of the fungus or that the population has recently emerged from a single, founder strain (points 3 and 4). In a clonal population, a recent population bottleneck would be expected to have a similar population genetic signature as the emergence of a virulent strain caused by strong selection, because such selective pressure should affect the whole genome of nonrecombining organisms through genetic hitchhiking.

An analysis of genetic relationships among strains showed low overall geographical structuring and host specificity of genotypes (Fig. 3). These data do not support the hypothesis that *B. dendrobatidis* has emerged from a pre-existing relationship between the fungus and amphibians via recent climatic change or other abiotic factors, and are more consistent with the recent introduction of the pathogen into naïve populations. If the relationship between frog and fungus had been established for many generations in the absence of gene flow, the MLST data would be predicted to show geographical population structure. Indeed, there does seem to be some geographical structuring of populations as evidenced by a significant genetic differentiation between North America and the rest of the world at loci *r6046* and *lsu35*. These loci suggest the North American population may form a distinct gene pool. In contrast, the finding of identical MLSTs in Panama and

Australia argues for recent dispersal or introduction of the pathogen into these two areas of amphibian population declines. *B. dendrobatidis* displays other characteristics of a virulent pathogen introduced into a naïve population, such as low host specificity and high mortality (Daszak *et al.* 1999). As with some human pathogens, [e.g. the agent of plague, *Yersinia pestis* (Achtman *et al.* 1999) and the fungal skin pathogen *Trichophyton rubrum* (Gräser *et al.* (1999)], low levels of intercontinental genetic variation in *B. dendrobatidis* could be due to recent dissemination of the disease by humans.

Despite the recent geographical expansion of the fungus, both asymptomatic and symptomatic frogs can share the same genotypes. Also, regions where populations are declining display nearly the same genotypes as regions where the fungus has not caused obvious population declines (e.g. eastern North America). This supports the argument that population declines are context-dependent (Blaustein & Kiesecker 2002), i.e. the disease interacts with both frog species and the environment in unique ways in different geographical areas.

The MLST data were also used to understand the mode of *B. dendrobatidis* reproduction. The detection of two loci for which frequencies of heterozygous individuals were either fixed or nearly fixed suggests that *B. dendrobatidis* has a predominately clonal population structure (Tibayrenc *et al.* 1990). Fixed heterozygosity arises from the absence of normal segregation of homologues at meiosis that breaks apart heterozygous genotypes.

Although we interpreted the double-peaked sites in the sequence chromatograms as heterozygous sites, we addressed alternative explanations for the presence of two sequence copies within a single strain. One possibility is that the strains, which are presumably single individuals, are actually a population of individuals that differ in alleles at the polymorphic loci (i.e. represent 'mixed' cultures). This can be dismissed because we sequenced the 10 loci from a strain derived from a single zoospore (JEL270), yet two of its 10 loci were heterozygous. Zoospores possess only a single nucleus (Longcore *et al.* 1999), therefore this strain derives from a single genetic individual. Double-peaked sites could also arise from the amplification of two paralogous loci that differ in sequence. This is obviously a possibility for ribosomal RNA gene *lsu35*. However, the hybridizing of probes of three polymorphic loci to single bands in Southern analyses suggests that these loci are present in a single copy in the *B. dendrobatidis* genome. Moreover, the low level of nucleotide variation between the two sequence types within a strain, approximately one silent polymorphism per 475-bp, suggests that the two sequence types are allelic rather than paralogous. These data strongly suggest that the observed double-peaked sites in the chromatograms are the result of heterozygosity; however, neither the chromosomal nor the sequence poly-

morphism data exclude the possibility that *B. dendrobatidis* is of higher than diploid ploidy. Hybridization of probes for the polymorphic loci to CHEF separated chromosomes or *in situ* chromosomal preparations are needed to more adequately determine ploidy.

In diploid organisms, all new point mutations arise in a heterozygous state. Homozygotes for the new mutation are created through sexuality or processes such as mitotic gene conversion or crossing over (collectively termed somatic recombination). A conflict arises because certain *B. dendrobatidis* loci suggest clonality through the extreme overrepresentation of heterozygous genotypes (*ctsyn1* and *aprt13*), whereas others suggest segregation/recombination by displaying appreciable frequencies of both homozygous types (*lsu35* and *r6046*). Such a pattern might emerge if the process of generating homozygous genotypes is a locus or chromosome-specific phenomenon and not a genome-wide phenomenon. Somatic recombination happens during vegetative growth and can affect single gene regions or chromosomes. Conversely, meiosis is a genome-wide recombination process that shuffles alleles as well as chromosomes. We suggest that the presence of both possible homozygous genotypes at a locus has arisen from somatic recombination within heterozygous progenitors. The difference between levels of heterozygosity at the four variable loci could be explained by differential rates of somatic recombination, perhaps as a result of the proximity to the centromere. For example loci *ctsyn1* and *aprt13*, which display fixed heterozygosity, may be near the centromere where recombination is inhibited. That somatic recombination may play a significant role in generating novel multilocus genotypes in *B. dendrobatidis* is supported by evidence of recombination in other fungi that are believed to reproduce exclusively asexually, such as *Coccidioides immitis* and *Aspergillus flavus* (Burt *et al.* 1996; Geiser *et al.* 1998).

B. dendrobatidis displayed extensive chromosome length polymorphism among strains as has been observed in other fungal species (Zolan 1995). Chromosome length polymorphism in ancient asexual organisms is consistent with the expectation that large length differences between homologues could hinder proper pairing at meiosis or result in nondisjunction (Birky 1996; Welch & Meselson 1998). Despite this, because both sexual and asexual fungi frequently have chromosome length polymorphisms, our documentation of chromosome length polymorphism in *B. dendrobatidis* does not lead directly to the conclusion that the fungus is primarily asexual (Zolan 1995). The high level of genetic variation suggested by chromosome polymorphisms contrasts with the near absence nucleotide variation. The elevated level of chromosome rearrangement in *B. dendrobatidis* may be related to the processes governing somatic recombination.

The finding of a largely clonal population structure suggests that *B. dendrobatidis* may lack a sexual stage,

therefore it is possible that a resting stage is absent. Predominately clonal reproduction in *B. dendrobatidis* would not be surprising because sexuality has not been observed in many chytridiales fungi, and has not been reported for any members of the Spizellomycetales or Neocallimastigales (Barr 2001). Although lack of sexuality does not preclude the generation of asexually produced resting spores, the only propagules that have been observed are zoospores, which cannot survive desiccation or marine travel. That drying can kill cultures of *B. dendrobatidis* (Berger *et al.* 1999) also supports the absence of a resting spore. An absence of a resting spore stage would suggest that natural causes such as airborne dispersal of spores are less likely to be the reason for the pathogen's entry into naïve populations than inadvertent dispersal by humans or perhaps other long distance travelers such as birds.

The MLST data are limiting because they do not provide high resolution of genetic relationships among the strains. In addition, our sampling is incomplete because it does not cover the entire range of the disease, and it is biased towards North American strains. Further work with more variable markers, such as microsatellite loci or amplified length fragment polymorphisms, and a more geographically diverse set of strains may be able to distinguish more genotypes than the MLST data and provide a better understanding of the origins of the disease.

Acknowledgements

This work would have not been possible without the gracious help of those who provided infected frogs from which fungal strains were isolated: Gregory A. Bradley, John Clarke, Elizabeth W. Davidson, Gary M. Fellers, Alastair Freeman, Mark S. Jones, Thomas R. Jones, Michael J. Linn, Jerry R. Longcore, Gerry Marantelli, Keith McDonald, Igor Mikaelian, Donald K. Nichols, Linda Northey, Martin Ouellet, Allan P. Pessier, Timothy Reichard, Michael J. Sredl, Benjamin Stubbs and Kathy Taylor. We thank Sudhir Kumar, Cynthia Riginos and two anonymous reviewers for reviews of earlier drafts, and Thomas G. Mitchell for generous sharing of laboratory space. This research was supported by the Howard Hughes Precollege Program, by IRCEB grant IBN-9977063 from the National Science Foundation (James P. Collins, PI), and a Clark Fellowship in Molecular Evolution to AG.

References

Achtman M, Zurth K, Morelli G, Torrea G, Guiryoule A, Carniel E (1999) *Yersinia pestis*, the cause of plague, is a recently emerged clone of *Yersinia pseudotuberculosis*. *Proceedings of the National Academy of Sciences USA*, **96**, 14043–14048.

Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K (1998) *Current Protocols in Molecular Biology*. Greene and Wiley Interscience, New York.

Barr DJS (2001) *Chytridiomycota*. In: *The Mycota VII. Part A, Systematics and Evolution* (eds McLaughlin DJ, McLaughlin EG, Lemke PA), pp. 93–112. Springer-Verlag, Berlin.

Berger L, Speare R, Daszak P, Green DE *et al.* (1998) Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proceedings of the National Academy of Sciences USA*, **95**, 9031–9036.

Berger L, Speare R, Hyatt AD (1999) Chytrid fungi and amphibian declines: overview, implications and future directions. In: *Declines and Disappearances of Australian Frogs* (ed. Campbell A), pp. 23–33. Environment Australia, Canberra.

Birky CW Jr (1996) Heterozygosity, heteromorphy, and phylogenetic trees in asexual eukaryotes. *Genetics*, **144**, 427–437.

Blaustein AR, Kiesecker JM (2002) Complexity in conservation: lessons from the global Decline of amphibian populations. *Ecology Letters*, **5**, 597–608.

Bosch J, Martínez-Solano I, García-París M (2001) Evidence of a chytrid fungus infection involved in the decline of the common midwife toad (*Alytes obstetricans*) in protected areas of central Spain. *Biological Conservation*, **97**, 331–337.

Bradley GA, Rosen PC, Sredl MJ, Jones TR, Longcore JE (2002) Chytridiomycosis in native Arizona frogs. *Journal of Wildlife Diseases*, **38**, 206–212.

Burt A, Carter DA, Koenig GL, White TJ, Taylor JW (1996) Molecular markers reveal cryptic sex in the human pathogen *Coccidioides immitis*. *Proceedings of the National Academy of Sciences USA*, **93**, 770–773.

Carey C, Cohen N, Rollins-Smith LA (1999) Amphibian declines: an immunological perspective. *Developmental and Comparative Immunology*, **23**, 459–472.

Clement M, Posada D, Crandall KA (2000) TCS: a computer program to estimate gene genealogies. *Molecular Ecology*, **9**, 1657–1660.

Daszak P, Berger L, Cunningham AA, Hyatt AD, Green DE, Speare R (1999) Emerging infectious diseases and amphibian population declines. *Emerging Infectious Diseases*, **5**, 735–748.

Daszak P, Cunningham AA, Hyatt AD (2000) Emerging infectious diseases of wildlife — threats to biodiversity and human health. *Science*, **287**, 443–449.

Forche A, Schönian G, Gräser Y, Vilgalys R, Mitchell TG (1999) Genetic structure of typical and atypical populations of *Candida albicans* from Africa. *Fungal Genetics and Biology*, **28**, 107–125.

Geiser DM, Pitt JI, Taylor JW (1998) Cryptic speciation and recombination in the aflatoxin-producing fungus *Aspergillus flavus*. *Proceedings of the National Academy of Sciences USA*, **95**, 388–393.

Gräser Y, Kühnisch J, Presber W (1999) Molecular markers reveal exclusively clonal reproduction in *Trichophyton rubrum*. *Journal of Clinical Microbiology*, **37**, 3713–3717.

Iadonato SP, Gnirke A (1996) RARE — cleavage analysis of Yacc. In: *Methods in Molecular Biology V. 54: YAC Protocols* (ed. Markie D), pp. 75–85. Humana Press, Inc, Totowa.

Iwaguchi S-I, Homma M, Tanaka K (1990) Variation in the electrophoretic karyotype analysed by the assignment of DNA probes in *Candida albicans*. *Journal of General Microbiology*, **136**, 2433–2442.

Jiménez-Gasco MM, Milgroom MG, Jiménez-Díaz RM (2002) Gene genealogies support *Fusarium oxysporum* f. sp. *ciceris* as a monophyletic group. *Plant Pathology*, **51**, 72–77.

Lips KR (1999) Mass mortality and population declines of anurans at an upland site in western Panama. *Conservation Biology*, **13**, 117–125.

Longcore JE, Pessier AP, Nichols DK (1999) *Batrachochytrium dendrobatidis* gen et sp. nov., a chytrid pathogenic to amphibians. *Mycologia*, **91**, 219–227.

- Nichols DK, Lamirande EW, Pessier AP, Longcore JE (2001) Experimental transmission of cutaneous chytridiomycosis in dendrobatid frogs. *Journal of Wildlife Diseases*, **37**, 1–11.
- O'Donnell K, Kistler HC, Tacke BK, Casper HH (2000) Gene genealogies reveal global phylogeographic structure and reproductive isolation among lineages of *Fusarium graminearum*, the fungus causing wheat scab. *Proceedings of the National Academy of Sciences USA*, **97**, 7905–7910.
- Pessier AP, Nichols DK, Longcore JE, Fuller MS (1999) Cutaneous chytridiomycosis in poison dart frogs (*Dendrobates* spp.) and White's tree frogs (*Litoria caerulea*). *Journal of Veterinary Diagnostic Investigation*, **11**, 194–199.
- Rozen S, Skaletsky HJ (1997) Primer3. Code available at http://www-genome.wi.mit.edu/genome_software/other/primer3.html.
- Sparrow FK (1960) *Aquatic Phycomycetes*, 2nd rev. edn. University of Michigan Press, Ann Arbor.
- Speare R, Core Working Group of Getting the Jump on Amphibian Disease (2001) Nomination for listing of amphibian chytridiomycosis as a key threatening process under the Environment Protection and Biodiversity Conservation Act 1999. In: *Developing Management Strategies to Control Amphibian Diseases: Decreasing the Risks Due to Communicable Diseases* (eds Speare R, Steering Committee of Getting the Jump on Amphibian Disease), pp. 186–208. School of Public Health and Tropical Medicine, James Cook University, Townsville.
- Sreevatsan S, Pan X, Stockbauer KE *et al.* (1997) Restricted structural gene polymorphism in the *Mycobacterium tuberculosis* complex indicates evolutionarily recent global dissemination. *Proceedings of the National Academy of Sciences USA*, **94**, 9869–9874.
- Tibayrenc M, Kjellberg F, Ayala FJ (1990) A clonal theory of parasitic protozoa: the population structures of *Entamoeba*, *Giardia*, *Leishmania*, *Naegleria*, *Plasmodium*, *Trichomonas*, and *Trypanosoma* and their medical and taxonomical consequences. *Proceedings of the National Academy of Sciences USA*, **87**, 2414–2418.
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* spp. *Journal of Bacteriology*, **172**, 4238–4246.
- Welch JLM, Meselson M (1998) Karyotypes of bdelloid rotifers from three families. *Hydrobiologia*, **387/388**, 403–407.
- Xu J, Vilgalys R, Mitchell TG (2000) Multiple gene genealogies reveal recent dispersion and hybridization in the human pathogenic fungus *Cryptococcus neoformans*. *Molecular Ecology*, **9**, 1471–1481.
- Zolan ME (1995) Chromosome-length polymorphism in fungi. *Microbiological Reviews*, **59**, 686–698.
- Zolan ME, Pukkila PJ (1986) Inheritance of DNA methylation in *Coprinus cinereus*. *Molecular and Cellular Biology*, **6**, 195–200.

This investigation is part of the ongoing, multi-institutional effort at understanding the role of disease in global amphibian declines. E. A. Morehouse, T. Y. James, A. R. D. Ganley and R. Vilgalys are members of the Duke Mycology Research Unit and are dedicated to understanding the evolutionary genetics of pathogenic and saprobic fungi. P. J. Murphy's ecological interests include the responses of amphibians to variable environments, including unpredictable climate and novel pathogens. L. Berger is an amphibian pathologist who works on chytridiomycosis in Australia. J. E. Longcore is dedicated to all aspects of chytrid fungi.
