



Multiple Phenotypic Changes Define Neutrophil Priming

Irina Miralda¹, Silvia M. Uriarte^{1,2} and Kenneth R. McLeish^{2,3*}

¹ Department of Microbiology, University of Louisville School of Medicine, Louisville, KY, United States, ² Department of Medicine, University of Louisville School of Medicine, Louisville, KY, United States, ³ Robley Rex VA Medical Center, Louisville, KY, United States

Exposure to pro-inflammatory cytokines, chemokines, mitochondrial contents, and bacterial and viral products induces neutrophils to transition from a basal state into a primed one, which is currently defined as an enhanced response to activating stimuli. Although, typically associated with enhanced generation of reactive oxygen species (ROS) by the NADPH oxidase, primed neutrophils show enhanced responsiveness of exocytosis, NET formation, and chemotaxis. Phenotypic changes associated with priming also include activation of a subset of functions, including adhesion, transcription, metabolism, and rate of apoptosis. This review summarizes the breadth of phenotypic changes associated with priming and reviews current knowledge of the molecular mechanisms behind those changes. We conclude that the current definition of priming is too restrictive. Priming represents a combination of enhanced responsiveness and activated functions that regulate both adaptive and innate immune responses.

Keywords: neutrophils, priming, cytokines, chemotaxis, apoptosis, phagocytosis, respiratory burst, exocytosis

OPEN ACCESS

Edited by:

Joao Santana Silva,
University of São Paulo, Brazil

Reviewed by:

George Hajishengallis,
University of Pennsylvania,
United States

Henrique Serezani,
Vanderbilt University Medical Center,
United States

*Correspondence:

Kenneth R. McLeish
k.mcleish@louisville.edu

Received: 20 February 2017

Accepted: 11 May 2017

Published: 29 May 2017


Citation:

Miralda I, Uriarte SM and McLeish KR
(2017) Multiple Phenotypic Changes
Define Neutrophil Priming.
Front. Cell. Infect. Microbiol. 7:217.
doi: 10.3389/fcimb.2017.00217

INTRODUCTION

Polymorphonuclear leukocytes, or neutrophils, account for 40–60% of peripheral blood leukocytes in humans (Summers et al., 2010). They play an essential role in the innate immune response, as demonstrated by the development of life-threatening infections or uncontrolled inflammation in individuals with severe neutropenia or genetic disruption of neutrophil anti-microbial capabilities (Kannengiesser et al., 2008; van de Vijver et al., 2012; Moutsopoulos et al., 2014; Nauseef and Borregaard, 2014). **Figure 1** shows the multistep process of neutrophil recruitment in response to microbial invasion, including adhesion to vascular endothelium, transmigration into the interstitial space, chemotaxis/chemokinesis toward the site of infection, phagocytosis of pathogens, destruction of microbes within phagosomes by release of antimicrobial granule contents following granule fusion and ROS generation at the phagosomal membrane, and amplification and organization of the inflammatory response. Uncontrolled or prolonged neutrophil activation uses antimicrobial responses to injure normal host cells, leading to pathologic changes to tissues and organs in autoimmune and inflammatory diseases (Nathan, 2006). Consequently, neutrophil activation is normally tightly regulated.

Circulating neutrophils exist in a basal state, characterized by non-adherence, a round morphology, minimal transcriptional activity, and a limited capacity to respond to activating stimuli. That limited response protects against unwarranted inflammatory responses and tissue injury (Sheppard et al., 2005). To effectively clear invading organisms, neutrophils must be capable of mounting rapid, vigorous responses to activating stimuli. The transition to a state of enhanced



responsiveness has been termed priming (Condliffe et al., 1998; El-Benna et al., 2008; Wright et al., 2013). It occurs *in vitro* following neutrophil exposure to pro-inflammatory lipids and cytokines, chemokines, mitochondrial contents, and bacterial and viral products (El-Benna et al., 2008). Neutrophil priming *in vitro* represents an *in vivo* phenomena, as primed neutrophils have been identified in humans with infections, rheumatoid arthritis, chronic kidney disease, traumatic injury, and acute respiratory distress syndrome (Bass et al., 1986; McLeish et al., 1996; Ogura et al., 1999; Naegel et al., 2012). Although, substantial circumstantial evidence suggests that primed neutrophils participate in a number of human diseases, direct evidence is lacking. The relative contribution of neutrophil priming to the severity of human inflammatory diseases is an important gap in knowledge that needs to be addressed.

Historically, the term “priming” was primarily used to describe the augmented reactive oxygen species (ROS) generation upon neutrophil stimulation because of the depth of knowledge of molecular mechanisms of NADPH oxidase complex assembly, the ease of measurement of ROS generation, and the importance of ROS to anti-microbial activity. **Figure 1**

illustrates that primed neutrophils demonstrate a number of phenotypic changes in addition to enhanced NADPH oxidase activation, including granule release, cytokine and lipid synthesis, adhesion and transmigration, enhanced chemotaxis, and delayed apoptosis. Thus, neutrophil priming is not just a transition state in which neutrophils become more responsive to activating stimuli. We believe a new definition of priming is required to include the activation of a subset of neutrophil functions as opposed solely to a heightened state of responsiveness. In this review of the recent advances in neutrophil priming, we will highlight the functional evidence for the activation of a subset of neutrophil functions during priming and review the current state of knowledge of the molecular basis for those phenotypic changes to illustrate this new definition. Our goal is to encourage research that will provide a more complete understanding of priming, leading to identification of new targets for treatment of inflammatory and infectious diseases. Much of our discussion focuses on the effects of TNF α , as studies frequently use that cytokine as a model priming agent. The large number of agents capable of initiating priming of neutrophil respiratory burst activity was recently reviewed (El-Benna et al., 2016). We compare current state of knowledge of the effects of

those priming agents on the various phenotypic changes to those induced by TNF α in **Table 1**.

PHENOTYPIC CHANGES DURING PRIMING

Respiratory Burst Activity

For decades, enhanced respiratory burst activity has defined a primed neutrophil. The respiratory burst generates ROS through conversion of molecular oxygen to superoxide by the multi-component NADPH oxidase complex. The oxidase is comprised of three membrane subunits (gp91^{phox}/NOX2, p22^{phox}, and Rap1A) and four cytosolic proteins (p47^{phox}, p67^{phox}, p40^{phox}, and Rac2). Spatial separation of the membrane and cytosolic components maintains enzymatic inactivity in resting neutrophils. Upon stimulation, the cytosolic components translocate to the membrane to form the catalytically active enzyme complex. Phosphorylation of cytosolic NADPH oxidase components is necessary for translocation of those components to the plasma membrane. One of the major targets of phosphorylation is the p47^{phox} subunit. Phosphorylation of a number of serines (Ser³⁰³–Ser³⁷⁹) early in the activation process facilitates p47^{phox} docking to membrane and cytosolic oxidase components, leading to assembly of the functional oxidase (El-Benna et al., 1994, 1996; Groemping et al., 2003).

Non-receptor tyrosine kinases and p38 mitogen-activated protein kinase (MAPK) are signaling molecules that participate in priming respiratory burst activity by TNF α (El-Benna et al., 1996; McLeish et al., 1998; Forsberg et al., 2001; Dewas et al., 2003; Boussetta et al., 2010). Inhibition of tyrosine kinase activity blocks the activation of p38 MAPK by TNF α (McLeish et al., 1998), indicating that tyrosine kinases participate in priming by activating p38 MAPK. TNF α -mediated activation of the p38 MAPK pathway contributes to priming by enhancing plasma membrane translocation of the cytosolic components of the NADPH oxidase and by increasing expression of the plasma membrane oxidase components. Enhanced translocation of cytosolic components results from p38 MAPK-dependent phosphorylation of Ser³⁴⁵ on p47^{phox}. Phosphorylation of Ser³⁴⁵ initiates a series of conformational changes in p47^{phox} that result in hyperactivation of the NADPH oxidase. The initial event is binding of the prolyl isomerase Pin1 to the phospho-Ser³⁴⁵ site (Boussetta et al., 2010). This produces a conformational change in p47^{phox} that exposes additional amino acids for phosphorylation by protein kinase C (PKC). Phosphorylation by PKC produces a second conformational change that promotes p47^{phox} binding to p22^{phox}. That interaction leads to translocation and assembly of all the cytosolic oxidase components with the membrane NADPH oxidase components. Pin1 is also involved in priming by GM-CSF and CL097, a TLR8 agonist (Makni-Maalej et al., 2012, 2015). Unlike TNF α , GM-CSF induces phosphorylation of Ser³⁴⁵ on p47^{phox} through activation of ERK1/2, not p38 MAPK (Boussetta et al., 2010; Makni-Maalej et al., 2015). This observation indicates that multiple signal transduction pathways induce the same molecular events required for priming. Those

redundant signal transduction pathways are unlikely to serve as effective therapeutic targets.

Over a decade ago, it was suggested that TNF α and LPS play a role in respiratory burst priming by influencing membrane trafficking (DeLeo et al., 1998; Ward et al., 2000). Direct confirmation was provided recently by selectively blocking exocytosis prior to priming through the use of cell-permeable, peptide inhibitors of SNARE protein interactions (Uriarte et al., 2011; McLeish et al., 2013). Those studies determined that exocytosis of secretory vesicles and gelatinase granules is required for priming by TNF α and platelet activating factor. Exocytosis could be contributing to priming by increasing plasma membrane expression of receptors, signaling molecules, and/or NADPH oxidase membrane components. The role of receptor and signaling molecule expression in priming was examined by measuring the activation of p38 MAPK and ERK1/2 in neutrophils primed during inhibition of exocytosis (Uriarte et al., 2011). The absence of granule exocytosis had no effect on activation of either MAPK, indicating that increased expression of receptors and signaling molecules does not contribute to priming (Uriarte et al., 2011). Inhibition of Pin1 activity had no effect on neutrophil granule exocytosis (McLeish et al., 2013). We interpret those studies to indicate that enhanced translocation of cytosolic oxidase components and increased expression of membrane oxidase components are independent events, both of which are required for priming.

A second membrane trafficking event that participates in priming respiratory burst activity is clathrin-mediated endocytosis. Moreland and colleagues reported that the NADPH oxidase assembles on endosomes, and the subsequent H₂O₂ production was required for neutrophil priming by endotoxin (Moreland et al., 2007; Volk et al., 2011; Lamb et al., 2012). We have confirmed those observations and determined that endocytosis is an upstream event in neutrophil granule exocytosis.

Neutrophil Granule Release

Neutrophil granules are divided into four classes based on granule density and contents (Borregaard and Cowland, 1997; Lominadze et al., 2005; Rørvig et al., 2013). Secretory vesicles are created by endocytosis, while gelatinase (tertiary), specific (secondary), and azurophilic (primary) granules are formed from the *trans*-Golgi network during neutrophil maturation (Borregaard, 2010). Granule subsets undergo an ordered release based on stimulus intensity, termed graded exocytosis (Sengelov et al., 1993, 1995). Secretory vesicles undergo exocytosis more easily and completely than gelatinase granules. Specific and azurophilic granules, which contain toxic anti-microbial components, undergo the most limited exocytosis. An *in vivo* study showed that neutrophils migrating into a skin blister created in normal human subjects release nearly 100% of their secretory vesicles, 40% of gelatinase granules, 20% of specific granules, and <10% of azurophilic granules (Sengelov et al., 1995).

We recently reported that TNF α directly stimulated exocytosis of secretory vesicles and gelatinase granules (McLeish et al., 2017). Those results support previous studies showing that

TABLE 1 | Known priming agents' capacity to induce phenotypic changes in neutrophils.

Priming Agent	Adhesion	Chemotaxis	Phagocytosis	Granule Release	NET formation	Apoptosis	Inflammatory Mediators
Chemoattractants	fMLF ↑ (El Azreg et al., 2011)	↑ Halpert et al., 2011	↑ Richardson and Patel, 1995	↑ Uriarte et al., 2011	?	No change Klein et al., 2001	↑ Browning et al., 1997
	C5a ↑ Jagels et al., 2000	↑ Halpert et al., 2011	↑/↓ Morris et al., 2011; Tsuboi et al., 2011	↑ DiSopio et al., 2006	?	↓ Perianayagam et al., 2002	↑ Finsterbusch et al., 2014
	LTB4 ↑ Eun et al., 2011	↑ Afonso et al., 2012	↑ Mancuso et al., 2001	↑ Kaman, 2002	?	↓ Klein et al., 2001	↑ Finsterbusch et al., 2014
	PAF ↑ Kulkarni et al., 2007	↑ Shalit et al., 1987	↑ Rosales and Brown, 1991	↑ Andreasson et al., 2013	?	↓ Khreiss et al., 2004	↑ Aquino et al., 2016
Cytokines	TNF- α ↑ Bouaouina et al., 2004	↑ Montecucco et al., 2008	↑ Della Bianca et al., 1995	↑ McLeish et al., 2013, 2017	↑ Hazeldine et al., 2014	↑/↓ Murray et al., 1997	↑ Bauldry et al., 1991; Jablonska et al., 2002b
	GM-CSF ↑ Yu et al., 1990	↑ Cheng et al., 2001	↑ Kletter et al., 1989	↑ Kowanko et al., 1991	↑ Yousefi et al., 2009	↓ Klein et al., 2000	↑ DiPersio et al., 1988b; Lindemann et al., 1988
	IFN- γ ↑ Klebanoff et al., 1992	↓ Aas et al., 1996	↑ Melby et al., 1982	↑ Cassatella et al., 1988	?	↑ Perussia et al., 1987	↑ Humphreys et al., 1989
	IL-1 β ↑ Brandolini et al., 1997	↑ Brandolini et al., 1997	?	↑ Brandolini et al., 1997	?	?	?
	IL-8 ↑ Detmers et al., 1990	↑ Baggolini and Clark-Lewis, 1992	↑ Richardson and Patel, 1995	↑ Baggolini and Clark-Lewis, 1992	↑ Hazeldine et al., 2014; Podaza et al., 2016	↓ Acorci et al., 2009	?
	IL-15 ?	↑ Mastroianni et al., 2000	↑ Musso et al., 1998	?	?	↓ Mastroianni et al., 2000	↑ Musso et al., 1998; Jablonska et al., 2001
	IL-18 ↑ Wyman et al., 2002	?	?	?	?	↑ Wyman et al., 2002	↑ Jablonska et al., 2002a
	IL-33 ?	↑ Le et al., 2012	↑ Lan et al., 2016	?	?	?	?
	Adiponectin ?	?	?	?	?	?	?
Microbial Products	LPS ↑ Hayashi et al., 2003; Sabroe et al., 2003	↓/↑ Fan and Malik, 2003; Hayashi et al., 2003	↑ Hayashi et al., 2003	↑ Fittschen et al., 1988; Ward et al., 2000	↑ Hazeldine et al., 2014	↓ Klein et al., 2001	↑ Cassatella, 1995)
	LAMs ?	No change Fietta et al., 2000	No change Fietta et al., 2000	↑ Faldt et al., 2001	?	?	?
	Lipopeptide ↑ Hayashi et al., 2003; Sabroe et al., 2003	↓/↑ Aomatsu et al., 2008	↑ Hayashi et al., 2003	↑/↓ (Whitmore et al., 2016)	?	Minimal effect Sabroe et al., 2003	↑/↓ Whitmore et al., 2016
	Flagellin ↑ Hayashi et al., 2003	↓ Hayashi et al., 2003	↑ Hayashi et al., 2003	?	?	No change/ Francois et al., 2005; Salamone et al., 2010	↑/↓ Hayashi et al., 2003
Others	ATP ?	↑ Ding et al., 2016	?	↑ Aziz et al., 1997; Meshki et al., 2004	?	?	?
	Substance P ↑ Dianzani et al., 2003	↑ Marasco et al., 1981; Perianin et al., 1989	?	↑ Marasco et al., 1981	?	↓ Bockmann et al., 2001	↑ Perianin et al., 1989; Wozniak et al., 1989
	CL097, CL075 ?	?	?	↑ Makni-Maalej et al., 2012	?	?	?
	Adhesion -	-	↑ Kasorn et al., 2009	↑ Xu and Hakansson, 2002	?	↓ Mayadas and Cullere, 2005	↑ Steadman et al., 1996

Table 1 shows the phenotypic changes induced by agents known to prime neutrophil respiratory burst activity. All the priming agents listed in this table are known inducers of enhanced NADPH oxidase activity. ↑ refers to an increase in activity compared to unprimed neutrophils, ↓ refers to a decrease in activity, and ? indicates an unknown effect of priming agent.

exocytosis of secretory vesicles and gelatinase granule is required for TNF α -induced priming (McLeish et al., 2013). Neither TNF α nor fMLF, alone, stimulated exocytosis of specific and azurophilic granules. However, TNF α primed the release of both granule subsets upon subsequent stimulation by fMLF (McLeish et al., 2017). The ability of TNF α to prime exocytosis of azurophilic granules was also reported by Potera et al. (2016). Thus, differential regulation of exocytosis of the four granule subsets by TNF α primes the two major neutrophil anti-microbial defense mechanisms for enhanced release of ROS and toxic granule contents, while protecting against cell injury from inappropriate release of those toxic products. On the other hand, Ramadass et al. showed that GM-CSF both stimulated and primed exocytosis of gelatinase, specific, and azurophilic granules in mouse neutrophils (Ramadass et al., 2017). The basis for differences between TNF α and GM-CSF could be due to disparate capabilities of priming agents or to species differences.

Proteins that control priming by regulating exocytosis have only recently been identified. As pharmacologic inhibition of p38 MAPK prevents TNF- α stimulated exocytosis (Mocsai et al., 1999; Uriarte et al., 2011; McLeish et al., 2013), we employed a phosphoproteomic analysis by mass spectrometry to identify proteins phosphorylated by the p38 MAPK pathway during TNF α stimulation (McLeish et al., 2017). Four of the proteins identified, Raf1, MARCKS, ABI1, and myosin VI, were previously shown to be involved in exocytosis in various cells. We confirmed that Raf1 participates in TNF α -stimulated exocytosis. Catz and colleagues used neutrophils from transgenic mice to identify Rab27a and its target, Munc13-4, as mediators of neutrophil exocytosis stimulated by GM-CSF (Ramadass et al., 2017). They showed that Rab27a, but not Munc13-4, was required for GM-CSF priming of exocytosis to subsequent stimulation by TLR agonists or formyl peptides. Thus, the mechanisms that control neutrophil exocytosis during priming offer potential targets for intervention in inflammatory processes in which neutrophil priming is involved.

Adhesion, Chemotaxis, and Phagocytosis

As shown in **Figure 1**, microbial invasion or tissue injury releases pathogen-associated molecular pattern (PAMPs) or damage-associated molecular pattern (DAMPs) molecules that induce sentinel immune cells to release pro-inflammatory cytokines. Those cytokines modify both endothelial cell and neutrophil adhesion molecule expression to facilitate the capture of circulating neutrophils and to mediate their migration into tissues. As shown in **Table 1**, all priming agents for which there are data directly activate neutrophil adhesion. However, differential regulation of adhesion molecule expression and activation by different priming agents may produce different rates of neutrophil adhesion and migration efficiency. For example, neutrophil exposure to TNF α increases plasma membrane expression of the β 2 integrin receptor, CD11b/CD18, through exocytosis of secretory vesicles; decreases expression of the selectin receptor CD62-L through receptor shedding; and induces sustained activation of CD11b/CD18 through inside-out signaling (Condliffe et al., 1996; Swain et al., 2002). On the other hand, PAF increases surface expression of the CD11b/CD18,

has no effect on selectin expression, and induces only transient activation of CD11b/CD18 (Berends et al., 1997; Khreiss et al., 2004). The *in vivo* significance of those differences in adhesion molecule expression and activation remains to be determined.

With the exception of IFN γ , neutrophil chemotaxis is enhanced by all priming agents for which there are data (**Table 1**). In addition to increased expression of adhesion molecules and receptors resulting from exocytosis, priming agents increase actin reorganization (Borgquist et al., 2002), and enhances chemokinesis and chemotaxis (Montecucco et al., 2008; Yao et al., 2015). For example, treatment of neutrophils with PAF, IL-8, or TNF α , alone, induces chemokinesis, while subsequent exposure to an fMLF gradient leads to enhanced neutrophil chemotaxis (Drost and MacNee, 2002). Additionally, TNF α -primed neutrophils gain the ability to migrate toward the chemokine CCL3, which is found in inflammatory sites, but is normally not a neutrophil chemo attractant (Montecucco et al., 2008).

Neutrophil adhesion through both the engagement of neutrophil β 2 integrin receptors with endothelial cell adhesion molecules and the binding of neutrophil receptors with extracellular matrix proteins primes respiratory burst activity (Stanislowski et al., 1990; Dapino et al., 1993; Liles et al., 1995). Neutrophil adhesion induces other priming phenotypes, including exocytosis of secretory vesicles and gelatinase granules and a reduced rate of apoptosis (Hu et al., 2004; McGettrick et al., 2006; Paulsson et al., 2010). Thus, transmigration of neutrophils into the extravascular space can be expected to directly induce some of the features of priming.

When neutrophils arrive at the site of infection, they demonstrate increased phagocytosis due to upregulation in the number and affinity of phagocytic receptors (Condliffe et al., 1998; Rainard et al., 2000; Le et al., 2012). **Table 1** lists the effects of specific priming agents on phagocytosis. Exposure of bovine neutrophils to the combination of two priming agents, TNF α and C5a each at suboptimal concentrations, enhanced both the rate of phagocytosis and the killing capacity toward serum opsonized *Staphylococcus aureus* (Rainard et al., 2000); and incubation of human neutrophils with insulin-like growth factor I (IGF-I) results in a significant increase in phagocytosis of both IgG-opsonized *S. aureus* and serum-opsonized *Candida albicans* (Bjerknes and Aarskog, 1995). Increased neutrophil phagocytosis is dependent on the concentration and incubation time with IGF-1, and is due to increased complement receptor (CR) 1 and CR3 expression. IGF-1 enhances Fc γ receptor-dependent phagocytosis through increased receptor function and activation, while Fc γ receptor expression is unchanged (Bjerknes and Aarskog, 1995). Thus, neutrophil exposure to the complex milieu of priming agents *in vivo* is likely to produce additive or synergistic changes in functional responses. Defining neutrophil responses in that complex environment will require application of systems biology methodologies.

Neutrophil Extracellular Trap (NET) Formation

Since their first description in 2004, neutrophil extracellular traps (NETs) have received intense investigation. Although, the

majority of studies have measured NET formation by resting neutrophils, neutrophils from normal subjects primed by TNF α *in vitro* demonstrated robust NET formation following a 3 h exposure to anti-neutrophil cytoplasmic antibodies (Kessenbrock et al., 2009). Enhanced NET formation in primed neutrophils is supported by other *in vitro* studies using GM-CSF and TNF α (Yousefi et al., 2009; Hazeldine et al., 2014). The effect of priming agents on NET formation is listed in **Table 1**.

Despite their original classification as the third bacterial killing mechanism, current opinion leans toward NETs being important contributors to autoimmunity and tissue injury, rather than antibacterial activity (Sorensen and Borregaard, 2016). *In vivo*, enhanced NET formation following a systemic change in levels of inflammatory cytokines has been described in cancers, multiple sclerosis, and diabetes (Chechlinska et al., 2010; Naegele et al., 2012; Fadini et al., 2016). Using a chronic myelogenous leukemia mouse model, Demers and colleagues reported that non-malignant neutrophils showed enhanced NET formation, leading to increased coagulation and thrombosis (Demers et al., 2012). Priming of NET formation was reproduced in control mice by sequential administration of granulocyte colony-stimulating factor (G-CSF) and LPS. The authors suggested that priming NET formation by systemic cytokines plays a role in cancer progression. While the current literature indicates that enhanced NET formation is a component of neutrophil priming, the functional consequences of that response remain to be determined.

Secretion of Lipid and Cytokine Mediators

As summarized in **Table 1**, primed neutrophils demonstrate increased metabolic and transcriptional activity that leads to synthesis of a number of pro- and anti-inflammatory chemokines, cytokines, and lipids. Although, the ability of neutrophils to synthesize those products is less than that of macrophages, the large number of neutrophils present at sites of inflammation is postulated to influence both innate and adaptive immune responses through release of those inflammatory mediators.

Pro-inflammatory lipid mediators like leukotriene B₄ (LTB₄) can be produced *de novo* by the arachidonate 5-lipoxygenase (5-LO) pathway in neutrophils and play important roles in aggregation, degranulation, and chemotaxis (O'Flaherty et al., 1979; Flamand et al., 2000). The production of these lipid mediators occurs through a series of biochemical events that primarily take place in the perinuclear region where membrane phospholipids are first converted to arachidonic acid (AA) by the calcium-dependent enzyme phospholipase A₂ (PLA₂) (Luo et al., 2003; Leslie, 2004). The newly synthesized AA is then converted by 5-LO into leukotriene A₄ (LTA₄), which is the immediate precursor of LTB₄. Neutrophil production of LTB₄ is responsible for a second wave of neutrophil recruitment during inflammation, a process termed "swarming" (Lammermann et al., 2013). This is one of many examples of amplification loops initiated by neutrophils (Nemeth and Mocsai, 2016).

Direct activation of neutrophils by fMLF does not lead to the detectable release of leukotrienes, but priming with GM-CSF, LPS, or TNF α followed by fMLF stimulation significantly

increases LTB₄ release (see **Table 1**; DiPersio et al., 1988a; Schatz-Munding and Ullrich, 1992; Palmantier et al., 1994; Seeds et al., 1998; Zarini et al., 2006). All three of these priming agents activate PLA₂ and increase AA release without increasing intracellular Ca²⁺ (DiPersio et al., 1988b; Schatz-Munding and Ullrich, 1992; Zarini et al., 2006). The elevation in available AA substrate leads to prolonged activation of 5-LO and enhanced production of downstream lipid mediators (Surette et al., 1993, 1998; Doerfler et al., 1994). Once produced, LTB₄ can exert autocrine effects. It primes neutrophil responses to toll-like-receptor (TLR) agonists, resulting in enhanced cytokine (IL-8, TNF α) secretion (Gaudreault et al., 2012). TLR9 mRNA levels are upregulated upon priming with LTB₄, but there is no increase in surface expression of TLR2, TLR4, or the co-receptors TLR1 and TLR6 following LTB₄ exposure (Gaudreault and Gosselin, 2009; Gaudreault et al., 2012). Instead, neutrophil LTB₄-induced hyper-responsiveness is mediated by the potentiation of TLR-induced intracellular signaling. TAK1 and p38 MAPK, which are essential in TLR-activated cytokine release, are phosphorylated and activated following LTB₄ interaction with its seven transmembrane-spanning receptor.

PAF is another lipid inflammatory mediator whose production is primed in neutrophils. Both LPS and GM-CSF enhance PAF synthesis in response to activating stimuli (Aglietta et al., 1990; Surette et al., 1998). After priming with GM-CSF, there is increased enzymatic activity of acetyl transferase, the enzyme responsible for the synthesis of PAF (Aglietta et al., 1990). However, the pattern of PAF synthesis after LPS priming is attributed to a biphasic, autocrine response. The early peak in production is due to the direct effect of LPS, while the delayed peak is a result of LPS-induced IL-8 and TNF- α release (Bussolati et al., 1997).

Neutrophils modulate inflammation through the release of stored or newly produced cytokines and chemokines (Cassatella, 1999). Exposure of neutrophils to priming agents leads to an increase in synthesis and release of IL-1 α , IL-1 β , IL-6, IL-8, TNF α , CXCL1, CXCL2, CCL3 (MIP-1 α), CCL4 (MIP-1 β) (Roberge et al., 1998; Zallen et al., 1999; Jablonska et al., 2002b; Choi et al., 2008; Wright et al., 2013). The inducible synthesis of the majority of cytokines and chemokines results from increased gene transcription (Marucha et al., 1991; Cassatella et al., 1995; Cassatella, 1996, 1999; Fernandez et al., 1996). TNF α , LPS, and GM-CSF increase intra-nuclear translocation of NF- κ B, C/EBP, or CREB transcription factors (Cloutier et al., 2007, 2009; Mayer et al., 2013). LPS induces a biphasic production of IL-8. For the first few hours (2–6 h) of exposure, LPS directly stimulates IL-8 synthesis, but the second wave of sustained IL-8 release (up to 18 h) is due to the endogenous release of TNF α and IL-1 β (Cassatella et al., 1993).

Release of Neutrophil Extracellular Vesicles

Cell-derived vesicles represent a mechanism for cell-cell communication. Exosomes are 50–100 nm vesicles released from multivesicular bodies that are involved in antigen presentation and cell-to-cell transfer of receptors or RNA (Gyorgy et al.,

2011). Larger vesicles, called microvesicles or microparticles express tissue factors on their surface that are capable of initiating coagulation. Neutrophils undergoing apoptosis or stimulated by chemotactic agents, opsonic receptors, or TNF α release microparticles. However, the microparticles have varying compositions and functional capabilities, depending on the stimulus (Dalli et al., 2013; Johnson et al., 2014; Lorincz et al., 2015). Microparticles obtained from neutrophils stimulated by chemotactic agents or phorbol esters activate cytokine (IL-6) secretion from endothelial cells and platelets (Mesri and Altieri, 1998; Pluskota et al., 2008). Chemotactic peptide-induced microparticles increase secretion of the anti-inflammatory cytokine transforming growth factor- β and interfere with the maturation of monocyte-derived dendritic cells (Gasser and Schifferli, 2004; Eken et al., 2010). Auto-antibody-stimulated release of neutrophil microparticles was suggested to be involved in the pathogenesis of vasculitis (Hong et al., 2012). Additional activities ascribed to neutrophil microparticles include suppression of bacterial growth, activation of endothelial cell cytokine production, altered cytokine profile of natural killer cells and monocytes, and increased coagulation (Mesri and Altieri, 1998; Timar et al., 2013a,b; Pliyev et al., 2014). An understanding of the stimuli and signal transduction pathways leading to formation and release of neutrophil extracellular vesicles and their roles in inflammation remains to be developed.

Rate of Apoptosis

Table 1 indicates that neutrophil apoptosis is variably affected in response to priming agents. While LPS, GM-CSF, IL-8, and LTB $_4$ have been found to extend neutrophil lifespan *in vitro*, PAF, fMLF, and IL-6 show no effect, and TNF α shows a biphasic response where it promotes apoptosis during the first 8 h of exposure, followed by a delayed rate of apoptosis at later times (Klein et al., 2000, 2001; Cowburn et al., 2002; Liu et al., 2005; Wright et al., 2014). Primed neutrophils from patients with multiple sclerosis, ANCA-associated vasculitis, and liver cirrhosis show increased apoptosis (Harper et al., 2001; Klimchenko et al., 2011; Naegele et al., 2012), while neutrophils from patients at risk of multiple-organ failure and individuals presenting with septic peritonitis, severe trauma, or septic trauma show a decrease in apoptosis (Ertel et al., 1998; Biffl et al., 1999, 2001; Nolan et al., 2000; Feterowski et al., 2001). Those conflicting reports of the effect of inflammation on apoptosis *in vivo* are likely due to different priming agents involved in different diseases, different responses during the time course of disease, and differences in the neutrophil micro-environment, such as cell density (Hannah et al., 1998).

The mechanisms underlying the effects of priming on neutrophil apoptosis have been partially characterized. As for TNF α , increased rates of apoptosis during the first hours of exposure are associated with activation of caspase cascades (Murray et al., 1997). TNF α also induces an early, PI-3K-mediated increase in mRNA levels for Bad, a member of the BCL2 family that regulates apoptosis. On the other hand, decreased neutrophil apoptosis observed at later time points is associated with a reduction in Bad mRNA levels (Cowburn et al., 2002). GM-CSF, IL-8, LPS, and LTB $_4$ decrease the rate

of neutrophil apoptosis through activation of ERK1/2 and/or PI-3K/Akt pathways (Klein et al., 2000, 2001). Incubation of neutrophils with fMLF had no effect on the rate of apoptosis, despite activation of both ERK1/2 and Akt (Klein et al., 2001). GM-CSF was also shown to decrease mRNA levels of Bad, while increasing its phosphorylation (Cowburn et al., 2002). A RNA seq study comparing TNF α and GM-CSF priming pathways showed that out of 580 genes differentially expressed between both agents, 58 were implicated in the delay of apoptosis. Thus, each priming agent produced a distinct profile of pro- and anti-apoptotic genes (Wright et al., 2013). The varying rates of neutrophil apoptosis may serve different functions in the inflammatory response. For example, a reduced rate of apoptosis early in the recruitment of neutrophils results in a brisk accumulation of primed neutrophils. On the other hand, an enhanced rate of apoptosis at later time points promotes resolution through loss of active neutrophils and a change in phenotype of monocytes engulfing apoptotic neutrophils.

CONCLUSIONS

The altered neutrophil functions described in this review indicate that priming is a complex phenomenon. Priming involves enhanced respiratory burst, exocytosis, NET formation, and chemotaxis in response to a second stimulus. Priming, however, is not just preparation for an enhanced response to a second stimulus. Priming involves activation of a subset of neutrophil responses, including adhesion, transcription, cytoskeletal reorganization, translocation and expression of receptors, and other molecules, the rate of constitutive apoptosis, metabolic activity, and phagocytosis. The altered neutrophil responses associated with priming primarily result in amplification of the inflammatory response. Although, recruitment of primed neutrophils improves the clearance of invading microbes, the risk of directly injuring surrounding cells is increased. Moreover, the increased synthesis and release of cytokines and lipids by primed neutrophils, combined with increased neutrophil recruitment and life-span, result in an increased local concentration of pro-inflammatory agents. Those agents recruit and prime additional neutrophils, leading to an enhanced innate immune response. Neutrophil-dependent recruitment and activation of dendritic cells and various lymphocyte subsets also enhances the adaptive immune response.

We propose that the current definition of priming, which focuses on a transition state to an enhanced responsiveness to a second stimulus, is too restrictive. Neutrophil priming also results in activation of a subset of neutrophil responses that regulate innate and adaptive immunity. Additionally, neutrophil responses to priming agents vary depending on concentration of the priming agent, time of exposure, and the specific priming agent (Potera et al., 2016; McLeish et al., 2017). It seems likely that neutrophils are exposed to graded concentrations of priming agents as they progress through the multistep process of recruitment, as occurs with chemoattractants. This leads to the hypothesis that, similar to graded granule exocytosis, priming

occurs in a graded manner during a neutrophil's journey to the site of inflammation. This graded response allows neutrophils to acquire functions in an ordered manner, as required during recruitment. A fully primed neutrophil that releases a maximal amount of toxic chemicals would occur when an optimal concentration of a priming stimulus is encountered. Combining knowledge of the molecular events with an understanding of priming at a systems level will identify therapeutic targets for neutrophil functions that exacerbate individual diseases, while preserving the functions that participation in host defense.

REFERENCES

- Aas, V., Lappégard, K. T., Siebke, E. M., and Benestad, H. B. (1996). Modulation by interferons of human neutrophilic granulocyte migration. *J. Interferon Cytokine Res.* 16, 929–935. doi: 10.1089/jir.1996.16.929
- Acorci, M. J., Dias-Melicio, L. A., Golim, M. A., Bordon-Graciani, A. P., Peracoli, M. T., and Soares, A. M. (2009). Inhibition of human neutrophil apoptosis by *Paracoccidioides brasiliensis*: role of interleukin-8. *Scand. J. Immunol.* 69, 73–79. doi: 10.1111/j.1365-3083.2008.02199.x
- Afonso, P. V., Janka-Junttila, M., Lee, Y. J., McCann, C. P., Oliver, C. M., Aamer, K. A., et al. (2012). LTB4 is a signal-relay molecule during neutrophil chemotaxis. *Dev. Cell* 22, 1079–1091. doi: 10.1016/j.devcel.2012.02.003
- Aglietta, M., Monzeglio, C., Apra, F., Mossetti, C., Stern, A. C., Giribaldi, G., et al. (1990). *In vivo* priming of human normal neutrophils by granulocyte-macrophage colony stimulating factor: effect on the production of platelet activating factor. *Br. J. Haematol.* 75, 333–339. doi: 10.1111/j.1365-2141.1990.tb04345.x
- Andreasson, E., Onnheim, K., and Forsman, H. (2013). The subcellular localization of the receptor for platelet-activating factor in neutrophils affects signaling and activation characteristics. *Clin. Dev. Immunol.* 2013:456407. doi: 10.1155/2013/456407
- Aomatsu, K., Kato, T., Fujita, H., Hato, F., Oshitani, N., Kamata, N., et al. (2008). Toll-like receptor agonists stimulate human neutrophil migration via activation of mitogen-activated protein kinases. *Immunology* 123, 171–180. doi: 10.1111/j.1365-2567.2007.02684.x
- Aquino, E. N., Neves, A. C., Santos, K. C., Uribe, C. E., Souza, P. E., Correa, J. R., et al. (2016). Proteomic analysis of neutrophil priming by PAF. *Protein Pept. Lett.* 23, 142–151. doi: 10.2174/0929866523666151202210604
- Aziz, K. A., Cawley, J. C., Treweeke, A. T., and Zuzel, M. (1997). Sequential potentiation and inhibition of PMN reactivity by maximally stimulated platelets. *J. Leukoc. Biol.* 61, 322–328.
- Baggiolini, M., and Clark-Lewis, I. (1992). Interleukin-8, a chemotactic and inflammatory cytokine. *FEBS Lett.* 307, 97–101. doi: 10.1016/0014-5793(92)80909-Z
- Bass, D. A., Olbrantz, P., Szejda, P., Seeds, M. C., and McCall, C. E. (1986). Subpopulations of neutrophils with increased oxidative product formation in blood of patients with infection. *J. Immunol.* 136, 860–866.
- Bauldry, S. A., McCall, C. E., Cousart, S. L., and Bass, D. A. (1991). Tumor necrosis factor-alpha priming of phospholipase A2 activation in human neutrophils. An alternative mechanism of priming. *J. Immunol.* 146, 1277–1285.
- Berends, C., Dijkhuizen, B., de Monchy, J. G., Dubois, A. E., Gerritsen, J., and Kauffman, H. F. (1997). Inhibition of PAF-induced expression of CD11b and shedding of L-selectin on human neutrophils and eosinophils by the type IV selective PDE inhibitor, rolipram. *Eur. Respir. J.* 10, 1000–1007. doi: 10.1183/09031936.97.10051000
- Biffl, W. L., Moore, E. E., Zallen, G., Johnson, J. L., Gabriel, J., Offner, P. J., et al. (1999). Neutrophils are primed for cytotoxicity and resist apoptosis in injured patients at risk for multiple organ failure. *Surgery* 126, 198–202. doi: 10.1016/S0039-6060(99)70155-8
- Biffl, W. L., West, K. E., Moore, E. E., Gonzalez, R. J., Carnaggio, R., Offner, P. J., et al. (2001). Neutrophil apoptosis is delayed by trauma patients' plasma via a mechanism involving proinflammatory phospholipids and protein kinase C. *Surg. Infect.* 2, 289–293; discussion: 294–285. doi: 10.1089/10962960152813322
- Bjerknes, R., and Aarskog, D. (1995). Priming of human polymorphonuclear neutrophilic leukocytes by insulin-like growth factor I: increased phagocytic capacity, complement receptor expression, degranulation, and oxidative burst. *J. Clin. Endocrinol. Metab.* 80, 1948–1955.
- Bockmann, S., Seep, J., and Jonas, L. (2001). Delay of neutrophil apoptosis by the neuropeptide substance P: involvement of caspase cascade. *Peptides* 22, 661–670. doi: 10.1016/S0196-9781(01)00376-X
- Borgquist, J. D., Quinn, M. T., and Swain, S. D. (2002). Adhesion to extracellular matrix proteins modulates bovine neutrophil responses to inflammatory mediators. *J. Leukoc. Biol.* 71, 764–774.
- Borregaard, N. (2010). Neutrophils, from marrow to microbes. *Immunity* 33, 657–670. doi: 10.1016/j.immuni.2010.11.011
- Borregaard, N., and Cowland, J. B. (1997). Granules of the human neutrophilic polymorphonuclear leukocyte. *Blood* 89, 3503–3521.
- Bouaouina, M., Blouin, E., Halbwachs-Mecarelli, L., Lesavre, P., and Rieu, P. (2004). TNF-induced beta2 integrin activation involves Src kinases and a redox-regulated activation of p38 MAPK. *J. Immunol.* 173, 1313–1320. doi: 10.4049/jimmunol.173.2.1313
- Boussetta, T., Gougerot-Pocidallo, M. A., Hayem, G., Ciappelloni, S., Raad, H., Arabi Derkawi, R., et al. (2010). The prolyl isomerase Pin1 acts as a novel molecular switch for TNF-alpha-induced priming of the NADPH oxidase in human neutrophils. *Blood* 116, 5795–5802. doi: 10.1182/blood-2010-03-273094
- Brandolini, L., Sergi, R., Caselli, G., Boraschi, D., Locati, M., Sozzani, S., et al. (1997). Interleukin-1 beta primes interleukin-8-stimulated chemotaxis and elastase release in human neutrophils via its type I receptor. *Eur. Cytokine Netw.* 8, 173–178.
- Browning, D. D., Pan, Z. K., Prossnitz, E. R., and Ye, R. D. (1997). Cell type- and developmental stage-specific activation of NF-kappaB by fMet-Leu-Phe in myeloid cells. *J. Biol. Chem.* 272, 7995–8001. doi: 10.1074/jbc.272.12.7995
- Bussolati, B., Mariano, F., Montrucchio, G., Piccoli, G., and Camussi, G. (1997). Modulatory effect of interleukin-10 on the production of platelet-activating factor and superoxide anions by human leukocytes. *Immunology* 90, 440–447. doi: 10.1111/j.1365-2567.1997.00440.x
- Cassatella, M. A. (1995). The production of cytokines by polymorphonuclear neutrophils. *Immunol. Today* 16, 21–26. doi: 10.1016/0167-5699(95)80066-2
- Cassatella, M. A. (1996). Interferon-gamma inhibits the lipopolysaccharide-induced macrophage inflammatory protein-1 alpha gene transcription in human neutrophils. *Immunol. Lett.* 49, 79–82. doi: 10.1016/0165-2478(95)02484-0
- Cassatella, M. A. (1999). Neutrophil-derived proteins: selling cytokines by the pound. *Adv. Immunol.* 73, 369–509. doi: 10.1016/S0065-2776(08)60791-9
- Cassatella, M. A., Cappelli, R., Della Bianca, V., Grzeskowiak, M., Dusi, S., and Berton, G. (1988). Interferon-gamma activates human neutrophil oxygen metabolism and exocytosis. *Immunology* 63, 499–506.
- Cassatella, M. A., Gasperini, S., Calzetti, F., McDonald, P. P., and Trinchieri, G. (1995). Lipopolysaccharide-induced interleukin-8 gene expression in human granulocytes: transcriptional inhibition by interferon-gamma. *Biochem. J.* 310(Pt 3), 751–755. doi: 10.1042/bj3100751

AUTHOR CONTRIBUTIONS

Contributed to the writing and editing of the manuscript IM, KM, SU; designed and illustrated the figure IM.

FUNDING

The authors were supported by a Merit Review Award (BX001838) from the Department of Veterans Affairs (KM) and by the National Institute of Dental and Craniofacial Research (DE024509 to SU).

- Cassatella, M. A., Meda, L., Bonora, S., Ceska, M., and Constantin, G. (1993). Interleukin 10 (IL-10) inhibits the release of proinflammatory cytokines from human polymorphonuclear leukocytes. Evidence for an autocrine role of tumor necrosis factor and IL-1 beta in mediating the production of IL-8 triggered by lipopolysaccharide. *J. Exp. Med.* 178, 2207–2211. doi: 10.1084/jem.178.6.2207
- Chechliniska, M., Kowalewska, M., and Nowak, R. (2010). Systemic inflammation as a confounding factor in cancer biomarker discovery and validation. *Nat. Rev. Cancer* 10, 2–3. doi: 10.1038/nrc2782
- Cheng, S. S., Lai, J. J., Lukacs, N. W., and Kunkel, S. L. (2001). Granulocyte-macrophage colony stimulating factor up-regulates CCR1 in human neutrophils. *J. Immunol.* 166, 1178–1184. doi: 10.4049/jimmunol.166.2.1178
- Choi, J. C., Jung, J. W., Kwak, H. W., Song, J. H., Jeon, E. J., Shin, J. W., et al. (2008). Granulocyte macrophage colony stimulating factor (GM-CSF) augments acute lung injury via its neutrophil priming effects. *J. Korean Med. Sci.* 23, 288–295. doi: 10.3346/jkms.2008.23.2.288
- Cloutier, A., Ear, T., Blais-Charron, E., Dubois, C. M., and McDonald, P. P. (2007). Differential involvement of NF-kappaB and MAP kinase pathways in the generation of inflammatory cytokines by human neutrophils. *J. Leukoc. Biol.* 81, 567–577. doi: 10.1189/jlb.0806536
- Cloutier, A., Guindi, C., Larivee, P., Dubois, C. M., Amrani, A., and McDonald, P. P. (2009). Inflammatory cytokine production by human neutrophils involves C/EBP transcription factors. *J. Immunol.* 182, 563–571. doi: 10.4049/jimmunol.182.1.563
- Condliffe, A. M., Chilvers, E. R., Haslett, C., and Dransfield, I. (1996). Priming differentially regulates neutrophil adhesion molecule expression/function. *Immunology* 89, 105–111. doi: 10.1046/j.1365-2567.1996.d01-711.x
- Condliffe, A. M., Kitchen, E., and Chilvers, E. R. (1998). Neutrophil priming: pathophysiological consequences and underlying mechanisms. *Clin. Sci.* 94, 461–471. doi: 10.1042/cs0940461
- Cowburn, A. S., Cadwallader, K. A., Reed, B. J., Farahi, N., and Chilvers, E. R. (2002). Role of PI3-kinase-dependent Bad phosphorylation and altered transcription in cytokine-mediated neutrophil survival. *Blood* 100, 2607–2616. doi: 10.1182/blood-2001-11-0122
- Dalli, J., Montero-Melendez, T., Norling, L. V., Yin, X., Hinds, C., Haskard, D., et al. (2013). Heterogeneity in neutrophil microparticles reveals distinct proteome and functional properties. *Mol. Cell. Proteomics* 12, 2205–2219. doi: 10.1074/mcp.M113.028589
- Dapino, P., Dallegri, F., Ottonello, L., and Sacchetti, C. (1993). Induction of neutrophil respiratory burst by tumour necrosis factor-alpha; priming effect of solid-phase fibronectin and intervention of CD11b-CD18 integrins. *Clin. Exp. Immunol.* 94, 533–538. doi: 10.1111/j.1365-2249.1993.tb08230.x
- DeLeo, F. R., Renee, J., McCormick, S., Nakamura, M., Apicella, M., Weiss, J. P., et al. (1998). Neutrophils exposed to bacterial lipopolysaccharide upregulate NADPH oxidase assembly. *J. Clin. Invest.* 101, 455–463. doi: 10.1172/JCI949
- Della Bianca, V., Dusi, S., Nadalini, K. A., Donini, M., and Rossi, F. (1995). Role of 55- and 75-kDa TNF receptors in the potentiation of Fc-mediated phagocytosis in human neutrophils. *Biochem. Biophys. Res. Commun.* 214, 44–50. doi: 10.1006/bbrc.1995.2254
- Demers, M., Krause, D. S., Schatzberg, D., Martinod, K., Voorhees, J. R., Fuchs, T. A., et al. (2012). Cancers predispose neutrophils to release extracellular DNA traps that contribute to cancer-associated thrombosis. *Proc. Natl. Acad. Sci. U.S.A.* 109, 13076–13081. doi: 10.1073/pnas.1200419109
- Detmers, P. A., Lo, S. K., Olsen-Egbert, E., Walz, A., Baggiolini, M., and Cohn, Z. A. (1990). Neutrophil-activating protein 1/interleukin 8 stimulates the binding activity of the leukocyte adhesion receptor CD11b/CD18 on human neutrophils. *J. Exp. Med.* 171, 1155–1162. doi: 10.1084/jem.171.4.1155
- Dewas, C., Dang, P. M., Gougerot-Pocidallo, M. A., and El-Benna, J. (2003). TNF-alpha induces a phosphorylation of p47(phox) in human neutrophils: partial phosphorylation of p47(phox) is a common event of priming of human neutrophils by TNF-alpha and granulocyte-macrophage colony-stimulating factor. *J. Immunol.* 171, 4392–4398. doi: 10.4049/jimmunol.171.8.4392
- Dianzani, C., Collino, M., Lombardi, G., Garbarino, G., and Fantozzi, R. (2003). Substance P increases neutrophil adhesion to human umbilical vein endothelial cells. *Br. J. Pharmacol.* 139, 1103–1110. doi: 10.1038/sj.bjp.0705344
- Ding, Q., Quah, S. Y., and Tan, K. S. (2016). Secreted adenosine triphosphate from *Aggregatibacter actinomycetemcomitans* triggers chemokine response. *Mol. Oral Microbiol.* 31, 423–434. doi: 10.1111/omi.12143
- DiPersio, J. F., Billing, P., Williams, R., and Gasson, J. C. (1988a). Human granulocyte-macrophage colony-stimulating factor and other cytokines prime human neutrophils for enhanced arachidonic acid release and leukotriene B4 synthesis. *J. Immunol.* 140, 4315–4322.
- DiPersio, J. F., Naccache, P. H., Borgeat, P., Gasson, J. C., Nguyen, M. H., and McColl, S. R. (1988b). Characterization of the priming effects of human granulocyte-macrophage colony-stimulating factor on human neutrophil leukotriene synthesis. *Prostaglandins* 36, 673–691. doi: 10.1016/0090-6980(88)90013-5
- DiScipio, R. G., Schraufstatter, I. U., Sikora, L., Zuraw, B. L., and Sriramarao, P. (2006). C5a mediates secretion and activation of matrix metalloproteinase 9 from human eosinophils and neutrophils. *Int. Immunopharmacol.* 6, 1109–1118. doi: 10.1016/j.intimp.2006.02.006
- Doerfler, M. E., Weiss, J., Clark, J. D., and Elsbach, P. (1994). Bacterial lipopolysaccharide primes human neutrophils for enhanced release of arachidonic acid and causes phosphorylation of an 85-kD cytosolic phospholipase A2. *J. Clin. Invest.* 93, 1583–1591. doi: 10.1172/JCI117138
- Drost, E. M., and MacNee, W. (2002). Potential role of IL-8, platelet-activating factor and TNF-alpha in the sequestration of neutrophils in the lung: effects on neutrophil deformability, adhesion receptor expression, and chemotaxis. *Eur. J. Immunol.* 32, 393–403. doi: 10.1002/1521-4141(200202)32:2<393::AID-IMMU393>3.0.CO;2-5
- Eken, C., Martin, P. J., Sadallah, S., Treves, S., Schaller, M., and Schifferli, J. A. (2010). Ectosomes released by polymorphonuclear neutrophils induce a MerTK-dependent anti-inflammatory pathway in macrophages. *J. Biol. Chem.* 285, 39914–39921. doi: 10.1074/jbc.M110.126748
- El Azreq, M. A., Garceau, V., and Bourgoin, S. G. (2011). Cytohesin-1 regulates fMLF-mediated activation and functions of the beta2 integrin Mac-1 in human neutrophils. *J. Leukoc. Biol.* 89, 823–836. doi: 10.1189/jlb.0410222
- El-Benna, J., Dang, P. M., and Gougerot-Pocidallo, M. A. (2008). Priming of the neutrophil NADPH oxidase activation: role of p47phox phosphorylation and NOX2 mobilization to the plasma membrane. *Semin. Immunopathol.* 30, 279–289. doi: 10.1007/s00281-008-0118-3
- El-Benna, J., Faust, L. P., and Babior, B. M. (1994). The phosphorylation of the respiratory burst oxidase component p47phox during neutrophil activation. Phosphorylation of sites recognized by protein kinase C and by proline-directed kinases. *J. Biol. Chem.* 269, 23431–23436.
- El-Benna, J., Han, J., Park, J. W., Schmid, E., Ulevitch, R. J., and Babior, B. M. (1996). Activation of p38 in stimulated human neutrophils: phosphorylation of the oxidase component p47phox by p38 and ERK but not by JNK. *Arch. Biochem. Biophys.* 334, 395–400. doi: 10.1006/abbi.1996.0470
- El-Benna, J., Hurtado-Nedelec, M., Marzaioli, V., Marie, J. C., Gougerot-Pocidallo, M. A., and Dang, P. M. (2016). Priming of the neutrophil respiratory burst: role in host defense and inflammation. *Immunol. Rev.* 273, 180–193. doi: 10.1111/imr.12447
- Ertel, W., Keel, M., Infanger, M., Ungeth, U., Steckholzer, U., and Trentz, O. (1998). Circulating mediators in serum of injured patients with septic complications inhibit neutrophil apoptosis through up-regulation of protein-tyrosine phosphorylation. *J. Trauma* 44, 767–775; discussion: 775–766. doi: 10.1097/00005373-199805000-00005
- Eun, J. C., Moore, E. E., Banerjee, A., Kelher, M. R., Khan, S. Y., Elzi, D. J., et al. (2011). Leukotriene b4 and its metabolites prime the neutrophil oxidase and induce proinflammatory activation of human pulmonary microvascular endothelial cells. *Shock* 35, 240–244. doi: 10.1097/SHK.0b013e3181faceb3
- Fadini, G. P., Menegazzo, L., Rigato, M., Scatolini, V., Poncina, N., Bruttocao, A., et al. (2016). NETosis delays diabetic wound healing in mice and humans. *Diabetes* 65, 1061–1071. doi: 10.2337/db15-0863
- Faldt, J., Dahlgren, C., Ridell, M., and Karlsson, A. (2001). Priming of human neutrophils by mycobacterial lipoarabinomannans: role of granule mobilisation. *Microbes Infect.* 3, 1101–1109. doi: 10.1016/S1286-4579(01)01470-8
- Fan, J., and Malik, A. B. (2003). Toll-like receptor-4 (TLR4) signaling augments chemokine-induced neutrophil migration by modulating cell surface expression of chemokine receptors. *Nat. Med.* 9, 315–321. doi: 10.1038/nm832
- Fernandez, M. C., Walters, J., and Marucha, P. (1996). Transcriptional and post-transcriptional regulation of GM-CSF-induced IL-1 beta gene expression in PMN. *J. Leukoc. Biol.* 59, 598–603.

- Feterowski, C., Weighardt, H., Emmanuilidis, K., Hartung, T., and Holzmann, B. (2001). Immune protection against septic peritonitis in endotoxin-primed mice is related to reduced neutrophil apoptosis. *Eur. J. Immunol.* 31, 1268–1277. doi: 10.1002/1521-4141(200104)31:4<1268::AID-IMMU1268>3.0.CO;2-C
- Fietta, A., Francioli, C., and Gialdroni Grassi, G. (2000). Mycobacterial lipoarabinomannan affects human polymorphonuclear and mononuclear phagocyte functions differently. *Haematologica* 85, 11–18.
- Finsterbusch, M., Voisin, M. B., Beyrau, M., Williams, T. J., and Nourshargh, S. (2014). Neutrophils recruited by chemoattractants *in vivo* induce microvascular plasma protein leakage through secretion of TNF. *J. Exp. Med.* 211, 1307–1314. doi: 10.1084/jem.20132413
- Fittschen, C., Sandhaus, R. A., Worthen, G. S., and Henson, P. M. (1988). Bacterial lipopolysaccharide enhances chemoattractant-induced elastase secretion by human neutrophils. *J. Leukoc. Biol.* 43, 547–556.
- Flamand, N., Boudreault, S., Picard, S., Austin, M., Surette, M. E., Plante, H., et al. (2000). Adenosine, a potent natural suppressor of arachidonic acid release and leukotriene biosynthesis in human neutrophils. *Am. J. Respir. Crit. Care Med.* 161(2 Pt 2), S88–S94. doi: 10.1164/ajrccm.161.supplement_1.lta-18
- Forsberg, M., Löfgren, R., Zheng, L., and Stendahl, O. (2001). Tumour necrosis factor- α potentiates CR3-induced respiratory burst by activating p38 MAP kinase in human neutrophils. *Immunology* 103, 465–472. doi: 10.1046/j.1365-2567.2001.01270.x
- Francois, S., El Benna, J., Dang, P. M., Pedruzzi, E., Gougerot-Pocidalo, M. A., and Elbim, C. (2005). Inhibition of neutrophil apoptosis by TLR agonists in whole blood: involvement of the phosphoinositide 3-kinase/Akt and NF- κ B signaling pathways, leading to increased levels of Mcl-1, A1, and phosphorylated Bad. *J. Immunol.* 174, 3633–3642. doi: 10.4049/jimmunol.174.6.3633
- Gasser, O., and Schifferli, J. A. (2004). Activated polymorphonuclear neutrophils disseminate anti-inflammatory microparticles by ectocytosis. *Blood* 104, 2543–2548. doi: 10.1182/blood-2004-01-0361
- Gaudreault, E., and Gosselin, J. (2009). Leukotriene B4 potentiates CpG signaling for enhanced cytokine secretion by human leukocytes. *J. Immunol.* 183, 2650–2658. doi: 10.4049/jimmunol.0804135
- Gaudreault, E., Paquet-Bouchard, C., Fiola, S., Le Bel, M., Lacerte, P., Shio, M. T., et al. (2012). TAK1 contributes to the enhanced responsiveness of LTB₄-treated neutrophils to Toll-like receptor ligands. *Int. Immunol.* 24, 693–704. doi: 10.1093/intimm/dxs074
- Groemping, Y., Lapouge, K., Smerdon, S. J., and Rittinger, K. (2003). Molecular basis of phosphorylation-induced activation of the NADPH oxidase. *Cell* 113, 343–355. doi: 10.1016/S0092-8674(03)00314-3
- Gyorgy, B., Szabo, T. G., Pasztoi, M., Pal, Z., Misjak, P., Aradi, B., et al. (2011). Membrane vesicles, current state-of-the-art: emerging role of extracellular vesicles. *Cell. Mol. Life Sci.* 68, 2667–2688. doi: 10.1007/s00018-011-0689-3
- Halpert, M. M., Thomas, K. A., King, R. G., and Justement, L. B. (2011). TLT2 potentiates neutrophil antibacterial activity and chemotaxis in response to G protein-coupled receptor-mediated signaling. *J. Immunol.* 187, 2346–2355. doi: 10.4049/jimmunol.1100534
- Hannah, S., Nadra, I., Dransfield, I., Pryde, J. G., Rossi, A. G., and Haslett, C. (1998). Constitutive neutrophil apoptosis in culture is modulated by cell density independently of beta2 integrin-mediated adhesion. *FEBS Lett.* 421, 141–146. doi: 10.1016/S0014-5793(97)01551-2
- Harper, L., Cockwell, P., Adu, D., and Savage, C. O. (2001). Neutrophil priming and apoptosis in anti-neutrophil cytoplasmic autoantibody-associated vasculitis. *Kidney Int.* 59, 1729–1738. doi: 10.1046/j.1523-1755.2001.0590051729.x
- Hayashi, F., Means, T. K., and Luster, A. D. (2003). Toll-like receptors stimulate human neutrophil function. *Blood* 102, 2660–2669. doi: 10.1182/blood-2003-04-1078
- Hazeldine, J., Harris, P., Chapple, I. L., Grant, M., Greenwood, H., Livesey, A., et al. (2014). Impaired neutrophil extracellular trap formation: a novel defect in the innate immune system of aged individuals. *Aging Cell* 13, 690–698. doi: 10.1111/accel.12222
- Hong, Y., Eleftheriou, D., Hussain, A. A., Price-Kuehne, F. E., Savage, C. O., Jayne, D., et al. (2012). Anti-neutrophil cytoplasmic antibodies stimulate release of neutrophil microparticles. *J. Am. Soc. Nephrol.* 23, 49–62. doi: 10.1681/ASN.2011030298
- Hu, M., Miller, E. J., Lin, X., and Simms, H. H. (2004). Transmigration across a lung epithelial monolayer delays apoptosis of polymorphonuclear leukocytes. *Surgery* 135, 87–98. doi: 10.1016/S0039-6060(03)00347-7
- Humphreys, J. M., Hughes, V., and Edwards, S. W. (1989). Stimulation of protein synthesis in human neutrophils by gamma-interferon. *Biochem. Pharmacol.* 38, 1241–1246. doi: 10.1016/0006-2952(89)90329-8
- Jablonska, E., Izycka, A., and Wawrusiewicz, N. (2002a). Effect of IL-18 on IL-1 β and sIL-1RII production by human neutrophils. *Arch. Immunol. Ther. Exp.* 50, 139–141.
- Jablonska, E., Kiluk, M., Markiewicz, W., and Jablonski, J. (2002b). Priming effects of GM-CSF, IFN- γ and TNF- α on human neutrophil inflammatory cytokine production. *Melanoma Res.* 12, 123–128. doi: 10.1097/00008390-200204000-00004
- Jablonska, E., Piotrowski, L., Kiluk, M., Jablonski, J., Grabowska, Z., and Markiewicz, W. (2001). Effect of IL-15 on the secretion of IL-1 β , IL-1Ra and sIL-1RII by PMN from cancer patients. *Cytokine* 16, 173–177. doi: 10.1006/cyto.2001.0931
- Jagels, M. A., Daffern, P. J., and Hugli, T. E. (2000). C3a and C5a enhance granulocyte adhesion to endothelial and epithelial cell monolayers: epithelial and endothelial priming is required for C3a-induced eosinophil adhesion. *Immunopharmacology* 46, 209–222. doi: 10.1016/S0162-3109(99)00178-2
- Johnson, B. L. III, Kueth, J. W., and Caldwell, C. C. (2014). Neutrophil derived microvesicles: emerging role of a key mediator to the immune response. *Endocr. Metab. Immune Disord. Drug Targets* 14, 210–217. doi: 10.2174/1871530314666140722083717
- Kannan, S. (2002). Amplification of extracellular nucleotide-induced leukocyte(s) degranulation by contingent autocrine and paracrine mode of leukotriene-mediated chemokine receptor activation. *Med. Hypotheses* 59, 261–265. doi: 10.1016/S0306-9877(02)00213-X
- Kannengiesser, C., Gerard, B., El Benna, J., Henri, D., Kroviarski, Y., Chollet-Martin, S., et al. (2008). Molecular epidemiology of chronic granulomatous disease in a series of 80 kindreds: identification of 31 novel mutations. *Hum. Mutat.* 29, E132–E149. doi: 10.1002/humu.20820
- Kasorn, A., Alcaide, P., Jia, Y., Subramanian, K. K., Sarraj, B., Li, Y., et al. (2009). Focal adhesion kinase regulates pathogen-killing capability and life span of neutrophils via mediating both adhesion-dependent and -independent cellular signals. *J. Immunol.* 183, 1032–1043. doi: 10.4049/jimmunol.0802984
- Kessenbrock, K., Krumbholz, M., Schonermarck, U., Back, W., Gross, W. L., Werb, Z., et al. (2009). Netting neutrophils in autoimmune small-vessel vasculitis. *Nat. Med.* 15, 623–625. doi: 10.1038/nm.1959
- Khreiss, T., Jozsef, L., Chan, J. S., and Filep, J. G. (2004). Activation of extracellular signal-regulated kinase couples platelet-activating factor-induced adhesion and delayed apoptosis of human neutrophils. *Cell. Signal.* 16, 801–810. doi: 10.1016/j.cellsig.2003.12.005
- Klebanoff, S. J., Olszowski, S., Van Voorhis, W. C., Ledbetter, J. A., Waltersdorff, A. M., and Schlechte, K. G. (1992). Effects of gamma-interferon on human neutrophils: protection from deterioration on storage. *Blood* 80, 225–234.
- Klein, J. B., Buridi, A., Coxon, P. Y., Rane, M. J., Manning, T., Kettritz, R., et al. (2001). Role of extracellular signal-regulated kinase and phosphatidylinositol-3 kinase in chemoattractant and LPS delay of constitutive neutrophil apoptosis. *Cell. Signal.* 13, 335–343. doi: 10.1016/S0898-6568(01)00151-6
- Klein, J. B., Rane, M. J., Scherzer, J. A., Coxon, P. Y., Kettritz, R., Mathiesen, J. M., et al. (2000). Granulocyte-macrophage colony-stimulating factor delays neutrophil constitutive apoptosis through phosphoinositide 3-kinase and extracellular signal-regulated kinase pathways. *J. Immunol.* 164, 4286–4291. doi: 10.4049/jimmunol.164.8.4286
- Kletter, Y., Bleiberg, I., Golde, D. W., and Fabian, I. (1989). Antibody to Mol abrogates the increase in neutrophil phagocytosis and degranulation induced by granulocyte-macrophage colony-stimulating factor. *Eur. J. Haematol.* 43, 389–396. doi: 10.1111/j.1600-0609.1989.tb00325.x
- Klimchenko, O., Di Stefano, A., Georger, B., Hamidi, S., Opolon, P., Robert, T., et al. (2011). Monocytic cells derived from human embryonic stem cells and fetal liver share common differentiation pathways and homeostatic functions. *Blood* 117, 3065–3075. doi: 10.1182/blood-2010-07-295246
- Kowanko, I. C., Ferrante, A., Harvey, D. P., and Carman, K. L. (1991). Granulocyte-macrophage colony-stimulating factor augments neutrophil killing of *Torulopsis glabrata* and stimulates neutrophil

- respiratory burst and degranulation. *Clin. Exp. Immunol.* 83, 225–230. doi: 10.1111/j.1365-2249.1991.tb05619.x
- Kulkarni, S., Woollard, K. J., Thomas, S., Oxley, D., and Jackson, S. P. (2007). Conversion of platelets from a proaggregatory to a proinflammatory adhesive phenotype: role of PAF in spatially regulating neutrophil adhesion and spreading. *Blood* 110, 1879–1886. doi: 10.1182/blood-2006-08-040980
- Lamb, F. S., Hook, J. S., Hilkin, B. M., Huber, J. N., Volk, A. P., and Moreland, J. G. (2012). Endotoxin priming of neutrophils requires endocytosis and NADPH oxidase-dependent endosomal reactive oxygen species. *J. Biol. Chem.* 287, 12395–12404. doi: 10.1074/jbc.M111.306530
- Lammermann, T., Afonso, P. V., Angermann, B. R., Wang, J. M., Kastentmuller, W., Parent, C. A., et al. (2013). Neutrophil swarms require LTB4 and integrins at sites of cell death *in vivo*. *Nature*, 498, 371–375. doi: 10.1038/nature12175
- Lan, F., Yuan, B., Liu, T., Luo, X., Huang, P., Liu, Y., et al. (2016). Interleukin-33 facilitates neutrophil recruitment and bacterial clearance in *S. aureus*-caused peritonitis. *Mol. Immunol.* 72, 74–80. doi: 10.1016/j.molimm.2016.03.004
- Le, H. T., Tran, V. G., Kim, W., Kim, J., Cho, H. R., and Kwon, B. (2012). IL-33 priming regulates multiple steps of the neutrophil-mediated anti-*Candida albicans* response by modulating TLR and dectin-1 signals. *J. Immunol.* 189, 287–295. doi: 10.4049/jimmunol.1103564
- Leslie, C. C. (2004). Regulation of arachidonic acid availability for eicosanoid production. *Biochem. Cell Biol.* 82, 1–17. doi: 10.1139/o03-080
- Liles, W. C., Ledbetter, J. A., Waltersdorff, A. W., and Klebanoff, S. J. (1995). Cross-linking of CD18 primes human neutrophils for activation of the respiratory burst in response to specific stimuli: implications for adhesion-dependent physiological responses in neutrophils. *J. Leukoc. Biol.* 58, 690–697.
- Lindemann, A., Riedel, D., Oster, W., Meuer, S. C., Blohm, D., Mertelsmann, R. H., et al. (1988). Granulocyte/macrophage colony-stimulating factor induces interleukin 1 production by human polymorphonuclear neutrophils. *J. Immunol.* 140, 837–839.
- Liu, J. J., Song, C. W., Yue, Y., Duan, C. G., Yang, J., He, T., et al. (2005). Quercetin inhibits LPS-induced delay in spontaneous apoptosis and activation of neutrophils. *Inflamm. Res.* 54, 500–507. doi: 10.1007/s00011-005-1385-2
- Lominadze, G., Powell, D. W., Luerman, G. C., Link, A. J., Ward, R. A., and McLeish, K. R. (2005). Proteomic analysis of human neutrophil granules. *Mol. Cell. Proteomics* 4, 1503–1521. doi: 10.1074/mcp.M500143-MCP200
- Lorincz, A. M., Schutte, M., Timar, C. I., Veres, D. S., Kittel, A., McLeish, K. R., et al. (2015). Functionally and morphologically distinct populations of extracellular vesicles produced by human neutrophilic granulocytes. *J. Leukoc. Biol.* 98, 583–589. doi: 10.1189/jlb.3VMA1014-514R
- Luo, M., Jones, S. M., Peters-Golden, M., and Brock, T. G. (2003). Nuclear localization of 5-lipoxygenase as a determinant of leukotriene B4 synthetic capacity. *Proc. Natl. Acad. Sci. U.S.A.* 100, 12165–12170. doi: 10.1073/pnas.2133253100
- Makni-Maalej, K., Boussetta, T., Hurtado-Nedelec, M., Belambri, S. A., Gougerot-Pocidallo, M. A., and El-Benna, J. (2012). The TLR7/8 agonist CL097 primes N-formyl-methionyl-leucyl-phenylalanine-stimulated NADPH oxidase activation in human neutrophils: critical role of p47phox phosphorylation and the proline isomerase Pin1. *J. Immunol.* 189, 4657–4665. doi: 10.4049/jimmunol.1201007
- Makni-Maalej, K., Marzaioli, V., Boussetta, T., Belambri, S. A., Gougerot-Pocidallo, M. A., Hurtado-Nedelec, M., et al. (2015). TLR8, but not TLR7, induces the priming of the NADPH oxidase activation in human neutrophils. *J. Leukoc. Biol.* 97, 1081–1087. doi: 10.1189/jlb.2A1214-623R
- Mancuso, P., Nana-Sinkam, P., and Peters-Golden, M. (2001). Leukotriene B4 augments neutrophil phagocytosis of *Klebsiella pneumoniae*. *Infect. Immun.* 69, 2011–2016. doi: 10.1128/IAI.69.4.2011-2016.2001
- Marasco, W. A., Showell, H. J., and Becker, E. L. (1981). Substance P binds to the formylpeptide chemotaxis receptor on the rabbit neutrophil. *Biochem. Biophys. Res. Commun.* 99, 1065–1072. doi: 10.1016/0006-291X(81)90727-0
- Marucha, P. T., Zeff, R. A., and Kreutzer, D. L. (1991). Cytokine-induced IL-1 beta gene expression in the human polymorphonuclear leukocyte: transcriptional and post-transcriptional regulation by tumor necrosis factor and IL-1. *J. Immunol.* 147, 2603–2608.
- Mastroianni, C. M., d'Ettore, G., Forcina, G., Lichtner, M., Mengoni, F., D'Agostino, C., et al. (2000). Interleukin-15 enhances neutrophil functional activity in patients with human immunodeficiency virus infection. *Blood* 96, 1979–1984.
- Mayadas, T. N., and Cullere, X. (2005). Neutrophil beta2 integrins: moderators of life or death decisions. *Trends Immunol.* 26, 388–395. doi: 10.1016/j.it.2005.05.002
- Mayer, T. Z., Simard, F. A., Cloutier, A., Vardhan, H., Dubois, C. M., and McDonald, P. P. (2013). The p38-MSK1 signaling cascade influences cytokine production through CREB and C/EBP factors in human neutrophils. *J. Immunol.* 191, 4299–4307. doi: 10.4049/jimmunol.1301117
- McGettrick, H. M., Lord, J. M., Wang, K. Q., Rainger, G. E., Buckley, C. D., and Nash, G. B. (2006). Chemokine- and adhesion-dependent survival of neutrophils after transmigration through cytokine-stimulated endothelium. *J. Leukoc. Biol.* 79, 779–788. doi: 10.1189/jlb.0605350
- McLeish, K. R., Klein, J. B., Lederer, E. D., Head, K. Z., and Ward, R. A. (1996). Azotemia, TNF alpha, and LPS prime the human neutrophil oxidative burst by distinct mechanisms. *Kidney Int.* 50, 407–416.
- McLeish, K. R., Knall, C., Ward, R. A., Gerwings, P., Coxon, P. Y., Klein, J. B., et al. (1998). Activation of mitogen-activated protein kinase cascades during priming of human neutrophils by TNF-alpha and GM-CSF. *J. Leukoc. Biol.* 64, 537–545.
- McLeish, K. R., Merchant, M. L., Creed, T. M., Tandon, S., Barati, M. T., Uriarte, S. M., et al. (2017). Frontline Science: tumor necrosis factor- α stimulation and priming of human neutrophil granule exocytosis. *J. Leukoc. Biol.* doi: 10.1189/jlb.3HI0716-293RR. [Epub ahead of print].
- McLeish, K. R., Uriarte, S. M., Tandon, S., Creed, T. M., Le, J., and Ward, R. A. (2013). Exocytosis of neutrophil granule subsets and activation of prolyl isomerase 1 are required for respiratory burst priming. *J. Innate Immun.* 5, 277–289. doi: 10.1159/000345992
- Melby, K., Midtvedt, T., and Degre, M. (1982). Effect of human leukocyte interferon on phagocytic activity of polymorphonuclear leukocytes. *Acta Pathol. Microbiol. Immunol. Scand. B* 90, 181–184. doi: 10.1111/j.1699-0463.1982.tb00102.x
- Meshki, J., Tuluc, F., Bredeteau, O., Ding, Z., and Kunapuli, S. P. (2004). Molecular mechanism of nucleotide-induced primary granule release in human neutrophils: role for the P2Y2 receptor. *Am. J. Physiol. Cell Physiol.* 286, C264–C271. doi: 10.1152/ajpcell.00287.2003
- Mesri, M., and Altieri, D. C. (1998). Endothelial cell activation by leukocyte microparticles. *J. Immunol.* 161, 4382–4387.
- Mocsai, A., Ligeti, E., Lowell, C. A., and Berton, G. (1999). Adhesion-dependent degranulation of neutrophils requires the Src family kinases Fgr and Hck. *J. Immunol.* 162, 1120–1126.
- Montecucco, F., Steffens, S., Burger, F., Da Costa, A., Bianchi, G., Bertolotto, M., et al. (2008). Tumor necrosis factor-alpha (TNF-alpha) induces integrin CD11b/CD18 (Mac-1) up-regulation and migration to the CC chemokine CCL3 (MIP-1alpha) on human neutrophils through defined signalling pathways. *Cell. Signal.* 20, 557–568. doi: 10.1016/j.cellsig.2007.11.008
- Moreland, J. G., Davis, A. P., Matsuda, J. J., Hook, J. S., Bailey, G., Nauseef, W. M., et al. (2007). Endotoxin priming of neutrophils requires NADPH oxidase-generated oxidants and is regulated by the anion transporter ClC-3. *J. Biol. Chem.* 282, 33958–33967. doi: 10.1074/jbc.M705289200
- Morris, A. C., Brittan, M., Wilkinson, T. S., McAuley, D. F., Antonelli, J., McCulloch, C., et al. (2011). C5a-mediated neutrophil dysfunction is RhoA-dependent and predicts infection in critically ill patients. *Blood* 117, 5178–5188. doi: 10.1182/blood-2010-08-304667
- Moutsopoulos, N. M., Konkel, J., Sarmadi, M., Eskan, M. A., Wild, T., Dutzan, N., et al. (2014). Defective neutrophil recruitment in leukocyte adhesion deficiency type I disease causes local IL-17-driven inflammatory bone loss. *Sci. Transl. Med.* 6, 229ra40. doi: 10.1126/scitranslmed.3007696
- Murray, J., Barbara, J. A., Dunkley, S. A., Lopez, A. F., Van Ostade, X., Condliffe, A. M., et al. (1997). Regulation of neutrophil apoptosis by tumor necrosis factor-alpha: requirement for TNFR55 and TNFR75 for induction of apoptosis *in vitro*. *Blood* 90, 2772–2783.
- Musso, T., Calosso, L., Zucca, M., Millesimo, M., Puliti, M., Bulfone-Paus, S., et al. (1998). Interleukin-15 activates proinflammatory and antimicrobial functions in polymorphonuclear cells. *Infect. Immun.* 66, 2640–2647.
- Naegele, M., Tillack, K., Reinhardt, S., Schippling, S., Martin, R., and Sospedra, M. (2012). Neutrophils in multiple sclerosis are characterized by a primed phenotype. *J. Neuroimmunol.* 242, 60–71. doi: 10.1016/j.jneuroim.2011.11.009

- Nathan, C. (2006). Neutrophils and immunity: challenges and opportunities. *Nat. Rev. Immunol.* 6, 173–182. doi: 10.1038/nri1785
- Nauseef, W. M., and Borregaard, N. (2014). Neutrophils at work. *Nat. Immunol.* 15, 602–611. doi: 10.1038/ni.2921
- Nemeth, T., and Mocsai, A. (2016). Feedback amplification of neutrophil function. *Trends Immunol.* 37, 412–424. doi: 10.1016/j.it.2016.04.002
- Nolan, B., Collette, H., Baker, S., Duffy, A., De, M., Miller, C., et al. (2000). Inhibition of neutrophil apoptosis after severe trauma is NF κ B dependent. *J. Trauma* 48, 599–604; discussion: 604–595.
- O'Flaherty, J. T., Showell, H. J., Becker, E. L., and Ward, P. A. (1979). Neutrophil aggregation and degranulation. Effect of arachidonic acid. *Am. J. Pathol.* 95, 433–444.
- Ogura, H., Tanaka, H., Koh, T., Hashiguchi, N., Kuwagata, Y., Hosotsubo, H., et al. (1999). Priming, second-hit priming, and apoptosis in leukocytes from trauma patients. *J. Trauma* 46, 774–781; discussion: 781–773. doi: 10.1097/00005373-199905000-00004
- Palmantier, R., Surette, M. E., Sanchez, A., Braquet, P., and Borgeat, P. (1994). Priming for the synthesis of 5-lipoxygenase products in human blood *ex vivo* by human granulocyte-macrophage colony-stimulating factor and tumor necrosis factor- α . *Lab. Invest.* 70, 696–704.
- Paulsson, J. M., Jacobson, S. H., and Lundahl, J. (2010). Neutrophil activation during transmigration *in vivo* and *in vitro* A translational study using the skin chamber model. *J. Immunol. Methods* 361, 82–88. doi: 10.1016/j.jim.2010.07.015
- Perianayagam, M. C., Balakrishnan, V. S., King, A. J., Pereira, B. J., and Jaber, B. L. (2002). C5a delays apoptosis of human neutrophils by a phosphatidylinositol 3-kinase-signaling pathway. *Kidney Int.* 61, 456–463. doi: 10.1046/j.1523-1755.2002.00139.x
- Perianin, A., Snyderman, R., and Malfroy, B. (1989). Substance P primes human neutrophil activation: a mechanism for neurological regulation of inflammation. *Biochem. Biophys. Res. Commun.* 161, 520–524. doi: 10.1016/0006-291X(89)92630-2
- Perussia, B., Kobayashi, M., Rossi, M. E., Anegón, I., and Trinchieri, G. (1987). Immune interferon enhances functional properties of human granulocytes: role of Fc receptors and effect of lymphotoxin, tumor necrosis factor, and granulocyte-macrophage colony-stimulating factor. *J. Immunol.* 138, 765–774.
- Pliyev, B. K., Kalintseva, M. V., Abdulaeva, S. V., Yarygin, K. N., and Savchenko, V. G. (2014). Neutrophil microparticles modulate cytokine production by natural killer cells. *Cytokine* 65, 126–129. doi: 10.1016/j.cyto.2013.11.010
- Pluskota, E., Woody, N. M., Szpak, D., Ballantyne, C. M., Soloviev, D. A., Simon, D. I., et al. (2008). Expression, activation, and function of integrin α M β 2 (Mac-1) on neutrophil-derived microparticles. *Blood* 112, 2327–2335. doi: 10.1182/blood-2007-12-127183
- Podaza, E., Sabbione, F., Risnik, D., Borge, M., Almejun, M. B., Colado, A., et al. (2016). Neutrophils from chronic lymphocytic leukemia patients exhibit an increased capacity to release extracellular traps (NETs). *Cancer Immunol. Immunother.* 66, 77–89. doi: 10.1007/s00262-016-1921-7
- Potera, R. M., Jensen, M. J., Hilkin, B. M., South, G. K., Hook, J. S., Gross, E. A., et al. (2016). Neutrophil azurophilic granule exocytosis is primed by TNF- α and partially regulated by NADPH oxidase. *Innate Immun.* 22, 635–646. doi: 10.1177/1753425916668980
- Rainard, P., Rioulet, C., Poutrel, B., and Paape, M. J. (2000). Phagocytosis and killing of *Staphylococcus aureus* by bovine neutrophils after priming by tumor necrosis factor- α and the des-arginine derivative of C5a. *Am. J. Vet. Res.* 61, 951–959. doi: 10.2460/ajvr.2000.61.951
- Ramadass, M., Johnson, J. L., and Catz, S. D. (2017). Rab27a regulates GM-CSF-dependent priming of neutrophil exocytosis. *J. Leukoc. Biol.* 101, 693–702. doi: 10.1189/jlb.3AB0416-189RR
- Richardson, M. D., and Patel, M. (1995). Stimulation of neutrophil phagocytosis of *Aspergillus fumigatus* conidia by interleukin-8 and N-formylmethionyl-leucylphenylalanine. *J. Med. Vet. Mycol.* 33, 99–104. doi: 10.1080/02681219580000221
- Roberge, C. J., McColl, S. R., Larochelle, B., and Gosselin, J. (1998). Granulocyte-macrophage colony-stimulating factor enhances EBV-induced synthesis of chemotactic factors in human neutrophils. *J. Immunol.* 160, 2442–2448.
- Rørvig, S., Øtergaard, O., Heegaard, N. H., and Borregaard, N. (2013). Proteome profiling of human neutrophil granule subsets, secretory vesicles, and cell membrane: correlation with transcriptome profiling of neutrophil precursors. *J. Leukoc. Biol.* 94, 711–721. doi: 10.1189/jlb.1212619
- Rosales, C., and Brown, E. J. (1991). Two mechanisms for IgG Fc-receptor-mediated phagocytosis by human neutrophils. *J. Immunol.* 146, 3937–3944.
- Sabroe, I., Prince, L. R., Jones, E. C., Horsburgh, M. J., Foster, S. J., Vogel, S. N., et al. (2003). Selective roles for Toll-like receptor (TLR)2 and TLR4 in the regulation of neutrophil activation and life span. *J. Immunol.* 170, 5268–5275. doi: 10.4049/jimmunol.170.10.5268
- Salamone, G. V., Petracca, Y., Fuxman Bass, J. I., Rumbo, M., Nahmod, K. A., Gabelloni, M. L., et al. (2010). Flagellin delays spontaneous human neutrophil apoptosis. *Lab. Invest.* 90, 1049–1059. doi: 10.1038/labinvest.2010.77
- Schatz-Munding, M., and Ullrich, V. (1992). Priming of human polymorphonuclear leukocytes with granulocyte-macrophage colony-stimulating factor involves protein kinase C rather than enhanced calcium mobilisation. *Eur. J. Biochem.* 204, 705–712. doi: 10.1111/j.1432-1033.1992.tb16685.x
- Seeds, M. C., Jones, D. F., Chilton, F. H., and Bass, D. A. (1998). Secretory and cytosolic phospholipases A2 are activated during TNF priming of human neutrophils. *Biochim. Biophys. Acta* 1389, 273–284. doi: 10.1016/S0005-2760(97)00151-3
- Sengelov, H., Follin, P., Kjeldsen, L., Løllike, K., Dahlgren, C., and Borregaard, N. (1995). Mobilization of granules and secretory vesicles during *in vivo* exudation of human neutrophils. *J. Immunol.* 154, 4157–4165.
- Sengelov, H., Kjeldsen, L., and Borregaard, N. (1993). Control of exocytosis in early neutrophil activation. *J. Immunol.* 150, 1535–1543.
- Shalit, M., Dabiri, G. A., and Southwick, F. S. (1987). Platelet-activating factor both stimulates and “primes” human polymorphonuclear leukocyte actin filament assembly. *Blood* 70, 1921–1927.
- Sheppard, F. R., Kelher, M. R., Moore, E. E., McLaughlin, N. J., Banerjee, A., and Silliman, C. C. (2005). Structural organization of the neutrophil NADPH oxidase: phosphorylation and translocation during priming and activation. *J. Leukoc. Biol.* 78, 1025–1042. doi: 10.1189/jlb.0804442
- Sorensen, O. E., and Borregaard, N. (2016). Neutrophil extracellular traps - the dark side of neutrophils. *J. Clin. Invest.* 126, 1612–1620. doi: 10.1172/JCI84538
- Stanislawski, L., Huu, T. P., and Perianin, A. (1990). Priming effect of fibronectin on respiratory burst of human neutrophils induced by formyl peptides and platelet-activating factor. *Inflammation* 14, 523–530. doi: 10.1007/BF00914273
- Steadman, R., Petersen, M. M., and Williams, J. D. (1996). CD11b/CD18-dependent stimulation of leukotriene B4 synthesis by human neutrophils (PMN) is synergistically enhanced by tumour necrosis factor alpha and low dose diacylglycerol. *Int. J. Biochem. Cell Biol.* 28, 771–776. doi: 10.1016/1357-2725(96)00015-5
- Summers, C., Rankin, S. M., Condliffe, A. M., Singh, N., Peters, A. M., and Chilvers, E. R. (2010). Neutrophil kinetics in health and disease. *Trends Immunol.* 31, 318–324. doi: 10.1016/j.it.2010.05.006
- Surette, M. E., Dallaire, N., Jean, N., Picard, S., and Borgeat, P. (1998). Mechanisms of the priming effect of lipopolysaccharides on the biosynthesis of leukotriene B4 in chemotactic peptide-stimulated human neutrophils. *FASEB J.* 12, 1521–1531.
- Surette, M. E., Palmantier, R., Gosselin, J., and Borgeat, P. (1993). Lipopolysaccharides prime whole human blood and isolated neutrophils for the increased synthesis of 5-lipoxygenase products by enhancing arachidonic acid availability: involvement of the CD14 antigen. *J. Exp. Med.* 178, 1347–1355. doi: 10.1084/jem.178.4.1347
- Swain, S. D., Rohn, T. T., and Quinn, M. T. (2002). Neutrophil priming in host defense: role of oxidants as priming agents. *Antioxid. Redox Signal.* 4, 69–83. doi: 10.1089/152308602753625870
- Timar, C. I., Lorincz, A. M., Csepanyi-Kömi, R., Valyi-Nagy, A., Nagy, G., Buzas, E. I., et al. (2013a). Antibacterial effect of microvesicles released from human neutrophilic granulocytes. *Blood* 121, 510–518. doi: 10.1182/blood-2012-05-431114
- Timar, C. I., Lorincz, A. M., and Ligeti, E. (2013b). Changing world of neutrophils. *Pflugers Arch.* 465, 1521–1533. doi: 10.1007/s00424-013-1285-1
- Tsuboi, N., Hernandez, T., Li, X., Nishi, H., Cullere, X., Mekala, D., et al. (2011). Regulation of human neutrophil Fc γ receptor IIa by C5a receptor promotes inflammatory arthritis in mice. *Arthritis Rheum.* 63, 467–478. doi: 10.1002/art.30141

- Uriarte, S. M., Rane, M. J., Luerman, G. C., Barati, M. T., Ward, R. A., Nauseef, W. M., et al. (2011). Granule exocytosis contributes to priming and activation of the human neutrophil respiratory burst. *J. Immunol.* 187, 391–400. doi: 10.4049/jimmunol.1003112
- van de Vijver, E., Maddalena, A., Sanal, O., Holland, S. M., Uzel, G., Madkaikar, M., et al. (2012). Hematologically important mutations: leukocyte adhesion deficiency (first update). *Blood Cells Mol. Dis.* 48, 53–61. doi: 10.1016/j.bcmd.2011.10.004
- Volk, A. P., Barber, B. M., Goss, K. L., Ruff, J. G., Heise, C. K., Hook, J. S., et al. (2011). Priming of neutrophils and differentiated PLB-985 cells by pathophysiological concentrations of TNF-alpha is partially oxygen dependent. *J. Innate Immun.* 3, 298–314. doi: 10.1159/000321439
- Ward, R. A., Nakamura, M., and McLeish, K. R. (2000). Priming of the neutrophil respiratory burst involves p38 mitogen-activated protein kinase-dependent exocytosis of flavocytochrome b558-containing granules. *J. Biol. Chem.* 275, 36713–36719. doi: 10.1074/jbc.M003017200
- Whitmore, L. C., Hook, J. S., Philip, A. R., Hilkin, B. M., Bing, X., Ahn, C., et al. (2016). A common genetic variant in TLR1 enhances human neutrophil priming and impacts length of intensive care stay in pediatric sepsis. *J. Immunol.* 196, 1376–1386. doi: 10.4049/jimmunol.1500856
- Wozniak, A., McLennan, G., Betts, W. H., Murphy, G. A., and Scicchitano, R. (1989). Activation of human neutrophils by substance P: effect on FMLP-stimulated oxidative and arachidonic acid metabolism and on antibody-dependent cell-mediated cytotoxicity. *Immunology* 68, 359–364.
- Wright, H. L., Cross, A. L., Edwards, S. W., and Moots, R. J. (2014). Effects of IL-6 and IL-6 blockade on neutrophil function *in vitro* and *in vivo*. *Rheumatology* 53, 1321–1331. doi: 10.1093/rheumatology/keu035
- Wright, H. L., Thomas, H. B., Moots, R. J., and Edwards, S. W. (2013). RNA-seq reveals activation of both common and cytokine-specific pathways following neutrophil priming. *PLoS ONE* 8:e58598. doi: 10.1371/journal.pone.0058598
- Wyman, T. H., Dinarello, C. A., Banerjee, A., Gamboni-Robertson, F., Hiester, A. A., England, K. M., et al. (2002). Physiological levels of interleukin-18 stimulate multiple neutrophil functions through p38 MAP kinase activation. *J. Leukoc. Biol.* 72, 401–409.
- Xu, X., and Hakansson, L. (2002). Degranulation of primary and secondary granules in adherent human neutrophils. *Scand. J. Immunol.* 55, 178–188. doi: 10.1046/j.1365-3083.2002.01041.x
- Yao, Y., Matsushima, H., Ohtola, J. A., Geng, S., Lu, R., and Takashima, A. (2015). Neutrophil priming occurs in a sequential manner and can be visualized in living animals by monitoring IL-1beta promoter activation. *J. Immunol.* 194, 1211–1224. doi: 10.4049/jimmunol.1402018
- Yousefi, S., Mihalache, C., Kozlowski, E., Schmid, I., and Simon, H. U. (2009). Viable neutrophils release mitochondrial DNA to form neutrophil extracellular traps. *Cell Death Differ.* 16, 1438–1444. doi: 10.1038/cdd.2009.96
- Yuo, A., Kitagawa, S., Ohsaka, A., Saito, M., and Takaku, F. (1990). Stimulation and priming of human neutrophils by granulocyte colony-stimulating factor and granulocyte-macrophage colony-stimulating factor: qualitative and quantitative differences. *Biochem. Biophys. Res. Commun.* 171, 491–497. doi: 10.1016/0006-291X(90)91420-W
- Zallen, G., Moore, E. E., Johnson, J. L., Tamura, D. Y., Aiboshi, J., Biffl, W. L., et al. (1999). Circulating postinjury neutrophils are primed for the release of proinflammatory cytokines. *J. Trauma* 46, 42–48. doi: 10.1097/00005373-199901000-00007
- Zarini, S., Gijon, M. A., Folco, G., and Murphy, R. C. (2006). Effect of arachidonic acid reacylation on leukotriene biosynthesis in human neutrophils stimulated with granulocyte-macrophage colony-stimulating factor and formyl-methionyl-leucyl-phenylalanine. *J. Biol. Chem.* 281, 10134–10142. doi: 10.1074/jbc.M510783200

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Miralda, Uriarte and McLeish. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.