

## LITERATURE CITED

1. ASSOCIATION OF OFFICIAL AGRICULTURAL CHEMISTS. Official and tentative methods of analyses. Method 6.59, 114. Ed. 8, Washington, D. C. 1955.
2. ATTOE, O. J. Leaf-burn of tobacco as influenced by content of potassium, nitrogen, and chlorine. *Jour. Amer. Soc. Agron.* 38: 186–196. 1946.
3. BEHRENS, J. Weitere beitrage zur kenntnis der tabakpflanze. V. Der anatomische ban und der destandteile des tabakblattes in ihrer beziehung zur brennbarkeit. *Landwirtsch. Vers. Sta.* 43: 271–301. 1894.
4. BROYER, T. C., CARLTON, A. B., JOHNSON, C. M. and STOUT, P. R. Chlorine, a micronutrient element for higher plants. *Plant Physiol.* 29: 526–532. 1954.
5. EATON, F. M. Toxicity and accumulation of chloride and sulfate salts in plants. *Jour. Agr. Res.* 64: 357–399. 1942.
6. GARNER, W. W., McMURTREY, J. E., JR., BOWLING, J. D. and MOSS, E. G. Role of chlorine in nutrition and growth in the tobacco plant and its effect on the quality of the cured leaf. *Jour. Agr. Res.* 40: 627–648. 1930.
7. GAUCH, H. G. and EATON, F. M. Effect of saline substrate on hourly levels of carbohydrates and inorganic constituents of barley plants. *Plant Physiol.* 17: 347–365. 1942.
8. HAYWARD, H. E. and LONG, E. M. Anatomical and physiological responses of the tomato to varying concentrations of sodium chloride, sodium sulphate and nutrient solutions. *Bot. Gaz.* 102: 437–462. 1941.
9. HAYWARD, H. E., and LONG, E. M. Some effects of sodium salts on the growth of the tomato. *Plant Physiol.* 18: 556–569. 1943.
10. HOAGLAND, D. R. Relation of the concentration and reaction of the nutrient medium to the growth and absorption of the plant. *Jour. Agr. Res.* 18: 73–117. 1919.
11. HOAGLAND, D. R. The absorption of ions by plants. *Soil Sci.* 16: 225–246. 1923.
12. HOAGLAND, D. R. and ARNON, D. I. The water culture method of growing plants without soil. *Agr. Expt. Sta., California, Cir.* 347. 1938.
13. JOHNSON, J., OGDEN, W. B. and ATTOE, O. J. Experiments on the leaf-burn of tobacco. *Agr. Expt. Sta., Wisconsin, Res. Bull.* 153. 1944.
14. KRETSCHMER, A. E., TOTH, S. S. and BEAR, F. E. Effect of chloride versus sulfate ions on nutrient-ion absorption by plants. *Soil Sci.* 76: 193–199. 1953.
15. McEVoy, E. T. Studies on the optimum nutrition of flue-cured tobacco. *Sci. Agr.* 25: 489–498. 1945.
16. McEVoy, E. T. The relation of ammonium and sulphate ions to magnesium deficiency in tobacco. *Can. Jour. of Agr. Sci.* 34: 281–287. 1954.
17. McMURTREY, J. E. Nutritional deficiency studies on tobacco. *Jour. Amer. Soc. Agron.* 21: 142–149. 1929.
18. MYRHRE, D. L., ATTOE, O. J. and OGDEN, W. B. Chlorine and other constituents in relation to tobacco leaf-burn. *Soil Sci. Soc. of Amer. Proc.* 20: 547–551. 1956.
19. NEAS, IVAN. Sulphur nutrition in flue-cured tobacco. *Agron. Jour.* 45: 472–477. 1953.
20. REISENAUER, H. M. and COLWELL, W. E. Some factors affecting the absorption of chlorine by tobacco. *Soil Sci. Soc. of Amer. Proc.* 15: 222–229. 1950.
21. ULRICH, A. and OHKI, K. Chlorine, bromine and sodium as nutrients for sugar beet plants. *Plant Physiol.* 31: 171–181. 1956.

## MULTIPLE SITE UPTAKE OF INDIVIDUAL CATIONS BY ROOTS AS AFFECTED BY HYDROGEN ION<sup>1</sup>

MAURICE FRIED AND J. C. NOGGLE

SOIL AND WATER CONSERVATION RESEARCH DIVISION, AGRICULTURAL RESEARCH SERVICE,  
U. S. DEPARTMENT OF AGRICULTURE, BELTSVILLE, MARYLAND

Many investigators have suggested that metabolic uptake of ions by cells involves a system of carriers. A comprehensive review is given by Epstein (6). The ion combines with the carrier to form an active intermediate. The active intermediate breaks down essentially irreversibly to release the ion inside the cell.

Cations that are similar chemically may compete with each other for the same binding site. However different carriers are involved for different groups of cations. Thus Collander (4), and Epstein and Hagen (7), and Menzel and Heald (19), have indicated that rubidium and cesium and potassium are competitive for the same site. Although calcium utilizes a different site than potassium, Epstein and Leggett


(8), and Menzel and Heald (19) have shown that strontium and barium compete with calcium. Neither sodium nor magnesium was found to be competitive for either of these sites.

Over a limited concentration range Epstein and Hagen (7), and Epstein and Leggett (8) noted one carrier site for each cation. Epstein and Hagen noted an anomaly at higher ion concentrations with sodium and Epstein and Leggett an anomaly with strontium. Over a wide concentration range Hagen and Hopkins (12) demonstrated two carrier sites for phosphate uptake.

Various investigators (14, 15, 16, 21) have shown that the hydrogen ion concentration of the bathing medium affects ion uptake. Others (1, 2, 3, 11) have minimized this effect within usual physiological ranges. Jacobsen et al (17), using the model of Ro-



<sup>1</sup> Received November 18, 1957.

senberg (22), suggested that the cation carrier is metabolically produced at the external surfaces as the hydrogen complex, HR, and that the anion carriers were metabolically produced at the external surfaces as the hydroxyl complex R'OH. They formalized this in equations a and b,



where HR and R'OH represent the metabolically produced cationic and anionic binding substances. MR and R'A serve as transporting agents, transferring ions from culture medium to the vacuole (17).

On the other hand, Epstein and Hagen (7) and Hagen and Hopkins (12) have described ion uptake by carriers without any detailed assumptions as to the nature of the carrier. Ion absorption is described by these authors in the following equations.



where R represents a metabolically produced site, M, the ion, MR, the active intermediate and k the rate constant for each reaction.

Hagen and Hopkins (12) have shown by kinetic analysis that hydroxyl ion is a competitive inhibitor of phosphorus uptake and behaves similarly to other competitive inhibitors such as ADP and pyrophosphate in phosphate uptake (13) and selenate in sulfate uptake (18).

The competitive inhibition of phosphate uptake by hydroxyl ion is consistent with the uptake mechanism described by both Hagen and Hopkins (12) and Jacobsen et al (17). However, the present authors prefer the description afforded in equations 1 and 2 since it makes no assumptions as to the location of the carrier nor the specific combination in which the carrier occurs. By analogous interpretation from kinetic analysis as developed for ion absorption (7, 12), the present results demonstrate that hydrogen ion is a competitive inhibitor of cation uptake. The cations used were sodium, potassium, rubidium and strontium. Although different sites are involved for various cations, two sites are shown to be involved in uptake of each of these cations. Competitive inhibition is demonstrated at one of the two sites of ion uptake found for each cation. This is the site that dominates ion uptake at lower ion concentrations. Although hydrogen ion effects are also demonstrated at higher ion concentrations where a second site dominates cation uptake the experiments were not designed to show whether this effect is competitive.

#### METHODS

**THEORY:** Where  $k_3$  in equation 2 is rate limiting in the forward direction and  $k_4$  is negligible, the ob-

served absorption of an ion is proportional to MR as shown in equation 3.

$$v = k_3 [\text{MR}] t \quad (3)$$

where  $v$  = the amount of ion, M, absorbed in time,  $t$ .

From these equations a velocity equation can be derived analogous to case III for enzyme action described by Hofstee (16) and utilized by Hagen, Leggett and Jackson (13) for phosphorus uptake.

$$v = -K_m \frac{v}{[\text{M}]} + V_{\text{max}} \quad (4)$$

where  $K_m$  equals  $\frac{k_2 + k_3}{k_1}$  analogous to the Michaelis constant (20), and  $V_{\text{max}}$  equals the amount absorbed when all of the carrier is saturated with the ion under study, i.e.,  $\text{MR} = \Sigma\text{R}$ .

If  $v$  is plotted against  $\frac{v}{[\text{M}]}$  a straight line should result for each site provided  $[\text{M}]$  remains essentially constant during the steady state uptake. The slope of this straight line is  $-K_m$ , the apparent dissociation constant of MR, when  $k_2 \gg k_3$  (16). The y intercept is  $V_{\text{max}}$  and  $V_{\text{max}} = k_3 [\Sigma\text{R}]t$ .

If another ion competes for the same site, the  $V_{\text{max}}$  is the same in the presence of the competing ion (12) but the slope of the line changes. This change in slope reflects a decrease in the amount of ion at a given concentration associated with the site in the presence of a competing ion. Although the presence of one ion may affect the rate of uptake of another ion without being competitive, such an effect always results in a changed  $V_{\text{max}}$  and may also result in a change in slope.

It is more convenient to determine the total ion sorbed (absorbed plus adsorbed) instead of the total absorbed. Since adsorption not involved in uptake is negligible for monovalent cations as shown by Fried, Noggle and Hagen (10) and for phosphate as shown by Hagen and Hopkins (12) this would change equation 3 to 3a below:

$$v = [\text{MR}]k_3t + [\text{MR}] \quad (3a)$$

However it would not affect equation 4 except that

$$V_{\text{max}} = [\Sigma\text{R}]k_3t + [\Sigma\text{R}] \quad \text{instead of} \quad V_{\text{max}} = [\Sigma\text{R}]k_3t \quad (3b)$$

Thus, whether amount of ion absorbed or total sorbed is measured a plot of  $v$  vs  $\frac{v}{[\text{M}]}$  will give a slope,  $-k_m$ , the apparent dissociation constant.

**TECHNIQUE:** The methods employed are similar to those described by Epstein and Hagen (7). Seeds of *Hordeum vulgare*, var. Atlas 46, were germinated in demineralized water and grown for 6 days in continuously aerated  $2 \times 10^{-4}$  M  $\text{CaSO}_4$ . Twelve hours before an experiment was to begin, the roots were rinsed and demineralized water was substituted for the  $\text{CaSO}_4$  solution. The roots were excised, rinsed and suspended in two liters of continuously aerated,

demineralized water just prior to making the sorption studies.

Chloride salts of Rb, Na, K and Sr were labeled with  $\text{Rb}^{86}$ ,  $\text{Na}^{24}$ ,  $\text{Na}^{22}$ ,  $\text{K}^{42}$ , and  $\text{Sr}^{89}$  respectively. The pH of Rb and K solutions were adjusted to the desired value with 0.1 N NaOH or 0.1 N HCl. The pH of Na solution was adjusted to desired value with 0.1 N KOH or 0.1 N HCl.

The excised roots were blotted on washed, dry cheesecloth. One half gram portions of roots were weighed for Rb, K, or Na studies and 1.5 g portions for Sr studies and placed in beakers of continuously aerated water. The water was decanted and replaced by the appropriate radioactive solution to begin the sorption period. The cation concentration was maintained essentially constant by proper selection of root : volume ratios. At the end of the sorption period for Rb, Na or K, the solution was decanted and the roots were rinsed four times with demineralized water to remove adhering solutions and "outer space" (5) ions. The ion still associated with the root was considered total ion sorbed. At the end of the sorption period for Sr the solution was decanted, roots were rinsed with demineralized water, and adhering ions desorbed for a period of 30 minutes in continuously aerated solution of nonradioactive  $10^{-2}$  M  $\text{CaCl}_2$ . The Sr still associated with the root was considered total Sr absorbed. After rinsing, the roots were placed into circular,  $\frac{1}{4}$ -ounce seamless tin boxes and dried under an infrared lamp prior to measurement of activity. Aliquots of the radioactive solutions were pipetted into tin boxes, dried, measured and used as a standard for computing ion absorption by roots.

### EXPERIMENTAL RESULTS

The uptake of cation in a given period of time was plotted against the uptake divided by the concentration of cation in solution. The times chosen were in the steady state region as shown by Fried, Noggle and Hagen (10). This was 20 minutes for rubidium and potassium, 30 minutes for sodium and one hour for strontium.

The uptake of rubidium at various concentrations from  $10^{-6}$  to  $10^{-2}$  M was measured in the presence and absence of  $10^{-5}$  and  $10^{-4}$  M potassium. The results are presented in figure 1.

One site, assuming the carrier concept of ion uptake, should give a straight line relationship. However a curve was obtained. Since the curved line can be resolved into at least two linear components a and b, by the use of polar coordinates more than one site must be involved in rubidium uptake. Because the relationship appeared to approach linearity at both extremes of the curve, the linear components can also be approximated by extrapolation to the intercepts at both coordinates (fig 1). This approximation is feasible because the apparent  $K_m$  for each component differs widely (12). Therefore, a likely explanation for the experimentally observed phenomenon was




FIG. 1. The effect of potassium concentration on the uptake of rubidium by 0.5 g (fresh wt) of excised barley roots in 20 minutes. The uptake from solutions of various RbCl concentrations is plotted as a function of uptake divided by Rb concentration. The b-component of the reactions is separated at all potassium concentrations and the a-component at the zero-k level.

that two different sites were operative for rubidium absorption, one dominant at lower concentrations of the ion and the other at higher concentrations.

The uptake of rubidium in the presence of different concentrations of potassium also gives curved lines when plotted in the same manner. Again these curves can be separated into two linear components. The linear component at lower concentration of rubidium are illustrated by lines b' and b''. These linear components have the same  $V_{max}$  as the linear component, b, in the absence of potassium. This indicates that the decrease in the uptake of rubidium represents direct competition by potassium. Apparently potassium combines with the same carrier as rubidium.

The uptake of sodium, potassium, rubidium and strontium at various concentrations was measured in presence of  $10^{-4}$ ,  $10^{-5}$ ,  $10^{-6}$  and  $10^{-7}$  M hydrogen ion. Hydrogen ion concentration was maintained constant during the sorption period. The uptake of ion was then plotted against the uptake divided by concentration. For the sake of simplicity only the results at pH 4 and 6 are shown. At these pH levels no adjustment of pH was necessary during the experiment. The results at pH 5 were intermediate and those at pH 7 did not differ appreciably from pH 6 except for sodium which continued to show a hydrogen ion effect at the higher pH levels. For rubidium, potassium and sodium the total ion sorbed was measured. For strontium only the total ion absorbed was measured. Results are presented in figures 2 a, b, c, and d.

Again curved lines were obtained at all pH levels. The two linear components were again approximated by extrapolation to the intercept of both coordinates.




FIG. 2 a-d. The effect of pH on the uptake of cations by excised barley roots. The uptake from various concentrations of the cation is plotted as a function of uptake divided by concentration. The component reactions are separated into lines a and b for pH 6 and a' and b' for pH 4. Fig. a. Moles of Rb sorbed ( $\times 10^7$ ) by 0.5 g (fresh wt) of roots in 20 minutes. Fig. b. Moles of K sorbed ( $\times 10^7$ ) by 0.5 g (fresh wt) of roots in 20 minutes. Fig. c. Moles of Sr absorbed ( $\times 10^7$ ) by 1.5 g (fresh wt) of roots in 60 minutes. Fig. d. Moles of Na sorbed ( $\times 10^7$ ) by 0.5 g (fresh wt) of roots in 30 minutes.

Apparently the uptake of the four cations studied takes place from at least two distinct sites for each of the cations.

The results recorded in figure 2 also show that hydrogen ion is strictly competitive with each of the cations for b, the site that dominates ion uptake at low concentrations of the cation and probably competitive for the other site, a, that dominates ion uptake at the higher cation concentrations. The strict competition of hydrogen ion is evident because the  $V_{\max}$  (y intercept) is the same at both pH levels and the slope of the component lines differs at the two pH levels.

#### DISCUSSION

The uptake of each cation, apparently involves at least two distinct sites. This is somewhat analogous to the finding that two sites are involved in phosphorus uptake by excised barley roots as described by Hagen and Hopkins (12). However, the two sites for phosphorus were shown to be related to two separate ion species,  $\text{HPO}_4^{2-}$  and  $\text{H}_2\text{PO}_4^-$ . No such ex-

planation is apparent for cations, particularly mono-valent cations.

At least two sites of ion uptake were demonstrated for strontium, rubidium, potassium and sodium. Other cations may also have multiple sites. The potassium sites may be identical to the rubidium sites as suggested by figure 1 and the work of Epstein and Hagen (7). The calcium sites may also be identical to the strontium sites as indicated by both Epstein and Leggett (8) and Menzel and Heald (19) at relatively high ion concentrations.

The slope of line a and b equal the respective  $K_m$  values, the apparent dissociation constant of MR (the active intermediate as discussed under "METHODS"). These  $K_m$  values were calculated for Rb, K and Sr. The results recorded in table I under the heading  $K_m$  are a mean of several experiments. The results indicate that potassium, rubidium, and strontium complexes have approximately the same apparent dissociation constants.

To calculate the dissociation constant,  $K_1$ , of the complex formed with the competing ion, requires a

TABLE I  
APPARENT DISSOCIATION CONSTANTS OF Na, K, Rb, AND Sr AND THE APPARENT DISSOCIATION CONSTANTS OF COMPETING IONS

CATION SORBED	SITE DESIGNATION	APPARENT DISSOCIATION CONSTANTS		
		$K_m$	$K_1$ (HYDROGEN ION)	$K_1$ (POTASSIUM)
Na	a	$4 \times 10^{-8}$		
Na	b	$1 \times 10^{-5}$	$2 \times 10^{-6}$	
K	a	$1 \times 10^{-8}$		
K	b	$3 \times 10^{-6}$	$1 \times 10^{-5}$	
Rb	a	$2 \times 10^{-8}$		
Rb	b	$4 \times 10^{-6}$	$1 \times 10^{-5}$	$4 \times 10^{-6}$
Sr	a	$3 \times 10^{-8}$		
Sr	b	$3 \times 10^{-6}$	$1 \times 10^{-5}$	

generalized relationship for uptake in the presence and absence of the competing ion.

The equation for a competing ion used by Hagen and Hopkins (12) and Epstein and Hagen (7) when operating in the same system for the ion under test is formally analogous to equation 1 and is shown below as equation 5



where R in equation 1 and 5 is the same carrier, I is the competing ion and IR a complex of the competing ion and the carrier. By considering the equations for the equilibrium constants of the two reactions and the conservation equation

$$[\Sigma R] = [R] + [IR] + [MR] \tag{6}$$

a mass action equation can be derived relating the concentration of the ions and the apparent dissociation constants to the amount of active intermediates.

$$[MR] = [\Sigma R] - \left( K_m + \frac{[K_m][I]}{K_1} \right) \frac{[MR]}{[M]} \tag{7}$$

By substituting for MR and  $\Sigma R$  the respective relationships obtained from kinetic equations of steady state, 3a and 3b, equation 8 is obtained:

$$v = V_{max} - \left( K_m + \frac{[K_m][I]}{K_1} \right) \frac{v}{[M]} \tag{8}$$

where a plot of  $v$  vs  $v/[M]$  has a slope =

$$-\left( K_m + \frac{[K_m][I]}{K_1} \right).$$

Thus the apparent dissociation constant for potassium at the site that dominates at lower rubidium concentrations can be calculated from slope of line b ( $K_m$ ), and either  $b'$  or  $b''$ ,  $-\left( K_m + \frac{[K_m][I]}{K_1} \right)$  in figure 1. The apparent dissociation constants for hydrogen ion at the sites dominant at lower ion concentrations of rubidium; potassium and strontium sites can also be calculated from the appropriate lines b, pH 6 and  $b'$ , pH 4 in figure 2. Each run gave a

slightly different value. The averages of several determinations are presented in table I.

The apparent dissociation constant,  $K_m$ , of potassium as determined by the uptake of potassium was approximately the same as the apparent dissociation constant of potassium,  $K_1$ , from the rubidium site. Again it appears likely that the rubidium site b and potassium site b are one and the same site.

The apparent dissociation constant of the hydrogen ion complex,  $K_1$ , at site b is approximately the same for the site complexes of the three cations K, Rb, and Sr. This is higher than the dissociation constants for the cations themselves but only by a factor of approximately three. The magnitudes of the dissociation constants indicate that both hydrogen ion and the cations are relatively tightly held.

The  $K_1$  (hydrogen ion) for sodium could not be calculated from data in figure 2 d. A  $10^{-4}$  molar solution of hydrogen ion almost eliminated the appearance of site b, figure 2 d, in the sodium uptake system, indicating that hydrogen ion was more tightly held than sodium at this site. The same hydrogen ion concentration also had a marked effect on site a. Since hydrogen ion had a greater affinity for the sodium site than did sodium, pH 6 was not adequate to eliminate the effect of hydrogen ion at lower concentrations of sodium.

The  $K_m$  for a cation must be calculated in a system in which the concentration of competing ion is negligible. The  $K_m$  for sodium was therefore calculated from uptake data at pH 7. The  $K_1$  (hydrogen ion) was calculated from comparative uptake data at pH 5.0. The results are recorded in table I.

The  $K_m$  for sodium at the site that dominates at lower concentrations appears to be somewhat greater than that for the other cations tested. This indicates a slightly lower affinity of sodium for the sodium site than of the other cations for their respective sites. The  $K_m$  of site a is similar to that found for the other cations. The  $K_1$  (hydrogen ion) is smaller for sodium than the  $K_m$  of sodium indicating again that hydrogen is more tightly held than sodium. Hydrogen ion also appears to be more tightly held to the sodium sites than to the other cation sites.

The hydrogen ion competes with rubidium, potassium, sodium and strontium, and possibly most other cations. However, the dissociation constants for hydrogen ion on the b site are of same order of magnitude as the dissociation constants for the cation. Thus, when cation concentration is appreciably higher than hydrogen ion concentration there is little, if any, pH effect due to competition. However, if the concentrations are similar, marked pH effects may be noted.

### SUMMARY

The uptake of rubidium, potassium, sodium and strontium by excised barley roots under steady state conditions indicates at least two distinct carrier sites of absorption for each of the cations. One site domi-

nates at higher ion concentrations while the second dominates at lower ion concentrations.

The apparent dissociation constants were determined at each of the two sites for each of the four cations studied. They were similar in magnitude for potassium, rubidium and strontium and slightly higher for sodium.

Hydrogen ion was shown to compete directly for each of the sites that dominates at lower concentrations of all cations tested. The apparent dissociation constant for hydrogen ion at the strontium, rubidium and potassium site that dominates at lower concentrations was  $1 \times 10^{-5}$ . The comparable dissociation constant,  $K_1$ , at the sodium site was  $2 \times 10^{-6}$ . The apparent dissociation constants for hydrogen were similar in magnitude to the apparent dissociation constants of the ions themselves. Rubidium and potassium were shown to compete for the same site.

#### LITERATURE CITED

1. ARNON, D. I., FRATZKE, W. E. and JOHNSON, C. M. Hydrogen ion concentration in relation to absorption of inorganic nutrients by higher plants. *Plant Physiol.* 17: 515-525. 1942.
2. ARNON, D. I. and JOHNSON, C. M. Influence of hydrogen ion concentration on the growth of higher plants under controlled conditions. *Plant Physiol.* 17: 525-539. 1942.
3. BROYER, T. C. The nature of the process of inorganic solute accumulation in roots. In: *Mineral Nutrition of Plants*, E. Truog, ed. Pp. 187-250. Univ. of Wisconsin Press, Madison 1951.
4. COLLANDER, R. Selective absorption of cations by higher plants. *Plant Physiol.* 16: 691-720. 1942.
5. EPSTEIN, E. Passive permeation and active transport of ions in plant roots. *Plant Physiol.* 30: 529-535. 1955.
6. EPSTEIN, E. Mineral nutrition of plants: Mechanism of uptake and transport. *Ann. Rev. Plant Physiol.* 7: 1-24. 1956.
7. EPSTEIN, E. and HAGEN, C. E. A kinetic study of the absorption of alkali cations by barley roots. *Plant Physiol.* 27: 457-474. 1951.
8. EPSTEIN, E. and LEGGETT, J. E. The absorption of alkaline earth cations by barley roots: Kinetics and mechanism. *Amer. Jour. Bot.* 4: 785-791. 1954.
9. FAWZY, H., OVERSTREET, R. and JACOBSON, L. The influence of hydrogen ion concentration on cation absorption by barley roots. *Plant Physiol.* 29: 234-237. 1954.
10. FRIED, M., NOGGLE, J. C. and HAGEN, C. E. The relationship between adsorption and absorption of cations. *Soil Sci. Soc. Amer. Proc.* (In press.)
11. GUEST, P. L. and CHAPMAN, H. D. Some effects of pH on growth of citrus in sand and solution cultures. *Soil Sci.* 58: 455-465. 1944.
12. HAGEN, C. E. and HOPKINS, H. T. Ionic species in orthophosphate absorption by barley roots. *Plant Physiol.* 30: 193-199. 1955.
13. HAGEN, C. E., LEGGETT, J. E. and JACKSON, P. C. The sites of orthophosphate uptake by barley roots. *Proc. Nat. Acad. Sci. U.S.* 43: 496-506. 1957.
14. HESLEP, J. M. A study of the infertility of two acid soils. *Soil Sci.* 72: 67-80. 1951.
15. HOAGLAND, D. R. and BROYER, T. C. Hydrogen-ion effects and the accumulation of salt by barley roots as influenced by metabolism. *Amer. Jour. Bot.* 27: 173-185. 1940.
16. HOFSTEE, B. H. J. On the evaluation of the constants  $V_m$  and  $K_m$  in enzyme reactions. *Science* 116: 329-331. 1952.
17. JACOBSON, L., OVERSTREET, R., KING, H. M. and HANDLEY, R. A study of potassium absorption by barley roots. *Plant Physiol.* 25: 639-647. 1950.
18. LEGGETT, J. E. and EPSTEIN, E. Kinetics of sulfate absorption by barley roots. *Plant Physiol.* 31: 222-226. 1956.
19. MENZEL, R. G. and HEALD, W. R. Distribution of potassium, rubidium, cesium, calcium and strontium within plants grown in nutrient solutions. *Soil Sci.* 80: 287-293. 1955.
20. MICHAELIS, L. and MENTEN, M. L. Die Kinetick der Invertinwirkung. *Biochem. Zeits.* 49: 333-369. 1913.
21. NIELSEN, T. R. and OVERSTREET, R. A study of the role of the hydrogen ion in the mechanism of potassium absorption by excised barley roots. *Plant Physiol.* 30: 303-309. 1955.
22. ROSENBERG, T. On accumulation and active transport in biological systems. *Acta Chem. Scand.* 2: 14-33. 1948.

#### GROWTH SUBSTANCES IN CORN POLLEN<sup>1,2,3</sup>

H. N. FUKUI, F. G. TEUBNER, S. H. WITTEWITZ AND H. M. SELL

DEPARTMENTS OF AGRICULTURAL CHEMISTRY AND HORTICULTURE,  
MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN

Growth substances have been detected in the pollen of apple (3), orchid (7), tobacco (5), hazelnut shrub and pine (14), and corn (6, 10, 13). Studies of the growth substances in corn pollen have been con-

cerned solely with crude ether soluble extracts. This communication describes the growth substances found in the acid and non-acidic or "neutral" fractions in ethyl ether extracts of corn pollen, *Zea mays*, var. Golden Cross.

Freshly harvested pollen (166 grams) was extracted with cold, peroxide-free ethyl ether at 2° C for four hours. The supernatant was decanted and extracted with 5% aqueous sodium bicarbonate. The

<sup>1</sup> Received revised manuscript November 20, 1957.

<sup>2</sup> This investigation was supported in part by a grant from the National Science Foundation.

<sup>3</sup> Journal Article No. 2107 of the Michigan Agricultural Experiment Station.