

# Musculoskeletal Injuries and United States Army Readiness. Part II: Management Challenges and Risk Mitigation Initiatives

COL (ret) Joseph M. Molloy, PT PhD\*† ; LTC (ret) Timothy L. Pendergrass, PT DSc ATC† ; LTC Ian E. Lee, PT DSc‡ ; LTC (ret) Keith G. Hauret, MSPH MPT§ ; Michelle C. Chervak, PhD MPH§ ; MAJ (ret) Daniel I. Rhon, PT DSc\*†¶¶

---

## ABSTRACT

### Introduction

Noncombat injuries (“injuries”) threaten soldier health and United States (U.S.) Army medical readiness, accounting for more than twice as many outpatient medical encounters among active component (AC) soldiers as behavioral health conditions (the second leading cause of outpatient visits). Noncombat musculoskeletal injuries (MSKIs) account for more than 80% of soldiers’ injuries and 65% of medically nondeployable AC soldiers. This review focuses on MSKI risk reduction initiatives, management, and reporting challenges within the Army. The authors will summarize MSKI risk reduction efforts and challenges affecting MSKI management and reporting within the U.S. Army.

### Materials/Methods

This review focuses on (1) initiatives to reduce the impact of MSKIs and risk for chronic injury/pain or long-term disability and (2) MSKI reporting challenges. This review excludes combat or battle injuries.

### Results

#### Primary risk reduction

Adherence to standardized exercise programming has reduced injury risk among trainees. Preaccession physical fitness screening may identify individuals at risk for injury or attrition during initial entry training. Forward-based strength and conditioning coaching (provided in the unit footprint) and nutritional supplementation initiatives are promising, but results are currently inconclusive concerning injury risk reduction.

#### Secondary risk reduction

Forward-based access to MSKI care provided by embedded athletic trainers and physical therapists within military units or primary care clinics holds promise for reducing MSKI-related limited duty days and nondeployability among AC soldiers. Early point-of-care screening for psychosocial risk factors affecting responsiveness to MSKI intervention may reduce risk for progression to chronic pain or long-term disability.

#### Tertiary risk reduction

Operational MSKI metrics enable commanders and clinicians to readily identify soldiers with nonresolving MSKIs. Monthly injury reports to Army leadership increase command focus on soldiers with nonresolving MSKIs.

### Conclusions

Standardized exercise programming has reduced trainee MSKI rates. Secondary risk reduction initiatives show promise for reducing MSKI-related duty limitations and nondeployability among AC soldiers; timely identification/evaluation and appropriate, early management of MSKIs are essential. Tertiary risk reduction initiatives show promise for identifying soldiers whose chronic musculoskeletal conditions may render them unfit for continued military service. Clinicians must document MSKI care with sufficient specificity (including diagnosis and external cause coding) to enable large-scale systematic MSKI surveillance and analysis informing focused MSKI risk reduction efforts. Historical changes in surveillance methods and injury definitions make it difficult to compare injury rates and trends over time. However, the U.S. Army’s standardized injury taxonomy will enable consistent classification of current and future injuries by mechanism of energy transfer and diagnosis. The Army’s electronic physical profiling system further enables

---

\*Oak Ridge Institute for Science and Education, Oak Ridge, TN 37830

†Physical Performance Service Line, G 3/5/7, U.S. Army Office of the Surgeon General, Falls Church, VA 22042

‡Solution Delivery Division, U.S. Defense Health Agency, Falls Church, VA 22042

§U.S. Army Public Health Center, Injury Prevention Program, Aberdeen Proving Ground, Aberdeen, MD 21005

¶Center for the Intrepid, Brooke Army Medical Center, Joint Base San Antonio Fort Sam Houston, San Antonio, TX 78234

¶¶Duke Clinical Research Institute, Duke University, Durham, NC 27701  
Invited platform presentation during the 2019 Military Health System Research Symposium conference (Management & Treatment of Warfighter

---

Neuromusculoskeletal Injuries: Musculoskeletal Injury Rates Across the U.S. Army: A Review of the Data and Risk Mitigation Initiatives).

The views expressed in this article are those of the authors and do not necessarily reflect the official policy of the Department of Defense, Department of the Army, U.S. Army Medical Department or the U.S. Government.  
doi:10.1093/milmed/usaa028

Published by Oxford University Press on behalf of the Association of Military Surgeons of the United States 2020. This work is written by US Government employees and is in the public domain in the US.

standardized documentation of MSKI-related duty/work restrictions and mechanisms of injury. These evolving surveillance tools ideally ensure continual advancement of military injury surveillance and serve as models for other military and civilian health care organizations.

**OVERVIEW**

Noncombat injuries (“injuries”) threaten soldier health and U.S. Army medical readiness, accounting for more than twice as many outpatient medical encounters among active component (AC) soldiers as behavioral health conditions (the second leading cause of outpatient visits) or 2.2 versus 1 million encounters in 2017.<sup>1</sup> Noncombat musculoskeletal injuries (MSKIs) account for more than 80% of soldiers’ injuries<sup>2,3</sup> and 65% of medically nondeployable AC soldiers.<sup>4</sup> MSKIs accounted for 59% of soldiers’ limited duty days during the first 6 months of 2019;<sup>5</sup> pregnancy/postpartum and behavioral health (the second and third leading conditions) each accounted for only 10% of limited duty days.<sup>5</sup>

The U.S. Army incorporates primary, secondary, and tertiary risk reduction initiatives to mitigate the impact of MSKIs. MSKIs are defined as musculoskeletal disorders resulting from mechanical energy transfer,<sup>2</sup> including traumatic and overuse injuries, which may cause pain and/or limit function. Mitigation efforts focus on preaccession populations, trainees, and healthy but injured soldiers in operational units. [Supplementary Table S1](#) addresses pertinent MSKI risk reduction initiatives.

Successful risk reduction strategies require understanding of the relative incidence of overuse versus traumatic MSKI, common injury sites, leading causes of injury, and long-term trends in MSKI rates. Overuse MSKIs (comprising at least 70% of AC soldiers’ injuries)<sup>2,3</sup> are a more relevant target for risk reduction strategies than traumatic injuries, given their (1) higher incidence across the Army and (2) greater potential for mitigation by physical training-related adaptations.

**PURPOSE**

To summarize MSKI risk reduction efforts and challenges affecting MSKI management and reporting within the U.S. Army.

**METHODS**

This review focuses on (1) initiatives to reduce the impact of MSKIs and risk for chronic injury/pain or long-term disability and (2) MSKI reporting challenges. This review excludes combat or battle injuries.

**PRIMARY RISK REDUCTION**

Primary risk reduction initiatives focus on decreasing risk for MSKI occurrence.<sup>6</sup> Standardized exercise programming for trainees has proven effective; other initiatives (eg, preaccession physical fitness screening and nutritional supplementation during basic training) show promise for reducing injury rates.

**TABLE I.** Occupational Physical Assessment Test (OPAT) Overview

<p><b>Measures</b></p> <ul style="list-style-type: none"> <li>Upper body power (seated power throw)</li> <li>Lower body power (standing long jump)</li> <li>Total body strength (deadlift)</li> <li>Aerobic capacity (interval aerobic run or beep test)</li> </ul> <p><b>Standards</b></p> <ul style="list-style-type: none"> <li>Sex/age-neutral</li> </ul> <p><b>Physical demand category qualifications</b></p> <ul style="list-style-type: none"> <li>Black (specialties with heavy physical demands)</li> <li>Gray (specialties with significant physical demands)</li> <li>Gold (specialties with moderate physical demands)</li> <li>White (not ready for any specialty)</li> </ul>
---

**PREACCESSION PHYSICAL FITNESS SCREENING**

The U.S. Army Recruiting Command currently performs preaccession physical fitness screening through the Occupational Physical Assessment Test (OPAT). There are minimum OPAT standards for each military occupational specialty, based on that specialty’s physical demand category. Career soldiers seeking to reclassify to more physically demanding specialties must also meet OPAT standards for their desired new specialty before beginning training.<sup>7</sup> [Table I](#) highlights OPAT measures, standards, and physical demand category qualifications.

The OPAT serves to predict an individual’s capacity to perform the physically demanding tasks of their desired military occupational specialty.<sup>8</sup> Although not designed to predict or screen for injury risk, the OPAT may provide a secondary effect in reducing injuries and attrition by (1) establishing minimum fitness standards to enter the Army and (2) pairing individuals with an occupational specialty based on their OPAT performance and physical demand category of the occupational specialty.<sup>9,10</sup>

Preliminary reports indicate that male trainees with gray (significant) or gold (moderate) overall OPAT scores had a 1.2 times greater injury risk (and were 1.4 or 1.6 times more likely, respectively, to attrit during Basic Combat Training [BCT]) versus men meeting the highest standard-black (heavy)-for overall OPAT scores.<sup>10</sup> Female trainees with gold overall OPAT scores had a 1.1 times greater injury risk versus women with black overall OPAT scores.<sup>10</sup> OPAT scores among female trainees were not associated with attrition during BCT;<sup>10</sup> lack of an association might be because of the relatively small numbers of women in the study. Regardless of overall scores, injury risk was also greater among male and female trainees who failed to meet the black standard for each of the four individual events.<sup>10</sup>

The Army strives to incorporate risk reduction initiatives based on the best current and actionable evidence. Low aero-

Downloaded from https://academic.oup.com/milmed/article/185/9-10/e1472/5762817 by guest on 21 August 2022

bic capacity is the fitness component most strongly and consistently associated with higher injury risk among trainees.<sup>11</sup> Low muscular endurance is also associated with increased risk, but not as strongly or consistently as low aerobic capacity.<sup>11</sup> Relatively few reports have evaluated the association of muscular strength with injury risk in military populations.<sup>11</sup> The Army's increasing emphasis on training and testing muscular strength and power (eg, OPAT, Army Combat Fitness Test [ACFT]) will enable assessment of potential associations of muscular strength and power with injury risk.

### STANDARDIZED EXERCISE PROGRAMMING

U.S. Army trainees have performed standardized exercise programming, known as Army Physical Readiness Training (PRT) since 2004. The intent of PRT is to provide a standardized, safe, and effective approach to fitness training. PRT incorporates at least two strength/mobility sessions and two endurance/mobility sessions weekly. This ideally ensures equal emphasis on strength, endurance, and mobility development while providing sufficient recovery between training sessions. PRT's exercise drills incorporate greater exercise variety than traditional Army physical training programs for optimal physical fitness development.<sup>12</sup> PRT's emphasis on interval runs, shuttle runs, and hill repeat runs enables aerobic and anaerobic fitness development with less running distance and frequency compared to traditional Army fitness programs that emphasized steady state runs.<sup>12</sup> PRT ideally ensures a gradual progression in training volume while providing standardized (1) strength, endurance, and mobility activities and (2) standardized preparation and recovery exercises.<sup>12</sup>

PRT is recognized as a major contributor to the decline in trainee injury rates over the past decade.<sup>13</sup> Three field studies evaluating PRT found that it reduced injury risk among trainees by 33 to 45% (while trainees maintained or improved overall fitness) when compared to traditional physical training programs.<sup>14,15</sup> PRT's effectiveness in reducing trainee injury risk is likely due in part to consistent command emphasis on and enforcement of correct PRT performance across BCT and One Station Unit Training (OSUT).

### NUTRITIONAL SUPPLEMENTATION

Nutritional deficiencies are potential risk factors for MSKI, and can negatively impact postinjury recovery.<sup>16</sup> The combined effects of increased nutritional energy demands and physical requirements may further increase injury risk during BCT or OSUT.

The Army is evaluating the impact of providing a calcium and vitamin D-enhanced Performance Readiness Bar daily to trainees. This bone health initiative is based on reports that calcium and vitamin D supplementation reduced stress fracture incidence by 20% among female U.S. Navy recruits<sup>17</sup> and improved tibial cortical bone mineral density, content, and thickness among male and female trainees.<sup>18</sup> It is

currently unknown whether Performance Readiness Bar supplementation has affected Army trainees' stress fracture risk; the U.S. Army Research Institute of Environmental Medicine has an ongoing longitudinal study addressing modifiable and nonmodifiable factors (including self-reported Performance Readiness Bar consumption) that may affect trainees' MSKI risk.<sup>19</sup>

The Army is also evaluating the effects of voluntary multivitamin with iron (MVI) supplementation on injury risk among female trainees. Iron deficiency prevalence has been reported as nearly 20% among women beginning Basic Combat Training (BCT); the prevalence may exceed 50% upon completing BCT.<sup>20</sup> Iron status decrements across BCT have been associated with decreased 2-mile run performance.<sup>21</sup> Overall injury incidence among female BCT trainees decreased by 3% across the first year of voluntary MVI supplementation when compared with the prior year.<sup>10</sup> However, MVI intake, trainee age, body composition, and physical fitness were not controlled across the year-long observation period. It is currently unclear whether voluntary MVI supplementation has affected female trainees' injury incidence.

### COMPREHENSIVE MSKI PREDICTION MODELING

MSKI risk and prognosis for recovery depend on multiple factors (eg, physical, psychosocial). Comprehensive injury prediction models account for multiple risk factors (eg, smoking status, lower aerobic capacity or muscular endurance, pain during Functional Movement Screen clearing tests),<sup>22,23</sup> unlike risk prediction efforts based on a single functional movement or physical performance test.<sup>24,25</sup> A comprehensive model's predictive accuracy is based on the number of risk factors/predictors present; probability of injury increases as more predictors are present. Comprehensive predictive modeling tools are potentially effective, time and resource-efficient, low-cost injury risk screens that direct soldier-specific risk reduction interventions. [Supplementary Table S2](#) highlights pertinent injury predictive model reports.

### HOLISTIC HEALTH AND FITNESS SYSTEM (H2F)

H2F is a "system of training and testing that optimizes and validates performance readiness across a Soldier's career."<sup>26</sup> As described by the Commanding General, U.S. Army Center for Initial Military Training, H2F includes "mental, spiritual, nutrition and sleep readiness taught in military education and in the unit by unit-owned performance experts" and incorporates "much broader physical training options and modalities."<sup>26</sup> Prospective data are needed to validate H2F's effectiveness for reducing MSKI risk and improving military readiness.

H2F's emerging physical training and testing doctrine includes a transition to the Army Combat Fitness Test (ACFT). The ACFT is a six-event test intended to (1) demonstrate a soldier's 'strength, power, agility, coordination, balance,

anaerobic capacity, and aerobic capacity<sup>27</sup> and (2) “better predict a soldier’s readiness for the demands of the modern battlefield.”<sup>26</sup>

## RUNNING FOOTWEAR

The Army transitioned from wearing combat boots to running shoes during physical training in 1982 to reduce risk for running-related injuries.<sup>28</sup> However, a recent historical comparison found similar overall and lower body injury incidences among trainees wearing boots versus running shoes during physical training in BCT.<sup>28</sup>

Army trainees in BCT and OSUT have been directed to running shoe types (eg, cushioned, stability, motion control) based on their static plantar foot shape for many years. However, running shoe prescription based on static foot shape is not proven to reduce trainees’ or recruits’ injury risk.<sup>15,29</sup> Based on the evidence to date, soldiers’ footwear selection should emphasize comfortable fit, proper sizing (length and width), and timely replacement (before obvious midsole wear/breakdown or decreased comfort occur).<sup>15,29</sup>

## SECONDARY RISK REDUCTION

Secondary risk reduction initiatives focus on decreasing the impact of existing MSKIs.<sup>6</sup> Reducing MSKI-related physical profiling (limited duty days), nondeployability, and risk for chronic pain or long-term disability postinjury likely improves readiness and reduces injury-related financial costs. A small percentage of soldiers is responsible for a disproportionately large number of limited duty days and MSKI costs. Four percent of AC soldiers were on chronic MSKI profiles (exceeding 90 days across the previous six months) in 2016; they accounted for 51% of all MSKI-related limited duty days (approximately 5 million days annually) that year.<sup>30</sup>

## FORWARD-BASED/EMBEDDED MSKI CARE WITHIN THE OPERATIONAL UNIT FOOTPRINT

Soldiers with unresolved injuries are an ideal focus for timely, forward-based MSKI care; prior injury is one of the strongest predictors of future (recurrent and new-site) injuries.<sup>15,22,31–37</sup> The effect of prior injury is likely due partly to incomplete healing or rehabilitation, failure to return to full physical function before returning to full military duties, unresolved biomechanical or structural issues, postinjury deficits in proprioception or strength, postinjury compensatory gait patterns and/or altered pain perception.<sup>15,31,33,34,36–40</sup> [Supplementary Table S3](#) highlights pertinent MSKI risk factor reports.

Forward-based/embedded MSKI care initiatives emphasizing timely identification of injuries and expert management of care and rehabilitation can potentially expedite return to duty while reducing risk of chronic pain or long-term disability postinjury. The Army is bolstering early and appropriate access to MSKI care within the operational unit footprint. Expanding access by embedding additional MSKI care clinicians and fitness experts in operational units was a key com-

ponent of the U.S. Army Forces Command’s (FORSCOM) multi-interventional Soldier Readiness Test and Training Program (SRT2P). This program will continue its forward-based care emphasis as it transitions into the H2F. Long-term outcome data are needed to validate forward-based MSKI care’s role in decreasing the impact of existing MSKIs.

## H2F FIELD TEST

The SRT2P included forward medical care provided by multidisciplinary teams outside of the traditional military treatment facility (MTF). The SRT2P team, embedded in the operational units, also included strength and conditioning (SC) coaches and athletic trainers (ATs). SRT2P transitioned to the H2F field test, which will continue through September 2020. The H2F field test shares many capabilities with the Army’s Ranger Athlete Warrior (RAW) and U.S. Army Special Operations Command’s Tactical Human Optimization, Rapid Rehabilitation and Reconditioning (THOR3) programs.

The H2F field test currently assigns an AT, registered dietitian (RD), physical therapist (PT), and two SC coaches to each of 30 operational battalions (typically containing 400–1000 soldiers).<sup>41</sup> The H2F field test also assigns occupational therapists (OT) or cognitive enhancement specialists to multiple operational battalions. The H2F field test is a bridging program to the H2F system.

The H2F system is currently envisioned to provide multifunctional, forward-based/embedded health care and fitness teams of varying staff sizes to all brigade combat teams (typically containing 4,400–4,700 soldiers)<sup>41,42</sup> or other brigade equivalent-sized units (typically containing 3,000–5,000 soldiers).<sup>41</sup> Each brigade combat team currently has an embedded PT team (one PT and PT technician). Special operations units are currently the only units with embedded teams incorporating ATs, RDs, OTs, and SC coaches. More robust, forward-based/embedded health care and fitness teams will ideally (1) facilitate access to clinical and fitness resources for improved medical readiness and soldier performance and (2) reduce disruptions to time-sensitive unit training schedules.

H2F’s embedding of forward-based SC coaches in operational units is a promising primary and secondary risk reduction initiative. SC coaches can impact primary risk reduction through “safe exercise technique, appropriate programming and facility design.”<sup>43</sup> SC coaches’ collaboration with unit leadership and embedded health care providers can optimize (1) soldiers’ timely injury reporting (secondary risk reduction), (2) transitions from clinical-based to occupational task-specific injury rehabilitation (secondary risk reduction), and (3) balancing of physical training and occupational duty stressors to maximize potential for adequate postactivity recovery (primary risk reduction).<sup>43</sup>

## MSKI SUPPORT TO SUSTAINMENT BRIGADES

U.S. Army sustainment brigades provide logistical, transportation, maintenance, and personnel services support to

brigade combat teams and other units operating in their assigned support areas.<sup>44</sup> Pending full implementation of H2F System and assignment of multi-functional health care teams to sustainment brigades, the U.S. Army Medical Command (MEDCOM) is collaborating with sustainment brigade leadership across the Army to provide early, convenient MSKI care through morning MSKI sick call conducted by PTs at the nearest MTF or within the brigade living and training area.

Physical profiles address soldiers' physical abilities to perform required tasks, and document specific task/duty limitations based on each soldier's injury. PTs assigned to installation MTFs now participate regularly in sustainment brigade-led physical profile review boards (where unit leaders and clinicians collaborate to track the medical and functional status of soldiers on limited duty, provide active MSKI surveillance and share MSKI readiness metrics with sustainment brigade leadership). Collaboration between MEDCOM and sustainment brigade leadership stems from leading practices implemented in 2016 by the First Armored Division and William Beaumont Army Medical Center leadership at Fort Bliss, Texas.<sup>30,45</sup>

**FORWARD-BASED MSKI CARE WITHIN THE INITIAL ENTRY TRAINING FOOTPRINT**

MEDCOM and the U.S. Army Training and Doctrine Command (TRADOC) have collaborated since 2010 to embed ATs in BCT and OSUT battalions. The ATs evaluate and treat trainees' MSKIs during early morning sick call within the trainees' living and training areas. ATs refer high-risk, complex and nonresponding patients for consultation by PTs, primary care providers, or orthopedic specialists at the nearest MTF. Per TRADOC leadership's estimate, forward-based ATs can evaluate injured trainees, establish treatment plans, provide initial treatment, and return trainees to the training unit before the start of their training day in approximately 85% of cases.

**EMBEDDED PTS IN ARMY MEDICAL HOMES**

MEDCOM also embeds PTs in primary care clinics within Army Medical Homes (multidisciplinary health care delivery sites).<sup>46</sup> Although not technically forward-based care, medical homes are located closer to the soldiers' living and training areas. The Army Medical Homes provide closer collaboration between primary care clinicians and their embedded PTs, thus optimizing early, convenient access to expert MSKI care.<sup>46</sup>

**EARLY POINT-OF-CARE SCREENING FOR PSYCHOSOCIAL RISK FACTORS AFFECTING RESPONSIVENESS TO MSKI INTERVENTION**

The Military Orthopedic Tracking Injuries Outcomes Network Musculoskeletal Data Portal (MOTION-MDP) is a

**TABLE II.** Military Orthopedic Tracking Injuries Outcomes Network Musculoskeletal Data Portal (MOTION-MDP) Overview

<p><b>Capabilities</b></p> <ul style="list-style-type: none"> <li>Captures temporal changes in MSKI-related pain, function, and disability</li> <li>Enables point-of-care risk factor screening during initial patient encounters</li> <li>Facilitates clinical outcomes assessment after treatment for MSKIs and orthopedic surgical procedures</li> </ul> <p><b>Components</b></p> <ul style="list-style-type: none"> <li>National Institute of Health's Patient-Reported Outcomes Measurement Information System (PROMIS) outcome tool</li> <li>Validated screening and outcomes measures</li> <li>Surveys completed by patient and clinician at specified intervals across episodes of MSKI care</li> </ul> <p><b>Focus</b></p> <ul style="list-style-type: none"> <li>Resiliency, psychosocial risk factors, recovery perceptions, health care-seeking behaviors and therapeutic alliance measures that are relevant to MSKI-related outcomes</li> </ul>
---

Defense Health Agency program of record that collects patient-reported data within the Military Health System (MHS). MOTION-MDP is becoming a standard clinical practice for MSKI surgical and nonsurgical patient management. MOTION-MDP focuses on psychosocial and related risk factors that may identify patients who would benefit from a holistic interventional approach to MSKI care.<sup>47-56</sup> Table II highlights MOTION-MDP's capabilities, individual components, and overall focus.

Soldiers with injuries or complex conditions that are not responding to medical care and rehabilitation as expected may respond to holistic interventional approaches. Allocating appropriate holistic health care resources early in episodes of care and collaboratively intervening with comprehensive, integrated pain management team members (including nonmedical leadership) can minimize risk for chronic pain, long-term disability and excessive health care resource consumption.<sup>47-56</sup> Supplementary Table S4 highlights psychosocial risk factors affecting responsiveness to MSKI intervention.

**ARMY MEDICAL LEADERSHIP POLICY GUIDANCE ON BONE STRESS INJURIES (BSIS)**

MEDCOM has issued policy guidance concerning clinical and administrative management of soldiers sustaining BSIs. The objective is to optimize clinical and functional outcomes by minimizing variance in BSI identification and management. The policy guidance incorporates leading practices in clinical BSI management, including imaging procedures, standardized grading criteria for advanced imaging techniques, recommended notifications, early intervention, and diagnostic coding procedures. Outcome data are necessary to determine this policy's effectiveness in mitigating the impact of BSIs across the Army.

**TABLE III.** Operational MSKI Metrics for Corps, Division and Brigade-Sized Units<sup>30</sup>

Percentages of soldiers on MSKI-related limited duty in prior 180 days
1–30 days (short-term)
31–90 days (long-term)
Beyond 90 days (chronic)

### TERTIARY RISK REDUCTION

Tertiary risk reduction emphasizes mitigating long-term effects of chronic MSKIs on Army readiness and the individual soldier. Operational MSKI metrics enable unit commanders and clinicians to readily identify soldiers with nonresolving MSKIs. Monthly injury reports to the Army's leadership have increased command focus on soldiers with nonresolving MSKIs.

These soldiers should be considered for (1) reassignment to occupational specialties they can perform within the constraints of their MSKI-related duty limitations or (2) medical separation from the Army. Reassignment to more appropriate occupational specialties or medical separation may benefit soldiers by minimizing exposures to activities that aggravate their chronic conditions.

### OPERATIONAL MSKI METRICS

MEDCOM has developed MSKI-based measures of effectiveness targeting soldiers' limited duty days, as documented electronically in physical profiles. Clinicians write MSKI-associated profiles in the Army's electronic physical profiling (eProfile) system to limit soldiers' physical activities and duty requirements during postinjury recovery periods. eProfile enables standardized documentation of MSKI-related duty/work restrictions and mechanisms of injury.

Operational metrics communicate the extent of MSKI burden on military units, accounting for noncontinuous, repetitive short-term profiling in the past 180 days. [Table III](#) highlights operational MSKI metrics for corps, division, and brigade-sized units. Monthly metrics reported to Army leadership facilitate installation-level MSKI monitoring. The U.S. Army Public Health Center (APHC) publishes Army and installation injury metrics,<sup>30,57</sup> develops tools for monitoring high risk populations,<sup>58</sup> and coordinates with public health assets<sup>59</sup> to support MSKI risk reduction at the military installation level.<sup>60</sup>

Army leadership emphasis and the above initiatives are contributing to a reduced MSKI burden. AC soldiers had nearly 17% or 1.6 million fewer days on limited duty because of MSKI in 2018, despite the Army AC changing minimally in size when compared with 2017.<sup>61</sup>

### IMPROVING MILITARY MSKI REPORTING

Timely, accurate MSKI reporting and consistent/mandatory external cause coding are critical for (1) assessing the extent of the Army's MSKI problem, (2) identifying underlying causes of MSKIs, (3) identifying MSKI trends across units and time, and (4) determining the relative effectiveness of MSKI risk reduction interventions. Clinicians must document MSKI care with sufficient specificity (including diagnosis and external cause coding) to enable large-scale systematic MSKI surveillance and analysis informing focused MSKI risk reduction efforts.

Varying MSKI definitions, inadequate medical coding in the electronic health records and the October 2015 transition in coding systems have hindered MSKI reporting and surveillance efforts. Standardized injury definitions and consistent coding are essential for addressing the MSKI problem through effective reporting and surveillance.

The MHS transitioned to the International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) medical coding system in October 2015, making it difficult to compare injury rates with earlier reports.<sup>62,63</sup> The ICD-10 system provides a much greater selection of detailed injury diagnostic codes than prior coding systems.<sup>64</sup> Numbers of reported injuries may increase with ICD-10 coding, given (1) variance among clinicians when choosing MSKI diagnostic codes and (2) repeated documentation of the same injury by multiple clinicians across multiple clinical encounters.<sup>2</sup> However, the ICD-10 transition should not impact other injury incidence reports (eg, numbers of individuals sustaining at least one injury across a set time).

The ICD-9/10 coding system addresses traumatic injuries and injury-related musculoskeletal conditions (primarily overuse injuries) in separate diagnosis code series. Military injury surveillance efforts must account for all injuries (overuse and traumatic);<sup>2</sup> failing to account for overuse injuries would underestimate injury-related, outpatient encounters by more than 50%.<sup>65</sup>

The APHC developed a comprehensive injury taxonomy because of challenges with ICD-9/10's injury classification system.<sup>2</sup> The taxonomy advances injury surveillance by identifying and categorizing overuse and traumatic MSKI, differentiating between MSKI and nonmusculoskeletal injuries and differentiating among causative sources of injuries (eg, via mechanical energy or environmental exposures to thermal or radiant energy).<sup>2</sup>

Infrequent external cause coding in electronic health records for MSKI-related outpatient encounters limits analysis of underlying causes of MSKIs.<sup>66</sup> Consistent/mandatory external cause coding for MSKI-related encounters is critical for identifying and targeting underlying causes for MSKIs.

Differences in MSKI assessments can hinder comparisons among injury reports and assessments of injury trends over time. Variability in MSKI definitions related to (1) activity

**TABLE IV.** Survey Research Capabilities<sup>66–68</sup>

<p>Determination of cause or mechanism of injury</p> <p>Determination of injury severity, functional loss, and/or limited duty time</p> <p>Measurement of degree of pain</p> <p>Deployment injury data capture</p> <p>Note: Future analysis of electronic physical profile data will enhance understanding of severity of military injuries.</p> <p>Note: MOTION-MDP implementation should improve surveillance efforts to account for pain and functional loss associated with injuries.</p>
---

or work restrictions and (2) injuries requiring medical intervention versus self-treatment can lead to differing MSKI estimates.<sup>2</sup> Table IV highlights situations requiring MSKI survey research because of current medical surveillance limitations.

Standardized definitions are essential for injury risk reduction measures. Incorporating a severity measure (eg, amount of work loss, number of recurrences, long-term costs) provides greater clarity, but results must be clearly stated. Reported injury rates can vary considerably, based on the presence or absence of a time-loss severity measure (such as requiring at least one limited duty day per injury).<sup>67–69</sup>

Varied definitions exist for “training-related injury.” The APHC categorizes all overuse and traumatic injuries of the lower back, pelvis and lower extremities sustained by trainees as training-related injuries. This subset of MSKI includes the most common injuries sustained by trainees. In contrast, other researchers have included all injuries (MSKI and non-MSKI) in their definition of training-related injuries sustained during BCT.<sup>70</sup>

Comparisons among reports are also influenced by the varied populations under study. Reports have addressed the entire Army AC (BCT, OSUT and operational units combined), operational Army units, BCT and OSUT combined or BCT and OSUT separately. When comparing injury rates and trends among surveillance reports, one must consider each report’s injury definition, study population, surveillance methods and surveillance period length. One must also account for variations in Army populations, such as (1) female trainees comprising a greater proportion of the BCT versus the OSUT trainee population<sup>3</sup> and (2) shorter BCT versus OSUT duration (10 weeks versus 13–19 weeks).

The military environment poses additional unique challenges. Frequent moves among soldiers and clinicians impact continuity of care, which can increase long-term medical costs.<sup>71</sup> Soldiers may return to full military duty before they are fully ready because of (1) unit training or operational requirements or (2) insufficient criteria for determining readiness to return to full duty. Efforts are ongoing to understand and establish optimal return to duty criteria after an MSKI.<sup>72</sup> Concerns about career advancement opportunities and avoidance of duty limitations can cause underreporting

of MSKIs.<sup>73,74</sup> Alternatively, pursuing medical documentation (potentially for disability compensation) or justifying substandard duty performance can cause over-reporting of MSKIs.<sup>74</sup> Supplementary Table S5 highlights military MSKI surveillance and reporting issues.

## SUMMARY

Standardized exercise programming has reduced trainee MSKI rates. Secondary risk reduction initiatives show promise for reducing MSKI-related duty limitations and nondeployability among AC soldiers; timely identification/evaluation and appropriate, early management of MSKIs are essential. Tertiary risk reduction initiatives show promise for identifying soldiers whose chronic musculoskeletal conditions may render them unfit for continued military service.

Clinicians must document MSKI care with sufficient specificity (including diagnosis and external cause coding) to enable large-scale systematic MSKI surveillance and analysis informing focused MSKI risk reduction efforts. Historical changes in surveillance methods and injury definitions make it difficult to compare injury rates and trends over time. However, the U.S. Army’s standardized injury taxonomy will enable consistent classification of current and future injuries by mechanism of energy transfer and diagnosis. The Army’s electronic physical profiling system further enables standardized documentation of MSKI-related duty/work restrictions and mechanisms of injury. These evolving surveillance tools ideally ensure continual advancement of military injury surveillance and serve as models for other military and civilian health care organizations.

## SUPPLEMENTARY MATERIALS

Supplementary materials are available at *Military Medicine* online.

## REFERENCES

1. U.S. Army Public Health Center internal analysis, Defense Medical Surveillance System database, completed 2018.
2. Hauschild V, Hauret K, Richardson M, Jones B, Lee T: A Taxonomy of Injuries for Public Health Monitoring and Reporting (Public Health Information Paper No. 12-01-0717), 2017. Available at <http://www.dtic.mil/dtic/tr/fulltext/u2/1039481.pdf>; accessed September 5, 2018.
3. Hauschild V, Richardson M, Lee T, Hauret K, Jones B: Application of the Taxonomy of Injuries: Analysis of Army Recruit Injuries, CY 2016. Public Health Information Paper 12-01-0118, January 2018. Available at <http://www.dtic.mil/dtic/tr/fulltext/u2/1049222.pdf>; accessed August 25, 2018.
4. U.S. Army Medical Command internal analysis, Medical Operational Data System eProfile data, completed March 2019.
5. U.S. Army Public Health Center internal analysis, Medical Operational Data System eProfile data, completed August 2019.
6. Institute for Work and Health. Primary, secondary and tertiary prevention, April 2015. Available at <https://www.iwh.on.ca/what-researchers-mean-by/primary-secondary-and-tertiary-prevention>; accessed July 18, 2019.

7. Vergun D: Army implements new fitness standards for recruits and MOS transfers, January 3, 2017. Available at [https://www.army.mil/article/180199/army\\_implements\\_new\\_fitness\\_standards\\_for\\_recruits\\_and\\_mos\\_transfers](https://www.army.mil/article/180199/army_implements_new_fitness_standards_for_recruits_and_mos_transfers); accessed August 22, 2017.
8. Foulis S, Redmond J, Warr B, Zambraski E, Frykman P, Sharp M: USARIEM Technical Report T16-2: Development of the Occupational Physical Assessment Test (OPAT) for combat arms soldiers, 2015. Available at <https://dod.defense.gov/Portals/1/Documents/wisr-studies/Army%20-%20MEDCOM%20USARIEM%20Task%20Assessment3.pdf>; accessed June 17, 2019.
9. Knapik J, Jones B, Sharp M, et al: The case for pre-enlistment physical fitness testing: research and recommendations, 2004. Available at <http://www.dtic.mil/dtic/tr/fulltext/u2/a426848.pdf>; accessed August 5, 2019.
10. Hauret K, Canham-Chervak M, Schuh-Renner A, et al: Injury surveillance and longitudinal studies for gender integration in the Army, 2017, Second Annual Assessment (PHR No. S.0047783-17), 2018. Available at <https://apps.dtic.mil/dtic/tr/fulltext/u2/1062039.pdf>; accessed March 3, 2019.
11. Jones BH, Hauschild VD: Physical training, fitness, and injuries: lessons learned from military studies. *J Strength Cond Res* 2015; 29: S57–64.
12. Department of the U.S. Army. Army Physical Readiness Training, Field Manual 7-22, 2012. Available at [https://armypubs.army.mil/epubs/DR\\_pubs/DR\\_a/pdf/web/ARN7938\\_FM%207-22%20INC%20C1%20Final.pdf](https://armypubs.army.mil/epubs/DR_pubs/DR_a/pdf/web/ARN7938_FM%207-22%20INC%20C1%20Final.pdf); accessed December 1, 2019.
13. U.S. Army Public Health Center (provisional). 2015 Health of the Force Report. Spotlight: Physical Training During Basic Combat Training. Available at [http://ec.militarytimes.com/static/pdfs/Health\\_of\\_the\\_Force\\_Report\\_Nov\\_2015.pdf](http://ec.militarytimes.com/static/pdfs/Health_of_the_Force_Report_Nov_2015.pdf); accessed June 17, 2019.
14. Knapik J, Rieger W, Palkoska F, Van Camp S, Darakjy S: United States Army physical readiness training: rationale and evaluation of the physical training doctrine. *J Strength Cond Res* 2009; 23(4): 1353–62.
15. Molloy J: Factors influencing running-related musculoskeletal injury risk among U.S. military recruits. *Milit Med* 2016; 181(6): 512–23.
16. McClung JP, Gaffney-Stomberg E: Optimizing performance, health, and well-being: nutritional factors. *Milit Med* 2016; 181(1 Suppl): 86–91.
17. Lappe J, Cullen D, Haynatzki G, Recker R, Ahlf R, Thompson K: Calcium and vitamin D supplementation decreases incidence of stress fractures in female navy recruits. *J Bone Miner Res: Off J Am Soc Bone Miner Res* 2008; 23(5): 741–9.
18. Gaffney-Stomberg E, Lutz LJ, Rood JC, et al: Calcium and vitamin D supplementation maintains parathyroid hormone and improves bone density during initial military training: a randomized, double-blind, placebo controlled trial. *Bone* 2014; 68: 46–56.
19. Hughes JM, Foulis SA, Taylor KM, et al: A prospective field study of U.S. Army trainees to identify the physiological bases and key factors influencing musculoskeletal injuries: a study protocol. *BMC Musculoskelet Dis* 2019; 20(1): 282.
20. McClung JP, Marchitelli LJ, Friedl KE, Young AJ: Prevalence of iron deficiency and iron deficiency anemia among three populations of female military personnel in the US Army. *J Am Coll Nutr* 2006; 25(1): 64–9.
21. McClung JP, Karl JP, Cable SJ, Williams KW, Young AJ, Lieberman HR: Longitudinal decrements in iron status during military training in female soldiers. *Brit J Nutr* 2009; 102(4): 605–9.
22. Teyhen D, Shaffer S, Butler R, et al: What risk factors are associated with musculoskeletal injury in US Army Rangers? A prospective prognostic study. *Clin Orthop Relat Res* 2015; 473(9): 2948–58.
23. Teyhen D, Shaffer S, Goffar S, et al: Identification of risk factors prospectively associated with musculoskeletal injury in a warrior athlete population. *Sports Health*. 2020 (in press).
24. Lehr ME, Plisky PJ, Butler RJ, Fink ML, Kiesel KB, Underwood FB: Field-expedient screening and injury risk algorithm categories as predictors of noncontact lower extremity injury. *Scand J Med Sci Sports* 2013; 23(4): e225–32.
25. Bittencourt NFN, Meeuwisse WH, Mendonca LD, Nettel-Aguirre A, Ocarino JM, Fonseca ST: Complex systems approach for sports injuries: moving from risk factor identification to injury pattern recognition-narrative review and new concept. *Brit J Sport Med* 2016; 50(21): 1309–14.
26. Army Combat Fitness Test: Initial Operation Capability (1 October 2019 to 30 September 2020), October 2019. Available at [https://www.army.mil/e2/downloads/rv7/acft/acft\\_ioc.pdf](https://www.army.mil/e2/downloads/rv7/acft/acft_ioc.pdf); accessed November 13, 2019.
27. Brading T: ACFT is only one part of holistic health and fitness, experts say, 2019. Available at [https://www.army.mil/article/227217/acft\\_is\\_only\\_one\\_part\\_of\\_holistic\\_health\\_and\\_fitness\\_experts\\_say](https://www.army.mil/article/227217/acft_is_only_one_part_of_holistic_health_and_fitness_experts_say); accessed November 13, 2019.
28. Knapik JJ, Jones BH, Steelman RA: Physical training in boots and running shoes: a historical comparison of injury incidence in basic combat training. *Milit Med* 2015; 180(3): 321–8.
29. Knapik J, Trone D, Tchanda J, Jones B: Injury-reduction effectiveness of prescribing running shoes on the basis of foot arch height: summary of military investigations. *J Orth Sports Phys Ther* 2014; 44(10): 805–12.
30. U.S. Army Public Health Center: 2017 Health of the Force Report. Spotlight: Developing better strategic musculoskeletal readiness metrics. Available at [https://phc.amedd.army.mil/Periodical%20Library%20HealthoftheForceReportWeb\\_Printer.pdf](https://phc.amedd.army.mil/Periodical%20Library%20HealthoftheForceReportWeb_Printer.pdf); accessed June 17, 2019.
31. Saragiotta B, Yamato T, Hespanhol Junior L, Rainbow M, Davis I, Lopes A: What are the main risk factors for running-related injuries? *Sports medicine (Auckland, NZ)*. 2014; 44(8): 1153–63.
32. Wen DY: Risk factors for overuse injuries in runners. *Curr Sport Med Rep* 2007; 6(5): 307–13.
33. Fuller CW, Bahr R, Dick RW, Meeuwisse WH: A framework for recording recurrences, reinjuries, and exacerbations in injury surveillance. *Clin J Sport Med: Off J Can Acad Sport Med* 2007; 17(3): 197–200.
34. Fulton J, Wright K, Kelly M, et al: Injury risk is altered by previous injury: a systematic review of the literature and presentation of causative neuromuscular factors. *Int J Sport Phys Ther* 2014; 9(5): 583–95.
35. Zambraski EJ, Yancosek KE: Prevention and rehabilitation of musculoskeletal injuries during military operations and training. *J Strength Cond Res* 2012; 26(Suppl 2): S101–6.
36. Hill OT, Kay AB, Wahi MM, McKinnon CJ, Bulathsinhala L, Haley TF: Rates of knee injury in the U.S. active duty Army, 2000–2005. *Milit Med* 2012; 177(7): 840–4.
37. Wallace RF, Wahi MM, Hill OT, Kay AB: Rates of ankle and foot injuries in active-duty U.S. Army soldiers. *Milit Med* 2011; 176(3): 2000, 283–6, 290.
38. Arendt-Nielsen L, Graven-Nielsen T, Svensson P: Disturbances of pain perception in myofascial pain syndrome and other musculoskeletal pains. In: *Pathophysiology of Pain Perception. Plenum Series in Rehabilitation and Health*, pp 93–106. Edited by Lautenbacher S, Fillingim R. Boston, MA, Springer, 2004.
39. Peters M: Disturbances of pain perception in chronic back pain. In: *Pathophysiology of Pain Perception. Plenum Series in Rehabilitation and Health*, pp 59–75. Edited by Lautenbacher S, Fillingim R. Boston, MA, Springer, 2004.
40. Hootman JM, Macera CA, Ainsworth BE, Martin M, Addy CL, Blair SN: Predictors of lower extremity injury among recreationally active adults. *Clin J Sport Med: Off J Can Acad Sport Med* 2002; 12(2): 99–106.
41. Chapter 2: Department of the Army. The U.S. Military's Force Structure: A Primer. In: U.S. Congressional Budget Office, July 2016. Available at <https://www.cbo.gov/sites/default/files/114th-congress-2015-2016/reports/51535-fsprimerbreakoutchapter2.pdf>; accessed August 27, 2018.
42. MCoE Supplemental Manual 3-90: Force Structure Reference Data (Brigade Combat Teams). In: Fort Benning, GA: U.S. Army Maneuver Center of Excellence, September 2012. Available at [https://www.globalsecurity.org/military/library/policy/army/other/msm3-90\\_2012.pdf](https://www.globalsecurity.org/military/library/policy/army/other/msm3-90_2012.pdf); accessed August 27, 2018.
43. Talpey S, Siesmaa E: Sports injury prevention: the role of the strength and conditioning coach. *Strength Cond J* 2017; 39(3): 14–9.

44. Headquarters, Department of the U.S. Army. Army Techniques Publication No. 4-93. Sustainment Brigade. Available at [https://armypubs.army.mil/epubs/DR\\_pubs/DR\\_a/pdf/web/ATP%204-93%20FINAL%20WEB.pdf](https://armypubs.army.mil/epubs/DR_pubs/DR_a/pdf/web/ATP%204-93%20FINAL%20WEB.pdf); accessed July 22, 2019.
45. U.S. Army Public Health Center. 2017 Health of the Force Report. Spotlight: Measuring and reducing the impact of musculoskeletal conditions on medical readiness. Available at [https://phc.amedd.army.mil/Periodical%20Library/2017HealthoftheForceReportWeb\\_Printer.pdf](https://phc.amedd.army.mil/Periodical%20Library/2017HealthoftheForceReportWeb_Printer.pdf); accessed June 17, 2019.
46. U.S. Army Public Health Center. 2017 Health of the Force Report. Spotlight: Forward musculoskeletal care. Available at [https://phc.amedd.army.mil/Periodical%20Library/2017HealthoftheForceReportWeb\\_Printer.pdf](https://phc.amedd.army.mil/Periodical%20Library/2017HealthoftheForceReportWeb_Printer.pdf); accessed June 17, 2019.
47. Clewley D, Rhon DI, Flynn T, Sissel CD, Cook C: Does healthcare utilization before hip arthroscopy predict healthcare utilization after surgery in the United States military health system? An investigation into health seeking behavior. *J Orthop Sports Phys Ther* 2018; 1–27.
48. Clewley D, Rhon D, Flynn T, Koppenhaver S, Cook C: Health seeking behavior as a predictor of healthcare utilization in a population of patients with spinal pain. *PloS One* 2018; 13(8): e0201348.
49. Becker A, Held H, Redaelli M, et al: Low back pain in primary care: costs of care and prediction of future health care utilization. *Spine* 2010; 35(18): 1714–20.
50. Keeley P, Creed F, Tomenson B, Todd C, Borglin G, Dickens C: Psychosocial predictors of health-related quality of life and health service utilisation in people with chronic low back pain. *Pain*. 2008; 135(1–2): 142–50.
51. Ritzwoller DP, Crouse L, Shetterly S, Rublee D: The association of comorbidities, utilization and costs for patients identified with low back pain. *BMC Musculoskelet Disord* 2006; 7: 72.
52. Rosemann T, Joos S, Szecsenyi J, Laux G, Wensing M: Health service utilization patterns of primary care patients with osteoarthritis. *BMC Health Serv Res* 2007; 7: 169.
53. Piva SR, Fitzgerald GK, Wisniewski S, Delitto A: Predictors of pain and function outcome after rehabilitation in patients with patellofemoral pain syndrome. *J Rehabil Med* 2009; 41(8): 604–12.
54. Foster NE, Delitto A: Embedding psychosocial perspectives within clinical management of low back pain: integration of psychosocially informed management principles into physical therapist practice: challenges and opportunities. *Phys Ther* 2011; 91(5): 790–803.
55. Nicholas MK, Linton SJ, Watson PJ, Main CJ: Early identification and management of psychological risk factors ("yellow flags") in patients with low back pain: a reappraisal. *Phys Ther* 2011; 91(5): 737–53.
56. Sanders T, Foster NE, Bishop A, Ong BN: Biopsychosocial care and the physiotherapy encounter: physiotherapists' accounts of back pain consultations. *BMC Musculoskelet Disord* 2013; 14: 65.
57. Schuh A, Canham-Chervak M, Jones BH: Statistical process control charts for monitoring military injuries. *Injury Prev: J Int Soc Child Adolesc Injury Prev* 2017; 23(6): 416–22.
58. Knapik J, Hauret K, Jones B: Primary prevention of injuries in initial entry training. In: *Recruit Medicine*, pp 125–46. Edited by Lenhart M, Lounsbury D, North R. Fort Sam Houston, TX, Borden Institute, 2006.
59. Courie AF, Rivera MS, Pompey A: Managing public health in the Army through a standard community health promotion council model. *US Army Med Depart J* 2014; 82–90.
60. Canham-Chervak M, Schuh-Renner A, Steelman R, Jones B: Execution of MEDCOM Operational Order 15-74 (Improving Readiness through Reduction of Unintentional Injuries), August 2016–December 2017. In: U.S. Army Public Health Center IPP, ed. Aberdeen Proving Ground, MD, 2017. Available at <https://apps.dtic.mil/dtic/tr/fulltext/u2/1057120.pdf>; accessed June 17, 2019.
61. U.S. Army Medical Command internal analysis, Soldier Outcomes Trajectory Assessment (SOTA) database, completed July 2019.
62. U.S. Army Public Health Center. 2018 Health of the Force Report. Health Metrics-Injury. Available at <https://phc.amedd.army.mil/Periodical%20Library/2018HealthoftheForceReport.pdf>; accessed March 14, 2019.
63. U.S. Army Public Health Center. 2018 Health of the Force Report. Welcome to the 2018 Health of the Force Report. Available at <https://phc.amedd.army.mil/Periodical%20Library/2018HealthoftheForceReport.pdf>; accessed March 14, 2019.
64. Hedegaard H, Johnson RL, Warner M, Chen LH, Annett JL: Proposed framework for presenting injury data using the international classification of diseases, tenth revision, clinical modification (ICD-10-CM) diagnosis codes. *Nat Health Stat Rep* 2016; 89: 1–20.
65. Jones B, Canham-Chervak M, Canada S, Mitchener T, Moore S: Medical surveillance of injuries in the U.S. military: descriptive epidemiology and recommendations for improvement. *Am J Prev Med*. 2010; 38(1 Suppl): S42–60.
66. Canham-Chervak M, Steelman RA, Schuh A, Jones BH: Importance of external cause coding for injury surveillance: lessons from assessment of overexertion injuries among U.S. Army soldiers in 2014. *MSMR* 2016; 23(11): 10–5.
67. Roy TC, Knapik JJ, Ritland BM, Murphy N, Sharp MA: Risk factors for musculoskeletal injuries for soldiers deployed to Afghanistan. *Aviat Space Environ Med* 2012; 83(11): 1060–6.
68. Roy TC, Ritland BM, Sharp MA: A description of injuries in men and women while serving in Afghanistan. *Milit Med* 2015; 180(2): 126–31.
69. Teyhen DS, Goffar SL, Shaffer SW, et al: Incidence of musculoskeletal injury in US Army unit types: a prospective cohort study. *J Orthopaed Sports Phys Ther* 2018; 48(10): 749–57.
70. Sulsky SI, Bulzacchelli MT, Zhu L, et al: Risk factors for training-related injuries during U.S. Army basic combat training. *Milit Med* 2018; 183(Suppl 1): 55–65.
71. Magel J, Kim J, Thackeray A, Hawley C, Petersen S, Fritz JM: Associations between physical therapy continuity of care and health care utilization and costs in patients with low Back pain: a retrospective cohort study. *Phys Ther* 2018; 98(12): 990–9.
72. Rhon DI, Teyhen DS, Shaffer SW, Goffar SL, Kiesel K, Plisky PP: Developing predictive models for return to work using the military power, performance and prevention (MP3) musculoskeletal injury risk algorithm: a study protocol for an injury risk assessment programme. *Injury Prev: J Int Soc Child Adolesc Injury Prev* 2018; 24(1): 81–8.
73. Sauers SE, Smith LB, Scofield DE, Cooper A, Warr BJ: Self-Management of Unreported Musculoskeletal Injuries in a U.S. Army Brigade. *Milit Med* 2016; 181(9): 1075–80.
74. Smith L, Westrick R, Sauers S, et al: Underreporting of musculoskeletal injuries in the US Army: findings from an infantry brigade combat team survey study. *Sports Health* 2016; 8(6): 507–13.