

## MUTATIONAL ANALYSIS OF THE BH3 DOMAINS OF PROAPOPTOTIC Bcl-2 FAMILY GENES *Bad*, *Bmf* AND *Bcl-G* IN LARYNGEAL SQUAMOUS CELL CARCINOMAS

Nam Jin Yoo, Young Hwa Soung, Sang Hak Lee, Eun Goo Jeong, and Sug Hyung Lee

*Department of Pathology, College of Medicine, The Catholic University of Korea, Seoul, Korea*

**Aims:** There is mounting evidence that deregulation of apoptosis is involved in the mechanisms of cancer development. Somatic mutations of apoptosis-related genes have been reported in many human cancers. The aim of this study was to explore the possibility that mutation of the BH3 domains of the proapoptotic Bcl-2 genes *Bad*, *Bmf* and *Bcl-G* might be involved in the development of laryngeal cancer.

**Methods:** We analyzed the BH3 domains of *Bad*, *Bmf* and *Bcl-G* for the detection of somatic mutations in 33 squamous cell carcinomas of the larynx by a polymerase chain

reaction-based single-strand conformation polymorphism assay.

**Results:** There were no somatic mutations of the BH3 domains of *Bad*, *Bmf* and *Bcl-G* in the laryngeal squamous cell carcinoma samples.

**Conclusions:** The data presented here indicate that BH3 domain mutation of the proapoptotic genes *Bad*, *Bmf* and *Bcl-G* is rare in laryngeal squamous cell carcinoma and may not contribute to the apoptosis-resistance mechanisms of laryngeal squamous cell carcinoma.

**Key words:** apoptosis, BH3 domain, laryngeal cancer, mutation.

---

*Correspondence to:* Dr Sug Hyung Lee, Department of Pathology, College of Medicine, The Catholic University of Korea, 505 Banpo-dong, Socho-gu, Seoul 137-701, Korea. Tel +82-2-5901188; fax +82-2-5376586; e-mail suhulee@catholic.ac.kr

*Acknowledgments:* This work was supported by funding from the Korea Research Foundation in the program year 2006 (E00049).

Received September 18, 2006; accepted November 21, 2006.