

Muttalib–Borodin ensembles in random matrix theory — realisations and correlation functions

Peter J. Forrester* Dong Wang†

Abstract

Muttalib–Borodin ensembles are characterised by the pair interaction term in the eigenvalue probability density function being of the form $\prod_{1 \leq j < k \leq N} (\lambda_k - \lambda_j)(\lambda_k^\theta - \lambda_j^\theta)$. We study the Laguerre and Jacobi versions of this model — so named by the form of the one-body interaction terms — and show that for $\theta \in \mathbb{Z}^+$ they can be realised as the eigenvalue PDF of certain random matrices with Gaussian entries. For general $\theta > 0$, realisations in terms of the eigenvalue PDF of ensembles involving triangular matrices are given. In the Laguerre case this is a recent result due to Cheliotis, although our derivation is different. We make use of a generalisation of a double contour integral formula for the correlation functions contained in a paper by Adler, van Moerbeke and Wang to analyse the global density (which we also analyse by studying characteristic polynomials), and the hard edge scaled correlation functions. For the global density functional equations for the corresponding resolvents are obtained; solving this gives the moments in terms of Fuss–Catalan numbers (Laguerre case — a known result) and particular binomial coefficients (Jacobi case). For $\theta \in \mathbb{Z}^+$ the Laguerre and Jacobi cases are closely related to the squared singular values for products of θ standard Gaussian random matrices, and truncations of unitary matrices, respectively. At the hard edge the double contour integral formulas provide a double contour integral form of the scaled correlation kernel obtained by Borodin in terms of Wright’s Bessel function.

Keywords: Muttalib–Borodin ensembles; determinantal processes; biorthogonal polynomials.

AMS MSC 2010: Primary 60B20, Secondary 15B52; 82D35; 30E10.

Submitted to EJP on October 8, 2016, final version accepted on April 26, 2017.

Supersedes arXiv:1502.07147.

*Department of Mathematics and Statistics, The University of Melbourne, Victoria 3010, Australia; ARC Centre of Excellence for Mathematical & Statistical Frontiers. E-mail: p.forrester@ms.unimelb.edu.au

†Department of Mathematics, National University of Singapore, Singapore, 119076. E-mail: matwd@nus.edu.sg

1 Introduction

Recent studies in random matrix theory [18, 34, 17, 14, 24] have drawn renewed attention to the class of eigenvalue probability density functions (PDFs) proportional to

$$\prod_{l=1}^N e^{-V(\lambda_l)} \prod_{1 \leq j < k \leq N} (\lambda_k - \lambda_j)(\lambda_k^\theta - \lambda_j^\theta), \quad \lambda_l > 0. \quad (1.1)$$

These PDFs were proposed by Muttalib [43] in the context of a simplified model of the joint distribution of the transmission eigenvalues for disordered conductors in the metallic regime, and with no time reversal symmetry. The latter is known to have its exact form proportional to [8]

$$\prod_{l=1}^N e^{-V(\lambda_l)} \prod_{1 \leq j < k \leq N} (\lambda_k - \lambda_j) \left(\operatorname{arsinh}^2 \lambda_k^{1/2} - \operatorname{arsinh}^2 \lambda_j^{1/2} \right), \quad \lambda_l > 0, \quad (1.2)$$

where for large λ , $V(\lambda) = Nc \operatorname{arsinh}^2 \lambda^{1/2} (1 + \mathcal{O}(N^{-1}))$, with $c = \ell/L$, ℓ denoting the mean free path length and L the length of the wire. In practice one has $1 \ll c \ll N$.

Recalling that $\operatorname{arsinh} z = \log(z + \sqrt{z^2 + 1})$ one sees (1.1) relates to (1.2) in the limit $\theta \rightarrow 0^+$ when we have

$$\frac{1}{\theta} (\lambda_k^\theta - \lambda_j^\theta) \rightarrow (\log \lambda_k - \log \lambda_j), \quad (1.3)$$

although this is still only an approximation to the corresponding factor in (1.2). Actually in [43] attention was restricted to θ a positive integer; on this point we remark that the change of variables

$$\lambda \rightarrow \lambda^{1/\theta} \quad (1.4)$$

maps θ to $1/\theta$ in (1.1) at the expense of altering $V(\lambda)$.

Our interest is two special cases of (1.1). The first is when

$$e^{-V(\lambda)} = \lambda^c e^{-\lambda}, \quad \lambda > 0, \quad c > -1. \quad (1.5)$$

This is referred to as the Laguerre weight, due to its appearance as the weight function in the orthogonality of the Laguerre polynomials in the theory of classical orthogonal polynomials. The choice (1.5), together with the choice of $e^{-V(\lambda)}$ as a Gaussian or Jacobi weight (for the latter see (1.6) below), was considered in some detail by Borodin [13]. Due to the significant advancement contained in [13], we will refer to the general class of PDFs (1.1) as Muttalib–Borodin ensembles, and the particular choice of weight (1.5) in (1.1) as the Laguerre Muttalib–Borodin ensemble.

Let us now describe our results. In relation to the Laguerre Muttalib–Borodin ensemble, we first realise the special cases $c, \theta \in \mathbb{Z}^+$ as a particular class of complex Wishart matrices isolated in [2]. For general parameters we give a new derivation of a recent result of Cheliotis [17] which gives a realisation in terms of a particular class of random upper-triangular matrices, and we furthermore develop working in [2] to generalise this result (Section 2). The differential equation satisfied by the characteristic polynomial of the ensemble under the mapping (1.4) is studied, and we relate this to the resolvent and global density (Section 3). We use results contained in [2], and take inspiration from the recent work [37], to obtain a derivation of the global density directly from a double contour formula for the one-point function using the saddle point method (Section 4). Furthermore the double contour integral form of the correlation kernel given in [2], suitably generalised from integer to real parameters, is used to rederive the hard edge scaled limit known from [13] (Section 5).

Parallel to the analysis of the Laguerre Muttalib–Borodin ensemble, we also undertake an analogous program of study in relation to the Jacobi weight

$$e^{-V(\lambda)} = \lambda^{c_1}(1 - \lambda)^{c_2}, \quad 0 < \lambda < 1, \quad c_1, c_2 > -1. \tag{1.6}$$

This substituted in (1.1) gives the Jacobi Muttalib–Borodin ensemble. Our realisation and double contour integral formula for the correlation kernel makes essential use of results contained in [2]. In relation to the global density, the resolvent is specified by a nonlinear equation which we solve using the Lagrange inversion formula to deduce that the moments of the global density are given in terms of particular binomial coefficients. A trigonometric parametrisation of the spectral variable is given which allows for the determination of an explicit functional form for the global density. The hard edge scaled limit gives the same double contour integral form as found for the Laguerre case, in keeping with the findings of [13].

We now give a precise statement of the main results in our paper for the Laguerre Muttalib–Borodin ensemble. All the results obtained for the Laguerre case have counterparts for the Jacobi case, described in the paragraph above and which are presented in the body of the paper subsequent to presentation of the Laguerre case. We do not state the results in their most general form below, and readers can find the generalisations in subsequent sections.

1.1 Main results for Laguerre Muttalib–Borodin ensemble

First, the Laguerre Muttalib–Borodin ensemble can be realised as the eigenvalues of a random matrix in the *upper-triangular random matrix ensemble*, which is defined in Section 2.1, and if $\theta, c \in \mathbb{Z}_{\geq 0}$, it can be realised as the eigenvalues of a random matrix in the *multiple Laguerre ensemble*, which is defined in [2] and also described in Section 2.1.

Proposition 1.1. Let Y be a upper-triangular $N \times N$ matrix with all entries independent, the strictly upper-triangular entries distributed as standard complex Gaussians, and the diagonal entries are real positive random variables with $|y_{k,k}|^2 \stackrel{d}{=} \Gamma[\theta(k - 1) + c + 1, 1]$, or equivalently, $2|y_{k,k}|^2 \stackrel{d}{=} \chi_{2(\theta(k-1)+c+1)}^2$, for $k = 1, \dots, N$. Then the eigenvalues $\lambda_1, \dots, \lambda_n$ of $Y^\dagger Y$ have the PDF proportional to (1.1) with V given by (1.5). Furthermore, if $\theta, c \in \mathbb{Z}_{\geq 0}$, the PDF of the eigenvalues of Y is the same as the PDF of the eigenvalues of $X^\dagger X$, where X is an $N \times N$ random matrix whose entries $x_{j,k}$ with $1 \leq j \leq \theta(k - 1) + c + k$ have independent standard complex normal distribution, while other entries are zero.

This result is covered by Proposition 2.1 and Corollary 2.3.

The statistical system defined by the Laguerre Muttalib–Borodin ensemble is a determinantal point process, and so is fully determined by its correlation kernel, for which we give a double contour integral formula.

Proposition 1.2. The correlation kernel for the Laguerre Muttalib–Borodin ensemble can be written

$$K^L(x, y) = \frac{1}{(2\pi i)^2} \frac{h(x)}{h(y)} \oint_{\Sigma} dz \oint_{\Gamma_\alpha} dw \frac{x^{-z-1} y^w \Gamma(z + 1) \prod_{k=1}^N (z - \alpha_k)}{(z - w) \Gamma(w + 1) \prod_{l=1}^N (w - \alpha_l)}, \tag{1.7}$$

where $\alpha_j = \theta(j - 1) + c$ is specified in (2.3), and the contours Σ and Γ_α are specified in Proposition 2.10. The function $h(x) = x^{c/2} e^{x/2}$, as specified in (4.3) up to a multiplicative constant.

This result is covered by Proposition 2.10 and the definition (4.2) of the kernel.

Next, we derive the limiting global density of the ensemble, defined in terms of the correlation kernel $K^L(x, x)$ according to (4.4).

Proposition 1.3. For all $\theta > 0$, the limiting global density of the Laguerre Muttalib–Borodin ensemble with change of variable (1.4) is the Fuss–Catalan distribution, that is,

$$\tilde{\rho}_{(1)}^L(x) = \rho^{\text{F-C}}(x). \tag{1.8}$$

Here the the Fuss–Catalan distribution is defined in (4.7). We remark that this result is not totally new, as we explain in Proposition 3.1, and after proper interpretation can be proved by a different method. But this method does not generalise simply to the Jacobi case, so we introduce two alternative methods to prove Proposition 1.3, one in Section 3.1 for positive integer θ and the other in Section 4.1 for general $\theta > 0$. Both of these two methods can be applied to the Jacobi case with little change.

Finally, we consider the local behaviour of $K^L(x, y)$ around 0, first obtained by Borodin [13] in terms of Wright’s Bessel function.

Proposition 1.4. We have

$$\begin{aligned} & \lim_{N \rightarrow \infty} N^{-1/\theta} K^L(N^{-1/\theta} x, N^{-1/\theta} y) \\ &= \left(\frac{y}{x}\right)^{c/2} \frac{\theta}{(2\pi i)^2} \oint_{\Sigma_{-1/2}^\delta} dz \oint_{\Gamma_0} dw \frac{x^{-\theta z-1} y^{\theta w}}{z-w} \frac{\Gamma(\theta z + c + 1) \Gamma(z + 1) \sin \pi z}{\Gamma(\theta w + c + 1) \Gamma(w + 1) \sin \pi w}, \end{aligned} \tag{1.9}$$

where Γ_0 is the Hankel loop contour starting at $\infty + i\epsilon$, running parallel to the positive real axis, looping around the origin, and finishing at $\infty - i\epsilon$ after again running parallel to the negative real axis, while $\Sigma_{-1/2}^\delta$ is the contour consisting of two rays, one is from $-1/2$ to $e^{(\pi/2+\delta)i} \cdot \infty$ and the other from $e^{-(\pi/2+\delta)i} \cdot \infty$ to $-1/2$, where $\delta \in (0, \pi/2)$, see Figure 7. If $\theta \geq 1$, we can also take $\delta = 0$ and let $\Sigma_{-1/2}^\delta$ be the upward vertical contour through $-1/2$. In the case $\theta \in \mathbb{Z}^+$ this can be rewritten

$$\begin{aligned} & x^{1/\theta-1} \lim_{N \rightarrow \infty} N^{-1/\theta} K^L(\theta(x/N)^{1/\theta}, \theta(y/N)^{1/\theta}) = \left(\frac{y}{x}\right)^{c/(2\theta)} \\ & \times \frac{1}{(2\pi i)^2} \oint_{\Sigma_{-1/2}^\delta} dz \oint_{\Gamma_0} dw \frac{x^{-z-1} y^w \prod_{k=0}^{\theta-1} \Gamma(z + (c + 1 + k)/\theta) \Gamma(z + 1) \sin \pi z}{z-w \prod_{k=0}^{\theta-1} \Gamma(w + (c + 1 + k)/\theta) \Gamma(w + 1) \sin \pi w}. \end{aligned} \tag{1.10}$$

This result is proved in Section 5.1.1. We remark that the $\theta \rightarrow 0_+$ limit of Propositions 1.3 and 1.4 are also obtained in this paper, see Sections 4.1.2 and 5.1.2.

2 Realisations and extensions

2.1 The Laguerre upper-triangular ensemble

We use the term complex Wishart matrix to refer to a random matrix of the form $W^\dagger W$ with W containing complex Gaussian entries with mean and standard deviation to be specified. Furthermore, for any $M \times N$ matrix A , we denote $A_{m \times n}$ as the $m \times n$ matrix consisting of the upper-left $m \times n$ block of A . We are interested in a particular complex Wishart matrix, due to Adler, van Moerbeke and Wang [2], which is parametrised by non-negative integers $\alpha_1, \dots, \alpha_N$ satisfying

$$1 + \alpha_1 \leq 2 + \alpha_2 \leq \dots \leq N + \alpha_N \leq M, \tag{2.1}$$

with $M \geq N$. One defines the $M \times N$ random matrix $X = [x_{j,k}]_{\substack{j=1,\dots,M \\ k=1,\dots,N}}$ to have entries

$$x_{j,k} \stackrel{\text{d}}{=} \begin{cases} \text{N}[0, 1/2] + i\text{N}[0, 1/2], & 1 \leq j \leq k + \alpha_k \\ 0, & \text{otherwise.} \end{cases}$$

All nonzero entries of X are therefore standard complex Gaussians. Moreover, the condition (2.1) implies all entries on and above the diagonal of X are non-zero. Due to

its relationship to a certain family of special functions by the same name, the ensemble of matrices $X_{M \times n}^\dagger X_{M \times n}$, ($n = 1, \dots, M$) was referred to in [2] as the multiple Laguerre ensemble. Our first result is to specify an upper-triangular matrix obtained from X by a sequence of Householder transformations. Such transformations were introduced into random matrix theory in [51], [48], and furthermore underpin the construction of β -ensembles as formulated in [21].

We denote by $\Gamma[k, \sigma]$ the gamma distribution, specified by the density function $(\sigma^{-k}/\Gamma(k))t^{k-1}e^{-t/\sigma}$, ($t > 0$). Let $Y = [y_{j,k}]_{j,k=1,\dots,N}$ be an upper-triangular $N \times N$ random matrix with all entries independent. The strictly upper-triangular entries are distributed as standard complex Gaussians, and the diagonal entries are real positive random variables with distributions depending on parameters $\alpha_k > -1$ specified by

$$|y_{k,k}|^2 \stackrel{d}{=} \Gamma[\alpha_k + 1, 1], \quad \text{or equivalently} \quad 2|y_{k,k}|^2 \stackrel{d}{=} \chi_{2(\alpha_k+1)}^2, \quad (k = 1, \dots, N). \quad (2.2)$$

We say that Y is a random matrix in the upper-triangular ensemble, and that $Y^\dagger Y$ belongs to the Laguerre upper-triangular ensemble.

Proposition 2.1. The random matrices $X_{M \times 1}^\dagger X_{M \times 1}$, $X_{M \times 2}^\dagger X_{M \times 2}$, \dots , $X_{M \times N}^\dagger X_{M \times N}$ and $Y_{N \times 1}^\dagger Y_{N \times 1}$, $Y_{N \times 2}^\dagger Y_{N \times 2}$, \dots , $Y_{N \times N}^\dagger Y_{N \times N}$ have the same joint distribution. In particular, the eigenvalues of $X_{M \times 1}^\dagger X_{M \times 1}$, $X_{M \times 2}^\dagger X_{M \times 2}$, \dots , $X_{M \times N}^\dagger X_{M \times N}$ have the same joint distribution as the eigenvalues of $Y_{N \times 1}^\dagger Y_{N \times 1}$, $Y_{N \times 2}^\dagger Y_{N \times 2}$, \dots , $Y_{N \times N}^\dagger Y_{N \times N}$.

Proof. Recall that a complex Householder reflection matrix acting on the left of the $M \times N$ matrix X has the form

$$U = \mathbb{I}_M - 2\vec{u}\vec{u}^\dagger,$$

where \dagger denotes the operation of complex conjugate and transpose and \vec{u} is a $M \times 1$ complex column vector with the property that $\vec{u}^\dagger \cdot \vec{u} = 1$. This latter requirement implies $U^\dagger U = \mathbb{I}_M$, so U is unitary. Geometrically U corresponds to a reflection in the complex hyperplane orthogonal to \vec{u}^\dagger .

To prove the proposition, we construct a sequence of $M \times N$ random matrices $X^{(0)} = X, X^{(1)}, \dots, X^{(N)}$ and a sequence of $M \times M$ random Householder reflection matrices $U^{(1)}, \dots, U^{(N)}$ inductively. The matrix $U^{(l)}$ is determined by $X^{(l-1)}$ and $X^{(l+1)} = U^{(l+1)}X^{(l)}$ ($l = 0, 1, \dots, N - 1$), where $X^{(l)}$ satisfies: (1) the (j, k) entry is zero for $k < j \leq M$ if $1 \leq k \leq l$, or $k + \alpha_k < j \leq M$ if $l < k \leq N$; (2) the diagonal (k, k) entry is real positive and its square is in $\Gamma[\alpha_k + 1, 1]$ distribution if $k \leq l$; (3) all other entries are in complex standard Gaussian distribution. By the method of construction to be detailed below, the block of $X^{(N)}$ consisting of the first N rows is such that $X_{N \times N}^{(N)} \stackrel{d}{=} Y$, and all entries in the last $(M - N)$ rows of $X^{(N)}$ are zeros. Thus the joint distribution of $(X_{M \times n}^{(N)})^\dagger X_{M \times n}^{(N)}$ for $n = 1, \dots, N$ is the same as that of $Y_{N \times n}^\dagger Y_{N \times n}$. On the other hand, $X^{(N)}$ is a function of X and for any n , $(X_{M \times n}^{(N)})^\dagger X_{M \times n}^{(N)} = X_{M \times n}^\dagger X_{M \times n}$. Except for the construction of $\{X^{(l)}\}$, this finishes the proof.

We now give an algorithm for the construction using induction. For any $l = 0, \dots, N - 1$, by the induction assumption $X^{(l)} = [x_{j,k}^{(l)}]_{j=1,\dots,M, k=1,\dots,N}$ is well defined and $x_{j,l+1}^{(l)}$ with $j = l + 1, \dots, l + 1 + \alpha_{l+1}$ are in independent complex standard Gaussian distribution. We denote the $(1 + \alpha_{l+1})$ -dimensional vector

$$\vec{x}^{(l+1)} = (x_{l+1,l+1}^{(l)}, x_{l+2,l+1}^{(l)}, \dots, x_{l+1+\alpha_{l+1},l+1}^{(l)})^T.$$

We construct the Householder reflection matrix $U^{(l+1)}$ using $\vec{u}^{(l+1)}$, which is a concatenation of an l -dimensional zero vector, an $(1 + \alpha_{l+1})$ -dimensional vector $\vec{v}^{(l+1)}$, and an

$(M - l - 1 - \alpha_{l+1})$ -dimensional zero vector. Here the vector $\vec{x}^{(l+1)}$ is defined as the unit vector

$$\frac{\vec{w}}{\|\vec{w}\|}, \quad \text{where } \vec{w} = \vec{x}^{(l+1)} - (\|\vec{x}^{(l+1)}\|, 0, \dots, 0)^T,$$

if $\vec{x}^{(l+1)} \neq 0$, and is defined simply as $(1, 0, \dots, 0)^T$ if $\vec{x}^{(l+1)} = 0$.

From the definition of $U^{(l+1)}$, and recalling $X^{(l+1)} = U^{(l+1)}X^{(l)}$, it is clear that $X^{(l+1)}$ and $X^{(l)}$ are identical in the upper block consisting of the first l rows and the lower block consisting of the last $(M - l - 1 - \alpha_{l+1})$ rows. In the middle block consisting of the remaining $1 + \alpha_{l+1}$ rows, the left part consisting of the middle part of the left-most l columns of $X^{(l)}$ are zeros, so they remain zero in $X^{(l+1)}$. The entries of $X^{(l)}$ in the right part of the middle block consisting of the right-most $(N - l - 1)$ columns are in independent complex standard Gaussian distribution, and they are all independent of $\vec{w}^{(l+1)}$. So after the left multiplication by $U^{(l+1)}$, the entries of $X^{(l+1)}$ in that part of the middle block are also in independent complex standard Gaussian distribution by the rotational invariance of random Gaussian vectors. The $(l + 1)$ -th column of the middle block becomes $(\|\vec{x}^{(l+1)}\|, 0, \dots, 0)^T$, so its first entry is positive and the square of the first entry is in $\Gamma[\alpha_k + 1, 1]$ distribution, while all other entries are zero. Hence each $X^{(l+1)}$ satisfies the required properties, so finishing the proof by induction. \square

Remark 2.2. Since we are interested in the spectral properties of $X_{M \times n}^\dagger X_{M \times n}$ in the multiple Laguerre ensemble of [2], and by Proposition 2.1 they are identical to $Y_{N \times n}^\dagger Y_{N \times n}$ with $\alpha_1, \dots, \alpha_N$ restricted to some subset to their domain, we can think of the upper-triangular ensemble as a generalisation of the multiple Laguerre ensemble.

The upper-triangular matrix Y distributed as in (2.2) with

$$\alpha_j = \theta(j - 1) + c \quad (j = 1, \dots, N), \tag{2.3}$$

has recently been shown by Cheliotis [17] to have the PDF for its squared singular values given by the Laguerre Muttalib–Borodin ensemble (1.1) with weight (1.5). For this to relate to our construction from complex Gaussian matrices according to Proposition 2.1 we must have θ and c non-negative integers. Thus in this circumstance we have identified a realisation of this ensemble as the eigenvalue PDF of a Wishart matrix.

Corollary 2.3. Consider the matrix X as defined below (2.1), and with $\theta, c \in \mathbb{Z}_{\geq 0}$, let α_j be specified as in (2.3). We have that the eigenvalue PDF of the Wishart matrix $X^\dagger X$ is given by the Laguerre Muttalib–Borodin ensemble (1.1) with weight (1.5).

Corollary 2.3 is a special case of Corollary 2.8(a) below. Additional details of the spectral properties of $X^\dagger X$, or equivalently according to Proposition 2.1, of $Y^\dagger Y$ with the parameters α_j of integer values, beyond the joint distribution of the eigenvalues have been given in [2]. In particular, we can read off from the multiple Laguerre part of [2, Thm. 1] the explicit functional form of the conditional distribution of the eigenvalues of $Y_{N \times n}^\dagger Y_{N \times n}$, given the eigenvalues of $Y_{N \times (n-1)}^\dagger Y_{N \times (n-1)}$, where $Y_{p \times q}$ denotes the top $p \times q$ sub-block of Y . Below we show that the results can be generalised to arbitrary upper-triangular random matrices Y with real-valued $\alpha_j > -1$. The proofs are similar to those in [2] and we mainly emphasis the differences.

Proposition 2.4. Let the $N \times N$ random matrix Y be in the upper-triangular ensemble with diagonal entries specified by (2.2). Denote by $\{\lambda_1, \dots, \lambda_n\}$ and $\{\mu_1, \dots, \mu_{n-1}\}$ the eigenvalues of $Y_{N \times n}^\dagger Y_{N \times n}$ and $Y_{N \times (n-1)}^\dagger Y_{N \times (n-1)}$ respectively in descending order. The

conditional PDF of $\{\lambda_1, \dots, \lambda_n\}$, with $\{\mu_1, \dots, \mu_{n-1}\}$ fixed and distinct, is equal to

$$p_{n,n-1}(\{\lambda_j\}_{j=1}^n, \{\mu_j\}_{j=1}^{n-1}) = \frac{1}{\Gamma(\alpha_n + 1)} \prod_{k=1}^n \lambda_k^{\alpha_n} e^{-\lambda_k} \prod_{l=1}^{n-1} \mu_l^{-\alpha_n-1} e^{\mu_l} \frac{\prod_{1 \leq j < k \leq n} (\lambda_j - \lambda_k)}{\prod_{1 \leq j < k \leq n-1} (\mu_j - \mu_k)}, \quad (2.4)$$

subject to the interlacing constraint

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \geq \lambda_n \geq 0. \quad (2.5)$$

Actually we can prove a slightly stronger result:

Lemma 2.5. *Let $A = (a_{i,j})$ be an $(n - 1) \times (n - 1)$ matrix with the eigenvalues of $A^\dagger A$ being $\{\mu_1, \dots, \mu_{n-1}\}$ in descending order. Define the $n \times n$ matrix $B = (b_{i,j})$ by letting: (1) the $(n - 1) \times (n - 1)$ upper-triangular block equal A ; (2) the bottom row has all entries but the rightmost one equal 0; (3) all entries of the rightmost row be independent random variables, such that the (n, n) -entry $b_{n,n}$ is real positive with $|b_{n,n}|^2 \stackrel{d}{=} \Gamma[\alpha_n + 1, 1]$, and all other entries in the row are in standard complex normal distribution. Then the eigenvalues of $B^\dagger B$, denoted by $\{\lambda_1, \dots, \lambda_n\}$ in descending order, satisfies the interlacing constraint (2.5) and have the distribution given by $p_{n,n-1}(\{\lambda_j\}_{j=1}^n, \{\mu_j\}_{j=1}^{n-1})$ in (2.4).*

Proof of Lemma 2.5. A key point is that

$$BB^\dagger = B_{n \times (n-1)} B_{n \times (n-1)}^\dagger + \vec{y} \vec{y}^\dagger, \quad (2.6)$$

where $\vec{y} = (b_{1,n}, b_{2,n}, \dots, b_{n,n})^T$. The distribution of $b_{k,n}$ implies that

$$|b_{k,n}|^2 \stackrel{d}{=} \Gamma[1, 1] \quad (k = 1, \dots, n - 1), \quad |b_{n,n}|^2 \stackrel{d}{=} \Gamma[\alpha_n + 1, 1]. \quad (2.7)$$

Moreover there exists an $(n - 1) \times (n - 1)$ unitary matrix V depending on $A = B_{(n-1) \times (n-1)}$ such that

$$(V \oplus \mathbb{I}_1) B_{n \times (n-1)} B_{n \times (n-1)}^\dagger (V \oplus \mathbb{I}_1)^\dagger = \text{diag}[\mu_1, \dots, \mu_{n-1}, 0], \quad (2.8)$$

where

$$V \oplus \mathbb{I}_1 = \begin{bmatrix} V & \mathbf{0}_{(n-1) \times 1} \\ \mathbf{0}_{1 \times (n-1)} & 1 \end{bmatrix}.$$

Using the property that the multiplication of a unitary matrix and a vector of independent standard complex Gaussians yields another vector of independent standard complex Gaussians, we have that the vector $\vec{z} = (z_1, \dots, z_n)^T$ defined as

$$\vec{z} = (V \oplus \mathbb{I}_1) \vec{y}$$

has the properties that all its components are independent, z_1, \dots, z_{n-1} are in standard complex normal distribution, and

$$|z_k|^2 \stackrel{d}{=} \Gamma[1, 1] \quad (k = 1, \dots, n - 1), \quad \text{and} \quad z_n = b_{n,n}.$$

Conjugating both sides of (2.6) as in (2.8), we have that

$$\text{ePDF } B^\dagger B = \text{ePDF } BB^\dagger = \text{ePDF} \left(\text{diag}[\mu_1, \dots, \mu_{n-1}, 0] + \vec{z} \vec{z}^\dagger \right) \quad (2.9)$$

where ePDF denotes the eigenvalue PDF.

By a standard manipulation of the characteristic polynomial, one can show that the eigenvalue equation for the matrix on the RHS of (2.9) is given by

$$0 = 1 + \left(-\frac{|z_n|^2}{\lambda} + \sum_{k=1}^{n-1} \frac{|z_k|^2}{\mu_k - \lambda} \right). \tag{2.10}$$

The distribution of the roots of this rational function, with residues distributed according to (2.7), and thus the conditional PDF of $\{\lambda_1, \dots, \lambda_n\}$, can now be read off as a special case of [26, Cor. 3], and (2.4) with interlacing constraint (2.5) follows. \square

Knowledge of Proposition 2.4 allows us to rederive the result of Cheliotis [17] noted below (2.3). We will require the use of a particular multiple integral evaluation.

Lemma 2.6. *Let $R_{\leq \lambda}$ denote the region (2.5) for $(\mu_1, \dots, \mu_{n-1})$, and suppose $\alpha_k \neq \alpha_n$ ($k = 1, \dots, n - 1$). We have*

$$\begin{aligned} & \prod_{l=1}^n \lambda_l^{\alpha_n} \int_{R_{\leq \lambda}} \prod_{l=1}^{n-1} \mu_l^{-\alpha_n-1} \det[\mu_j^{\alpha_k}]_{j,k=1,\dots,n-1} d\mu_1 \cdots d\mu_{n-1} \\ &= \prod_{k=1}^{n-1} \frac{1}{\alpha_k - \alpha_n} \det[\lambda_j^{\alpha_k}]_{j,k=1,\dots,n}. \end{aligned} \tag{2.11}$$

Proof. We have

$$\prod_{l=1}^{n-1} \mu_l^{-\alpha_n-1} \det[\mu_j^{\alpha_k}]_{j,k=1,\dots,n-1} = \det[\mu_j^{\alpha_k - \alpha_n - 1}]_{j,k=1,\dots,n-1}. \tag{2.12}$$

Since the dependence on μ_j is entirely in row j , the integration over $\lambda_{j+1} > \mu_j > \lambda_j$ can be done row-by-row. Furthermore, by adding row $1, \dots, j - 1$ to row j the integration can be taken to be over $\lambda_{j+1} > \mu_j > \lambda_1$. Applying this operation to (2.12) gives

$$\prod_{k=1}^{n-1} \frac{1}{\alpha_k - \alpha_n} \det[\lambda_{j+1}^{\alpha_k - \alpha_n} - \lambda_1^{\alpha_k - \alpha_n}]_{j,k=1,\dots,n-1}.$$

But

$$\det[\lambda_{j+1}^{\alpha_k - \alpha_n} - \lambda_1^{\alpha_k - \alpha_n}]_{j,k=1,\dots,n-1} = \det[\lambda_j^{\alpha_k - \alpha_n}]_{j,k=1,\dots,n},$$

as can be seen by subtracting the first row from each of the next rows in the determinant on the RHS, then expanding by the final column to obtain the LHS. Thus (2.11) now follows. \square

Remark 2.7. Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_N)$ denote a partition [50, Chap. 7]. The Schur polynomial can be defined by

$$s_\lambda(x_1, \dots, x_N) = \frac{\det[x_j^{\lambda_{N-k+1}+k-1}]_{j,k=1,\dots,N}}{\prod_{1 \leq j < k \leq N} (x_k - x_j)}, \tag{2.13}$$

which in fact is well defined for any N -array λ . With $N = n - 1$, set $\lambda_{N-k+1} + k = \alpha_k - \alpha_n$ and define $\tilde{\alpha}^{(n-1)} = (\alpha_{n-1} - \alpha_n - n - 1, \alpha_{n-2} - \alpha_n - n + 2, \dots, \alpha_1 - \alpha_n + 1)$. Substituting (2.13) in (2.11) gives

$$\begin{aligned} & \int_{R_{\leq \lambda}} \prod_{1 \leq j < k \leq n-1} (\mu_k - \mu_j) s_{\tilde{\alpha}^{(n-1)}}(\mu_1, \dots, \mu_{n-1}) d\mu_1 \cdots d\mu_{n-1} \\ &= \prod_{k=1}^{n-1} \frac{1}{\alpha_k - \alpha_n} \prod_{1 \leq j < k \leq n} (\lambda_k - \lambda_j) s_{\tilde{\alpha}^{(n)}}(\lambda_1, \dots, \lambda_n), \end{aligned} \tag{2.14}$$

where $\tilde{\alpha}^{(n)}$ is the n -array formed from the $(n - 1)$ -array $\tilde{\alpha}^{(n-1)}$ by appending 0 to the end. This is a special case of an integration formula from the theory of Jack polynomials [46, 36, 32]; see also [23, Eq. (12.210)].

Corollary 2.8. *Let the $N \times N$ random matrix Y be in the upper-triangular ensemble with diagonal entries specified by (2.2).*

(a) [17] *For all $n = 1, \dots, N$, the eigenvalue PDF of $Y_{N \times n}^\dagger Y_{N \times n} = Y_{n \times n}^\dagger Y_{n \times n}$, denoted by $\lambda^{(n)} = \{\lambda_1^{(n)}, \dots, \lambda_n^{(n)}\}$ in descendent order, is equal to*

$$p_n(\lambda_1^{(n)}, \dots, \lambda_n^{(n)}) = \frac{C_n^{-1}}{\prod_{1 \leq j < k \leq n} (\alpha_j - \alpha_k)} \left(\prod_{k=1}^n e^{-\lambda_k^{(n)}} \right) \prod_{1 \leq j < k \leq n} (\lambda_j^{(n)} - \lambda_k^{(n)}) \det[(\lambda_j^{(n)})^{\alpha_k}]_{j,k=1, \dots, n}, \tag{2.15}$$

where

$$C_n = \prod_{l=1}^n \Gamma(\alpha_l + 1). \tag{2.16}$$

In the special case the α_j are given by (2.3) this reduces to

$$p_n(\lambda_1^{(n)}, \dots, \lambda_n^{(n)}) = \frac{1}{C_{n,\theta,c}} \prod_{k=1}^n (\lambda_k^{(n)})^c e^{-\lambda_k^{(n)}} \prod_{1 \leq j < k \leq n} (\lambda_j^{(n)} - \lambda_k^{(n)}) ((\lambda_j^{(n)})^\theta - (\lambda_k^{(n)})^\theta), \tag{2.17}$$

where

$$C_{n,\theta,c} = \prod_{l=1}^n \Gamma(\theta(l - 1) + c + 1) \theta^{n(n-1)/2} \prod_{l=1}^{n-1} l!. \tag{2.18}$$

(b) *The joint probability density function of $\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(N)}$ is equal to*

$$p(\lambda^{(1)}, \dots, \lambda^{(n)}) = \frac{1}{C_N} \prod_{1 \leq j < k \leq N} (\lambda_j^{(N)} - \lambda_k^{(N)}) \prod_{n=1}^N (\lambda_n^{(N)})^{\alpha_N} e^{-\lambda_n^{(N)}} \times \prod_{n=1}^{N-1} \left(\prod_{i=1}^n (\lambda_i^{(n)})^{\alpha_n - \alpha_{n+1} - 1} \right) 1_{\lambda^{(n)} \preceq \lambda^{(n+1)}}, \tag{2.19}$$

where C_n is defined in (2.16), and $1_{\mu \preceq \lambda}$ is the indicator function of the region that satisfies inequality (2.5).

In case that some α_i are identical, we understand the formulas in the limiting sense with l’Hôpital’s rule.

Proof of Part (a). We prove the case that α_i are distinct, and the general result follows by analytical continuation.

In the case $n = 1$ we see from (2.2) that the PDF of the unique eigenvalue $\lambda_1^{(1)}$ is equal to $\Gamma(\alpha_1 + 1)^{-1} (\lambda_1^{(1)})^{\alpha_1} e^{-\lambda_1^{(1)}}$, which is the $n = 1$ case of (2.15). Let us now assume that (2.15) is valid in the case $n - 1$. Next we prove the $n > 1$ cases by induction. For notational simplicity, we denote $\lambda_i^{(n)}$ by λ_i and $\lambda_i^{(n-1)}$ by μ_i . Recall the conditional PDF $p_{n,n-1}(\{\lambda_j\}_{j=1}^n, \{\mu_j\}_{j=1}^{n-1})$ of $\{\lambda_1, \dots, \lambda_n\}$ with fixed $\{\mu_1, \dots, \mu_{n-1}\}$, such that the interlacing condition (2.5) is satisfied, defined in (2.4). Our task is to show that with $R_{\preceq \lambda}$ the domain of $\{\mu_1, \dots, \mu_{n-1}\}$ given by (2.5),

$$\int_{R_{\preceq \lambda}} p_{n,n-1}(\{\lambda_j\}_{j=1}^n, \{\mu_j\}_{j=1}^{n-1}) p_{n-1}(\mu_1, \dots, \mu_{n-1}) d\mu_1 \cdots d\mu_{n-1} = p_n(\lambda_1, \dots, \lambda_n). \tag{2.20}$$

Substituting for the integrand, then making use of Lemma 2.6 shows that the LHS is equal to

$$\frac{1}{C_{n-1}} \frac{1}{\Gamma(\alpha_n + 1)} \prod_{1 \leq j < k \leq n} \frac{1}{\alpha_j - \alpha_k} \prod_{k=1}^n e^{-\lambda_k} \prod_{1 \leq j < k \leq n} (\lambda_j - \lambda_k) \det[\lambda_j^{\alpha_k}]_{j,k=1,\dots,n}.$$

Comparison with (2.15) and recalling (2.16) shows that this is precisely the RHS.

To deduce (2.17) from (2.15) we make use of the Vandermonde determinant identity

$$\prod_{1 \leq j < k \leq n} (x_k - x_j) = \det[x_j^{k-1}]_{j,k=1,\dots,n}. \quad \square$$

Proof of Part (b). By Lemma 2.5, we have that the eigenvalues $\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(N)}$ constitute an inhomogeneous Markov chain with the transition probability density function from time $n - 1$ to time n given by (2.4). Thus the joint distribution function of $\lambda^{(n)}$ ($n = 1, \dots, N$) is obtained by multiplying (2.4) repeatedly. The argument is the same as the proof of [2, Cor. 1] and we omit the details. \square

Remark 2.9. In the special case $\theta = 0$ of (2.3) we have from (2.2) that $2|y_{k,k}|^2 \stackrel{d}{=} \chi_{2(c+1)}^2$ independent of k . Taking the limit $\theta \rightarrow 0^+$ in (2.17), we have that the corresponding eigenvalue PDF is equal to [17]

$$\frac{1}{\Gamma(c + 1)^n \prod_{l=1}^{n-1} l!} \prod_{k=1}^n \lambda_k^c e^{-\lambda_k} \prod_{1 \leq j < k \leq n} (\lambda_j - \lambda_k)(\log \lambda_j - \log \lambda_k). \quad (2.21)$$

From the joint probability distribution function (2.19), we have, as a natural generalisation of [2, Thm. 3(c)]:

Proposition 2.10. For matrices in the the upper-triangular ensemble, let the eigenvalues $\lambda^{(1)}, \dots, \lambda^{(N)}$ be defined as in Corollary 2.8. The eigenvalues $\lambda^{(1)}, \dots, \lambda^{(N)}$ constitute a determinantal process, and the correlation kernel of $\lambda^{(n_1)}$ and $\lambda^{(n_2)}$ is given by

$$K(n_1, x; n_2, y) = \frac{-1}{2\pi i} \oint_{\Gamma_\alpha} \frac{x^{-w-1} y^w}{\prod_{l=n_1+1}^{n_2} (w - \alpha_l)} dw 1_{x < y} 1_{n_1 < n_2} + \frac{1}{(2\pi i)^2} \oint_{\Sigma} dz \oint_{\Gamma_\alpha} dw \frac{x^{-z-1} y^w \Gamma(z + 1) \prod_{k=1}^{n_1} (z - \alpha_k)}{(z - w) \Gamma(w + 1) \prod_{l=1}^{n_2} (w - \alpha_l)}, \quad (2.22)$$

where Γ_α is a contour enclosing $\alpha_1, \dots, \alpha_N$, while Σ is a Hankel like contour, starting at $-\infty - i\epsilon$, running parallel to the negative real axis, looping around the point $z = -1$ and the contour Γ_α , and finishing at $-\infty + i\epsilon$ after again running parallel to the negative real axis, see Figure 1.




Figure 1: Schematic figures of Γ_α and Σ .

The proof of the proposition is by a standard argument for determinantal processes based on the joint probability density function (2.19). In [2, Thm. 3(c)], the proposition for non-negative integer α_i under condition (2.1) is proved. Since the proof does not use these additional conditions, it is also a complete proof to Proposition 2.10.

2.2 The Jacobi upper-triangular ensemble

Adler, van Moerbeke and Wang [2] considered the joint eigenvalue PDF of a sequence of random matrices

$$\begin{aligned} \tilde{J}_n &= (X_{M \times n}^\dagger X_{M \times n})(A_{M' \times n}^\dagger A_{M' \times n} + X_{M \times n}^\dagger X_{M \times n})^{-1} \\ &= (A_{M' \times n}^\dagger A_{M' \times n}(X_{M \times n}^\dagger X_{M \times n})^{-1} + \mathbb{I}_n)^{-1}, \quad n = 1, \dots, N \end{aligned} \tag{2.23}$$

where $X_{M \times n}$ is the left $M \times n$ sub-block of the $M \times N$ matrix X specified by (2.2), while $A_{M' \times n}$ is the top $M' \times n$ sub-block of the $M' \times N$ matrix A which has all elements independently distributed as standard complex Gaussians. Here it is required that $M \geq n$ and $M' \geq n$. Due to its relationship to particular multiple orthogonal polynomials, this was referred to as the Jacobi-Piñeiro ensemble.

We consider in this section a Jacobi-type counterpart of the random matrix ensembles $Y_{N \times n}^\dagger Y_{N \times n}$ in Section 2.1. To this end, we use the $N \times N$ random matrix Y specified in Section 2.1 above Proposition 2.1, and denote the $N \times N$ random matrix Z , which is also in the same upper-triangular ensemble as Y . Let $Z = [z_{j,k}]_{j,k=1,\dots,N}$ be an upper-triangular $N \times N$ random matrix with all upper-triangular entries independent, the diagonal entries be real positive with distributions depending on parameters $\beta_k > -1$

$$|z_{k,k}|^2 \stackrel{d}{=} \Gamma[\beta_k + 1, 1], \quad \text{or equivalently} \quad 2|z_{k,k}|^2 \stackrel{d}{=} \chi_{2(\beta_k+1)}^2, \quad (k = 1, \dots, N), \tag{2.24}$$

and all entries strictly above the diagonal be complex and in standard complex Gaussian distribution. Then define for all $n = 1, \dots, N$

$$\begin{aligned} J_n &= (Y_{N \times n}^\dagger Y_{N \times n})(Z_{N \times n}^\dagger Z_{N \times n} + Y_{N \times n}^\dagger Y_{N \times n})^{-1} \\ &= (Z_{N \times n}^\dagger Z_{N \times n}(Y_{N \times n}^\dagger Y_{N \times n})^{-1} + \mathbb{I}_n)^{-1}, \end{aligned} \tag{2.25}$$

where $Y_{N \times n}$ (resp. $Z_{N \times n}$) is the left $N \times n$ sub-block of the $N \times N$ matrix Y (resp. Z). Note that both \tilde{J}_n and J_n depend on parameters $\alpha_1, \dots, \alpha_n > -1$, J_n also depends on $\beta_1, \dots, \beta_n > -1$ and for \tilde{J}_n it is further assumed that $\alpha_1, \dots, \alpha_n$ are non-negative integers satisfying inequality (2.1). We say that the matrices J_n form the Jacobi upper-triangular ensemble.

Parallel to Proposition 2.1, we have

Proposition 2.11. Fix nonnegative integers $\alpha_1, \dots, \alpha_N$ such that (2.1) is satisfied and fix $\beta_i = M' - i$ ($i = 1, \dots, N$), and consider J_n and \tilde{J}_n defined by (2.25) and (2.23) respectively with the same parameters $\alpha_1, \dots, \alpha_n$. Then the joint distribution of $\tilde{J}_1, \tilde{J}_2, \dots, \tilde{J}_N$ is the same as the joint distribution of J_1, J_2, \dots, J_N . In particular, the joint distribution of the eigenvalues of $\tilde{J}_1, \dots, \tilde{J}_n$ is the same as the joint distribution of the eigenvalues of J_1, \dots, J_n .

Proof. The proof is analogous to that of Proposition 2.1, and we divide it into two steps. As a bridge between \tilde{J}_n and J_n , for all $n = 1, \dots, N$ we define

$$\hat{J}_n = (Y_{N \times n}^\dagger Y_{N \times n})(A_{M' \times n}^\dagger A_{M' \times n} + Y_{N \times n}^\dagger Y_{N \times n})^{-1}. \tag{2.26}$$

Recall the sequence of random matrices $X^{(0)} = X, X^{(1)}, \dots, X^{(N)}$ and random Householder matrices $U^{(1)}, \dots, U^{(N)}$ constructed in the proof of Proposition 2.1. We have that for all $n = 1, \dots, N$, $(X_{M \times n}^{(N)})^\dagger X_{M \times n}^{(N)}$ are independent of A , and they have the same joint distribution as $Y_{N \times n}^\dagger Y_{N \times n}$. So the joint distribution of \hat{J}_n is the same as that of

$$\tilde{J}_n^{(N)} = ((X_{M \times n}^{(N)})^\dagger X_{M \times n}^{(N)})(A_{M' \times n}^\dagger A_{M' \times n} + (X_{M \times n}^{(N)})^\dagger X_{M \times n}^{(N)})^{-1}.$$

On the other hand, $X^{(N)}$ is a function of X and $(X_{M \times n}^{(N)})^\dagger X_{M \times n}^{(N)} = X_{M \times n}^\dagger X_{M \times n}$, so $\tilde{J}_n^{(N)}$ is identical to \tilde{J}_n given that X and A are the same. Thus we have showed that the joint distribution of \tilde{J}_n is the same as that of \hat{J}_n .

Next, since A is also in the upper-triangular ensemble, we use the same algorithm in the proof of Proposition 2.1 to construct a sequence of random matrices $A^{(0)} = A, A^{(1)}, \dots, A^{(N)}$ and random Householder matrices $V^{(1)}, \dots, V^{(N)}$, such that $A^{(n+1)} = V^{(n+1)}A^{(n)}$ and $A^{(n)}$ satisfy the same conditions as $X^{(n)}$ but with the parameters $M, \alpha_1, \dots, \alpha_N$ replaced by $M', \beta_1 = M' - 1, \dots, \beta_N = M' - N$. We have that for all $n = 1, \dots, N$, each $(A_{M' \times n}^{(N)})^\dagger A_{M' \times n}^{(N)}$ is independent of Y , and they have the same joint distribution as $Z_{N \times n}^\dagger Z_{N \times n}$. So the joint distribution of J_n is the same as that of

$$\hat{J}_n^{(N)} = (Y_{N \times n}^\dagger Y_{N \times n})((A_{N \times n}^{(N)})^\dagger A_{N \times n}^{(N)} + Y_{N \times n}^\dagger Y_{N \times n})^{-1}.$$

On the other hand, $A^{(N)}$ is a function of A and $(A_{M' \times n}^{(N)})^\dagger A_{M' \times n}^{(N)} = A_{M' \times n}^\dagger A_{M' \times n}$. So $\hat{J}_n^{(N)}$ is identical to \hat{J}_n given that Y and A are identical. Thus we show that the joint distribution of \hat{J}_n is the same as that of J_n . \square

The following proposition is the Jacobi counterpart of Proposition 2.4, and its proof is analogous to that of the Jacobi–Piñeiro part of [2, Thm. 1].

Proposition 2.12. Let the $N \times N$ random matrices Y and Z be in the upper-triangular ensemble with diagonal entries specified by (2.2) and (2.24), and thus with parameters $\alpha_1, \dots, \alpha_N$ and β_1, \dots, β_N respectively. Denote by $\{\lambda_1, \dots, \lambda_n\}$ and $\{\mu_1, \dots, \mu_{n-1}\}$ the eigenvalues of J_n and J_{n-1} respectively in descending order, where J_n and J_{n-1} are defined in (2.25). The conditional PDF of $\{\lambda_1, \dots, \lambda_n\}$, with $\{\mu_1, \dots, \mu_{n-1}\}$ fixed and distinct, is equal to

$$p_{n,n-1}(\{\lambda_j\}_{j=1}^n, \{\mu_j\}_{j=1}^{n-1}) = \frac{\Gamma(\alpha_n + \beta_n + n + 1)}{\Gamma(\alpha_n + 1)\Gamma(\beta_n + 1)} \times \prod_{k=1}^n \lambda_k^{\alpha_n} (1 - \lambda_k)^{\beta_n} \prod_{l=1}^{n-1} \mu_l^{-\alpha_n - 1} (1 - \mu_l)^{-\beta_n - 1} \frac{\prod_{1 \leq j < k \leq n} (\lambda_j - \lambda_k)}{\prod_{1 \leq j < k \leq n-1} (\mu_j - \mu_k)}, \quad (2.27)$$

subject to the interlacing constraint

$$1 \geq \lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \geq \lambda_n \geq 0. \quad (2.28)$$

Analogous to Proposition 2.4, we actually prove a slightly stronger result:

Lemma 2.13. Let $C_{n-1} = (c_{i,j})$ and $B_{n-1} = (b_{i,j})$ be $(n-1) \times (n-1)$ invertible matrices such that the eigenvalues of $(C_{n-1}^\dagger C_{n-1})(B_{n-1}^\dagger B_{n-1} + C_{n-1}^\dagger C_{n-1})^{-1}$ are $\{\mu_1, \dots, \mu_{n-1}\}$ in descending order. Define the $n \times n$ matrix C_n (resp. B_n) by letting: (1) the $(n-1) \times (n-1)$ upper-triangular block equal C_{n-1} (resp. B_{n-1}); (2) the bottom row has all entries but the rightmost one equal 0; (3) all entries of the rightmost column be independent random variables, such that the (n, n) -entry $c_{n,n}$ (resp. $b_{n,n}$) is real positive with $|c_{n,n}|^2 \stackrel{d}{=} \Gamma[\alpha_n + 1, 1]$ (resp. $|b_{n,n}|^2 \stackrel{d}{=} \Gamma[\beta_n + 1, 1]$), and all other entries in the column are in standard complex normal distribution. Then the eigenvalues of $(C_n^\dagger C_n)(B_n^\dagger B_n + C_n^\dagger C_n)^{-1}$, denoted by $\{\lambda_1, \dots, \lambda_n\}$ in descending order, satisfy the interlacing constraint (2.28) and have distribution given by $p_{n,n-1}(\{\lambda_j\}_{j=1}^n, \{\mu_j\}_{j=1}^{n-1})$ in (2.27).

Proof of Lemma 2.13. It is more convenient to consider for $* = n$ or $n - 1$,

$$T_* = (B_*^\dagger B_*)^{-1} C_*^\dagger C_* \quad \text{and} \quad R_* = (C_* B_*^{-1})^\dagger (C_* B_*^{-1}),$$

and their eigenvalues $\{\tilde{\lambda}_1, \dots, \tilde{\lambda}_n\}$ for T_n and R_n , and $\{\tilde{\mu}_1, \dots, \tilde{\mu}_{n-1}\}$ for T_{n-1} and R_{n-1} , assumed in ascending order, noting that T_* and R_* have the same eigenvalues. The relations between $\tilde{\lambda}_i, \tilde{\mu}_i$ and λ_i, μ_i are

$$\tilde{\lambda}_i = \frac{\lambda_i}{1 - \lambda_i}, \quad \tilde{\mu}_i = \frac{\mu_i}{1 - \mu_i}, \quad \text{if } \lambda_i \neq 1 \text{ and } \mu_i \neq 1.$$

It is straightforward to check that the proposition is equivalent to the statement that the conditional PDF of $\{\tilde{\lambda}_1, \dots, \tilde{\lambda}_n\}$ is equal to

$$\begin{aligned} \tilde{p}_{n,n-1}(\{\tilde{\lambda}_j\}_{j=1}^n, \{\tilde{\mu}_j\}_{j=1}^{n-1}) &= \frac{\Gamma(\alpha_n + \beta_n + n + 1)}{\Gamma(\alpha_n + 1)\Gamma(\beta_n + 1)} \\ &\times \prod_{k=1}^n \tilde{\lambda}_k^{\alpha_n} (1 + \tilde{\lambda}_k)^{-(\alpha_n + \beta_n + n + 1)} \prod_{l=1}^{n-1} \tilde{\mu}_l^{-\alpha_n - 1} (1 + \tilde{\mu}_l)^{\alpha_n + \beta_n + n} \frac{\prod_{1 \leq j < k \leq n} (\lambda_j - \lambda_k)}{\prod_{1 \leq j < k \leq n-1} (\mu_j - \mu_k)}, \end{aligned} \tag{2.29}$$

subject to the interlacing constraint

$$0 < \tilde{\lambda}_1 \leq \tilde{\mu}_1 \leq \tilde{\lambda}_2 \leq \tilde{\mu}_2 \leq \dots \leq \tilde{\lambda}_n. \tag{2.30}$$

For notational simplicity, we denote $\vec{y} = (c_{1,n}, c_{2,n}, \dots, c_{n-1,n})^T$, $\vec{z} = (b_{1,n}, b_{2,n}, \dots, b_{n-1,n})^T$, $\eta = c_{n,n}$ and $\zeta = b_{n,n}$. Then

$$R_n = Q^\dagger Q, \quad \text{where } Q = \begin{bmatrix} C_{n-1} B_{n-1}^{-1} & \zeta^{-1}(\vec{y} - C_{n-1} B_{n-1}^{-1} \vec{z}) \\ \mathbf{0}_{1 \times (n-1)} & \eta \zeta^{-1} \end{bmatrix}.$$

Taking the singular value decomposition to $C_{n-1} B_{n-1}^{-1}$, we have $(n - 1)$ -dimensional unitary matrices U, V such that

$$C_{n-1} B_{n-1}^{-1} = U D V^{-1}, \quad \text{where } D = \text{diag}[\sqrt{\tilde{\mu}_1}, \dots, \sqrt{\tilde{\mu}_{n-1}}].$$

Introducing

$$U \oplus \mathbb{I}_1 = \begin{bmatrix} U & \mathbf{0}_{(n-1) \times 1} \\ \mathbf{0}_{1 \times (n-1)} & 1 \end{bmatrix}, \quad V \oplus \mathbb{I}_1 = \begin{bmatrix} V & \mathbf{0}_{(n-1) \times 1} \\ \mathbf{0}_{1 \times (n-1)} & 1 \end{bmatrix},$$

we have

$$Q = (U \oplus \mathbb{I}_1) \begin{bmatrix} D & \zeta^{-1} \vec{w} \\ \mathbf{0}_{1 \times (n-1)} & \eta \zeta^{-1} \end{bmatrix} (V \oplus \mathbb{I}_1)^{-1}, \quad \text{where } \vec{w} = U^{-1} \vec{y} - D V^{-1} \vec{z}.$$

Analogous to (2.10), the eigenvalue equation for the matrix $R_n = Q^\dagger Q$ is given by

$$0 = \lambda + \left(-\frac{\eta^2}{\zeta^2} + \sum_{k=1}^{n-1} \frac{|w_k|^2}{\zeta^2} \frac{\lambda}{\tilde{\mu}_k - \lambda} \right), \tag{2.31}$$

where the w_k are components of \vec{w} . Note that $|w_1|^2, \dots, |w_{n-1}|^2, \eta^2, \zeta^2$ are independent, and their distribution functions are positive with densities

$$|w_k|^2 \stackrel{d}{=} \frac{1}{1 + \tilde{\mu}_k} e^{-\frac{t}{1 + \tilde{\mu}_k}}, \quad \eta^2 \stackrel{d}{=} \frac{1}{\Gamma(\alpha_n + 1)} t^{\alpha_n} e^{-t}, \quad \zeta^2 \stackrel{d}{=} \frac{1}{\Gamma(\beta_n + 1)} t^{\beta_n} e^{-t}.$$

Comparing (2.31) with [2, Eq. (131)], and using the calculations in [2, Eq. (139)–(141)], we prove (2.29). (In [2, Eqs. (139)–(141)] the calculations are done for integer valued α_n and β_n which are denoted as $M' - n$. But the method works also for real-valued α_n and β_n .) \square

This result can be used to derive the analogue of Corollary 2.8 in the Jacobi case.

Corollary 2.14. *Let $\alpha_1, \dots, \alpha_N$ be arbitrary real numbers greater than -1 and*

$$\beta_k = \beta + N - k \quad \text{for } k = 1, \dots, N \text{ where } \beta > -1. \quad (2.32)$$

Let J_n be defined in (2.25) for all $n = 1, \dots, N$.

(a) *Then the eigenvalues of the random matrix J_n , denoted by $\lambda^{(n)} = \{\lambda_1^{(n)}, \dots, \lambda_n^{(n)}\}$ in descending order, is equal to*

$$p_n(\lambda_1^{(n)}, \dots, \lambda_n^{(n)}) = \frac{C_n^{-1}}{\prod_{1 \leq j < k \leq n} (\alpha_j - \alpha_k)} \times \prod_{k=1}^n (1 - \lambda_k^{(n)})^{\beta_n} \prod_{1 \leq j < k \leq n} (\lambda_j^{(n)} - \lambda_k^{(n)}) \det[(\lambda_j^{(n)})^{\alpha_k}]_{j,k=1, \dots, n}, \quad (2.33)$$

where

$$C_n = \prod_{l=1}^n \frac{\Gamma(\alpha_l + 1)\Gamma(\beta_l + 1)}{\Gamma(\alpha_l + \beta + N + 1)}. \quad (2.34)$$

In the special case that α_j is given by (2.3) this reduces to

$$p_n(\lambda_1^{(n)}, \dots, \lambda_n^{(n)}) = \frac{1}{C_{n,\theta,c}} \prod_{k=1}^n (\lambda_k^{(n)})^c (1 - \lambda_k^{(n)})^{\beta_n} \prod_{1 \leq j < k \leq n} (\lambda_j^{(n)} - \lambda_k^{(n)}) ((\lambda_j^{(n)})^\theta - (\lambda_k^{(n)})^\theta), \quad (2.35)$$

where

$$C_{n,\theta,c} = \prod_{l=1}^n \frac{\Gamma(\theta(l-1) + c + 1)\Gamma(\beta_l + 1)}{\Gamma(\theta(l-1) + c + \beta + N + 1)} \theta^{n(n-1)/2} \prod_{l=1}^{n-1} l!. \quad (2.36)$$

(b) *The joint probability distribution function of $\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(N)}$ is equal to*

$$p(\lambda^{(1)}, \dots, \lambda^{(N)}) = \frac{1}{C_N} \prod_{1 \leq j < k \leq N} (\lambda_j^{(N)} - \lambda_k^{(N)}) \prod_{n=1}^N (\lambda_n^{(N)})^{\alpha_N} (1 - \lambda_n^{(N)})^\beta \times \prod_{n=1}^N \left(\prod_{i=1}^n (\lambda_i^{(n)})^{\alpha_n - \alpha_{n+1} - 1} \right) 1_{\lambda^{(n)} \preceq \lambda^{(n+1)}}, \quad (2.37)$$

where C_n is defined in (2.34), and $1_{\mu \preceq \lambda}$ is the indicator function of the region that satisfies inequality (2.30).

In case that some α_i are identical, we understand the formulas in the limiting sense with l'Hôpital's rule.

Proof of Part (a). We prove the case that the α_i are distinct, and the general result follows by analytical continuation.

In the case $n = 1$, J_1 defined in (2.25) is equal in distribution to $\chi_{2(\alpha_1+1)}^2 / (\chi_{2(\beta_1+1)}^2 + \chi_{2(\alpha_1+1)}^2)$. It is a classical result [30, Sec. 25.2] that this combination of random variables is distributed according to the beta distribution $B[\alpha_1 + 1, \beta_1 + 1]$ and thus has for its PDF

$$\frac{\Gamma(\alpha_1 + \beta_1 + 2)}{\Gamma(\alpha_1 + 1)\Gamma(\beta_1 + 1)} x^{\alpha_1} (1 - x)^{\beta_1},$$

in agreement with (2.33) with $n = 1$. Proceeding by induction, let us now assume that (2.33) is valid in the case $n - 1$. For notational simplicity, we denote $\lambda_i^{(n)}$ by λ_i and $\lambda_i^{(n-1)}$ by μ_i . Our task is to check the validity of

$$\int_{R_{\leq \lambda}^J} p_{n,n-1}(\{\lambda_j\}_{j=1}^n, \{\mu_j\}_{j=1}^{n-1}) p_{n-1}(\mu_1, \dots, \mu_{n-1}) d\mu_1 \cdots d\mu_{n-1} = p_n(\lambda_1, \dots, \lambda_n),$$

which is analogous to (2.20), but the integral domain $R_{\leq \lambda}^J$ for $\{\mu_1, \dots, \mu_{n-1}\}$ is defined by (2.28), and the p_n , p_{n-1} and $p_{n,n-1}$ are defined differently. Recall the conditional PDF $p_{n,n-1}(\{\lambda_j\}_{j=1}^n, \{\mu_j\}_{j=1}^{n-1})$ of $\{\lambda_1, \dots, \lambda_n\}$ with fixed $\{\mu_1, \dots, \mu_{n-1}\}$, such that the interlacing condition (2.28) is satisfied, defined in (2.27). Substituting for the integrand, then making use of Lemma 2.6 shows that the LHS is equal to

$$\frac{1}{C_{n-1}} \frac{\Gamma(\alpha_n + \beta_n + n + 1)}{\Gamma(\alpha_n)\Gamma(\beta_n + 1)} \times \prod_{1 \leq j < k \leq n} \frac{1}{\alpha_j - \alpha_k} \prod_{k=1}^n (1 - \lambda_k)^{\beta_n} \prod_{1 \leq j < k \leq n} (\lambda_j - \lambda_k) \det[\lambda_j^{\alpha_k}]_{j,k=1,\dots,n}.$$

Comparison with (2.33) and (2.34) shows that this is precisely the RHS.

We deduce (2.35) from (2.33) in the same way as we deduced (2.17) from (2.15). \square

Proof of Part (b). By Lemma 2.13, we have that the eigenvalues $\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(N)}$ constitute an inhomogeneous Markov chain with the transition probability density function from time $n - 1$ to time n given by (2.27). Thus the joint distribution function of $\lambda^{(n)}$ ($n = 1, \dots, N$) is obtained by multiplying (2.27) repeatedly. The argument is the same as the proof of [2, Cor. 1] and we omit the details. \square

Remark 2.15. The assumption (2.32) that β_k are in arithmetic progression with common difference -1 is crucial in the application of Lemma 2.6. By the symmetry of the model, it is also possible to let β_k be arbitrary and α_k in arithmetic progression with common difference -1 .

From the joint probability distribution function (2.37), we have, as a natural generalisation of [2, Thm. 3(d)]:

Proposition 2.16. Let the matrices J_n and the eigenvalues $\lambda^{(1)}, \dots, \lambda^{(N)}$ be defined as in Corollary 2.14. The eigenvalues $\lambda^{(1)}, \dots, \lambda^{(N)}$ constitute a determinantal process, and the correlation kernel of $\lambda^{(n_1)}$ and $\lambda^{(n_2)}$ is given by

$$K(n_1, x; n_2, y) = \frac{-1}{2\pi i} \oint_{\Gamma_\alpha} \frac{x^{-w-1} y^w}{\prod_{l=n_1+1}^{n_2} (w - \alpha_l)} dw 1_{x < y} 1_{n_1 < n_2} + \frac{1}{(2\pi i)^2} \oint_{\Sigma} dz \oint_{\Gamma_\alpha} dw \frac{x^{-z-1} y^w}{(z - w)} \frac{\Gamma(w + \beta + N + 1)\Gamma(z + 1)}{\Gamma(z + \beta + N + 1)\Gamma(w + 1)} \frac{\prod_{k=1}^{n_1} (z - \alpha_k)}{\prod_{l=1}^{n_2} (w - \alpha_l)}, \quad (2.38)$$

where

1. if $\beta \in \mathbb{Z}$, Γ_σ is a positively oriented contour enclosing $\alpha_1, \dots, \alpha_N$, while Σ is a contour going counterclockwise enclosing $-1, -2, \dots, -(\beta + N)$ and the contour Γ_α , and
2. if $\beta \notin \mathbb{Z}$, Γ_α is a Hankel like contour, starting at $-\infty - i\epsilon$, running parallel to the negative real axis, enclosing the poles $w = -(\beta + N + k)$ with $k \in \mathbb{Z}_+$ and $w = \alpha_1, \dots, \alpha_N$, and finishing at $-\infty + i\epsilon$ after again running parallel to the negative real axis, while Σ is a Hankel like contour that loops around Γ_α .

The proof of the proposition is by a standard argument of determinantal process based on the joint probability density function (2.37). In [2, Thm. 3(d)], the proposition for non-negative integer β and non-negative integer α_i under condition (2.1) is proved. Since the proof does not use these additional conditions, it is also a complete proof to Proposition 2.16. Note that in [2], only case (1) of the contours occurs.

It is possible to use Corollary 2.8(a) to give an alternative derivation of Corollary 2.14(a), in the case that if β_1, \dots, β_N satisfies (2.32) with $\beta \in \mathbb{Z}^+$. In this case we note that the eigenvalue PDF of J_n is the same as the eigenvalue PDF of \hat{J}_n defined in (2.26), where the height of the random matrix A is $M' = \beta + N$. We also need a recent result due to Kuijlaars and Stivigny [34]. Below we give the derivation without the tedious calculation of the normalisation constant of the PDF.

Proposition 2.17 (Special case of [34, Thm. 2.1]). Let the matrix W be an $n \times n$ random matrix such that $W^\dagger W$ has an eigenvalue PDF proportional to

$$\prod_{1 \leq j < k \leq n} (x_k - x_j) \det[f_{k-1}(x_j)]_{j,k=1}^n \tag{2.39}$$

for some $\{f_{k-1}(x)\}_{k=1, \dots, n}$. For $\nu \geq 0$, let G be an $(n + \nu) \times n$ random matrix whose entries are in independent standard complex Gaussian distribution. The squared singular values of GW , or equivalently the eigenvalues of $(GW)^\dagger GW$, have PDF proportional to

$$\prod_{1 \leq j < k \leq n} (y_k - y_j) \det[g_{k-1}(y_j)]_{j,k=1}^n, \tag{2.40}$$

where

$$g_k(y) = \int_0^\infty x^\nu e^{-x} f_k\left(\frac{y}{x}\right) \frac{dx}{x}, \quad (k = 0, \dots, n - 1). \tag{2.41}$$

In the application of Proposition 2.17, we let $W = Y_{n \times n}^{-1}$ and $G = A_{M' \times n}$, such that $\nu = M' - n = \beta_n = \beta + N - n$. In Corollary 2.8 we obtained the eigenvalue PDF of $Y_{n \times n}^\dagger Y_{n \times n}$. From this, by the change of variables $\lambda_j \mapsto 1/x_j$, we have that the eigenvalue PDF of $W^\dagger W = (Y_{n \times n} Y_{n \times n}^\dagger)^{-1}$ is proportional to

$$\prod_{k=1}^n x_k^{-(n+1)} e^{-1/x_k} \prod_{1 \leq j < k \leq n} (x_k - x_j) \det[x_j^{-\alpha_k}]_{j,k=1, \dots, n},$$

and so we can take

$$f_k(x) = x^{-(n+1)} e^{-1/x} x^{-\alpha_k}.$$

Then substituting this in (2.41) gives that

$$g_k(y) \propto y^\nu (1 + y)^{-(\nu + \alpha_k + n + 1)}.$$

Hence we deduce that the eigenvalue PDF of $(GW)^\dagger GW = (A_{M' \times n} Y_{n \times n}^{-1})^\dagger A_{M' \times n} Y_{n \times n}^{-1}$, which is the same as the eigenvalue PDF of

$$\hat{S}_n = (Y_{n \times n}^\dagger Y_{n \times n})^{-1} A_{M' \times n}^\dagger A_{M' \times n},$$

has its eigenvalue PDF $p_n^{\hat{S}}(\sigma_1, \dots, \sigma_n)$ proportional to

$$\prod_{l=1}^n \sigma_l^\nu (1 + \sigma_l)^{-(\nu + n + 1)} \prod_{1 \leq j < k \leq n} (\sigma_k - \sigma_j) \det[(1 + \sigma_j)^{-\alpha_k}]_{j,k=1}^n.$$

The eigenvalues $\{\lambda_j\}$ of \hat{J}_n and the eigenvalues $\{\sigma_j\}$ of \hat{S}_n are related by

$$\lambda_j = \frac{1}{\sigma_j + 1}. \tag{2.42}$$

Changing variables to $\{\lambda_j\}$ according to (2.42) gives (2.33) up to the normalisation constant C_n .

3 The global density — characteristic polynomial approach

3.1 The Laguerre Muttalib–Borodin ensemble

Recent results [34, 24] have revealed an intimate relationship between random matrix products and the Laguerre Muttalib–Borodin ensemble. To explain this requires the introduction of a family of integer sequences — the Fuss–Catalan numbers — parametrised by $s \in \mathbb{R}^+$ and specified by

$$C_s(k) = \frac{1}{sk+1} \binom{sk+k}{k}, \quad k = 0, 1, 2, \dots \quad (3.1)$$

For general $s > 0$ these are known to be moments of a PDF — the Fuss–Catalan density — with compact support $[0, L]$, $L > 0$ [7, 41], and they uniquely define the PDF.

Consider first a product of s $N \times N$ matrices X_1, \dots, X_s with each containing independent, identically distributed zero mean, unit standard deviation random variables. Alternatively, for one such matrix X_1 say, consider the power X_1^s . In either case, ask for the limiting spectral density of the squared singular values after dividing by N^s — what results is precisely the Fuss–Catalan density with parameter s [4, 7, 45].

Consider now the Laguerre Muttalib–Borodin ensemble defined by (1.1) and (1.5). Let $\lambda_1^{(N)}, \dots, \lambda_N^{(N)}$ be the eigenvalues in the N -dimensional ensemble. As $N \rightarrow \infty$, by standard techniques we have that the empirical distribution of the scaled eigenvalues $N^{-1}\lambda_1^{(N)}, \dots, N^{-1}\lambda_N^{(N)}$ converges in distribution to a limiting probability distribution, also known as the equilibrium measure of the model; see [20, Sec. 6.4]. The equilibrium measure is characterised as the minimum of a variation problem; see Claeys and Romano [18, Eq. (1.22)]. By interpreting the recent results of [18], Forrester and Liu [24] have identified the global density (i.e. density scaled by an appropriate power of N to have compact support) for the Laguerre Muttalib–Borodin ensemble in terms of the Fuss–Catalan density for general $s \geq 1$.

Proposition 3.1. Suppose $\theta \geq 1$ (this is for technical reasons in the working of [18]; in [24] it is commented that the same result is expected to hold for all $\theta > 0$ and in fact this has recently been established in [25]). After changing variables $x_k^{(N)} = (\lambda_k^{(N)}/(N\theta))^\theta$ where $\lambda_1^{(N)}, \dots, \lambda_N^{(N)}$ are the eigenvalues in the Laguerre Muttalib–Borodin ensemble defined by (1.1) and (1.5), the empirical distribution of $x_1^{(N)}, \dots, x_N^{(N)}$ converges to the Fuss–Catalan density with parameter θ as $N \rightarrow \infty$.

In particular, this shows a relationship between the product of s random matrices and the Laguerre Muttalib–Borodin ensemble with $\theta = s$ (see also [34] and the appendix in [25]). Here we will demonstrate the relationship in a different way, by considering the characteristic polynomial of the latter after the change of variables (1.4). The proof of Proposition 3.1 via Claeys and Romano’s approach is valid for all $\theta \geq 1$ at least, but depends on the construction of a mapping $J(s)$ [18, Eq. (1.25)], so it is unclear if it can be applied to the Jacobi case. On the other hand, the approach presented below is valid only for $\theta \in \mathbb{Z}^+$, but it does generalise to the Jacobi case.

Crucial for the alternative proof is knowledge of certain biorthogonal polynomials associated with (1.1) where $V(x)$ is given in (1.5). Thus for given $j = 0, 1, 2, \dots$ let $p_j(x), q_j(x)$ be monic polynomials of degree j , and suppose these polynomials have the biorthogonal property

$$\int_0^\infty e^{-V(x)} p_j(x) q_k(x^\theta) dx = h_j \delta_{j,k}, \quad h_j > 0, \quad (3.2)$$

where the positivity of h_j follows from the positivity of the integral over (1.1), because $n!h_0h_1 \cdots h_{n-1}$ is equal to the integral over (1.1). Let $\mathcal{P}_{N,\theta}$ denote the PDF specified by

(1.1). Straightforward working (see e.g. [23, Prop. 5.1.3]) shows that

$$\left\langle \prod_{l=1}^N (x - \lambda_l^{(N)}) \right\rangle_{\mathcal{P}_{N,\theta}} = p_N(x), \quad \left\langle \prod_{l=1}^N (x - (\lambda_l^{(N)})^\theta) \right\rangle_{\mathcal{P}_{N,\theta}} = q_N(x). \quad (3.3)$$

Since Proposition 3.1 requires the change of variables (1.4), we see that $q_N(x)$ is equal to the corresponding averaged characteristic polynomial. For the Laguerre weight (1.5), the explicit form of $q_N(x)$ is known from a result of Konhauser [33]. A relation between $q_N(x)$ and generalised hypergeometric functions is observed in [49]. We will use the standard notation ${}_pF_q$ to denote the generalized hypergeometric function defined by a series as presented in e.g. [39, Sec. 3].

Proposition 3.2. For the Laguerre weight (1.5), the biorthogonal polynomials $\{q_j(x)\}$ in (3.2) are given by

$$q_j(x) = (-1)^j \Gamma(\theta j + c + 1) \sum_{l=0}^j (-1)^l \binom{j}{l} \frac{x^l}{\Gamma(\theta l + c + 1)}. \quad (3.4)$$

In the case that $\theta \in \mathbb{Z}^+$, use of the duplication formula for the gamma function shows that

$$q_j(x) = C \cdot {}_1F_\theta \left(\begin{matrix} -j \\ (c+1)/\theta, (c+2)/\theta, \dots, (c+\theta)/\theta \end{matrix} \middle| \frac{x}{\theta^\theta} \right), \quad (3.5)$$

where C is independent of x and chosen so that $q_j(x)$ is monic.

Below we will make use of the standard fact (see e.g. [39, Sec. 5.1]) that the generalised hypergeometric function ${}_pF_q \left(\begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix} \middle| x \right)$ satisfies the differential equation

$$x \prod_{n=1}^p \left(x \frac{d}{dx} + a_n \right) f = x \frac{d}{dx} \prod_{n=1}^q \left(x \frac{d}{dx} + b_n - 1 \right) f. \quad (3.6)$$

Also, we require a technical result relating to the convergence of the Stieltjes transforms of the empirical distributions of $\lambda_1^{(N)}, \dots, \lambda_N^{(N)}$.

Lemma 3.3. For all z in a compact subset of $\mathbb{C} \setminus [0, \infty)$, as $N \rightarrow \infty$, uniformly

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log \left\langle \prod_{n=1}^N \left(z - \left(\frac{\lambda_n^{(N)}}{N\theta} \right)^\theta \right) \right\rangle &= \int_J \log \left(z - \left(\frac{x}{\theta} \right)^\theta \right) d\mu^L(x) \\ &= \int_{\tilde{J}} \log(z - \tilde{x}) d\tilde{\mu}^L(\tilde{x}), \end{aligned} \quad (3.7)$$

where $d\mu^L$ denotes the equilibrium measure, J is the corresponding support, $d\tilde{\mu}^L$ is the measure transformed from $d\mu^L$ by the change of variable $\tilde{x} = (x/\theta)^\theta$, and \tilde{J} is the support of $d\tilde{\mu}^L$.

The proof of Lemma 3.3 is similar to [20, Lem. 6.77]. Note that in [20] it is required that the function ϕ , corresponding to the function $\log(z - (x/\theta)^\theta)$, is a bounded function in x , while our $\log(z - (x/\theta)^\theta)$ is not. But since the growth of the function as $x \rightarrow \infty$ is mild, the argument there can be applied. Note that for z in a compact subset of $\mathbb{C} \setminus [0, \infty)$, the convergence in (3.7) is uniform, since the functions are uniformly bounded and equi-continuous. So if we take derivative on both sides of (3.7), the convergence still holds. Comparing the left-hand side of (3.7) with the formula (3.3) for q_N , we have that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \frac{d}{dz} \frac{q_N((N\theta)^\theta z)}{q_N((N\theta)^\theta z)} = \lim_{N \rightarrow \infty} \frac{(N\theta)^\theta}{N} \frac{q'_N(x)}{q_N(x)} \Big|_{x=(N\theta)^\theta z} = \tilde{G}^L(z), \quad z \in \mathbb{C} \setminus [0, \infty), \quad (3.8)$$

where

$$\tilde{G}^L(z) = \int_{\tilde{x}} \frac{d\tilde{\mu}^L(\tilde{x})}{z - \tilde{x}} \tag{3.9}$$

is the limiting resolvent (or equivalently Stieltjes transform) of the measure $d\tilde{\mu}$. Below we show that $\tilde{G}^L(z)$ satisfies a polynomial equation which uniquely characterises the Fuss–Catalan distribution.

Proposition 3.4. Transform the Laguerre Muttalib–Borodin ensemble according to (1.4). Define $\tilde{G}^L(z)$ by (3.9) so that it is equal to the resolvent corresponding to the global scaled density, scaling $x = (N\theta)^\theta z$. For $\theta \in \mathbb{Z}^+$ we have that

$$z(z\tilde{G}^L(z) - 1) = (z\tilde{G}^L(z))^{\theta+1}. \tag{3.10}$$

Proof. First we note that the convergence (3.8) and (3.9) yields that as $N \rightarrow \infty$, for $z \in \mathbb{C} \setminus [0, \infty)$,

$$q_N((N\theta)^\theta z) = \exp\left(N\left(\int \tilde{G}^L(z)dz + C + o(1)\right)\right),$$

where $o(1)$ is an analytic function in z that vanishes uniformly for z in any compact set of $\mathbb{C} \setminus [0, \infty)$, and C is a constant. The asymptotic formulas below are thus justified.

Suppose $x = \mathcal{O}(N^\theta)$ and $\arg x \in (0, 2\pi)$, then to leading order in N , after substituting for $\{a_i\}, \{b_j\}$ as implied by (3.5), we see that (3.6) reads

$$\frac{x}{\theta^\theta} \left(x \frac{d}{dx} - N\right) q_N = \left(x \frac{d}{dx}\right)^{\theta+1} q_N + o(N^{\theta+1}). \tag{3.11}$$

Now change variables $x = (N\theta)^\theta z$, and let $q_N^{(k)}(x)$ denote the k -th derivative of $q_N(x)$. Working contained in [24, above Prop. 5.1], establishes that under the validity of (3.8),

$$\frac{\frac{d^k}{dz^k} q_N((N\theta)^\theta z)}{q_N((N\theta)^\theta z)} = (N\theta)^{k\theta} \frac{q_N^{(k)}(x)}{q_N(x)} \Big|_{x=(N\theta)^\theta z} \sim N^k (\tilde{G}^L(z))^k \tag{3.12}$$

(note that (3.8) itself is the case $k = 1$; the general k case follows by expressing higher derivatives in terms of the logarithmic derivatives). Using this in (3.11) gives (3.10). \square

Remark 3.5. (i) The large z expansion

$$z\tilde{G}^L(z) = \sum_{k=0}^{\infty} \frac{\tilde{m}_k^L}{z^k}, \tag{3.13}$$

where \tilde{m}_k^L denotes the k -th moment of the global scaled spectral measure, substituted in (3.10) shows that \tilde{m}_k^L is equal to the k -th Fuss–Catalan number (3.1) with $s = \theta$. The details of the required calculation can be found in e.g. [24, paragraph beginning with eq. (2.4)].

(ii) The spectral density is the global scaled limit of the one-point correlation function. Let $\lambda_1^{(N)} = \lambda, \lambda_2^{(N)}, \dots, \lambda_N^{(N)}$ be the eigenvalues in the N -dimensional Laguerre Muttalib–Borodin ensemble. Then the one-point correlation function is (we suppress the N -dependence in the notation $\rho_{(1)}$)

$$\rho_{(1)}(\lambda) = N \int_0^\infty d\lambda_2^{(N)} \dots \int_0^\infty d\lambda_N^{(N)} p_N(\lambda, \lambda_2^{(N)}, \dots, \lambda_N^{(N)}), \tag{3.14}$$

where p_N is the PDF implied by (1.1) with the Laguerre weight (1.5). The scaled spectral density, as $N \rightarrow \infty$, converges to the equilibrium measure in distribution:

$$d\mu^L(x) = \lim_{N \rightarrow \infty} \rho_{(1)}(Nx)dx, \tag{3.15}$$

where $d\mu^L$ is the same as in (3.7).

(iii) The resolvent $\tilde{G}^L(z)$ defined in (3.9) can be expressed by the limiting spectral density/equilibrium measure of the Laguerre Muttalib–Borodin ensemble as

$$\tilde{G}^L(z) = \int_J \frac{d\mu^L(x)}{z - (x/\theta)^\theta}. \tag{3.16}$$

There is another viewpoint on deducing the global spectral density from knowledge of (3.5). This makes use of a recent result of Hardy [29], which subject to a mild technical condition [29, Eq. (1.12)] states that characteristic polynomials coming from a class of determinantal point processes including the multiple orthogonal polynomial ensemble under present discussion (see [29, Sec. 1.4]), the limiting density of zeros equals the limiting global spectral density. On the other hand, the limiting density of zeros for the generalised hypergeometric function ${}_1F_\theta$ in (3.5) has been shown by Neuschel [44] to be given by the Fuss–Catalan density with parameter $s = \theta$. (Strictly speaking the result of [44] assumes the bottom line of parameters in (3.5) to be positive integers. Since the leading asymptotics are independent of these parameters, it is expected that this assumption in [44] is not necessary.)

3.2 The Jacobi Muttalib–Borodin ensemble

As with the Laguerre weight (1.5), for the Muttalib–Borodin ensemble with Jacobi weight (1.6), we also let $\lambda_1^{(N)}, \dots, \lambda_N^{(N)}$ be the eigenvalues in the N -dimensional ensemble. As $N \rightarrow \infty$, by standard techniques, we have that the empirical distribution of the eigenvalues converges in distribution to a limiting probability distribution, also known as the equilibrium measure of this model. The equilibrium measure can be characterized as the minimum of a variation problem analogous to that for the Laguerre case, and from this, in the recent work [25] the moments of the corresponding density have been given in terms of certain binomial coefficients (see Proposition 3.10 below). Analogous to the Laguerre ensemble, there is a corresponding system of biorthogonal polynomials. For $j = 0, 1, 2, \dots$, let $p_j(x), q_j(x)$ be monic polynomials of degree j , and with weight $V(x)$ defined in (1.6) suppose these polynomials have the biorthogonal property analogous to (3.2),

$$\int_0^1 e^{-V(x)} p_j(x) q_k(x^\theta) dx = h_j \delta_{j,k}, \quad h_j > 0. \tag{3.17}$$

Let $\mathcal{P}_{N,\theta}$ denote the PDF specified by (1.1) with V specified by (1.6) and $0 < \lambda_l < 1$. Then (3.3) also holds in the Jacobi ensemble with corresponding different meanings of p_N, q_N and $\mathcal{P}_{N,\theta}$. Below we state algebraic results on the biorthogonal polynomials analogous to Proposition 3.2. Again we will focus on the polynomials $q_j(x)$.

Proposition 3.6. [16, 15, 40] For the Jacobi weight (1.6), the biorthogonal polynomials $\{q_j(x)\}$ in (3.2) are given by

$$q_j(x) = (-1)^j \frac{(1 + c_1)_{\theta j}}{(1 + c_1 + c_2 + j)_{\theta j}} \sum_{l=0}^j (-1)^l \binom{j}{l} \frac{(1 + c_1 + c_2 + j)_{\theta l}}{(1 + c_1)_{\theta l}} x^l, \tag{3.18}$$

where $(a)_p := \Gamma(a + p)/\Gamma(a)$. In the case that $\theta \in \mathbb{Z}^+$, use of the duplication formula for the gamma function shows that

$$q_j(x) \propto {}_{\theta+1}F_\theta \left(\begin{matrix} -j, \delta/\theta, (\delta + 1)/\theta, \dots, (\delta + \theta - 1)/\theta \\ (c_1 + 1)/\theta, (c_1 + 2)/\theta, \dots, (c_1 + \theta)/\theta \end{matrix} \middle| x \right), \tag{3.19}$$

where $\delta := 1 + c_1 + c_2 + j$.

Also we have the analogue of Lemma 3.3.

Lemma 3.7. For all $z \in \mathbb{C} \setminus [0, 1]$, as $N \rightarrow \infty$,

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \log \left\langle \prod_{n=1}^N \left(z - (\lambda_n^{(N)})^\theta \right) \right\rangle &= \int_{[0,1]} \log(z - x^\theta) d\mu^J(x) \\ &= \int_{[0,1]} \log(z - \tilde{x}) d\tilde{\mu}^J(\tilde{x}), \end{aligned} \tag{3.20}$$

where $d\mu^J$ denote the equilibrium measure of the Jacobi ensemble and $d\tilde{\mu}^J$ denote the measure transformed from $d\mu^J$ by the change of variable $\tilde{x} = x^\theta$. Note that both $d\mu^J$ and $d\tilde{\mu}^J$ have support $[0, 1]$.

The proof of Lemma 3.7 is similar to [20, Lem. 6.77], and we omit the details. From Lemma 3.7, we derive the counterpart of (3.8), that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \frac{q'_N(z)}{q_N(z)} = \tilde{G}^J(z), \quad \lambda \in \mathbb{C} \setminus [0, 1], \tag{3.21}$$

where

$$\tilde{G}^J(z) = \int_{[0,1]} \frac{d\tilde{\mu}^J(\tilde{x})}{z - \tilde{x}} \tag{3.22}$$

is the limiting resolvent of the measure $d\tilde{\mu}^J$. Below we show that $\tilde{G}^J(z)$ satisfies a polynomial equation that is similar to (3.10) characterising the Fuss–Catalan distribution.

Proposition 3.8. Transform the Jacobi Muttalib–Borodin ensemble according to (1.4). Define $\tilde{G}^J(z)$ by (3.22) so that it is equal to the resolvent corresponding to the global density. For $\theta \in \mathbb{Z}^+$ we have that

$$z(z\tilde{G}^J(z) - 1)(z\tilde{G}^J(z) + 1/\theta)^\theta = (z\tilde{G}^J(z))^{\theta+1}. \tag{3.23}$$

Proof. Applying the identity (3.6) to (3.19) with $j = N$, we have, to leading order in N ,

$$z \left(z \frac{d}{dz} - N \right) \left(z \frac{d}{dz} + \frac{N}{\theta} \right)^\theta q_N = \left(z \frac{d}{dz} \right)^{\theta+1} q_N, \tag{3.24}$$

while the analogue of (3.12) is

$$\frac{q_N^{(k)}(z)}{q_N(z)} \sim N^k (\tilde{G}^J(z))^k. \tag{3.25}$$

Use of (3.25) in (3.24) gives (3.23). □

Remark 3.9. (i) Taking the limit $\theta \rightarrow \infty$ in (3.23) gives the nonlinear equation

$$\left(\frac{1}{z\tilde{G}^J(z)} - 1 \right) \exp \left(\frac{1}{z\tilde{G}^J(z)} - 1 \right) = -\frac{1}{ze}. \tag{3.26}$$

By definition, the Lambert W -function $W(z)$ is the principal branch of the functional equation $z = W(z)e^{W(z)}$ defined in $\mathbb{C} \setminus (-\infty, -e^{-1})$, and thus we have in this case

$$\lim_{\theta \rightarrow \infty} \frac{1}{z\tilde{G}^J(z)} - 1 = W \left(-\frac{1}{ze} \right). \tag{3.27}$$

(ii) Generally we expect the global density associated with the weight (1.6) to be independent of c_1 and c_2 provided those parameters are themselves independent of N . On the other hand, the change of variables (1.4) shows that for finite N the density $\rho_{(1)}(x)$, defined by (3.14) with p_N the PDF implied by (1.1) with the Jacobi weight (1.6), has the functional property

$$\rho_{(1)}(x) \Big|_{c_2=0} = \theta x^{\theta-1} \rho_{(1)}(x^\theta) \Big|_{\substack{c_1 \rightarrow (c_1+1)/\theta-1, \\ c_2=0, \theta \rightarrow 1/\theta}}. \tag{3.28}$$

(iii) For $\theta = 1$ (3.23) reduces to the simple quadratic equation $(z - 1)(z\tilde{G}^J(z))^2 = z$ and thus $z\tilde{G}^J(z) = 1/(1 - 1/z)^{1/2}$, where we have imposed the requirement that $zG(z) \rightarrow 1$ as $z \rightarrow \infty$ in choosing the root of the quadratic. Recalling (3.16), this implies the well known functional form for the density in the classical Jacobi ensemble (see e.g. [23, Prop. 3.6.3])

$$d\tilde{\mu}^J(y) = d\mu^J(y) = \frac{1}{\pi} \frac{1}{\sqrt{y(1-y)}} dy, \quad 0 < y < 1. \tag{3.29}$$

A feature of (3.29) is that the moments are given in terms of a binomial coefficient,

$$\frac{1}{\pi} \int_0^1 \frac{y^p}{\sqrt{y(1-y)}} dy = \frac{1}{2^{2p}} \binom{2p}{p}. \tag{3.30}$$

In fact it is possible to show that the moments of the density implied by the appropriate solution of (3.23) are given in terms of binomial coefficients for general $\theta > 0$.

By the definition (3.22) of $\tilde{G}^J(z)$, for $|z| > 1$ we have the expansion

$$z\tilde{G}^J(z) = \sum_{p=0}^{\infty} \frac{\tilde{m}_p^J}{z^p}, \tag{3.31}$$

where \tilde{m}_p^J denotes the p -th moment of $\tilde{\mu}^J(x)$, analogous to (3.13). To compute $\{\tilde{m}_p^J\}$ we are guided by the knowledge that the moments (3.1) corresponding to the density for the Laguerre Muttalib–Borodin ensemble can be computed from the functional equation (3.10) by using the Lagrange inversion formula (see e.g. [24, Sec. 2]). The setting of the latter requires two analytic functions $f(z)$ and $\phi(z)$ in a neighbourhood Ω of a point a , and t to be small enough so that $|t\phi(z)| < |z - a|$, $z \in \Omega$. It tells us that the equation in ζ

$$\zeta = a + t\phi(\zeta) \tag{3.32}$$

has one solution in Ω and furthermore

$$f(\zeta) = f(a) + \sum_{n=1}^{\infty} \frac{t^n}{n!} \frac{d^{n-1}}{da^{n-1}} (f'(a)(\phi(a))^n). \tag{3.33}$$

Proposition 3.10. Let $\theta > 0$ and $L := (1 + \theta)^{1+\theta}\theta^{-\theta}$, and define $\tilde{G}^J(z)$ as the solution of (3.23) with the expansion (3.31) for $\theta > 0$. We have

$$\tilde{m}_p^J = L^{-p} \binom{(1 + \theta)p}{p}. \tag{3.34}$$

Proof. Let $1/z = Lt$ and $X = z\tilde{G}^J(z)$. Then by (3.31), X is a power series in $1/z$ and also a power series in t . Simple manipulation of (3.23) shows

$$X = t\phi(X), \quad \text{where} \quad \phi(z) = (1 + \theta) \frac{(1 + z)^{\theta+1}}{(1 + \theta z/(1 + \theta))^{\theta}}, \tag{3.35}$$

which is of the form (3.32) with $a = 0$. Applying (3.33) with $f(\zeta) = \zeta$ and recalling (3.31) shows that

$$\tilde{m}_p^J = \frac{L^{-p}}{p!} \frac{d^p X}{dt^p} \Big|_{t=0} = L^{-p} \frac{(1 + \theta)^p}{p!} \frac{d^{p-1}}{dz^{p-1}} \left(\frac{(1 + z)^{\theta+1}}{(1 + \theta z/(1 + \theta))^{\theta}} \right)^p \Big|_{z=0}. \tag{3.36}$$

Using the binomial theorem to expand the two main factors on the right-hand side in (3.36) into power series in z , then combining the coefficients appropriately to form a single power series shows

$$\tilde{m}_p^J = L^{-p} \frac{(1 + \theta)^p}{p} \sum_{q=0}^{p-1} \binom{(\theta + 1)p}{q} \binom{-\theta p}{p - 1 - q} \left(\frac{\theta}{1 + \theta} \right)^{p-1-q}. \tag{3.37}$$

The sum can be recognised as a polynomial example of a particular ${}_2F_1$ Gaussian hypergeometric function, allowing us to write

$$\tilde{m}_p^J = L^{-p} \frac{\theta^p (1 + \theta)}{p - \theta} \binom{-\theta p}{p - 1} {}_2F_1 \left(\begin{matrix} 1 - p, -p(1 + \theta) \\ 2 - p(1 + \theta) \end{matrix} \middle| \frac{1 + \theta}{\theta} \right). \tag{3.38}$$

The functional equation for the gamma function shows

$$\binom{-\theta p}{p - 1} = \frac{(-1)^{p-1} p}{p(\theta + 1) - 1} \binom{p(1 + \theta) - 1}{p}. \tag{3.39}$$

Also, in general, it is a simple exercise to verify from the series definition of ${}_2F_1$ that

$$\begin{aligned} {}_2F_1(a, b; c; z) &= \frac{2b - c + 2 + (a - b - 1)z}{b - c + 1} {}_2F_1(a, b + 1; c; z) \\ &\quad + \frac{(b + 1)(z - 1)}{b - c + 1} {}_2F_1(a, b + 2; c; z). \end{aligned}$$

In the case of the ${}_2F_1$ in (3.38) we have $2b - c + 2 + (a - b - 1)z = 0$, $b + 2 = c$. Thus on the RHS, only the second term contributes, and furthermore it can be simplified from the fact that ${}_2F_1(a, b + 2; b + 2; z) = (1 - z)^{-a}$, implying the result

$${}_2F_1 \left(\begin{matrix} 1 - p, -p(1 + \theta) \\ 2 - p(1 + \theta) \end{matrix} \middle| \frac{1 + \theta}{\theta} \right) = (-1)^{p-1} \frac{p(1 + \theta) - 1}{\theta^p}. \tag{3.40}$$

Substituting (3.39) and (3.40) into (3.38) we obtain (3.34). □

Remark 3.11. (i) With $1/z = Lt$, the large z expansion of $z\tilde{G}^J(z)$ is thus seen to be a special case of the function

$$F(t) = \sum_{n=0}^{\infty} \binom{\alpha + \beta n}{n} t^n. \tag{3.41}$$

This is intimately related to Lambert’s solution of the trinomial equation $x = q + x^m$ in a power series in q [28]. In mathematical physics, there are applications of (3.41), and its multivariable analogue, in the theory of anyons [5, 6].

(ii) As $z \rightarrow 0$, we read off from (3.23) that $(z\tilde{G}^J(z))^{\theta+1} \sim -z(1/\theta)^\theta$. Since the corresponding global density is given in terms of the resolvent by $d\tilde{\mu}^J(x) = \frac{1}{\pi} \text{Im} \tilde{G}^J(x - i0) dx$, it follows that

$$d\tilde{\mu}^J(x) \underset{x \rightarrow 0^+}{\sim} \frac{\sin \left(\frac{\pi}{\theta+1} \right)}{\pi \theta^{\frac{\theta}{\theta+1}}} x^{-\frac{\theta}{\theta+1}} dx. \tag{3.42}$$

Up to the scale factor $(1/\theta)^{\theta/(\theta+1)}$, this is identical to the known $x \rightarrow 0^+$ behaviour of the global density in the Laguerre case [24, Cor. 2.4 with $p = \theta + 1$, $r = 1$].

(iii) For $\theta = 2$ we have the explicit functional form [42]

$$d\tilde{\mu}^J(x)|_{\theta=2} = \left(\frac{3(1 + \sqrt{1-x})^{1/3}}{4\pi\sqrt{3}(1-x)} x^{-2/3} + \frac{3(1 + \sqrt{1-x})^{-1/3}}{4\pi\sqrt{3}(1-x)} x^{-1/3} \right) dx.$$

The corresponding leading term behaviour for $x \rightarrow 0^+$ agrees with that implied by (3.42).

In addition to the above remarks, we draw attention to a relationship between the Jacobi Muttalib–Borodin ensemble and products of truncations of Haar distributed unitary matrices. Let U_j be a Haar distributed unitary random matrix of size $m_j \times m_j$ and let T_j be the corresponding $(n + \nu_j) \times (n + \nu_{j-1})$, with $\nu_j \geq 0$ and $\nu_0 = 0$, and $m_j \geq 2n$ upper left block such that $\ell \geq 2n$. Let G_j denote a standard complex Gaussian matrix of size $(n + \nu_j) \times (n + \nu_{j-1})$. According to the recent work [27], in the limit $n \rightarrow \infty$ the singular values squared of the random matrix product

$$G_r \cdots G_{s+1} T_s \cdots T_1 \tag{3.43}$$

has a density such that its moments $J_{r,s,1}(n)$ are given by

$$J_{r,s,a}(n) = \frac{a}{n} \left(\frac{a^r}{(1+a)^s} \right)^n P_{n-1}^{(\alpha_{n-1}, \beta_{n-1})} \left(\frac{1-a}{1+a} \right), \tag{3.44}$$

where

$$\alpha_n = rn + r + 1, \quad \beta_n = -(r + 1 - s)n - (r + 2 - s)$$

and $P_k^{(\alpha, \beta)}(x)$ denotes the Jacobi polynomial. It is also required that $r > s$ and thus at least one Gaussian matrix in the product. The immediate relevance of this work is due to the fact that the moments (3.44) are shown in [27] to be such that $w = azG(z)$, where $G(z)$ denotes the corresponding resolvent, satisfy

$$w^{r+1} - z(w - a)(w + 1)^s = 0. \tag{3.45}$$

This is the same as our equation (3.23) with $w/a = z\tilde{G}^J$, $r = s = a = \theta$. Indeed for these parameters we can check that (3.44) reduces to (3.34), up to the form of the scale factor L .

Subsequent to [27], the work [31] has considered integrability and exactly solvable features of the random matrix product (3.43) with $r = s$. In particular, it has been shown that the corresponding characteristic polynomial for the squared singular values is given by [31, Eq. (2.31) with $r \mapsto s$]

$$P_n(x) = G_{s+1, s+1}^{0, s+1} \left(\begin{matrix} n + 1, n - m_1, \dots, n - m_s \\ 0, -\nu_1, \dots, -\nu_s \end{matrix} \middle| x \right), \tag{3.46}$$

where $G_{s+1, s+1}^{0, s+1}$ is a particular Meijer G -function and instead of the constraint $m_j \geq 2n$ as required in [27], it is required $m_1 \geq 2n_1 + \mu_1$ and $m_j \geq n + \nu_j + 1$ ($j = 2, \dots, n$). According to the strategy introduced in [24], and applied in the derivation of Propositions 3.4 and 3.8 above, the significance of this result is that the Meijer G -function satisfies a linear differential equation (see e.g. [39, Sec. 5.8]) allowing us to deduce a polynomial equation for the corresponding resolvent $G(z)$. Specifically, this procedure gives

$$z(zG(z) - 1) \prod_{j=1}^s (zG(z) - 1 + \alpha_j) = zG(z) \prod_{j=1}^s (zG(z) + \beta_j), \tag{3.47}$$

where $\alpha_j = \lim_{n \rightarrow \infty} m_j/n$, $\beta_j = \lim_{n \rightarrow \infty} \nu_j/n$. We see that (3.47) reduces to (3.23) in the case $s = \theta$, $\beta_j = 0$ ($j = 1, \dots, s$), $\alpha_j = 1 + 1/\theta$ ($j = 1, \dots, s$). Thus we learn that the global spectral density for the Jacobi Muttalib–Borodin ensemble in the case $\theta = s \in \mathbb{Z}^+$ is the same as the global spectral density for the squared singular values of the random matrix product $T_s \cdots T_1$ where each T_j is an $n \times n$ sub-block of a Haar distributed unitary matrix of size $(n + p) \times (n + p)$ where $\lim_{n \rightarrow \infty} p/n = 1/s$.

Choosing instead $\beta_j = 0$ ($j = 1, \dots, s$) and $\alpha_j = 1 + 1/a$ we see that (3.47) is the same equation as (3.45) with $w/a = zG$ in the latter. Thus $J_{s,s,a}(n)$ has an interpretation as

the moments of the spectral density of the squared singular values of the random matrix product $T_s \cdots T_1$ where each T_j is an $n \times n$ sub-block of a Haar distributed unitary matrix of size $(n+p) \times (n+p)$ where $\lim_{n \rightarrow \infty} p/n = 1/a$ for general $a > 0$. The case $a \rightarrow 0$ exhibits further structure. Then the underlying unitary matrices are of infinite size, and after rescaling the sub-blocks have a Gaussian distribution. More specifically, an $n \times n$ sub-block U_n of an $(n+p) \times (n+p)$ Haar distributed unitary matrix has a distribution proportional to $(\det(\mathbb{I}_n - U_n^\dagger U_n))^{p-n}$ [53]. It follows that for $p \rightarrow \infty$, and with n fixed, the distribution of $G_n = \sqrt{p}U_n$ is proportional to $e^{-G_n^\dagger G_n}$. Hence G_n is a standard complex Gaussian matrix. For $a \rightarrow 0$ we see from (3.44) that

$$a^{-1-sn} J_{s,s,a}(n) \rightarrow \frac{1}{n} P_{n-1}^{(sn+1,-n)}(1) = \frac{1}{sn+1} \binom{sn+n}{n}. \quad (3.48)$$

These are Fuss–Catalan numbers (3.1), which we know are moments of global spectral limit for the squared singular values of a product of s standard complex Gaussian matrices.

4 The global density — saddle point method

It has already been remarked that the Borodin–Muttalib ensemble (1.1) is intimately related to biorthogonal polynomials of one variable as specified by (3.2). Moreover, as noted in [43] (1.1) is an example of a determinantal point process. This means that the general k -point correlation function $\rho_{(k)}(x_1, \dots, x_k)$ is fully determined by a function $K(x, y) = K_N(x, y)$, independent of k and referred to as the correlation kernel, according to the formula

$$\rho_{(k)}(x_1, \dots, x_k) = \det[K(x_i, x_j)]_{i,j=1,\dots,k}. \quad (4.1)$$

Moreover, the correlation kernel is expressed in terms of the biorthogonal polynomials p_l, q_l defined and corresponding normalisation h_l in (3.2) according to

$$K(x, y) = e^{-(V(x)+V(y))/2} \sum_{l=0}^{N-1} \frac{1}{h_l} p_l(x) q_l(y^\theta). \quad (4.2)$$

Since we focus on the classical Laguerre and Jacobi cases, we denote the correlation kernel by $K^L(x, y)$ for V defined in (1.5) and $K^J(x, y)$ for V defined in (1.6).

In [13], an indirect way to transform the summation in (4.2) was devised for the classical Laguerre and Jacobi weights. This transformed summation enabled the computation of the so called hard edge scaled limit. This refers to the limit $N \rightarrow \infty$ with x, y scaled so that in the neighbourhood of the origin the spacing between eigenvalues is of order unity. Our present interest is in the global limit of the one-point function. The transformed summation of [13] is not suited for that purpose. Fortunately Propositions 2.10 and 2.16 provide the double contour integral formulas for both the Laguerre and Jacobi cases of the Borodin–Muttalib ensemble, which is suited. In Section 2 we showed that they are spectrally equivalent to the upper-triangular ensemble (defined in (2.2)) and its Jacobi counterpart (defined in (2.24)) respectively, but with restricted values of α_i and β_i .

4.1 The Laguerre Muttalib–Borodin ensemble

In the Laguerre case, comparing the correlation kernel formula (4.2) with the correlation formula (2.22) with $n_1 = n_2 = N$, we derive the kernel formula (1.7) in Proposition 1.2, modulo the factor $h(x)/h(y)$. A multiplicative factor in the form of $h(x)/h(y)$ in the correlation kernel does not change the joint probability density function expressed in (4.1), so the kernel formula in (1.7) is valid. This factor account for the different

meaning of $K(n_1, x; n_2, y)$ adopted in Proposition 2.10 in consistency with [2, Eq. (21)], which has the factor $e^{-(V(x)+V(y))/2}$ in (4.2) replaced by e^{-xy^c} , see also [2, Eq. (108)] and derivations above it. Thus the function h can be determined, up to a multiplicative constant, by the requirement

$$\frac{h(x)}{h(y)} = \left(\frac{x}{y}\right)^{c/2} e^{(x-y)/2}. \tag{4.3}$$

We recall from (2.21) that the upper-triangular ensemble is well defined for $\theta = 0$ and $c > -1$ and it is the $\theta \rightarrow 0^+$ limit of the Laguerre Muttalib–Borodin ensemble. We call it the $\theta = 0$ case of the ensemble. The double contour integral formula (1.7) is still valid in this case, where $\alpha_j = c$ for all $j = 1, \dots, N$. Note that this case is degenerate in the sense that the biorthogonal polynomials p_i and q_i are not well defined by (3.2).

Below we consider the limiting global density in two cases, first for $\theta > 0$ and next for $\theta = 0$. We give details of the derivation in the former case, while point out the differences of the argument in the latter.

4.1.1 The $\theta > 0$ case

We seek to use (1.7) to compute the density function of the the transformed limiting counting measure $d\tilde{\mu}$ in (3.7). By definition the density function is given by (3.15) and the required change of variables is $\tilde{x} = (x/\theta)^\theta$ (recall the text below (3.7)), so (4.2) gives

$$\tilde{\rho}_{(1)}^L(x) = \lim_{N \rightarrow \infty} \tilde{K}^L(x) := \lim_{N \rightarrow \infty} x^{1/\theta-1} K^L(N\theta x^{1/\theta}, N\theta x^{1/\theta}), \tag{4.4}$$

and is the global density appearing in (3.9). The scale factor θ is chosen with the benefit of hindsight as it allows (3.10) to be reclaimed without the need for rescaling.

Under the assumption $\theta \in \mathbb{Z}^+$, we showed in Section 3.1 that the resolvent $\tilde{G}^L(z)$ defined in (3.9) corresponding to $d\tilde{\mu}^L$, satisfies the identity (3.10) characterizing the Fuss–Catalan density. Consequently

$$\tilde{\rho}_{(1)}^L(x) = \lim_{\epsilon \rightarrow 0^+} \frac{1}{2\pi i} \left(\tilde{G}^L(x - i\epsilon) - \tilde{G}^L(x + i\epsilon) \right), \tag{4.5}$$

is the Fuss–Catalan distribution supported on $I = (0, (1 + \theta)^{1+\theta}\theta^{-\theta})$. The parametrization by Biane and independently Neuschel [9, 44] gives that for $x \in I$, there is a unique $\varphi \in (0, \pi/(\theta + 1))$ such that

$$x = \frac{(\sin((\theta + 1)\varphi))^{\theta+1}}{\sin \varphi (\sin(\theta\varphi))^\theta}, \quad \text{where } 0 < \varphi < \frac{\pi}{\theta + 1}, \tag{4.6}$$

and which allows for the simple functional form

$$\rho^{\text{F-C}}(x) = \frac{1}{\pi x} \frac{\sin((\theta + 1)\varphi) \sin \varphi}{\sin(\theta\varphi)} = \frac{\sin(\theta\varphi)^{\theta-1} (\sin \varphi)^2}{\pi \sin((\theta + 1)\varphi)^\theta}. \tag{4.7}$$

With the meaning of $\rho^{\text{F-C}}(x)$ in (1.8) so established, we now turn to our main task of this subsection.

Proof of Proposition 1.3. Our analysis, which is based on the method of steepest descents, is guided by a very similar calculation carried out recently by Liu, Wang and Zhang [37] in relation to the global density for the squared singular values of a product M standard complex Gaussian matrices. That these two computations should be closely related is not surprising upon recalling from Section 3.1 that the global density in the case $\theta = M \in \mathbb{Z}^+$ coincides with the global density for the squared singular values of

a product M standard complex Gaussian matrices. It turns out that if $\theta \geq 1$, our proof follows that in [37] closely, and if $\theta \in (0, 1)$, we need to construct the contour Σ in an alternative way, which we explain in the proof. As well as guiding our overall strategy, [37] will be referred to for the proof of some technical bounds, which we omit below.

Substituting $N\theta x^{1/\theta}$ for x and $N\theta x^{1/\theta}$ for y in (1.7) gives

$$\tilde{K}^L(x) := x^{\frac{1}{\theta}-1} K^L(N\theta x^{\frac{1}{\theta}}, N\theta x^{\frac{1}{\theta}}) = \frac{1}{N\theta x} \frac{1}{(2\pi i)^2} \oint_{\Sigma} dz \oint_{\Gamma_{\alpha}} dw \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w}, \tag{4.8}$$

where

$$F(z; x) = \log \left(\frac{\Gamma(z+1)}{(N\theta x^{\frac{1}{\theta}})^z} \prod_{k=1}^N (z - c - (k-1)\theta) \right). \tag{4.9}$$

The logarithm function takes the principal branch and we assume that the value of $\log z$ for $z \in (-\infty, 0)$ is continued from above, to remove ambiguity. To derive the asymptotics of $F(z; x)$, we denote

$$z = N\theta u, \quad w = N\theta v, \tag{4.10}$$

and let ϵ be a positive constant. We begin by noting that uniformly in z such that for all z satisfying

$$\text{dist}(z, [0, N\theta]) > \epsilon N, \quad \text{or equivalently} \quad \text{dist}(u, [0, 1]) > \epsilon/\theta, \tag{4.11}$$

we have

$$\begin{aligned} & \log \left(\prod_{k=1}^N (z - c - (k-1)\theta) \right) \\ &= N(\log N + \log \theta) + N \log u + N \int_0^1 \log \left(1 - \frac{c}{N\theta u} - \frac{x}{u} \right) dx \\ & \quad + \sum_{k=1}^N \left[\log \left(1 - \frac{c}{\theta} + \frac{(k-1)\theta}{Nu} \right) - N \int_{\frac{k-1}{N}}^{\frac{k}{N}} \log \left(1 - \frac{c}{N\theta u} - \frac{x}{u} \right) dx \right] \\ &= N \left((1-u) \log \left(1 - \frac{1}{u} \right) + \log u \right) \\ & \quad + \left(\frac{c}{\theta} - \frac{1}{2} \right) \log \left(1 - \frac{1}{u} \right) + N(\log N + \log \theta - 1) + \mathcal{O}(N^{-1}). \end{aligned} \tag{4.12}$$

Next using Stirling’s formula [1, 6.1.37], we have that for z satisfying (4.11) and $\arg z = \arg u \in (-\pi + \epsilon, \pi - \epsilon)$,

$$\log \Gamma(z+1) = N\theta u(\log N + \log \theta + \log u - 1) + \frac{1}{2} \log u + \log \sqrt{2\pi N\theta} + \mathcal{O}(N^{-1}).$$

This allows us to write

$$F(z; x) = N\hat{F}_1(u; x) + \hat{F}_0(u) + N \log N + \log \sqrt{2\pi N\theta} + \mathcal{O}(N^{-1}), \tag{4.13}$$

where

$$\hat{F}_1(u; x) = (\theta + 1)u(\log u - 1) - (u - 1)(\log(u - 1) - 1) - u \log x, \tag{4.14}$$

$$\hat{F}_0(u) = \left(\frac{c}{\theta} - \frac{1}{2} \right) \log \left(1 - \frac{1}{u} \right) + \frac{1}{2} \log u, \tag{4.15}$$

and the error bound $\mathcal{O}(N^{-1})$ holds uniformly for all z satisfying (4.11) and $\arg z = \arg u \in (-\pi + \epsilon, \pi - \epsilon)$. Substituting in (4.8) shows that if we can deform Σ and Γ_{α} such that $\arg z, \arg w \in (-\pi + \epsilon, \pi - \epsilon)$ and $|z|, |w| > \epsilon N$, then

$$\tilde{K}^L(x) = \frac{1}{N\theta x} \frac{1}{(2\pi i)^2} \oint_{\Sigma} dz \oint_{\Gamma_{\alpha}} dw \frac{e^{N\hat{F}_1(u;x) + \hat{F}_0(u;x)}}{e^{N\hat{F}_1(v;x) + \hat{F}_0(v;x)}} \frac{1}{z-w} \left(1 + \mathcal{O}(N^{-1}) \right). \tag{4.16}$$

In the case $\theta = M \in \mathbb{Z}^+$ this contour integral is comparable to [37, Eq. (2.7)], and our $\hat{F}_1(u; x)$ is defined the same as the $\hat{F}(z; a)$ occurring in [37, Eq. (2.6)] with $u = z$ and $x = a$.

It follows from (4.14) that

$$\frac{d}{du} \hat{F}_1(u; x) = \log \frac{u^{\theta+1}}{(u-1)x},$$

and thus stationary points occur for u such that

$$u^{\theta+1} = (u-1)x. \tag{4.17}$$

Note that this is precisely the equation (3.10) after the identification $z \mapsto x, z\tilde{G}^L(z) \mapsto u$ in the latter. It is known that (4.17) permits a pair of complex conjugate solutions for $x \in (0, \theta(1 + 1/\theta)^{\theta+1})$, u_+^L, u_-^L say, which merge to real solutions for $x = 0$ and $x = \theta(1 + 1/\theta)^{\theta+1}$. With x parametrized by φ as in (4.6), we can check that

$$u_+^L = \frac{\sin((\theta+1)\varphi)}{\sin(\theta\varphi)} e^{i\varphi}, \quad u_-^L = \frac{\sin((\theta+1)\varphi)}{\sin(\theta\varphi)} e^{-i\varphi}. \tag{4.18}$$

Note that our u_{\pm}^L are the same as the w_{\pm} defined in [37, Eq. (2.10)]. In terms of this parametrisation we have (comparing with [37, Eq. (2.11)])

$$C_{\pm} := \frac{d^2}{du^2} \hat{F}_1(u_{\pm}^L; x) = \frac{1}{u_{\pm}^L} \left(\theta + 1 - \frac{\sin((\theta+1)\varphi)}{\sin \varphi} e^{\mp i\theta\varphi} \right), \tag{4.19}$$

and in particular $C_{\pm} \neq 0$ for φ in the range given in (4.6).

Next we deform the contours Σ and Γ_{α} for the steepest-descent analysis. We consider the cases $\theta \geq 1$ and $\theta \in (0, 1)$ separately.

Construction of the contours: $\theta \geq 1$ case We assume that $\Re N\theta u_{\pm}^L$ is different from all α_k ($k = 1, \dots, N$). If this assumption is not satisfied, see Remark 4.3 below.

It is clear that the Hankel like contour Σ can be deformed to an infinite contour that is from $-i \cdot \infty$ to $i \cdot \infty$, as long as it keeps the poles $-1, -2, \dots$ of z to its left and Γ_{α} to its right. Actually this deformation has more freedom. By the residue theorem, if Σ is deformed into the infinite vertical contour to the right of Γ_{α} , the double contour integral in (4.8) remains the same. Moreover, we can split Γ_{α} into two positively oriented contours that jointly enclose all the poles $\alpha_1, \dots, \alpha_N$, and let Σ be an infinite vertical contour passing between them. See [37, Eq. (2.8)] for the explicit computation in a similar case. Specifically, deform Σ from the Hankel contour into the upward vertical contour

$$\Sigma = \{ \Re N\theta u_{\pm}^L + it \mid t \in \mathbb{R} \}. \tag{4.20}$$

To express the shape of the deformed contour Γ_{α} , we define first the contours $\tilde{\Gamma}$ and $\tilde{\Gamma}^{\epsilon}$ where $\epsilon > 0$. Thus define

$$\tilde{\Gamma} = \left\{ \frac{\sin((\theta+1)\phi)}{\sin(\theta\phi)} e^{i\phi} \mid \phi \in \left[-\frac{\pi}{\theta+1}, \frac{\pi}{\theta+1} \right] \right\}, \tag{4.21}$$

and for a small enough $\epsilon > 0$, (see Figure 2)

$$\begin{aligned} \tilde{\Gamma}^{\epsilon} = \{ z \in \tilde{\Gamma} \mid |z| \geq \epsilon \} \\ \cup \text{ the arc of } \{ |z| = \epsilon \} \text{ connecting } \tilde{\Gamma} \cap \{ |z| = \epsilon \} \text{ and through } -\epsilon. \end{aligned} \tag{4.22}$$

Then we define Γ_α , depending on two small positive constants ϵ and ϵ' . Here we take the notational convention that if C is a contour and $r > 0$, then rC is the contour consisting of $\{z \mid z/r \in C\}$ and with the same orientation. In terms of this notation

$$\Gamma_\alpha = \Gamma_{\text{curved}} \cup \Gamma_{\text{vertical}}, \quad \text{where} \quad \Gamma_{\text{curved}} = \Gamma_1 \cup \Gamma_2, \quad \Gamma_{\text{vertical}} = \Gamma_3 \cup \Gamma_4, \quad (4.23)$$

and

$$\Gamma_1 = N\theta\tilde{\Gamma}^r \cap \{z \mid \Re z \leq \Re N\theta u_{\pm}^L - \epsilon\}, \quad (4.24)$$

$$\Gamma_2 = N\theta\tilde{\Gamma}^r \cap \{z \mid \Re z \geq \Re N\theta u_{\pm}^L + \epsilon\}, \quad \text{with } r = \frac{[\epsilon'N] + \frac{1}{2}}{N}, \quad (4.25)$$

$$\Gamma_3 = \text{vertical bar connecting the two ending points of } \Gamma_1, \quad (4.26)$$

$$\Gamma_4 = \text{vertical bar connecting the two ending points of } \Gamma_2. \quad (4.27)$$




Figure 2: The schematic shape of $\tilde{\Gamma}^\epsilon$ for the Laguerre Muttalib–Borodin ensemble with $\theta \geq 1$.




Figure 3: The schematic shapes of Σ and Γ_α for the Laguerre Muttalib–Borodin ensemble with $\theta \geq 1$.




Figure 4: The schematic shape of $\tilde{\Gamma}^\epsilon$ for the Laguerre Muttalib–Borodin ensemble with $\theta \in (0, 1)$.




Figure 5: The schematic shapes of Σ and Γ_α for the Laguerre Muttalib–Borodin ensemble with $\theta \in (0, 1)$.

Note that $\Gamma_1 \cup \Gamma_3$ and $\Gamma_2 \cup \Gamma_4$ are disjoint closed contours, and we assume that they are both oriented counterclockwise. Here we choose ϵ small enough so that $\Re N\theta u_{\pm}^L \pm \epsilon$ lie between two poles of the integral α_j and α_{j+1} , and then $\Gamma_1 \cup \Gamma_3$ and $\Gamma_2 \cup \Gamma_4$ jointly cover all the poles $\alpha_1, \dots, \alpha_N$. Our contours Σ and Γ_α are identical to the contours \mathcal{C} and Σ respectively defined in [37, Sec. 2.2] up to the factor θ and our $\tilde{\Gamma}$ and $\tilde{\Gamma}^\epsilon$ are identical

to $\tilde{\Sigma}$ and $\tilde{\Sigma}^\epsilon$ in [37, Sec. 3.1] respectively, if $\theta = M \in \mathbb{Z}^+$. See Figure 3 for the shapes of Σ and Γ_α .

Construction of contours: $\theta \in (0, 1)$ case The construction of the contours in the $\theta \geq 1$ case obviously is not valid if $\theta < 1$ and x is close to 0, since if the vertical line through u_\pm^L will intersect $\tilde{\Gamma}^\epsilon$ defined in (4.22) at other points, see Figure 4. So Σ needs to be deformed into a more complicated shape, and the construction becomes less straightforward. We use method of elementary dynamical systems to construct the contour Σ .

We construct $\Sigma = N\theta\tilde{\Sigma}$, where $\tilde{\Sigma}$ is a contour passing through u_\pm^L . Consider the gradient field generated by the function $\Re\hat{F}_1(z; x)$, denoted by $\nabla\Re\hat{F}_1(z; x)$. Through a regular point, there is a unique flow line associated to $\nabla\Re\hat{F}_1(z; x)$, and the value of $\Re\hat{F}_1(z; x)$ increases as z moves along the flow line. Since $\Re\hat{F}_1(z; x)$ is a harmonic function, a flow line will not stop at a local maximum or start from a local minimum. At a saddle point, we can concatenate a flow line into it and a flow line out of it, and $\Re\hat{F}_1(z; x)$ increases along the concatenated flow line. Hence we have that any flow line can be prolonged so that it starts from either infinity or a singular point of $\nabla\Re\hat{F}_1(z; x)$, (that is, 0 or 1,) although the prolonged flow line may not be unique if it goes through a saddle point.

By the definition (4.18) of u_+^L , we have that u_+^L is a critical point of $\nabla\Re\hat{F}_1(z; x)$ of second order, so from this point there are two flow lines going into this point. Note that the curve $\tilde{\Gamma} \cap \mathbb{C}_+$ divides \mathbb{C}_+ into two disconnected parts. Since $\Re\hat{F}_1(z; x)$ attains its minimum over $\tilde{\Gamma} \cap \mathbb{C}_+$ at u_+^L , the two flow lines into u_+^L cannot intersect $\tilde{\Gamma} \cap \mathbb{C}_+$. We denote γ_1 the flow line coming to u_+^L from below $\tilde{\Gamma} \cap \mathbb{C}_+$, and γ_2 the flow line coming to u_+^L from above it.

Since the gradient field $\nabla\Re\hat{F}_1(z; x)$ does not have a singularity within the region enclosed by $\tilde{\Gamma} \cap \mathbb{C}_+$ and the interval $[0, 1 + \theta^{-1}]$, flow line γ_1 does not start within the region, and has to enter the region before reaching u_+^L . Since the value of $\Re\hat{F}_1(z; x)$ on $\tilde{\Gamma} \cap \mathbb{C}_+$ attains its minimum at u_+^L , γ_1 has to enter the region from the $[0, 1 + \theta^{-1}]$ side. Hence we let σ_1 be the part of γ_1 between its (last) entrance to the region and u_+^L . It is clear that σ_1 lies in the region and connects u_+^L and a point on $(0, 1 + \theta^{-1})$. (We can further show that this point is 1, but it is not relevant to our construction.)

On the other hand, we want to show that the flow line γ_2 does not intersect the real axis. If this holds, it has to be from infinity to u_+^L , and from the limiting behaviour of $\hat{F}_1(z; x)$ we know that it comes from the direction $e^{i\pi} \cdot \infty$. We construct $\tilde{\Sigma} \cap \mathbb{C}_+$ as $\gamma_2 \cup \sigma_1$, and then have $\tilde{\Sigma}$ by taking the reflection about the real axis. In the end, we take $\Sigma = N\theta\tilde{\Sigma}$ and orient Σ as from $N\theta u_-^L$ to $N\theta u_+^L$, and finish the construction of Σ .

To show that γ_2 does not intersect the real axis, or more specifically, $\mathbb{R} \setminus [0, 1 + \theta^{-1}]$, we note that the ray $(1 + \theta^{-1}, +\infty)$ is a flow line, so γ_2 does not intersect with it since γ_2 does not overlap with it. Let $x^* \in (-\infty, 0)$ be the unique critical point of $\nabla\Re\hat{F}_1(z; x)$ on $(-\infty, 0)$, then $(-\infty, x^*)$ and $(x^*, 0)$ are flow lines, so γ_2 can intersect with $(-\infty, 0)$ only at x^* . However, we can check that $\Re\hat{F}_1(x^*; x) > \Re\hat{F}_1(0; x) > \Re\hat{F}_1(u_+^L; x)$, so γ_2 cannot pass through x^* before reaching u_+^L .

Now we briefly describe the construction of Γ_α in the $\theta \in (0, 1)$ case. We still define $\tilde{\Gamma}^\epsilon$ by (4.22), and then define Γ_1 and Γ_2 by (4.24) and (4.25), and also define Γ_{curved} as the union of Γ_1 and Γ_2 , as in (4.23). Next, analogous to Γ_3 and Γ_4 defined in (4.26) and (4.27), we define, in our case, Γ_3 as a contour connecting the two ending points of Γ_1 such that all points of Γ_3 are within distance ϵ to Σ , and Γ_4 as a contour connecting the two ending points of Γ_2 such that all points of Γ_4 are within distance ϵ to Σ . Finally we denote the union of Γ_3 and Γ_4 as Γ_{vertical} , although Γ_3 and Γ_4 are no longer vertical, and

let $\Gamma = \Gamma_{\text{curved}} \cup \Gamma_{\text{vertical}}$ as in (4.23). Hence we finish the construction of the contours in the $\theta \in (0, 1)$ case, and see Figure 5 for the shape of the contours.

Remark 4.1. The construction of contours Σ and Γ_α above also applies for the $\theta \geq 1$ case. We still keep the construction with vertical Σ , because it is conceptually simpler and more analogous to the construction in [37].

Our choice of the contours Σ and $\tilde{\Gamma}_\alpha$, in either the $\theta \geq 1$ case or the $\theta \in (0, 1)$ case, allows $N\theta u_\pm^L$ to be identified as maximum points of $\Re F(z; x)$ for $z \in \Sigma$ and minimum points of $\Re F(z; x)$ for $z \in \Gamma_{\text{curved}}$, which is key to the subsequent asymptotic analysis. To be precise, we state the result as follows, where we denote $D_r(z) = \{w \in \mathbb{C} \mid |w - z| < r\}$.

Lemma 4.2. *There exists $\delta > 0$ such that for all N large enough,*

$$\Re F(z; x) \geq \Re F(N\theta u_\pm^L) + \delta N \left| \frac{z}{N\theta} - u_\pm^L \right|^2, \quad z \in \Gamma_{\text{curved}} \cap D_{N^{\frac{3}{5}}}(N\theta u_\pm^L), \tag{4.28}$$

$$\Re F(z; x) > \Re F(N\theta u_\pm^L) + \delta N^{\frac{1}{5}}, \quad z \in \Gamma_{\text{curved}} \setminus \left(D_{N^{\frac{3}{5}}}(N\theta u_+^L) \cup D_{N^{\frac{3}{5}}}(N\theta u_-^L) \right), \tag{4.29}$$

$$\Re F(z; x) \leq \Re F(N\theta u_\pm^L) - \delta N \left| \frac{z}{N\theta} - u_\pm^L \right|^2, \quad z \in \Sigma \cap D_{N^{\frac{3}{5}}}(N\theta u_\pm^L), \tag{4.30}$$

$$\Re F(z; x) < \Re F(N\theta u_\pm^L) - \delta N^{\frac{1}{5}}, \quad z \in \Sigma \setminus \left(D_{N^{\frac{3}{5}}}(N\theta u_+^L) \cup D_{N^{\frac{3}{5}}}(N\theta u_-^L) \right), \tag{4.31}$$

$$\Re F(z; x) < \Re F(N\theta u_\pm^L) - \delta |z| \quad z \in \Sigma \cap \{|z| > \delta^{-1}N\}. \tag{4.32}$$

The most important ingredient of the proof of Lemma 4.2 is the estimate of the leading term $\hat{F}_1(z; x)$, which is identical to $\hat{F}(z; x)$ in [37] if $\theta = M$. In the $\theta \geq 1$ case, the estimate of $\hat{F}_1(z; x)$ is stated in [37, Lem. 3.1 and 3.2], where the results and the proofs hold for general $\theta > 1$. Then Lemma 4.2 is proved analogously to the proof of [37, Lem. 2.1] in [37, Sec. 3.2]. We omit the detail. In the $\theta \in (0, 1)$ case, (4.28) and (4.29) are the same as the $\theta \geq 1$ case, and (4.30), (4.31), (4.32) are due to the flow line definition of Σ . We also omit the detail.

Remark 4.3. In the $\theta \geq 1$ case, if $\Re Nu_\pm^L$ happens to be identical to a pole α_j , we replace $\Re Nu_\pm^L$ into $\Re Nu_\pm^L + 1/2$ in formulas (4.20), (4.24) and (4.25). All later arguments are valid with notational changes. In the $\theta \in (0, 1)$ case, if Σ hits a pole α_j , we deform Σ and Γ_{vertical} slightly in an analogous way to avoid the pole.

Taking the limit $\epsilon \rightarrow 0$, we have

$$\tilde{K}^L(x) = I_1 + I_2, \tag{4.33}$$

where, with p.v. denoted the principal value integral,

$$\begin{aligned} I_1 &= \lim_{\epsilon \rightarrow 0} \frac{(N\theta x)^{-1}}{(2\pi i)^2} \int_\Sigma dz \int_{\Gamma_1 \cup \Gamma_2} dw \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w} \\ &= \frac{(N\theta x)^{-1}}{(2\pi i)^2} \text{p.v.} \int_{N\theta \tilde{\Gamma}^r} \left(\int_\Sigma dz \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w} \right) dw, \end{aligned} \tag{4.34}$$

and

$$\begin{aligned} I_2 &= \lim_{\epsilon \rightarrow 0} \frac{(N\theta x)^{-1}}{(2\pi i)^2} \int_\Sigma dz \int_{\Gamma_3 \cup \Gamma_4} dw \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w} \\ &= \frac{(N\theta x)^{-1}}{2\pi i} \int_{N\theta u_-^L}^{N\theta u_+^L} \frac{e^{F(z;x)}}{e^{F(z;x)}} dz \\ &= \frac{\Im u_+^L}{\pi x} = \frac{(\sin \varphi)^2 (\sin(\theta \varphi))^{\theta-1}}{\pi (\sin((\theta + 1)\varphi))^\theta}. \end{aligned} \tag{4.35}$$

To evaluate I_1 , we define

$$\Sigma_{\text{local}}^{\pm} = \Sigma \cap D_{N^{\frac{3}{5}}}(N\theta u_{\pm}^L), \quad \Gamma_{\text{local}}^{\pm} = \tilde{\Gamma}_{\alpha} \cap D_{N^{\frac{3}{5}}}(N\theta u_{\pm}^L). \quad (4.36)$$

Our strategy is to consider the Cauchy principal integral (4.34) on $\Sigma_{\text{local}}^+ \times \Gamma_{\text{local}}^+$ and $\Sigma_{\text{local}}^- \times \Gamma_{\text{local}}^-$ first, and then to show that the remaining part of the integral is negligible in the asymptotic analysis.

Make the change of variables

$$z = N\theta u_+^L + N^{\frac{1}{2}}\theta s, \quad w = N\theta u_+^L + N^{\frac{1}{2}}\theta t. \quad (4.37)$$

Then by (4.13)

$$\begin{aligned} F(z; x) &= N\hat{F}_1(u_+^L + N^{-\frac{1}{2}}s; x) + \hat{F}_0(u_+^L + N^{-\frac{1}{2}}s) + N \log N + \log \sqrt{2\pi N\theta} + \mathcal{O}(N^{-1}) \\ &= \frac{C_+}{2}s^2 + N\hat{F}_1(u_+^L; x) + N \log N + \log \sqrt{2\pi N\theta} + \hat{F}_0(u_+^L) + \mathcal{O}(N^{-\frac{1}{5}}), \end{aligned} \quad (4.38)$$

where C_+ is defined in (4.19). This gives

$$\begin{aligned} \text{p. v.} \int_{\Gamma_{\text{local}}^+} dw \int_{\Sigma_{\text{local}}^+} dz \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w} \\ = \frac{1}{\sqrt{N\theta}} \text{p. v.} \int_{\Gamma_{\text{local}}^+} dw \int_{\Sigma_{\text{local}}^+} dz \frac{e^{\frac{C_+}{2}s^2}}{e^{\frac{C_+}{2}t^2}} \frac{1 + \mathcal{O}(N^{-\frac{1}{5}})}{s-t}. \end{aligned} \quad (4.39)$$

Note that the $\mathcal{O}(N^{-1/5})$ term in the integrand on the right-hand side of (4.39) is uniform and analytic in $N_{N^{3/5}}(N\theta u_+^L)$. Using (4.28) of (4.29) in Lemma 4.2, we find that there exists $c_1, c_2 > 0$ such that for $z \in \Sigma_{\text{local}}^+$ and $w \in \Gamma_{\text{local}}^+$,

$$\left| e^{\frac{C_+}{2}s^2} \right| \leq e^{-c_1 s^2}, \quad \left| e^{\frac{C_+}{2}t^2} \right| \geq e^{c_2 t^2}. \quad (4.40)$$

Hence standard application of the saddle point method yields

$$\text{p. v.} \int_{\Gamma_{\text{local}}^+} dw \int_{\Sigma_{\text{local}}^+} dz \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w} = \mathcal{O}(N^{\frac{1}{2}}). \quad (4.41)$$

Analogous reasoning gives

$$\text{p. v.} \int_{\Gamma_{\text{local}}^-} dw \int_{\Sigma_{\text{local}}^-} dz \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w} = \mathcal{O}(N^{\frac{1}{2}}). \quad (4.42)$$

Finally, by (4.30), (4.31) and (4.32) in Lemma 4.2, there exists $c_3 > 0$ such that for large enough N

$$\left| e^{-F(w;x)} \right| < \left| e^{-F(N\theta u_{\pm}^L; x)} \right| e^{-c_3 N^{\frac{1}{5}}} \quad \text{if } w \in \Gamma_{\alpha} \setminus \Gamma_{\text{local}}^{\pm}, \quad (4.43)$$

$$\left| e^{F(z;x)} \right| < \begin{cases} \left| e^{F(N\theta u_{\pm}^L; x)} \right| e^{-c_3 N^{\frac{1}{5}}} & \text{if } z \in \Sigma \setminus \Sigma_{\text{local}}^{\pm}, \\ \left| e^{F(N\theta u_{\pm}^L; x)} \right| e^{-c_3 |z|} & \text{if } z \in \Sigma \setminus \{|z| > \frac{N}{c_3}\}. \end{cases} \quad (4.44)$$

With the help of estimates (4.43), (4.44), (4.38) and (4.40), we obtain

$$\begin{aligned} \text{p. v.} \int_{\tilde{\Gamma}_{\alpha}} dw \int_{\Sigma_{-1}} dz \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w} - \text{p. v.} \int_{\Gamma_{\text{local}}^+} dw \int_{\Sigma_{\text{local}}^+} dz \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w} \\ - \text{p. v.} \int_{\Gamma_{\text{local}}^-} dw \int_{\Sigma_{\text{local}}^-} dz \frac{e^{F(z;x)}}{e^{F(w;x)}} \frac{1}{z-w} = \mathcal{O}(e^{-\epsilon n^{\frac{1}{5}}}). \end{aligned} \quad (4.45)$$

Plugging (4.41), (4.42) and (4.45) into (4.34), we have that

$$I_1 = \mathcal{O}(N^{-\frac{1}{2}}). \tag{4.46}$$

Therefore we have proved Proposition 1.3 upon combining (4.33), (4.35) and (4.46). \square

Remark 4.4. The asymptotic analysis above confirms a simple relation that

$$\tilde{\rho}_{(1)}^L(x) = \lim_{N \rightarrow \infty} \tilde{K}^L(x) = \frac{\Im u_+^L}{\pi x}. \tag{4.47}$$

This pattern persists in the computation of the global density for the Jacobi Muttalib–Borodin ensemble given in Section 4.2.

Remark 4.5. The asymptotic analysis above can also prove the bulk local universality of the limiting distribution of the eigenvalues, as in [37]. The result, which is unsurprisingly the sine universality, is also the same as in [37]. Since the local universality is not our focus in this paper, we omit further discussion on it. In the case $\theta = 2$ this problem, along with Airy kernel universality at the soft edge, was solved some time ago [38]. Very recently, these universalities have been established for general $\theta > 0$ by Zhang [52] according to the method of [37]. In [52], an equivalent form of (1.7) is also derived.

4.1.2 The $\theta = 0$ case

In Proposition 3.4, we do not use the most straightforward scaling transform

$$x \mapsto \frac{x}{N} \tag{4.48}$$

to compute the limiting global density but rather we took the combined change of variables and scaling transform

$$x \mapsto \left(\frac{x}{N\theta}\right)^\theta, \tag{4.49}$$

like in Proposition 1.3, to conform our result to the Fuss–Catalan distribution. It is clear that the combined change of variables and scaling transform (4.49) is not well defined for $\theta = 0$. So in the $\theta = 0$ case, which is the particularly interesting $\theta \rightarrow 0$ limit of the Laguerre Muttalib–Borodin ensemble (2.21), we use the scaling transform (4.48) instead.

According to (1.7), we then have

$$K^L(x) := K^L(Nx, Nx) = \frac{1}{(2\pi i)^2} \oint_{\Sigma} dz \oint_{\Gamma_\alpha} dw \frac{x^{-z-1} x^w \Gamma(z+1)(z-c)^N}{(z-w)\Gamma(w+1)(w-c)^N} N^{w-z-1}. \tag{4.50}$$

Changing variables $z = Nu$, $w = Nv$ and using Stirling’s formula shows that for large N , if we can deform Σ and Γ_α such that $\arg u, \arg v \in (-\pi + \epsilon, \pi - \epsilon)$ and $|u|, |v| > \epsilon > 0$,

$$K^L(x) = \frac{1}{Nx(2\pi i)^2} \oint_{\Sigma} dz \oint_{\Gamma_\alpha} dw \frac{1}{z-w} \frac{e^{NH_1(u;x) + \frac{1}{2} \log u - \frac{c}{u}}}{e^{NH_1(v;x) + \frac{1}{2} \log v - \frac{c}{v}}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right),$$

where $H_1(u; x) := u(\log u - 1) - u \log x + \log u$. Noting that

$$\frac{d}{du} H_1(u; x) = \log\left(\frac{u}{x} e^{1/u}\right)$$

we see that the stationary points of $H_1(u; x)$ occur when

$$u e^{1/u} = x, \quad \text{or equivalently} \quad \left(-\frac{1}{u}\right) e^{-\frac{1}{u}} = -\frac{1}{x}. \tag{4.51}$$

This equation is solved by the Lambert W function [47, Sec. 4.13] — recall too Remark 3.5(i) — and the two solutions are complex conjugates

$$u_+^L = -W\left(-\frac{1}{x}\right), \quad u_-^L = -\overline{W\left(-\frac{1}{x}\right)}. \tag{4.52}$$

Then using steepest descent arguments like in the $\theta > 0$ case, we have

Proposition 4.6. As $\theta \rightarrow 0$, the limiting global density of the Muttalib–Borodin ensemble is

$$\lim_{N \rightarrow \infty} K^L(x) = -\Im \frac{1}{\pi x W(-x^{-1})}. \tag{4.53}$$

The proof is omitted since it is similar to the $\theta > 0$ case. Note that the relation (4.47) in Remark 4.4 still holds. The result (4.53) was derived in [17, Cor. 1] by a determination of the moments, with the latter also known from [22].

4.2 The Jacobi Muttalib–Borodin ensemble

In the case of the Jacobi Muttalib–Borodin ensemble, comparing the correlation kernel formula (4.2) with the correlation kernel (2.38), we have, analogous to (1.7),

$$K^J(x, y) = \frac{1}{(2\pi i)^2} \frac{\tilde{h}(x)}{\tilde{h}(y)} \times \oint_{\Sigma} dz \oint_{\Gamma_\alpha} dw \frac{x^{-z-1} y^w \Gamma(w + c_2 + N + 1) \Gamma(z + 1) \prod_{k=1}^N (z - \alpha_k)}{(z - w) \Gamma(z + c_2 + N + 1) \Gamma(w + 1) \prod_{l=1}^N (w - \alpha_l)}, \tag{4.54}$$

where $\alpha_j = \theta(j - 1) + c_1$ as specified in (2.3) and $c = c_1$. Analogous to (1.7) we have that the factor $\tilde{h}(x)/\tilde{h}(y)$ in (4.54) cancels out of the determinant (4.1), and the reasoning leading to (4.3) tells us that we can take

$$\tilde{h}(x) = x^{c_1/2} (1 - x)^{-c_2/2}, \quad \text{because} \quad \frac{\tilde{h}(x)}{\tilde{h}(y)} = \left(\frac{x}{y}\right)^{c_1/2} \left(\frac{1 - x}{1 - y}\right)^{-c_2/2}. \tag{4.55}$$

Here we assume $\theta > 0$. The $\theta = 0$ case of the Jacobi upper-triangular ensemble, that is, $\alpha_1 = \dots = \alpha_N = c_1$, is well defined and can be thought as the $\theta \rightarrow 0^+$ limit of the Jacobi Muttalib–Borodin ensemble, but we omit it in this paper. Our aim is to use a saddle point analysis to compute

$$\tilde{\rho}_{(1)}^J(x) = \lim_{N \rightarrow \infty} \tilde{K}^J(x) := \lim_{N \rightarrow \infty} \frac{1}{N\theta} x^{1/\theta-1} K^J(x^{1/\theta}, x^{1/\theta}), \tag{4.56}$$

which is the limiting density of the eigenvalues under the change of variables $x \rightarrow x^\theta$. The required workings is structurally identical to that just given to derive Proposition 1.3; only a brief sketch will be given below.

Recall that in the special case that $\theta \in \mathbb{Z}^+$, the resolvent $\tilde{G}^J(z)$ defined in (3.22), satisfies the identity (3.23), and then the global density $\tilde{\rho}^J(x)$ satisfies

$$\tilde{\rho}_{(1)}^J(x) = \lim_{\epsilon \rightarrow 0^+} \frac{1}{2\pi i} \left(\tilde{G}^J(x - i\epsilon) - \tilde{G}^J(x + i\epsilon) \right), \quad x \in (0, 1). \tag{4.57}$$

Below we show that the global density $\tilde{\rho}^J(x)$ has an explicit formula given as follows. For all $\theta > 0$ and all $x \in (0, 1)$, there is a unique $\varphi \in (0, \pi/(\theta + 1))$ such that

$$x = \frac{\theta^\theta}{(1 + \theta)^{1+\theta}} \frac{\sin((\theta + 1)\varphi)^{\theta+1}}{\sin \varphi \sin(\theta\varphi)^\theta}. \tag{4.58}$$

We define

$$v(\varphi) = \frac{\theta \sin((\theta + 1)\varphi)}{(1 + \theta) \sin(\theta\varphi)} e^{i\varphi}, \tag{4.59}$$

and then the density function

$$\begin{aligned} \rho^{\text{J-M-B}}(x) &= \frac{1}{\pi x} \Im \left(\frac{1}{\theta} \frac{v(\varphi)}{1 - v(\varphi)} \right) \\ &= \frac{1}{(\pi x) \sin(\theta\varphi)^2 + \theta^2 \sin(\varphi)^2 - 2\theta \cos((\theta + 1)\varphi) \sin(\theta\varphi) \sin(\varphi)}. \end{aligned}$$

Thus we have, parallel to Proposition 1.3

Proposition 4.7. For all $\theta > 0$, the limiting global density of the Jacobi Muttalib–Borodin ensemble with change of variable (1.4) is

$$\tilde{\rho}_{(1)}^{\text{J}}(x) = \rho^{\text{J-M-B}}(x).$$

Remark 4.8. The distribution $\rho^{\text{J-M-B}}(x)$ is the counterpart of the Fuss–Catalan distribution $\rho^{\text{F-C}}(x)$ in the Laguerre case. According to the results of Section 3.2, for $\theta \in \mathbb{Z}^+$, the measure $d\mu(x) = \rho^{\text{J-M-B}}(x)dx$ is the unique probability measure supported on $[0, 1]$ that makes the following identity hold:

$$z(zG(z) - 1)(zG(z) + 1/\theta)^\theta = (zG(z))^{\theta+1}, \quad \text{where } G(z) = \int_{[0,1]} \frac{\rho^{\text{J-M-B}}(x)dx}{z - x}. \tag{4.60}$$

This is analogous to the resolvent for the Fuss–Catalan distribution satisfying the identity $z(zG(z) - 1) = (zG(z))^{\theta+1}$ hold. Of interest is to extend the characterisation (4.60) to general $\theta > 0$. Such a characterisation was established in [27] for the measures corresponding to the moments (3.44) with $r > s$, whereas as remarked below (3.45) the moments (3.34) deduced from (4.60) require $r = s$.

The method of the proof to Proposition 4.7 is the same as the proof of Proposition 1.3 for the Laguerre case. So we only give a sketch of the proof and point out the main differences.

Sketch of the proof of Proposition 4.7. First we consider the case that $c_2 \in \mathbb{Z}$. Making the replacements $x \mapsto x^{1/\theta}$, $y \mapsto y^{1/\theta}$ in (4.54) gives

$$\tilde{K}^{\text{J}}(x) := \frac{1}{N\theta} x^{\frac{1}{\theta}-1} K^{\text{J}}(x^{\frac{1}{\theta}}, x^{\frac{1}{\theta}}) = \frac{1}{\theta x} \frac{1}{(2\pi i)^2} \oint_{\Sigma} dz \oint_{\Gamma_\alpha} dw \frac{e^{G(z;x)}}{e^{G(w;x)}} \frac{1}{z - w}, \tag{4.61}$$

where

$$G(z; x) = \log \left(\frac{\Gamma(z + 1)}{\Gamma(z + N + c_2 + 1)} x^{\frac{z}{\theta}} \prod_{k=1}^N (z - c_1 - (k - 1)\theta) \right). \tag{4.62}$$

Changing variables in the integrand according to (4.10), then use of Stirling’s formula and (4.12) shows that for large N

$$G(z; x) = N\hat{G}_1(u; x) + \hat{G}_0(u) + N \log \theta - c_2 \log N + \mathcal{O}(N^{-1}), \tag{4.63}$$

where, after the change of variables of z, w into u, v as in (4.10),

$$\hat{G}_1(u; x) = \theta u \log \frac{\theta u}{1 + \theta u} - \log(1 + \theta u) - u \log x + (1 - u) \log(1 - \frac{1}{u}) + \log u, \tag{4.64}$$

$$\hat{G}_0(u) = \frac{1}{2} \log \frac{\theta u}{1 + \theta u} - c_2 \log(1 + \theta u) + \left(\frac{c}{\theta} - \frac{1}{2} \right) \log(1 - \frac{1}{u}), \tag{4.65}$$

and the validity of the error bound $\mathcal{O}(N^{-1})$ is the same as in (4.13), i.e., all z satisfying (4.11) and $\arg z = \arg u \in (-\pi + \epsilon, \pi - \epsilon)$. Substituting in (4.61) gives

$$\tilde{K}^J(x) = \frac{1}{N\theta x} \frac{1}{(2\pi i)^2} \oint_{\Sigma} dz \oint_{\Gamma_{\alpha}} dw \frac{e^{N\hat{G}_1(u;x)+\hat{G}_0(u)}}{e^{N\hat{G}_1(v;x)+\hat{G}_0(v)}} \frac{1}{z-w} \left(1 + \mathcal{O}(N^{-1})\right). \tag{4.66}$$

This is structurally identical to (4.16), and is analysed accordingly. For the sake of steepest-descent analysis, we need to deform the contours Σ and Γ_{α} . Since we assume that $c_2 \in \mathbb{Z}$, Γ_{α} is a finite contour similar to the Γ_{α} in (4.8). It is clear that Σ can be deformed to an infinite contour that is from $-i \cdot \infty$ to $i \cdot \infty$, as long as it keeps the poles $-1, \dots, -(\beta + N)$ and Γ_{α} to the left. Then we see that we can split Γ_{α} into two positively oriented contours that jointly enclose all the poles $\alpha_1, \dots, \alpha_N$, and let Σ be an infinite vertical contour passing between them. Note that the precise shapes of Σ and Γ_{α} in the Laguerre case depend on the computation of the critical points of $\hat{F}_1(u; x)$ that is the counterpart of our $\hat{G}_1(u; x)$, see (4.18) and (4.21). Below we explain the analogous construction of Σ and Γ_{α} in the Jacobi case.

A difference in detail is now that

$$\frac{d}{du} \hat{G}_1(u; x) = \log \left(\left(\frac{\theta u}{1 + \theta u} \right)^{\theta} \left(\frac{u}{u - 1} \right) \frac{1}{x} \right),$$

so the stationary points now occur for u such that

$$u^{\theta+1} = x(1/\theta + u)^{\theta}(u - 1). \tag{4.67}$$

Analogous to the situation with (4.17), we observe that this is precisely the equation (3.23) after the identification $z \mapsto x$, $z\tilde{G}^J(z) \mapsto u$ in the latter. It is straightforward to verify that if $x \in (0, 1)$ is parametrised by $\varphi \in (0, \pi/(\theta + 1))$ in (4.58), then with $v(\varphi)$ defined in (4.59)

$$u_+^J = \frac{1}{\theta} \frac{v(\varphi)}{1 - v(\varphi)}, \quad u_-^J = \overline{u_+^J} = \frac{1}{\theta} \frac{\overline{v(\varphi)}}{1 - \overline{v(\varphi)}}$$

are two solutions to (4.67). As x runs over $(0, 1)$, the locus of u_+^J (resp. u_-^J) is a curve in \mathbb{C}_+ (resp. \mathbb{C}_-) whose two ends are on the real line. Thus we let Σ be the upward vertical contour through $N\theta u_{\pm}^J$, analogous to (4.20) in the Laguerre case. To define Γ_{α} , we first define

$$\tilde{\Gamma}^J = \left\{ \frac{1}{\theta} \frac{v(\phi)}{1 - v(\phi)} \mid \phi \in \left[-\frac{\pi}{\theta + 1}, \frac{\pi}{\theta + 1} \right] \right\}$$

analogous to $\tilde{\Gamma}$ in (4.21), and then deform it to the desired Γ_{α} parallel to the deformation carried out in the Laguerre case. We omit the detail, but only point out that the shapes of Σ and Γ_{α} is like those in Figure 3 if $\theta \geq 1$ or Figure 5 if $\theta \in (0, 1)$.

By saddle-point analysis analogous to that in the Laguerre case, we derive, as expected in Remark 4.4,

$$\tilde{\rho}_{(1)}^J(x) = \lim_{N \rightarrow \infty} \tilde{K}^J(x) = \frac{\Im u_+^J}{\pi x} = \rho^{J\text{-M-B}}(x),$$

and hence finish the proof.

In the case that $c_2 \notin \mathbb{Z}$, the contour Γ_{α} in (4.54) is infinite. We express it as the combination of $\Gamma'_{\alpha} \cup \Gamma''_{\alpha}$. Here Γ'_{α} is an infinite, Hankel like contour starting at $-\infty - i\epsilon$, running parallel to the negative real axis, looping around the poles $w = -(c_2 + N + k)$ with $k \in \mathbb{Z}_+$, and finishing at $-\infty + i\epsilon$ after again running parallel to the negative real axis. Γ''_{α} is a finite positive oriented contour enclosing $\alpha_1, \dots, \alpha_N$. Then by argument of

contour deformation used in the $c_2 \in \mathbb{Z}$ case and the Laguerre case, we can deform Σ into a contour going from $-i \cdot \infty$ to $i \cdot \infty$ between Γ'_α and Γ''_α . We write

$$\tilde{K}^J(x) = \tilde{K}^{J,'}(x) + \tilde{K}^{J, ''}(x), \quad \text{where}$$

$$\tilde{K}^{J,*}(x) = \frac{1}{\theta x} \frac{1}{(2\pi i)^2} \oint_{\Sigma} dz \oint_{\Gamma_\alpha^*} dw \frac{e^{G(z;x)}}{e^{G(w;x)}} \frac{1}{z-w}, \quad * = ' \text{ or } ''.$$

In the computation of $\tilde{K}^{J, ''}(x)$, we further deform the contours Γ''_α into two parts as the deformation of Γ_α in the $c_2 \in \mathbb{Z}$ case, and let Σ go between the two parts, like the deformed shapes of Γ''_α , Γ'_α and Σ in the $c_2 \in \mathbb{Z}$ case. See Figure 6 if $\theta \geq 1$. By the argument in the $c_2 \in \mathbb{Z}$ case we have that

$$\lim_{N \rightarrow \infty} \tilde{K}^{J, ''}(x) = \rho^{J\text{-M-B}}(x).$$

On the other hand, by estimating the factors of the integrand

$$x^{-z-1} \Gamma(z+1) \prod_{k=1}^N (z - \alpha_k) \Gamma(z + c_2 + N + 1)^{-1}$$

by Stirling’s formula on $\Sigma_{-1/2}$ and

$$y^w \Gamma(w + c_2 + N + 1) \left(\Gamma(w + 1) \prod_{l=1}^N (w - \alpha_l) \right)^{-1}$$

on Γ'_α separately, we find that

$$\lim_{N \rightarrow \infty} \tilde{K}^{J,'}(x) = 0.$$

Thus we prove the proposition by combining the two limits above. □




Figure 6: The deformed shape of contours Σ and Γ_α for the computation of the global density with $\theta \geq 1$ and $c_2 \notin \mathbb{Z}$.




Figure 7: The shape of contours $\Sigma_{-1/2}^\delta$ and Γ_0 for the hard edge correlation kernel.

5 Hard edge density

5.1 The Laguerre Muttalib–Borodin ensemble

5.1.1 The $\theta > 0$ case

The meaning of the hard edge scaling has already been noted in Section 4.1.1. It is the $N \rightarrow \infty$ limit with x, y scaled in the neighbourhood of the origin so that the spacing between eigenvalues is of order unity. The limiting kernel with proper scaling is stated in Proposition 1.4, and below we give the proof.

Proof of Proposition 1.4. First we use the argument in Section 4.1, more specifically the paragraph above (4.20), to justify that the double contour integral formula (1.7) still holds with Σ deformed into a contour from $e^{-(\pi/2+\delta)i} \cdot \infty$ to $e^{(\pi/2+\delta)i} \cdot \infty$ and going on the left of Γ_α . Thus analogous to (4.20) we can take Σ to be $\Sigma_{-1/2}^\delta$ in (1.7). Furthermore it is clear that the closed contour Γ_α can be deformed to the infinite Hankel loop contour Γ_0 .

Substituting (2.3) and making use of the functional and reflection formula for the gamma function shows that

$$\frac{\prod_{k=1}^N (z - \alpha_k)}{\prod_{k=1}^N (w - \alpha_k)} = \frac{\Gamma((z - c)/\theta + 1)\Gamma(N - (z - c)/\theta) \sin \pi(z - c)/\theta}{\Gamma((w - c)/\theta + 1)\Gamma(N - (w - c)/\theta) \sin \pi(w - c)/\theta}.$$

It is standard result that for z and w fixed and $N \rightarrow \infty$,

$$\frac{\Gamma(N - (z - c)/\theta)}{\Gamma(N - (w - c)/\theta)} = N^{(w-z)/\theta} \left(1 + O\left(\frac{1}{N}\right)\right).$$

Taking the limit $N \rightarrow \infty$ inside the integral (a step which is justified by estimating the integrand and using dominated convergence; see [35, Sec. 5.2] for the details in a very similar setting) we thus see that

$$\begin{aligned} \lim_{N \rightarrow \infty} N^{-1/\theta} K^L(N^{-1/\theta} x, N^{-1/\theta} y) &= \left(\frac{x}{y}\right)^{c/2} \\ &\times \frac{1}{(2\pi i)^2} \oint_{\Sigma_{-1/2}^\delta} dz \oint_{\Gamma_0} dw \frac{x^{-z-1} y^w}{z - w} \frac{\Gamma(z + 1)\Gamma((z - c)/\theta + 1) \sin \pi(z - c)/\theta}{\Gamma(w + 1)\Gamma((w - c)/\theta + 1) \sin \pi(w - c)/\theta}. \end{aligned}$$

The result (1.9) now follows by the change of variables $z \mapsto \theta z + c$, $w \mapsto \theta w + c$, and then deforming the contours back to $\Sigma_{-1/2}^\delta$ and Γ_0 . (Since the deformation does not cross any pole, it does not change the value of integral).

In the special case $\theta \in \mathbb{Z}^+$ the duplication formula can be used to rewrite $\Gamma(\theta z + c + 1)/\Gamma(\theta w + c + 1)$ and (1.10) results. \square

Borodin has previously given a different formula for the hard edge scaled limit in (1.9).

Proposition 5.1. [13] We have

$$\lim_{N \rightarrow \infty} N^{-1/\theta} K^L(N^{-1/\theta} x, N^{-1/\theta} y) = \left(\frac{y}{x}\right)^{c/2} K^{(c,\theta)}(x, y) \tag{5.1}$$

where ¹

$$K^{(c,\theta)}(x, y) = \theta x^c \int_0^1 J_{(c+1)/\theta, 1/\theta}(xu) J_{c+1, \theta}(yu)^\theta u^c du, \tag{5.2}$$

with $J_{a,b}$ denoting Wright’s generalisation of the Bessel function given by

$$J_{a,b}(x) = \sum_{j=0}^\infty \frac{(-x)^j}{j! \Gamma(a + jb)}. \tag{5.3}$$

Equating (5.1) and (1.9) gives us a double contour integral form of the kernel (5.2). Before doing this, we note that the term in the second line on the RHS of (1.10) can be identified with the hard edge scaled kernel $K_{\nu_1, \dots, \nu_M}(x, y)$ of Kuijlaars and Zhang [35], which came about from the hard edge scaled limit of the correlation kernel for the product of complex standard Gaussian rectangular random matrices [3]. We also note

¹Our notation $K^{(c,\theta)}(x, y)$ agrees with the notation $K^{(\alpha,\theta)}(x, y)$ defined in [34, Eq. (5.3)], but differs from the $\mathcal{K}^{(\alpha,\theta)}(x, y)$ in [13, Eq. (3.6)] by a factor.

that the hard edge scaled kernel is a component of the kernel for the Meijer G random point field, see [10], [11] and [12]. This kernel reads

$$K_{\nu_1, \dots, \nu_M}(x, y) = \frac{1}{(2\pi i)^2} \int_{\Sigma_{-1/2}^\delta} dz \oint_{\Gamma_0} dw \frac{x^w y^{-z-1}}{z-w} \prod_{j=0}^M \frac{\Gamma(z+1+\nu_j) \sin \pi z}{\Gamma(w+1+\nu_j) \sin \pi w}, \quad (5.4)$$

where $\nu_0 := 0$, $\nu_j > -1$ ($j = 1, \dots, M$). Denoting the second line on the RHS of (1.10) by $\tilde{K}^{(c, \theta)}(x, y)$, for $\theta \in \mathbb{Z}^+$ we see that

$$\tilde{K}^{(c, \theta)}(x, y) = K_{\nu_1, \dots, \nu_\theta}(y, x), \quad \nu_j = \frac{c}{\theta} - 1 + \frac{j}{\theta} \quad (j = 1, \dots, \theta). \quad (5.5)$$

Recalling (5.1) and the LHS of (1.10) we thus have

$$x^{1/\theta-1} K^{(c, \theta)}(\theta x^{1/\theta}, \theta y^{1/\theta}) = K_{\nu_1, \dots, \nu_\theta}(y, x), \quad (5.6)$$

for $\{\nu_j\}$ as in (5.5). This equation has been deduced using a rewrite of (5.2) in [34, Eq. (5.7)].

We now specify the double contour integral formula for Borodin’s kernel (5.2).

Corollary 5.2. *We have*

$$K^{(c, \theta)}(x, y) = \frac{\theta}{(2\pi i)^2} \int_{\Sigma_{-1/2}^\delta} dz \oint_{\Gamma_0} dw \frac{x^{-\theta z-1} y^{\theta w}}{z-w} \frac{\Gamma(\theta z + c + 1) \Gamma(z + 1) \sin \pi z}{\Gamma(\theta w + c + 1) \Gamma(w + 1) \sin \pi w}. \quad (5.7)$$

Remark 5.3. (i) Our proof of (5.7) is indirect. A direct proof can be obtained via the Meijer-G function representation of $J_{a,b}(x)$ in (5.3), analogous to the proof of [34, Thm. 5.1].

(ii) Here we use a nearly vertical contour $\Sigma_{-1/2}^\delta$ to be in consistent with [35]. It is also possible to deform Σ into a Hankel-like form, which is symmetric to Γ_0 . With the help of this symmetry, and under the change of variables $z \mapsto -(z + (c + 1)/\theta)$, $w \mapsto -(w + (c + 1)/\theta)$ in (5.7) the contours Σ and Γ_0 interchange. Making use of the reflection formula for the gamma function then allows us to deduce that

$$\frac{1}{\theta} x^{1/\theta-1} K^{(\alpha, \theta)}(x^{1/\theta}, y^{1/\theta}) = \left(\frac{x}{y}\right)^{\alpha'} K^{(\alpha', 1/\theta)}(y, x), \quad (5.8)$$

which as noted in [34] also follows from the original form (5.2).

(iii) When our article was almost complete, we received a preprint from Zhang [52], containing amongst other things an independent analysis of the hard edge scaling of the Laguerre Muttalib–Borodin ensemble.

5.1.2 The $\theta = 0$ case

We could take the limit of the double contour integral formula (1.7) with $\alpha_1 = \dots = \alpha_N = c$ to derive the limiting correlation kernel near 0 for the $\theta = 0$ case of the Laguerre Muttalib–Borodin ensemble, but here we introduce an alternative approach. Note that the joint PDF of the $\theta = 0$ Laguerre Muttalib–Borodin ensemble is given in (2.21). Changing variables $\log \lambda_j \mapsto \mu_j$, we have that the joint PDF for $\mu_1 \geq \mu_2 \geq \dots \geq \mu_N$ is proportional to

$$\prod_{k=1}^N e^{-V(\mu_k)} \prod_{1 \leq j < k \leq N} (e^{\mu_j} - e^{\mu_k})(\mu_j - \mu_k), \quad \text{where } V(x) = -(c + 1)x + e^x.$$

Then the results in [19] indicate that the correlation functions for the smallest variables μ_N, μ_{N-1}, \dots converge to the correlation functions in the Tracy-Widom distribution, upon proper scaling. More specifically, the results in [19] are only concerned with the asymptotics of biorthogonal polynomials. In principle, by summing up the products of the biorthogonal polynomials, the asymptotics of the correlation kernel results, but technically this is nontrivial. Note that in [19] there is a technical assumption that $V(x) \rightarrow +\infty$ faster than any linear equation as $x \rightarrow \pm\infty$. But it is not hard to see that in the $-\infty$ direction this requirement can be relaxed to a linear growth to $+\infty$.

5.2 The Jacobi Muttalib–Borodin ensemble

The hard edge scaled limit of the Jacobi Muttalib–Borodin ensemble is the same as that for the Laguerre Muttalib–Borodin ensemble, although the specific scale is different [13].

Proposition 5.4. Let $c_1 = c$ in (1.6), and specify the ratio $\tilde{h}(x)/\tilde{h}(y)$ occurring in the definition (4.54) by (4.55). We have

$$\lim_{N \rightarrow \infty} N^{-1-1/\theta} K^J(N^{-1-1/\theta}x, N^{-1-1/\theta}y) = \lim_{N \rightarrow \infty} N^{-1/\theta} K^L(N^{-1/\theta}x, N^{-1/\theta}y). \tag{5.9}$$

Proof. First we consider the case that $c_2 \in \mathbb{Z}$. Then the contour Γ_α in the integral (4.54) is a finite contour enclosing $\alpha_1, \dots, \alpha_N$. By the argument similar to that in Section 4.2, we can deform the contour Σ in (4.54) into a vertical contour going from $e^{-(\pi/2+\delta)i} \cdot \infty$ to $e^{(\pi/2+\delta)i} \cdot \infty$ on the left of Γ_α . Furthermore we can deform the closed contour Γ_α into the infinite contour Γ_0 . Thus the double integral formula (4.54) still holds with Σ and Γ_α replaced by $\Sigma_{-1/2}^\delta$ and Γ_0 respectively.

Comparing the integrands of K^L (1.7) and K^J (4.54) we see that the only difference is that in the latter there is an additional factor $\Gamma(w + c_2 + N + 1)/\Gamma(z + c_2 + N + 1)$. For z and w fixed and $N \rightarrow \infty$, this simplifies to

$$\frac{\Gamma(w + c_2 + N + 1)}{\Gamma(z + c_2 + N + 1)} = N^{(w-z)} \left(1 + \mathcal{O}(1/N) \right),$$

so the only difference in the present case is an additional factor of $N^{(w-z)}$. The additional factor is accounted for by the different scalings: $x \mapsto N^{-1/\theta}x$ for the Laguerre case, and $x \mapsto N^{-1-1/\theta}x$ for the Jacobi case.

In the case that $c_2 \notin \mathbb{Z}$, the contour Γ_α in (4.54) is infinite. We express $\Gamma_\alpha = \Gamma'_\alpha \cup \Gamma''_\alpha$ and Σ a contour from $e^{-(\pi/2+\delta)i} \cdot \infty$ to $e^{(\pi/2+\delta)i} \cdot \infty$ in between Γ'_α and Γ''_α as in the proof of Proposition 4.7. Furthermore, we deform Γ''_α into the infinite contour Γ_0 , and then take Σ as $\Sigma_{-1/2}^\delta$. We write

$$K^J(x, y) = K^{J,'}(x, y) + K^{J, ''}(x, y), \quad \text{where} \quad K^{J,*}(x, y) = \frac{1}{(2\pi i)^2} \frac{\tilde{h}(x)}{\tilde{h}(y)} \\ \times \oint_{\Sigma} dz \oint_{\Gamma_\alpha^*} dw \frac{x^{-z-1}y^w}{(z-w)} \frac{\Gamma(w + c_2 + N + 1)\Gamma(z + 1) \prod_{k=1}^N (z - \alpha_k)}{\Gamma(z + c_2 + N + 1)\Gamma(w + 1) \prod_{l=1}^N (w - \alpha_l)}, \quad * = ' \text{ or } ''.$$

By the argument in the $c_2 \in \mathbb{Z}$ case, we have that

$$\lim_{N \rightarrow \infty} N^{-1-1/\theta} K^{J, ''}(N^{-1-1/\theta}x, N^{-1-1/\theta}y) = \lim_{N \rightarrow \infty} N^{-1/\theta} K^L(N^{-1/\theta}x, N^{-1/\theta}y). \tag{5.10}$$

On the other hand, by estimating the factors of the integrand

$$x^{-z-1}\Gamma(z + 1) \prod_{k=1}^N (z - \alpha_k)\Gamma(z + c_2 + N + 1)^{-1}$$

by Stirling’s formula on $\Sigma_{-1/2}$ and

$$y^w \Gamma(w + c_2 + N + 1) \left(\Gamma(w + 1) \prod_{l=1}^N (w - \alpha_l) \right)^{-1}$$

on Γ'_α separately, we find that

$$\lim_{N \rightarrow \infty} N^{-1-1/\theta} K^{J, \prime}(N^{-1-1/\theta} x, N^{-1-1/\theta} y) = 0. \quad (5.11)$$

Thus we prove (5.9) by combining (5.10) and (5.11). \square

Acknowledgments. The work of PJF was supported by the Australian Research Council. The work of DW was partially supported by the start-up grant R-146-000-164-133. Thanks are also given to the Department of Mathematics, National University of Singapore, for hosting a visit during July 2014 when this work was begun. We thank Arno Kuijlaars and Lun Zhang for spotting misprints in an earlier version. We thank Dongzhou Huang for noticing that the $\theta \in (0, 1)$ case of Proposition 1.3 requires a deformation of Σ other than a vertical line.

References

- [1] Milton Abramowitz and Irene A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series, vol. 55, For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964. MR-MR0167642
- [2] Mark Adler, Pierre van Moerbeke, and Dong Wang, *Random matrix minor processes related to percolation theory*, *Random Matrices Theory Appl.* **2** (2013), no. 4, 1350008, 72. MR-3149438
- [3] Gernot Akemann, Jesper R. Ipsen, and Mario Kieburg, *Products of rectangular random matrices: Singular values and progressive scattering*, *Phys. Rev. E* **88** (2013), no. 5, 052118, 13.
- [4] N. Alexeev, F. Götze, and A. Tikhomirov, *Asymptotic distribution of singular values of powers of random matrices*, *Lith. Math. J.* **50** (2010), no. 2, 121–132. MR-2653641
- [5] Kazuhiko Aomoto and Kazumoto Iguchi, *On quasi-hypergeometric functions*, *Methods Appl. Anal.* **6** (1999), no. 1, 55–66, Dedicated to Richard A. Askey on the occasion of his 65th birthday, Part I. MR-1803881
- [6] Kazuhiko Aomoto and Kazumoto Iguchi, *Wu’s equations and quasi-hypergeometric functions*, *Comm. Math. Phys.* **223** (2001), no. 3, 475–507. MR-1866164
- [7] T. Banica, S. T. Belinschi, M. Capitaine, and B. Collins, *Free Bessel laws*, *Canad. J. Math.* **63** (2011), no. 1, 3–37. MR-2779129
- [8] C. W. J. Beenakker and B. Rejaei, *Nonlogarithmic repulsion of transmission eigenvalues in a disordered wire*, *Phys. Rev. Lett.* **71** (1993), 3689–3692.
- [9] Hari Bercovici and Vittorino Pata, *Stable laws and domains of attraction in free probability theory*, *Ann. of Math. (2)* **149** (1999), no. 3, 1023–1060, With an appendix by Philippe Biane. MR-1709310
- [10] M. Bertola, M. Gekhtman, and J. Szmigielski, *The Cauchy two-matrix model*, *Comm. Math. Phys.* **287** (2009), no. 3, 983–1014. MR-2486670
- [11] M. Bertola, M. Gekhtman, and J. Szmigielski, *Cauchy-Laguerre two-matrix model and the Meijer-G random point field*, *Comm. Math. Phys.* **326** (2014), no. 1, 111–144. MR-3162486
- [12] Marco Bertola and Thomas Bothner, *Universality conjecture and results for a model of several coupled positive-definite matrices*, *Comm. Math. Phys.* **337** (2015), no. 3, 1077–1141. MR-3339172
- [13] Alexei Borodin, *Biorthogonal ensembles*, *Nuclear Phys. B* **536** (1999), no. 3, 704–732. MR-1663328

- [14] Gaëtan Borot, Alice Guionnet, and Karol K. Kozłowski, *Large- N asymptotic expansion for mean field models with Coulomb gas interaction*, Int. Math. Res. Not. IMRN (2015), no. 20, 10451–10524. MR-3455872
- [15] L. Carlitz, *Problem 72-17, “Biorthogonal conditions for a class of polynomials”*, SIAM Rev. **15** (1973), 670–672. MR-0330557
- [16] W. A. Chat, *Biorthogonal conditions for a class of polynomials*, SIAM Rev. **14** (1972), no. 3, 494–495.
- [17] Dimitris Cheliotis, *Triangular random matrices and biorthogonal ensembles*, 2014, arXiv:1404.4730.
- [18] Tom Claeys and Stefano Romano, *Biorthogonal ensembles with two-particle interactions*, Nonlinearity **27** (2014), no. 10, 2419–2444. MR-3265719
- [19] Tom Claeys and Dong Wang, *Random matrices with equispaced external source*, Comm. Math. Phys. **328** (2014), no. 3, 1023–1077. MR-3201219
- [20] P. A. Deift, *Orthogonal polynomials and random matrices: a Riemann-Hilbert approach*, Courant Lecture Notes in Mathematics, vol. 3, New York University Courant Institute of Mathematical Sciences, New York, 1999. MR-MR1677884
- [21] Ioana Dumitriu and Alan Edelman, *Matrix models for beta ensembles*, J. Math. Phys. **43** (2002), no. 11, 5830–5847. MR-1936554
- [22] Ken Dykema and Uffe Haagerup, *DT-operators and decomposability of Voiculescu’s circular operator*, Amer. J. Math. **126** (2004), no. 1, 121–189. MR-2033566
- [23] P. J. Forrester, *Log-gases and random matrices*, London Mathematical Society Monographs Series, vol. 34, Princeton University Press, Princeton, NJ, 2010. MR-2641363
- [24] Peter J. Forrester and Dang-Zheng Liu, *Raney distributions and random matrix theory*, J. Stat. Phys. **158** (2015), no. 5, 1051–1082. MR-3313617
- [25] Peter J. Forrester, Dang-Zheng Liu, and Paul Zinn-Justin, *Equilibrium problems for Raney densities*, Nonlinearity **28** (2015), no. 7, 2265–2277. MR-3366643
- [26] Peter J. Forrester and Eric M. Rains, *Interpretations of some parameter dependent generalizations of classical matrix ensembles*, Probab. Theory Related Fields **131** (2005), no. 1, 1–61. MR-2105043
- [27] Wolfgang Gawronski, Thorsten Neuschel, and Dries Stivigny, *Jacobi polynomial moments and products of random matrices*, Proc. Amer. Math. Soc. **144** (2016), no. 12, 5251–5263. MR-3556269
- [28] Ronald L. Graham, Donald E. Knuth, and Oren Patashnik, *Concrete mathematics*, second ed., Addison-Wesley Publishing Company, Reading, MA, 1994, A foundation for computer science. MR-1397498
- [29] Adrien Hardy, *Average characteristic polynomials of determinantal point processes*, Ann. Inst. Henri Poincaré Probab. Stat. **51** (2015), no. 1, 283–303. MR-3300971
- [30] Norman L. Johnson, Samuel Kotz, and N. Balakrishnan, *Continuous univariate distributions. Vol. 2*, second ed., Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics, John Wiley & Sons, Inc., New York, 1995, A Wiley-Interscience Publication. MR-1326603
- [31] Mario Kieburg, Arno B. J. Kuijlaars, and Dries Stivigny, *Singular value statistics of matrix products with truncated unitary matrices*, Int. Math. Res. Not. IMRN (2016), no. 11, 3392–3424. MR-3556414
- [32] Heiner Kohler, *Exact diagonalization of 1D interacting spinless fermions*, J. Math. Phys. **52** (2011), no. 3, 032107, 24. MR-2814695
- [33] Joseph D. E. Konhauser, *Biorthogonal polynomials suggested by the Laguerre polynomials*, Pacific J. Math. **21** (1967), 303–314. MR-0214825
- [34] Arno B. J. Kuijlaars and Dries Stivigny, *Singular values of products of random matrices and polynomial ensembles*, Random Matrices Theory Appl. **3** (2014), no. 3, 1450011, 22. MR-3256862

- [35] Arno B. J. Kuijlaars and Lun Zhang, *Singular values of products of Ginibre random matrices, multiple orthogonal polynomials and hard edge scaling limits*, *Comm. Math. Phys.* **332** (2014), no. 2, 759–781. MR-3257662
- [36] Vadim B. Kuznetsov, Vladimir V. Mangazeev, and Evgeny K. Sklyanin, *Q-operator and factorised separation chain for Jack polynomials*, *Indag. Math. (N.S.)* **14** (2003), no. 3-4, 451–482. MR-2083086
- [37] Dang-Zheng Liu, Dong Wang, and Lun Zhang, *Bulk and soft-edge universality for singular values of products of Ginibre random matrices*, *Ann. Inst. Henri Poincaré Probab. Stat.* **52** (2016), no. 4, 1734–1762. MR-3573293
- [38] T. Lueck, H.-J. Sommers, and M. R. Zirnbauer, *Energy correlations for a random matrix model of disordered bosons*, *J. Math. Phys.* **47** (2006), no. 10, 103304, 24. MR-2268865
- [39] Yudell L. Luke, *The special functions and their approximations, Vol. I*, *Mathematics in Science and Engineering*, Vol. 53, Academic Press, New York-London, 1969. MR-0241700
- [40] H. C. Madhekar and N. K. Thakare, *Biorthogonal polynomials suggested by the Jacobi polynomials*, *Pacific J. Math.* **100** (1982), no. 2, 417–424. MR-669334
- [41] Wojciech Młotkowski, *Fuss-Catalan numbers in noncommutative probability*, *Doc. Math.* **15** (2010), 939–955. MR-2745687
- [42] Wojciech Młotkowski and Karol A. Penson, *Probability distributions with binomial moments*, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **17** (2014), no. 2, 1450014, 32. MR-3212684
- [43] K. A. Muttalib, *Random matrix models with additional interactions*, *J. Phys. A* **28** (1995), no. 5, L159–L164. MR-1327636
- [44] Thorsten Neuschel, *Plancherel–Rotach formulae for average characteristic polynomials of products of Ginibre random matrices and the Fuss-Catalan distribution*, *Random Matrices Theory Appl.* **3** (2014), no. 1, 1450003, 18. MR-3190209
- [45] Alexandru Nica and Roland Speicher, *Lectures on the combinatorics of free probability*, *London Mathematical Society Lecture Note Series*, vol. 335, Cambridge University Press, Cambridge, 2006. MR-2266879
- [46] A. Okounkov and G. Olshanski, *Shifted Jack polynomials, binomial formula, and applications*, *Math. Res. Lett.* **4** (1997), no. 1, 69–78. MR-1432811
- [47] Frank W. J. Olver, Daniel W. Lozier, Ronald F. Boisvert, and Charles W. Clark (eds.), *NIST handbook of mathematical functions*, U.S. Department of Commerce, National Institute of Standards and Technology, Washington, DC; Cambridge University Press, Cambridge, 2010, With 1 CD-ROM (Windows, Macintosh and UNIX). MR-2723248
- [48] Jack W. Silverstein, *The smallest eigenvalue of a large-dimensional Wishart matrix*, *Ann. Probab.* **13** (1985), no. 4, 1364–1368. MR-806232
- [49] H. M. Srivastava, *Some biorthogonal polynomials suggested by the Laguerre polynomials*, *Pacific J. Math.* **98** (1982), no. 1, 235–250. MR-644951
- [50] Richard P. Stanley, *Enumerative combinatorics. Vol. 2*, *Cambridge Studies in Advanced Mathematics*, vol. 62, Cambridge University Press, Cambridge, 1999, With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin. MR-MR1676282
- [51] Hale F. Trotter, *Eigenvalue distributions of large Hermitian matrices; Wigner’s semicircle law and a theorem of Kac, Murdock, and Szegő*, *Adv. in Math.* **54** (1984), no. 1, 67–82. MR-761763
- [52] Lun Zhang, *Local Universality in Biorthogonal Laguerre Ensembles*, *J. Stat. Phys.* **161** (2015), no. 3, 688–711. MR-3406705
- [53] Karol Życzkowski and Hans-Jürgen Sommers, *Truncations of random unitary matrices*, *J. Phys. A* **33** (2000), no. 10, 2045–2057. MR-1748745

Electronic Journal of Probability

Electronic Communications in Probability

Advantages of publishing in EJP-ECP

- Very high standards
- Free for authors, free for readers
- Quick publication (no backlog)
- Secure publication (LOCKSS¹)
- Easy interface (EJMS²)

Economical model of EJP-ECP

- Non profit, sponsored by IMS³, BS⁴, ProjectEuclid⁵
- Purely electronic

Help keep the journal free and vigorous

- Donate to the IMS open access fund⁶ (click here to donate!)
- Submit your best articles to EJP-ECP
- Choose EJP-ECP over for-profit journals

¹LOCKSS: Lots of Copies Keep Stuff Safe <http://www.lockss.org/>

²EJMS: Electronic Journal Management System <http://www.vtex.lt/en/ejms.html>

³IMS: Institute of Mathematical Statistics <http://www.imstat.org/>

⁴BS: Bernoulli Society <http://www.bernoulli-society.org/>

⁵Project Euclid: <https://projecteuclid.org/>

⁶IMS Open Access Fund: <http://www.imstat.org/publications/open.htm>