

## Research Article

### SPECIAL ISSUE: The Role of Below-Ground Processes in Mediating Plant Invasions

## Mutualism-disrupting allelopathic invader drives carbon stress and vital rate decline in a forest perennial herb

Nathan L. Brouwer\*, Alison N. Hale and Susan Kalisz

Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260, USA

**Received:** 30 July 2014; **Accepted:** 30 January 2015; **Published:** 27 February 2015

**Associate Editor:** Inderjit

**Citation:** Brouwer NL, Hale AN, Kalisz S. 2015. Mutualism-disrupting allelopathic invader drives carbon stress and vital rate decline in a forest perennial herb. *AoB PLANTS* 7: plv014; doi:10.1093/aobpla/plv014

**Abstract.** Invasive plants can negatively affect belowground processes and alter soil microbial communities. For native plants that depend on soil resources from root fungal symbionts (RFS), invasion could compromise their resource status and subsequent ability to manufacture and store carbohydrates. Herbaceous perennials that depend on RFS-derived resources dominate eastern North American forest understories. Therefore, we predict that forest invasion by *Alliaria petiolata*, an allelopathic species that produces chemicals that are toxic to RFS, will diminish plant carbon storage and fitness. Over a single growing season, the loss of RFS could reduce a plant's photosynthetic physiology and carbon storage. If maintained over multiple growing seasons, this could create a condition of carbon stress and declines in plant vital rates. Here we characterize the signals of carbon stress over a short timeframe and explore the long-term consequence of *Alliaria* invasion using *Maianthemum racemosum*, an RFS-dependent forest understory perennial. First, in a greenhouse experiment, we treated the soil of potted *Maianthemum* with fresh leaf tissue from either *Alliaria* or *Hesperis matronalis* (control) for a single growing season. *Alliaria*-treated plants exhibit significant overall reductions in total non-structural carbohydrates and have 17 % less storage carbohydrates relative to controls. Second, we monitored *Maianthemum* vital rates in paired experimental plots where we either removed emerging *Alliaria* seedlings each spring or left *Alliaria* at ambient levels for 7 years. Where *Alliaria* is removed, *Maianthemum* size and vital rates improve significantly: flowering probability increases, while the probability of plants regressing to non-flowering stages or entering prolonged dormancy are reduced. Together, our results are consistent with the hypothesis that disruption of a ubiquitous mutualism following species invasion creates symptoms of carbon stress for species dependent on RFS. Disruption of plant-fungal mutualisms may generally contribute to the common, large-scale declines in forest biodiversity observed in the wake of allelopathic invaders.

**Keywords:** Allelochemicals; *Alliaria petiolata*; carbon stress/carbon starvation; *Maianthemum racemosum*; mutualism disruption; root fungal symbiont; species invasion; vital rates.

\* Corresponding author's e-mail address: brouwern@gmail.com

Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

## Introduction

The majority of flowering plant species form mutualisms with root fungal symbionts (RFS) such as arbuscular mycorrhizal fungi (AMF; 74 % of angiosperms; Brundrett 2009) and dark septate endophytes (DSE;  $\geq 600$  species; Jumpponen and Trappe 1998). Arbuscular mycorrhizal fungi and DSE live inside plant roots and deploy hyphae outside the root that increase water, nitrogen, phosphorus and other soil nutrients' availability to their plant partner (Smith and Read 2008; Newsham 2011). The RFS receive a substantial fraction of the plant partner's fixed carbon (for AMF up to 20 %; Smith and Read 2008).

Recent work highlights how anthropogenic changes in the environment, such as invasion, can negatively affect mutualisms (Tylianakis et al. 2008; Kiers et al. 2010). Invasive species can impact belowground processes and directly or indirectly alter soil microbial communities, including RFS. Mechanisms through which belowground impacts can occur (summarized in part by Wolfe and Klironomos 2005) include alterations in the quality, quantity and timing of litter inputs and subsequent changes in soil nutrient status (reviewed by Ehrenfeld 2003), direct changes to soil nutrient status through novel nutrient fixation strategies by the invader (e.g. Vitousek and Walker 1989), mutualist degradation (Vogelsang and Bever 2009) and allelopathy (e.g. Callaway et al. 2008; Grove et al. 2012). Specifically, allelochemicals can act as novel weapons that are directly toxic to plants or act indirectly on their associated microbes (Callaway and Ridenour 2004; Weir et al. 2004).

The invasion of North American forests by *Alliaria petiolata* (Brassicaceae, garlic mustard) is an emerging model system for investigations of allelopathic effects on belowground processes (Rodgers et al. 2008a). This species produces a suite of allelochemicals (Vaughn and Berhow 1999; Cipollini and Gruner 2007) that are toxic to RFS (Roberts and Anderson 2001; Stinson et al. 2006; Koch et al. 2011) even at low concentrations (Callaway et al. 2008; Cantor et al. 2011). Field studies document that areas infested with *Alliaria* exhibit shifts in soil fungal community composition with frequent reductions in AMF species richness (Burke et al. 2011; Lankau 2011a; Lankau et al. 2014), declines in total soil hyphal abundances (Cantor et al. 2011; Koch et al. 2011) and changes in the within-root community of AMF-dependent plants (Burke 2008; Bongard et al. 2013). Together, these studies suggest that within *Alliaria*-invaded ecosystems the function of the mutualistic fungal community can be compromised and that these changes contribute to *Alliaria*'s invasive success.

Herbaceous perennials dominate the temperate forest understories that *Alliaria* invades and these species as a

group are typically highly- to obligately-dependent on RFS (Brundrett and Kendrick 1988; Whigham 2004). The fact that temperate forest soils are strongly resource limited (Whigham 2004; Gilliam 2014) likely drives the obligate nature of the relationship for many understory herbaceous perennials. Typically these species are slow growing (Gilliam 2014), exhibit high rates of RFS colonization (e.g. Brundrett and Kendrick 1988; Boerner 1990; Burke 2008) and have long-lived arbuscules (Brundrett and Kendrick 1990). Many also lack fine roots or root hairs (e.g. LaFrankie 1985) perhaps because their associated RFS hyphae fulfil this soil resource-gathering role. Since resources supplied by RFS are intimately tied to many plant metabolic functions (Schweiger et al. 2014), disruption of soil mutualisms is expected to severely limit the physiological rates of forest species (Hale et al. 2011). In the absence of RFS, plants generally exhibit reduced photosynthetic rates (Allen et al. 1981; Wright et al. 1998; Zhu et al. 2011) and subsequent carbon stress can curb their ability to carry out carbon-demanding functions such as growth (Lu and Koide 1994) and flowering (Koide et al. 1994).

Carbon stress is the reduction of a plant's pool of total non-structural carbohydrates (NSCs) (*sensu* Anderegg et al. 2012). In herbaceous perennials, chronic carbon stress can alter key vital rates including survival (Gremer and Sala 2013), flowering (Crone et al. 2009) and prolonged dormancy (Gremer et al. 2010). Invaders like *Alliaria* that alter the soil environment and essential RFS functions could induce carbon stress or 'carbon starvation' (*sensu* McDowell et al. 2008), ultimately diminishing the stability of populations of RFS-dependent native species.

Our prior experiments on the RFS-dependent understory perennial, *Maianthemum racemosum* (Ruscaceae, false Solomon's seal) confirm the dramatic physiological consequences of short-term RFS disruption by *Alliaria*'s allelochemicals. Key physiological traits including stomatal conductance, which is known to be highly dependent on RFS colonization (Augé et al. 2014), and photosynthetic rate both significantly declined in plants exposed to fresh *Alliaria* leaf litter (Hale et al. 2011). Soil respiration, to which fungi are the primary contributors (Anderson and Domsch 1975), was also reduced with *Alliaria* treatment. Importantly, in field plots invaded by *Alliaria* and in pot experiments with an *Alliaria* litter treatment, we demonstrated significant declines in the abundance of soil fungal hyphae relative to controls (37 % decline, Cantor et al. 2011; 29–38 % decline, A. N. Hale et al., submitted for publication). Together these data strongly support the idea that the observed physiological declines are driven by the inhibition of the RFS hyphal network in the soil (Hale et al. 2011).

Here we explore how the physiological stress of RFS-mutualism disruption in *Alliaria*-invaded forests could result in performance declines in an RFS-dependent forest perennial across two time scales. First, we ask: Given that *Alliaria*'s allelochemicals cause detectable shifts in the soil fungal community and alternative plant physiological rates, do they also cause declines in carbon storage in plants within a single growing season? In a greenhouse experiment we show that *Alliaria*-treated *Maianthemum* store significantly less carbon in their rhizome over one growing season relative to controls. Second, to determine the potential for short-term effects to scale up over time and affect population processes, we conducted a 7-year field experiment in an *Alliaria*-invaded forest in which *Alliaria* was weeded or left at ambient levels. We test whether *Maianthemum* exhibit lower growth rates consistent with carbon stress in the *Alliaria*-ambient plots. We also ask if *Alliaria* reduces size-based vital rates of *Maianthemum* and if so, how quickly these changes occur. We show that where *Alliaria* is present, *Maianthemum* have suppressed growth and vital rates relative to adjacent plot where *Alliaria* is removed.

## Methods

### Greenhouse study: assessing potential for carbon stress

The greenhouse study was conducted during the summer of 2010 in the greenhouse facilities at the University of Pittsburgh. In May, we obtained bare-root adult *Maianthemum* plants ( $N = 42$ ) from a native plant nursery (Prairie Moon Nursery, Winona, MN, USA). Rhizomes ranged in size from 6.7 to 39.7 g fresh weight. We potted each rhizome in a 3 : 1 mixture of autoclaved Fafard potting soil and Turface. We inoculated plants with RFS by adding 150 g of field soil collected from areas adjacent to *Maianthemum* plants at our experimental field site (see details below). Pots were then placed in the greenhouse and watered every 2–3 days for 1 month, allowing the plants to complete stem elongation and establish the RFS mutualism.

In June, we assigned each plant to either an *Alliaria* treatment or a control treatment. To control for potential differences in initial carbohydrate status due to differences in plant age and/or size (e.g. [Olano et al. 2006](#)), we stratified the randomized assignment of rhizomes into the treatments to ensure that mean rhizome mass was the same in the *Alliaria* and control treatments. Plants in the *Alliaria* treatment were then exposed to *Alliaria* allelochemicals by placing 25 g of fresh *Alliaria* leaf tissue collected from a population with a recent history of invasion (<20 years) on top of the soil. When these plants were watered, the glucosinolates leached out of

the *Alliaria* leaves and into the soil (A. N. Hale et al., submitted for publication). As in previous experiments ([Hale et al. 2011](#)), plants in the control treatment received 25 g of fresh *Hesperis matronalis* (dame's rocket; Brassicaceae) leaf tissue. Like *Alliaria*, *Hesperis* is an invasive mustard in eastern North America ([Leicht-Young et al. 2012](#)). While *Hesperis* produces some glucosinolates ([Larsen et al. 1992](#)), RFS hyphae and vesicles have been observed within its root system ([DeMars and Boerner 1995](#)), indicating that *Hesperis* chemicals are less toxic to RFS than *Alliaria*. In the field, the high mortality rates of *Alliaria* seedlings and rosettes throughout the year ([Davis et al. 2006](#)) and the mortality of adults in the summer ([Anderson et al. 1996](#)) likely result in a sustained supply of allelochemicals into the soil. Thus, we re-applied fresh leaf tissue in both treatments every 2 weeks until the end of August to simulate a season-long supply of *Alliaria* allelochemicals.

We destructively harvested plants three times during the growing season (9 July, 6 August and at senescence) to assess the effect of the treatments on the carbohydrate status. For the last time point, we classified plants as being senesced when 40 % of the leaf tissue had yellowed and photosynthetic rates were  $<1.0 \mu\text{mol m}^{-2} \text{s}^{-1}$ . Details of the leaf gas exchange protocol for *Maianthemum* can be found in [Hale et al. \(2011\)](#). To harvest the plants, we carefully clipped the shoot and roots away from the rhizome. We also stained the roots of a subset of plants per treatment following [Brundrett et al. \(1984\)](#) to confirm RFS colonization. We then weighed the rhizome and immediately flash-froze it in liquid nitrogen. We stored samples at  $-80^\circ\text{C}$  until they could be lyophilized and ground. We followed the protocol of [Zuleta and Sambucetti \(2001\)](#) to analyse rhizome inulin (storage carbohydrate) and sucrose (mobile carbohydrate) content via high-performance liquid chromatography (HPLC). [Note: Starch is not present in the rhizome of *Maianthemum* (A. N. Hale et al., submitted for publication).] In brief, a 0.03 g dried sample for each plant is boiled while stirring with a magnetic stir bar. Once samples cool to room temperature, they are filtered through a  $0.20 \mu\text{m}$  filter, and run on HPLC (Aminex HPX-87C anion-exchange column, deionized water at  $85^\circ\text{C}$  was set as the mobile phase with a flux rate of  $0.6 \text{ mL min}^{-1}$ ). Standards are used (inulin from dahlia tubers, Sigma-Aldrich; sucrose, Sigma-Aldrich) to confirm the identity of the sample peaks and to create standard curves to determine inulin and sucrose concentrations. Here, we express inulin and sucrose concentrations as a percentage of the HPLC dry sample mass. We also sum each plant's inulin and sucrose content to determine total NSC concentration (%).

To explore the effect of our treatments on rhizome carbohydrate status, we use a multivariate analysis of covariance (MANCOVA). Following a significant MANCOVA,

individual ANCOVA tests are conducted for inulin, sucrose and total NSC. For all models, we include harvest date as a main effect because rhizome carbohydrate concentration varies over the growing season in perennial herbs (e.g. Lapointe 1998; Wyka 1999; Kleijn et al. 2005). We also include initial plant mass as a covariate to account for differences in carbohydrate storage that are related to plant size/age (MANCOVA model: total NSC + inulin + sucrose = treatment + harvest date + initial plant mass; ANCOVA models: carbohydrate = treatment + harvest date + initial plant mass). We calculate least squares means and standard errors for all ANCOVA models with a significant ( $P < 0.05$ ) treatment effect. All analyses were conducted in SAS (v. 9.3, SAS Institute, Cary, NC, USA).

### Field study: measuring impacts on vital rates of native plant populations

**Study site.** Our experimental plots are located in a beech-maple forest in southwest Pennsylvania [Trillium Trail Nature Reserve (hereafter TT), Allegheny County, PA, USA: 40°52′01.40″N; 79°90′10.75″W] with a rich herbaceous perennial understory flora (Knight et al. 2009). Based on previous work at TT (Burke 2008) and other temperate deciduous forests (e.g. Brundrett and Kendrick 1988), we estimate that 73 % of TT herbaceous perennials are AMF-dependent (Hale et al. 2011). We detected *Alliaria* allelochemicals in the soil of TT in concentrations that are toxic to AMF spores in lab assays (Cantor et al. 2011). Additionally, we showed that in soils where *Alliaria* occurs at TT, the density of fungal hyphae is lower (Cantor et al. 2011) and the fungal community composition shifts (Burke et al. 2011) relative to paired, non-invaded areas. *Maianthemum* plants collected at TT are heavily colonized by RFS, but their intra-root AMF community is significantly altered where *Alliaria* is present (Burke 2008). These results motivate further investigation of mutualism disruption by *Alliaria* in understanding mechanisms driving native plant performance declines.

**Field experiment.** We collected data on naturally occurring individuals of *M. racemosum* within six 14 × 14 m plots in TT from 2003 through 2013. Our six plots are split in half longitudinally so that each contains two experimental treatments: *Alliaria* removal (= low or no allelochemicals) or *Alliaria* present at ambient levels (= allelochemicals present). Annual removal of *Alliaria* from half of each plot (i.e. a 14 × 7 m area) began in spring 2006, ~15 years after *Alliaria* became established at this site (L. Smith, pers. comm.) This time frame for TT invasion coincides with the estimated *Alliaria* invasion history in the region that indicates that this invader has been present locally for <25 years (Lankau et al. 2009). We remove *Alliaria* concurrent with the onset of emergence of the perennial

herb community. *Alliaria* individuals are removed as tiny seedlings, minimizing disturbance to the soil and other plants. Removed plants are discarded off site. In June of each year prior to *Alliaria* seed dispersal we erect a barrier at the border of the two treatments to block seed dispersal from the ambient into the *Alliaria* removal treatment. All *Maianthemum* plants emerging in the plots are permanently tagged and have annually been scored for individual size, stage (i.e. seedling, non-flowering, flowering and dormant) and deer browse status. Prior to initiation of the *Alliaria* removal treatment in 2006 there was no difference in *Alliaria* per cent cover between the plots ( $\chi^2 = 0.11$ ,  $P = 0.74$ ) or total per cent cover of all species ( $\chi^2 = 0.038$ ,  $P = 0.85$ ).

**Plant vital rates.** We assess the effect of *Alliaria* removal on *Maianthemum* growth and three vital rates: annual flowering frequency, retrogression of flowering plants to non-flowering the following year and the frequency of prolonged vegetative dormancy (Shefferson 2009). We test for differences using data collected prior to the implementation of the removal treatment (2003–06) and after the removal treatment began (2007–13). All models have the general form: response variable = treatment + year + treatment × year. To estimate differences in growth rate, we investigate the differences in average size between treatments for the initial cohort of plants first observed when the experiment began in 2003. The mean size of this cohort is estimated with a linear mixed model for each year since 2006 (Zuur et al. 2009). We model log(plant size) to improve normality of the residuals.

Annual flowering frequencies are modelled using a logistic mixed model. Retrogression frequencies were modelled without random effects for the years 2008–13 because of limited sample size. Our retrogression model, stated in terms of probability, is

$$\text{Pr}(\text{Not Flowering}_{\text{time } t} | \text{Flowered}_{\text{time } t-1} \text{ and Not dormant}_{\text{time } t}).$$

Our sample for retrogression was therefore set by the number of plants that flowered the previous year (time  $t - 1$ ) that emerged as either flowering or non-flowering the next year (time  $t$ ).

Growth and vital rate analyses are conducted in R 3.1.0 (R Development Core Team 2014) using the *lme4* package (Bates et al. 2014). To account for repeated measures and blocking effects, we include random intercepts for individual plants and pairs of treatments within a plot. For each response variable we test for significant differences between annual means using the *multcomp* package in R (Bretz et al. 2010). We test for the presence of a long-term trend since 2006 in each treatment mean by specifying a trend contrast (Rosenthal and Rosnow 1985; Gurevitch



and Chester 1986). All tests are planned contrasts so we do not correct for multiple comparisons. To further investigate trends in flowering frequencies, we also analyse these data using a two-level hierarchical model with time as a continuously varying main effect and year as a random effect.

Results of flowering and retrogression analyses are reported as effect sizes using odds ratios (OR) (Rita and Komonen 2008). Odds ratios have a lower bound of zero and no upper bound. Odds ratios of 1 indicate no difference between two treatments in the odds of an event happening. Statistical tests for OR therefore test whether they are different from 1. Odds ratios and their 95 % confidence intervals (CIs) are given in the text on their normal scale but graphed on a log scale to improve interpretation (*sensu* Galbraith 1988).

**Mark-recapture models.** We use mark-recapture models, a modified logistic regression approach (Kéry et al. 2005), to estimate the probability of prolonged vegetative dormancy. To test for pre-existing differences in dormancy rates, we conduct separate mark-recapture analyses of the 3 years prior to implementation of the removal treatment (2003–05) and the 7 years after the treatment began (2007–13). Mark-recapture results are assessed using the small sample size corrected information criteria AICc ( $AICc = AIC + 2k(k+1)/(n-k-1)$ , where  $k$  = the number of parameters and  $n$  = sample size) to rank the explanatory ability of different models (Anderson 2010). To summarize the data we also analyse the entire data set (2003–13) and calculate the mean difference in dormancy rates between treatments. We first calculate dormancy rates for each treatment in each year, calculate the difference between these means and average the differences for the pre- and post-treatment time periods. We use the delta method (Powell 2007) in the R package *msm* to combine multiple standard errors and construct 95 % CIs around our final effect size estimates. Mark-recapture models are run in the R package *marked* (Laake et al. 2013).

**Missing data due to herbivory.** Deer browse compromised our ability to gain information on some individuals. Deer preferentially browse flowering *Maianthemum* and flowering individuals are of larger size than non-flowering individuals (N. L. Brouwer and S. Kalisz, unpubl. data). Accordingly, in the cases where an individual was browsed before its reproductive status was determined during the 10 annual censuses ( $n = 103$  instances across 10 years), we assumed the browsed individual was flowering. Further, if browse occurred before an individual's size data was collected or size was otherwise unavailable, we used linear imputation (Gelman and Hill 2006) to estimate its


size (412 instances of size imputation out of 1481 total size records). Including imputed size data for the browsed plants prevents biasing our results against detecting a treatment effect (Hadfield 2008; Nakagawa and Freckleton 2008).

We imputed missing size data using estimates generated from multiple rounds of linear regression based on observed size data from the years prior to and after the missing data. We averaged these multiple estimates to arrive at a final imputed size estimate for each browsed individual. Linear regression models included all available covariates, including previous size, current status, treatment and reproductive output for flowering plants. We validated our imputations by comparing mean plant size and the overall size distribution in the population with and without imputed data [see Supporting Information—Table S1].

## Results

### Greenhouse study: assessing potential for carbon stress

All *M. racemosum* plants examined exhibit colonization by internal RFS structures. However, *Maianthemum*'s rhizome carbohydrates were significantly affected by the *Alliaria* treatment (MANCOVA; Roy's greatest root = 7.57,  $P = 0.002$ ), with plants in the *Alliaria* treatment experiencing a significant reduction in total NSC (Fig. 1; ANCOVA  $F_{1,36} = 7.31$ ,  $P = 0.01$ ). Specifically, plants treated with *Alliaria* stored, on average, 17 % less inulin relative



**Figure 1.** *Maianthemum racemosum* rhizome carbohydrate content (%) from *Alliaria* (yellow) and *Hesperis* (control; black) treatments in the greenhouse experiment. Total NSC content is shown in solid-coloured bars. Total NSC is a composite measure of stored sugars (inulin; bars with diagonal shading) and mobile sugars (sucrose; stippled bars). Values are least squares means from ANCOVAs  $\pm 1$  standard error. \* $P < 0.05$ ; \*\* $P < 0.005$ .

to plants in the *Hesperis* treatment (Fig. 1; ANCOVA  $F_{1,36} = 9.28$ ,  $P = 0.004$ ). While plants in the *Alliaria* treatment had fewer stored sugars, they had higher sucrose concentrations in their rhizomes compared with plants in the *Hesperis* treatment (Fig. 1; ANCOVA  $F_{1,36} = 12.88$ ,  $P = 0.001$ ). The increase in mobile sugars did not compensate for the dramatic difference in stored sugars between treatments as total NSC in the *Alliaria*-treated plants was 13 % lower than that of *Hesperis*-treated plants. Harvest date was not a significant predictor of total NSC, inulin or sucrose.

### Field study: impact on vital rates

**Growth.** Prior to implementation of the removal treatment, there was no difference in the mean size of plants in the initial 2003 cohort (Fig. 2;  $P = 0.55$ ). By 2013 plants in the removal treatment are significantly larger than those in the ambient *Alliaria* treatment (mean difference = 6.70 cm, SE = 2.96;  $P = 0.02$ ). There is a significant positive linear trend in size from 2006 to 2013 (trend contrast  $P = 0.0056$ ) in the *Alliaria* removal plots but no trend in the ambient plots ( $P = 0.91$ ).


**Flowering.** There is no significant difference in flowering probability across treatments for the first 6 years of the *Alliaria* removal (e.g. Fig. 3;  $P_{2006} = 0.65$ ,  $P_{2007} = 0.29$ ,  $P_{2008} = 0.42$ ). However, by 2012 the flowering probability is 'leaning' (*sensu* Tukey 1991) in the predicted direction (OR = 1.72, CI<sub>95%</sub> = 0.84–3.52,  $P = 0.14$ ) and by 2013 is significantly higher (OR = 1.96 CI<sub>95%</sub> = 1.0–3.87,  $P = 0.051$ ) in the removal treatment. Across all years

(2006–13) there is an increasing trend in flowering probability in the removal treatment (trend contrast  $P = 0.00008$ ) but no increase in the ambient treatment ( $P_{\text{trend}} = 0.57$ ).


Analyses using time as a continuous variable and year as a random effect confirmed that flowering frequencies diverged between the treatments (treatment  $\times$  time  $\chi^2 = 6.81$ ,  $P = 0.009$ ) with a significant positive linear trend in the removal treatment ( $\beta_{\text{removal} \times \text{time}} = 0.18$ , SE = 0.069) contrasted with evidence of a decrease in flowering probability in *Alliaria*-ambient plots ( $\beta_{\text{time}} = -0.10$ , SE = 0.072).

**Retgression.** The number of flowering individuals was too low in 2005 and 2006 to accurately estimate retrogression of flowering plants in 2006 and 2007. By 2011, there was evidence that removal-treatment plants were less likely to retrogress (OR = 0.28 CI<sub>95%</sub> = 0.052–1.57,  $P = 0.15$ ) and in 2012 they were significantly less likely to retrogress (OR = 0.14 CI<sub>95%</sub> = 0.021–0.96,  $P = 0.045$ ). There was a significant decreasing trend in retrogression in the removal treatment from 2008 until 2013 ( $P_{\text{trend}} = 0.011$ ) but no trend in the ambient treatment ( $P_{\text{trend}} = 0.90$ ).

**Dormancy.** Dormancy rates were highly variable between years, ranging from <10 to >30 %, but estimated to be lower in the *Alliaria* removal treatment in six out of 7 years [see Supporting Information—Table S2]. For years prior to the implementation of the *Alliaria* removal treatment (2003–06) the best-ranked model contains only a year effect (Table 1) while for models of post-treatment years



**Figure 2.** Effect of *Alliaria* on plant size of *Maianthemum* marked in the initial 2003 survey of the field experiment. (A) Mean difference (effect size) in plant size between *Alliaria* in ambient and removal treatments. (B) Annual mean plant sizes in both treatments and ANOVA trend contrasts. Error bars represent  $\pm 95\%$  CIs. Asterisk indicates a significant difference in plant size between the two treatments ( $P < 0.05$ ). Size data were not available for 2008 and 2009.



**Figure 3.** Effect of *Alliaria* on *Maianthemum* flowering frequency. (A) Mean difference (effect size, ES) in flowering frequency in *Alliaria*-ambient and removal plots. Effect size is expressed as an OR and plotted on the log scale. (B) Annual mean flowering frequencies for both treatments and ANOVA trend contrasts. Error bars represent  $\pm 95\%$  CIs. Asterisk indicates a significant effect of *Alliaria* removal ( $P < 0.05$ ).

**Table 1.** Ranking of mark-recapture models testing the effects of *Alliaria* removal on prolonged vegetative dormancy. Three sets of models were run over different time periods during the study: Set 1: years before *Alliaria* removal began (Pre-treatment); Set 2: years after the annual weeding treatment was initiated (post-treatment) and Set 3: all years. *N*, number of plants tracked over each time period; *K*, number of parameters in a model; Ln(lik), log likelihood. To calculate the mean pre-treatment and post-treatment effect size (Fig. 5) we used the parameters from the ‘Removal  $\times$  Year’ model in the ‘All years’ model Set 3.


Set	Period	Model	<i>N</i>	<i>K</i>	AICc	$\Delta$ AICc	Ln(lik)
1	Pre- <i>Alliaria</i> removal (2003–06)	Year	158	5	452.8	0.00	–216.21
		Removal + Year		6	454.6	1.74	–215.00
		Removal $\times$ Year		9	466.2	11.59	–214.47
2	Post- <i>Alliaria</i> removal (2007–13)	Removal + Year	210	9	1166.4	0.00	–564.73
		Year		8	1172.4	6.03	–569.84
		Removal $\times$ Year		15	1187.2	14.76	–562.34
3	All years (2003–13)	Removal + Year	236	12	1646.3	0.00	–798.46
		Year		11	1652.5	6.23	–803.68
		Removal $\times$ Year		21	1680.3	27.74	–795.98

(2007–13) and the entire dataset (2003–13) the best models contain an effect of *Alliaria* removal, indicating that dormancy rates were typically lower in this treatment. There was an initially large difference in dormancy rates between plots that would be allocated to the two treatments in the first year of the study [see Supporting Information—Table S2], potentially resulting in the model of the pre-treatment years containing an *Alliaria* removal effect (AICc = 454.6) ranked almost as high as a year-only model (AICc = 452.8). However, since the year-only model has a lower AICc and fewer parameters, the larger model is not considered competitive (Arnold 2010). Moreover, in the other two pre-treatment years (2004 and 2005), there is no difference between dormancy estimates

[see Supporting Information—Table S2]. The results of model selection are reinforced by the calculation of average effect sizes for the period prior to *Alliaria* removal and after removal (Fig. 5). Prior to removal there is no significant difference between dormancy rates (ES = –0.05, CI<sub>95%</sub> = –0.13–0.03) but after removal dormancy rates are  $\sim 7\%$  lower than in the *Alliaria*-ambient treatments (ES = –0.069, CI<sub>95%</sub> = –0.12 to –0.2).

## Discussion


To our knowledge this is the first study to explore the connections between an allelopathic invasive species’ impacts on the soil biotic environment and changes in



**Figure 4.** Effect of *Alliaria* on *Maianthemum* retrogression from flowering to non-flowering. (A) Annual mean difference in retrogression frequency (ES) in *Alliaria*-ambient and removal plot. Effect size is expressed as an OR and plotted on the log scale. (B) Mean retrogression frequencies in both treatments and ANOVA trend contrasts. Error bars represent  $\pm$  95 % CIs. Asterisk indicates a significant effect of *Alliaria* removal ( $P < 0.05$ ). Retrogression is calculated conditional on a plant being observed above-ground and not dormant. Sample sizes for 2006 and 2007 were insufficient for vital rate calculation.

individual plants' carbon status and vital rates. The results presented here in conjunction with prior studies substantiate multiple steps in a physiologically based causal pathway between invasion and population-level impacts on native plants. Our prior work demonstrates that *Alliaria* treatment of soil around *Maianthemum* reduces the density of soil fungal hyphae (A. N. Hale et al., submitted for publication) and plant photosynthetic rates (Hale et al. 2011). Here, our results demonstrate that treatment with *Alliaria* across the entire growing season results in negative effects on season-long carbon storage (Fig. 1). Relative to control plants, *Maianthemum* exposed to *Alliaria* stored 17 % less inulin in their rhizomes and experienced an overall reduction in total NSCs at the end of the season. Stomatal conductance modulates carbon fixation and is a key physiological rate affected by *Alliaria* exposure (Hale et al. 2011). Interestingly, a recent meta-analysis (Augé et al. 2014) comparing the effects of AMF inoculation on stomatal conductance ( $g_s$ ) in field vs. greenhouse studies indicates that greenhouse experiments have smaller effect sizes than field studies. Thus, our carbon storage results are likely conservative estimates of the carbon impacts of mutualism disruption in the field.

Over time, chronic exposure to *Alliaria* was predicted to compound this carbon deficit and affect plant growth and vital rates. Results from our long-term field study of *Alliaria* removal are consistent with this prediction. Individual aboveground plant size (Fig. 2) and multiple carbon-intensive and size-dependent vital rates (Figs 3–5)



**Figure 5.** Effect size of *Alliaria* removal on the frequency of prolonged vegetative dormancy in *Maianthemum* before (2003–06; yellow) and after the treatment began (2007–13; black). Calculated with mark-recapture models; error bars represent  $\pm$  95 % CIs. Asterisk indicates a significant effect of *Alliaria* removal ( $P < 0.05$ ).

are positively affected in *Alliaria* removal relative to *Alliaria*-ambient plots.

Other experimental studies where *Alliaria* and native plants are grown together in pots (Meekins and McCarthy 1999; Wixted and McGraw 2010; Lankau 2012; Smith and Reynolds 2014) or in the field (McCarthy 1997; Carlson and Gorchov 2004; Cipollini et al. 2008; Lankau 2011b) also find negative effects of *Alliaria* on native species. Competition, direct allelopathic phytotoxicity and



allelopathic RFS-mutualism disruption are all mechanisms that could contribute to these results. Our greenhouse experiment adds support to the idea that it is *Alliaria*'s disruption of key belowground mutualists (RFS) rather than competition or direct phytotoxicity that accounts for its success as an invader. Below we discuss the general support or lack thereof for the likelihood of all three mechanisms.

### Competition

We are aware of only two studies that have attempted to quantify reciprocal competition between *Alliaria* and focal plants. These pot studies found that *Alliaria* was equal to or weaker in competitive ability than three of four species tested (Meekins and McCarthy 1999; Leicht-Young et al. 2012). However, these studies are problematic in that they cannot separate competition from phytotoxicity or mutualism disruption. Bossdorf et al. (2004) found that *Alliaria* individuals from the native range outcompete *Alliaria* plants from the invaded range, supporting the hypothesis that invasive *Alliaria* express a different trade-off relative to their source populations. Invasive *Alliaria* are armed with novel allelochemical weapons but have evolved to be less competitive (Bossdorf et al. 2004). Further, field experiments demonstrate that native competitors can suppress *Alliaria* performance and abundance when the natives are not experiencing overabundant herbivore pressure (Eschtruth and Battles 2009a), as deer preferentially consume native plants and facilitate the high population growth and spread of *Alliaria* (Kalisz et al. 2014). In experimental studies that exclude deer from invaded sites, *Alliaria* abundance rapidly declines (Eschtruth and Battles 2009b; Knight et al. 2009; Kalisz et al. 2014). In total, these results underscore the widely held view that *Alliaria* is a relatively poor competitor (Rodgers et al. 2008a).

### Direct phytotoxicity

Glucosinolates are known antimicrobial chemicals produced by members of the mustard family as defences against pathogens (Tierens et al. 2001). While *Alliaria*'s allelochemicals can be inhibitory to germinating seeds and inhibit new seedling root growth (lettuce and radish seed experiments: Vaughn and Berhow 1999; Roberts and Anderson 2001; Pisula and Meiners 2010; *Impatiens* and *Viola* seed experiments: Prati and Bossdorf 2004; Barto et al. 2010; Cipollini and Flint 2013), to our knowledge direct toxicity of *Alliaria* on mature plant tissues has never been demonstrated. *Alliaria* invades forest understories dominated by adult perennial plants dependent on RFS. The direct effect of allelochemicals is inversely proportional to target plant density or biomass (Weidenhamer 2006). Single-celled fungal spores and thin fungal hyphae

should be much more susceptible to *Alliaria* allelochemicals than mature plant tissues. Thus, while we cannot rule out direct phytotoxic effects of *Alliaria* on adult *Maianthemum* performance in our field or greenhouse experiments, a direct allelochemical effect is likely of small magnitude relative to indirect effects on RFS.

### RFS-mutualism disruption

Mounting evidence shows that *Alliaria* can exert potent indirect effects on plants by suppressing RFS. Glucosinolates, like those produced by *Alliaria*, have a short half-life in the soil (<15 h; Gimsing et al. 2006). Yet, native plants grown in soils conditioned by *Alliaria*, treated with *Alliaria* tissue extracts, or collected from *Alliaria*-invaded sites all express reduced growth (Stinson et al. 2006; Callaway et al. 2008; Wolfe et al. 2008) despite the fact that the volatile allelochemicals were likely no longer present. Importantly, these studies demonstrate that *Alliaria* impacts are similar in magnitude to soil sterilization and that experimental soils result in lower colonization of roots by mycorrhizae (Stinson et al. 2006; Callaway et al. 2008; Wolfe et al. 2008). Finally, *Maianthemum* plants treated with *Alliaria* retain RFS structures internal to their roots, while exhibiting significant declines in soil hyphae (A. N. Hale et al., submitted for publication). Together these experiments provide strong support for RFS-mutualism disruption and that its effects are of large magnitude relative to competition or direct phytotoxicity.

Mechanistically, our working model linking RFS-mutualism disruption to carbon stress is based on the following premises: If *Alliaria*'s allelochemicals destroy the hyphal network, yet the normally long-lived internal structures (Brundrett and Kendrick 1990) remain intact, then we would predict that the plant would increase carbon allocation to its RFS to provision the regrowth of the soil hyphal network, resulting in significant carbon stress for the plant. Loss of the hyphal network severely limits available soil nutrients and water to the plant (Newsham 2011; Augé et al. 2014). As a result, the plants photosynthesize less (Hale et al. 2011) and fix less carbon (NSC; Fig. 1). With this limited carbon pool, we suggest that plants may maintain concentrations of mobile sugars in the rhizome and roots to re-establish a functional RFS hyphal network that is repeatedly destroyed by our application of fresh *Alliaria* tissue. While our results are consistent with this working model (e.g. we observe greater sucrose concentrations in the rhizome of *Alliaria* vs. *Hesperis*-treated plants (Fig. 1)), additional experiments are needed to fully explore this hypothesis.

We note that the effects of allelopathic mutualism disruption by *Alliaria* could be amplified by additional factors. Like other invasive species of deciduous forests (Ehrenfeld et al. 2001; Poulette and Arthur 2012; Smith and Reynolds 2012; Kuebbing et al. 2014; Schuster and

Dukes 2014), *Alliaria* can affect multiple components of the soil environment. *Alliaria* increases soil nutrient availability (Rodgers et al. 2008b), litter decomposition rates and nitrogen loss (Ashton et al. 2005). Since the RFS community in general (Van Diepen et al. 2011) and specific RFS–plant interactions (e.g. Klironomos 2002) are sensitive to soil conditions, multiple invader-mediated changes to the soil environment could magnify the impacts of allelopathic RFS–mutualism disruption. These diverse and widespread consequences of invasive species for soil environments and RFS communities are alarming given the potentially central role RFS and other microbes play in the diversity, productivity and functioning of plant communities (van der Heijden et al. 2008).

Our greenhouse study indicates that *Maianthemum* carbon storage declines significantly in response to *Alliaria* treatment in just one growing season. In contrast, we observe a relatively slow recovery of individual size, growth and vital rates following *Alliaria* removal in our field study. The predicted significant trends indicative of recovery (Figs 2–4) emerged after a few years of *Alliaria* removal while significant differences within the single-year comparisons were not seen until ~6–7 years post removal (2012 or 2013). Two, non-mutually exclusive mechanisms could underlie this lag. First, the lag could be due to *Maianthemum*'s habit (LaFrankie 1985). In general, forest understory herbaceous perennials are light-limited, slow-growing, long-lived species (Whigham 2004) with slow responses to perturbation (Morris et al. 2008). Our data are consistent with the idea that following *Alliaria* removal, *Maianthemum* may take multiple years to re-gain sufficient carbon stores to allow size growth, sustain flowering and maintain low dormancy rates. Second, the observed lag in *Maianthemum* vital rate responses may be due to slow recovery of the RFS soil community following *Alliaria* removal, a phenomenon observed by Anderson et al. (2010) and Lankau et al. (2014). If populations of beneficial RFS have gone locally extinct and low dispersal distance limits RFS re-colonization (Rout and Callaway 2012), then the observed time lag of *Maianthemum* could be due to the low abundance of effective fungal partners. Given the reciprocal obligate dependence of AMF and forest herbaceous perennial plants, declines in the native understory community may drive reciprocal declines in the RFS soil community (Lankau et al. 2014).

## Conclusions

Increases in invasive species are generally correlated with declines in native biodiversity (e.g. Butchart et al. 2010). However, the mechanistic underpinnings leading to native population collapse are rarely understood yet are the subject of numerous studies and invasion hypotheses

(Levine et al. 2003; Hulme et al. 2013). The disruption of plant soil feedbacks and root fungal symbioses are common aspects of plant invasions (i.e. Grove et al. 2012; Meinhardt and Gehring 2012; Ruckli et al. 2014; Shannon et al. 2014). As suggested by Hale and Kalisz (2012), chronic RFS–mutualism disruption could act as the first step in native plant biodiversity loss. In our system, the disruption of RFS by an allelopathic invader appears to begin a downward spiral in the physiological function (Hale et al. 2011), carbon status (Fig. 1) and ultimately vital rates (Figs 2–5) of a common native forest plant. Loss of these critical belowground mutualisms may be the proximate cause of plant mortality that is instead attributed to second-order effects (e.g. drought or herbivory) that are easier to observe (*sensu* McDowell 2011). Additional studies in invaded communities that explore the links between plant physiology, carbon allocation and population demographic performance are needed to determine the generality of these results. Mutualism disruption may be a widespread mechanism that helps explain how invasive species can cause large-scale changes to forest biodiversity observed in the wake of invasion (e.g. Rodgers et al. 2008a).

## Sources of Funding

Funding was supplied by a United States National Science Foundation award DEB-0958676 to S.K., a NSF predoctoral fellowship to N.L.B. and a Phipps Conservatory Botany-in-Action award and an Andrew K. Mellon predoctoral fellowship to A.N.H.

## Contributions by the Authors

S.K. conceived, designed, implemented and led data collection of the field experiment and assisted with the conception, design and implementation of the greenhouse experiment. A.N.H. designed, implemented and analysed data from the greenhouse experiment and assisted in data collected for the field experiment from 2008 to 2011. N.L.B. managed and analysed data from the field experiment and assisted with data collection for the field experiment since 2010. All three authors collaborated on the conception and writing of this article.

## Conflicts of Interest Statement

None declared.

## Acknowledgements

We thank the members of the Kalisz lab for field, lab and greenhouse assistance, and two anonymous reviewers whose detailed suggestions improved the manuscript.

## Supporting Information

The following additional information is available in the online version of this article –

**Table S1.** Validation of imputed *Maianthemum* size data from field experiment. Original and imputed size data are compared using *t*-tests and Kolmogorov–Smirnov tests.

**Table S2.** Estimated frequency of prolonged vegetative dormancy of *Maianthemum* from field experiment using a Mark-Recapture model.

## Literature Cited

- Allen MF, Smith WK, Moore TS Jr, Christensen M. 1981. Comparative water relations and photosynthesis of mycorrhizal and non-mycorrhizal *Bouteloua gracilis* H.B.K. Lag ex steud. *New Phytologist* **88**:683–693.
- Anderegg WRL, Berry JA, Smith DD, Sperry JS, Anderegg LDL, Field CB. 2012. The roles of hydraulic and carbon stress in a widespread climate-induced forest die-off. *Proceedings of the National Academy of Sciences of the USA* **109**:233–237.
- Anderson DR. 2010. *Model based inference in the life sciences: a primer on evidence*. New York: Springer.
- Anderson JPE, Domsch KH. 1975. Measurement of bacterial and fungal contributions to respiration of selected agricultural and forest soils. *Canadian Journal of Microbiology* **21**:314–322.
- Anderson RC, Dhillon SS, Kelley TM. 1996. Aspects of the ecology of an invasive plant, garlic mustard (*Alliaria petiolata*), in central Illinois. *Restoration Ecology* **4**:181–191.
- Anderson RC, Anderson MR, Bauer JT, Slater M, Herold J, Baumhardt P, Borowicz V. 2010. Effect of removal of garlic mustard (*Alliaria petiolata*, Brassicaceae) on arbuscular mycorrhizal fungi inoculum potential in forest soils. *The Open Ecology Journal* **3**:41–47.
- Arnold TW. 2010. Uninformative parameters and model selection using Akaike's Information Criterion. *The Journal of Wildlife Management* **74**:1175–1178.
- Ashton IW, Hyatt LA, Howe KM, Gurevitch J, Lerdau MT. 2005. Invasive species accelerate decomposition and litter nitrogen loss in a mixed deciduous forest. *Ecological Applications* **15**:1263–1272.
- Augé R, Toler HD, Saxton AM. 2014. Arbuscular mycorrhizal symbiosis alters stomatal conductance of host plants more under drought than under amply watered conditions: a meta-analysis. *Mycorrhiza* **25**:13–24.
- Barto EK, Friese C, Cipollini D. 2010. Arbuscular mycorrhizal fungi protect a native plant from allelopathic effects of an invader. *Journal of Chemical Ecology* **36**:351–360.
- Bates D, Maechler M, Bolker B, Walker S. 2014. *lme4: linear mixed-effect models using Eigen and S4*. R package version 1.1-7. <http://CRAN.R-project.org/package=lme4>.
- Boerner REJ. 1990. Role of mycorrhizal fungus origin in growth and nutrient uptake by *Geranium robertianum*. *American Journal of Botany* **77**:483–489.
- Bongard CL, Fulthorpe RR, Bongard CL, Fulthorpe RR. 2013. Invasion by two plant species affects fungal root colonizers. *Ecological Restoration* **31**:253–263.
- Bossdorf O, Prati D, Auge H, Schmid B. 2004. Reduced competitive ability in an invasive plant. *Ecology Letters* **7**:346–353.
- Bretz F, Hothorn T, Westfall P. 2010. *Multiple comparisons using R*. London: CRC Press.
- Brundrett MC. 2009. Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. *Plant and Soil* **320**:37–77.
- Brundrett MC, Kendrick B. 1988. The mycorrhizal status, root anatomy, and phenology of plants in a sugar maple forest. *Canadian Journal of Botany* **66**:1153–1173.
- Brundrett MC, Kendrick B. 1990. The roots and mycorrhizas of herbaceous woodland plants. I. Quantitative aspects of morphology. *New Phytologist* **114**:457–468.
- Brundrett MC, Piché Y, Peterson RL. 1984. A new method for observing the morphology of vesicular-arbuscular mycorrhizae. *Canadian Journal of Botany* **62**:2128–2134.
- Burke DJ. 2008. Effects of *Alliaria petiolata* (garlic mustard; Brassicaceae) on mycorrhizal colonization and community structure in three herbaceous plants in a mixed deciduous forest. *American Journal of Botany* **95**:1416–1425.
- Burke DJ, Weintraub MN, Hewins CR, Kalisz S. 2011. Relationship between soil enzyme activities, nutrient cycling and soil fungal communities in a northern hardwood forest. *Soil Biology and Biochemistry* **43**:795–803.
- Butchart SHM, Walpole M, Collen B, van Strien A, Scharlemann JPW, Almond REA, Baillie JEM, Bomhard B, Brown C, Bruno J, Carpenter KE, Carr GM, Chanson J, Chenery AM, Csirke J, Davidson NC, Dentener F, Foster M, Galli A, Galloway JN, Genovesi P, Gregory RD, Hockings M, Kapos V, Lamarque JF, Leverington F, Loh J, McGeoch MA, McRae L, Minasyan A, Morcillo MH, Oldfield TEE, Pauly D, Quader S, Revenga C, Sauer JR, Skolnik B, Spear D, Stanwell-Smith D, Stuart SN, Symes A, Tierney M, Tyrrell TD, Vié JC, Watson R. 2010. Global biodiversity: indicators of recent declines. *Science* **328**:1164–1168.
- Callaway RM, Ridenour WM. 2004. Novel weapons: invasive success and the evolution of increased competitive ability. *Frontiers in Ecology and the Environment* **2**:436–443.
- Callaway RM, Cipollini D, Barto K, Thelen GC, Hallett SG, Prati D, Stinson K, Klironomos J. 2008. Novel weapons: invasive plant suppresses fungal mutualists in America but not in its native Europe. *Ecology* **89**:1043–1055.
- Cantor A, Hale A, Aaron J, Traw MB, Kalisz S. 2011. Low allelochemical concentrations detected in garlic mustard-invaded forest soils inhibit fungal growth and AMF spore germination. *Biological Invasions* **13**:3015–3025.
- Carlson AM, Gorchoff DL. 2004. Effects of herbicide on the invasive biennial *Alliaria petiolata* (Garlic Mustard) and initial responses of native plants in a southwestern Ohio forest. *Restoration Ecology* **12**:559–567.
- Cipollini D, Gruner B. 2007. Cyanide in the chemical arsenal of garlic mustard, *Alliaria petiolata*. *Journal of Chemical Ecology* **33**:85–94.
- Cipollini KA, Flint WN. 2013. Comparing allelopathic effects of root and leaf extracts of invasive *Alliaria petiolata*, *Lonicera maackii*, and *Ranunculus ficaria* on germination of three native woodland plants. *Ohio Journal of Science* **112**:37–43.
- Cipollini KA, McClain GY, Cipollini D. 2008. Separating above- and belowground effects of *Alliaria petiolata* and *Lonicera maackii* on the performance of *Impatiens capensis*. *The American Midland Naturalist* **160**:117–128.



- Crone EE, Miller E, Sala A. 2009. How do plants know when other plants are flowering? Resource depletion, pollen limitation and mast-seeding in a perennial wildflower. *Ecology Letters* **12**:1119–1126.
- Davis AS, Landis DA, Nuzzo V, Blossey B, Gerber E, Hinz HL. 2006. Demographic models inform selection of biocontrol agents for garlic mustard (*Alliaria petiolata*). *Ecological Applications* **16**: 2399–2410.
- DeMars BG, Boerner REJ. 1995. Arbuscular mycorrhizal development in three crucifers. *Mycorrhiza* **5**:405–408.
- Ehrenfeld JG. 2003. Effects of exotic plant invasions on soil nutrient cycling processes. *Ecosystems* **6**:503–523.
- Ehrenfeld JG, Kourtev P, Huang W. 2001. Changes in soil functions following invasions of exotic understory plants in deciduous forests. *Ecological Applications* **11**:1287–1300.
- Eschtruth AK, Battles JJ. 2009a. Assessing the relative importance of disturbance, herbivory, diversity, and propagule pressure in exotic plant invasion. *Ecological Monographs* **79**:265–280.
- Eschtruth AK, Battles JJ. 2009b. Acceleration of exotic plant invasion in a forested ecosystem by a generalist herbivore. *Conservation Biology* **23**:388–399.
- Galbraith RF. 1988. A note on graphical presentation of estimated odds ratios from several clinical trials. *Statistics in Medicine* **7**: 889–894.
- Gelman A, Hill J. 2006. *Data analysis using regression and multilevel/hierarchical models*. Cambridge: Cambridge University Press.
- Gilliam F. 2014. *The herbaceous layer in forests of eastern North America*. Oxford: Oxford University Press.
- Gimsing AL, Sorensen JC, Tovgaard L, Jorgensen AMF, Hansen HCB. 2006. Degradation kinetics of glucosinolates in soil. *Environmental Toxicology and Chemistry* **25**:2038–2044.
- Gremer JR, Sala A. 2013. It is risky out there: the costs of emergence and the benefits of prolonged dormancy. *Oecologia* **172**:937–947.
- Gremer JR, Sala A, Crone EE. 2010. Disappearing plants: why they hide and how they return. *Ecology* **91**:3407–3413.
- Grove S, Haubensak KA, Parker IM. 2012. Direct and indirect effects of allelopathy in the soil legacy of an exotic plant invasion. *Plant Ecology* **213**:1869–1882.
- Gurevitch J, Chester ST Jr. 1986. Analysis of repeated measures experiments. *Ecology* **67**:251–255.
- Hadfield JD. 2008. Estimating evolutionary parameters when viability selection is operating. *Proceedings of the Royal Society B: Biological Sciences* **275**:723–734.
- Hale AN, Kalisz S. 2012. Perspectives on allelopathic disruption of plant mutualisms: a framework for individual- and population-level fitness consequences. *Plant Ecology* **213**:1991–2006.
- Hale AN, Tonsor SJ, Kalisz S. 2011. Testing the mutualism disruption hypothesis: physiological mechanisms for invasion of intact perennial plant communities. *Ecosphere* **2**:art110.
- Hulme PE, Pysěk P, Jarošík V, Pergl J, Schaffner U, Vilà M. 2013. Bias and error in understanding plant invasion impacts. *Trends in Ecology and Evolution* **28**:212–218.
- Jumpponen A, Trappe JM. 1998. Dark septate endophytes: a review of facultative biotrophic root-colonizing fungi. *New Phytologist* **140**:295–310.
- Kalisz S, Spigler RB, Horvitz CC. 2014. In a long-term experimental demography study, excluding ungulates reversed invader's explosive population growth rate and restored natives. *Proceedings of the National Academy of Sciences of the USA* **111**: 4501–4506.
- Kéry M, Gregg KB, Schaub M. 2005. Demographic estimation methods for plants with unobservable life-states. *Oikos* **108**:307–320.
- Kiers ET, Palmer TM, Ives AR, Bruno JF, Bronstein JL. 2010. Mutualisms in a changing world: an evolutionary perspective. *Ecology Letters* **13**:1459–1474.
- Kleijn D, Treier UA, Müller-Schärer H. 2005. The importance of nitrogen and carbohydrate storage for plant growth of the alpine herb *Veratrum album*. *New Phytologist* **166**:565–575.
- Klironomos JN. 2002. Feedback with soil biota contributes to plant rarity and invasiveness in communities. *Nature* **417**:67–70.
- Knight TM, Dunn JL, Smith LA, Davis J, Kalisz S. 2009. Deer facilitate invasive plant success in a Pennsylvania forest understory. *Natural Areas Journal* **29**:110–116.
- Koch AM, Antunes PM, Barto EK, Cipollini D, Mummey DL, Klironomos JN. 2011. The effects of arbuscular mycorrhizal (AM) fungal and garlic mustard introductions on native AM fungal diversity. *Biological Invasions* **13**:1627–1639.
- Koide RT, Shumway DL, Mabon SA. 1994. Mycorrhizal fungi and reproduction of field populations of *Abutilon theophrasti* Medic. (Malvaceae). *New Phytologist* **126**:123–130.
- Kuebbing SE, Classen AT, Simberloff D. 2014. Two co-occurring invasive woody shrubs alter soil properties and promote subdominant invasive species. *Journal of Applied Ecology* **51**:124–133.
- Laake JL, Johnson DS, Conn PB. 2013. *marked*: an R package for maximum likelihood and Markov Chain Monte Carlo analysis of capture–recapture data. *Methods in Ecology and Evolution* **4**:885–890.
- LaFrankie JV Jr. 1985. Morphology, growth, and vasculature of the sympodial rhizome of *Smilacina racemosa* (Liliaceae). *Botanical Gazette* **146**:534–554.
- Lankau RA. 2011a. Resistance and recovery of soil microbial communities in the face of *Alliaria petiolata* invasions. *New Phytologist* **189**:536–548.
- Lankau RA. 2011b. Interpopulation variation in allelopathic traits informs restoration of invaded landscapes. *Evolutionary Applications* **5**:270–282.
- Lankau RA. 2012. Coevolution between invasive and native plants driven by chemical competition and soil biota. *Proceedings of the National Academy of Sciences of the USA* **109**: 11240–11245.
- Lankau RA, Nuzzo V, Spyreas G, Davis AS. 2009. Evolutionary limits ameliorate the negative impact of an invasive plant. *Proceedings of the National Academy of Sciences of the USA* **106**: 15362–15367.
- Lankau RA, Bauer JT, Anderson MR, Anderson RC. 2014. Long-term legacies and partial recovery of mycorrhizal communities after invasive plant removal. *Biological Invasions* **16**:1979–1990.
- Lapointe L. 1998. Fruit development in *Trillium*: dependence on stem carbohydrate reserves. *Plant Physiology* **117**:183–188.
- Larsen LM, Nielsen JK, Sørensen H. 1992. Host plant recognition in monophagous weevils: specialization of *Ceutorhynchus inaffectatus* to glucosinolates from its host plant *Hesperis matronalis*. *Entomologia Experimentalis et Applicata* **64**:49–55.
- Leicht-Young SA, Pavlovic NB, Adams JV. 2012. Competitive interactions of garlic mustard (*Alliaria petiolata*) and damesrocket (*Hesperis matronalis*). *Invasive Plant Science and Management* **5**:27–36.
- Levine JM, Vilà M, Antonio CMD, Dukes JS, Grigulis K, Lavorel S. 2003. Mechanisms underlying the impacts of exotic plant invasions. *Proceedings of the Royal Society B: Biological Sciences* **270**: 775–781.



- Lu X, Koide RT. 1994. The effects of mycorrhizal infection on components of plant growth and reproduction. *New Phytologist* **128**:211–218.
- McCarthy BC. 1997. Response of a forest understory community to experimental removal of an invasive nonindigenous plant (*Alliaria petiolata*, Brassicaceae). In: Luken JO, Thieret JW, eds. *Assessment and management of plant invasions*. New York: Springer, 117–130.
- McDowell NG. 2011. Mechanisms linking drought, hydraulics, carbon metabolism, and vegetation mortality. *Plant Physiology* **155**:1051–1059.
- McDowell NG, Pockman WT, Allen CD, Breshears DD, Cobb N, Kolb T, Plaut J, Sperry J, West A, Williams DG, Yezpe EA. 2008. Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought? *New Phytologist* **178**:719–739.
- Meekins JF, McCarthy BC. 1999. Competitive ability of *Alliaria petiolata* (Garlic Mustard, Brassicaceae), an invasive, nonindigenous forest herb. *International Journal of Plant Sciences* **160**:743–752.
- Meinhardt KA, Gehring CA. 2012. Disrupting mycorrhizal mutualisms: a potential mechanism by which exotic tamarisk outcompetes native cottonwoods. *Ecological Applications* **22**:532–549.
- Morris WF, Pfister CA, Tuljapurkar S, Haridas CV, Boggs CL, Boyce MS, Bruna EM, Church DR, Coulson T, Doak DF, Forsyth S, Gaillard J-M, Horvitz CC, Kalisz S, Kendall BE, Knight TM, Lee CT, Menges ES. 2008. Longevity can buffer plant and animal populations against changing climatic variability. *Ecology* **89**:19–25.
- Nakagawa S, Freckleton RP. 2008. Missing inaction: the dangers of ignoring missing data. *Trends in Ecology and Evolution* **23**:592–596.
- Newsham KK. 2011. A meta-analysis of plant responses to dark septate root endophytes. *New Phytologist* **190**:783–793.
- Olano JM, Menges ES, Martínez E. 2006. Carbohydrate storage in five resprouting Florida scrub plants across a fire chronosequence. *New Phytologist* **170**:99–106.
- Pisula NL, Meiners SJ. 2010. Relative allelopathic potential of invasive plant species in a young disturbed woodland. *The Journal of the Torrey Botanical Society* **137**:81–87.
- Poulette MM, Arthur MA. 2012. The impact of the invasive shrub *Lonicera maackii* on the decomposition dynamics of a native plant community. *Ecological Applications* **22**:412–424.
- Powell LA. 2007. Approximating variance of demographic parameters using the delta method: a reference for avian biologists. *The Condor* **109**:949–954.
- Prati D, Bossdorf O. 2004. Allelopathic inhibition of germination by *Alliaria petiolata* (Brassicaceae). *American Journal of Botany* **91**:285–288.
- R Development Core Team. 2014. *R: a language and environment for statistical computing*. Vienna, Austria.
- Rita H, Komonen A. 2008. Odds ratio: an ecologically sound tool to compare proportions. *Annales Zoologici Fennici* **45**:66–72.
- Roberts KJ, Anderson RC. 2001. Effect of garlic mustard [*Alliaria petiolata* (Beib. Cavara & Grande)] extracts on plants and arbuscular mycorrhizal (AM) fungi. *The American Midland Naturalist* **146**:146–152.
- Rodgers VL, Stinson KA, Finzi AC. 2008a. Ready or not, garlic mustard is moving in: *Alliaria petiolata* as a member of Eastern North American forests. *BioScience* **58**:426–436.
- Rodgers VL, Wolfe BE, Werden LK, Finzi AC. 2008b. The invasive species *Alliaria petiolata* (garlic mustard) increases soil nutrient availability in northern hardwood-conifer forests. *Oecologia* **157**:459–471.
- Rosenthal R, Rosnow R. 1985. *Contrast analysis: focused comparisons in the analysis of variance*. Cambridge: Cambridge University Press.
- Rout ME, Callaway RM. 2012. Interactions between exotic invasive plants and soil microbes in the rhizosphere suggest that ‘everything is not everywhere’. *Annals of Botany* **110**:213–222.
- Ruckli R, Rusterholz HP, Baur B. 2014. Invasion of an annual exotic plant into deciduous forests suppresses arbuscular mycorrhiza symbiosis and reduces performance of sycamore maple saplings. *Forest Ecology and Management* **318**:285–293.
- Schuster MJ, Dukes JS. 2014. Non-additive effects of invasive tree litter shift seasonal N release: a potential invasion feedback. *Oikos* **123**:1101–1111.
- Schweiger R, Baier MC, Persicke M, Müller C. 2014. High specificity in plant leaf metabolic responses to arbuscular mycorrhiza. *Nature Communications* **5**:Article 3886.
- Shannon SM, Bauer JT, Anderson WE, Reynolds HL. 2014. Plant-soil feedbacks between invasive shrubs and native forest understory species lead to shifts in the abundance of mycorrhizal fungi. *Plant and Soil* **382**:317–328.
- Shefferson RP. 2009. The evolutionary ecology of vegetative dormancy in mature herbaceous perennial plants. *Journal of Ecology* **97**:1000–1009.
- Smith LM, Reynolds HL. 2012. Positive plant-soil feedback may drive dominance of a woodland invader, *Euonymus fortunei*. *Plant Ecology* **213**:853–860.
- Smith LM, Reynolds HL. 2014. Light, allelopathy, and post-mortem invasive impact on native forest understory species. *Biological Invasions* **16**:1131–1144.
- Smith SE, Read DJ. 2008. *Mycorrhizal symbiosis*, 3rd edn. London: Academic Press.
- Stinson KA, Campbell SA, Powell JR, Wolfe BE, Callaway RM, Thelen GC, Hallett SG, Prati D, Klironomos JN. 2006. Invasive plant suppresses the growth of native tree seedlings by disrupting belowground mutualisms. *PLoS Biology* **4**:e140.
- Tierens K, Thomma BPH, Brouwer M, Schmidt J, Kistner K, Porzel A, Mauch-Mani B, Cammue BPA, Broekaert WF. 2001. Study of the role of antimicrobial glucosinolate-derived isothiocyanates in resistance of *Arabidopsis* to microbial pathogens. *Plant Physiology* **125**:1688–1699.
- Tukey JW. 1991. The philosophy of multiple comparisons. *Statistical Science* **6**:100–116.
- Tylianakis JM, Didham RK, Bascompte J, Wardle DA. 2008. Global change and species interactions in terrestrial ecosystems. *Ecology Letters* **11**:1351–1363.
- van der Heijden MGA, Bardgett RD, van Straalen NM. 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology Letters* **11**:296–310.
- Van Diepen LTA, Lilleskov EA, Pregitzer KS. 2011. Simulated nitrogen deposition affects community structure of arbuscular mycorrhizal fungi in northern hardwood forests. *Molecular Ecology* **20**:799–811.
- Vaughn SF, Berhow MA. 1999. Allelochemicals isolated from tissues of the invasive weed garlic mustard (*Alliaria petiolata*). *Journal of Chemical Ecology* **25**:2495–2504.
- Vitousek PM, Walker LR. 1989. Biological invasion by *Myrica Faya* in Hawaii: plant demography, nitrogen fixation, ecosystem effects. *Ecological Monographs* **59**:247–265.

- Vogelsang KM, Bever JD. 2009. Mycorrhizal densities decline in association with nonnative plants and contribute to plant invasion. *Ecology* **90**:399–407.
- Weidenhamer JD. 2006. Distinguishing allelopathy from resource competition: the role of density. In: Reigosa MJ, Pedrol N, Gonzalez L, eds. *Allelopathy: a physiological process with ecological implications*. The Netherlands: Springer, 85–103.
- Weir TL, Park SW, Vivanco JM. 2004. Biochemical and physiological mechanisms mediated by allelochemicals. *Current Opinion in Plant Biology* **7**:472–479.
- Whigham DF. 2004. Ecology of woodland herbs in temperate deciduous forests. *Annual Review of Ecology, Evolution, and Systematics* **35**:583–621.
- Wixted KL, McGraw JB. 2010. Competitive and allelopathic effects of garlic mustard (*Alliaria petiolata*) on American ginseng (*Panax quinquefolius*). *Plant Ecology* **208**:347–357.
- Wolfe BE, Klironomos JN. 2005. Breaking new ground: soil communities and exotic plant invasion. *BioScience* **55**:477–487.
- Wolfe BE, Rodger VL, Stinson KA, Pringle A. 2008. The invasive plant *Alliaria petiolata* (garlic mustard) inhibits ectomycorrhizal fungi in its introduced range. *Journal of Ecology* **96**:777–783.
- Wright DP, Read DJ, Scholes JD. 1998. Mycorrhizal sink strength influences whole plant carbon balance of *Trifolium repens* L. *Plant, Cell and Environment* **21**:881–891.
- Wyka T. 1999. Carbohydrate storage and use in an alpine population of the perennial herb, *Oxytropis sericea*. *Oecologia* **120**:198–208.
- Zhu X-C, Song F-B, Liu S-Q, Liu T-D. 2011. Effects of arbuscular mycorrhizal fungus on photosynthesis and water status of maize under high temperature stress. *Plant and Soil* **346**:189–199.
- Zuleta A, Sambucetti ME. 2001. Inulin determination for food labeling. *Journal of Agricultural and Food Chemistry* **49**:4570–4572.
- Zuur A, Ieno EN, Walker N, Saveliev AA, Smith GM. 2009. *Mixed effects models and extensions in ecology with R*. New York: Springer Science & Business Media.