



Mycosphere Essay 8: A review of genus *Agaricus* in tropical and humid subtropical regions of Asia

Karunarathna SC^{1, 2, 3, 4}, Chen J^{3, 4}, Mortimer PE^{1, 2}, Xu JC^{1, 2}, Zhao RL⁵, Callac P^{6*} and Hyde KD^{1, 2, 3, 4*}

¹Key Laboratory of Economic Plants and Biotechnology, Kunming Institute of Botany, Chinese Academy of Sciences, 132 Lanhei Road, Kunming 650201, China.

²World Agroforestry Centre, China & East-Asia Office, 132 Lanhei Road, Kunming 650201, China.

³Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand.

⁴Mushroom Research Foundation, 128 M.3 Ban Pa Deng T. Pa Pae, A. Mae Taeng, Chiang Mai 50150, Thailand.

⁵State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China.

⁶INRA, MYCSA (Mycologie et sécurité des aliments), CS20032, 33882, Villenave d'Ornon, France.

Karunarathna SC, Chen J, Mortimer PE, Xu JC, Zhao RL, Callac P, Hyde KD 2016 – Mycosphe Essay 8: A review of genus *Agaricus* in tropical and humid subtropical regions of Asia. Mycosphe 7(4), 417–439, Doi 10.5943/mycosphe/7/4/3

Abstract

The genus *Agaricus* includes both edible and poisonous species, with more than 400 species worldwide. This genus includes many species, which are enormously important as sources of food and medicine, such as the button mushroom (*Agaricus bisporus*) and the almond mushroom (*Agaricus subrufescens*). This paper reviews the genus *Agaricus* in tropical and humid subtropical regions of Asia, including the history, characteristics, pertinent morphological and organoleptic taxonomic traits, molecular phylogeny and taxonomy advances, toxicity and edibility. This review includes *Agaricus* species that are known to be highly valued as edible and medicinal mushrooms, and provides a comprehensive checklist of species described from the tropical and humid subtropical regions of Asia until the end of 2015.

Keywords – check list – edible mushroom– medicinal mushroom – molecular phylogeny – poisonous mushroom

Introduction to the genus *Agaricus*

The history of the denomination of the genus *Agaricus* is relatively complex and is reported in detail in the monograph of Parra (2008), from which we briefly resume here the last major nomenclatural events. Donk (1962) proposed "*Agaricus* L. ex Fries" as "*nomen conservandum*" with "*Agaricus campestris* Linnaeus ex Fries" as the conserved type. In the 2006 ICBN edition (Vienna Code 2006), the starting point for the nomenclature of fungi is *Species plantarum* (Linnaeus, 1753) and, accordingly, the name of the genus is *Agaricus* L. (1753). This name has priority over two homotypic synonyms, which are *Psalliota* (Fr.) P. Kumm. (1871) and *Pratella* (Pers.) Gray (1821). There are about 6000 records with the name of "*Agaricus*" in Index Fungorum (Index Fungorum 2016). This is due to the fact that in the past, many species were designated as *Agaricus* added because the species had lamellae.

Submitted 24 June 2016, Accepted 19 July 2016, Published online 26 July 2016

Corresponding Author: Hyde KD – e-mail – kdhyde3@gmail.com

Callac P – e-mail – philippe.callac@bordeaux.inra.fr

Agaricus L. is a large and well-known edible mushroom genus that also includes a small number of species that are toxic if eaten (Kerrigan et al. 2006, Zhao et al. 2012a, Bau et al. 2014). *Agaricus* species are saprotrophic fungi and often gregarious in woods, forests, gardens, on roadside, fields, pasture-land, grass-land, rubbish dumps, manure heaps and alluvial soils. *Agaricus* can be found from sea level up to the vegetation limit in mountainous areas (Cappelli 1984), and also occurs in some arid areas (Xu et al. 1998; Lebel & Syme 2012; Lebel 2013).

According to Bas (1991), the number of *Agaricus* species worldwide is close to 400. Zhao et al. (2011) recognized 386 species in the genus, and from this publication to the end of 2015, 48 new species have been introduced. This includes 18 species new from Thailand (Chen et al. 2012, 2015, Zhao et al. 2012a, b, Karunaratna et al. 2014, Thongklang et al. 2014a, Ariyawansa et al. 2015, Liu et al. 2015), four from China (Li et al. 2014, Gui et al. 2015), one from Sri Lanka (Liu et al. 2015), 16 from Europe (Parra et al. 2011, 2014, Parra 2013), and nine new sequestrate species from Australasia (Lebel & Syme 2012, Lebel 2013). Accordingly, the total number of species is presently 434, and includes 200 tropical species. Even though the genus *Agaricus* is well-known as important edible mushrooms, the taxonomy and delimitation of the species within the genus is very complex (Cappelli 1984).

Taxonomic monographs and other taxonomic studies of *Agaricus* species are mostly from the Americas, Australia, Europe, India and New Zealand (Murrill 1912, 1918, 1941, Hotson 1938, Schäffer & Møller 1938, Smith 1944, Møller 1950, 1952, Pilát 1951, Huijsman 1960, Orton 1960, Bohus 1975, 1990, 1995, Heinemann 1978, 1986, Freeman 1979a, b, Pegler 1983, 1990, Cappelli 1984, Kerrigan 1985, 1989, Wasser 1989, Callac et al. 1993, Alberto & Wright 1994, Grgurinovic 1997, Saini et al. 1997, Valenzuela et al. 1997, Alberto 1998, Esteve-Raventós 1998, Flower et al. 1997, Mitchell & Walter 1999; Nauta 1999, 2000, Peterson et al. 2000, Lanconelli 2002, Parra 2003, 2008, 2013, Lacheva & Stoichev 2004, Natarajan et al. 2005, Geml et al. 2007, Ludwig 2007).

The most commonly referenced monographs on tropical *Agaricus* are those of Heinemann (Heinemann 1953–1996, especially those of 1978 and 1980). Heinemann's 1978 and 1980 treatments of *Agaricus* ranked species traditionally included in the genus at the subgeneric level. He added two predominantly subtropical and tropical subgenera, *Lanagaricus* Heinem. and *Conioagaricus* Heinem.

With DNA sequencing and rapidly expanding databases of genetic and taxonomic information, researchers are developing new, efficient tools for the identification and classification of fungi (Geml et al. 2004). Phylogenetic analyses based on ribosomal DNA sequences are now widely exploited to deduce evolutionary relationships among agaric taxa (Challen et al. 2003, Kerrigan et al. 1999, 2006, 2008, Geml et al. 2004, Zhao et al. 2011). The concept (or delimitation) of sections initially described from temperate species has further evolved in recent years (Parra 2008, Parra 2013), but these changes may not be sufficient to incorporate tropical diversity. Up until 2010, most reports concerning the systematics and phylogeny of *Agaricus* species were based on European and North American taxa, and failed to include those originating in Asia.

Pertinent morphological and organoleptic taxonomic traits in *Agaricus*

The sporocarp of *Agaricus* is generally described as having a white to brown more or less yellowish or rufescent pileus; free lamellae with a regular trama when young, later becoming irregular and producing a dark brown spore print; and partial veil which forms a ring on the stipe. Basidiospores are smooth with a compound wall and not visibly pseudoamyloid.

The taxonomic system of *Agaricus* is complex, and several treatments have been proposed by different mycologists based on their judgments of how significant each morphological characteristic is in the recognition of species and sections in the genus (Møller 1950, Pilát 1951, Konrad & Maublanc 1952, Kühner & Romagnesi 1953, Moser 1967–1983, Heinemann 1978, Wasser 1980, Cappelli 1984, Kerrigan 1986, Singer 1986, Parra 2008, 2013). All recent systematic treatments of the genus used four taxonomically relevant characteristics: 1) macrochemical reactions: Schaeffer's cross-reaction – a chemical test with aniline and concentrated Nitric acid; and

the alkali test (application of strong alkali, either NaOH or KOH); 2) basidiosporocarp colour change when bruised or cut; 3) odour, which requires that the researcher smelling the samples is trained beforehand (Heinemann 1987); 4) the structure of the annulus, which can be superous or inferous, simple or double, has more recently been recognized as a crucial characteristic.

Some characteristics that are generally not pertinent at the sectional level remain useful at the species level, although in certain cases traits such as the pileus colour or the microscopic features are sometimes more variable within species than between species. However, at the species level and within certain sections the following traits remain useful for species identification: pileus colour, aspect of on pileus (squamules), base of the stipe (marginate bulbous), spore size, cheilocystidia (catenulate), vacuolar pigment in the pileipellis hyphae and also habit, habitat or the geographic/climatic distribution.

Molecular phylogeny and taxonomy advances in *Agaricus*

Molecular phylogeny based on the internal transcribed spacers (ITS1 and ITS2) segments of the nuclear ribosomal DNA region of *Agaricus* was first carried out by Mitchell & Bresinsky (1999). This phylogeny gives a base to identify species and to develop their classification. It is also helpful to detect species that possess interesting traits such as edibility, ability to fruit on compost, odour and other biochemical properties, which are generally shared by members of the same section. Based on ITS data, *Agaricus* sect. *Bivelares* (previously named *A. sect. Duploannulati*) and *A. sect. Xanthodermatei*, have been phylogenetically reconstructed using temperate specimens from North America and Europe (Challen et al. 2003, Kerrigan et al. 2006, 2008). *Agaricus* sect. *Bivelares* includes edible species such as *A. bisporus* (J.E. Lange) Imbach and *A. bitorquis* (Qué.) Sacc., which are both cultivated, while species of *A. sect. Xanthodermatei* are confirmed or suspected to be toxic. Some species or varieties were reported as being new to science based on morphological, molecular and/or biological data (Callac et al. 1993, 2003, Callac & Guinberteau 2005, Geml et al. 2007, Kerrigan et al. 2008, Lebel 2012, Lebel & Syme 2013).

All the classical temperate sections are monophyletic except *A. sect. Spissicaules* and *A. sect. Sanguinolenti*. The latter appeared as a paraphyletic group comprising three clades in the ITS phylogenetic analyses of Zhao et al. (2011). Recently, Parra et al. (2014) has shown that one of these three clades that already contained *A. pattersoniae* Peck and *A. boisseletii* Heinem. corresponded to *A. sect. Nigrobrunnescens*, since the type species of the section, *A. nigrobrunnescens*, belongs to this clade. Many species belonging to *A. sect. Sanguinolenti* (e.g. *A. padanus* Lancon.) were transferred to section *Nigrobrunnescens*; moreover *A. caballeroi* L.A. Parra, G. Muñoz & Callac, a new species described from Spain (Parra et al. 2014) also belongs to this section. The two other clades representing subsections *Bohusia* and *Sylvatici* remain in *A. sect. Sanguinolenti*. With these recent research findings, *Agaricus* species of temperate areas are now distributed in nine sections that are all represented in Figure 1, except the section *Nigrobrunnescens*. Future investigation will be needed to characterize the tropical clades that have not yet been resolved, except *A. sect. Brunneopicti* also represented in Figure 1.

The first phylogenetic analysis including tropical *Agaricus* species was based on ITS data and published by Zhao et al. (2011). The authors aimed to examine the extent to which the current system of classification is appropriate for tropical species. The analysis included 38 temperate species representing the eight classical sections of the genus, and 86 putative species of *Agaricus* from tropical areas of Africa, Asia and the Americas, but mostly from northern Thailand. Samples of approximately 50 putatively novel species from a small area of northern Thailand were included. It was shown that (i) only about one-third of tropical species belong to the classical sections based on temperate species, and that therefore the systematics of the genus needed to be expanded; (ii) most of the remaining two-thirds of tropical species grouped in 11 clades generally exclusively neotropical (Americas) or palaeotropical (Africa + Asia), suggesting that secondary species diversification had occurred separately in these two areas; (iii) in contrast, several clades of classical sections contained both American and African or Asian species along with temperate




Fig. 1 ML phylogram based on ITS1+2 sequences showing 29 species of *Agaricus* described or reported from tropical and humid subtropical regions of Asia, which have been sequenced before 2016. Species originally described from this region are indicated in red while those only reported are in green. T = type specimen (a recent epitype in the case of *A. arvensis*). *A. brunneopunctatus* has been recently proposed to replace the illegitimate name *A. brunneopictus*. The two outgroup species are *Heinemannomyces* cf. *splendidissima* Watling and *Hymenagaricus ardosicolor* (Heinem.) Heinem. Eight classical sections and the palaeotropical section *Brunneopicti* are shown. For each sample the species name, the sample number, and the GenBank accession number are indicated.

species. Seven of the 11 tropical clades were statistically well supported and numbered from TR I to TR VII. It was hypothesized that clade TR I might represent *A. sect. Brunneopicti*.

This first investigation of tropical species has been followed by several studies using both morphological approaches and phylogenetic analyses based on ITS1+2 sequence data. These studies focused on tropical species in different groups such as the *A. sect. Xanthodermatei*, the *A. sect. Arvenses* and the clade TRI. Based on collections from Thailand, two new species *A. megacystidiatus* Karun., Guinb. & K.D. Hyde and *A. Chiangmaiensis* Karun., Guinb. & K.D. Hyde belonging to the clade TRI have been described (Karunaratna et al. 2014, Figure 1). More recently, Chen et al. (2015) confirmed that this clade corresponds to *A. sect. Brunneopicti*. The reconstructed section included 16 species grouped in four strongly supported subclades and two isolated branches. Only the six species of one of these clades exhibit punctiform squamules with remains of the veil on the pileus and stipe, which initially characterized this section. In addition, the new name *A. brunneopunctatus* L.J. Chen, Callac & L.A. Parra was introduced to replace the illegitimate name *A. brunneopictus* Berk. & Broome of the type species and four of the 16 species were described as new to science, *A. brunneosquamulosus* L.J. Chen, R.L. Zhao, K.D. Hyde & Callac, *A. niveogranulatus* L.J. Chen, R.L. Zhao, Callac & K.D. Hyde, *A. sordidocarpus* L.J. Chen, Callac & K.D. Hyde, and *A. toluenolens* Callac, L.J. Chen & K.D. Hyde (Figure 1). Studies on *A. sect. Xanthodermatei* including Thai and Pakistani tropical samples revealed five new species (*A. atrodiscus* L.J. Chen, Callac, R.L. Zhao & K.D. Hyde, *A. bisporiticus* Nawaz, Callac, Thongklang & Khalid, *A. exilissimus* L.J. Chen, Callac, R.L. Zhao & K.D. Hyde, *A. fuscopunctatus* Thongklang, L.J. Chen, Callac & K.D. Hyde, and *A. murinocephalus* R.L. Zhao, Desjardin & K.D. Hyde) and *A. microvolvatulus* Heinem., which was originally described from Africa, was reported from Thailand (Zhao et al. 2011; Thongklang et al. 2014a, Ariyawansa et al. 2015). Similarly, Gui et al. (2015) reported eight species of *Agaricus* section *Arvenses* from highland subtropical Southwest China and three of them (*A. guizhouensis* Y. Gui, Zuo Y. Liu & K.D. Hyde, *A. longistipes* Y. Gui, Zuo Y. Liu, Callac, L.A. Parra & K.D. Hyde, and *A. megalocarpus* Y. Gui, Zuo Y. Liu, Callac, L.A. Parra & K.D. Hyde) were described as new to science (Fig. 1). However, for certain species the correct classification remains uncertain (*A. hanthanaensis* Karun. & K.D. Hyde, Sri Lanka, Liu et al. 2015) or unresolved (*A. haematinus* K.D. Hyde & R.L. Zhao and *A. pseudolangei* K.D. Hyde & R.L. Zhao, Ariyawansa et al. 2015). We predict that the species richness of somewhat forgotten or new tropical sections will increase in coming years.

Toxicity and unknown edibility

Agaricus is a genus of saprobic mushrooms that includes economically important species such as *A. bisporus*, the button mushroom (Savoie et al. 2013, Thawthong et al. 2014). There has been significant ecological, nutritional and medicinal interest in the genus (Dai et al. 2009, 2011), yet the extent of its diversity remains poorly known, particularly in subtropical and tropical areas (Zhao et al. 2011). Excellent edible frequently consumed species belong to the sections *Agaricus*, *Arvenses*, *Bivelaes*, *Nigrobrunnescetes* and *Sanguinolenti*, nevertheless, heavy metal accumulation (cadmium, lead and mercury) has been reported in species of *A. sect. Arvenses* (Kalac & Svoboda 2000). In the five remaining sections most species are toxic or their edibility is generally unknown for different reasons; for example, they may be unattractive (unpleasant odour, small basidiosporocarps) or simply too rare, hard to identify, or recently recognized.

In *A. sect. Spissicaules*, *A. litoralis* (Wakef. & A. Pearson) Pilát is edible. However, *A. bresadolanus* Bohus is regarded as toxic causing gastro-intestinal syndrome. Other species are suspect and also difficult to identify.

Species of *A. sect. Minores* are not popular because they are hard to identify, even for experts, and they are generally small. However, it can be noted that intoxication by such species has not been reported and that they generally have a pleasant almond-like odour, as do species of the closely related *A. sect. Arvenses*. *Agaricus brunneolus* (I.E. Lange) Pilát (syn. *A. porphyizon* P.D. Orton) is one of the relatively large and sometimes abundant species of this section which is consumed in Europe (Cappelli 2011).

Species of *A. sect. Chitonioides* are sometimes eaten, although certain species can exhibit an unpleasant odour that disappears with cooking, such as *A. bernardii* Quél. which is widely consumed and highly prized by many families in certain places of Denmark. Bernard (1882) reported that *A. bernardii* is edible and sold at the market in La Rochelle (France). Møller (1950) pointed out that it is often eaten and is not toxic, even though many people do not like it because of its unpleasant smell, and because it is tough and somewhat indigestible.

The edibility of species of *A. sect. Brunneopicti* (Chen et al. 2015) is generally unknown except for the two species *A. subsaharianus* L.A. Parra, Hama & De Kesel and *A. bingensis* Heinem., which are consumed in Africa (Hama et al. 2010, Pegler 1977). This section contains species with odours ranging from pleasant bitter almond-like smells, to unpleasant odours similar to phenol or the solvents used in marker pens. Species exhibiting these kinds of unpleasant odours are potentially toxic, similar to the species of *A. sect. Xanthodermatei*, which give off similar odours.

Species of the *A. sect. Xanthodermatei* are recognized or suspected to be toxic. They are characterized by a negative Schäffer's reaction, a positive KOH reaction (yellow), an odour of phenol (sometimes faint or absent), and a yellow (sometimes absent or reddish) discoloration when cut or bruised (Kerrigan et al. 2006, Parra 2013, Thongklang et al. 2014a). Despite their potential or recognized toxicity, some species of this section are regularly eaten and tolerated by some people from Eastern Europe and Russia (Pilát 1951, Bohus 1974, Kerrigan et al. 2006, Parra 2008) or Spain (Parra 2008) without gastro-intestinal symptoms. Genevier (1876) reported that *A. xanthodermus* Genev. was sold in the markets of Nantes and noted three *A. xanthodermus* poisoning cases, of which one was serious. Menier (1893) noted that *A. menieri* Bon was not tasty, and though not much prized, it was eaten in Siant-Brevin. Spegazzini (1919) pointed out *A. iodoformicus* Speg. (a synonym of *A. xanthodermus*) is edible, despite its strong odour and taste of iodoform perceivable even in dried specimens. Regarding the edibility of *A. phaeolepidotus* F.H. Møller, Pilát & Ušák (1961) mentioned it is a good edible fungus. Genevier (1876) and Jordan (2000) indicated that some people can eat *A. xanthodermus* and *A. placomyces* Peck without any ill-effects at all, while Passecker (1930) reported that inhabitants of Rosemberg, Austria, cultivate *A. xanthodermus* by cutting the stipes and planting the caps of *A. xanthodermus* in gardens, after which basidiomes appear in the following year which are collected and eaten without any problem; moreover Passecker (1930) cultivated *A. xanthodermus* mycelia artificially and noted that the mycelia smelled of phenol. Bon (1987) and Jordan (2000) reported that eating *A. xanthodermus* can cause alarming symptoms including coma, vomiting or diarrhea, but recovery is complete within a few days. According to Walton (1986) three families suffered by consuming *A. xanthodermus*. First detected in *A. xanthodermus* (Gill & Strauch 1984), the phenol is also the major volatile component in *A. moelleri* Wasser (syn. *A. praeclaresquamosus* A.E. Freeman) (Wood et al. 1998) and likely responsible for the poisonous symptoms. In the medical poison center in Marseille, France, 20% of poisonings by identified mushrooms are due to ingestion of species of *A. sect. Xanthodermatei* (De Haro et al. 1999).

Agaricus species having high value as edible and medicinal mushrooms

The main cultivated species are in section *Bivelares* and *Arvenses*, while it remains difficult to cultivate species of sections *Sanguinolenti*, *Nigrobrunnescentes* and *Agaricus* because mycelium growth rate is generally too slow (Callac 2007). The consumed species in section *Sanguinolenti* (e.g. *A. sylvaticus* Schaeff.) and in section *Nigrobrunnescentes* (e.g. *A. liliceps* Zeller) are considered very tasty (Wang et al. 2015). Unfortunately, cultivation of *A. liliceps* on standard substrates has been limited or has failed due a slow mycelium growth rate (Hildén et al. 2013).

In *Agaricus* sect. *Agaricus*, *A. campestris* (the field or meadow mushroom) is widely collected and eaten, but this mushroom is not commercially cultivated on account of its fast maturing and short shelf-life (Grigson 1975). In fact, the cultivation of this species has never been formally reported. Studies reported on *A. campestris* must be carefully considered knowing that this species was in the past frequently confused with other species of *A. sect. Agaricus* or even with

species of other sections such as *A. bisporus*. This possibly results from the perpetuation of an outdated and incorrect concept confusing samples of different species exhibiting a white pileus. In fact, the colour of the pileus can vary from white to dark brown within the populations of *A. bisporus* (Callac et al. 2002). For example, it is unfortunate that Gray and Flatt (1998) report in their “materials and methods” section of their study: “Plant material. Dried fruiting bodies of mushroom (*A. campestris*) obtained from a commercial source”. The antihyperglycemic effects reported by these authors and subsequently by Gallagher et al. (2003), Patrick (2008), and De Silva et al. (2012b) do not concern *A. campestris*, but likely *A. bisporus*.

In *A. sect. Bivelares* most species seem edible and cultivatable, but only *A. bisporus* and *A. bitorquis* are commercially cultivated. Species closely related to *A. bisporus* (*A. subfloccosus* (J.E. Lange) Hlaváček and *A. agrinferus* Kerrigan & Callac) have been cultivated on an experimental basis in several laboratories and are regarded as high-quality mushrooms (Noble et al. 1995, Kerrigan et al. 1999, 2008, Callac et al. 2005). Interestingly, *A. cupressicola* Bon & Grilli and many strains of *A. devoniensis* P.D. Orton, the sand dune mushroom, have been successfully cultivated at INRA, France (Callac et al. 2005, Callac 2007).

In *A. sect. Arvenses* most species seem to be edible, but not all of them are cultivatable, and only *A. subrufescens* Peck. and *A. arvensis* Schaeff. are commercially cultivated. There are reports of cultivation experiments of *A. macrosporoides* Bohus (Bohus 1978, 1987), *A. crocodilinus* Murrill (syn. *A. macrosporus* (F.H. Møller & Jul. Schäff.) Pilát and *A. urinascens* (Jul. Schäff. & F.H. Møller) Singer) (Fermor 1982) and *A. sylvicola* (Vittad.) Peck (Heinemann 1977). *Agaricus flocculosipes* R.L. Zhao, Desjardin, Guinb. & K.D. Hyde was described from northern Thailand, and is expected to be a good edible species that may have potential commercial value for Thailand and other tropical countries. It also has an almond flavor (Zhao et al. 2012b) and some fruit bodies have been obtained through cultivation (Thongklang et al. 2014b). *Agaricus guizhouensis*, which is consumed in China, and *A. megalocarpus* are potentially valuable for cultivation because of their relatively large sized sporocarps. Although *A. longistipes* has smaller sporocarps, it may be of medicinal interest since it is a relative of the robust *A. crocodilinus*, in which analgesic triglycerides have been discovered (Stadler et al. 2005, Gui et al. 2015). *Agaricus augustus* Fr. is mainly collected in Europe and North America, and is highly sought after for its flavor (Boa 2006); however, despite its popularity it has never been successfully cultivated. Wild edible mushrooms in the section *Arvenses* include *A. arvensis*, *A. augustus* and *A. sylvicola*, all of which are collected from the wild for human consumption in Europe and the Americas (Kalač & Svoboda 2000, Wisitrassameewong et al. 2012a, Thongklang et al. 2014b). Nutritional studies have been carried out on *A. augustus*, *A. crocodilinus* and *A. essettei* Bon (Elliott 1978, Fermor 1982, Geml & Rimoczi 1999, Dabbour & Takruri 2002, Geml & Royse 2002, Barros et al. 2007, Ozturk et al. 2011, Marekov et al. 2012); volatile components responsible for almond or aniseed odours have been identified in several species such as *Agaricus essettei* (Rapior et al. 2002). *Agaricus crocodilinus* has been reported as a potential bioremediation agent (Garcia et al. 2005).

Commercially cultivated species of *Agaricus* and their medicinal potential

Commercially cultivated *Agaricus* species belong to the sections *Bivelares* or *Arvenses*. The species of *A. sect. Bivelares* are known to be highly nutritious, an example being *A. bisporus*. This mushroom was first cultivated three and half centuries ago in France, and remains the most widely cultivated species of edible mushrooms, representing about 32% of world production in 1997 and 21% in 2013 (Chang 1999; Challen et al. 2003; Anonymous 2004, Zhang et al. 2015). *Agaricus bisporus* has a broad geographic distribution all along the temperate areas of the north hemisphere. A highly diversified germplasm is available (Callac et al. 2002) and the biology and genetics of *A. bisporus* has been well documented (Savoie et al. 2013). Great attention has been focused on the immune modulating and antitumor properties of *A. bisporus* (Yu et al. 1993, Chen et al. 2006, Ren et al. 2008, Adams et al. 2008, Jeong et al. 2010). This mushroom has potential anti-inflammatory, hypoglycemic and hypocholesterolemic effects due to the presence of high amounts of acidic

polysaccharides, dietary fibre, antioxidants (folate, ergothioneine, polyphenol) and vitamins (C, B12, D) (Fukushima et al. 2000, Mattila et al. 2001, Koyyalamudi et al. 2009, Geösel 2011). It also suggests that a high intake of button mushrooms may promote innate immunity against tumors and viruses (Wu et al. 2007, Adotey et al. 2011, De Silva et al. 2012a, b). Lectins are abundant in *A. bisporus* and dehydrated fruiting body extracts of *A. bisporus* have been shown to lower blood glucose and cholesterol levels (Jeong et al. 2010, Yamac et al. 2010, Smiderle et al. 2011, De Silva et al. 2012a). Rats fed with *A. bisporus* fruiting bodies exhibited significant anti-glycemic and anti-hypercholesterolemic effects (Jeong et al. 2010, Volman et al. 2010, De Silva et al. 2013), and significant decrease in plasma total cholesterol (TC) and low-density lipoprotein (LDL) (22.8 % and 33.1 %) accompanied by a significant increase in plasma high-density lipoprotein concentrations. This was observed after oral administration of *A. bisporus* for four weeks in an animal model (Jeong et al. 2010, De Silva et al. 2013). Moreover, this mushroom shows a positive influence on lipid metabolism and liver function.

Another commercially cultivated species of *A. sect. Bivelares* is *A. bitorquis* (the pavement mushroom), which is frequently found on hard-packed soils. It was first introduced for cultivation by Poppe (1972) in the 1970s and its cultivation has persisted in regions with hotter climates, including India (Cappelli 1984, Heinemann 1978, Kerrigan 1986). In Spain *A. bitorquis* is grown instead of *A. bisporus* in hot summers as it has a slightly higher growth temperature. However, recent studies have shown that certain strains of *A. bisporus* can also fructify at the relatively high temperature of 25 °C or more (Largeteau et al. 2011).

Agaricus subrufescens, the almond mushroom, is a species of *A. sect. Arvenses* which was first discovered in eastern North America and commercially cultivated in USA from 1890 to 1910 (Peck 1893, Kerrigan 2005, Firenzuoli et al. 2008). This species was ‘rediscovered’ and cultivated in Brazil but misidentified as *Agaricus blazei* Murrill (or ABM) or *Agaricus sylvaticus* (misapplied names, Dias et al. 2004). This species is also used as a health food for alternative medicines and cultivated in Asia, mainly in Indonesia, China, Japan, and Taiwan. For taxonomy and synonymy of this taxon we follow Kerrigan (2005), Arrillaga and Parra (2006), Ludwig (2007), Cappelli (2011) and Thongklang et al. (2016). Main synonyms are *A. blazei sensu* Heinemann (1993), *A. bambusae* Beeli var. *bambusae* (1928), *A. rufotegulis* Nauta (1999), *A. brasiliensis* Wasser, M. Didukh, Amazonas & Stamets (2002) nom. illegit., and *A. albopersistens* Zuccher (2006). *Agaricus subrufescens* is a cosmopolitan species reported from the Americas, Oceania (Hawaii) and Europe (Kerrigan 2005), and more recently from Thailand (Wisitrassameewong et al. 2012a), China (Gui et al. 2015) and Africa (Thongklang et al. 2016). This sort of large geographical distribution range is infrequent in the genus *Agaricus* and thus the question of potential interfertility between specimens originating from different continents arose. Kerrigan (2005) reported hybridization between Brazilian and Californian strains, and recently, Thongklang et al. (2014c) showed that samples from Brazil, France and Thailand are amphithallic (i.e. both heterothallic and pseudohomothallic) and interfertile in all pairwise combinations. In addition, Brito de Rocha et al. (2016) found that the sporocarps of certain isolates produce a proportion of heterokaryotic spores giving rise to heterokaryons, which are infertile (i.e. unable to fruit), but can easily cross with homokaryons. Recent genetic analyses revealed a duplication of the ITS in a French isolate suggesting a possible introgression from Asian or African populations (Chen et al. 2016a). A genetic linkage map of *A. subrufescens*, which exhibits a highly conserved macrosyteny with *A. bisporus* was published recently (Foulongne-Oriol et al. 2016). Such a rich available germplasm and so much available information relating to the genetics and biology of *A. bisporus* should facilitate the genetic improvement of *A. subrufescens* in the future.

However, currently, with the exception of a Brazilian-Californian hybrid, strains of *A. subrufescens* that are commercially cultivated in Asia have likely been introduced from Brazil. Thongklang et al. (2014b) showed the successful results of preliminary fruiting test of a Thai wild strain, while Jatuwong et al. (2014) described the development of a Thai-French hybrid strain of *A. subrufescens* recently developed through a joint program between INRA, France and Mae Fah

Luang University. This strain has the potential to grow in northern Thailand, as its optimal growth conditions are 25–30°C with 91–95 % relative humidity. *Agaricus subrufescens* has many health promoting benefits or medicinal properties such as anti-cancer and tumor suppressive activity, anti-genotoxicity activities, cytokine induction activity, lymphocyte activation activity, antimicrobial activity, anti-allergy effect, and biological responses on the immune system (Wisitrassameewong et al. 2012b). There has also been research on the nutraceutical properties of *A. subrufescens* and the protective effects of β -glucans (Angeli et al. 2006; Mantovani et al. 2007). The same fungus has also been reported to have anti-mutagenic activity (Menoli et al. 2001, Machado et al. 2005), anti-clastogenic properties (Luiz et al. 2003), anti-genotoxic activity (Martins de Oliveira et al. 2002) and anti-tumor effects (Kawagishi et al. 1989, Mizuno et al. 1990, 1999, Lee et al. 2003). *Agaricus subrufescens* has previously been reported to have pharmaceutical potential (Firenzuoli 2008) and is used in cosmetics (Hyde et al. 2010). Under the misapplied name *A. sylvaticus* some research has investigated its effects on cancer and AIDS, and other potential medicinal properties (Gennari 2000a, 2000b, Gennari et al. 2001, Taveira et al. 2008; Fortes et al. 2008, 2009, Fortes and Carvalho 2011, De Silva et al. 2012b, 2013). In spite of these useful properties, the reason *A. subrufescens* has not found wide commercial application is possibly its difficulty of digestion and strong almond flavour (Wisitrassameewong et al. 2012b). Although this is a well-known edible species, we queried northern Thai locals in local markets and concluded that this species is not popularly consumed.

Agaricus arvensis, the horse mushroom, is also a species of *A. sect. Arvenses* cultivated in Europe (Galli 2004), where it is produced by mushroom cultivators in France and Holland (Couvy 1974). A recent study, however, indicated that a strain commercialized under this name belongs to the closely related species *A. fissuratus* F.H. Møller (Parra 2013).

A checklist of species reported from tropical and humid subtropical regions of Asia

In Table 1, we listed 84 species of *Agaricus* described before 2016 from tropical and humid subtropical regions of Asia. Historically, 31 species were described in the nineteenth century, mostly (27) from Sri Lanka by the British mycologists M. J. Berkeley and C. E. Broome. These species have relatively short descriptions and only a few of them have been further studied, again by P. Heinemann or D. Pegler. A single species is well known today, *A. endoxanthus* Berk. & Broome, which is distributed in most tropical areas and has even been introduced to tropical niches of temperate regions such as greenhouses (Chen et al. 2016b). In the twentieth century, 31 new species were described including eight, mainly from Singapore, by the Belgium mycologist P. Heinemann. Only one, *A. duplocingulatus*, has been recently reported again, in Thailand (Chen et al. 2015). From 2000 to 2015, 22 new species have been described, mostly (18 species) from Thailand.

Sequence data (ITS) are currently available in GenBank for 24 of the 84 (29 %) species originally described from tropical and subtropical humid areas of Asia. This includes *A. endoxanthus*, *A. duplocingulatus* Heinem. and the 22 new species described from 2000 to 2015. In addition, sequences of specimens collected in this Asian region, but belonging to five species initially described from other continents are also available. Two of them, *abruptibulbus* Peck (1905) and *A. subrufescens* Peck (1894), are cosmopolitan species which have been originally described from North America, then reported from Europe and more recently from subtropical China (Gui et al. 2015). Moreover, *A. subrufescens*, which is one of the rare species of the genus to be distributed in both temperate and tropical areas, has also been reported from Thailand (Wisitrassameewong et al. 2012a). The three remaining species, *A. heterocystis* Heinem. & Gooss. Font., *A. microvolvatulus* and *A. xanthosarcus* Heinem. & Gooss.-Font., are exclusively tropical and were originally described from tropical Africa by Heinemann (1956, 1971). Zhao et al. (2011) reported Thai samples having ITS sequences identical to their respective African type specimens. Because morphological comparative studies have not yet been done except for *A. microvolvatulus* (Thongklang et al. 2014c), these three species are represented by their African type specimen in the

phylogenetic tree of Figure 1. In sum, 29 species from tropical or humid subtropical regions of Asia are represented in the phylogenetic tree of Figure 1: 24 species originally described from these regions, two cosmopolitan species (*A. subrufescens* and *A. abruptibulbus*), and three tropical species distributed in both Africa and Asia (*A. heterocystis*, *A. microvolvatulus* and *A. xanthodermatei*). It must be noted that many unnamed species or species newly described since 2016 are not included in this review.

Climate was noted as a major factor in the evolution of *Agaricus* by Zhao et al. (2011). Among the 29 sequenced species, 25 have been described or reported from tropical areas, mainly in Thailand. Among these 25 species, only the cosmopolitan species *A. subrufescens* and the “great traveler” species *A. endoxanthus* are also reported in temperate areas but the latter only because it was introduced to tropical greenhouses (Chen et al. 2016b). Another species, *A. flocculosipes*, is reported from the subtropical region of China. It can be noted that the two non-strictly tropical species *A. subrufescens* and *A. flocculosipes*, both belonging to *A. sect. Arvenses*, are also reported from tropical Africa by Zhao et al. (2012b) and Thongklang et al. (2016). Among the remaining 22 species having a known strictly tropical distribution range, 17 belong to the three sections *A. sect. Xanthodermatei*, *Minores*, and *Brunneopicti*, one to *A. sect. Agaricus*, one to *A. sect. Spissicaules*, and the three remaining species are unclassified. This count roughly reflects the phylogenetic diversity and species richness of *Agaricus* in tropical Asia where two traditional sections (*A. sect. Xanthodermatei* and *A. sect. Minores*) and tropical sections (*A. sect. Brunneopicti* and unnamed groups) dominate. Few members of *A. sect. Arvenses* are also present but they are not exclusively tropical, except for *A. subtilipes* Thongklang, L.J. Chen, Callac & K.D. Hyde, a new species recently described in 2016 (Thongklang et al. 2016) and thus not included in our checklist.

We, members of Center of Excellence in Fungal Research collected 400 *Agaricus* specimens in a small area of Northern Thailand between May 2005 and October 2015. These samples potentially include more than 70 novel species, indicating that *Agaricus* is a species-rich genus in the tropics as well as in temperate regions (Zhao 2008, Zhao et al. 2011, 2012a, b, Chen et al. 2012, 2015, Karunaratna et al. 2014, Thongklang et al. 2014a, b, 2016, Thongklang 2016). In agreement with the checklist and the tree presented above, no species of *A. sect. Bivelares* or *A. sect. Chitonioides* and only a few species of *A. sect. Agaricus*, *A. sect. Sanguinolenti* and *A. sect. Spissicaules*, were found. We did not list all the species originally described from other continents and reported from tropical or humid subtropical Asia, except those that were molecularly confirmed. The reason is that most identification of such species are dubious. For example, some species originally described in Europe such as *A. campestris*, *A. bisporus* and *A. bitorquis* have been inventoried from Thailand in local reports (Zhao 2008, Chandrasrikul et al. 2011). In absence of published detailed descriptions or ITS sequences, these identifications will remain doubtful, given that during ten years of field collecting in Thailand we never found any species which was originally described in Europe. However, the reports of some species originally described from Africa, such as *A. rufolanosus*, are more plausible.

Numerous new species, mostly of *A. sect. Xanthodermatei*, *A. sect. Minores* and *A. sect. Brunneopicti* remain to be named. However, before describing a new species, we generally expect to have several samples and to know its phylogenetic position. This allows us to take into consideration intraspecific variability and to identify characteristics that are distinct from neighboring species. Certain species of *A. sect. Minores* and *A. sect. Brunneopicti* would merit investigation of their edibility and cultivability. Some species remain unclassified and certain putative tropical new sections remain to be named. However, new species and sections have been recently proposed in 2016 (Zhao et al. 2016), including, for example, the new *A. sect. Amoeni* Callac & Zhao corresponding to the tropical clade TRIII in Zhao et al. (2011). An inventory of *Agaricus* species in tropical and humid subtropical regions of Asia will be a first step towards describing numerous species new to science. Further research could build on the present study by providing DNA sequences and developing the classification of species in tropical regions of Africa and the Americas.

Table 1 *Agaricus* species originally described until end 2015 from tropical and humid subtropical regions of Asia

Country	Species	Reference
China		
Yunnan prov.	<i>A. dimorphosquamatus</i> Yu Li	Li 1990
Yunnan prov.	<i>A. lepiotiformis</i> Yu Li	Li 1990
Yunnan prov.	<i>A. compressipes</i> W.F. Chiu	Wei-Fan 1973
Yunnan prov.	<i>A. yunnanensis</i> W.F. Chiu	Wei-Fan 1973
Guizhou prov.	<i>A. guizhouensis</i> Y. Gui et al.	Gui et al. 2015
Guizhou prov.	<i>A. longistipes</i> Y. Gui et al.	Gui et al. 2015
Guizhou prov.	<i>A. megalocarpus</i> Y. Gui et al.	Gui et al. 2015
India		
Bengal	<i>A. burkillii</i> (Masse) Sacc. & Trotter Basionym <i>Psallota burkillii</i>	Saccardo & Trotter 1913 Masse 1907
Calcutta	<i>A. squalidus</i> Masse Nom. illegit.	Masse 1912
Bombay	<i>A. woodrowii</i> Masse <i>A. fulviceps</i> Berk.	Masse 1901 Berkeley 1854
Karnataka	<i>A. heinemanniensis</i> K. Natarajan & Purush.	Natarajan & Purushothama 1996
Kerala	<i>A. basianulosus</i> Paracer & Chahal	Paracer & Chahal 1963 [1962]
Japan		
	<i>A. semotellus</i> S. Imai. <i>A. comtulellus</i> S. Imai. <i>A. jezoensis</i> S. Imai.	Imai 1938 Imai 1938 Imai 1938
Bonin Islands	<i>A. primipilus</i> Berk. & M. A. Curtis	Berkeley & Curtis 1860 [1858]
Java		
	<i>A. rhinocerotis</i> Jungh. <i>A. inoxydabilis</i> ^a Heinem.	Junghuhn 1840 Heinemann 1980
Malaysia		
Penang	<i>A. inoxydabilis</i> ^a Heinem.	Heinemann 1980
Sabah	<i>A. albidoperonatus</i> Heinem.	Heinemann 1980
Pakistan		
Punjab	<i>A. bisporiticus</i> ^b Nawaz et al.	Thongklang et al. 2014a
Philippines		
	<i>A. argyrotectus</i> Copel. <i>A. boltoni</i> Copel. <i>A. philippinensis</i> Berk. <i>A. merrillii</i> Copel. <i>A. manilensis</i> Copel. <i>A. perfuscus</i> Copel.	Copeland 1905 Copeland 1905 Berkeley 1842 Copeland 1905 Copeland 1905 Copeland 1905
Singapore		
	<i>A. inedulius</i> Heinem. <i>A. pleurocystidiatus</i> Heinem. <i>A. inoxydabilis</i> ^a Heinem. ^a <i>A. hypophaeus</i> Heinem. <i>A. duplocingulatus</i> Heinem. (in Thailand) <i>A. oenotrichus</i> Heinem. <i>A. singaporensis</i> Heinem. <i>A. tenuiceps</i> Masse	Heinemann 1980 Heinemann 1980 Heinemann 1980 Heinemann 1980 Heinemann 1980 Chen et al. 2015 Heinemann 1980 Heinemann 1990 Masse 1914
Sri Lanka		
	<i>A. simulans</i> Berk. <i>A. rufo-albus</i> Berk. <i>A. crocopeplus</i> Berk. & Broome (in Singapore) (in Sri Lanka; subg. <i>Lanagaricus</i>) <i>A. didactylus</i> Berk. & Broome <i>A. spilocephalus</i> Berk. & Broome <i>A. lasiophrys</i> Berk. & Broome <i>A. hemilasius</i> Berk. & Broome <i>A. endoxanthus</i> Berk. & Broome (in Thailand)	Berkeley 1847 Berkeley 1847 Berkeley & Broome 1871 Heinemann 1980 Peggler 1986 Berkeley & Broome 1871 Berkeley & Broome 1870 [1871] Berkeley & Broome 1871 Berkeley & Broome 1871 Berkeley & Broome 1871 Zhao et al. 2012a

Country	Species	Reference
	(in Malaysia)	Chen et al. 2016b
	<i>A. actinorachis</i> Berk. & Broome	Berkeley & Broome 1871
	(in Sri Lanka; subg. <i>Contoagaricus</i>)	Pegler 1986
	<i>A. tornocephalus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. nymphidius</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. bolorhizus</i> Berk. & Broome	Berkeley & Broome 1871
	(in Sri Lanka)	Pegler 1986
	<i>A. dyspines</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. lituratus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. celidotus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. chrysocyclus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. lepiotoides</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. rhodochrous</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. argineus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. microcosmus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. plumarius</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. callipeplus</i> Berk. & Broome	Berkeley & Broome 1871
	(in Sri Lanka)	Pegler 1986
	<i>A. chloroconius</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. illotus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. illotus</i> var. <i>thyranophurus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. subcitrinus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. erythrospilus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. subaeruginosus</i> Berk. & Broome	Berkeley & Broome 1871
	<i>A. stadii</i> (Petch) Cash	Cash, in Trotter 1972
	(basonym: <i>Psalliota stadii</i>)	Petch 1925
	<i>A. petchii</i> Pegler nom nov.	Pegler 1986
	(syn. <i>Psalliota zeylanica</i> Petch)	Petch 1917
	<i>A. hanthanaensis</i> Karun. & K.D. Hyde	Liu et al. 2015
Thailand		
Chiang Rai	<i>A. flocculosipes</i> ^c R.L. Zhao et al.	Zhao et al. 2012b
	(in Guizhou, China)	Gui et al. 2015
Chiang Rai	<i>A. megalosporus</i> J. Chen et al.	Chen et al. 2012
Chiang Rai	<i>A. bisporiticus</i> ^b Nawaz et al.	Thongklang et al. 2014a
Chiang Rai	<i>A. fuscopunctatus</i> Thongklang et al.	Thongklang et al. 2014a
Chiang Mai	<i>A. murinocephalus</i> R.L. Zhao et al.	Zhao et al. 2012a
Chiang Mai	<i>A. megacystidiatus</i> Karunarathna et al.	Karunarathna et al. 2014
Chiang Mai	<i>A. chiangmaiensis</i> Karunarathna et al.	Karunarathna et al. 2014
Chiang Mai	<i>A. niveogranulatus</i> L.J. Chen et al.	Chen et al. 2015
Chiang Mai	<i>A. brunneosquamulosus</i> L.J. Chen et al.	Chen et al. 2015
Chiang Mai	<i>A. sordidocarpus</i> L.J. Chen et al.	Chen et al. 2015
Chiang Mai	<i>A. toluenolens</i> Callac et al.	Chen et al. 2015
Chiang Rai	<i>A. flavicentrus</i> Karunarathna & K.D. Hyde	Liu et al. 2015
Chiang Mai	<i>A. parvibicolor</i> L.J. Chen et al.	Liu et al. 2015
Chiang Mai	<i>A. sodalis</i> L.J. Chen et al.	Liu et al. 2015
Chiang Mai	<i>A. atrodiscus</i> L.J. Chen et al.	Ariyawansa et al. 2015
Chiang Mai	<i>A. exilissimus</i> L.J. Chen et al.	Ariyawansa et al. 2015
Chiang Mai	<i>A. haematinus</i> K.D.Hyde & R.L.Zhao	Ariyawansa et al. 2015
Chiang Mai	<i>A. pseudolangei</i> K.D.Hyde & R.L.Zhao	Ariyawansa et al. 2015
Vietnam		
	<i>A. iocephalus</i> Pat., (syn. <i>A. iocephalusiopsis</i>)	Patouillard 1913
	<i>A. phaeocyclus</i> Pat.	Patouillard 1913
	<i>A. rhopalopodius</i> Pat.	Patouillard 1913

^a Described from Java, Malaysia and Singapore

^b Described from both Thailand and Pakistan

^c Described from both Thailand and Mayotte Island (Africa)

Available ITS GenBank accession number of species in bold type are indicated in Fig. 1

Acknowledgements

Andrew Stevenson in the World Agroforestry Centre (ICRAF), East and Central Asia Office, Kunming Institute of Botany, Kunming, Yunnan, China is thanked for English editing. Saranyaphat Boonmee and Thatsanee Luangharn (MFLU) are thanked for their assistance in the herbarium. Kevin D. Hyde thanks the Chinese Academy of Sciences, project number 2013T2S0030, for the award of Visiting Professorship for Senior International Scientists at Kunming Institute of Botany. This study was financially supported by Thailand Research Fund grant - Taxonomy, Phylogeny and biochemistry of Thai Basidiomycetes (BRG 5580009), the National Research Council of Thailand (NRCT), projects - Taxonomy, Phylogeny and cultivation of *Lentinus* species in northern Thailand (NRCT/55201020007), Mae Fah Luang University, the project—Taxonomy, Phylogeny and cultivation of *Lentinus* species in northern Thailand (MFU/54 1 01 02 00 48), National Natural Science Funds of China (Project No: 31000013, 31360014 and 31470152), and the Chinese Ministry of Science and Technology, under the 12th 5-year National Key Technology Support Program (NKTSP)2013BAB07B06 integration and comprehensive demonstration of key technologies on Green Phosphate-mountaion Construction. We would also like to thank the CGIAR Research Program 6: Forest, Trees and Agroforestry, for partially funding this work.

References

- Adams LS, Phung S, Wu X, Ki L, Chen S. 2008 – White button mushroom (*Agaricus bisporus*) exhibits antiproliferative and proapoptotic properties and inhibits prostate tumor growth in athymic mice. *Nutrition and Cancer* 60(6), 744–56.
- Adotey G, Quarcoo A, Holliday JC, Fofie S, Saaka B. 2011 – Effect of immunomodulation and antiviral agent of medicinal mushroom (immune assist 24/7) on CD4 + T-lymphocyte counts of HIV-infected patients. *International Journal of Medicinal Mushrooms* 13, 109–113.
- Albertó E. 1998 – *Agaricus santacatalinensis* a new species from Argentina. *Mycotaxon* 66, 205–214.
- Albertó E, Wright JE. 1994 – *Agaricus pseudoargentinus* n. sp. from Argentina. *Mycotaxon* 50, 271–278.
- Angeli, JP, Ribeiro LR, Gonzaga ML, Soares Sde A, Ricardo MP, Tsuboy MS, Stidl R, Knasmueller S, Linhares RE, Mantovani MS. 2006 – Protective effects of beta-glucan extracted from *Agaricus brasiliensis* against chemically induced DNA damage in human lymphocytes. *Cell Biology and Toxicology* 22, 285–291.
- Anonymous 2004 – Mushroom production falling in European Union. *Mushroom Business (The Netherlands)* 4, 1.
- Ariyawansa HA, Hyde KD, Jayasiri SC, Buyck B, Chethana KWT, Dai DQ, Dai YC, Daranagama DA, Jayawardena RS, Lücking R, Ghobad-Nejhad M, Niskanen T, Thambugala KM, Voigt K, Zhao RL, Li GJ, Doilom M, Boonmee S, Yang ZL, Cai Q, Cui Y-Y, Bahkali AH, Chen J, Cui BK, Chen JJ, Dayarathne MC, Dissanayake AJ, Ekanayaka AH, Hashimoto A, Hongsanan S, Jones EBG, Larsson E, Li WJ, Li Q-R, Liu JK, Luo ZL, Maharachchikumbura SSN, Mapook A, McKenzie EHC, Norphanphoun C, Konta S, Pang KL, Perera RH, Phookamsak R, Phukhamsakda C, Pinruan U, Randrianjohany E, Singtripop C, Tanaka K, Tian CM, Tibpromma S, Abdel-Wahab MA, Wanasinghe DN, Wijayawardene NN, Zhang J-F, Zhang H, Abdel-Aziz FA, Wedin M, Westberg M, Ammirati JF, Bulgakov TS, Lima DX, Callaghan TM, Callac P, Chang C-H, Coca LF, Dal-Forno M, Dollhofer V, Fliegerová K, Greiner K, Griffith GW, Ho H-M, Hofstetter V, Jeewon R, Kang JC, Wen T-C, Kirk PM, Kytövuori I, Lawrey JD, Xing J, Li H, Liu ZY, Liu XZ, Liimatainen K, Lumbsch HT, Matsumura M, Moncada B, Nuankaew S, Parnmen S, de Azevedo Santiago ALCM, Sommai S, Song Y, de Souza CAF, de Souza-Motta CM, Su HY, Suetrong S, Wang YW, Wei S-F, Wen TC, Yuan HS, Zhou LW, Réblová M,

- Fournier J, Camporesi E, Luangsa-ard JJ, Tasanathai K, Khonsanit A, Thanakitpipattana D, Somrithipol S, Diederich P, Millanes AM, Common RS, Stadler M, Yan JY, Li XH, Lee HW, Nguyen TTT, Lee HB, Battistin E, Marsico O, Vizzini A, Vila J, Ercole E, Eberhardt U, Simonini G, Wen H-A, Chen X-H, Miettinen O, Spirin V, Hernawati 2015 – Fungal diversity notes 111–252 taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 75(1), 27–274.
- Arrillaga P, Parra LA. 2006 – género *Agaricus* L. en España. XI, *Agaricus subrufescens*, primera cita para España. *Bol. Soc. Micol. Madrid* 25: 201-207 - *Revue du Cercle de Mycologie de Bruxelles* 7, 48–51.
- Barros L, Baptista P, Correia DM, Casal S, Oliveira B, Ferreira ICFR. 2007 – Fatty acid and sugar compositions and nutritional value of five wild edible mushrooms from Northeast Portugal. *Food Chemistry* 105, 140–145.
- Bas C. 1991 – A short introduction to the ecology, taxonomy and nomenclature of the genus *Agaricus*. In: Van Grievsen LJLD editor, *Genetics and breeding of Agaricus*. Proceedings of the First International Seminar on Mushroom Science, Pudoc Wageningen, The Netherlands, 21–24.
- Bau T, Bao HY, Li Y. 2014 – A revised checklist of poisonous mushrooms in China. *Mycosystema* 33: 517–548.
- Berkeley MJ. 1842 – Enumeration of fungi, collected by H. Cuming, Esq. F.L.S. in the Philippine Islands. *London Journal of Botany* 1(3), 142–157.
- Berkeley MJ. 1847 – Decades of fungi. Decade XV–XIX. Ceylon fungi. *London Journal of Botany* 6, 479–514.
- Berkeley MJ. 1854 – Decades of fungi, Decades XLI– XLIII, Indian fungi. *Hooker's Journal of Botany and Kew Garden Miscellany* 6, 129–143.
- Berkeley MJ, Broome CE. 1870 [1871] – On some species of the genus *Agaricus* from Ceylon. *Journal of the Linnean Society of London, Botany* 27, 149–152.
- Berkeley MJ, Broome CE. 1871 – The Fungi of Ceylon. *Journal of the Linnean Society of London, Botany* 11(56), 494–567.
- Berkeley MJ, Curtis MA. 1860 – Characters of new fungi, collected in the North Pacific Exploring Expedition by Charles Wright. *Proceedings of the American Academy of Arts and Sciences* 4, 111–130.
- Bernard G. 1882 – Champignons observés à la Rochelle et dans les environs, 1–300
- Boa ER. 2006 – "Champignons Comestibles Sauvages" [Edible Wild Mushrooms] (in French), Food and Agriculture Organization of the United Nations, 141.
- Bohus G. 1974 – *Agaricus* studies IV. *Annales Historico–Natureles Musei Nationalis Hungarici* 66, 77–85.
- Bohus G. 1975 – *Agaricus* studies V (Basidiomycetes, Agaricaceae). *Annales Historico Naturales Musei Nationalis Hungarici* 67, 37–40.
- Bohus, G. 1978 – The introduction of *Agaricus macrosporoides* into cultivation. *Acta Agronomica Academiae Scientiarum Hungaricae* 27, 282–315.
- Bohus G. 1987 – Supplementary investigations concerning the introduction of *Agaricus macrosporoides* into cultivation. *Acta Agron* 36, 245–248.
- Bohus G. 1990 – *Agaricus* studies, X (Basidiomycetes, Agaricaceae). *Annales Historico Naturales Musei Nationalis Hungarici* 81, 37–44.
- Bohus G. 1995 – *Agaricus* Studies XIII. *Hungarian Mycological Society* 34, 5–37.
- Bon M. 1987 – The Mushrooms and Toadstools of Britain and North–Western Europe. Hodder & Stoughton, 278.
- Callac P. 2007 – El género *Agaricus*, In JE Sanchez, D. Royse, and H Leal Lara (eds). *Cultivo mercadotecnia e inocuidad alimenticia de Agaricus bisporus* 19–36.
- Callac P, Billette C, Imbernon M, Kerrigan RW. 1993 – Morphological, genetic, and interfertility analyses reveal a novel, tetrasporic variety of *Agaricus bisporus* from the Sonoran Desert of California. *Mycologia* 85, 335–351.

- Callac P, Guinberteau J. 2005 – Morphological and molecular characterization of two novel species of *Agaricus* section *Xanthodermatei*. *Mycologia* 97, 416–424.
- Callac P, Jacobe HI, Imbernon M, Guinberteau J, Theochari I. 2003 – A novel homothallic variety of *Agaricus bisporus* comprises rare tetrasporic isolates from Europe. *Mycologia* 95, 222–231.
- Callac P, Spataro C, Lataillade E, Blasi P, Guinberteau J. 2005 – *Agaricus devoniensis* complex comprises a group of heterothallic isolates constituting a basis for breeding in: *Genetics and Cellular Biology of Basidiomycetes VI*. Pisabarro AG, Ramirez L (Eds). Universidad Pública de Navarra. Pamplona Spain, 273.
- Callac P, Theochari I, Kerrigan RW. 2002 – The germplasm of *Agaricus bisporus*: main results after ten years of collecting in France, in Greece, and in North America. *Acta Horticulturae* 579, 49–55.
- Cappelli A. 1984 – *Agaricus* L.: Fr. incl. colour plates. Libreria editrice Biella Giovanna, I – 21047, Saronno, Italy, 558.
- Cappelli A. 2011 – Approccio al genere *Agaricus* – V *Rivista di Micologia* 2, 99–119.
- Challen MP, Kerrigan RW, Callac P. 2003 – A phylogenetic reconstruction and emendation of *Agaricus* section *Duploannulatae*. *Mycologia* 95, 61–73.
- Chandrasrikul A, Suwanarit P, Sangwanit U, Lumyong S, Payapanon A, Sanoamuang N, Pukahuta C, Petcharat V, Sardud U, Duengkae K, Klinhom U, Thongkantha S, Thongklam S. 2011 – Checklist of Mushrooms (Basidiomycetes) in Thailand. Office of Natural Resources and Environmental Policy and Planning, Bangkok, Thailand, 448.
- Chang ST. 1999 – World production of cultivated edible and medicinal mushrooms in 1997 with emphasis on *Lentinus edodes* (Berk.) Sing. in China. *International Journal Medicinal Mushrooms* 1, 291–300.
- Chen J, Moinard M, Xu J, Wang S, Foulongne-Oriol M, Zhao R, Hyde KD, Callac P. 2016a – Genetic analyses of the internal transcribed spacer sequences suggest introgression and duplication in the medicinal mushroom *Agaricus subrufescens*. *Plos One* 11(5), e0156250. <http://dx.doi.org/10.1371/journal.pone.0156250>
- Chen J, Parra LA, De Kesel A, Khalid AN, Qasim T, Aisha A, Bahkali AH, Hyde KD, Zhao RL, Callac P. 2016b – Inter- and intra-specific diversity in *Agaricus endoxanthus* and allied species reveals a new taxon, *A. punjabensis*. *Phytotaxa* 252, 1–16.
- Chen J, Parra LA, Guelly A, Rapior S, Hyde KD, Zao RL, Callac P 2015 – *Agaricus* section *Brunneopicti*: a phylogenetic reconstruction and emendation with commentary on taxa. *Phytotaxa*, 192, 145–168.
- Chen J, Zhao RL, Karunarathna SC, Callac P, Raspé O, Bahkali AL, Hyde KD. 2012 – *Agaricus megalosporus*: a new species in section *Minores*. *Cryptogamie Mycologie* 33(2), 145–155.
- Chen S, Oh S, Phung S, Hur G, Ye JJ, Kwok SL, Williams D. 2006 – Anti-aromatase activity of phytochemicals in white button mushrooms (*Agaricus bisporus*). *Cancer Research* 66, 12026–12034.
- Copeland EB. 1905 – Fungi esculentes Philippinenses. *Annales Mycologici* 3, 25–29.
- Couvy J. 1974 – La fructification d'*Agaricus bisporus* en milieu aseptique: un modele experimental pour l'etude des substances impliquees dans l'initiation fructifere. *Mushroom Science* 9/1, 157–164.
- Dabbour I, Takruri HR. 2002 – Protein Digestibility using Corrected Amino Acid Score method (PDCAAS) of four types of mushrooms in Jordan. *Plant Foods for Human Nutrition* 57(1), 13–24.
- Dai YC, Yang ZL, Cui BK, Yu CJ, Zhou LW, 2009 – Species diversity and utilization of medicinal mushrooms and fungi in China (Review). *International Journal of Medicinal Mushrooms* 11: 287–302.
- Dai YC, Zhou LW, Yang ZL, Wen HA, Bao T, Li TH, 2010 – A revised checklist of edible fungi in China. *Mycosystema* 29: 1–21.
- De Haro L, Prost N, Perringue C, Arditti J, David JM, Drouet G, Thomas M, Valli M. 1999 –

- Intoxication par champignons. Expérience du Centre anti-poisons de Marseille en 1994 et 1998. *Bulletin Epidémiologique Hebdomadaire* 30, 125–127
- De Silva DD, Rapior S, Fons F, Bahkali AH, Hyde KD. 2012a – Medicinal mushrooms in supportive cancer therapies: an approach to anticancer effects and putative mechanisms of action: a review. *Fungal Diversity* 55, 1–35.
- De Silva DD, Rapior S, Hyde KD, Bahkali AH. 2012b – Medicinal mushrooms in prevention and control of diabetes mellitus; a review. *Fungal Diversity* 56, 1–29.
- De Silva DD, Rapior S, Sudarman E, Stadler M, Xu JC, Alias SA, Hyde KD. 2013 – Bioactive metabolites from macrofungi: ethnopharmacology, biological activities and chemistry. *Fungal Diversity* 62, 1–40.
- Dias ES, Abe C, Schwan RF. 2004 – Truths and myths about the mushroom *Agaricus blazei*. *Scientia Agricola (Piracicaba, Braz.)* 61(5), 545–549.
- Donk MA 1962 – The generic names proposed for Agaricaceae. *Beihefte zur. Nova Hedwigia* 5, 1–320.
- Elliott TJ. 1978 – Comparative sexuality in *Agaricus* species. *Journal of General Microbiology* 107, 115–122.
- Esteve-Raventós F. 1998 – *Agaricus heinemannianus*, a new species of section *Minores*. *Belgian Journal of Botany* 131(2), 163–168.
- Fermor TR. 1982 – *Agaricus macrosporus*: An edible fungus with commercial potential. *Scientia Horticulturae* 16, 273–282.
- Firenzuoli F, Gori L, Lombardo G. 2008 – The medicinal mushroom *Agaricus blazei* Murrill: Review of literature and pharmaco-toxicological problems. *Evidence-Based Complementary and Alternative Medicine* 5(1), 3–15.
- Flower L, Hosagoudar VB, Abraham TK. 1997 – *Xanthagaricus*, a new generic name in the family Agaricaceae. *New Botanist* 24, 93–100.
- Fortes RC, Carvalho MR. 2011 – The effects of *Agaricus sylvaticus* fungi dietary supplementation on the metabolism and blood pressure of patients with colorectal cancer during post surgical phase. *Nutrición Hospitalaria*, núm. 1, enero–febrero 26, 176–186.
- Fortes RC, Novaes MRCG, Recova VL, Melo AL. 2009 – Immunological, hematological, and glycemia effects of dietary supplementation with *Agaricus sylvaticus* on patients' colorectal cancer. *Experimental Biology and Medicine* 234, 53–62.
- Fortes RC, Recôva VL, Melo AL, Novaes MRCG. 2008 – Effects of Dietary Supplementation with Medicinal Fungus in Fasting Glycemia Levels of Patients with colorectal Cancer: a Randomized, Double-blind, Placebo-controlled Clinical Study. *Nutrición Hospitalaria* 23(6), 591–598.
- Foulongne-Oriol M, Rocha de Brito M, Cabannes D, Clément A, Spataro C, Moinard M, Souza Dias E, Callac P, Savoie J-M. 2016 – The genetic linkage map of the medicinal mushroom *Agaricus subrufescens* reveals highly conserved macrosynteny with the congeneric species *Agaricus bisporus*. *G3* 6(5), 1217 – 1226.
- Freeman AEH. 1979a – *Agaricus* in North America: type studies. *Mycotaxon* 8, 1–49.
- Freeman AEH. 1979b – *Agaricus* in the southeastern United States. *Mycotaxon* 8, 50–118.
- Fukushima M, Nakano Y, Morii Y, Ohashi T, Fujiwara Y, Sonoyama K. Hepatic LDL. 2000 – receptor mRNA in rats is increased by dietary mushroom (*Agaricus bisporus*) fiber and sugar beet fiber. *Journal of Nutrition* 130, 2151–2156.
- Gallagher AM, Flatt PR, Duffy G, Abdel-Wahab YHA. 2003 – The effects of traditional antidiabetic plants on in vitro glucose diffusion. *Nutrition Research* 23, 413–424.
- Galli R. 2004 – *Gli Agaricus* : atlante pratico-monografico per la determinazione del genere *Agaricus* L.: Fr., Milano. Dalla Natura, 247.
- Garcia MA, Alonso J, Melgar MJ. 2005 – *Agaricus macrosporus* as a potential bioremediation agent for substrates contaminated with heavy metals. *Journal of Chemical Technology and Biotechnology* 80, 325–330.
- Geml J, Geiser DM, Royse DJ. 2004 – Molecular evolution of *Agaricus* species based on ITS and

- LSU rDNA sequences. *Mycological Progress* 3(2), 157–176.
- Geml J, Laursen GA, Nusbaum HC, Taylor DL. 2007 – Two new species of *Agaricus* from the Subantarctic. *Mycotaxon* 100:193–208.
- Geml J, Rimóczi I. 1999 – Attempts at cultivating wild strains of various *Agaricus* species. *Fungal Genetics Newsletter* 46, 38.
- Geml J, Royse DJ. 2002 – Molecular phylogeny and cultivation of *Agaricus* species. In: *Proceedings of the IV International Conference on Mushroom Biology and Mushroom*.
- Genevier LG. 1876 – Étude sur les champignons consommés à Nantes sous le nom de champignon rose ou de couche (*Agaricus campestris* L.). *Bulletin de la Société Botanique de France* (in French) 23, 28–35.
- Gennari JL. 2000a – Casos clínicos sobre o emprego do cogumelo *Agaricus sylvaticus* no câncer e na AIDS. *Journal of Biomolecular Medicine & Free Radicals* 6, 9–11.
- Gennari JL. 2000b – *Agaricus sylvaticus*. *Journal of Biomolecular Medicine & Free Radicals* 6, 35–36.
- Gennari JL, Gennari M, Felipe JO. 2001 – *Agaricus sylvaticus* aumenta o número de células Natural Killer em pacientes com câncer. *Revista de Medicina Complementar* 7, 42.
- Geösel A. 2011 – New developments in integrated pest management for mushroom culture, challenges and opportunities in quality mushroom production. *7th International Conference on Mushroom Biology and Mushroom Products, Volume 1*.
- Gill, Strauch. 1984 – Constituents of *Agaricus xanthodermus* Genevier: the first naturally endogenous azo compound and toxic phenolic metabolites, *Zeitschrift Fur Naturforschung. C Journal of Biosciences* 39, 1027–1029.
- Gray AM, Flatt PR. 1998 – Insulin-releasing and insulin-like activity of *Agaricus campestris* (mushroom). *The Journal of Endocrinology* 157 (2), 259–66.
- Gray, SF. 1821 – *A Natural Arrangement of British Plants*, volume 1. London: Baldwin, Cradock and Joy, 627.
- Grgurinovic. 1997 – *Lepiota albofibrillosa* Cleland in *Trans. Proc. roy. Soc. Australia* [type study Grgurinovic. *South Australia* 55, 154.
- Grigson J. 1975 – *The Mushroom Feast*. London, UK: Penguin, ISBN 0–14–046273–2.
- Gui Y, Zhu GS, Callac P, Hyde KD, Parra LA, Chen J, Yang TJ, Huang WB, Gong GL, Liu ZY. 2015 – *Agaricus* section *Arvenses*: three new species in highland subtropical Southwest China. *Fungal Biology* 119(2–3), 79–94.
- Hama O, Maes E, Guissou ML, Ibrahim D, Baragé M, Parra LA, Raspé O, De Kesel A, 2010 – *Agaricus subsaharianus*, une nouvelle espèce comestible et consommée au Niger, au Burkina Faso et en Tanzanie. *Cryptogamie Mycologie* 31, 221–234.
- Heinemann P. 1956 – Champignons récoltés au Congo Belge par madame M. Goossens-Fontana. II. *Agaricus* Fries s.s. *Bulletin du Jardin Botanique National de Belgique*. 26(1), 1–127.
- Heinemann P. 1971 – In: *Cahiers de La Maboké* 9(1), 6.
- Heinemann P. 1977 – Les Psalliotes. *Nat Belges* 58, 145–165.
- Heinemann P. 1978 – Essai d'une clé de détermination des genres *Agaricus* et *Micropsalliota*. *Sydowia* 30, 6–37.
- Heinemann P. 1980 – Les genres *Agaricus* et *Micropsalliota* en Malaisie et en Indonésie. *Bulletin du Jardin Botanique National de Belgique* 50, 3–68.
- Heinemann P. 1986 *Agarici austroamericani* VI. Aperçu sur les *Agaricus* de Patagonie et de la terre de Feu. *Bulletin du Jardin Botanique National de Belgique* 56: 417–446.
- Heinemann P. 1987 – Clave para la determinacion de las especies de *Agaricus* (Agaricales) de la Patagonia y Tierra del Fuego. *Darwiniana* 28, 283–291.
- Heinemann P. 1990 – *Agaricus singaporensis* sp. nov. *Bulletin du Jardin Botanique National de Belgique* 60, 417–419.
- Hildén K, Mäkelä MR, Lankinen P, Lundell T. 2013 – *Agaricus bisporus* and related *Agaricus* species on lignocellulose production of manganese peroxidase and multicopper oxidases. *Fungal Genetics and Biology* 55, 32–41.

- Hotson JW, Stuntz DE. 1938 – The genus *Agaricus* in western Washington. *Mycologia* 30, 204–234.
- Huijsman HSC. 1960 – Notes sur le genre *Agaricus*. *Persoonia* 1(3), 321–324.
- Hyde KD, Bahkali AH, Moslem MA. 2010 – Fungi—an unusual source for cosmetics, *Fungal Diversity* 43, 1–9.
- Imai S. 1938 – Studies on the Agaricaceae of Hokkaido. 1. *Journal of the Faculty of Agriculture of the Hokkaido Imperial University* 43, 1–378.
- Index Fungorum. 2016 – <http://www.indexfungorum.org/Names/Names.asp>
- Jatuwong K, Kakumyan P, Chamyuang S, Chukeatirote E, Hyde KD. 2014 – Optimization condition for cultivation of *Agaricus subrufescens* hybrid strains. The 26th Annual meeting of the Thai Society for Biotechnology and International Conference, Mae Fah Luang University, Chiang Rai, Thailand.
- Jeong SC, Jeong YT, Yang BK, Islam R, Koyyalamudi SR, Pang G, Cho KY, Song CH. 2010 – White button mushroom (*Agaricus bisporus*) lowers blood glucose and cholesterol levels in diabetic and hypercholesterolemic rats. *Nutrition Research* 30, 49–56.
- Jordan P. 2000 – *The Mushroom Guide and Identifier: The Ultimate Guide to Identifying, Picking and Using Mushrooms*. London: Hermes House, 100.
- Junghuhn FW 1840 – Nova genera et species plantarum florae javanicae. *Tijdschrift voor Natuurlijke Geschiedenis en Physiologie* 7, 285–317.
- Kalač P, Svoboda L. 2000 – A review of trace element concentrations in edible mushrooms. *Food Chemistry* 69, 273–281.
- Karunarathna SC, Guinberteau J, Chen J, Vellinga EC, Zhao R, Chukeatirote E, Yan J, Hyde KD, Callac P. 2014 – Two new species in *Agaricus* tropical clade I. *Chiang Mai Journal of Science* 41(4), 771–780.
- Kawagishi H, Inagaki R, Kanao T, Mizuno T, Shimura K, Ito H, Hagiwara T, Nakamura T 1989 – Fractionation and antitumor activity of the water-insoluble residue of *Agaricus blazei* fruiting bodies. *Carbohydrate Research* 186, 267–273.
- Kerrigan RW 1985 – Studies in *Agaricus* III: new species from California. *Mycotaxon* 20(2), 419–434.
- Kerrigan RW. 1986 – *Agaricales of California. Agaricaceae*, vol 6. Mad River Press, Eureka.
- Kerrigan RW. 1989 – Studies in *Agaricus* IV: new species from colourado. *Mycotaxon* 34, 119–128.
- Kerrigan RW. 2005 – *Agaricus subrufescens*, a cultivated edible and medicinal mushroom, and its synonyms. *Mycologia* 97(1), 12–24.
- Kerrigan RW, Callac P, Guinberteau J, Challen MP, Parra LA. 2006 – *Agaricus* section *Xanthodermatei*: a phylogenetic reconstruction with commentary on taxa. *Mycologia* 97, 1292–1315.
- Kerrigan R, Callac P, Parra L. 2008 – New and rare taxa in *Agaricus* section *Bivelares* (*Duploannulati*). *Mycologia* 100(6), 876–892.
- Kerrigan RW, Callac P, Xu J, Noble R. 1999 – Population and phylogenetic structure within the *Agaricus subfloccosus* complex. *Mycological Research* 103, 1515–1523.
- Konrad P, Maublanc A. 1952 – *Les Agaricales*. Lechevalier, Paris.
- Koyyalamudi SR, Jeong SC, Song CH, Cho KY, Pang G. 2009 – Vitamin D2 formation and bioavailability from *Agaricus bisporus* button mushrooms treated with ultraviolet irradiation. *Journal of Agricultural and Food Chemistry* 57, 3351–3355.
- Kühner P, Romagnesi H. 1953 – *Flore analytique des champignons superieurs (Agarics, Boletes, Chanterelles)*. Masson et CIE, Paris.
- Lacheva MN, Stoichev GT. 2004 – New species (records) of the genus *Agaricus* (Agaricaceae) for Bulgaria. *Mycologia Balcanica* 1, 35–40.
- Lanconelli L. 2002 – *Agaricus padanus* sp. nov. *Rivista di Micologia* 1, 29–37.
- Largeteau ML, Callac P, Navarro-Rodríguez AM, Savoie JM. 2011 – Diversity in the ability of *Agaricus bisporus* wild isolates to fruit at high temperature (25°C). *Fungal Biology* 115,

1186–1195.

- Lebel T. 2013 – Two new species of sequestrate *Agaricus* (section *Minores*) from Australia. *Mycological Progress* 12, 699–707.
- Lebel T, Syme A. 2012 – Sequestrate species of *Agaricus* and *Macrolepiota* from Australia: new species and combinations and their position in a calibrated phylogeny. *Mycologia* 104(2), 496–520.
- Lee YL, Kim HJ, Lee MS, Kim JM, Han JS, Hong EK, Kwon MS, Lee MJ. 2003 – Oral administration of *Agaricus blazei* (H1 strain) inhibited tumor growth in a sarcoma 180 inoculation model. *Experimental Animals* 52, 371–375.
- Li SF, Xi YL, Qi CX, Liang QQ, Wei SL, Li GJ, Zhao D, Li SJ, Wen HA. 2014 – *Agaricus taeniatus* sp. nov., a new member of *Agaricus* sect. *Bivelares* from northwest China. *Mycotaxon* 129(1), 187–196.
- Li Y. 1990 – New species and records of the genus *Agaricus* from China. *Acta Botanica Yunnanica* 12(2), 154–160.
- Linnaeus C. 1753 – *Species plantarum: exhibentes plantas rite cognitatas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas*, ed. 1. Laurentius Salvius, Stockholm, 1200.
- Liu JK, Hyde KD, Gareth EBG, Ariyawansa HA, Bhat DJ, Boonmee S, Maharachchikumbura S, McKenzie EHC, Phookamsak R, Phukhamsakda R, Abdel-Wahab MA, Buyck B, Chen J, Chethana KWT, Singtripop C, Dai DQ, Dai YC, Daranagama DA, Dissanayake AJ, Doilom M, D'souza MJ, Fan LX, Goonasekara ID, Hirayama K, Hongsan S, Jayasiri SC, Jayawardena RS, Karunarathna SC, Li WJ, Mapook A, Norphanphoun C, Pang KL, Perera RH, Peršoh D, Pinruan U, Senanayake IC, Somrithipol S, Satinee S, Tanaka K, Thambugala KM, Tian Q, Tibpromma S, Udayanga D, Wijayawardene NN, Wanasinghe D, Abdel-Aziz FA, Adamčík S, Bahkali AH, Boonyuen N, Bulgakov T, Callac P, Chomnunti P, Greiner K, Hashimoto A, Hofstetter V, Kang JC, Li XH, Liu ZY, Matumura M, Mortimer PE, Rambold R, Randrianjohany E, Sato G, Indrasudthi VS, Verbeken A, Brackel W, Wang Y, Wen TC, Xu JC, Yan JY, Zhao RL, Camporesi E. (2015) Fungal diversity notes 1 – 110: taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity* 72, 1 – 197
- Ludwig E. 2007 *Pilzkompendium*, Band 2, Beschreibungen, Fungicon verlag, Berlin, 723.
- Luiz RC, Jordão BQ, da Eira AF, Ribeiro LR, Mantovani MS. 2003 – Mechanism of anticlastogenicity of *Agaricus blazei* Murill mushroom organic extracts in wild type CHO (K(1)) and repair deficient (xrs5) cells by chromosome aberration and sister chromatid exchange assays. *Mutation Research* 528, 75–79.
- Machado MP, Filho ER, Terezan AP, Ribeiro LR, Mantovani MS. 2005 – Cytotoxicity, genotoxicity and antimutagenicity of hexane extracts of *Agaricus blazei* determined in vitro by the comet assay and CHO/HGPRT gene mutation assay. *Toxicology In Vitro* 19, 533–539.
- Mantovani MS, Bellini MF, Angeli JP, Oliveira RJ, Silva AF, Ribeiro LR. 2007 – Betaglucans in promoting health: Prevention against mutation and cancer. *Mutation Research* 658(3), 154–161.
- Marekov I, Momchilova S, Grung B, Damyanova BN. 2012 – Fatty acid composition of wild mushroom species of order Agaricales—examination by gas chromatography-mass spectrometry and chemometrics. *Journal of Chromatography B* 910, 54–60.
- Martins de Oliveira J, Jordão BQ, Ribeiro LR, Ferreira da Eira A, Mantovani MS. 2002 – Anti-genotoxic effect of aqueous extracts of sun mushroom (*Agaricus blazei* Murill lineage 99/26) in mammalian cells in vitro. *Food and Chemical Toxicology* 40, 1775–1780.
- Massee GE. 1901 – *Fungi exotici*, III. *Bulletin of Miscellaneous Informations of the Royal Botanical Gardens Kew* 1901, 150–169.
- Massee GE. 1907 – *Fungi exotici*, VI. *Bulletin of Miscellaneous Informations of the Royal Botanical Gardens Kew*, 123.
- Massee GE. 1912 – *Fungi exotici*, XV. *Bulletin of Miscellaneous Informations of the Royal Botanical Gardens Kew* 1912, 357–359.

- Massee GE. 1914 – Fungi exotici, XVII. Bulletin of Miscellaneous Informations of the Royal Botanical Gardens Kew 1914, 72–76.
- Mattila P, Konko K, Euroala M, Pihlava JM, Astola J, Vahteristo L, Hietaniemi V, Kumpulainen J, Valtonen M, Piironen V. 2001 – Contents of vitamins, mineral elements, and some phenolic compounds in cultivated mushrooms. *Journal of Agricultural and Food Chemistry* 49(5), 2343–2348.
- Menier 1893 – Une Nouvelle Psalliote, *Psalliota ammophila* découverte dans la Loire-Inférieure. *Bulletin de la Société des sciences naturelles de l'Ouest de la France* 3(2), 67–69.
- Menoli RC, Mantovani MS, Ribeiro LR, Speit G, Jordão BQ. 2001 – Antimutagenic effects of the mushroom *Agaricus blazei* Murrill extracts on V79 cells. *Mutation Research* 496, 5–13.
- Mitchell AD, Bresinsky A. 1999 – Phylogenetic relationships of *Agaricus* species based on ITS-2 and 28S ribosomal DNA sequences. *Mycologia* 91, 811–819.
- Mitchell AD, Walter M. 1999 – Species of *Agaricus* occurring in New Zealand. *New Zealand Journal of Botany* 37, 715–725.
- Mizuno M, Minato K, Ito H, Kawade M, Terai H, Tsuchida H. 1999 – Antitumor polysaccharide from the mycelium of liquid-cultured *Agaricus blazei* mill. *Biochemistry and Molecular Biology International* 47, 707–714.
- Mizuno T, Hagiwara T, Nakamura T, Ito H, Shimura K, Sumiya T, Asakura A. 1990 – Antitumor activity and some properties of water-soluble polysaccharides from "Himmematsutake," the fruiting of *Agaricus blazei* Murrill. *Agricultural and Biological Chemistry* 54, 2889–2896.
- Møller FH. 1950 – Danish *Psalliota* species. Preliminary Studies for a Monograph on the Danish Psalliotae. Part 1. *Friesia* 4(1–2), 1–60.
- Møller FH. 1952 – Danish *Psalliota* Species. Preliminary Studies for a Monograph on the Danish Psalliotae. Part 2. *Friesia* 4(3), 135–242.
- Moser M. 1967–1983 – Kleine Kryptogamenflora: Die Röhrlinge und Blatterpilze (Polyporales, Boletales, Agaricales, Russulales). Gustav Fischer Verlag, Stuttgart.
- Murrill WA. 1912 – The Agaricaceae of the Pacific Coast III. *Mycologia* 4, 294–300.
- Murrill WA. 1918 – The Agaricaceae of tropical North America, VIII. *Mycologia* 10, 62–85.
- Murrill WA. 1941 – More Florida novelties. *Mycologia* 33, 434–448.
- Natarajan K, Kumaresan V, Narayanan K. 2005 – A checklist of Indian Agarics and Boletes (1984–2002). *Kavaka* 33, 61–128.
- Natarajan K, Purushothama KB. 1996 – South Indian Agaricales XXIII, *Kavaka* 22/23, 47–51.
- Nauta MM. 1999 – Notulae ad floram Agaricinam neerlandicam–XXXIII: notes on *Agaricus* section *Spissicaules*. *Persoonia* 17(2), 221–233.
- Nauta MM. 2000 – Notulae ad floram Agaricinam neerlandicam–XXXVII: notes on *Agaricus* section *Arvenses*. *Persoonia* 17(3), 457–463.
- Noble R, Grogan H, Elliott T. 1995 – Variation in morphology, growth, and fructification of isolates in the *Agaricus subfloccosus* complex. *Mycological Research* 99, 1453–1461.
- Orton PD. 1960 – New check-list of British Agarics and Boleti. Part iii. Notes on genera and species in the list. *Transactions of the British mycological Society* 43(2), 159–439.
- Öztürk M, Duru ME, Kivrak S, Mercan-Doğan N, Türkoglu A, Özler MA. 2011 – In vitro antioxidant, anticholinesterase and antimicrobial activity studies on three *Agaricus* species with fatty acid compositions and iron contents: a comparative study on the three most edible mushrooms. *Food and Chemical Toxicology* 49, 1353–1360
- Paracer CS, Chahal DS. 1963 [1962] – A new edible species of genus *Agaricus* in the Punjab (India), *Mycopathologia et Mycologia Applicata* 18(4), 267–270.
- Parra LA. 2003 – Contribution to the knowledge of the genus *Agaricus* [Fingi Delineati. Pars XXIV]. Edizioni Candusso, Alassio.
- Parra LA. 2008 – Fungi Europaei, Volume 1: *Agaricus: Allopsalliota*, Nauta and Bas. Candusso Edizioni, Alassio, 824.
- Parra LA. 2013 – Fungi Europaei, Volume 1A: *Agaricus: Allopsalliota*, Nauta and Bas. Candusso Edizioni s.a.s. Alassio, 1168.

- Parra LA, Mua A, Cappelli A, Callac P. 2011 – *Agaricus biannulatus* sp. nov., a new species of the section *Xanthodermatei* collected in Sardinia and Sicily. *Micologia e Vegetazione Mediterranea* 26 (1), 3–20.
- Parra LA, Muñoz G, Callac P. 2014 – *Agaricus caballeroi* sp. nov., una nueva especie de la sección *Nigrobrunnescentes* recolectada en España. *Micologia e Vegetazione Mediterranea* 29(1), 21–38.
- Passecker. 1932 – *Zeitschr. Pilzkunde*, N. F. 11, 39.
- Patouillard NT 1913 – Quelques champignons du Tonkin. *Bulletin de la Société Mycologique de France* 29, 206–228.
- Patrick H. 2008 – *Mushroom Miscellany*. Collins. ISBN 978–0–00–728464–1.
- Peck CH. 1893 – *New York State. Mushroom Annual Report* 46, 83–152.
- Pegler DN. 1977 – A preliminary Agaric flora of east Africa. *Kew Bulletin Additional Series* 6, 1–615.
- Pegler DN. 1983 – *The Genus Lentinus: A World Monograph* HMSO, London.
- Pegler DN. 1986 – Agaric flora of Sri Lanka. *Kew Bulletin Additional Series* 12, 1–519.
- Pegler DN. 1990 – Agaricales of Brazil described by J.P.F.C. Montagne. *Kew Bulletin* 45, 161–177.
- Petch T. 1917 – Revision of Ceylan fungi. *Annals of the Royal Botanic Gardens Peradeniya* 6, 307–355.
- Petch T. 1925 – Additions to Ceylon fungi. III. *Annals of the Royal Botanic Gardens Peradeniya* 9(3), 313–328.
- Peterson KR, Desjardin ED, Hemmes Don E. 2000 – Agaricales of Hawaiian Islands. 6. Agaricaceae L. Agaricaceae: *Agaricus* and *Melanophyllum*. *Sydowia* 52(2), 204–257.
- Pilat A. 1951 – The Bohemian species of the genus *Agaricus*. *Acta Entomologica Musei Nationalis Pragae*, VII B.
- Pilát A, Ušák O. 1961 – Champignons Comestibles et Veneneux by Marcel Locquin, Bengt Cortin; *A Handbook of Mushrooms* 53(3), 314–316.
- Poppe J. 1972 – Un excellent *Agaricus* tetra-sporique ultivable commercialement avec succès. *Mushroom Science* 8, 517–525.
- Rapier S, Breheret S, Talou T, Pelissier Y, Bessiere JM, 2002 – The anise-like Odour of *Clitocybe Odoura*, *Lentinellus cochleatus* and *Agaricus essettei*. *Mycologia* 94, 373–376.
- Ren Z, Guo Z, Meydani SN, Wu D. 2008 – White button mushroom enhances maturation of bone marrow-derived dendritic cells and their antigenpresenting function in mice. *Journal of Nutrition* 138, 544–50.
- Saccardo PA, Trotter A. 1913 – *Supplementum Universale, Pars IX. Sylloge Fungorum* 22, 1–1612.
- Saini SS, Atri NS, Gupta AK. 1997 – Studies on the genus *Agaricus* L.: Fr.: the subgenus *Agaricus* section *Sanguinolenti* Schaeff et Motter from northwest India. *Mushroom Research* 6(2), 53–58.
- Savoie JM, Foulongne-Oriol M, Barroso G, Callac P. 2013 – Genetics and Genomics of cultivated mushrooms, Application to breeding of *Agarics*. In: *The Mycota, Agricultural Applications* Second ed, Kempken, F., vol. ed., Springer-Verlag, Berlin, Heidelberg 11, 3–33.
- Singer R. 1986 – *The Agaricales in modern taxonomy*. 4th edn. Königstein, Koeltz, 981.
- Smiderle FR, Andrea CR, Arkel J, Chanpu W, Iacomini M, Wichers HJ, Leo JLD. 2011 – Polysaccharides from *Agaricus bisporus* and *Agaricus brasiliensis* show similarities in their structures and their immunomodulatory effects on human monocytic THP-1 cells. *BMC Complementary and Alternative Medicine* 11, 58.
- Smith. 1944 – Interesting North American agarics. *Bulletin of the Torrey Botanical Club* 71, 390–409.
- Spegazzini C. 1919 – *Reliquiae mycologicae tropicae*. *Boletín de la Academia Nacional de Ciencias en Córdoba* 23(3–4), 365–541.

- Stadler M, Hellwig V, Mayer–Barttschmid A, Denzer D, Wiese B, Burkhardt N. 2005 – Novel analgesic triglycerides from cultures of *Agaricus macrosporus* and other basidiomycetes as selective inhibitors of neurolysin, *The Journal of Antibiotics* 58, 775–786.
- Taveira VC, Novaes MRCG, Reis MA, Silva MF. 2008 – Hematologic and Metabolic Effects of Dietary Supplementation with *Agaricus sylvaticus* Fungi on Rats Bearing Solid Walker 256 Tumor. *Experimental Biology and Medicine* 233(11), 1341–1347.
- Thawthong A, Karunarathna SC, Thongklang N, Chukeatirote E, Kakumyan P, Chamyuang S, Rizal M, Mortimer, PE, Xu, JC, Callac P, Hyde KD. 2014 – Discovering and Domesticating Wild Tropical Cultivable Mushrooms. *Chiang Mai Journal of Science* 41, 731–764.
- Thongklang N. 2016a – Phylogeny of *Agaricus* section *Arvenses*, Biochemistry and Domestication of selected species, PhD thesis, Mae Fah Luang University, Chiang Rai, Thailand.
- Thongklang N, Chen J, Bandara AR, Hyde KD, Raspé O, Parra LA, Callac P. 2016b – Studies on *Agaricus subtilipes*, a new cultivatable species from Thailand, incidentally reveal the presence of *Agaricus subrufescens* in Africa. *Mycoscience* 4, 239–250
- Thongklang N, Nawaz R, Khalid AN, Chen J, Hyde KD, Zhao RL, Parra LA, Hanif M, Moinard M, Callac P. 2014a – Morphological and molecular characterization of three *Agaricus* species from tropical Asia (Pakistan, Thailand) reveals a new group in section *Xanthodermatei*. *Mycologia* 106(6):1220-32.
- Thongklang N, Sysouphanthong P, Callac P, Hyde KD. 2014b – First cultivation of *Agaricus flocculosipes* and a novel Thai strain of *A. subrufescens*. *Mycosphere* 5(6), 814–820.
- Thongklang N, Sysouphanthong P, Hyde KD, Moinard M, Hoang E, Rodriguez A, Kerrigan RW, Foulongne–Oriol M, Callac P. 2014c – Evidence for amphithallism and broad geographical hybridization potential among *Agaricus subrufescens* isolates from Brazil, France and Thailand. *Fungal Biology* 118(12), 1013–1023.
- Trotter A. 1972 – *Sylloge Fungorum*, Johnson Reprint Corporation, New York, London 26, 1–1563.
- Valenzuela E, Heinemann P, Esteve–raventos FS, Polette GM, Castells W. 1997 – A new contribution to the knowledge of the genus *Agaricus* in Chile. *Agaricus wrightii* sp. nov. *Mycotaxon* 63, 71–76.
- Vienna Code 2006 – International Code of Botanical Nomenclature (Vienna Code), Regnum Vegetabile 146. A.R.G. Gantner Verlag KG. ISBN 3-906166-48-1
- Volman JJ, Helsper JPF, Wei S, Baars JJP, Van Griensven LJLD, Sonnenberg ASM, Mensink RP, Plat J. 2010 – Effect of mushroom–derived β -glucan–rich polysaccharide extracts on nitric oxide production by bone marrow–derived macrophages and nuclear factor– κ B transactivation in Caco–2 receptor cells: Can effects be explained by structure? *Molecular Nutrition and Food Research* 54, 268–276.
- Walton GJ. 1986 – *Edible and poisonous mushrooms of Canada*, 2nd Edition, Ottawa: Research Branch, Agriculture Canada, 325. ISBN 10:066010136X.
- Wasser SP. 1980 – *Agaricaceae Cohn.*, *Flora Fungorum RSS Ucrainicae*, Naukova Dumka. Kiev.
- Wasser SP. 1989 – *Tribe Agariceae Pat. of the Soviet Union*. Koenigstein, Koeltz Scientific Books, 120.
- Wang ZR, Parra L, Callac P, Zhou JL, Fu WJ, Dui SH, Hyde KD, Zhao RL. 2015 – Edible species of *Agaricus* (Agaricaceae) from Xinjiang Province (Western China). *Phytotaxa*, 202, 185–197.
- Wei-Fan C. 1973 – Ten new species of Agaricales from Yunnan. *Acta Microbiologica Sinica* 13(2), 130.
- Wisitrasameewong K, Karunarathna SC, Thongklang N, Zhao RL, Callac P, Chukeatirote E, Bahkali AH, Hyde KD. 2012a – *Agaricus subrufescens*: new records to Thailand. *Chiang Mai University Journal of Science* 39(2), 281–291.
- Wisitrasameewong K, Karunarathna SC, Thongklang N, Zhao RL, Callac P, Moukha S, Férandon C, Chukeatirote E, Hyde KD. 2012b – *Agaricus subrufescens*: A review. *Saudi Journal of Biological Sciences* 19(2), 131–146.

- Wood WF, Watson RL, Largent DL. 1998 – Phenol, the odour compound from *Agaricus praeclaresquamosus*. *Biochemical Systematics and Ecology* 26, 793–794.
- Wu D, Pae M, Ren Z, Guo Z, Smith D, Meydani SN. 2007 – Dietary supplementation with white button mushroom enhances natural killer cell activity in C57BL/6 mice. *Journal of Nutrition* 137(6), 1472–1477.
- Xu J, Kerrigan RW, Sonnenberg A, Callac P, Horgen PA, Anderson JB. 1998 – Mitochondrial DNA variation in natural populations of the mushroom *Agaricus bisporus*. *Molecular Ecology* 7, 19–33.
- Yamac M, Kanbak G, Zeytinoglu M, Senturk H, Bayramoglu G, Dokumacioglu A, Van Griensven LJLD. 2010 – Pancreas protective effect of antioxidative *Agaricus bisporus* extract on rats with streptozotocin induced diabetes. *International Journal of Medicinal Mushrooms* 12 (4), 379–389.
- Yu L, Fernig DG, Smith JA, Milton JD, Rhodes JM. 1993 – Reversible inhibition of proliferation of epithelial cell lines by *Agaricus bisporus* (edible mushroom) lectin. *Cancer Research* 53, 4627–32.
- Zhang JX, Chen Q, Huang CY, Gao W, Qu JB. 2015 – History, current situation and trend of edible mushroom industry development. *Mycosystema* 34: 524–540.
- Zhao RL. 2008 – Systematics of *Agaricus*, *Cyathus* and *Micropsalliota* in Northern Thailand. PhD thesis, KMITL, Bangkok, Thailand.
- Zhao RL, Desjardin DE, Callac P, Parra LA, Guinberteau J, Soyong K, Karunarathna S, Zhang Y, Hyde KD. 2012a – Two species of *Agaricus* sect. *Xanthodermatei* from Thailand. *Mycotaxon* 122, 187–195.
- Zhao RL, Hyde KD, Desjardin DE, Raspe O, Soyong K, Guinberteau J, Karunarathna SC, Callac P. 2012b – *Agaricus flocculosipes* sp. nov., a new potentially cultivatable species from the palaeotropics. *Mycoscience* 53, 300–311.
- Zhao RL, Karunarathna SC, Raspé O, Parra LA, Guinberteau J, Moinard M, De Kesel A, Barroso G, Courtecuisse R, Hyde KD, Gelly AK, Desjardin ED, Callac P. 2011 – Major clades in tropical *Agaricus*. *Fungal Diversity* 51, 279–296.
- Zhao RL, Zhou JL, Chen J, Margaritescu S, Sánchez-Ramírez S, Hyde KD, Callac P, Parra L A, Li GJ, Moncalvo JM. 2016 – Towards standardizing taxonomic ranks using divergence times – a case study for reconstruction of the *Agaricus* taxonomic system. *Fungal diversity* 78, 239–292.