

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***N'*-(1*E*)-(4-Fluorophenyl)methylene]-6-methoxy-2-naphthohydrazide**H. S. Yathirajan,^a B. Narayana,^a K. Sunil,^a B. K. Sarojini^c and Michael Bolte^{d*}^aDepartment of Studies in Chemistry, University of Mysore, Manasagangothri, Mysore 570 006, India, ^bDepartment of Studies in Chemistry, Mangalore University, Mangalagangothri 574 1999, India, ^cDepartment of Chemistry, P. A. College of Engineering, Nadupadavu, Mangalore 574 153, India, and ^dInstitut für Anorganische Chemie, J. W. Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, 60438 Frankfurt/Main, Germany

Correspondence e-mail: bolte@chemie.uni-frankfurt.de


Received 5 April 2007; accepted 12 April 2007

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.063; wR factor = 0.127; data-to-parameter ratio = 7.1.

The title compound, $\text{C}_{19}\text{H}_{15}\text{FN}_2\text{O}_2$, is a Schiff base which has been synthesized by a condensation reaction of 6-methoxy-2-naphthohydrazide and 4-fluorobenzaldehyde. The molecule is almost planar with the $\text{C}=\text{N}$ double bond in a *trans* configuration. The crystal packing is stabilized by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds that link molecules into chains running along the c axis. There are also weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds as well as $\text{C}-\text{H}\cdots\pi$ -ring interactions in the crystal structure.

Related literature

Some related structures of hydrazides such as 2'-(4-dimethylaminobenzylidene)pyrazine-2-carbohydrazide (Shi & Yuan, 2006), *N'*-(3-ethoxy-4-hydroxybenzylidene)isonicotinohydrazide (Qian *et al.*, 2006) and 2'-(1,3-benzodioxol-5-ylmethylene)-2-methoxybenzo-hydrazide (Jing & Yu, 2007) and a Schiff base (Yathirajan *et al.*, 2007) have been reported.



Experimental

Crystal data

 $\text{C}_{19}\text{H}_{15}\text{FN}_2\text{O}_2$
 $M_r = 322.33$ Monoclinic, Pc
 $a = 15.681$ (2) Å $b = 5.6574$ (7) Å
 $c = 9.2758$ (10) Å
 $\beta = 104.746$ (9)°
 $V = 795.79$ (17) Å³
 $Z = 2$ Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 173$ (2) K
 $0.24 \times 0.09 \times 0.08$ mm

Data collection

Stoe IPDSII two-circle diffractometer
Absorption correction: none
9090 measured reflections1531 independent reflections
1246 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.069$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.063$
 $wR(F^2) = 0.127$
 $S = 1.57$
1531 reflections
217 parameters2 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.25$ e Å⁻³**Table 1**
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^{\text{i}}$	0.88	2.11	2.778 (4)	132
$\text{C23}-\text{H23}\cdots\text{O2}^{\text{ii}}$	0.95	2.55	3.428 (6)	154
$\text{C26}-\text{H26}\cdots\text{O1}^{\text{iii}}$	0.95	2.50	3.403 (6)	158
$\text{C14}-\text{H14}\cdots\text{Cg1}^{\text{i}}$	0.98	2.89	3.666 (5)	140
$\text{C19}-\text{H19}\cdots\text{Cg2}^{\text{iv}}$	0.93	2.95	3.697 (5)	136

Symmetry codes: (i) $x, -y + 1, z - \frac{1}{2}$; (ii) $x - 1, y - 1, z$; (iii) $x, -y, z - \frac{1}{2}$; (iv) $x, -y + 2, z + \frac{1}{2}$. Cg1 and Cg2 are the centroids of the rings $\text{C11/C12/C13/C18/C19/C20}$ and $\text{C13/C14/C15/C16/C17/C18}$, respectively.

Data collection: *X-Area* (Stoe & Cie, 2001); cell refinement: *X-Area*; data reduction: *X-Area*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1990); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON* (Spek, 2003) and *XP* in *SHELXTL-Plus* (Sheldrick, 1991); software used to prepare material for publication: *SHELXL97*.

BN thanks Mangalore University for research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FB2048).

References

- Jing, Z.-L. & Yu, M. (2007). *Acta Cryst.* **E63**, o509–o510.
Qian, H.-Y., Yin, Z.-G., Jia, J., Liu, S.-M. & Feng, L.-Q. (2006). *Acta Cryst.* **E62**, o3623–o3624.
Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
Sheldrick, G. M. (1991). *SHELXTL-Plus*. Release 4.1. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
Shi, X.-F. & Yuan, C.-C. (2006). *Acta Cryst.* **E62**, o3290–o3291.
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
Stoe & Cie (2001). *X-Area*. Stoe & Cie, Darmstadt, Germany.
Yathirajan, H. S., Sarojini, B. K., Narayana, B., Sunil, K. & Bolte, M. (2007). *Acta Cryst.* **E63**, o1398–o1399.

supplementary materials

Acta Cryst. (2007). E63, o2565 [doi:10.1107/S160053680701820X]

N'-[(1*E*)-(4-Fluorophenyl)methylene]-6-methoxy-2-naphthohydrazide

H. S. Yathirajan, B. Narayana, K. Sunil, B. K. Sarojini and M. Bolte

Comment

The title compound, C₁₉H₁₅FN₂O₂, is a Schiff base which has been synthesized by a condensation reaction of 6-methoxy-2-naphthohydrazide and 4-fluorobenzaldehyde.

A view of (I) is shown in Fig. 1. Bond lengths and angles can be regarded as normal (Cambridge Crystallographic Database, Version 5.28, November 2006; updated January 2007 (Allen, 2002); Mogul Version 1.1 (Bruno *et al.*, 2004)). The molecule is almost planar (r.m.s. deviation for all non-H atoms = 0.123 Å) with the C=N double bond in a trans configuration. The crystal packing is stabilized by N—H···O hydrogen bonds that link molecules into chains running along the *c*-axis (Tab. 1 and Fig. 2). There are also C—H···O weak hydrogen bonds (Tab. 1) as well as C—H··· π -ring interactions in the structure (Tab. 2).

Experimental

A mixture of 6-methoxy-2-naphthohydrazide (1.08 g, 0.005 mol) and 4-fluorobenzaldehyde (0.62 ml, 0.005 mol) in 15 ml of absolute ethanol containing 2 drops of 4 M sulfuric acid was refluxed for about 3 hours. On cooling to room temperature, a solid precipitated. The solid was filtered off, redissolved in 10 ml of a mixture (1:1) of dimethylformamide and acetone, and then recrystallized. (M. p.: 492-494 K). Analysis for C₁₉H₁₅FN₂O₂: Found (Calculated): C: 70.68 (70.80); H: 4.63 (4.69); N: 8.62 (8.69) weight %.

Refinement

All the H atoms were found in a difference Fourier map. Nevertheless, the hydrogens were refined using a riding model with N—H = 0.88 Å, C—H = 0.95 Å or C_{methyl}—H = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$. In the absence of anomalous scatterers 1517 Friedel pairs have been merged prior to refinement.

Figures




Fig. 1. A view of the title compound with the atom numbering. The displacement ellipsoids are at the 50% probability level.




Fig. 2. Packing diagram of the title compound. H atoms bonded to C are not shown. The N—H...O hydrogen bonds are shown as dashed lines.

***N'*-'[(1*E*)-(4-Fluorophenyl)methylene]-6-methoxy-2-naphthohyrazide**

Crystal data

$C_{19}H_{15}FN_2O_2$

$M_r = 322.33$

Monoclinic, *Pc*

Hall symbol: *P* -2yc

$a = 15.681$ (2) Å

$b = 5.6574$ (7) Å

$c = 9.2758$ (10) Å

$\beta = 104.746$ (9)°

$V = 795.79$ (17) Å³

$Z = 2$

$F_{000} = 336$

$D_x = 1.345$ Mg m⁻³

Mo *K*α radiation

$\lambda = 0.71073$ Å

Cell parameters from 6963 reflections

$\theta = 3.7$ – 25.3 °

$\mu = 0.10$ mm⁻¹

$T = 173$ (2) K

Needle, colourless

$0.24 \times 0.09 \times 0.08$ mm

Data collection

Stoe IPDSII two-circle diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 173$ (2) K

ω scans

Absorption correction: none

9090 measured reflections

1531 independent reflections

1246 reflections with $I > 2\sigma(I)$

$R_{int} = 0.069$

$\theta_{max} = 25.9$ °

$\theta_{min} = 3.6$ °

$h = -19 \rightarrow 19$

$k = -6 \rightarrow 6$

$l = -11 \rightarrow 11$

Refinement

Refinement on F^2

Least-squares matrix: full

Hydrogen site location: difference Fourier map

H-atom parameters constrained

$R[F^2 > 2\sigma(F^2)] = 0.063$	$w = 1/[\sigma^2(F_o^2) + (0.0088P)^2]$
$wR(F^2) = 0.127$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.57$	$(\Delta/\sigma)_{\max} < 0.001$
1531 reflections	$\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$
217 parameters	$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$
2 restraints	Extinction correction: none
60 constraints	
Primary atom site location: structure-invariant direct methods	
Secondary atom site location: difference Fourier map	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.3049 (2)	-0.5672 (6)	0.2289 (4)	0.0710 (10)
N1	0.6850 (2)	0.3354 (7)	0.3540 (4)	0.0459 (10)
H1	0.7100	0.3124	0.2803	0.055*
N2	0.6140 (2)	0.1982 (8)	0.3648 (4)	0.0454 (10)
O1	0.6834 (2)	0.5483 (7)	0.5623 (3)	0.0560 (10)
O2	1.1811 (2)	0.9463 (7)	0.3792 (4)	0.0596 (10)
C1	0.7163 (3)	0.5057 (9)	0.4572 (4)	0.0451 (12)
C2	0.5913 (3)	0.0397 (9)	0.2636 (5)	0.0443 (11)
H2	0.6241	0.0238	0.1909	0.053*
C3	1.2147 (4)	1.1340 (11)	0.4833 (7)	0.0630 (15)
H3A	1.1768	1.2735	0.4573	0.095*
H3B	1.2748	1.1740	0.4791	0.095*
H3C	1.2152	1.0822	0.5844	0.095*
C11	0.7977 (3)	0.6295 (9)	0.4426 (4)	0.0417 (11)
C12	0.8549 (3)	0.5335 (9)	0.3667 (5)	0.0426 (11)
H12	0.8393	0.3898	0.3136	0.051*
C13	0.9355 (3)	0.6423 (9)	0.3658 (5)	0.0402 (11)
C14	0.9964 (3)	0.5461 (9)	0.2891 (5)	0.0446 (11)
H14	0.9813	0.4052	0.2327	0.054*
C15	1.0744 (3)	0.6503 (9)	0.2950 (5)	0.0478 (12)
H15	1.1129	0.5842	0.2416	0.057*

supplementary materials

C16	1.0994 (3)	0.8597 (9)	0.3811 (5)	0.0451 (12)
C17	1.0422 (3)	0.9647 (9)	0.4528 (5)	0.0467 (12)
H17	1.0585	1.1073	0.5068	0.056*
C18	0.9591 (3)	0.8602 (8)	0.4460 (5)	0.0408 (11)
C19	0.8981 (3)	0.9577 (9)	0.5207 (5)	0.0442 (11)
H19	0.9120	1.1021	0.5737	0.053*
C20	0.8208 (3)	0.8489 (9)	0.5175 (5)	0.0454 (12)
H20	0.7811	0.9205	0.5665	0.055*
C21	0.5165 (3)	-0.1173 (8)	0.2571 (4)	0.0415 (11)
C22	0.4518 (3)	-0.0650 (10)	0.3318 (5)	0.0470 (12)
H22	0.4567	0.0750	0.3899	0.056*
C23	0.3808 (3)	-0.2141 (10)	0.3223 (5)	0.0528 (13)
H23	0.3366	-0.1758	0.3719	0.063*
C24	0.3750 (3)	-0.4206 (10)	0.2395 (6)	0.0517 (13)
C25	0.4372 (3)	-0.4777 (10)	0.1636 (5)	0.0508 (12)
H25	0.4319	-0.6195	0.1071	0.061*
C26	0.5081 (3)	-0.3244 (9)	0.1707 (5)	0.0484 (12)
H26	0.5506	-0.3602	0.1170	0.058*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.065 (2)	0.074 (2)	0.079 (2)	-0.0181 (16)	0.0265 (17)	-0.0044 (18)
N1	0.043 (2)	0.071 (3)	0.030 (2)	-0.009 (2)	0.0204 (17)	-0.0011 (18)
N2	0.045 (2)	0.063 (3)	0.0297 (18)	0.0003 (18)	0.0123 (16)	0.0062 (18)
O1	0.060 (2)	0.083 (3)	0.0327 (17)	-0.0114 (18)	0.0258 (15)	-0.0084 (16)
O2	0.048 (2)	0.082 (3)	0.055 (2)	-0.0157 (19)	0.0231 (17)	-0.0125 (19)
C1	0.053 (3)	0.057 (3)	0.028 (2)	0.003 (2)	0.015 (2)	0.004 (2)
C2	0.036 (3)	0.069 (3)	0.031 (2)	0.002 (2)	0.0141 (19)	0.003 (2)
C3	0.058 (3)	0.074 (4)	0.059 (3)	-0.017 (3)	0.018 (3)	-0.013 (3)
C11	0.041 (3)	0.057 (3)	0.030 (2)	0.002 (2)	0.0143 (19)	0.004 (2)
C12	0.045 (3)	0.060 (3)	0.025 (2)	0.000 (2)	0.0127 (19)	0.004 (2)
C13	0.039 (3)	0.055 (3)	0.028 (2)	0.001 (2)	0.0111 (18)	0.002 (2)
C14	0.042 (3)	0.061 (3)	0.035 (2)	0.000 (2)	0.017 (2)	-0.004 (2)
C15	0.044 (3)	0.066 (3)	0.037 (3)	0.005 (3)	0.017 (2)	0.000 (2)
C16	0.043 (3)	0.061 (3)	0.033 (2)	-0.002 (2)	0.012 (2)	0.004 (2)
C17	0.056 (3)	0.050 (3)	0.036 (2)	-0.006 (2)	0.014 (2)	-0.005 (2)
C18	0.042 (3)	0.051 (3)	0.033 (2)	-0.002 (2)	0.0151 (19)	0.005 (2)
C19	0.051 (3)	0.052 (3)	0.033 (2)	0.000 (2)	0.018 (2)	-0.001 (2)
C20	0.049 (3)	0.060 (3)	0.030 (2)	0.007 (2)	0.015 (2)	0.007 (2)
C21	0.035 (2)	0.064 (3)	0.026 (2)	0.008 (2)	0.0078 (18)	0.006 (2)
C22	0.044 (3)	0.066 (3)	0.033 (2)	0.002 (2)	0.015 (2)	-0.001 (2)
C23	0.039 (3)	0.073 (4)	0.051 (3)	0.007 (2)	0.022 (2)	0.011 (3)
C24	0.038 (3)	0.068 (4)	0.049 (3)	-0.003 (3)	0.011 (2)	0.009 (3)
C25	0.051 (3)	0.059 (3)	0.042 (3)	0.002 (2)	0.011 (2)	0.000 (2)
C26	0.044 (3)	0.064 (3)	0.038 (2)	0.009 (2)	0.014 (2)	0.006 (2)

Geometric parameters (Å, °)

F1—C24	1.360 (6)	C14—H14	0.9500
N1—C1	1.359 (6)	C15—C16	1.427 (7)
N1—N2	1.381 (5)	C15—H15	0.9500
N1—H1	0.8800	C16—C17	1.379 (7)
N2—C2	1.281 (6)	C17—C18	1.417 (7)
O1—C1	1.238 (5)	C17—H17	0.9500
O2—C16	1.377 (6)	C18—C19	1.428 (6)
O2—C3	1.442 (7)	C19—C20	1.353 (7)
C1—C11	1.492 (7)	C19—H19	0.9500
C2—C21	1.460 (7)	C20—H20	0.9500
C2—H2	0.9500	C21—C22	1.399 (6)
C3—H3A	0.9800	C21—C26	1.407 (7)
C3—H3B	0.9800	C22—C23	1.381 (8)
C3—H3C	0.9800	C22—H22	0.9500
C11—C12	1.384 (6)	C23—C24	1.389 (7)
C11—C20	1.424 (7)	C23—H23	0.9500
C12—C13	1.408 (7)	C24—C25	1.378 (7)
C12—H12	0.9500	C25—C26	1.399 (8)
C13—C14	1.436 (6)	C25—H25	0.9500
C13—C18	1.439 (6)	C26—H26	0.9500
C14—C15	1.346 (7)		
C1—N1—N2	120.1 (3)	O2—C16—C15	114.3 (4)
C1—N1—H1	120.0	C17—C16—C15	120.5 (4)
N2—N1—H1	120.0	C16—C17—C18	120.1 (5)
C2—N2—N1	114.8 (4)	C16—C17—H17	119.9
C16—O2—C3	115.8 (4)	C18—C17—H17	119.9
O1—C1—N1	123.0 (4)	C17—C18—C19	122.6 (5)
O1—C1—C11	121.2 (4)	C17—C18—C13	119.6 (4)
N1—C1—C11	115.7 (4)	C19—C18—C13	117.7 (4)
N2—C2—C21	121.7 (4)	C20—C19—C18	121.4 (5)
N2—C2—H2	119.1	C20—C19—H19	119.3
C21—C2—H2	119.1	C18—C19—H19	119.3
O2—C3—H3A	109.5	C19—C20—C11	121.5 (4)
O2—C3—H3B	109.5	C19—C20—H20	119.3
H3A—C3—H3B	109.5	C11—C20—H20	119.3
O2—C3—H3C	109.5	C22—C21—C26	118.9 (4)
H3A—C3—H3C	109.5	C22—C21—C2	121.9 (4)
H3B—C3—H3C	109.5	C26—C21—C2	119.2 (4)
C12—C11—C20	118.2 (4)	C23—C22—C21	120.9 (5)
C12—C11—C1	122.8 (5)	C23—C22—H22	119.5
C20—C11—C1	118.8 (4)	C21—C22—H22	119.5
C11—C12—C13	122.0 (5)	C22—C23—C24	119.2 (4)
C11—C12—H12	119.0	C22—C23—H23	120.4
C13—C12—H12	119.0	C24—C23—H23	120.4
C12—C13—C14	123.3 (5)	F1—C24—C25	118.9 (5)
C12—C13—C18	119.0 (4)	F1—C24—C23	119.4 (4)

supplementary materials

C14—C13—C18	117.7 (4)	C25—C24—C23	121.6 (5)
C15—C14—C13	121.8 (5)	C24—C25—C26	119.2 (5)
C15—C14—H14	119.1	C24—C25—H25	120.4
C13—C14—H14	119.1	C26—C25—H25	120.4
C14—C15—C16	120.2 (4)	C25—C26—C21	120.1 (4)
C14—C15—H15	119.9	C25—C26—H26	120.0
C16—C15—H15	119.9	C21—C26—H26	120.0
O2—C16—C17	125.1 (5)		
C1—N1—N2—C2	-178.4 (4)	C16—C17—C18—C13	0.9 (7)
N2—N1—C1—O1	-0.9 (7)	C12—C13—C18—C17	176.5 (5)
N2—N1—C1—C11	175.0 (4)	C14—C13—C18—C17	-3.1 (6)
N1—N2—C2—C21	-178.8 (4)	C12—C13—C18—C19	-1.2 (5)
O1—C1—C11—C12	154.8 (5)	C14—C13—C18—C19	179.2 (5)
N1—C1—C11—C12	-21.2 (7)	C17—C18—C19—C20	-177.1 (4)
O1—C1—C11—C20	-21.0 (7)	C13—C18—C19—C20	0.5 (6)
N1—C1—C11—C20	162.9 (4)	C18—C19—C20—C11	1.5 (7)
C20—C11—C12—C13	2.0 (6)	C12—C11—C20—C19	-2.7 (6)
C1—C11—C12—C13	-173.9 (4)	C1—C11—C20—C19	173.4 (4)
C11—C12—C13—C14	179.5 (4)	N2—C2—C21—C22	17.9 (7)
C11—C12—C13—C18	-0.1 (6)	N2—C2—C21—C26	-163.6 (4)
C12—C13—C14—C15	-177.5 (4)	C26—C21—C22—C23	0.5 (6)
C18—C13—C14—C15	2.1 (7)	C2—C21—C22—C23	179.0 (4)
C13—C14—C15—C16	1.2 (7)	C21—C22—C23—C24	1.2 (7)
C3—O2—C16—C17	12.4 (8)	C22—C23—C24—F1	-179.5 (4)
C3—O2—C16—C15	-170.3 (4)	C22—C23—C24—C25	-1.6 (7)
C14—C15—C16—O2	179.2 (4)	F1—C24—C25—C26	178.2 (4)
C14—C15—C16—C17	-3.5 (7)	C23—C24—C25—C26	0.3 (7)
O2—C16—C17—C18	179.4 (4)	C24—C25—C26—C21	1.4 (7)
C15—C16—C17—C18	2.4 (7)	C22—C21—C26—C25	-1.8 (6)
C16—C17—C18—C19	178.5 (4)	C2—C21—C26—C25	179.7 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots O1 ⁱ	0.88	2.11	2.778 (4)	132
C23—H23 \cdots O2 ⁱⁱ	0.95	2.55	3.428 (6)	154
C26—H26 \cdots O1 ⁱⁱⁱ	0.95	2.50	3.403 (6)	158
C14—H14 \cdots Cg1 ⁱ	0.98	2.89	3.666 (5)	140
C19—H19 \cdots Cg2 ^{iv}	0.93	2.95	3.697 (5)	136

Symmetry codes: (i) $x, -y+1, z-1/2$; (ii) $x-1, y-1, z$; (iii) $x, -y, z-1/2$; (iv) $x, -y+2, z+1/2$.

Fig. 1




Fig. 2

