

N-H Bond Activation by Palladium(II) and Copper(I) Complexes Featuring a Reactive Bidentate PN-Ligand

Sandra Y. de Boer,^a Yann Gloaguen,^a Joost N. H. Reek,^a Martin Lutz^b
and Jarl Ivar van der Vlugt^{a,*}

^a *van 't Hoff Institute for Molecular Sciences, University of Amsterdam
Science Park 904, 1098 XH Amsterdam, The Netherlands.*

E-mail: J.I.vanderVlugt@uva.nl

Fax: +31 20 5256422; Tel: +31 20 5256437

^b *Crystal and Structural Chemistry, Faculty of Science, Utrecht University,
Padualaan 8, 3584 CH Utrecht, The Netherlands.*

NMR and MS Spectra of New Ligands and Complexes




Fig. S1. ¹H NMR spectrum of ligand L1H (C₆D₆, 400 MHz, 298 K).




Fig. S2. ¹³C NMR spectrum of ligand L1H (C₆D₆, 100 MHz, 298 K).




Fig. S3. ^{31}P NMR spectrum of ligand **L1H** (C_6D_6 , 162 MHz, 298 K).




Fig. S4. ^1H NMR spectrum of complex **1** (CDCl_3 , 500 MHz, 298 K).




Fig. S5. ^{13}C NMR spectrum of complex **1** (CDCl_3 , 126 MHz, 298 K).




Fig. S6. ^{31}P NMR spectrum of complex **1** (CDCl_3 , 202 MHz, 298 K).




Fig. S7. ¹H NMR spectrum of complex X (acetone-*d*₆, 400 MHz, 298 K).




Fig. S8. ³¹P NMR spectrum of complex X (acetone-*d*₆, 162 MHz, 298 K).




Fig. S9. ¹H NMR spectrum of complex Y (C₆D₆, 400 MHz, 298 K). X at δ 1.06 is *tert*-butyl alcohol, at δ 0.30 grease.




Fig. S10. ¹³C NMR spectrum of complex Y (C₆D₆, 100 MHz, 298 K). X at δ 60 is an artefact.




Fig. S11. ^{31}P NMR spectrum of complex Y (C_6D_6 , 162 MHz, 298 K).




Fig. S12. Mass spectrum of complex Y.




Fig. S13. ¹H NMR spectrum of complex 2 (acetone-*d*₆, 400 MHz, 298 K).




Fig. S14. ¹³C NMR spectrum of complex 2 (acetone-*d*₆, 100 MHz, 298 K).




Fig. S15. ^{31}P NMR spectrum of complex 2 (acetone- d_6 , 162 MHz, 298 K).




Fig. S16. Mass spectrum of complex 2.




Fig. S17. ¹H NMR spectrum of complex 4 (acetone-*d*₆, 400 MHz, 298 K). X at δ 3.63 and 1.81 is THF.




Fig. S18. ¹³C (APT) NMR spectrum of complex 4 (acetone-*d*₆, 100 MHz, 298 K).




Fig. S19. ^{31}P NMR spectrum of complex 4 (acetone- d_6 , 162 MHz, 298 K).




Fig. S20. Mass spectrum of complex 4.

