

N-Tosyl-(*S_a*)-binam-L-prolinamide as Highly Efficient Bifunctional Organocatalyst for the General Enantioselective Solvent-Free Aldol Reaction

Gabriela Guillena,* Carmen Nájera,* Santiago F. Vióquez

Dpto. Química Orgánica and Instituto de Síntesis Orgánica, Universidad de Alicante, Apdo. 99, 03080 Alicante, Spain
Fax +34(965)903549; E-mail: gabriela.guillena@ua.es; E-mail: cnajera@ua.es

Received 30 July 2008

Abstract: *N*-Tosyl-(*S_a*)-binam-L-prolinamide (5 mol%) and benzoic acid (1 mol%) were used as catalysts in the enantioselective direct aldol reaction between different ketones and aldehydes under solvent-free conditions in the presence or absence of water. Under these reaction conditions it was possible to reduce the amount of required ketone to two equivalents to give the corresponding aldol products with high yields, regio-, diastereo- and enantioselectivities. The aldol reaction between aldehydes or the intramolecular aldol reaction can be also performed with excellent results.

Key words: organocatalyzed, aldol reactions, solvent-free, enantioselective, ketone

Bisprolinamides derived from 1,1'-binaphthyl-2,2'-diamine (binam) have been studied as organocatalysts¹ in the aldol reaction² under different reaction conditions using alkyl³ and α -functionalized⁴ ketones as source of nucleophile, with recoverable catalyst (*S_a*)-binam-L-Pro **1** (Figure 1),^{3a,c} providing the best results. For this prolinamide, the addition of an equimolecular amount of benzoic acid as co-catalyst in the reaction has been crucial to get a great acceleration in the reaction. Under these reaction conditions, less reactive ketones, such as butanone^{3c} and α -functionalized ketones can be used as donors, even in water.⁴ Recently, we have described the aldol reaction under solvent-free conditions, reducing the loading of catalyst **1** and the amount of required ketone to two equivalents to give the corresponding products with a high level of selectivity.⁵ Mechanistic studies showed that the derived enamines from one or both proline moieties are formed but only one prolinamide works as catalyst. In addition, the presence of benzoic acid as co-catalyst increased the reaction rate but decreased the enantioselectivity.

With these previous results in our hands, we thought that it would be of interest to achieve the replacement of one of the proline residue in catalyst **1**, by an acidic moiety such as a sulfonamide group, which might allow the reduction of the amount of acid co-catalyst. In addition, the sulfonamide moiety can activate the carbonyl group of the acceptor aldehydes through a hydrogen bond acting as a bifunctional catalyst. Here, we report the application of the unsymmetrical (*S_a*)-binam-L-prolinamide sulfonamide





Figure 1 Binam-L-prolinamide-derived catalysts

derivative **2**⁶ (Figure 1) in the aldol reaction under solvent-free reaction conditions.⁷

Catalyst **2** was prepared by a modified procedure of the standard method⁶ using *p*-toluenesulfonyl chloride and commercially available (*S*)-1,1'-binaphthyl-2,2'-diamine [(*S_a*)-binam], and subsequently coupling with the in situ generated Fmoc-L-Pro chloride followed by deprotection with piperidine, giving compound **2** in 80% overall yield after purification.

Then, catalyst **2** was tested in the reaction between cyclohexanone and 4-nitrobenzaldehyde under solvent-free conditions, to optimize the reaction conditions. The addition of a small amount of benzoic acid (1 mol%) was required to obtain the aldol product **4aa** in shorter reaction times (Table 1, compare entries 1 and 2). Under the same reaction conditions, catalysts **1** showed to be less efficient, giving after three hours, 90% of conversion, with the aldol products being obtained with a slightly lower diastereo- and enantioselectivity (Table 1, entry 3). In some organocatalyzed aldol reactions, the addition of water to an organic solvent has shown to have a positive effect on the reaction rates.⁸ Thus, the effect of the use of some amount of water in the reaction catalyzed by **2** was studied (Table 1). First, the addition of 1 mL of water was evaluated. Under these conditions, the reaction was much slower, giving the aldol product **4aa** with lower conversion, diastereo- and enantioselectivity (Table 1, compare entries 2 and 4). Decreasing gradually the amount of added water, caused an acceleration in the reaction rate as well as an increase in the conversion, diastereo- and enantioselectivity ~~was achieved~~ (Table 1, entries 5–7), with the best results being obtained by adding only 33 μ L (7 equiv) of water to the reaction mixture (Table 1, entry 7). This effect was in contrast to that observed when proline was used as catalyst for the aldol reaction under solvent-free conditions, in which the addition of small amounts of wa-

SYNLETT 2008, No. x, pp 0001–0005
Advanced online publication: xx.xx.2008
DOI: 10.1055/s-0028-xxxxx; Art ID: G24208ST
© Georg Thieme Verlag Stuttgart · New York

Imprimatur: 
Date, Signature 08/10/2008

g24208st.fm, 10/7/08

Table 1 Effect of the Addition of Water in the Reaction between 4-Nitrobenzaldehyde and Cyclohexanone under Solvent-Free Conditions^a

Entry	H ₂ O (μL)	T (°C)	Time (h)	Conversion (%) ^c	<i>anti/syn</i> ^d	ee (%) ^e
1 ^b	–	0	12	97	97:3	99
2	–	0	6	99	99:1	97
3 ^f	–	0	3	90	96:4	94
4	1000	0	26	44	86:16	94
5	500	0	23	39	93:7	96
6	125	0	18	70	86:11	98
7	33	0	4	99	99:1	98
8 ^f	33	0	3	97	97:3	92

^a Reaction conditions: 4-nitrobenzaldehyde (0.25 mmol), cyclohexanone (2 equiv), **2** (5 mol%), BzOH (1 mol%), 0 °C, unless stated otherwise.

^b The reaction was carried out in the absence of BzOH.

^c Conversion based on the unreacted aldehyde.

^d Determined by ¹H NMR of the crude product.

^e Determined by chiral-phase HPLC analysis for the *anti* isomer.

^f Reaction conditions: catalyst **1** (5 mol%), BzOH (1 mol%).

ter caused a decrease in the reaction rate.^{7c} Applying these reaction conditions using catalyst **1** (5 mol%), benzoic acid (1 mol%), the achieved result in terms of conversion, diastereo- and enantioselectivity were poorer than those obtained using catalyst **2** (Table 1, compare entries 7 and 8) where longer reaction time was required.

The scope of catalyst **2** under anhydrous and wet solvent-free reaction conditions was studied using several aromatic aldehydes and cyclic, acyclic and α -functionalized ketones (Scheme 1 and Table 2). For cyclic ketones the reaction proceeded smoothly in all cases giving mainly the *anti* isomers **4** (Table 2, entries 1–8), except for cyclopentanone, with isomer *syn*-**4ba** being in this case the major isomer (Table 2, entries 9 and 10). The reaction times were shorter under wet than under anhydrous conditions. Whereas 4-nitrobenzaldehyde gave the corresponding aldol product in a short reaction time, nonactivated aldehydes such as benzaldehyde or tolualdehyde needed longer reaction time to give the aldol product in moderate yields (36–76%) but in excellent diastereo- and enantioselectivities (96% de and 96% ee; Table 2, entries 3–6). Also heteroaromatic aldehydes such as furfural were suitable donor for this process yielding mainly *anti*-**4ad** in better yield (52%) under wet than under anhydrous conditions and in high diastereo- and enantioselectivity (Table 2, entries 7 and 8). Cyclopentanone gave mainly aldol *syn*-**4ba** with slightly better diastereo- and enantioselectivity under anhydrous than under wet conditions (Table 2, entries 9 and 10). Surprisingly, 4-pyranone,⁹

gave the expected *anti* isomer (major isomer) in only 50% yield under wet conditions compared to 93% yield obtained under anhydrous conditions, with comparable diastereo- and enantioselectivities achieved for both cases (Table 2, entries 11 and 12). When acetone was allowed to react with 4-nitrobenzaldehyde the aldol **4da** was obtained in good yield and enantioselectivity in only 16 hours under wet conditions, whereas 48 hours were required under anhydrous conditions (Table 2, entries 13 and 14). Longer reaction times were required with less reactive 2-butanone giving mainly the *iso* isomer **5ea** with excellent enantioselectivity under wet conditions (Table 2, entries 15 and 16).

In all cases, with the exception of 2-butanone, the achieved diastereo- and enantioselectivities were better than those previously obtained using catalyst **1** (5 mol%) and benzoic acid (10 mol%) under solvent-free conditions, with longer reaction times being required with catalyst **2**.⁵ Finally, several α -functionalized ketones were reacted with 4-nitrobenzaldehyde under wet and anhydrous solvent-free conditions. Whereas α -methoxyacetone gave mainly the *anti* isomer **4fa** in shorter reaction time under anhydrous conditions (Table 2, entries 17 and 18), α -methylsulfanylacetone gave as main product *iso*-**5ga** with good enantioselectivity, with catalyst **2** (10 mol%) and benzoic acid (5 mol%) being needed to obtain 87% yield under anhydrous conditions (Table 2, entries 19 and 20).













**Scheme 1** Reaction between 4-nitrobenzaldehyde and different ketones under solvent-free conditions

Table 2 Reaction of Aldehydes with Cyclohexanone under Anhydrous and Wet Solvent-Free Conditions^a

Entry	Major product	Time (h)	Yield (%) ^b	4/5	<i>anti/syn</i> ^c	ee (%) ^d
1		6	94	--	99:1	98
2 ^e		4	98	--	99:1	98
<i>anti-4aa</i>						
3		120	76	--	99:1	98
4 ^e		72	74	--	99:1	98
<i>anti-4ab</i>						
5		120	36	--	98:2	96
6 ^e		91	52	--	>99:1	98
<i>anti-4ac</i>						
7		26	27	--	96:4	92
8 ^e		91	52	--	97:3	94
<i>anti-4ad</i>						
9		15	90	--	32:68	84
10 ^e		7.5	88	--	37:63	78
<i>syn-4ba</i>						
11		24	93	--	97:3	89
12 ^e		19	50	--	98:2	94
<i>anti-4fa</i>						
13		48	86	--	--	84
14 ^e		16	88	--	--	86
<i>anti-4da</i>						
15		96	62	23:77	>99:1	90
16 ^e		68	81	36:64	>99:1	98
<i>iso-5ea</i>						
17		48	73	78:22	82:18	93
18 ^e		65	76	72:28	84:16	94
<i>anti-4fa</i>						
19 ^f		120	87	17:83	>99:1	88
20 ^{e,f}		67	40	19:81	>99:1	86
<i>iso-5ga</i>						


^a Reaction conditions: 4-nitrobenzaldehyde (0.25 mmol), ketone (2 equiv), **2** (5 mol%), BzOH (1 mol%), 0 °C, unless stated otherwise.^b Yield after purification by column chromatography.^c Determined by ¹H NMR of the crude product.^d Determined by chiral-phase HPLC analysis for the major isomer.^e H₂O (33 μL) was added to the reaction mixture.^f Catalyst **2** (10 mol%) and BzOH (5 mol%) were used.



Scheme 2 Reaction between 4-nitrobenzaldehyde and propanal under solvent-free conditions

Aldehydes can also act as donors in this transformation. In this case catalyst **2** (10 mol%) and benzoic acid (5 mol%) have to be used at 25 °C to perform the reaction. When propanal (5 equiv) and 4-nitrobenzaldehyde were allowed to react under these conditions 1,3-diol **7** was obtained under anhydrous conditions in seven days with 80% yield, 66% de and 94% ee, after in situ reduction of the corresponding aldol with NaBH₄ in MeOH (Scheme 2). When the same reaction was carried out by adding 33 μL of water (wet conditions), compound **7** was obtained after 2.5 days with 86% yield, 78 de and 92% ee (Scheme 2). These results are superior in terms of reaction time, yield, diastereo- and enantioselectivity to those reported previously using catalyst **1** (5 mol%) and benzoic acid (10 mol%).^{5b} In this case six days of reaction time was required to yield 45% of compound **7** with 56% de and 85% ee after final reduction.

-/ anhydrous Finally, these ~~2.5 days of reaction time~~ and wet solvent free-conditions were applied to the intramolecular aldol reaction (Robinson-type annulation),¹⁰ which has been extensively used in the synthesis of natural products (Scheme 3 and Table 3).^{10,2} Using catalyst **2** (5 mol%) and benzoic acid (1 mol%) at room temperature under anhydrous solvent-free conditions the 6-(enolendo)-*exo, trig* cyclization of triketone **8a** (n = 2), led directly to the corresponding Wieland–Miescher ketone¹¹ **9a** in only 27 hours with excellent yield and enantioselectivity (Table 3, entry 1).



Scheme 3 Intramolecular aldol reaction under solvent-free conditions

A shorter reaction time was required when a small amount of water (14 μL, 3 equiv) was added to the reaction mixture (wet conditions), giving ketone **9a** with similar yield and ee (Table 3, entry 2). These results are better when compared to those reported in the literature using MeCN as solvent in the presence of HClO₄ and 50 mol% of L-proline at 80 °C,^{10c} affording compound **9a** in 73% yield

Table 3 Intramolecular Aldol Reaction under Anhydrous and Wet Solvent-Free Conditions^a

Entry	Product	n	Time (h)	Yield (%) ^b	9/10 ^c	ee (%) ^d
1	9a	2	27	95	100:0	90
2 ^e	9a	2	16	90	100:0	88
3	9b	1	48	84	67:36	83
4 ^e	9b	1	48	85	70:30	90

^a Reaction conditions: triketone **8** (0.25 mmol), **2** (5 mol%), BzOH (1 mol%), 25 °C.

^b Yield after purification by column chromatography for compound **9**.

^c Determined by ¹H NMR of the crude product.

^d Determined by chiral-phase HPLC analysis of the isolated product.

^e H₂O (14 μL, 3 equiv) was added to the reaction mixture.

and 71% ee. The synthesis of bicyclic compound **9b**, which is an important intermediate in the total synthesis of steroids was optimized using L-proline (3 mol%) in DMF as solvent at room temperature.^{10a} Under these conditions product **10a** (n = 1) was obtained with 100% yield and 93% ee, with a subsequent dehydration step being necessary to achieve the desired product. Under solvent-free conditions and using catalyst **2**, the main product observed was the indanone derivative **9b**. After column chromatography purification, the only product isolated with similar yield, under both anhydrous and wet conditions, was compound **9b**, with better enantioselectivity obtained under wet conditions (Table 3, entries 3 and 4).


In conclusion, sulfonamide **2** in the presence of a small amount of benzoic acid has shown to be a useful catalyst to perform the intramolecular and the ketone–aldehyde or aldehyde–aldehyde intermolecular aldol reaction under solvent-free conditions.¹² The aldol products were obtained with good yields and excellent diastereo- and enantioselectivities, generally, with the addition of a small amount of water (3–7 equiv) to the reaction mixture leading to an increase in the reaction rates. In general, these results are superior in terms of diastereo- and enantioselectivities to those previously reported using L-proline or prolinamide **1** as catalyst under solvent-free conditions. Further scope and application of sulfonamide catalyst **2** in this and other related transformations are currently being studied.

Acknowledgment

This work was financially supported by the Dirección General de Investigación of the Ministerio de Educación y Ciencia of Spain (Proyectos CTQ2007-62771/BQU and Consolider Ingenio 2010 CSD2007-00006), and the University of Alicante (GRJ06-05).

References and Notes

- For comprehensive books and selected reviews, see: (a) Dalko, P. I.; Moisan, L. *Angew. Chem. Int. Ed.* **2004**, *43*, 5138. (b) Berkessel, A.; Gröger, H. *Asymmetric Organocatalysis: From Biomimetic Concepts to Applications in Asymmetric Synthesis*; Wiley-VCH: Weinheim, **2005**. (c) Seayad, J.; List, B. *Org. Biomol. Chem.* **2005**, *3*, 719. (d) Kočovský, P.; Malkov, A. V. *Tetrahedron* **2006**, *62*, 255. (e) Lelais, G.; McMillan, D. W. C. *Aldrichimica Acta* **2006**, *39*, 79. (f) Marigo, M.; Jørgensen, K. A. *Chem. Commun.* **2006**, 2001. (g) Guillena, G.; Ramón, D. J. *Tetrahedron: Asymmetry* **2006**, *17*, 1465. (h) List, B. *Chem. Commun.* **2006**, 819. (i) Gaunt, M. J.; Johansson, C. C. C.; McNally, A.; Vo, N. T. *Drug Discovery Today* **2007**, *12*, 8. (j) Pellissier, H. *Tetrahedron* **2007**, *63*, 9267. (k) *Enantioselective Organocatalysis*; Dalko, P. I., Ed.; Wiley-VCH: Weinheim, **2007**. (l) List, B. *Chem. Rev.* **2007**, *107*, 5413. (m) Enders, D.; Grondal, C.; Hüttl, M. R. M. *Angew. Chem. Int. Ed.* **2007**, *46*, 1570. (n) Guillena, G.; Ramón, D. J.; Yus, M. *Tetrahedron: Asymmetry* **2007**, *18*, 693. (o) de Figueiredo, R. M.; Christmann, M. *Eur. J. Org. Chem.* **2007**, 2575. (p) Kotsuki, H.; Ikishima, H.; Okuyama, A. *Heterocycles* **2008**, *75*, 493. (q) Kotsuki, H.; Ikishima, H.; Okuyama, A. *Heterocycles* **2008**, *75*, 757.
- Guillena, G.; Nájera, C.; Ramón, D. J. *Tetrahedron: Asymmetry* **2007**, *18*, 2249.
- (a) Guillena, G.; Hita, M. C.; Nájera, C. *Tetrahedron: Asymmetry* **2006**, *17*, 729. (b) Gryko, D.; Kowalczyk, B.; Zawadzki, L. *Synlett* **2006**, 1059. (c) Guillena, G.; Hita, M. C.; Nájera, C. *Tetrahedron: Asymmetry* **2006**, *17*, 1493. (d) *Corrigendum*: Guillena, G.; Hita, M. C.; Nájera, C. *Tetrahedron: Asymmetry* **2007**, *18*, 1031. (e) Guizzetti, S.; Benaglia, M.; Pignataro, L.; Puglisi, A. *Tetrahedron: Asymmetry* **2006**, *17*, 2754. (f) Ma, G.-N.; Zhang, Y.-P.; Shi, M. *Synthesis* **2007**, 197. (g) Guizzetti, S.; Benaglia, M.; Raimondi, L.; Celentano, G. *Org. Lett.* **2007**, *9*, 1247.
- (a) Guillena, G.; Hita, M. C.; Nájera, C. *Tetrahedron: Asymmetry* **2006**, *17*, 1027. (b) *Corrigendum*: Guillena, G.; Hita, M. C.; Nájera, C. *Tetrahedron: Asymmetry* **2007**, *18*, 1030. (c) Guillena, G.; Hita, M. C.; Nájera, C. *Arkivoc* **2007**, (iv), 260. (d) *Corrigendum*: Guillena, G.; Hita, M. C.; Nájera, C. *Arkivoc* **2007**, (i), 146. (e) Guillena, G.; Hita, M. C.; Nájera, C. *Tetrahedron: Asymmetry* **2007**, *18*, 1272.
- (a) Guillena, G.; Hita, M. C.; Nájera, C.; Vióquez, S. F. *Tetrahedron: Asymmetry* **2007**, *18*, 2300. (b) Guillena, G.; Hita, M. C.; Nájera, C.; Vióquez, S. F. *J. Org. Chem.* **2008**, *73*, 5933.
- Chen, T.; Gao, J.; Shi, M. *Tetrahedron* **2006**, *62*, 6289.
- For some other organocatalyzed enantioselective aldol reactions under solvent-free conditions, see: (a) Rodríguez, B.; Rantanen, T.; Bolm, C. *Angew. Chem. Int. Ed.* **2006**, *45*, 6924. (b) Hayashi, Y.; Aratake, S.; Itoh, T.; Okano, T.; Sumiya, T.; Shoji, M. *Chem. Commun.* **2007**, 957. (c) Rodríguez, B.; Bruckmann, A.; Bolm, C. *Chem. Eur. J.* **2007**, *13*, 4710.
- Nyberg, A. I.; Usano, A.; Pinko, P. M. *Synlett* **2004**, 1891.
- Chen, J.-R.; Li, X.-Y.; Xing, X.-N.; Xiao, W.-J. *J. Org. Chem.* **2006**, *71*, 8198.
- (a) Eder, U.; Wiechert, R.; Sauer, G. Ger. Patent, DE 2014757, **1971**; *Chem. Abstr.* **1972**, *76*, 14180. (b) Hajos, Z. G.; Parrish, D. R. Ger. Patent, DE 2102623, **1971**; *Chem. Abstr.* **1972**, *76*, 59072. (c) Eder, U.; Sauer, G.; Wiechert, R. *Angew. Chem., Int. Ed. Engl.* **1971**, *10*, 496. (d) Hajos, Z. G.; Parrish, D. R. *J. Org. Chem.* **1974**, *39*, 1615. (e) Hajos, Z. G.; Parrish, D. R. In *Org. Synth., Coll. Vol. VII*; Freeman, J. P., Ed.; Wiley: New York, **1990**, 363. (f) Buchschacher, P.; Fürst, A.; Gutzwiller, J. In *Org. Synth., Coll. Vol. VII*; Freeman, J. P., Ed.; Wiley: New York, **1990**, 368. (g) Kwiatkowski, S.; Syed, A.; Brock, C. P.; Watt, D. S. *Synthesis* **1989**, 818. (h) Tietze, L. F.; Utecht, J. *Synthesis* **1993**, 927.
- Wieland, P.; Miescher, K. *Helv. Chim. Acta* **1950**, *33*, 2215.
- Representative Experimental Procedure:**
Anhydrous Solvent-Free Conditions: To a mixture of the corresponding aldehyde (0.25 mmol), catalyst **2** (0.0125 mmol, 6.5 mg) and benzoic acid (0.0025 mmol, 0.3 mg) at 0 °C was added the corresponding ketone (0.5 mmol). The reaction was stirred until the aldehyde was consumed (monitored by TLC). Then, the crude product was diluted in CH₂Cl₂ (10 mL), silica gel was added and the solvent was evaporated in vacuo. The resulting residue was purified by flash chromatography (hexanes–EtOAc) to yield the pure aldol product.
Wet Solvent-Free Conditions: To a mixture of the corresponding aldehyde (0.25 mmol), catalyst **2** (0.0125 mmol, 6.5 mg), benzoic acid (0.0025 mmol, 0.3 mg), H₂O (33 μL) was added at 0 °C followed by the corresponding ketone (0.5 mmol). The reaction was stirred until the aldehyde was consumed. The crude product was diluted in CH₂Cl₂ (10 mL), MgSO₄ was added and filtered. To the filtrate, silica gel was added and the above purification procedure was followed to obtain the aldol product.



change by
 inser
 the new
 scheme

Journal: **SYNLETT**

MS-No.: G 24 2018 RT Author: G. Guilleua

License between Georg Thieme Verlag and the Author of the Manuscript mentioned above

- The Publisher (Georg Thieme Verlag) hereby grants to the author of the article mentioned above, the right and license to make copies from the electronic reprint and to distribute it to individuals who request a reprint of the author's work. Such copying shall be limited to making single copies of the article.
- Georg Thieme Verlag warrants that it is the owner of the exclusive copyright in the licensed material or that it has been granted the copyrights in the licensed material. The author acknowledges and agrees that the licensed material under this agreement is copyright protected and that the exclusive copyright to the licensed material shall remain with Georg Thieme Verlag.
- The electronic reprint of the article may not be used for any (i) fee-for-service use including providing access to or selling copies of items, (ii) systematic supply or distribution of the article in any form to anyone other than individuals who requested a reprint of the article (iii) or any commercial or marketing activity. The electronic reprint or a printout of the electronic reprint of the article or portions of the article shall not be transmitted to libraries or any other organizations. The author is not allowed to post the publication on his personal web site or any other web site that is open to the public.
- The author or any individual receiving an electronic reprint of the mentioned article is not allowed to modify, adapt, transform, translate or create any derivative work based on any materials included in the article, or otherwise use any such materials, in a manner that would infringe the copyright therein. Any copyright notices, other notices or disclaimers included by the Publisher in the online form of the Publications or any accompanying screen displays may not be removed, obscured or modified in any way.
- The author shall promptly notify the Publisher of any infringements of copyrights in the article or any unauthorized use of the electronic reprint of which they become aware. It is the responsibility of the author to alert their colleagues that they may not further distribute the article. Publisher reserves the right to investigate occurrences of suspected unauthorized use or other contract violations and to take appropriate action or to deny the author access to the electronic reprint. Publisher reserves the right to terminate this Agreement for cause or claim damages.

Copyright Statement As stated in the journal imprint, by accepting the author's manuscript, the Publisher shall acquire the exclusive exploitation rights as laid down by the German copyright law. This shall cover in particular - unlimited with respect to the territory or duration - the exclusive right for the first and all subsequent editions/updates (irrespective of the number of pieces of reproduction), also for exploitation in part or in edited form or in the form of a combination with other works, to duplicate the contribution in printed form or in electronic media forms (databases, online network systems, Internet, broadcasting or telecasting, DVD, CD-Rom, hard disk storage, microcopy edition, photomechanical or other reproduction methods, etc.) and to make it accessible to users of such media (e.g., as online or offline backfile) including the right of translation and the right to grant rights to third parties.

Insofar as the above rights can be exercised by a clearing house, the publishers are entitled to conclude relevant agreements with the said clearing house; the publishers are likewise the conclude agreements with third parties in respect of the usufruct of these rights.

Will you please forward any inquiries received by you concerning any of the utilization rights mentioned above, to the publishers for proper settlement.

Attention: Liability The new product liability legislation makes increased demands on the duty of care to be exercised by authors of scientific research. Will you therefore please examine with particular care also in your own interests the factual correctness of contents of the correction and galley proof. You may make this easier for you by arranging to have your statements counter-checked in accordance with the risks theoretically involved in the same, by expert colleagues and co-workers.

During the technical production of books and journals checks are repeatedly made for print errors. Failing expert knowledge on the part of proofreaders, however, all that can be done is to see whether printed matter and manuscript agree. The responsibility for the correctness of data and statements made in the manuscript rests entirely with you as the author.

We should join hands in reducing the risk of right of recourse or claim to damages, and in this regard your responsible cooperation is fundamental.

GEORG THIEME VERLAG KG – STUTTGART • NEW YORK

I agree to the terms of the license agreement for electronic reprints and the copyright statement.

Dr. Gabriela Guilleua
Opto. Química Orgánica
Universidad de Alicante
03080 ANICANTE (Spain)

Name and address (complete and legible, typed or stamped, if possible)

Gabriela Guilleua
Signature