

Natural colorants from filamentous fungi

Fábio Aurélio Esteves Torres¹ · Bruna Regina Zaccarim¹ · Letícia Celia de Lencastre Novaes² ·
Angela Faustino Jozala³ · Carolina Alves dos Santos⁴ · Maria Francisca Simas Teixeira⁵ ·
Valéria Carvalho Santos-Ebinuma¹

Received: 29 October 2015 / Revised: 15 December 2015 / Accepted: 19 December 2015 / Published online: 18 January 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract In the last years, there is a trend towards the replacement of synthetic colorants by natural ones, mainly due to the increase of consumer demand for natural products. The natural colorants are used to enhance the appearance of pharmaceutical products, food, and different materials, making them preferable or attractive. This review intends to provide and describe a comprehensive overview of the history of colorants, from prehistory to modern time, of their market and their applications, as well as of the most important aspects of the fermentation process to obtain natural colorants. Focus is given to colorants produced by filamentous fungal species, aiming to demonstrate the importance of these microorganisms and biocompounds, highlighting the production performance to get high yields and the aspects of conclusion that should be taken into consideration in future studies about natural colorants.

Keywords Natural colorants · Microorganisms · Filamentous fungi · Biomolecules · Bioprocess

Introduction

Color has always fascinated humans and plays an important role in the acceptability of products of several industrial segments, since consumers first judge the quality of a product by its color (Wrolstad and Culve 2011). For centuries, industries use additives to enhance or restore the original color of products and to ensure its uniformity between batches (Medeiros et al. 2012; Fernández-Lopez et al. 2013). Additives used to provide color to substances are called colorants and can be classified as pigments or dyes. The basic difference between them is their size and solubility (Saron and Felisberti 2006). Pigments are practically insoluble in the medium in which they are incorporated while dyes dissolve during application, losing their crystal or particulate structure in the process. Regarding the size, pigment particles are about 1–2 µm and dyes are smaller than it. Furthermore, the difference between pigments and dyes is therefore due to physical characteristics rather than chemical composition (Patel et al. 2012). Natural colorants from microorganisms can be pigments or dyes. To this reason, in this review, it will be called colorants independent of the source.

From prehistoric times, humans have left their mark on environment in the form of painted images, whether in the form of simple handprints, works of fine art, or spray-can graffiti. Many American indigenous civilizations used body and hair painting to communicate. Prehistoric painters used the colorants available next to their homes. The colors yellow ochre, red ochre, and black were obtained from soot from burning animal fat and charcoal from fire (Barnett et al. 2006). Natural colorants, such as bixin, genipin, and

✉ Valéria Carvalho Santos-Ebinuma
valeriac@fcfar.unesp.br

¹ Department of Bioprocess and Biotechnology, School of Pharmaceutical Sciences, UNESP-Universidade Estadual Paulista, Rodovia Araraquara - Jaú Km 1, Araraquara, SP 14801-902, Brazil

² Institute of Biotechnology, RWTH Aachen University, Worringerweg 3, 52074 Aachen, Germany

³ Department of Technological and Environmental Processes, Sorocaba University – UNISO, Sorocaba, SP, Brazil

⁴ Laboratory of Biomaterials and Nanotechnology, University of Sorocaba (UNISO), Sorocaba, SP, Brazil

⁵ Culture Collection DPUA/UFAM, Federal University of Amazonas, Manaus, AM, Brazil

andirobin, were used for esthetic, religious, and protection purposes (Saron et al. 2006).

Egyptians began to develop colorants in 4000 BC. They introduced washed colorants so they could obtain a more colorful and a more pure compound. In 3000 BC, they create the Egyptian blue, derived from calcium copper silicate. They also used malachite and azurite, green and greenish-blue colorants, respectively. These colorants were used in the Egyptian tomb paintings from the fourth dynasty (2613 BC to 2495 BC) and appeared too in European paintings from the fifteenth and sixteenth centuries (Barnett et al. 2006).

Tyrian purple was the first global colorant and was traded by the Phoenicians (Bechtold and Mussak 2009). Roman centuries used red colorants from the marine mollusk *Murex* sp. to dye their tunics, while the Egyptians and the Bretons used natural indigo, extracted from the plant *Isatis tinctoria*. Other colorants used at the same period to color clothes were saffron, extracted from plants, and cochineal carmine, obtained from animals (Zhang et al. 2006). Until the mid-nineteenth century, colorants were essentially obtained from animals and plants (Hill 1997).

In 1856, the English chemist William H. Perkin synthesized Mauve—also known as aniline purple, phenamine, or indisine—the first organic colorant. In 1862, the establishment of a factory that produced this colorant commercially marked the beginning of the modern dye industry (Sousa et al. 2008; Cañamares and Lombardi 2015). Since then, and with the development of alchemy and chemistry, various strongly colored inorganic compounds have been created and used extensively as colorants (Zhang et al. 2006). Therewith, natural colorants were rapidly replaced by chemically synthesized colorants at the end of the nineteenth century and continued until the twentieth century, due to their low cost, large-scale production, flexible location, homogeneity of composition, among others (Bechtold and Mussak 2009).

However, the chemically synthesized colorants were used during many years with little concern or tests regarding their safety. Later, toxicity tests performed by regulatory agencies, such as Food and Drug Administration (FDA), showed the toxic effect of these compounds in the dose administrated by the industries to give color to several substances. So as a result, synthetic colorants can cause significant environmental pollution and have been suggested that they can cause adverse toxicological effects (Malik et al. 2012), such as allergic reactions and intolerance (Wang et al. 2006) and carcinogenic effect (Pan et al. 2009), and affect children's behavior, being hyperactivity the most common disorder (Masone and Chanforan 2015).

As result, concerns about the safety of artificial colorants have been growing, which resulted in the implementation of numerous regulations in several countries, the reduction of the number of synthetic colorants approved for food use by the regulatory agencies (Pan et al. 2009; Rymbai et al. 2011), as well as the return of natural colorants to the market,

particularly in the food sector (Mapari et al. 2010; Mirjalili et al. 2011; Sivakumar et al. 2011). According to a report by “ReportBuyer,” the food color market is projected to reach \$2.5 billion by 2020, growing at a compound annual growth rate (CAGR) of 4.5 %. The European food and beverage colorant market is estimated to grow at a CAGR of 7 % from 2015 to 2020 (<https://www.reportbuyer.com/product/3024599/uk-food-colorants-market-growth-trends-and-forecast-2015-2020.html>). In this way, there is an intense market to be explored not only to food industries as demonstrated but also to other segments such as pharmaceutical, chemistry, among others.

Natural colorant is, by definition, synthesized by and accumulated and/or excreted by living cells (Dabas et al. 2011). It can be extracted from plants, insects, microorganisms, animals, and minerals. They constitute a group of chemically heterogeneous and biosynthetically unrelated molecules that are united by a common feature: their electronic structure contains a chromophore that is responsible for the characteristic colors of these compounds (Pereira et al. 2014). Among them, five are considered as the most important in the global market: bixin, paprika, turmeric, anthocyanins, and cochineal carmine (Constant et al. 2002). However, there are several studies using microorganisms as producers of natural colorants, and efforts have been made in order to reduce the production costs of these compounds compared to synthetic colorants or colorants extracted from natural sources (Dufossé et al. 2014).

Furthermore, the use of fungi for producing colorants is increasing, since many ascomycetes naturally synthesize and secrete colorants and may be considered as a reliable source of natural organic colorants with improved functionality (Mapari et al. 2009a). The diversity of fungal colorants is because of the range of colors that they can produce in some cases with interesting properties such as antioxidant and antitumor characteristics (Mapari et al. 2009a). So, this review will explore the diversity of filamentous fungi producer of colorants, some aspects of their production process, and future applications.

Natural colorants from microbial origin

The production of biomolecules by microorganisms is related to the conditions of the bioprocess. It is important to find the best way to conduct the process and define the nutrient concentrations, not only to reduce production cost but also to increase the colorant production. Therefore, most studies in the literature focus on the improvement of production conditions (Velmurugan et al. 2010a, Hailei et al. 2012, Hu et al. 2012a, Arai et al. 2013, Santos-Ebinuma et al. 2013a, Santos-Ebinuma et al. 2014).

Natural colorants from microorganisms have production advantages compared with similar products extracted from plants or animals, since they do not exhibit the problem of

seasonal availability and are often more stable and soluble (Gunasekaran and Poomiammal 2008). The synthesis of colorants by plant species is slower than that by microorganisms and algae, since the fermentation processes are inherently faster and more productive than other chemical processes (Velmurugan et al. 2010a).

For the industrial perspective, the colorants produced by microorganisms are relevant because it is possible to guarantee that the metabolite of interest will be produced under controlled conditions, regardless of external factors and seasonal supply of raw materials, and the batch-to-batch variability can be minimized (Mapari et al. 2010). Besides, there is a diversity of microorganisms in the tropical and subtropical regions which offers a promising variety of colorant compounds that may be used for different applications (Ventura et al. 2013).

Food industries from Europe and the USA have already obtained natural colorants from microorganisms due to their advantages such as stability, large-scale production, high growth rate, high throughput and wide range of colors and also due to their biological activities, such as antimicrobial, antioxidant, and anticancer properties (Mapari et al. 2009a; Pangestuti and Kim 2011; Teixeira et al. 2012; Tuli et al. 2014). The microorganisms produce a diversity of biocompounds, such as carotenoids, melanins, flavins, quinones, monascines, violaceins, phycocyanins, and indigo (Mapari et al. 2009a; Dufossé et al. 2014). Among the microorganisms, fungi play an important role as colorant producer. So, they will be the focus of this review. Table 1 shows some examples of natural colorants produced by filamentous fungi.

Production process of natural colorants from fungi

Fungi produce a large amount of secondary metabolites that are important for the industry, including the colorant industry. The colorants naturally produced by fungi can be considered as a reliable source of natural organic colorants with improved functionality (Mapari et al. 2009a). The diversity of fungal colorants is not only due to their chemical structure but also due to their range of colors, which may add new or additional hues to the color palette of the existing colorants arising from contemporary sources (Mapari et al. 2009a, Santos-Ebinuma et al. 2013b).

The colorants from fungi can be broadly classified chemically as carotenoids and polyketides (Mapari et al. 2010). Fungal polyketide pigments range in structure from tetraketides to octaketides, which have four or eight C₂ units that contribute to the polyketide chain. Representative classes include the anthraquinones, hydroxyanthraquinones, naphthoquinones, and azaphilone structures, each of which exhibits an array of color hues (Mapari et al. 2010). The structure of the polyketides does not exhibit localized negatively charged ions. These molecules often have polyunsaturated

function, i.e., a ring system, one or more carbonyl groups, carboxylic acid, and functional ester or amide groups that absorb in the UV-visible spectrum (Mapari et al. 2009a). Moreover, polyketides can be bioactive molecules, such as antibiotics, anticancer agents, immunosuppressors, antifungals, and cardiovascular agents (Mapari et al. 2009a).

The oldest colorant in use is angkak, also called red koji, from fermented rice by the fungus *Monascus*. In Europe, the first success story of colorant production by microorganisms was the fungus *Blakeslea* to produce β-carotene. Successful production of carotenoids has been achieved with two microalgae, namely β-carotene using *Dunaliella* and astaxanthin using *Haematococcus* (Dufossé et al. 2005). Fungi *Blakeslea trispora*, *Mucor circinelloides*, and *Phycomyces blakesleeanus* are microorganisms that produce β-carotene colorant, which is a yellowish carotenoid colorant also known as pro-vitamin A (Kumar et al. 2015).

Fungi of the genus *Monascus* are known as great producers of natural colorants. The colorants produced by the species of *Monascus* are the most part azaphilone ones; the six major azaphilone colorants produced by *Monascus* species are monascin and ankaflavin (yellow colorants), monascorubrin and rubropunctatin (orange colorants), and monascorubramine and rubropunctamine (red ones) (Frisvad et al. 2013). Among the genus *Monascus*, the species *M. purpureus* (Shi and Pan 2011; Prajapati et al. 2014) and *M. ruber* (Meinicke et al. 2012, Vendruscolo et al. 2013) are the most studied in terms of colorant production. Previous studies have shown that colorants produced by fungus of the genus *Monascus* exhibit biological activities, such as antimutagenic and anticancer properties, antimicrobial activity, potential anti-obesity activity (Feng et al. 2012), anti-inflammatory (Cheng et al. 2012), antioxidant activities (Hsu and Pan 2012) and as a dietary supplement to hyperlipidemia, ameliorate hypertension, and hypercholesterolemia (Lin et al. 2008). However, certain strains of *Monascus* have also been reported as co-producers of mycotoxin citrinin and other potentially toxic metabolites (Mapari et al. 2010) under certain conditions of cultivation (Lin et al. 2008). Despite of the efforts from several research groups to minimize the mycotoxin production, it is still present in the fermentation process, increasing the urge to find new producers.

Mapari et al. (2008) performed a screening of *Talaromyces* (formerly *Penicillium*) strains that are potential producers of natural colorants with a polyketide structure and chromophores similar to the *Monascus* colorants. The species under analysis were *T. purpurogenus* (formerly *P. purpurogenum*), *T. pinophilus* (formerly *P. pinophilum*), *T. aculeatus* (formerly *P. aculeatum*), *T. funiculosus* (formerly *P. funiculosum*), and *T. minioluteus* (formerly *P. minioluteum*). Among the produced metabolites, monascorubrin, xanthomonasin A, and derivatives of rubropunctatin were found in the extract of *T. aculeatus* (formerly *P. aculeatum*), and monascorubrin was

Table 1 Examples of microbial colorants

Molecule	Color	Microorganism
Ankaflavin	Yellow	<i>Monascus</i> sp.
Antraquinone	Red	<i>Penicillium oxalicum</i> <i>Isaria farinosa</i> (formerly <i>Paecilomyces farinosus</i>)
β-Carotene	Cream	<i>Blakeslea trispora</i>
	Yellow-orange	<i>Fusarium sporotrichioides</i> <i>Mucor circinelloides</i> <i>Neurospora crassa</i> <i>Phycomyces blakesleeanus</i>
	Orange-pink	<i>Monascus roseus</i>
	Red	<i>Blakeslea trispora</i> <i>Fusarium sporotrichioides</i>
Monascin	Red-yellow	<i>Monascus purpureus</i>
Monascorubramin	Red-orange	<i>Monascus</i> sp.
Riboflavin	Yellow	<i>Ashbya gossypii</i> (also known as <i>Eremothecium gossypii</i>)
Rubropunctatin	Orange	<i>Monascus</i> sp.
Unknown	Orange-red	<i>Aspergillus</i> sp.
	Dark-red	<i>Aspergillus glaucus</i>
	Red	<i>Talaromyces purpurogenus</i> (formerly <i>Penicillium purpurogenum</i>) <i>Isaria cicadae</i> (formerly <i>Paecilomyces sinclairii</i>)
	Orange	<i>Penicillium cyclopium</i>
	Yellow	<i>Penicillium nalgiovense</i>

SOURCE: Dufossé 2006; Malik et al. 2012

identified in the extract of *T. pinophilus* (formerly *P. pinophilum*). None of the extracts contained citrinin. Figure 1 shows the chemical structure of some colorants produced by filamentous fungi. Later on, Mapari et al. (2009b) identified ten strains of fungi belonging to four species *T. purpurogenus* (formerly *P. purpurogenum*), *T. aculeatus* (formerly *P. aculeatum*), *T. funiculosus* (formerly *P. funiculosum*), and *T. pinophilus* (formerly *P. pinophilum*) as potential pigment producers that produce *Monascus*-like pigment but with no known mycotoxins.

Frisvad et al. (2013) also studied some *Talaromyces* species as producer of polyketide azaphilone *Monascus* red colorants: *T. aculeatus*, *T. pinophilus*, *T. purpurogenus*, *T. funiculosus*, *T. atroseus*, *T. amestolkiae*, *T. ruber*, and *T. funiculosus*. These species did not produce citrinin; however, some strains can be human pathogens (*T. amestolkiae* and *T. stollii*) or producers of other class of mycotoxins such as rubratoxins A and B by *T. purpurogenus* (Frisvad et al. 2013).


Many colorants with non-carotenoid characteristics are produced by fungi, including quinones, dihydroxynaphthalene melanin (a complex aggregate of polyketides), and flavin components. From quinone class, the pigments anthraquinone and erioglaucine are produced by *Eutotium* spp., *Fusarium* spp., *Curvularia lunata*, and *Drechslera* spp. The first commercial product of the anthraquinone family was the colorant *Arpink*

*Red*TM (Natural RedTM), produced by the company ASCOLOR BIOTECH, Czech Republic, by the fungal strain *Penicillium oxalicum* var. *Armeniaca* CCM 8242 (Dufossé et al. 2014).

Among the species of *Penicillium*, *T. purpurogenus* (formerly *P. purpurogenum*) produces colorants not only in solid medium but also in liquid (Méndez et al. 2011). In studies conducted by Teixeira et al. (2012), natural colorants produced by this strain exhibited significant antimicrobial activity and absence of toxicity against the microcrustacea *Artemia salina*. Figure 2 shows the reverse of the colony of *T. purpurogenus* (formerly *P. purpurogenum*) DPUA 1275 which secreted red colorants according the media used.

There are several submerged culture techniques involving microorganisms. In the batch procedure (also called discontinuous procedure), all the nutrients required for the culture, with the exception of oxygen in aerobic processes or chemical components for pH adjustment, are added before the start of the cultivation, and the product is only removed at the end of each procedure. In the continuous procedure, nutrients are added continuously, and the culture medium is removed at the same speed of the feeding flow, with the volume inside the bioreactor remaining constant (Schmidell et al. 2001). Furthermore, increasing the scale of submerged cultures for further industrial application requires maintaining the uniformity of various conditions of the process (temperature, pH,

Fig. 1 Chemical structure of some colorants produced by filamentous fungi (Mapari et al. 2008)



oxygen concentration, pressure, nutrient concentration, and shear stress), regardless of the volume of the bioreactor.

At bioreactor, the process of natural colorant production using filamentous fungi is a complex process that involves numerous variables, such as aeration rate, mass and heat transfer, and pH (Singh and Kaur 2014). In the submerged culture, the fungal morphology is usually divided into two types: individual filamentous mycelia and dense spherical colonies or pellets. The filamentous mycelia create a highly viscous and pseudoplastic culture medium, which reduces the gas–liquid mass transfer and the homogeneity in the bioreactor (Cui et al. 1998). Although the shape of the pellets generates a less viscous medium (Berovic et al. 1991), limited internal mass transfer can occur (Cui et al. 1998). In addition, the

hydrodynamic conditions in the bioreactor, the tension of the dissolved oxygen, the fungal morphology and growth, the rheology of the medium, and the formation and secretion of products affect the bioprocess (Cui et al. 1998).

Therefore, the rheological characteristics of the liquid medium are affected by the submerged culture of filamentous fungi because of the changes in the composition of the substrate, the concentration of the biomass, and the morphology of microorganism (Cho et al. 2002a). The non-Newtonian behavior is an essential characteristic of the fungal submerged culture systems, in which either filaments or mycelia can predominate. At high microbial concentrations, the hypha can become entangled, resulting in a highly viscous suspension with a non-Newtonian behavior. Many studies have demonstrated that a culture medium containing high concentrations of filamentous microorganisms is highly viscous and is characterized by a shear rate that depends on viscosity and by drainage tension (Cho et al. 2002b). However, the production of natural colorants by fungi has been performed by many authors. Table 2 presents some works found in the literature showing the microorganism and colorant production. It can be observed that the red colorants are the most studied because of its high potential to be applied industrially. Following the works are described in more detail, mainly the production methodology used.

Cho et al. (2002a) studied the influence of the carbon source and the aeration rate on the rheology of the fermentation broth and the mycelial morphology; they also investigated the production of red pigments by *Isaria cicadae* (formerly *Paecilomyces sinclairii*) in a stirred tank-type bioreactor. It




Fig. 2 Reverse of colony of *T. purpurogenus* (formerly *P. purpurogenum*) grown at Czapek Yeast Extract Agar—CYA (a), and agar potato dextrose—BDA (b)

Table 2 Summary of works presented showing microorganism and production of colorants

Colorant	Colorant production	Microorganism	Reference
Red pigments	4.73 g/L in a stirRed tank-type bioreactor	<i>Isaria cicadae</i> (formerly <i>Paecilomyces sinclairii</i>)	Cho et al. (2002a)
Red pigments	4.4 g/L in 5-L batch fermenter	<i>Isaria cicadae</i> (formerly <i>Paecilomyces sinclairii</i>)	Cho et al. (2002b)
Red pigment	2.75 g/L in shaker and 7.7 g/L in a 15-L bioreactor	<i>Candida tropicalis</i> with <i>Penicillium</i> sp. HSD07B	Hailei et al. (2011)
Not described	36.75 OD/g ^a	<i>Monascus purpureus</i>	Velmurugan et al. (2010a)
Red pigment	In orbital shaker was 4.25 and 6.40 g/L in batch reactor	<i>P. novae-zeelandiae</i>	Hailei et al. (2012)
Red pigment	2.46 g/L in orbital shaker	<i>T. purpurogenus</i> (formerly <i>Penicillium purpurogenum</i>)	Mendez et al. (2011)
Red pigments	7.38 UA _{480 nm} in rotary shaker ^a	<i>Monascus ruber</i>	Meinicke et al. (2012)
Red pigments	7.18 ODU/mL in rotary shaker ^a	<i>Monascus purpureus</i>	Sharmila et al. (2013)
Violet pigment PP-V [(10Z)-12-carboxyl- monascorubramine]	– ^b	<i>T. purpurogenus</i> (formerly <i>P. purpurogenum</i>)	Arai et al. (2013)
Red pigment	14.5 U/mL in rotary shaker ^a	<i>Monascus purpureus</i>	Prajapati et al. (2014)
Red colorants	3.19 UA _{490 nm} in rotary shaker ^a	<i>T. purpurogenus</i> (formerly <i>Penicillium purpurogenum</i>)	Santos-Ebinuma et al. (2013a)
Red colorants	2.04 UA _{490 nm} in rotary shaker ^a	<i>T. purpurogenus</i> (formerly <i>Penicillium purpurogenum</i>)	Santos-Ebinuma et al. (2014)
Red pigment	20.44 abs, 500 nm/mg dry fungal biomass in orbital shaker ^a	<i>Monascus purpureus</i>	Singh et al. (2015)
Red pigments	96 mg/L in fed-batch bioreactor	<i>Monascus ruber</i>	Hajjaj et al. (2015)

^a The colorant production can be expressed in different ways. The more used is measuring the corresponding colorant absorbance in the wavelength at maximum absorption of colorants. In this case, the unit used is U/UA which means unity of absorbance or ODU which means optical density

^b Not described

was observed that the specific production rate of the colorant was higher in the medium containing starch. The maximum production of red pigment, 4.73 g/L, was achieved with aeration rate of 3.5 vvm (volume of air per volume of liquid per minute).

Cho et al. (2002b) studied the optimization of culture conditions for colorant production by *I. cicadae* (formerly *P. sinclairii*). They concluded that peptone meat gave higher yield of the colorant production. When performed in a bioreactor batch, production increased almost nine times the production in bioreactor achieving a production of 4.4 g/L of red colorants in a 5-L batch fermenter.

Hailei et al. (2011) cultivated *Candida tropicalis* with *Penicillium* sp. HSD07B and observed the production of a non-toxic and non-mutagenic red colorant. This colorant was stable at pH 2 to 10 and temperatures of 10–100 °C. In addition, the colorant proved to be resistant to oxidation by hydrogen peroxide and to reduction by sodium sulfite, and also a good photostability. Under optimized conditions, the colorant production was of 2.75 g/L in a shaker and 7.7 g/L in a 15-L bioreactor.

Given the high cost of production of red colorant, due to the complexity of the junction of cultures, later on Hailei et al. (2012) studied the production of red colorant through self

immobilization from the biomembrane, formed on the surface of the liquid culture of *Penicillium novae-zeelandiae*. Statistical analysis showed that the concentration of glucose and the diameter of the membrane are the factors that influence colorant production. The strain growth occurred in three phases, wherein the colorant is secreted initially within the membrane, in the deceleration phase, whereby the exhaustion of glucose occurred. After optimization, the maximum red colorant production in orbital shaker was 4.25 g/L and it was found that production increased 61.5 % when subjected to a continuous reactor.

The same authors (Hailei et al. 2012) investigated the mechanism induction of red colorant production in the co-culture and developed a process for the production of it in a pure culture. It was observed that the exhaustion of the first carbon source is the fundamental reason for the production of colorant. *Candida tropicalis* converts glucose to glycerol, ethanol, organic acids, and other substances resulting in the depletion of glucose and secretion of the pigment. The new system of cultivation was the eutrophic-oligotrophic, developed to produce the red colorant during the monoculture of *Penicillium* sp. HSD07B. Although some of pigment components from the monoculture have appeared only with the supply of glycerol, it showed an increase in the production of the

same. Therefore, it was noted that the glucose metabolite accumulation is an important factor in red pigment production in co-culture.

Santos-Ebinuma et al. (2013a) optimized the production of red colorants by *T. purpurogenus* (formerly *P. purpurogenum*), increasing the production by 78 % using statistical tool. To evaluate the stability of colorants, the authors studied several conditions: pH, temperature, and the presence of salts or polymers. The best conditions were pH 8.0 and 70 °C; the colorant also showed good stability in the presence of the salts NaCl and Na₂SO₄ and the polymers polyethylene glycol and sodium polyacrylate.

In another study, Santos-Ebinuma et al. (2014) investigated the production of natural colorants yellow, orange, and red in submerged culture of *T. purpurogenus* (formerly *P. purpurogenum*). Different conditions were evaluated as follows: the effect of incubation time, type and size of the inoculum, and different sources of carbon and nitrogen. The authors obtained the best kinetics growing for all three colorants under conditions of five agar discs with mycelia at 336 h of fermentation. In addition, the most suitable sources of carbon and nitrogen were sucrose and yeast extract, respectively.

Arai et al. (2013) investigated the importance of assimilation of ammonia in the production of amino derivative colorants homologous from *Monascus* by *T. purpurogenus* (formerly *P. purpurogenum*) (PP-V). They also investigated the expression of glutamine synthetase (glnB) and glutamine dehydrogenase (gdh1), two enzymes associated with the conversion of nitrate or nitrite to amino acids and inhibition of the pigment production using L-methionine-DL-sulfoximine (MSX). The genes of gdh1 and glnB were expressed in the PP-V production conditions, and the glnB expression was slightly higher. PP-V biosynthesis was more efficient with Gln use as a nitrogen source than with the use of Glu. With the addition of MSX in half the amount of PP-V decreases markedly. Finally, it is concluded that the ammonia synthesis Gln glutamine synthetase is important for the production of PP-V.

In order to optimize the production of pigments by *M. ruber*, Meinicke et al. (2012) used culture media containing glycerol as a carbon source and monosodium glutamate (MSG) as a nitrogen source. It was found that both the concentrations of MSG and glycerol are important factors, but the concentration of MSG is the most significant variable in the production of the colorant. The best production of red colorants was obtained between 40–70 g/L glycerol and 7–9 g/L MSG. Moreover, it is noted that it is possible to use waste glycerol from biodiesel production for colorants.

Sharmila et al. (2013) examined variables and optimized the production of red pigments by *M. purpureus* by using potato powder as an alternative source of carbon. The maximum absorbance of the pigment was 7.18 ODU/mL and was foreseen for an optimum of

potato powder of 2.5 % (w/v), 0.48 % (w/v) K₂HPO₄, and 0.60 % (w/v) MSG. They concluded that potato powder can be used as a substrate for low-cost production of red colorant on a large scale.

In study performed by Hu et al. (2012b) the perstractive fermentation of intracellular colorants produced by *Monascus* in aqueous non-ionic surfactant Triton X-100 was examined, in which intracellular products were transported to the extracellular micelles in aqueous solution of non-ionic surfactants. It was noted that the relatively low density of the colorant in the microbial cells eliminates the inhibition of the product and prevents it from being degraded. Kang et al. (2013) examined the effects of pH and non-ionic surfactant in the profile of intra- and extracellular *Monascus* colorants. Results showed that extracellular pH influences the absorption of MSG and the exportation of water-soluble red pigment during fermentation of *Monascus*. Regarding the effect of the non-ionic surfactant, it was found that the export of hydrophobic colorants was affected by intracellular perstractive fermentation and independent of pH. In this fermentation, the production of red and yellow colorants extracellular was higher when the initial pH were, respectively, higher and lower.

Mendez et al. (2011) studied the effect of temperature and pH on the production of red colorants by *T. purpurogenus* (formerly *P. purpurogenum*) in submerged culture. It was observed that maximum production of red pigment (2.46 g/L) was obtained at pH 5 and 34 °C, and the maximum concentration of biomass was obtained at pH 7 and 34 °C. From these results, it was found that there is a direct relationship between the production of colorant and biomass concentration. Furthermore, the maximum pigment production was associated with strong combination of pH and temperature effects.

Prajapati et al. (2014) studied the increase in the production of red colorant by submerged culture of *M. purpureus* through optimization. The results indicated that glucose, tryptone, and pH of the medium are highly important in colorant production. Moreover, unlike other researchers reported, in this study, the best nitrogen source is tryptone, instead of MSG. The maximum production of the colorant (OD 14.5 U/mL) was achieved with 58 g/L glucose, 1 g/L tryptone, and pH 8.

Singh et al. (2015) investigated the application of artificial neural network (ANN) in modeling a liquid state fermentation for red colorant production by *M. purpureus* MTCC 369 using rice water base medium. They concluded that rice water medium is an effective substrate for producing the red colorant without any external nitrogen source and, also, that the ANN analysis provides a dynamic model for successful prediction of optimal operating conditions.

Hamano et al. (2005) showed that the specific production of extracellular red colorant by different strains of *Monascus* sp. is 1.7–3.5 higher than the specific production of intracellular red colorant in semi-synthetic medium on a rotary shaker.

Mukherjee and Singh (2011) purified and characterized a new red colorant from *M. purpureus* in submerged fermentation. The colorant was extracted using hexane and purified by thin layer chromatography, column chromatography, and semi-preparative HPLC. By UV-vis, IR, GC-MS, and NMR analyses, the pigment was characterized as 9-(1-hydroxyhexyl)-3-(2-hydroxypropyl)-6a-methyl-9,9a-dihydrofuro[2,3-h]isoquinoline-6,8(2*H*,6*aH*)-dione; this colorant presented similarities with rubropunctamine and monascorubramine, but it presented an hydroxyl alkane substitutions at C-3 and C-9 position. This colorant also showed antibacterial activity against Gram-positive bacteria. The authors indicate that in higher pH, consolidation of red colorant production was observed and found that 18 g/L of glucose is optimum for red colorant production.

Velmurugan et al. (2010a) studied the effects of total darkness and different shades of light in the extra- and intracellular production of colorants and biomass of five fungi: *M. purpureus*, *Isaria farinosa*, *Emericella nidulans*, *Fusarium verticillioides*, and *T. purpurogenus* (formerly *P. purpurogenum*). It was noted that in all fungi, incubation in the dark increased biomass and colorant production. In addition, it was concluded that in all of them, there is no effect on growth under red light, and under green or yellow light, the growth decreases. Maximum production of extra- and intracellular red colorants by *M. purpureus* occurred in the dark and was, respectively, 36.75 and 18.27 OD/g.

Hajjaj et al. (2015) focused on the influence of nitrogen/carbon ratio and mode of cultivation on the production of red colorants and citrinin in *M. ruber*. Using monosodium glutamate and glucose, they confirmed that the max production of red colorant was at a ratio around 10 of C/N; citrin production increased with the increase of the C/N ratio. They showed, too, that the successive feed of glucose with low C/N ratio induced the accumulation of L-malic acid, which is unfavorable for red colorant production.

Velmurugan et al. (2011) investigated the production of red colorants by *M. purpureus* using corn cob powder as a substrate for solid-state fermentation. The higher production of pigment was 25.42 OD/g after 7 days of incubation at 30 °C, and inoculated with 4 mL spores/g and initial moisture content of 60 %. The pigments are stable in acidic pH, at high temperature, and in saline solutions. It was found that the production of colorants using corn cob powder is much higher than with the use of other agricultural waste.

Application of natural colorants

Natural colorants are widely used in the textile (Islam et al. 2013), paper (Malik et al. 2012), leather (Guillén et al. 2012), and timber industries (Ali, 2010). Foods are often colored with natural or synthetic colorants approved by a regulatory agency (Carocho et al. 2014). It is also used to provide color to petroleum derivatives, such as waxes, lubricant oils, ointments, and gasoline; plastics; resins; and rubber products (Kumari et al. 2012). One of the important issues in the application of colorants is safety, so there are many limitations and legislation on the use of dyes in different countries (Nikfar and Jaberidoost, 2014).

Also, there is an increasing demand for natural non-toxic and/or eco-friendly colorants on industry, particularly those to be applied in the healthcare area, food, children's toys, and leather elements (Velmurugan et al. 2010b).

It is hard to estimate the market for natural colorants produced by bioprocess, although there is an increasing preference for the use of natural additives in food and cosmetic products. According to Mapari et al. (2010), natural colorants accounted for 31 % of the colorants market, whereas synthetic colorants made up 40 % and nature-identical colorants accounted for 29 % of the market. In addition, the Leatherhead Food International points out that the natural colorants market is growing and that these products will replace synthetic colorants in the future. In 2007, the colorants market in the food sector was estimated at USD 1.15 billion, a 2.5 % increase compared to 2004 (USD 1.07 billion); from this, USD 465 million was from natural colorants, a 4.6 % increase compared to 2004 (Mapari et al. 2010).

Natural colorants derived from algae or extracted from flowering plants have been used in food and nutritional supplements showing the importance of niche markets in which consumers are willing to pay a premium for “natural healthy ingredients” (Dufossé et al. 2014). In the pharmaceutical industry, the colorants produced by filamentous fungi can be interesting because they can not only promote color but also have potential clinical application since they can act as anticancer, antiproliferative, antibiotics, etc. In the textile area, the use of non-allergic, non-toxic, and eco-friendly natural colorants has increased due to environmental awareness in order to avoid some synthetic colorants with hazardous characteristics (Samanta and Agarwal 2009).

Velmurugan et al. (2010b) studied the dyeing potential of five different water-soluble colorants extracted from *M. purpureus*, *Isaria* spp., *Emericella* spp., *Fusarium* spp., and *Penicillium* spp. The authors optimized the dyeing process in leather resulting in uniform dyeing and intense and bright shade. According to the authors, the results obtained indicated that the fungal colorants could be an alternate for leather dyeing as a natural dye.

Some species of filamentous fungi such as *P. oxalicum* and *Monascus* species, which produce colorants that have been used as food colorants for hundreds of years in the Far East, *Phaffia rhodozoa*, *Ashbya gossypii*, and *Blaskelea trispora* are in the stage of industrial production. However, the most part of natural colorants produced by fungus is at development or research stage, and in order to obtain success in the application of these natural colorants, the approval of their application by the regulatory agencies is essential. In this way, there is a vast field of study to be explored involving new producing species and new molecules with biological activity and without the associated production of mycotoxins and studies of toxicity.

Conclusion

Since the health-conscious consumer has had an upward growth, the natural colorant application has been increasing in the last years in different industries from food until textile coloration. For this reason, the natural colorants produced by microorganisms have been deeply studied. Although *Monascus* is the more studied specie, other filamentous fungi have been evaluated resulting in colorants with diversified chemical structure, properties, and applications. This review has highlighted studies that utilized different filamentous fungi species to produce high yields of the natural colorants, indicating the biological activity and its application. However, there is a lack of deeper studies of the use of bioreactors for natural colorant production, as well as their toxicology studies, which are necessary to the industrialization and introduction of these biotechnological products on the market.

Acknowledgments This research was supported by grants from the Coordination for Higher Level Graduate Improvements (CAPES/Brazil), National Council for Scientific and Technological Development (CNPq/Brazil), State of São Paulo Research Foundation (FAPESP/Brazil, Process number 2014/01580-3), and Development Support Programme Science of Pharmaceutical Sciences College of UNESP Araraquara (PADC, Process number 22/2014).

Compliance with ethical standards

Ethical statement The authors, whose names appear on the submission, declare that they have contributed sufficiently to the scientific work and therefore share collective responsibility and accountability for the results. This manuscript has not been published or presented elsewhere in part or in entirety and is not under consideration by another journal. This research does not involve human participants or animals.

All the authors have approved the manuscript and agree with submission to your esteemed journal.

Conflict of interest The authors declare that they have no competing interests.

References

- Ali H (2010) Biodegradation of synthetic dyes—a review. *Water Air Soil Poll* 2013:251–273. doi:10.1007/s11270-010-0382-4
- Arai T, Koganei K, Umemura S, Kojima R, Kato J, Kasumi T, Ogihara J (2013) Importance of the ammonia assimilation by *Penicillium purpurogenum* in amino derivative *Monascus* pigment, PP-V, production. *AMB Express* 3:19. doi:10.1186/2191-0855-3-19
- Barnett JR, Miller S, Pearce E (2006) Colour and art: a brief history of pigments. *Opt Laser Technol* 38:445–453. doi:10.1016/j.optlastec.2005.06.005
- Bechtold T, Mussak R. (2009) Front Matter, in Handbook of natural colorants. John Wiley & Sons, Ltd, Chichester, UK. doi:10.1002/9780470744970.fmatter
- Berovic M, Cimerman A, Steiner W, Koloin T (1991) Submerged citric acid fermentation: rheological properties of *Aspergillus niger* broth in a stirred tank reactor. *Appl Microbiol Biotechnol* 34:579–581. doi:10.1007/BF00167902
- Cañamares MV, Lombardi JR (2015) Raman, SERS, and DFT of mauve dye: adsorption on Ag nanoparticles. *J Phys Chem C* 119:14297–14303. doi:10.1021/acs.jpcc.5b02619
- Carocho M, Barreiro MF, Morales P, Ferreira ICFR (2014) Adding molecules to food, pros and cons: a review of synthetic and natural food additives. *Compr Rev Food Sci F* 13:377–399. doi:10.1111/1541-4337.12065
- Cheng MJ, Wua MD, Yuan GF, Su YS, Yanai H (2012) Secondary metabolites produced by the fungus *Monascus pilosus* and their anti-inflammatory activity. *Phytochem Lett* 5:567–571. doi:10.1016/j.phytol.2012.05.015
- Cho YJ, Hwang HJ, Kim SW, Song CH, Yun JW (2002a) Effect of carbon source and aeration rate on broth rheology and fungal morphology during red pigment production by *Paecilomyces sinclairii* in a batch bioreactor. *J Biotech* 95:13–23. doi:10.1016/S0168-1656(01)00445-X
- Cho YJ, Park JP, Hwang HJ, Kim SW, Choi JW, Yun JW (2002b) Production of red pigment by submerged culture of *Paecilomyces sinclairii*. *Lett Appl Microbiol* 35:195–202. doi:10.1046/j.1472-765X.2002.01168.x
- Constant PBL, Strungheta PC, Sandi D (2002) Corantes alimenticios. *Bol Centro Pesqui Process Aliment* 20:203–220. doi:10.5380/cep.v20i2.1248
- Cui YQ, Van der Lans RG, Luyben KC (1998) Effects of dissolved oxygen tension and mechanical forces on fungal morphology in submerged fermentation. *Biotechnol Bioeng*. 57:409–419. doi:10.1002/(SICI)1097-0290(19980220)
- Dabas D, Elias RJ, Lambert JD, Ziegler GR (2011) A colored avocado seed extract as a potential natural colorant. *J Food Sci* 76:1335–1341. doi:10.1111/j.1750-3841.2011.02415.x
- Dufossé L, Fouillaud M, Caro Y, Mapari SAS, Sutthiwong N (2014) Filamentous fungi are large-scale producers of pigments and colorants for the food industry. *Curr Opin Biotechnol* 56:56–61. doi:10.1016/j.copbio.2013.09.007
- Dufossé L (2006) Microbial production of food grade pigments. *Food Tech Biotechnol* 44:313–321. doi:10.1016/j.copbio.2013.09.007
- Dufossé L, Galaup P, Yaron A, Arad SM, Blanc P, Murthy KNC, Ravishankar GA (2005) Microorganisms and microalgae as sources of pigments for food use: a scientific oddity or an industrial reality? *Trends Food Sci Tech* 16:389–406. doi:10.1016/j.tifs.2005.02.006
- Feng Y, Shao Y, Chen F (2012) *Monascus* pigments. *Appl Microbiol Biotechnol* 96:1421–1440. doi:10.1007/s00253-012-4504-3
- Fernández-Lopez JA, Angosto JM, Giménez PJ, León G (2013) Thermal stability of selected natural red extracts used as food colorants. *Plant Foods Hum Nutr* 68:11–17. doi:10.1007/s11130-013-0337-1
- Frisvad JC, Yilmaz N, Thrane U, Rasmussen KB, Houbraken J, Samson RA (2013) *Talaromyces atroroseus*, a new species efficiently

- producing industrially relevant red pigments. *PlosONE* 8:e84102. doi:10.1371/journal.pone.0084102
- Guillén D, Ginebreda A, Petrovic M, Barceló D, Darbra RM, Rydin S (2012) Global risk-based management of chemical additives I. The handbook of environmental chemistry. Springer, London
- Gunasekaran S, Poorniammal R (2008) Optimization of fermentation conditions for red pigment production from *Penicillium* sp. under submerged cultivation. *Afr J Biotechnol* 7:1894–1898. doi:10.1111/jam.12110
- Hailei W, Ping L, Yufeng L, Zhifang R, Gang W (2012) Overproduction of a potential red pigment by a specific self-immobilization biomembrane-surface liquid culture of *Penicillium novae-zeelandiae*. *Bioproc Biosyst Eng* 35:1407–1416. doi:10.1007/s00449-012-0729-x
- Hailei W, Zhifang R, Ping L, Yanchang G, Guosheng L, Jianming Y (2011) Improvement of the production of a red pigment in *Penicillium* sp. HSD07B synthesized during co-culture with *Candida tropicalis*. *Bioresour Technol* 102:6082–6087. doi:10.1016/j.biortech.2011.01.040
- Hajjaj H, Goma G, François JM (2015) Effect of the cultivation mode on red pigments production from *Monascus ruber*. *Int J Food Sci Tech* 50:1731–1736. doi:10.1111/ijfs.12803
- Hamano PS, Orozco SFB, Kilikian BV (2005) Concentration determination of extracellular and intracellular red pigments produced by *Monascus* sp. *Braz Arch Biol Technol* 48:43–49. doi:10.1590/S1516-89132005000400006
- Hill DJ (1997) Is there a future for natural dyes? *Rev Prog Color* 27:18–25. doi:10.1111/j.1478-4408.1997.tb03771.x
- Hsu WH, Pan TM (2012) *Monascus purpureus* fermented products and oral cancer: a review. *Appl Microbiol Biotechnol* 93:1831–1842. doi:10.1007/s00253-012-3891-9
- Hu Z, Zhang X, Wu Z, Qi H, Wang Z (2012a) Perstraction of intracellular pigments by submerged cultivation of *Monascus* in nonionic surfactant micelle aqueous solution. *Appl Microbiol Biotechnol* 94:81–89. doi:10.1007/s00253-011-3851-9
- Hu Z, Zhang X, Wu Z, Qi H, Wang Z (2012b) Export of intracellular *Monascus* pigments by two-stage microbial fermentation in nonionic surfactant micelle aqueous solution. *J Biotechnol* 162:202–209. doi:10.1016/j.jbiotec.2012.10.004
- Islam S, Shahid M, Mohammad F (2013) Perspectives for natural products based agents derived from industrial plants in textile applications—a review. *J Clean Prod* 57:2–18. doi:10.1016/j.jclepro.2013.06.004
- Kang B, Zhang X, Wu Z, Qi H, Wang Z (2013) Effect of pH and nonionic surfactant on profile of intracellular and extracellular *Monascus* pigment. *Process Biochem* 48:759–767. doi:10.1016/j.procbio.2013.03.020
- Kumari LS, Rao PP, Sameera S, Koshy P (2012) Synthesis and optical properties of $Ce_{0.95}Pr_{0.05-x}M_xO_2$ (M = Mn, Si) as potential ecological red pigments for coloration of plastics. *Ceram Int* 38:4009–4016. doi:10.1016/j.ceramint.2012.01.057
- Kumar A, Vishwakarma HS, Singh J, Dwivedi S, Kumar M (2015) Microbial pigments: production and their applications in various industries. *Int J Pharm Chem Biol Sci* 5:203–212
- Lin YL, Wang TH, Lee MH (2008) Biologically active components and nutraceuticals in the *Monascus*-fermented rice: a review. *Appl Microbiol Biotechnol* 77:965–973. doi:10.1007/s00253-007-1256-6
- Malik K, Tokka J, Goyal S (2012) Microbial pigments: a review. *International J Microbial Resource Technol* 1(4):361–365
- Mapari SAS, Hansen ME, Meyer AS, Thrane U (2008) Computerized screening for novel producers of *Monascus*-like food pigments in *Penicillium* species. *J Agric Food Chem* 56:9981–9989. doi:10.1021/jf801817q
- Mapari SAS, Meyer AS, Thrane U (2009a) Photostability of natural orange-red and yellow fungal pigments in liquid food model systems. *J Agr Food Chem* 57:6253–6261. doi:10.1021/jf900113q
- Mapari SAS, Meyer AS, Thrane U, Frisvad JC (2009b) Identification of potentially safe promising fungal cell factories for the production of polyketide natural food colorants using chemotaxonomic rationale. *Microb Cell Factories* 8:24. doi:10.1186/1475-2859-8-24
- Mapari SAS, Thrane U, Meyer AS (2010) Fungal polyketide azaphilone pigments as future natural food colorants? *Trends Biotechnol* 28:300–307. doi:10.1016/j.tibtech.2010.03.004
- Masone D, Chanforan C (2015) Study on the interaction of artificial and natural food colorants with human serum albumin: a computational point of view. *Comput Biol Chem* 56:152–158. doi:10.1016/j.compbiolchem.2015.04.006
- Medeiros RA, Lourenço BC, Rocha-Filho RC, Fatibello-Filho O (2012) Simultaneous voltammetric determination of synthetic colorants in food using a cathodically pretreated boron-doped diamond electrode. *Talanta* 97:291–297. doi:10.1016/j.talanta.2012.04.033
- Meinicke RM, Vendruscolo F, Esteves Moritz D, Oliveira D, Schmidell W, Samohyl RW, Ninow JL (2012) Potential use of glycerol as substrate for the production of red pigments by *Monascus ruber* in submerged fermentation. *Biocatal Agric Biotechnol* 1:238–242. doi:10.1016/j.cbab.2012.03.001
- Méndez A, Pérez C, Montañez JC, Martínez G, Aguilar CN (2011) Red pigment production by *Penicillium purpurogenum* GH2 is influenced by pH and temperature. *J Zhejiang Univ Sci B* 12:961–968. doi:10.1631/jzus.B1100039
- Mirjalili M, Nazarpour K, Karimi L (2011) Eco-friendly dyeing of wool using natural dye from weld as co-partner with synthetic dye. *J Clean Prod* 19:1045–1051. doi:10.1016/j.jclepro.2011.02.001
- Mukherjee G, Singh SK (2011) Purification and characterization of a new red pigment from *Monascus purpureus* in submerged fermentation. *Proc Biochem* 46:188–192. doi:10.1016/j.procbio.2010.08.006
- Nikfar S, Jaberidoost M (2014) Dyes and Colorants. *Reference Module in Biomedical Sciences. Encyclopedia of Toxicology*, third edition, Elsevier, p. 252–261
- Pan Y, Zhu Z, Huang Z, Wang H, Liang Y, Wang K, Lei Q, Liang M (2009) Characterisation and free radical scavenging activities of novel red pigment from *Osmanthus fragrans* seeds. *Food Chem* 112:909–913. doi:10.1016/j.foodchem.2008.06.077
- Pangestuti R, Kim SK (2011) Biological activities and health benefit effects of natural pigments derived from marine algae. *J Func Foods* 3:2255–2266. doi:10.1016/j.jff.2011.07.001
- Patel SK, Saurabh KK, Kamlesh K, Mali MD (2012) Studies on synthesis of novel low molecular weight anthraquinone disperse dyes and their application on polyester and nylon. *J Ind Chem Soc* 89:789–795
- Pereira DN, Valentão P, Andrade PB (2014) Marine natural pigments: chemistry, distribution and analysis. *Dyes Pigments* 111:124–134. doi:10.1016/j.dyepig.2014.06.011
- Prajapati VS, Soni N, Trivedi UB, Patel KC (2014) An enhancement of red pigment production by submerged culture of *Monascus purpureus* MTCC 410 employing statistical methodology. *Biocatal Agric Biotechnol* 3:140–145. doi:10.1016/j.cbab.2013.08.008
- Rymbai H, Sharma RR, Srivastav M (2011) Biocolorants and its implications in health and food industry—a review. *Int J PharmTech Res* 3:2228–2244
- Samanta AK, Agarwal P (2009) Application of natural dyes on textiles. *Indian J Fibre Text* 34:384–399.
- Santos-Ebinuma VC, Roberto IC, Teixeira MFS, Pessoa A (2014) Improvement of submerged culture conditions to produce colorants by *Penicillium purpurogenum*. *Braz J Microbiol* 45:731–742. doi:10.1590/S1517-83822014000200049
- Santos-Ebinuma VC, Roberto IC, Teixeira MFS, Pessoa A (2013a) Improving of red colorants production by a new *Penicillium purpurogenum* strain in submerged culture and the effect of different

- parameters in their stability. *Biotechnol Prog* 29:778–785. doi:10.1002/btpr.1720
- Santos-Ebinuma VC, Teixeira MFS, Pessoa A (2013b) Submerged culture conditions for the production of alternative natural colorant by a new isolated *Penicillium purpurogenum* DPUA 1275. *J Microbiol Biotechnol* 23:802–810. doi:10.4014/jmb.1211.11057
- Saron C, Felisberti MI (2006) Ação de colorantes na degradação e estabilização de polímeros. *Quím Nov*. 29:124–128. doi:10.1590/S0100-40422006000100022
- Sharmila G, Nidhi B, Muthukumaran C (2013) Sequential statistical optimization of red pigment production by *Monascus purpureus* (MTCC 369) using potato powder. *Ind Crop Prod* 44:158–164. doi:10.1016/j.indcrop.2012.11.007
- Schmidell W, Lima AU, Aquarone E, Borzani W (2001) *Biocatal Industrial*. Blücher, São Paulo
- Shi YC, Pan TZ (2011) Beneficial effects of *Monascus purpureus* NTU568-fermented products: a review. *Appl Microbiol Biotechnol* 90:1207–1217. doi:10.1007/s00253-011-3202-x
- Singh N, Goel G, Singh N, Pathak BK, Kaushik D (2015) Modeling the red pigment production by *Monascus purpureus* MTCC 369 by artificial neural network using rice water based medium. *Food Bioscience II* 11:17–22. doi:10.1016/j.fbio.2015.04.001
- Singh D, Kaur G (2014) Swainsonine, a novel fungal metabolite: optimization of fermentative production and bioreactor operations using evolutionary programming. *Bioproc Biosyst Eng* 37:1599–1607. doi:10.1007/s00449-014-1132-6
- Sivakumar V, Vijaaeswarri J, Anna JL (2011) Effective natural dye from different plant materials using ultrasound. *Ind Crop Prod* 33:116–122. doi:10.1016/j.indcrop.2010.09.007
- Sousa MM, Melo MJ, Parola AJ, Morris PJ, Rzepa HS, de Melo JS (2008) A study in mauve: unveiling Perkin's dye in historic sample. *Chem Eur J* 14:8507–8513. doi:10.1002/chem.200800718
- Teixeira MFS, Martins MS, Silva JC, Kirsch LS, Fernandes OCC, Carneiro ALB, Conti R, Durán N (2012) Amazonian biodiversity: pigments from *Aspergillus* and *Penicillium*-characterizations, antibacterial activities and their toxicities. *Curr Trends Biotechnol Pharm* 6:300–311
- Tuli HS, Chaudhary P, Beniwal V, Sharma AK (2014) Microbial pigments as natural color sources: current trends and future perspectives. *J Food Sci Technol* 52:4669–4678. doi:10.1007/s13197-014-1601-6
- Velmurugan P, Hur H, Balachandar V, Kamala-Kannan S, Lee KJ (2011) *Monascus* pigment production by solid-state fermentation with corn cob substrate. *J Biosci Bioeng* 116:590–594. doi:10.1016/j.jbiosc.2011.08.009
- Velmurugan P, Lee YH, Venil CK, Lakshmanaperumalsamy P, Chae JC, Oh BT (2010a) Effect of light on growth, intracellular and extracellular pigment production by five pigment-producing filamentous fungi in synthetic medium. *J Biosci Bioeng* 109:346–350. doi:10.1016/j.jbiosc.2009.10.003
- Velmurugan P, Kamala-Kannan S, Balachandar V, Lakshmanaperumalsamy P, Chae JC, Oh BT (2010b) Natural pigment extraction from five filamentous fungi for industrial applications and dyeing of leather. *Carbohydr Polym* 79:262–268. doi:10.1016/j.carbpol.2009.07.058
- Vendruscolo F, Müller BL, Moritz DE, Oliveira D, Schmidell W, Ninow JL (2013) Thermal stability of natural pigments produced by *Monascus ruber* in submerged fermentation. *Biocatal Agric Biotechnol* 2:278–284. doi:10.1016/j.bcab.2013.03.008
- Ventura SPM, Santos-Ebinuma VC, Pereira JF, Teixeira MF, Pessoa A, Coutinho JA (2013) Isolation of natural red colorants from fermented broth using ionic liquid-based aqueous two-phase systems. *J Ind Microbiol Biot* 40:507–516. doi:10.1007/s10295-013-1237-y
- Wang HS, Pan YM, Tang XJ, Huang ZQ (2006) Isolation and characterization of melanin from *Osmanthus fragrans*' seeds. *LWT-Food Sci Technol* 39:496–502. doi:10.1016/j.lwt.2005.04.001
- Wrolstad RE, Culve CA (2011) Alternatives to those artificial FD&C food colorants. *Annu Rev Food Sci Technol* 3:59–77. doi:10.1146/annurev-food-022811-101118
- Zhang H, Zhan J, Su K, Zhang Y (2006) A kind of potential food additive produced by *Streptomyces coelicolor*: characteristics of blue pigment and identification of a novel compound, λ -actinorhodin. *Food Chem* 95:186–192. doi:10.1016/j.foodchem.2004.12.028