

Natural genetic variation in *Arabidopsis* for responsiveness to plant growth-promoting rhizobacteria

Paul C. A. Wintermans¹ · Peter A. H. M. Bakker¹ · Corné M. J. Pieterse¹

Received: 10 December 2015 / Accepted: 15 January 2016 / Published online: 30 January 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract The plant growth-promoting rhizobacterium (PGPR) *Pseudomonas simiae* WCS417r stimulates lateral root formation and increases shoot growth in *Arabidopsis thaliana* (*Arabidopsis*). These plant growth-stimulating effects are partly caused by volatile organic compounds (VOCs) produced by the bacterium. Here, we performed a genome-wide association (GWA) study on natural genetic variation in *Arabidopsis* for the ability to profit from rhizobacteria-mediated plant growth-promotion. To this end, 302 *Arabidopsis* accessions were tested for root architecture characteristics and shoot fresh weight in response to exposure to WCS417r. Although virtually all *Arabidopsis* accessions tested responded positively to WCS417r, there was a large variation between accessions in the increase in shoot fresh weight, the extra number of lateral roots formed, and the effect on primary root length. Correlation analyses revealed that the bacterially-mediated increase in shoot fresh weight is related to alterations in root architecture. GWA mapping for WCS417r-stimulated changes in root and shoot growth characteristics revealed 10 genetic loci highly associated with the responsiveness of *Arabidopsis* to the plant growth-promoting activity of WCS417r. Several of the underlying candidate genes have been implicated in important plant growth-related processes. These results demonstrate that plants possess

natural genetic variation for the capacity to profit from the plant growth-promoting function of a beneficial rhizobacterium in their rhizosphere. This knowledge is a promising starting point for sustainable breeding strategies for future crops that are better able to maximize profitable functions from their root microbiome.

Keywords *Arabidopsis thaliana* · Genome-wide association (GWA) · Plant growth-promoting rhizobacteria (PGPR) · *Pseudomonas simiae* WCS417r · Volatile organic compounds (VOCs)

Introduction

Beneficial microbes in the microbiome of plant roots provide important services to the plant as they improve plant nutrition and provide protection against plant pathogens (Berendsen et al. 2012; Bulgarelli et al. 2013; Philippot et al. 2013). In recent years, microarray and next-generation sequencing technologies provided detailed insight into the composition and assembly of microbiota communities on and in plants roots (Bai et al. 2015; Bulgarelli et al. 2012; Hardoim et al. 2015; Lareen et al. 2016; Lebeis et al. 2015; Lundberg et al. 2012; Mendes et al. 2011). Genotypic variations in host plants can influence the composition of the root microbiome with important adaptive consequences for plant health (Haney et al. 2015; Pieterse et al. 2016). Evidence is accumulating that domestication of crop species changed root microbiome characteristics in a plant genotype-dependent manner (Bulgarelli et al. 2015; Peiffer et al. 2013). It has been postulated that during the process of plant domestication, plant breeding for high yields under conditions of high fertilizer input in soils has resulted in the erosion of plant genotypic traits involved in

Electronic supplementary material The online version of this article (doi:10.1007/s11103-016-0442-2) contains supplementary material, which is available to authorized users.

✉ Corné M. J. Pieterse
C.M.J.Pieterse@uu.nl

¹ Plant–Microbe Interactions, Department of Biology, Faculty of Science, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands

beneficial plant–microbe interactions (Pérez-Jaramillo et al. 2016). Hence, the identification of loci associated with the capacity of plants to maximize profitable functions from their root microbiome will be highly instrumental for sustainable plant breeding strategies that are directed towards optimizing root microbiome functions in crops.

Plant root-colonizing bacteria that have a positive effect on plant growth are collectively known as plant growth-promoting rhizobacteria (PGPR). Mechanisms of growth promotion have been studied in detail and different PGPR may have different modes of action. Many PGPR stimulate plant growth through the inhibition of plant pathogens. For some PGPR this is due to direct inhibition of growth and activity of pathogens (Doornbos et al. 2012; Raaijmakers and Weller 1998), whereas others can suppress diseases indirectly by eliciting induced systemic resistance (ISR), a process that is characterized by priming plant-borne defense mechanisms to respond faster and stronger to pathogen attack (Conrath et al. 2015; Pieterse et al. 2014). Numerous rhizosphere inhabiting bacteria can elicit ISR in plants, including *Pseudomonas* spp. (Bakker et al. 2007; Pieterse et al. 2014) and *Bacillus* ssp. (Kloepper et al. 2004). Also abiotic stresses like salinity and drought stress can be alleviated by PGPR (Yang et al. 2009). Finally, PGPR can exhibit a direct influence on plant growth and development (Lugtenberg and Kamilova 2009) and this phenomenon is the focus of this study.

A common way by which PGPR directly promote plant growth is by aiding in the uptake of essential nutrients. For example, *Rhizobium* species stimulate the formation of root nodules in which atmospheric nitrogen is fixed and made available to the plant (Van Rhijn and Vanderleyden 1995). Other PGPR free up scarce nutrients in the soil, like phosphate (Browne et al. 2009), or induce plant uptake mechanisms for nitrate (Mantelin and Touraine 2004) or iron (Zamioudis et al. 2014, 2015; Zhang et al. 2009). PGPR can also influence plant growth by the production of plant hormones like auxin, abscisic acid, cytokinins, and gibberellins (Hayat et al. 2010), or influence plant ethylene concentrations by production of the enzyme 1-aminocyclopropane-1-carboxylic acid (ACC) deaminase that breaks down ACC, the direct precursor of ethylene (Glick et al. 1998; Penrose et al. 2001). The latter process can effectively lower local ethylene concentrations resulting in increased root and shoot biomass (Glick et al. 2007).

More recently, volatile organic compounds (VOCs) have been reported to be involved in plant–microbe interactions, including rhizobacteria-mediated plant growth promotion (Blom et al. 2011; Junker and Tholl 2013). An early study reporting on VOCs and plant growth promotion deals with a *Bacillus subtilis* strain that promotes growth of *Arabidopsis thaliana* (Arabidopsis) via the bacterial VOCs acetoin and 2,3-butanediol (Ryu et al. 2003). Also root

system architecture in Arabidopsis can be modulated by VOCs (Gutierrez-Luna et al. 2010). For instance, bacterial indole promotes root growth by interfering with auxin signaling in Arabidopsis (Bailly et al. 2014; Bhattacharyya et al. 2015). Moreover, Meldau et al. (2013) reported that the VOC dimethyl disulfide produced by *Bacillus* sp. B55 promotes growth in *Nicotiana attenuata*. Whereas most VOCs seem to act as inter-kingdom signaling molecules, dimethyl disulfide was suggested to be metabolized by the plant, thus reducing the need for energy costly sulfur uptake.

In Arabidopsis, colonization of the roots by the PGPR *Pseudomonas simiae* WCS417r (formerly known as *Pseudomonas fluorescens* WCS417r; Berendsen et al. 2015) can increase shoot fresh weight by approximately 30 % when co-cultivated in soil (Pieterse and Van Loon 1999). Using a split-plate system, it was shown that this growth-promoting effect is partly mediated by bacterial VOCs and unrelated to the ISR-inducing capacity of WCS417r (Zamioudis et al. 2013). Co-cultivating Arabidopsis accession Col-0 with WCS417r leads to inhibition of primary root elongation and promotion of lateral root and root hair formation, resulting in significant changes in the root architecture and increased shoot fresh weight (Zamioudis et al. 2013). This bacterially-induced process requires the action of the plant growth regulator auxin.

The involvement of microbial VOCs in plant growth promotion is typically studied by co-cultivating plants and PGPR in sealed Petri dishes in which microbially-produced CO₂ can accumulate. Hence, it has been hypothesized that CO₂ produced by the bacteria causes the growth promotion through the enhanced availability of this photosynthesis substrate. Whereas there is some evidence that CO₂ can be partially involved (Kai and Piechulla 2009), the growth response stimulated by PGPR is far greater than can be explained by elevated CO₂ alone (Blom et al. 2011). Elevated CO₂ levels can increase plant biomass by up to 25 % (Sun et al. 2002; Ward and Strain 1997), but many PGPR easily surpass this as increases in plant biomass of over eightfold have been reported (Park et al. 2015). For WCS417r VOCs, Arabidopsis shoot fresh weight increases of up to fourfold have been reported (Zamioudis et al. 2013), indicating that also WCS417r VOCs have the capacity to stimulate plant growth beyond that caused solely by enhanced CO₂ levels.

To investigate whether plant genotypic traits are involved in the plant's capacity to profit from the plant growth-promoting activity of PGPR, we performed a genome-wide association (GWA) study. GWA mapping is a method initially utilized in human population studies to identify the genetic basis of complex traits (Hirschhorn and Daly 2005). GWA mapping has also been successfully utilized in plant studies (e.g. Aranzana et al. 2005; Atwell

et al. 2010; Bac-Molenaar et al. 2015; Baxter et al. 2010; Chan et al. 2010; Kloth et al. 2012; Li et al. 2010). The underlying rationale of GWA studies is that natural variation in phenotypic traits in a population is caused by genetic differences that via single nucleotide polymorphisms (SNPs) in the genomes of the phenotyped genotypes can be linked to genetic loci and ideally candidate genes. Because plant growth promotion by PGPR is known to display host variation (Smith and Goodman 1999), GWA mapping may be a promising tool to investigate the genetic basis of PGPR-mediated plant growth promotion. In this study, we tested whether *Arabidopsis* has natural genetic variation in the ability to profit from the plant growth-promoting capacity of beneficial rhizosphere bacteria. If so, then this would hold promise for plant breeding strategies in crops that are aimed at maximizing profitable functions from the root microbiome. To test this, we used a global population of 302 *Arabidopsis* accessions that have been genotyped for 214 k SNPs (Baxter et al. 2010; Li et al. 2010) and performed a GWA study for growth-promotion characteristics in response to WCS417r. Virtually all accessions showed enhanced shoot fresh weight and increased lateral root formation upon exposure to WCS417r, while effects on primary root length were more variable. However, the magnitude of the plant growth-promoting effects was significantly different between *Arabidopsis* accessions and was independent of the intrinsic growth rate of the accessions. GWA mapping resulted in the identification of genetic loci associated with the plant's capacity to benefit from growth-promoting effects of WCS417r rhizobacteria.

Materials and methods

Cultivation of bacteria

Pseudomonas simiae WCS417r (formerly known as *Pseudomonas fluorescens* WCS417r; Berendsen et al. 2015) was isolated in the 1980s from the rhizosphere of wheat (Lamers et al. 1988) and served since as a model *Pseudomonas* spp. strain for studying plant growth promotion (Van Peer and Schippers 1989; Zamioudis et al. 2013) and rhizobacteria-induced systemic resistance (Pieterse et al. 1996; Van Peer et al. 1991; Zamioudis and Pieterse 2012). WCS417r was grown for 24 h at 28 °C on King's medium B agar medium supplemented with 50 µg mL⁻¹ rifampicin as described (Van Wees et al. 2013). Subsequently, the WCS417r bacteria were suspended in 10 mL of 10 mM MgSO₄ and centrifuged at 3200g for 5 min. The pellet was washed twice by resuspension in 10 mM MgSO₄ and subsequent centrifuging at 3200g for 5 min after which the

density was adjusted to 2×10^6 colony-forming units (cfu) mL⁻¹ of 10 mM MgSO₄ (OD₆₀₀ = 0.002).

Plant growth conditions and PGPR treatment

A collection of 302 natural accessions of *Arabidopsis thaliana* (Supplemental Table S1) was used to investigate their responsiveness to growth-promoting effects of *P. simiae* WCS417r. Seeds were gas sterilized for 4 h in a desiccator as described (Van Wees et al. 2013). Sterilized seeds were sown on square Petri dishes with agar-solidified Murashige and Skoog (MS) medium supplemented with 0.5 % sucrose as described (Zamioudis et al. 2013). The seeds on MS agar were stratified for 2 days at 4 °C and subsequently incubated in a vertical position in a plant growth chamber at 21 °C with a 16-h day (100 µmol m⁻² s⁻¹) and 8-h night cycle. After 4 days of growth, seedlings of similar size were carefully transferred to 9 cm diameter Petri dishes containing MS agar medium. For the bacterial treatment, 240 µl of a WCS417r suspension containing 2×10^6 cfu mL⁻¹ was applied on the agar medium 5 cm below the roots of the seedlings. For the mock treatment, 240 µl of 10 mM MgSO₄ was applied in a similar manner. The Petri dishes were briefly dried in a laminar flow cabinet, sealed with a lid and two layers of Parafilm, and placed in a vertical position in the growth room.

Phenotyping responsiveness of *Arabidopsis* accessions to WCS417r

Per biological replicate, 4 seedlings of a tester *Arabidopsis* accession were grown on a MS agar plate alongside 4 seedlings of the reference accession Col-0. For each of the 302 tested accessions, 3 replicate plates with WCS417r-treated seedlings were compared to 3 replicate plates with mock-treated seedlings. After 8 days of growth, plates were photographed after which average primary root length per seedling and average number of lateral roots per seedling was determined for each plate. After 10 days of growth, average shoot fresh weight per seedling was determined for each plate of the mock- and WCS417r-treated seedlings. Data were obtained from three replicate plates per accession for both the control and the WCS417r treatment.

GWA mapping, SNP selection, and statistics

A collection of 302 accessions was used to investigate the genetic variation present within *Arabidopsis* (Baxter et al. 2010; Li et al. 2010; Platt et al. 2010). Each of these accessions were genotyped versus the Col-0 accession with

~214 k single nucleotide polymorphism markers (Kim et al. 2007). GWA mapping was performed on the WCS417r-mediated changes in shoot fresh weight (Δ SW), number of lateral roots formed (Δ LRF), and primary root length (Δ PRL). For all traits, means per seedling ($n = 4$) per biological replicate ($n = 3$) were used to calculate the mean per treatment per accession. Values obtained after subtracting the mean value of the mock treatment from that of the WCS417r treatment were used for the GWA analysis (Δ SW, Δ LRF, Δ PRL). The online tool GWAPP (<http://gwas.gmi.oeaw.ac.at/>) was used for GWA mapping as described (Seren et al. 2012). The data sets were normally distributed and the AMM algorithm was employed for mapping. This algorithm closely resembles the commonly used EMMAX (Kang et al. 2010). The GWAPP Geneviewer was used to zoom in on trait-associated SNPs and reveal their position in the genome to pinpoint candidate genes within 10 kb up- and downstream of the identified SNP. For each of the candidate genes, the annotations were retrieved from TAIR10 (arabidopsis.org). With the exception of the GWA mapping, all statistics were carried out with IBM SPSS Statistics 20.

Results

Natural variation in Arabidopsis for responsiveness to PGPR-mediated growth promotion

In soil and plate assays, colonization of the roots of Arabidopsis accession Col-0 by *P. simiae* WCS417r has previously been shown to result in an increase in shoot fresh weight (Pieterse and Van Loon 1999; Zamioudis et al. 2013). On plates, it becomes visible that co-cultivation of Col-0 seedlings with WCS417r not only results in enhanced shoot growth, it also coincides with a faster development of lateral roots (Fig. 1), confirming previous findings (Zamioudis et al. 2013). In order to investigate the natural variation in responsiveness of Arabidopsis to the growth-promoting activity of WCS417r, we tested 302 Arabidopsis accessions for changes in shoot fresh weight and root architecture in response to exposure to WCS417r (Supplemental Table S1). PGPR-induced gain in shoot fresh weight (Δ SW), increase in the number of lateral roots formed (Δ LRF), and effects on primary root length (Δ PRL) were determined after an 8-day (root architecture) or 10-day (shoot fresh weight) period of co-cultivation with WCS417r by subtracting the average values of mock-treated plants from those of the WCS417r treatments. Figure 2a–c show that for all three parameters tested, the Arabidopsis accessions displayed extensive variation in the capacity to respond to WCS417r. Upon exposure to WCS417r, the reference accession Col-0 showed an

average increase in shoot fresh weight of 10 mg, which is a 2.1-fold increase relative to the shoot fresh weight of mock-treated Col-0 plants. Among the 302 accessions, shoot fresh weight in response to WCS417r exposure changed between 0.9 and 4.2 fold, with Col-0 around the 65th percentile (Fig. 2a). For the increase in the number of lateral roots over the 8-day period of exposure to WCS417r, Col-0 performed in the upper range (Fig. 2b). While Col-0 formed on average 7 additional lateral roots in comparison to mock-treated seedlings, other accessions formed up to 14 additional lateral roots or none at all. The primary root length remained virtually unchanged in Col-0 in response to WCS417r, but in the other accessions it ranged from significant decreases to significant increases in length (Fig. 2c). Overall, these results demonstrate that Arabidopsis possesses extensive natural variation in responsiveness to the PGPR WCS417r.

PGPR-mediated growth promotion capacity of Arabidopsis accessions is not linked to their intrinsic growth rates

To investigate whether the observed variation in the WCS417r-induced changes in the shoot and root growth parameters is related to intrinsic growth rates among the accessions, we plotted the absolute values of shoot fresh weight, number of lateral roots, and primary root length for both the mock and the WCS417r treatment and ranked the accessions ascendingly for the mock treatment (Fig. 2d–f). Figure 2d, e show that the values of the mock and the WCS417r treatment have a low correlation coefficient for the parameters shoot fresh weight and number of lateral roots ($R^2 = 0.313$ and 0.025 , respectively), indicating that the magnitude of these WCS417r-induced growth responses are, at the most, weakly related to the intrinsic growth capacity for the parameters tested under these conditions. Hence, faster growing accessions, or accessions that form more lateral roots in the experimental setup are not necessarily stronger responders to WCS417r. For the parameter primary root length, the correlation between mock and WCS417r treatment is stronger ($R^2 = 0.620$; Fig. 2f).

PGPR-mediated enhancement of shoot fresh weight is correlated with root architecture parameters

In order to investigate whether the natural variation in the measured shoot and root growth characteristics of the 302 tested Arabidopsis accessions is correlated, we performed a correlation analysis for both the mock and the WCS417r data sets. Figure 3 shows that in mock-treated plants, shoot fresh weight only weakly correlates with the number of lateral roots ($R^2 = 0.228$; Fig. 3a) and primary root length ($R^2 = 0.301$; Fig. 3b), while primary root length and




Fig. 1 Phenotyping of Arabidopsis accessions for responsiveness to the plant growth-promoting effect of *P. simiae* WCS417r. Shown are photographs of MS agar plates with 4 Col-0 seedlings (left half of the plate) and 4 seedlings of a tester accession (right half of the plate),

8 days after spotting 240 μ l of 10 mM MgSO_4 (Mock) or a WCS417r bacterial suspension (2×10^6 cfu mL^{-1}) on the plate below the plant roots

number of lateral roots do not correlate at all ($R^2 = 0.032$; Fig. 3c). However, in WCS417r-treated plants, the correlation between number of lateral roots and shoot fresh weight ($R^2 = 0.515$; Fig. 3a), primary root length and shoot fresh weight ($R^2 = 0.604$; Fig. 3b), and primary root length and number of lateral roots ($R^2 = 0.385$; Fig. 3c) is markedly higher than in the mock-treated plants. These results indicate that the PGPR-mediated increase in shoot fresh weight is at least partly related to PGPR-mediated changes in root architecture.

GWA mapping reveals loci related to the capacity of Arabidopsis to respond to PGPR

To study the genetic basis of PGPR-mediated plant growth promotion, we performed GWA mapping on the increase in shoot fresh weight data (ΔSFW), the increase in number of lateral roots formed data (ΔLRF), and the change in primary root length data (ΔPRL), using the 214 k SNP set that is commonly used for GWA studies in Arabidopsis (Atwell et al. 2010; Bac-Molenaar et al. 2015; Horton et al. 2012; Kim et al. 2007; Li et al. 2010). Data for ΔSFW , ΔLRF , and ΔPRL (Supplementary Table S1) were uploaded to the online tool GWAPP (<http://gwas.gmi.oeaw.ac.at/>) (Seren et al. 2012) to analyze the quality of the data and perform GWA mapping. The (pseudo-)heritabilities for the recorded phenotypic variation in ΔSFW , ΔLRF , ΔPRL , were 0.28, 0.62, and 0.37, respectively. After confirming that all datasets were normally distributed, they were processed with the AMM algorithm. The GWA mapping results for all three parameters are depicted in Fig. 4. Of the loci with high SNP-trait associations [$-\log_{10}(P) > 5.0$], 5 loci were associated with PGPR-stimulated shoot fresh weight and 5

loci with PGPR-mediated increase in lateral root formation. No loci SNPs were highly associated with PGPR-mediated changes in primary root length (Table 1; Supplemental Table S2). When using a lower threshold level [$-\log_{10}(P) > 4.0$], 18 additional loci were detected for PGPR-induced increase in shoot fresh weight, 20 for the PGPR-mediated increase number of lateral roots, and 20 for the PGPR-induced changes in primary root length (Supplemental Table S2). Although we observed a clear level of correlation between WCS417r-mediated changes in shoot fresh weight and the root architecture parameters (Fig. 3), none of the identified trait-associated SNPs overlapped.

The linkage disequilibrium of the natural population of Arabidopsis accessions used for GWA mapping is on average 10 kb (Kim et al. 2007). Therefore, all genes within 10 kb up- or downstream of the highly [$-\log_{10}(P) > 5.0$] and moderately [$-\log_{10}(P) > 4.0$] associated SNPs were identified. Their identities, gene annotation, and functional information are provided in Supplemental Table S2. In Table 1, the highly-associated SNPs are given along with their closest candidate gene. Among the candidate genes in the closest vicinity of the high SNP-trait associations for the PGPR-mediated changes in shoot and root growth are genes encoding proteins that are associated with important plant growth processes, such as the cytokinin transporter PUP19 (Cedzich et al. 2008), the photosynthesis-related protein Golden2-like 1 (GLK1) (Waters et al. 2009), the Rubisco biogenesis-related DEAD box protein RH39 (Nishimura et al. 2010), carbohydrate transmembrane transport protein At2G18480, Nitrilase-like protein 1 (NLPI) (Kusano et al. 2008), and the photorespiration-related mitochondrial serine hydroxymethyltransferase SHM2 (Engel et al. 2011; Voll et al. 2006)




Fig. 2 Natural variation of 302 *A. thaliana* accessions in their responsiveness to the growth-promoting effect of *P. simiae* WCS417r. **a** Accessions sorted for increase in shoot fresh weight (Δ SFW) in response to exposure to WCS417r for 10 days (Col-0 is indicated with arrow and black dot). **b** Accessions sorted for increase in the number of lateral roots formed (Δ LRF) in response to exposure to seedlings to WCS417r for 8 days. **c** Accessions sorted for difference in the length of the primary root (Δ PRL) after exposure of the seedlings to WCS417r for 8 days. **d** Average shoot fresh weight of mock-treated (grey dots) and WCS417r-treated (green dots) plants. **e** Number of

lateral roots formed in mock-treated and WCS417r-treated (red dots) plants. **f** Primary root length of mock- and WCS417r-treated (purple dots) plants. Each dot represents the average of 3 biological replicates. Values in **a–c** were calculated by subtracting the mean of the mock treatment from the mean of the WCS417r treatment. In each panel, accessions were sorted for increasing values. The order of the accessions differs per panel. The Pearson correlation coefficient (R^2) indicates the degree of correlation between values of control and WCS417r-treated plants


(Table 1). Overall, these results provide insight in the genetic components that may contribute to the plant's capacity to benefit from growth-promoting rhizobacteria in the rhizosphere.

Discussion

Natural accessions of *Arabidopsis* have a high degree of variation in plant development, physiology, and adaptation to their biotic and abiotic environment (Alonso-Blanco et al. 2009). Using a small collection of *Arabidopsis* accessions, we previously explored the natural genetic variation in *Arabidopsis* for its capacity to develop PGPR-mediated ISR and identified quantitative trait loci related to this phenomenon (Ton et al. 1999, 2001). Haney et al. (2015) recently showed that genotypic variations in

Arabidopsis accessions can influence the ability to associate with specific PGPR with consequences for plant fitness. Here we made use of the natural variation within 302 *Arabidopsis* accessions to investigate to what extent the different genotypes differ in their capacity to respond to growth-promoting effects of beneficial rhizosphere bacteria and to gain insight into the genetic basis of the profitable functions by which plants can benefit from their root microbiome. Analysis of the growth promotion parameters shoot fresh weight, lateral root formation, and primary root length in response to WCS417r treatment showed large natural variation among the tested accessions for all three parameters (Fig. 2). The capacity to respond to WCS417r only weakly correlated with the intrinsic growth rate of the accessions. Hence, accessions that gained more shoot fresh weight or produced more lateral roots in the mock-treatment, were not necessarily better responders to the PGPR

Fig. 3 Correlation analysis of shoot fresh weight, number of lateral roots, and primary root length for 302 mock- and *P. simiae* WCS417r-treated *Arabidopsis* accessions. **a** Correlation between number of lateral roots and shoot fresh weight. **b** Correlation between primary root length and shoot fresh weight. **c** Correlation between primary root length and number of lateral roots. The correlation coefficient (R^2) is presented in the graphs and is statistically significant at 0.01 (Pearson's r , 2-tailed) unless stated otherwise (*ns* not significant)



treatment. By contrast, WCS417r-mediated changes in the growth parameters were moderately to highly correlated with each other. Especially the WCS417r-mediated increase in shoot fresh weight showed a high correlation with PGPR-induced changes in the root architecture parameters (Fig. 3), suggesting that changes in root architecture and shoot fresh weight are linked. Previously, Zamioudis et al. (2013) demonstrated that root colonization by WCS417r enhances the auxin response in the root of *Arabidopsis* and stimulates auxin-dependent

developmental programs related to primary root, lateral root, and root hair development. Collectively, these root architectural changes enlarge the capacity of the root system to take up water and nutrients (Lopez-Bucio et al. 2003), which may contribute to the observed correlation between extra number of lateral roots formed and increased shoot fresh weight in WCS417r-treated *Arabidopsis* seedlings. It should, however, be noted that *Arabidopsis* seedlings were grown on plates in which nutrient availability was not limiting, hence we cannot rule out the possibility




Fig. 4 GWA mapping of *P. simiae* WCS417r-mediated effects on shoot fresh weight (Δ SFW), lateral roots formation (Δ LRF), and primary root length (Δ PRL). Manhattan plots of the $-\log_{10}(P)$ SNP

marker–trait associations are shown. From left to right, different colors represent Arabidopsis chromosomes I–V. The dotted grey lines indicate the arbitrary thresholds of $-\log_{10}(P) = 4$ and $-\log_{10}(P) = 5$

that the correlation between root architectural changes and increased shoot fresh weight is caused by another, so far unknown process.

In this study, we chose to treat the Arabidopsis seedlings with PGPR by co-cultivating them on the same MS agar plate while avoiding physical contact (Fig. 1). Previously, we showed that in this setup the effects on root and shoot growth are predominantly mediated by VOCs produced by the bacteria (Zamioudis et al. 2013). We realize that in this way we mainly capture the natural variation in VOCs-mediated plant growth effects in the Arabidopsis population, but because the magnitude of VOCs effects are very similar to those mediated by direct application of PGPR to the root system, we chose for this more simplified system. Also in other systems VOCs produced by PGPR have been shown to have major impacts on plant growth and development (Blom et al. 2011; Farag et al. 2013; Gutierrez-Luna et al. 2010; Park et al. 2015; Ryu et al. 2003). Microbial VOCs have been shown to stimulate auxin responses in the roots (Bailly et al. 2014; Bhattacharyya

et al. 2015; Zamioudis et al. 2013), or are suggested to be taken up and utilized as e.g. sulfur nutrition (Meldau et al. 2013).

Although the PGPR-mediated increase in shoot fresh weight and changes in root architecture seem to be related in the population of natural Arabidopsis accessions tested (Fig. 3), the GWA analysis did not yield common candidate genes (Fig. 4, Table 1; Supplemental Table S2). Among the identified candidate genes for the PGPR-mediated increase in shoot fresh weight were genes associated with important plant growth processes. For instance, the *GLK1* gene encoding transcription factor Golden2-like 1 (GLK1), which together with *GLK2* plays a role in the regulation of photosynthetic genes to optimize photosynthetic capacity in varying environmental and developmental conditions (Waters et al. 2009). Or the gene encoding DEAD box protein RH39, which is required for maturation of chloroplast 23S rRNA and the biogenesis of ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) and other chloroplast-encoded, photosynthetic proteins (Nishimura

Table 1 List of candidate genes from the GWA analysis of the *P. simiae* WCS417r-mediated effects on shoot fresh weight, lateral root formation, and primary root length of 302 natural Arabidopsis accessions

Candidate gene (AGI)	Gene annotation	Gene ontology	Growth parameter	$-\log_{10}P$
At1G47603	Purine permease 19 (PUP19)	Purine transporter; may be involved in the transport of purine and derivatives such as cytokinins across the plasma membrane	Δ SFW	5.22
At2G18480	Major facilitator superfamily protein	Carbohydrate transmembrane transporter activity; plant structure	Δ SFW	5.58
At2G20570	Golden2-like 1 (GLK1)	Transcription factor activity; regulates expression of the photosynthetic apparatus (with GLK2)	Δ SFW	5.46
At4G09730	DEAD-box protein RH39	Protein involved in chloroplast rRNA structure; involved in RUBISCO biogenesis	Δ SFW	5.17
At5G59190	Subtilase family protein	Serine-type endopeptidase activity	Δ SFW	6.42
At1G22080	Cysteine proteinases superfamily protein	Cysteine proteinase activity, unknown protein	Δ LRF	5.28
At1G58330	ZW2, unknown protein	Similarities with RESPONSE TO ABA AND SALT 1 (RAS1)	Δ LRF	6.06
At2G27450	Nitrilase-like protein 1 (NLP1)	N-carbamoylputrescine amidohydrolase; Involved in polyamine biosynthesis, nitrogen compound metabolic process	Δ LRF	5.14
At3G14075	Mono-/di-acylglycerol lipase	Involved in lipid metabolic process	Δ LRF	5.24
At5G26780	Serine hydroxymethyltransferase 2 (SHM2)	Serine hydroxymethyltransferase activity; involved in photorespiration	Δ LRF	6.46

Shown are Arabidopsis AGI numbers of candidate genes that are located in the closest proximity of the identified highly-associated SNPs [$-\log_{10}(P) > 5$] for each of the measured growth parameters, their annotation and ontology in TAIR10. Δ SFW, WCS417r-induced increase in shoot fresh weight; Δ LRF, WCS417r-mediated increase in number of lateral roots formed

et al. 2010). Also the cytokinin transporter gene *PUP19* (Cedzich et al. 2008) is among the candidate genes associated with enhanced shoot fresh weight. Among the candidate genes related to the PGPR-mediated increase in lateral root formation is *SHM2*, which encodes the mitochondrial serine hydroxymethyltransferase SHM2 (Engel et al. 2011). The isozyme SHM1 catalyzes an essential step of the photorespiratory C2 cycle, but because *SHM2* does not complement the mutant phenotype of *shm1* and is relatively highly expressed in roots, its function is currently unknown (Voll et al. 2006). Another candidate gene related to increase in lateral root formation encodes Nitrilase-like protein 1 (NLP1). NLP1 is a N-carbamoylputrescine amidohydrolase involved in biosynthesis of polyamines that play an essential role in plant growth and survival (Kusano et al. 2008). Considering all candidate genes with SNP-trait associations in the GWA analysis, several have known or predicted functions that hold promise for being functional in mediating the plant growth-promoting effect of PGPR. Future analysis of knock-out mutants and ectopic over expressors of these genes and those with a currently unknown function should provide insight into the validity of the candidate genes and their role in the capacity of the plant to benefit from the plant growth-promoting activity of PGPR in their rhizosphere.

In summary, our survey of the natural genetic variation within Arabidopsis for the responsiveness to the plant growth-promoting activity of rhizosphere bacteria revealed that the capacity of plants to profit from their root-associated microbes has a clear genetic component. Hence, identification of genes and loci related to PGPR responsiveness in economically relevant crop species holds promise for breeding strategies that are aimed at introducing traits by which crops maximize profitable functions from their root microbiome.

Acknowledgments The authors thank Joel Boerefijn, Silvia Coolen, Silvia Proietti, and Christos Zamioudis for assisting in the GWA analysis and the growth promotion assays. This work was supported by ERC Advanced Grant 269072 of the European Research Council.

Authors contribution P. C. A. W. designed and performed experiments, analyzed the data, and wrote the manuscript. P. A. H. M. B. designed experiments and wrote the manuscript. C. M. J. P. designed experiments and wrote the manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Alonso-Blanco C, Aarts MGM, Bentsink L, Keurentjes JJB, Reymond M, Vreugdenhil D, Koornneef M (2009) What has natural variation taught us about plant development, physiology, and adaptation? *Plant Cell* 21:1877–1896
- Aranzana MJ, Kim S, Zhao K, Bakker E, Horton M, Jakob K, Lister C, Molitor J, Shindo C, Tang C, Toomajian C, Traw B, Zheng H, Bergelson J, Dean C, Marjoram P, Nordborg M (2005) Genome-wide association mapping in *Arabidopsis* identifies previously known flowering time and pathogen resistance genes. *PLoS Genet* 1:531–539
- Atwell S, Huang YS, Vilhjalmsón BJ, Willems G, Horton M, Li Y, Meng DZ, Platt A, Tarone AM, Hu TT, Jiang R, Muiyati NW, Zhang X, Amer MA, Baxter I, Brachi B, Chory J, Dean C, Debieu M, de Meaux J, Ecker JR, Faure N, Kniskern JM, Jones JDG, Michael T, Nemri A, Roux F, Salt DE, Tang CL, Todesco M, Traw MB, Weigel D, Marjoram P, Borevitz JO, Bergelson J, Nordborg M (2010) Genome-wide association study of 107 phenotypes in *Arabidopsis thaliana* inbred lines. *Nature* 465:627–631
- Bac-Molenaar JA, Fradin EF, Becker FFM, Rienstra JA, van der Schoot J, Vreugdenhil D, Keurentjes JJB (2015) Genome-wide association mapping of fertility reduction upon heat stress reveals developmental stage-specific QTLs in *Arabidopsis thaliana*. *Plant Cell* 27:1857–1874
- Bai Y, Müller DB, Srinivas G, Garrido-Oter R, Potthoff E, Rott M, Dombrowski N, Münch PC, Spaepen S, Remus-Emsermann M, Hüttel B, McHardy AC, Vorholt JA, Schulze-Lefert P (2015) Functional overlap of the *Arabidopsis* leaf and root microbiota. *Nature* 528:364–369
- Bailly A, Groenhagen U, Schulz S, Geisler M, Eberl L, Weisskopf L (2014) The inter-kingdom volatile signal indole promotes root development by interfering with auxin signalling. *Plant J* 80:758–771
- Bakker PAHM, Pieterse CMJ, Van Loon LC (2007) Induced systemic resistance by fluorescent *Pseudomonas* spp. *Phytopathology* 97:239–243
- Baxter I, Brazelton JN, Yu DN, Huang YS, Lahner B, Yakubova E, Li Y, Bergelson J, Borevitz JO, Nordborg M, Vitek O, Salt DE (2010) A coastal cline in sodium accumulation in *Arabidopsis thaliana* is driven by natural variation of the sodium transporter AtHKT1;1. *PLoS Genet* 6:e1001193
- Berendsen RL, Pieterse CMJ, Bakker PAHM (2012) The rhizosphere microbiome and plant health. *Trends Plant Sci* 17:478–486
- Berendsen RL, Van Verk MC, Stringlis IA, Zamioudis C, Tommassen J, Pieterse CMJ, Bakker PAHM (2015) Unearthing the genomes of plant-beneficial *Pseudomonas* model strains WCS358, WCS374 and WCS417. *BMC Genom* 16:539
- Bhattacharyya P, Garladinne M, Lee YH (2015) Volatile indole produced by rhizobacterium *Proteus vulgaris* JBL5202 stimulates growth of *Arabidopsis thaliana* through auxin, cytokinin, and brassinosteroid pathways. *J Plant Growth Regul* 34:158–168
- Blom D, Fabbri C, Connor EC, Schiestl FP, Klauser DR, Boller T, Eberl L, Weisskopf L (2011) Production of plant growth modulating volatiles is widespread among rhizosphere bacteria and strongly depends on culture conditions. *Environ Microbiol* 13:3047–3058
- Browne P, Rice O, Miller SH, Burke J, Dowling DN, Morrissey JP, O’Gara F (2009) Superior inorganic phosphate solubilization is linked to phylogeny within the *Pseudomonas fluorescens* complex. *Appl Soil Ecol* 43:131–138
- Bulgarelli D, Rott M, Schlaeppi K, Loren Ver, van Themaat E, Ahmadinejad N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner FO, Amann R, Eickhorst T, Schulze-Lefert P (2012) Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature* 488:91–95
- Bulgarelli D, Schlaeppi K, Spaepen S, Van Themaat EVL, Schulze-Lefert P (2013) Structure and functions of the bacterial microbiota of plants. *Annu Rev Plant Biol* 64:807–838
- Bulgarelli D, Garrido-Oter R, Munch PC, Weiman A, Droge J, Pan Y, McHardy AC, Schulze-Lefert P (2015) Structure and function of the bacterial root microbiota in wild and domesticated barley. *Cell Host Microbe* 17:392–403
- Cedzich A, Stransky H, Schulz B, Frommer WB (2008) Characterization of cytokinin and adenine transport in *Arabidopsis* cell cultures. *Plant Physiol* 148:1857–1867
- Chan EKF, Rowe HC, Kliebenstein DJ (2010) Understanding the evolution of defense metabolites in *Arabidopsis thaliana* using genome-wide association mapping. *Genetics* 185:991–1007
- Conrath U, Beckers GJM, Langenbach CJG, Jaskiewicz MR (2015) Priming for enhanced defense. *Annu Rev Phytopathol* 53:97–119
- Doornbos RF, Van Loon LC, Bakker PAHM (2012) Impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere. *Rev Agron Sustain Dev* 32:227–243
- Engel N, Ewald R, Gupta KJ, Zrenner R, Hagemann M, Bauwe H (2011) The presequence of *Arabidopsis* serine hydroxymethyltransferase SHM2 selectively prevents import into mesophyll mitochondria. *Plant Physiol* 157:1711–1720
- Farag MA, Zhang H, Ryu CM (2013) Dynamic chemical communication between plants and bacteria through airborne signals: induced resistance by bacterial volatiles. *J Chem Ecol* 39:1007–1018
- Glick BR, Penrose DM, Li JP (1998) A model for the lowering of plant ethylene concentrations by plant growth-promoting bacteria. *J Theor Biol* 190:63–68
- Glick BR, Todorovic B, Czarny J, Cheng ZY, Duan J, McConkey B (2007) Promotion of plant growth by bacterial ACC deaminase. *Crit Rev Plant Sci* 26:227–242
- Gutierrez-Luna FM, Lopez-Bucio J, Altamirano-Hernandez J, Valencia-Cantero E, de la Cruz HR, Macias-Rodriguez L (2010) Plant growth-promoting rhizobacteria modulate root-system architecture in *Arabidopsis thaliana* through volatile organic compound emission. *Symbiosis* 51:75–83
- Haney CH, Samuel BS, Bush J, Ausubel FM (2015) Associations with rhizosphere bacteria can confer an adaptive advantage to plants. *Nat Plants* 1:1–9
- Hardoim PR, Van Overbeek LS, Berg G, Pirttila AM, Compant S, Campisano A, Doring M, Sessitsch A (2015) The hidden world within plants: ecological and evolutionary considerations for defining functioning of microbial endophytes. *Microbiol Mol Biol Rev* 79:293–320
- Hayat R, Ali S, Amara U, Khalid R, Ahmed I (2010) Soil beneficial bacteria and their role in plant growth promotion: a review. *Ann Microbiol* 60:579–598
- Hirschhorn JN, Daly MJ (2005) Genome-wide association studies for common diseases and complex traits. *Nat Rev Genet* 6:95–108
- Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Auton A, Muiyati NW, Platt A, Sperone FG, Vilhjalmsón BJ, Nordborg M, Borevitz JO, Bergelson J (2012) Genome-wide patterns of genetic variation in worldwide *Arabidopsis thaliana* accessions from the RegMap panel. *Nat Genet* 44:212–216
- Junker RR, Tholl D (2013) Volatile organic compound mediated interactions at the plant-microbe interface. *J Chem Ecol* 39:810–825
- Kai M, Piechulla B (2009) Plant growth promotion due to rhizobacterial volatiles—an effect of CO₂? *FEBS Lett* 583:3473–3477
- Kang HM, Sul JH, Service SK, Zaitlen NA, Kong SY, Freimer NB, Sabatti C, Eskin E (2010) Variance component model to account

- for sample structure in genome-wide association studies. *Nat Genet* 42:348–354
- Kim S, Plagnol V, Hu TT, Toomajian C, Clark RM, Ossowski S, Ecker JR, Weigel D, Nordborg M (2007) Recombination and linkage disequilibrium in *Arabidopsis thaliana*. *Nat Genet* 39:1151–1155
- Kloepper JW, Ryu C-M, Zhang SA (2004) Induced systemic resistance and promotion of plant growth by *Bacillus* spp. *Phytopathology* 94:1259–1266
- Kloth KJ, Thoen MPM, Bouwmeester HJ, Jongsma MA, Dicke M (2012) Association mapping of plant resistance to insects. *Trends Plant Sci* 17:311–319
- Kusano T, Berberich T, Tateda C, Takahashi Y (2008) Polyamines: essential factors for growth and survival. *Planta* 228:367–381
- Lamers JG, Schippers B, Geels FP (1988) Soil-borne diseases of wheat in the Netherlands and results of seed bacterization with pseudomonads against *Gaeumannomyces graminis* var. *tritici*, associated with disease resistance. In: Jorna ML, Sloomaker LAJ (eds) *Cereal breeding related to integrated cereal production*. Pudoc, Wageningen, pp 134–139
- Lareen A, Burton F, Schäfer P (2016) Plant root-microbe communication in shaping root microbiomes. *Plant Mol Biol*. doi:10.1007/s11103-015-0417-8
- Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, Malfatti S, del Rio TG, Jones CD, Tringe SG, Dangl JL (2015) Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science* 349:860–864
- Li Y, Huang Y, Bergelson J, Nordborg M, Borevitz JO (2010) Association mapping of local climate-sensitive quantitative trait loci in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 107:21199–21204
- Lopez-Bucio J, Cruz-Ramirez A, Herrera-Estrella L (2003) The role of nutrient availability in regulating root architecture. *Curr Opin Plant Biol* 6:280–287
- Lugtenberg B, Kamilova F (2009) Plant-growth-promoting rhizobacteria. *Annu Rev Microbiol* 63:541–556
- Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J, Engelbrektson A, Kunin V, del Rio TG, Edgar RC, Eickhorst T, Ley RE, Hugenholtz P, Tringe SG, Dangl JL (2012) Defining the core *Arabidopsis thaliana* root microbiome. *Nature* 488:86–90
- Mantelin S, Touraine B (2004) Plant growth-promoting bacteria and nitrate availability: impacts on root development and nitrate uptake. *J Exp Bot* 55:27–34
- Meldau DG, Meldau S, Hoang LH, Underberg S, Wunsche H, Baldwin IT (2013) Dimethyl disulfide produced by the naturally associated bacterium *Bacillus* sp B55 promotes *Nicotiana attenuata* growth by enhancing sulfur nutrition. *Plant Cell* 25:2731–2747
- Mendes R, Kruijt M, De Bruijn I, Dekkers E, Van der Voort M, Schneider JHM, Piceno YM, DeSantis TZ, Andersen GL, Bakker PAHM, Raaijmakers JM (2011) Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science* 332:1097–1100
- Nishimura K, Ashida H, Ogawa T, Yokota A (2010) A DEAD box protein is required for formation of a hidden break in *Arabidopsis* chloroplast 23S rRNA. *Plant J* 63:766–777
- Park YS, Dutta S, Ann M, Raaijmakers JM, Park K (2015) Promotion of plant growth by *Pseudomonas fluorescens* strain SS101 via novel volatile organic compounds. *Biochem Biophys Res Commun* 461:361–365
- Peiffer J, Spor A, Koren O, Jin Z, Tringe S, Dangl J, Buckler E, Ley R (2013) Diversity and heritability of the maize rhizosphere microbiome under field conditions. *Proc Natl Acad Sci USA* 110:6548–6553
- Penrose DM, Moffatt BA, Glick BR (2001) Determination of 1-aminocyclopropane-1-carboxylic acid (ACC) to assess the effects of ACC deaminase-containing bacteria on roots of canola seedlings. *Can J Microbiol* 47:77–80
- Pérez-Jaramillo JE, Mendes R, Raaijmakers JM (2016) Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant Mol Biol*. doi:10.1007/s11103-015-0337-7
- Philippot L, Raaijmakers JM, Lemanceau P, van der Putten WH (2013) Going back to the roots: the microbial ecology of the rhizosphere. *Nat Rev Microbiol* 11:789–799
- Pieterse CMJ, Van Loon LC (1999) Salicylic acid-independent plant defence pathways. *Trends Plant Sci* 4:52–58
- Pieterse CMJ, Van Wees SCM, Hoffland E, Van Pelt JA, Van Loon LC (1996) Systemic resistance in *Arabidopsis* induced by biocontrol bacteria is independent of salicylic acid accumulation and pathogenesis-related gene expression. *Plant Cell* 8:1225–1237
- Pieterse CMJ, Zamioudis C, Berendsen RL, Weller DM, Van Wees SCM, Bakker PAHM (2014) Induced systemic resistance by beneficial microbes. *Annu Rev Phytopathol* 52:347–375
- Pieterse CMJ, de Jonge R, Berendsen RL (2016) The soil-borne supremacy. *Trends Plant Sci*. doi:10.1016/j.tplants.2016.1001.1018
- Platt A, Horton M, Huang YS, Li Y, Anastasio AE, Mulyati NW, Agren J, Bossdorf O, Byers D, Donohue K, Dunning M, Holub EB, Hudson A, Le Corre V, Loudet O, Roux F, Warthmann N, Weigel D, Rivero L, Scholl R, Nordborg M, Bergelson J, Borevitz JO (2010) The scale of population structure in *Arabidopsis thaliana*. *PLoS Genet* 6:e1000843
- Raaijmakers JM, Weller DM (1998) Natural plant protection by 2,4-diacetylphloroglucinol-producing *Pseudomonas* spp. in take-all decline soils. *Mol Plant-Microbe Interact* 11:144–152
- Ryu C-M, Farag MA, Hu CH, Reddy MS, Wei HX, Pare PW, Kloepper JW (2003) Bacterial volatiles promote growth in *Arabidopsis*. *Proc Natl Acad Sci USA* 100:4927–4932
- Seren U, Vilhjalmsón BJ, Horton MW, Meng DZ, Forai P, Huang YS, Long Q, Segura V, Nordborg M (2012) GWAPP: a web application for genome-wide association mapping in *Arabidopsis*. *Plant Cell* 24:4793–4805
- Smith KP, Goodman RM (1999) Host variation for interactions with beneficial plant-associated microbes. *Annu Rev Phytopathol* 37:473–491
- Sun JD, Gibson KM, Kiirats O, Okita TW, Edwards GE (2002) Interactions of nitrate and CO₂ enrichment on growth, carbohydrates, and rubisco in *Arabidopsis* starch mutants. Significance of starch and hexose. *Plant Physiol* 130:1573–1583
- Ton J, Pieterse CMJ, Van Loon LC (1999) Identification of a locus in *Arabidopsis* controlling both the expression of rhizobacteria-mediated induced systemic resistance (ISR) and basal resistance against *Pseudomonas syringae* pv. *tomato*. *Mol. Plant-Microbe Interact* 12:911–918
- Ton J, Davison S, Van Wees SCM, Van Loon LC, Pieterse CMJ (2001) The *Arabidopsis* *ISR1* locus controlling rhizobacteria-mediated induced systemic resistance is involved in ethylene signaling. *Plant Physiol* 125:652–661
- Van Peer R, Schippers B (1989) Plant growth responses to bacterization and rhizosphere microbial development in hydroponic cultures. *Can J Microbiol* 35:456–463
- Van Peer R, Niemann GJ, Schippers B (1991) Induced resistance and phytoalexin accumulation in biological control of fusarium wilt of carnation by *Pseudomonas* sp. strain WCS417r. *Phytopathology* 81:728–734
- Van Rhijn P, Vanderleyden J (1995) The *Rhizobium*-plant symbiosis. *Microbiol Rev* 59:124–142
- Van Wees SCM, Van Pelt JA, Bakker PAHM, Pieterse CMJ (2013) Bioassays for assessing jasmonate-dependent defenses triggered

- by pathogens, herbivorous insects, or beneficial rhizobacteria. *Methods Mol Biol* 1011:35–49
- Voll LM, Jamai A, Renne P, Voll H, McClung CR, Weber APM (2006) The photorespiratory *Arabidopsis shm1* mutant is deficient in *SHM1*. *Plant Physiol* 140:59–66
- Ward JK, Strain BR (1997) Effects of low and elevated CO₂ partial pressure on growth and reproduction of *Arabidopsis thaliana* from different elevations. *Plant Cell Environ* 20:254–260
- Waters MT, Wang P, Korkaric M, Capper RG, Saunders NJ, Langdale JA (2009) GLK transcription factors coordinate expression of the photosynthetic apparatus in *Arabidopsis*. *Plant Cell* 21:1109–1128
- Yang J, Klopper JW, Ryu CM (2009) Rhizosphere bacteria help plants tolerate abiotic stress. *Trends Plant Sci* 14:1–4
- Zamioudis C, Pieterse CMJ (2012) Modulation of host immunity by beneficial microbes. *Mol Plant-Microbe Interact* 25:139–150
- Zamioudis C, Mastranesti P, Dhonukshe P, Blilou I, Pieterse CMJ (2013) Unraveling root developmental programs initiated by beneficial *Pseudomonas* spp. bacteria. *Plant Physiol* 162:304–318
- Zamioudis C, Hanson J, Pieterse CMJ (2014) β -Glucosidase BGLU42 is a MYB72-dependent key regulator of rhizobacteria-induced systemic resistance and modulates iron deficiency responses in *Arabidopsis* roots. *New Phytol* 204:368–379
- Zamioudis C, Korteland J, Van Pelt JA, Van Hamersveld M, Dombrowski N, Bai Y, Hanson J, Van Verk JC, Ling HQ, Schulze-Lefert P, Pieterse CMJ (2015) Rhizobacterial volatiles and photosynthesis-related signals coordinate MYB72 in *Arabidopsis* roots during onset of induced systemic resistance and iron deficiency responses. *Plant J*. 84:308–322
- Zhang H, Sun Y, Xie X, Kim M-S, Dowd SE, Paré PW (2009) A soil bacterium regulates plant acquisition of iron via deficiency-inducible mechanisms. *Plant J* 58:568–577