



BMJ Open Natural population cohort study on long-lived adults: West China longevity and ageing procedure (WCLAP)

Xiaochu Wu ,¹ Tianyao Zhang,² Yan Zhang,¹ Yan She,³ Li Wang,⁴ Yanling Gao,⁵ Yiping Deng,¹ Meijuan Chen,⁴ Yan He,⁵ Xiaoyan Chen ,¹ Qiukui Hao,¹ Jirong Yue,¹ Birong Dong¹

To cite: Wu X, Zhang T, Zhang Y, *et al.* Natural population cohort study on long-lived adults: West China longevity and ageing procedure (WCLAP). *BMJ Open* 2022;**12**:e055407. doi:10.1136/bmjopen-2021-055407

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-055407>).

Received 17 July 2021
Accepted 07 April 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹National Clinical Research Center for Geriatrics (WCHSCU), Sichuan University West China Hospital, Chengdu, Sichuan, China

²The No.1 Affiliated Hospital of Chengdu Medical College, Chengdu, Sichuan, China

³Mei Shan Shi Peng Shan Qu Chineses Medicine Hospital, Meishan City, Sichuan, China

⁴People's Hospital of Leshan, Leshan, Sichuan, China

⁵Sichuan University West China Hospital of Yibin Hospital, Yibin, Sichuan, China

Correspondence to
Birong Dong;
birongdong@163.com

ABSTRACT

Purpose The West China longevity and ageing procedure (WCLAP) cohort study aims to provide guidance for older adults in western China with the aim of improving quality of life, reducing the burden of family care, summarising the characteristics of longevity lifestyles, building a Chinese-longevity-population biobank and exploring the mechanisms underlying population ageing.

Participants Since the establishment of the WCLAP research baseline in 2018, a population of 1537 adults aged 80 years and above, living in the community, have been enrolled in the programme as research participants. Of these, 231 are aged 100 years and above. Participants are followed up every year.

Finding to data WCLAP data are collected in five hospital research subcentres strategically located adjacent to the national 'Longevity Townships' of Chengdu Ziyang, Leshan, Yibin and Pengshan. Data collection included a comprehensive assessment of the participant's health (including physical, psychological, social and common chronic disease assessments), instrumental tests (body composition and muscle percentage) and the collection of biomedical-biobank samples (include blood, urine, faeces, hair and urine).

Future plans Through the annual cohort follow-up, survival-related information is collected at a group level. Analysis of biological samples facilitates biological characterisation at the microscopic level through proteomics, metabolomics, genomics and other techniques. Baseline data, group-level follow-up data and microbiological examination data are integrated together to provide an evaluation tool, exploring sarcopenia, disability, dementia, caregiver burden, ageing biomarkers and other influencing factors.

Trial registration numbers 2018-463; ChiCTR1900020754.

INTRODUCTION

The increasing age of the population has become one of the main factors affecting the quality of life globally. As the elderly population increases, age-related chronic conditions contribute to the global healthcare burden and are anticipated to become the next global public health challenge.¹ There is much research ongoing to determine how best to

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The first multicentre study conducted in the elderly Chinese people living in longevity townships.
- ⇒ Exploring longevity-related behaviours in western China, and characterising chronic diseases incident in this population (sarcopenia, disability, mild cognitive impairment frailty, etc.) and the present prevalence of these diseases.
- ⇒ Establishing a biological sample bank of people living in these areas, and plan to find and verify longevity-related biomarkers by multiomics.
- ⇒ Much of the data are collected through self-report, creating a potential for recall bias. To counter this, data evaluators underwent comprehensive training, but there may still be investigator bias in the evaluation stage.
- ⇒ The target population of this research is elderly adults, so a selective survival bias may exist in the disease epidemiological aspects of the research, as may the loss to follow-up bias.

achieve healthy ageing, improve quality of life in old age and reduce the burden on family caregivers of elderly disabled relatives. As such, a great investment is being driven into the field of ageing health in various countries. In 2018, the population of people aged over 64 surpassed that of children under 5 for the first time. By 2050, it is predicted that nearly 20% of the world's population will be over 65.² Consequently, the workforce in social production is likely to face increased life pressure and social responsibility. It is important to establish a balance between social burden and social productivity, and a good strategy is required to achieve this.

In 2000, China was acknowledged to have an ageing society. A national survey conducted by the National Bureau of Statistics reported China's elderly population (defined as those aged 60 and over) to have reached 249 million at the end of 2018, comprising 17.9% of the total population; with those aged 65 and over

numbering 167 million, accounting for 11.9% of the total population.³ This rate of ageing is projected to increase further to 24.7% in the next 25 years.⁴ In 2010, Chinese scholars reported the total number of people over 60 to be 178 million, accounting for 13.32% of the total population and predicted that by 2030, this will reach 359 million. By 2050, the total number of people over 60 is estimated to reach 448 million, accounting for about 40% of the total population, with a serious impact on quality of life.^{5,6} In the future, with further advancements in technology and human intelligence, China is expected to face more severe consequences from this ageing society.

The elderly population of longevity townships in China has different behavioural characteristics to those in Europe and America, including the variety and structure of dietary intake, economic level, educational background, psychological health (anxiety and depression), the provision of family care and physical activity. The present research aims to focus on the behavioural characteristics associated with high-quality survival and longevity of the elderly population in China's longevity areas. This will be achieved by assessing the lifestyles at a large scale and the biological multiomics at the microscopic scale. This study aims to summarise the characteristics associated with longevity in this special population of elderly Chinese people. By starting from the concept of primary prevention, this study aims to characterize behaviours associated with a long and healthy and longevity life, providing government departments with evidence supporting public health primary prevention strategies. Conversely, through the testing and analysis of biological samples with the help of metabolomics, proteomics and other methods, biomarkers of healthy ageing will be identified to provide a basis for the future development of antiageing drugs and the treatment of age-related diseases.

COHORT DESCRIPTION

Study design and setting

The design and baseline of West China longevity and ageing procedure (WCLAP) were formed in 2017 and 2018, respectively, with the plan of conducting an annual cohort follow-up. WCLAP was designed to be a prospective dynamic cohort study across four medical and health institutions in the national longevity region in western China. From 10 October 2018 to 1 December 2019, WCLAP successively established baseline research subcentres in five longevity regions in western China; Chengdu (The West China Hospital of Sichuan University), Ziyang (Ziyang Zhonghua Hospital), Leshan (The First People's Hospital of Leshan City), Meishan (Pengshan Traditional Chinese Medicine Hospital of Meishan City) and Yibin (The Second People's Hospital of Yibin City). The data collected by each subcentre were collated at the West China Hospital of Sichuan University (WCHSCU). All biological samples were collected

according to standardised operational processes and stored in the biological specimen bank of the West China Hospital of Sichuan University.

Data collection

Data collection comprised (1) the completion of a questionnaire survey, (2) a full physical examination and (3) biological specimen collection and laboratory tests.

The design of the questionnaire was based on the National Health and Ageing Trends Study (NHATS) created by the Johns Hopkins University⁷ and applied to the population of western China. The contents of the questionnaire survey comprised basic demographic characteristics, social activity, family longevity, health and disease status, lifestyle behaviours and evaluation scales. Before the formal use of the questionnaire, our team carried out pretest among the old adult in the community to continuously verify the reliability and validity of the questionnaire. The internal consistency was determined from Cronbach's alpha calculation. Our questionnaire had a Cronbach's alpha of 0.91, and finally was confirmed the final version.

The physical examination included measures of grip strength, pace, body mass index (BMI), blood pressure, skinfold thickness at the triceps, the sit-up test and body composition. Before data collection, all test instruments were calibrated according to manufacturer guidelines.

Biological specimen collection for laboratory testing included blood, urine, stool, saliva and hair. Subjects were provided with a free routine screening of their blood and urine samples within 1 month of their physical examination.

Study subjects

Participants were recruited from the national longevity region in western China. The eligibility criteria were as follows:

1. Participants agreed to participate in the study and signed an informed consent form.
2. Participants were aged 80 years old or older (according to their identification card).
3. Participants had lived in the locality for at least 1 year.
4. Participants were willing and able to cooperate with the evaluator in the local language to complete the 30-min evaluation.

Exclusion criteria:

1. Unwillingness to sign the informed consent form, complete all assessment content independently.
2. The presence of metal implants in the body, such as stents, artificial joints, pins, plates, or cardiac pacemakers.
3. Being under the age of 80 (according to their identification card).
4. Having a life expectancy of under 6 months (diagnosed by medical institution).

Sampling method

A multistage cluster sampling method was used as follows:

1. The study population comprised the population of elderly adults across the varied topography and land-forms of the national longevity area in western China, including hills, basins, mountains and plains.
2. Given the two factors of the geographical environment and the extent of the national longevity areas, four cities in western China were selected as subcentres for the study, namely Meishan, Ziyang, Yibin and Leshan.
3. Given the influence of many factors, including the convenience of transport routes, the willingness of municipal/county/village-level government bodies to cooperate, the sizes of local populations, the distances between collection points and local subcentres and local acceptance of modern medicine, districts were selected adjacent to each subcentre. These were Pengshan District (adjacent to Meishan City), Yanjiang District (adjacent to Ziyang City), Cuiping District (adjacent to Yibin City) and Shizhong District (adjacent to Leshan City).
4. In each of the above districts, a number of towns (10 towns from Pengshan, 17 towns from Yanjiang, 12 towns from Cuiping 12 towns from Shizhong) were randomly selected based on factors such as accessibility, population size and structure.
5. Data collection was conducted at each research site, facilitated by preliminary communication with government departments, subcentres and community organisations at all levels of the project site, to initiate social mobilisation and publicity for the study. Residents of each of the selected towns were invited to voluntarily participate in the study. Our team will provide each participant with free medical examination service (about US\$90) and long-term health follow-up service (free hospital referral service will be provided if necessary) as compensation.

Baseline evaluation

The baseline assessment was completed in 2019, whereby 1546 elderly adults participated. The baseline assessment included the survey, biological sample collection, and physical examination as detailed above.

The questionnaire included basic personal information, information about social and support networks, social microenvironment, religious beliefs, family longevity, chronic disease, health self-assessment, eating habits, household drinking water, smoking status, alcohol/tea intake, physical exercise and daily and leisure activity. This was achieved using validated assessment tools, and where possible those designed specifically for use in elderly populations; Activities of Daily Living (ADL),^{8–10} Instrumental Activities of Daily Living (IADL),^{9,10} Clinical Frailty Scale,^{11,12} Social Support Rating Scale (SSRS),¹³ Pittsburgh Sleep Quality Index (PSQI),^{14,15} the short form of the multinutritional assessment (MNA-SF),^{16,17} mini mental-state exam (MMSE),^{18,19} the Generalised Anxiety Disorder questionnaire (GAD-7)²⁰ and the Geriatric Depression Scale (GDS-15).²¹ The questionnaire contents are detailed in [table 1](#).

Biological samples were collected from participants, and in return they received a free routine blood and urine screening including liver and kidney function tests. All specimens were preprocessed on the day of collection. For convenience, samples were labelled with the participant's basic information, including their sample number, name, gender and age. All specimens except hair specimens were stored temporarily at 4°C before being moved (using dry ice to ensure ultra-low temperature refrigeration throughout the transportation) to the main –80°C storage facility at the Biological Specimen Bank of West China Hospital of Sichuan University. The description of the specimen preprocessing procedure and long-term storage conditions at each subcentre are shown in [table 2](#).

A full physical examination was conducted to characterise the basic physical fitness level and incidence of age-related disease (such as sarcopenia) in the study population. The procedure for the full physical examination is shown in [table 3](#).

Measurement of key variables

Mental state and cognitive assessment

The assessment of the psychological state mainly comprised the evaluation of anxiety and depression in participants using the GAD-7 and GDS-15 assessment tools, respectively. The GAD-7 comprises seven questions to assess the participant's generalised anxiety in the past 2 weeks. The GAD-7 scale is scored from 0 to 21, with a higher score representing a more severe anxiety level. The evaluation standards are 0–4 points signifying a normal level, 5–9 points for mild anxiety, 10–14 points for moderate anxiety and 15–21 points signifying severe anxiety.²² The GDS-15 scale is used in the evaluation of depression in elderly adults over the past week. The scale consists of four questions that are scored directly and 11 reverse-scored questions. Again, a higher score represents the more severe depression. The evaluation standards are 0–4 signifying a normal level, 5–8 for mild depression, 9–11 for moderate depression and 12–15 signifying severe depression.²³

Considering the characteristics of China's elderly population, such as low education level, poor economic level and the need for an assessment that was easy to administer, this study used the MMSE to evaluate cognitive function. The MMSE scale evaluates participants' orientation, memory, recall ability, language ability, attention and calculation ability through 30 questions. The highest achievable score is 30, with scores of 0–26 points representing cognitive dysfunction.²⁴

Sarcopenia

Following the recommendations of the 'Asian Working Group for Sarcopenia: 2019 Consensus Update on Sarcopenia Diagnosis and Treatment', the assessment of sarcopenia in used this study mainly comprised the measurement of grip strength, pace and bioelectrical impedance analysis (BIA; Inbody S10, BioSpace, Seoul, Korea). Previous studies have shown no statistical

**Table 1** The main contents of the questionnaire survey

Questionnaire frame	Content description
Basic personal information	Name, gender, age, ID number, place of birth, length of local residence, local residential address, telephone number (her/himself+relatives), educational background, nationality, language (minority language), main occupation before 60 years old, etc.
Social network and support	Childbirth (number of sons and daughters), family situation, family financial control, family respect, home care, family function, neighbour/friend relationship, support and assistance, marriage/spouse situation, etc.
SRSS scale	Social support evaluation, including friends, neighbours, family, asking for help, talking, etc.
Social microenvironment	Family living area, source of income, annual family income and economic satisfaction.
Religious belief	Religious type, religious activities.
Longevity family survey	Long-lived relatives, family history of genetic disease.
History of chronic disease	Types of chronic diseases, hospitalisation, medical insurance, medical expenses in the past year, satisfaction with medical conditions, timely medical treatment, etc.
Health self-assessment	Limitation of activity, physical state, emotional state, pain, self-feeling, etc.
Eating habits	The number of meals per day, the amount of food, the combination of meals, whether or not breakfast, salt intake, cooking methods, eating speed, taste preference and dietary intake structure/frequency/type.
MNA-SF scale	BMI, psychology, calf circumference, mid-arm circumference diet, etc.
Household drinking water	Type/source of drinking water.
Smoking	Smoking history, age of first smoking, cigarette type, smoking cessation history, smoking status of family members, secondhand smoke inhalation status.
Alcohol intake	Drinking history, first drinking age, drinking frequency and type.
Tea intake	Whether to drink tea now and in the past, age of first drinking, tea type, daily intake (mL).
Physical exercise	Whether physical exercise is performed now or in the past, age at physical exercise start and end.
Daily and leisure activity	Housework, farming, raising poultry, reading, playing mahjong, TV, radio, chat in teahouses, etc.
ADL scale	Daily living ability assessment, including eating, walking, dressing, bathing, etc.
IADL scale	Instrumental assessment of activities of daily living, including cooking, taking medicine, shopping, calling, etc.
Frail scale	Fatigue, endurance, walking, illness, weight loss.
PSQI scale	Sleep assessment.
MMSE scale	Cognitive function assessment, including orientation, memory, attention, calculation, meeting ability, language ability.
GAD-7 scale	Anxiety assessment, including nervousness, worry, irritability, fear, etc.
GDS-15 scale	Depression assessment, including feelings of helplessness, memory, getting help, difficulty, boredom, emptiness, etc.

ADL, Activities of Daily Living; BMI, body mass index; GAD-7, Generalised Anxiety Disorder-7; GDS-15, Geriatric Depression Scale-15; IADL, Instrumental Activities of Daily Living; MMSE, Mini-mental State Examination; MNA-SF, Mini Nutritional Assessment-Short Form; PSQI, Pittsburgh Sleep Quality Index; SSRS, Social Support Rating Scale.

differences between the use of MRI and BIA for the assessment of sarcopenia.²⁵ Muscle mass was defined using the appendicular skeletal muscle mass index (ASMI), calculated similar to BMI, using the formula, appendicular skeletal mass (ASM)/height². A threshold of 7.0 kg/m² in men and 5.7 kg/m² in women was considered to signify low muscle mass. Low grip strength for men and women was defined as 26 and 18kg, respectively.²⁶ The participant's usual walking speed was measured over 4m, with a gait speed of less than 0.8 m/s considered to signify sarcopenia.²⁷

Activities of daily life

Participants' daily living activity assessment used health self-assessment measures (body, emotion and feeling) together with the ADL (level of basic self-care) and IADL (ability to use basic appliances) scales.⁸⁻¹⁰ Based on previous research findings, we additionally used some simple, independently designed health self-evaluation questions covering the previous month of the participant's life condition, including overall self-evaluation, limitations of the ability to undertake activity, the impact of health status on daily life, the impact of emotional

Table 2 Specimen preprocessing procedure and long-term storage conditions

Sample type	Preprocess standard method	Storage condition
Blood	Two tubes (5 mL/tube) of blood per person were collected, which had been centrifuged by 3500 rpm in 15 min, and divided into 8 EP tubes (2 tubes of albuginea and 6 tubes of plasma).	−80°C ultra-low temperature refrigerator.
Urine	On the day of collection, the urine was divided into 2 eppendorf (EP) tubes (2 and 15 mL each) at room temperature.	−80°C ultra-low temperature refrigerator.
Stool	Take a small amount, place in the preservation solution and mix well.	−80°C ultra-low temperature refrigerator.
Saliva	Centrifuge, 2500 rpm, 1 min, the supernatant and residue were placed in two 5 mL pointed EP tubes.	−80°C ultra-low temperature refrigerator.
Hair	Put 10–20 hairs (10 cm/piece) in a sealed bag.	Store at room temperature.

All EP tubes are suitable for ultra-low temperature storage.
EP, eppendorf.

status on daily life, pain status, self-assessed psychological state, the influence of health and mental state on daily social interaction and other issues.

The ADL scale assesses participants' ability to take care of themselves by asking the ability of participants to defecate, urinate, groom, use the toilet, eat, move, dress, use

Table 3 Physical examination procedure

Item	Description
Physical condition	
Height	Stand up straight, measure two times in a row and ask about last year's weight.
Weight	Take off the coat and other heavy clothes, repeat the measurement twice.
Upper arm circumference	Measure the circumference of the upper arm at the midpoint of the line between the shoulder and the elbow, repeat the measurement twice.
Triceps skinfold thickness	Use a cortical thickness gauge to measure the fat thickness at the midpoint of the shoulder and elbow joint, repeat the measurement twice.
Calf circumference	Keep leg upright, measure the circumference of the thickest part of the calf, repeat the measurement twice.
Waist circumference	Measure the waist circumference at a point 2 cm above the belly button, repeat the measurement twice.
Hip circumference	The circumference between the symphysis pubis and the most convex part of the back gluteus maximus, repeat the measurement twice.
Knee height	The length between the knee joint and the heel, repeat the measurement twice.
Finger distance	The length between the index fingers of both hands, repeat the measurement twice.
Disease-related condition	
Blood pressure and pulse	After sitting and resting for 2–5 min, start the measurement, record the diastolic and systolic blood pressure (electronic sphygmomanometer), and repeat the measurement twice.
ECG bioelectrical impedance analysis (BIA)	After confirming that there is no metal in the participant's body, use INBODY S10 (BioSpace, Seoul, Korea) to measure the BIA data, enter the participant's information into the host, adopt a sitting posture and connect all limbs with electrodes. No talking is allowed during the measurement.
Gait speed (s/m)	Measure the pace of the participants and calculate the pace (seconds/metre) at the start and end times of 3 segments of 4 m.
Grip strength	Ask the participant's habit of using hands and test the grip strength of both hands twice.
Balance test	Participants were tested for balance in three stances, including feet side by side, staggered feet, and front and rear feet. Each stance held for 10 s was deemed qualified.
Blood sugar	Fingertip blood sugar test.
Sit-up ability tests	Participants hold their shoulders, complete 1 and 5 consecutive sit-up tests, respectively, and record the completion status (whether it can be completed and when it is completed).



the stairs and bathe. The full scale of the scale comprises 100 points, with 100 points representing perfect ability with no need to rely on others; 61–99 points representing a mild dysfunction, but basically having the ability to take care of themselves; 41–60 points representing moderate dysfunction, and needing some help; 21–40 points representing severe dysfunction and needing to be dependent on others and less than 20 points representing complete dependence on others.

The IADL scale comprehensively evaluates the participants' ability to use basic appliances. The evaluation content includes cooking, housework, taking medicine, walking, shopping, financial management, using the telephone and washing clothes. The scale has a maximum score of 16, with a score of less than eight indicating impaired instrumental life ability.⁹

Sleep quality

The PSQI scale is used to evaluate the sleep of participants through sleep quality, the time taken to fall asleep, duration of sleep, sleep efficiency, sleep disorders, use of hypnotic drugs and daytime dysfunction. The scale has a maximum score of 21 points, with a score of 0–5 representing the best sleep quality; 6–10 good sleep quality; 11–15 average sleep quality; and 16–21 poor sleep quality. In addition, we also asked questions about 'siesta'/napping and the Sleep Self-Assessment.²⁸

Medical history and social support

Participants were asked to provide a full medical history of chronic diseases (diagnosed by professional medical institutions), including cardiovascular and cerebrovascular diseases, respiratory diseases, nervous system diseases, hearing disorders, bone and joint diseases, liver and kidney diseases and incontinence. The assessor asked about the symptoms experienced, including whether the participant was ill, the number of years they experienced the illness and the treatment undertaken.

The SSRS was used to evaluate the social support of the participants through factors such as the social interaction between the participants and friends, neighbours, family members and social groups.²⁹ The maximum score of this scale is 40 points, with a higher score representing better social support. A score of 30–40 was considered to represent a good degree of social support; a score of 20–29 a general degree of social support and a score of less than 20 a lower degree of social support. In addition, we investigated the family microenvironment, including the housing structure, family economic control/discourse power, family respect, family decision-making, family support, family conflict response, and spousal relationship.

Nutrition and behaviour

The MNA-SF was used to conduct a nutritional assessment.³⁰ This assessment tool consists of two sections, with all participants completing the first section and participants only progressing to the second section if they scored

less than 11 points on the first, indicating possible malnutrition. A combined score (scores from both sections added together) of greater than 24 points was interpreted as a good nutritional status; 17–23.5 points was interpreted as a risk of malnutrition and a score of less than 17 indicated the definite presence of malnutrition.

In addition, based on the characteristics of the traditional Chinese diet and previous research findings, we independently designed a series of survey questions related to eating habits, diet types and the frequency with which they ate certain foods. The survey of eating habits included the number of meals per day, meal times, the amount eaten at each meal, the types of food eaten at each meal (meat and vegetables), breakfast, flavourings, cooking methods and speed of eating. Dietary content was assessed by asking how frequently (ie, daily, 3–5 times a week, 1–2 times a week, 1–3 times a month or never) participants consumed grains, vegetables, fruits, meat, eggs, seafood, dairy products, vegetable oil, animal oil, nuts, candy, cakes, etc.

Participant behaviour was captured by asking about smoking, alcohol intake, tea intake, physical labour and leisure activities. Participants were asked whether they smoke (at least one cigarette per day for more than 6 months), for how many years, what type of cigarettes, how many cigarettes per day, whether they had attempted to quit smoking, whether they experienced secondhand smoke inhalation and how often. Similarly, alcohol consumption was defined as drinking alcohol at least once a week. Participants were asked at what age they started drinking alcohol, their drinking frequency, type of alcohol consumed and the volume of a single drink. Participants were asked if they drank tea frequently (more than three times a week), whether their habits had been different in the past, the age they started drinking tea, the type of tea, and the amount of tea consumed daily.

Manual labour was assessed by asking participants whether they undertake moderate or severe manual labour, the number of days they did this per week, and the number of hours per day. Leisure activities were captured including growing vegetables, raising poultry, raising pets, reading books and newspapers, educational activities (Mahjong, etc.), watching television or listening to the radio, socialising in tea houses, etc.

Follow-up

The follow-up tool was designed taking into account the characteristics/results of the baseline data (October 2018–December 2019), key issues (frailty, cognition, disability, sarcopenia, psychology, etc.) and follow-up methods (phone or face-to-face follow-up). In 2020, the follow-up work needed to incorporate further external factors such as COVID-19, and the government's policy response. The initial plan was to conduct a telephone follow-up with all participants recruited at baseline. Later, depending on the COVID-19 situation in China, it may be possible to conduct an in-person follow-up.

Table 4 Participant characteristics

Characteristics	All (n=1537)	Men (n=643)	Women (n=894)	P value	Missing number	
Age(years), mean(±SD)	88.7 (7.36)	87.7 (6.39)	89.40 (7.91)	<0.001	1	
80–85	682 (44.4)	295 (46.0)	387 (43.3)	<0.001	1	
86–90	293 (19.1)	142 (22.1)	151 (16.9)			
91–99	328 (21.4)	150 (23.4)	178 (19.9)			
≥100	233 (15.2)	55 (8.6)	178 (19.9)			
Ethnics, n (%)					3	
Han	1530 (99.7)	639 (99.7)	891 (99.8)			
Others	4 (0.3)	2 (0.4)	2 (0.2)			
Educational level, n (%)				<0.001	6	
No formal education	915 (59.8)	232 (36.3)	683 (76.6)			
Elementary school	432 (28.2)	281 (44.0)	151 (16.9)			
Middle school	98 (6.4)	66 (10.3)	32 (3.6)			
Technical secondary school	35 (2.3)	23 (3.6)	12 (1.3)			
High school and above	51 (3.3)	37 (5.8)	14 (1.6)			
Status of spouse, n (%)				<0.001	20	
Alive	512 (33.8)	352 (55.3)	160 (18.2)			
Divorced	5 (0.3)	2 (0.3)	3 (0.3)			
Widowed	1000 (65.9)	282 (44.3)	718 (81.5)			
Annual household income per capita, n (%)					8	
<1000 rmb/year	77 (5.0)	32 (5.0)	45 (5.1)			
1000–3000 rmb/year	299 (19.6)	121 (18.9)	178 (20.0)			
3001–6000 rmb/year	257 (16.8)	101 (15.8)	156 (17.5)			
6001–8000 rmb/year	104 (6.8)	41 (6.4)	63 (7.1)			
8001–10000 rmb/year	75 (4.9)	29 (4.5)	46 (5.2)			
>10000 rmb/year	582 (38.1)	269 (42.1)	313 (35.2)			
Unknown	135 (8.8)	46 (7.2)	89 (10.0)			
Anthropometric measures						
Height (cm)	148.4 (10.7)	156.1 (7.7)	142.8 (8.9)	<0.001	12	
Weight (kg)	49.4 (10.4)	54.3 (9.8)	45.9 (9.4)	<0.001	12	
BMI (kg/m ²)	22.3 (3.7)	22.2 (3.3)	22.4 (4.0)	<0.001	15	
Underweight (<18.5)	201 (13.2)	62 (9.7)	139 (15.7)			
Normal (18.5–24.0)	863 (56.7)	391 (61.2)	472 (53.4)			
Overweight (24.0–27.9)	356 (23.4)	158 (24.7)	198 (22.4)			
Obese (>28.0)	103 (6.8)	28 (4.4)	75 (8.5)			
Grip strength	17.5 (11.2)	22.2 (11.0)	14.0 (10.0)	<0.001	47	
4-m gait speed	0.58 (0.23)	0.63 (0.22)	0.58 (0.23)	<0.001	103	
Life-styles						
Drinking tea history	Yes	435 (28.4)	329 (51.4)	106 (11.9)	<0.001	6
	No	1096 (71.6)	311 (48.6)	785 (88.1)		
Drinking alcohol history	Yes	476 (31.1)	196 (46.2)	180 (20.2)	<0.001	6
	No	1055 (68.9)	344 (53.8)	711 (79.8)		
Smoking history	Yes	429 (28.1)	332 (52.1)	97 (10.9)	<0.001	9
	No	1099 (71.9)	305 (47.9)	794 (89.1)		
Scale evaluation						
ADL scale result	93.3 (14.3)	95.3 (12.0)	91.9 (15.6)	<0.001	17	

Continued

**Table 4** Continued

Characteristics	All (n=1537)	Men (n=643)	Women (n=894)	P value	Missing number
Good (100 points)	934 (61.7)	443 (70.0)	491 (55.7)		
Mild dysfunction (>60)	514 (33.9)	170 (26.9)	344 (39.0)		
Moderate dysfunction (41–60)	36 (2.4)	12 (1.9)	24 (2.7)		
Severe dysfunction (21–40)	17 (1.1)	4 (0.6)	13 (1.5)		
Completely disabled (<20)	14 (0.9)	4 (0.6)	10 (1.1)		
IADL scale result	11.4 (4.6)	12.1 (4.2)	10.9 (4.9)	<0.001	16
Good function (≥8)	1214 (79.8)	542 (85.6)	672 (75.7)		
Impaired function (<20)	307 (20.2)	91 (14.4)	216 (24.3)		
MMSE scale result	16.3 (8.1)	19.0 (7.7)	14.3 (7.7)	<0.001	174
Normal	151 (11.1)	100 (17.3)	51 (6.5)		
Cognitive impairment	1212 (88.9)	478 (82.7)	734 (93.5)		

ADL, Activities of Daily Living; IADL, Individual Activities of Daily Living.

The telephone follow-up comprised a phone call of no longer than 10 min. For any participants who had died, the investigator recorded the date of death, cause of death, and other information and comforted the family members of the deceased. For participants with the wrong phone number, the investigator tried to contact their family members or community staff to minimise the rate of loss to follow-up. For participants who were temporarily unavailable/busy, the investigator reattempted contact on at least five occasions at different times of the working day/week/weekend. Telephone follow-up investigators were mostly medical college students, and all received comprehensive training.

Following completion of the telephone follow-up on all participants recruited at baseline, the attrition rate of the study was assessed. Then, according to the sample size requirements of the prospective dynamic cohort study, new participants were recruited to join the cohort, in line with the recruitment criteria outlined above.

Data quality control and management

Due to the participants being over 80 years old, and the locations of longevity areas being typically in marginal mountainous areas, this study chose to conduct data capture using paper questionnaires. Paper questionnaires were verified on the day of data collection; 30%–50% of the questionnaires were randomly selected for verification by two independent investigators on the same day. Any missing or ambiguous responses were confirmed by the investigator telephoning the participant.

The database was established using Epidata V.3.0. Independent double entry was used, and the two independent databases were compared using the ‘consistency check’ function of the software. Any inconsistencies between the two databases were modified on a case-by-case basis until the two databases were completely consistent.

All biological specimens were preprocessed and marked on the day of collection (adding marking information, such as name, gender, age and code, based on

the original barcode and QR code of each cryopreservation tube), and stored in the ultra-low temperature refrigerator medium (−80°C). The transport conditions of biological specimens in intracity were with the help of a transfer box, which was kept at 4°C to avoid haemolysis of biological specimens; for the transfer of biological specimens between cities, dry ice was used to maintain an ultra-low temperature environment and the entire process of transport had a temperature control record; all biological specimens according to ethical requirements were stored in the biological specimen bank of West China Hospital of Sichuan University for long-term preservation. At the same time, after each transfer was completed, whether it was intra-city transfer or inter-city transfer, random inspections (1%–3%) of the location information of cryopreservation tubes were required to ensure that the storage location of biological specimens will not change due to the transfer work (change, move and loss of storage location information, etc.).

All evaluation results would be fed back to the person or family members within 1 month, laying a good foundation of trust for the next follow-up work.

The principal investigator has overall responsibility for data management, including data storage, application and use. The data management plan follows guidance on medical ethics, fairness and bias. Therefore, before using the data, the data were deidentified by removing sensitive information such as the fields of participant name, gender, age, identification number and home address.

Patient and public involvement

No patient involved.

Statistical strategy and findings to date

Data analyses were conducted using SPSS V.22.0 and R V.3.6.1. Descriptive statistical analyses were conducted on longevity-related characteristics and medical examination data to provide percentages, means and SD. For interrogation of independent samples by region, gender, ages

and disease, χ^2 tests were used. If the conditions for using χ^2 tests were not met, the Fisher test was used instead. For variables such as scores on any of the psychometric or behavioural scales, the rank-sum test was used.

Participant characteristics are displayed in [table 4](#).

Collaboration

Welcome geriatric medicine and longevity-related researchers through our centre website (<http://www.wchscu.cn/scientific/clinical/platform/55440.html>) for more data information (database name: a natural cohort study of the old adult), if researchers have any requirements, you can also contact us by e-mail (hxncrcg@163.com) for more details and cooperation.

Future details

It is planned to continue to carry out annual participant follow-up and collection of biological samples. Participants in the existing cohort will be followed up continuously, and new participants will be recruited into the prospective dynamic cohort study based on maintaining a constant sample size (ie, to compensate for dropout). Participants who did not have the opportunity to contribute biological samples at the baseline time point will be followed up as soon as possible, and annual sample collections will occur as appropriate.

Global ageing problem has been coming. How to improve the longevity and quality of life is a great challenge for modern medicine, biology and sociology. Centenarians are an important model to study longevity and 'healthy ageing'. The project will conduct experiments on gene and protein levels of centenarians from blood samples to explore the mechanism of longevity and ageing, and provide a theoretical basis for the prevention and treatment of ageing-related diseases, which would be reasonable for extension of life and realisation of healthy ageing.

Frailty syndrome is a systemic change, which companions with multisystem dysfunction, especially the decline of capacity of physiological reserve in neuromuscular, metabolic and immune systems in the elderly. Frailty could reduce the ability to fight stress and significantly increasing the risk of adverse events in the elderly. In order to evaluate the diagnostic accuracy of frailness, the selection, detection, validation and clinical application of biomarkers representing different stages of frailness based on biological theory were established from the perspective of genomics and epigenetics.

We are going to further explore the biological mechanisms of elderly health ageing, reveal changes in the longevous elderly, discoversome novelimportant longevity-related genes and their related functions and signal path, from genomics, apparent genome, transcriptome, proteomics, metabolomics and microbic genomics level. Furthermore, confirmatory researches need to be extensive based on biomarkers associated with longevity among population. In combination with animal models,

antiageing drugs, cells and other therapeutic strategies would be discovered.

The findings of these complex analyses will be exploited for any factors that can be translated into community benefit. With reference to China's ageing society, non-pharmaceutical health intervention and promotion programmes will be explored and formulated with the intention of meeting the needs of the current and future Chinese population. We will propose primary prevention measures suitable for community health promotion, which will be beneficial for a healthy lifestyle, incorporating what we have learnt about the behaviours of the elderly Chinese inhabitants of longevity townships.

Acknowledgements We would like to thank Pengshan District Hospital of Traditional Chinese Medicine, Yibin Second People's Hospital, Leshan People's Hospital and Ziyang Zhonghua Hospital for their strong support and assistance in this study. At the same time, thanks for the help from Ru Gao, Hanlin Liu, Lingling Shen, Yi Lv, Aiguo Yang, Shuli Jia, Meiling Ge.

Contributors Writing: XC Wu and TY Zhang; study design: QK Hao, JR Yue and BR Dong; project coordination: XC Wu, YD, TZ and BD; data acquisition: XC Wu, YZ, YS, LW, YG, MC and YH; data quality control: XC Wu and BR Dong; data management: XC Wu and BR Dong ; guarantor: BR Dong.

Disclaimer Our sources of funding had no role in the design of cohort profile, and will not be any impact on data collection, analysis, writing and decision to submit or publish the research results.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting or dissemination plans of this research.

Patient consent for publication Consent obtained directly from patient(s).

Ethics approval This study involves human participants. The WCLAP was approved by the Medical Ethics Committee of the West China Hospital of Sichuan University (reference number: 2018-463), and registered with the China Clinical Trial Registration Centre (registration number: ChiCTR1900020754). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Welcome geriatric medicine and longevity-related researchers through our centre website (<http://www.wchscu.cn/scientific/clinical/platform/55440.html>) for more data information (database name: a natural cohort study of the old adult), if researchers have any requirements, you can also contact us by e-mail (hxncrcg@163.com) for more details and cooperation.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Xiaochu Wu <http://orcid.org/0000-0001-7055-9449>

Xiaoyan Chen <http://orcid.org/0000-0001-8109-4691>

REFERENCES

- 1 Rudnicka E, Napierala P, Podfigurna A, *et al*. The world Health organization (who) approach to healthy ageing. *Maturitas* 2020;139:6–11.
- 2 Dzau VJ, Finkelman EM, Balatbat CA, *et al*. Achieving healthy human longevity: a global grand challenge. *Sci Transl Med* 2020;12. doi:10.1126/scitranslmed.abd3816. [Epub ahead of print: 21 10 2020].
- 3 He X, Song M, Qu J, *et al*. Basic and translational aging research in China: present and future. *Protein Cell* 2019;10:476–84.

- 4 Chen R, Xu P, Song P, *et al.* China has faster PACE than Japan in population aging in next 25 years. *Biosci Trends* 2019;13:287–91.
- 5 Chen Z, Jiang X, Shi G, *et al.* Social frailty and longitudinal risk of depressive symptoms in a Chinese population: the Rugao longevity and aging study. *Psychogeriatrics* 2021;21:483–90.
- 6 Zhang YS, Strauss JA, Hu P, *et al.* Links between mortality and socioeconomic characteristics, disease burden, and biological and physical functioning in the aging Chinese population. *J Gerontol B Psychol Sci Soc Sci* 2021.
- 7 Freedman VA, Kasper JD. Cohort profile: the National health and aging trends study (NHATS). *Int J Epidemiol* 2019;48:1044–5.
- 8 Friedman B, Li Y. Using Rasch analysis to evaluate accuracy of individual activities of daily living (ADL) and instrumental activities of daily living (IADL) for disability measurement. *J Appl Meas* 2015;16:315–35.
- 9 Pashmdarfard M, Azad A. Assessment tools to evaluate activities of daily living (ADL) and instrumental activities of daily living (IADL) in older adults: a systematic review. *Med J Islam Repub Iran* 2020;34:33.
- 10 Henskens M, Nauta IM, Drost KT, *et al.* The effects of movement stimulation on activities of daily living performance and quality of life in nursing home residents with dementia: a randomized controlled trial. *Clin Interv Aging* 2018;13:805–17.
- 11 Xu L, Zhang J, Shen S, *et al.* Clinical frailty scale and biomarkers for assessing frailty in elder inpatients in China. *J Nutr Health Aging* 2021;25:77–83.
- 12 Afilalo J. The clinical frailty scale: upgrade your eyeball test. *Circulation* 2017;135:2025–7.
- 13 Lu S, Wu Y, Mao Z, *et al.* Association of formal and informal social support with health-related quality of life among Chinese rural elders. *Int J Environ Res Public Health* 2020;17. doi:10.3390/ijerph17041351. [Epub ahead of print: 19 02 2020].
- 14 Han Q, Liu B, Lin S, *et al.* Pittsburgh sleep quality index score predicts all-cause mortality in Chinese dialysis patients. *Int Urol Nephrol* 2021;53:2369–76.
- 15 Purani H, Friedrichsen S, Allen AM. Sleep quality in cigarette smokers: associations with smoking-related outcomes and exercise. *Addict Behav* 2019;90:71–6.
- 16 Kaiser MJ, Bauer JM, Ramsch C, *et al.* Validation of the mini nutritional assessment short-form (MNA-SF): a practical tool for identification of nutritional status. *J Nutr Health Aging* 2009;13:782–8.
- 17 Dent E, Hoogendijk EO, Visvanathan R, *et al.* Malnutrition screening and assessment in hospitalised older people: a review. *J Nutr Health Aging* 2019;23:431–41.
- 18 Trivedi D. Cochrane review summary: mini-mental state examination (MMSE) for the detection of dementia in clinically unevaluated people aged 65 and over in community and primary care populations. *Prim Health Care Res Dev* 2017;18:527–8.
- 19 Pinto TCC, Machado L, Bulgacov TM, *et al.* Is the Montreal cognitive assessment (MoCA) screening superior to the Mini-Mental state examination (MMSE) in the detection of mild cognitive impairment (MCI) and Alzheimer's disease (AD) in the elderly? *Int Psychogeriatr* 2019;31:491–504.
- 20 Creese B, Khan Z, Henley W, *et al.* Loneliness, physical activity, and mental health during COVID-19: a longitudinal analysis of depression and anxiety in adults over the age of 50 between 2015 and 2020. *Int Psychogeriatr* 2021;33:505–14.
- 21 Shin C, Park MH, Lee S-H, *et al.* Usefulness of the 15-item geriatric depression scale (GDS-15) for classifying minor and major depressive disorders among community-dwelling elders. *J Affect Disord* 2019;259:370–5.
- 22 Vasiliadis H-M, Chudzinski V, Gontijo-Guerra S, *et al.* Screening instruments for a population of older adults: the 10-item Kessler psychological distress scale (K10) and the 7-item generalized anxiety disorder scale (GAD-7). *Psychiatry Res* 2015;228:89–94.
- 23 Dias FLdaC, Teixeira AL, Guimarães HC, *et al.* Accuracy of the 15-item geriatric depression scale (GDS-15) in a community-dwelling oldest-old sample: the Pietà study. *Trends Psychiatry Psychother* 2017;39:276–9.
- 24 Kvitting AS, Fällman K, Wressle E, *et al.* Age-Normative MMSE data for older persons aged 85 to 93 in a longitudinal Swedish cohort. *J Am Geriatr Soc* 2019;67:534–8.
- 25 Chien M-Y, Huang T-Y, Wu Y-T. Prevalence of sarcopenia estimated using a bioelectrical impedance analysis prediction equation in community-dwelling elderly people in Taiwan. *J Am Geriatr Soc* 2008;56:1710–5.
- 26 Landi F, Calvani R, Tosato M, *et al.* Age-Related variations of muscle mass, strength, and physical performance in Community-Dwellers: results from the Milan EXPO survey. *J Am Med Dir Assoc* 2017;18:88.e17–88.e24.
- 27 Cruz-Jentoft AJ, Landi F, Schneider SM, *et al.* Prevalence of and interventions for sarcopenia in ageing adults: a systematic review. Report of the International sarcopenia initiative (EWGSOP and IWGS). *Age Ageing* 2014;43:748–59.
- 28 Zhang H, Li Y, Zhao X, *et al.* The association between PSQI score and hypertension in a Chinese rural population: the Henan rural cohort study. *Sleep Med* 2019;58: :27–34.
- 29 Xiao J, Huang B, Shen H, *et al.* Association between social support and health-related quality of life among Chinese seafarers: A cross-sectional study. *PLoS One* 2017;12:e0187275.
- 30 Montejano Lozoya R, Martínez-Alzamora N, Clemente Marín G, *et al.* Predictive ability of the mini nutritional assessment short form (MNA-SF) in a free-living elderly population: a cross-sectional study. *PeerJ* 2017;5:e3345.