

Nature and Extent of the Deep Biosphere

Frederick S. Colwell

*College of Earth, Ocean, and Atmospheric Sciences
Oregon State University
Corvallis, Oregon 97331-5503, U.S.A.
rcolwell@coas.oregonstate.edu*

Steven D'Hondt

*Graduate School of Oceanography
University of Rhode Island
Narragansett, Rhode Island 02882, U.S.A.
dhondt@gso.uri.edu*

INTRODUCTION

In the last three decades we have learned a great deal about microbes in subsurface environments. Once, these habitats were rarely examined, perhaps because so much of the life that we are concerned with exists at the surface and seems to pace its metabolic and evolutionary rhythms with the overt planetary, solar, and lunar cycles that dictate our own lives. And it certainly remains easier to identify with living beings that are in our midst, most obviously struggling with us or against us for survival over time scales that are easiest to track using diurnal, monthly or annual periods. Yet, research efforts are drawn again and again to the subsurface to consider life there. No doubt this has been due to our parochial interests in the resources that exist there (the water, minerals, and energy) that our society continues to require and that in some cases are created or modified by microbes. However, we also continue to be intrigued by the scientific curiosities that might only be solved by going underground and examining life where it does and does not exist.

But really, is life underground just a peculiarity of most life on the planet and only a recently discovered figment of life? Or is it actually a more prominent and fundamental, if unseen, theme for life on our planet? Our primary purpose in this chapter is to provide an incremental assembly of knowledge of subsurface life with the aim of moving us towards a more complete conceptual model of deep life on the planet. We aim to merge the consideration of the subseafloor and the continental subsurface because it is only through such a unified treatment that we can reach a comprehensive view of this underground life. We also provide some thoughts on a way forward with what we consider to be interesting new research areas, along with the methods by which they might be addressed as we seek new knowledge about life in this Stygian realm.

EARLY STUDIES AND COMPREHENSIVE REVIEWS

The earliest studies attempted to tease out the nature of subsurface life, albeit in qualitative ways. Edson Sunderland Bastin and his team, intrigued by the formation of sulfides in oil wells, examined oil and water pumped from wells in eastern Illinois (Bastin et al. 1926). Using classical methods of cultivating and detecting sulfate-reducing microbes, they established that

cells were present in fluids produced from oil reservoirs even if their original provenance could not be ascertained. These land-based studies only preceded Claude ZoBell's first look into the seafloor by a few years. Even with short core lengths, ZoBell and Anderson discerned a trend towards increasing numbers of anaerobes relative to aerobes as the samples came from deeper and deeper in the top two meters of sediment (ZoBell and Anderson 1936). Of course, much detail has been laid atop these early observations but this glimpse of the effect of plummeting redox still holds as a principle of subsurface studies.

A number of useful reviews of progress in the science of subsurface microbiology should be consulted to grasp the origin of this field and some of the important directions. Early work was described in a special issue of the journal *Microbial Ecology* (cf., Balkwill et al. 1988 and other papers in this volume) and an update to these findings was reported with a distinctive emphasis on continental habitats in the deep subsurface (Fredrickson and Onstott 1996). Shortly thereafter, two books similarly focused on terrestrial systems (Amy and Haldeman 1997; Fredrickson and Fletcher 2001). Perhaps prompted by the estimates presented by Whitman and colleagues (Whitman et al. 1998) and the first microbiological findings reported by scientists associated with the Ocean Drilling Program (ODP; cf., Whelan et al. 1986; Parkes et al. 1994) the last ten years has seen an upswing in papers published on the microbiology of subsurface of marine systems (D'Hondt et al. 2002b; Smith and D'Hondt 2006). Recent overview papers consider past research progress and directions for the future and begin to express the need to bind together our consideration of deep terrestrial and deep seafloor life in a more inclusive light (Fredrickson and Balkwill 2006; Onstott et al. 2009a; Schrenk et al. 2010, 2013; Edwards et al. 2012; Anderson et al. 2013; Meersman et al. 2013). These works were all paralleled by the five editions of the book *Geomicrobiology* (cf., Ehrlich 1990), now completed by the publication of H.L. Ehrlich's memoir (Ehrlich 2012).


WHERE WE ARE NOW – THE TERROIR OF SUBSURFACE LIFE

And what do we know now? In a broad sense we might start by examining the range of chemical and physical processes that sustain gradients in Earth and therefore may allow the establishment of microbial communities at depth. Regardless of where it exists, metabolically active life—as we know it—seems to require a gradient (Kappler et al. 2005). Living cells may transit places where there is no gradient and manage to survive if these cells can make the passage through harsh conditions. Cells that are locked in a permanent deep freeze (Vorobyova et al. 1997), although some manage activity under frigid conditions (Bakermans et al. 2003), or encased within materials such as amber (Cano and Borucki 1995) or in halite (Satterfield et al. 2005) may be unwitting travelers through a hostile geological medium that is thermodynamically static. However, if cells are shown to be active in any sense then they must be taking advantage of a thermodynamic disequilibrium within their geological setting (cf., Gaidos et al. 1999) even if detection of such disequilibria is beyond our current methods of measure and may only be computationally modeled.

Thermodynamic disequilibria are established in any environment by one or several chemical or physical regimes that determine the presence of some even furtive supply of electron donors and acceptors. Under such conditions microbes can make a living. And if conditions are ample in a large volume of earth then there may be enough active cells in such pockets to make a biogeochemical difference through their collective activities.

Numerous events or processes in and on Earth permit the conditions under which life has found or might find a way to be active (Fig. 1). Research has confirmed many of the processes depicted here; however, many of the processes are conjectural (e.g., that microbes respond to the long cycles of glacial compression and rebound). In addition to the type of event or process we might consider the time over which the respective phenomenon occurs or the spatial

Human-induced processes



Natural processes

Figure 1. Subsurface phenomena that create chemical, physical, or mixed chemical-physical gradients within which microbial metabolic activity is known to occur or might occur. Various processes are plotted along an arbitrary axis that defines whether the process is primarily chemical or physical in origin. Processes are classified according to whether they are mainly natural associated with Earth (non-bold), mainly human-induced (bold), or best described as being either natural or human-induced (shaded).

extent of its influence as additional attributes pertinent to any microbes exploiting a subsurface niche. For example, the acute effect of a meteorite impact or nuclear test would be followed by a chronic stage of alteration of the surrounding medium that plays out over millennia as the system stabilizes. By comparison, processes governed by the diurnal sighing of tidal action or (on a much longer period) the compression and rebound associated with advance and retreat of continental glaciers are related in a temporally cyclic manner and lack a concussive onset. The spatial boundaries over which these phenomena occur range from submicron perturbations that any of these processes could impose on individual cells to continental- or global-scale processes or events such as sea level rise or seismicity.

Natural processes that are reasonably well studied and that have been demonstrated to stimulate life include hydrothermal vent systems, geothermal activity or volcanism, and serpentinized environments. Noteworthy here is that many of these sites have been sampled at the surface, as windows to the subsurface; however, deep coring rarely occurs either because of the expense of seagoing expeditions or because the fragility and aesthetic value of sites like the thermal features of Yellowstone National Park is too much to risk. It is also exceedingly difficult to collect competent cores of representative quality from deep within porous and fractured rocks of volcanic provenance and where thermal fluids circulate, because the same properties that make these fluidically active habitats also lead to the likelihood that drilling fluids will contaminate the interiors of these cores. For a number of the natural processes we can only speculate that there are microbes in the subsurface that are managing to use to their benefit the chemical or physical energy inherent to the system. These processes would include roll front development for various ore bodies (e.g., uranium), plate subduction, hydrate formation and decomposition. Receiving attention recently, some in deep subsurface environments, are processes that include permafrost thawing (Mackelprang et al. 2011), infrared radiation (Beatty et al. 2005), seismic activity (Hirose et al. 2011), and asteroid or comet impact (Cockell et al. 2012).

And now, in the Anthropocene, many human-induced processes intentionally or unintentionally have manipulated the subsurface to favor microbial activity and survival. Probably the most prominent example is the remediation of contaminants in the subsurface with resulting changes in microbial activities. With thirty years of experimentation, scientists and engineers have encouraged bioremediation of numerous human-introduced chemicals by introducing to the subsurface nutrients, electron acceptors, and electron donors and by altering the direction and rates of groundwater flow (Hazen 1997). Numerous refinements of these processes include organic destruction, metal redox changes, and precipitation in mineral form, and even so-called natural attenuation in which the contaminants are monitored as they disappear in the presence of naturally occurring microbes shown to adapt and degrade the waste. Other purposeful geomicrobial processes include microbial enhanced oil recovery, which aims to alter the mobility of hydrocarbons in porous media, and biohydrometallurgy, whereby oxygenated acidic fluids circulated through crushed metal ore bodies stimulate microbes that oxidize the iron and sulfide minerals and thereby leach metals from the decomposing rocks.

Human-induced processes that are conducted irrespective of whether they alter subsurface microbial activities almost certainly do alter such activities. Little research has been performed to determine how underground microbes respond to our engineering of repositories for nuclear or carbon-based wastes, hydrocarbon retort, nuclear weapons testing, unconventional oil and gas recovery, or geothermal energy exploitation. In these cases, the array of geotechnical phenomena that overwhelmingly relate to our extraction of energy from Earth or our storage of waste products of energy production also quite appropriately relate to the ways in which microbial processes can derive energy in the Earth. By engineering Earth, humans have imposed redox stratification, fluid movement, fracturing, seismicity, and groundwater fluctuation some of which are mirrored by natural phenomena; these changes most likely reform subsurface deserts into oases for microbial life. Whether life pre-existed in these deep settings may not ever be discovered. However, it has almost certainly become richer in these locations due to our efforts.

THE TOOLS THAT WE NEED

Over thirty years of concerted effort scientists have examined different locations of the planet's subsurface and have adapted and acquired a range of tools for conducting this research (Table 1). As for many system-science disciplines today, a step back to examine the approaches used for sample collection, characterization, and description reveals impressive progress.

Because access to the subsurface is a necessity in order to study it, scientists should consider how they can obtain deep samples. Once sample depths are beyond the reach of simple push cores or augers, coring while using circulating drilling fluids is required. Although this process is expensive and dirty it cannot be avoided if certain environments are to be explored. Coring remains a mainstay of subsurface sampling and numerous reviews summarize the approaches that should be used (cf., Kieft et al. 2007).

When sea- or land-based coring is used, researchers encounter the dual problems of low sample quantity and high cost per quantity of sample material recovered. To avoid drilling, many new sampling efforts have turned to new means of getting underground or maintaining a lasting sampling opportunity in the form of observatories or long-term sampling devices. Many researchers have taken advantage of deep mines that reach into Earth for minerals or to find secure geological strata that serve as repositories (Onstott et al. 1997; Pedersen 1999; Satterfield et al. 2005; Edwards et al. 2006; Rastogi et al. 2009). This approach means that some of our surveys of deep-Earth microbes are canted toward formations that are rich in economic minerals or are proximal to such formations, or else are abnormally quiescent (i.e., locations likely to be undisturbed as required for waste repositories). Elsewhere, sampling devices are placed into drill holes or squeezed into casing and allowed to incubate in place in order to

Table 1. Methods that have advanced or will advance our understanding and communication of subsurface microbiological processes. Shown are example studies from the mid-1980s to present including both subsurface and surface investigations. References are provided in the text.

Method	Enabling technologies	Some key studies
Sample collection	Drill ships, mine access, observatories, CORKS, SCIMPIs	IODP cruises, South African gold mines, long-term ecological research sites
Field analysis and manipulation	Stable isotope probing, “mark and recapture”, biogeophysics	Methanotrophy, denitrification, biodegradation, transport
Molecular science instrumentation	Extraction of nucleic acids, lipids or proteins; amplification, sequencing, comparison of nucleic acids; mass spectrometry of lipids and proteins; flow cytometry; single-cell techniques	-omics (e.g., genomics, transcriptomics, proteomics) studies of environmental microbial communities in soils, anoxic methane-rich sediments, thawing permafrost, ultra-low pH mine drainage
Cultivation	New bioreactor designs, high-throughput, miniaturization	anammox; SAR11 clade and <i>Pelagibacter</i> ; Iron Mountain, CA
Imaging	Fluorescent <i>in situ</i> hybridization, NanoSIMS; CT scanning, synchrotrons	Anaerobic methane oxidation, anammox
Computational simulation science	Bioinformatics, reaction-path modeling, thermodynamic modeling	Yellowstone hot spring mat communities, U mill tailings remediation
Data science	Internet	Census of Marine Life, the Tetherless World Constellation
Visualization, engagement	Internet, GoogleEarth	International Census of Marine Microbes

sample indigenous microbes. So-called CORKs or SCIMPIs (Davis et al. 1992; Moran et al. 2006) are used in the seafloor and parallel the multi-level samplers (Smith et al. 1991; Lehman et al. 2004), *in situ* flow cells (Nielsen et al. 2006), and flow through *in situ* reactors (FTISR) (Lehman 2007) used in the continental subsurface. Hybrids of these technologies and ingenious smaller systems that are deployed using clever means by which to sample microbes and chemistry include passive gas samplers (Spalding and Watson 2006), U-tube systems (Freifeld et al. 2005), and the ever-changing osmosampler (Orcutt et al. 2010). And, with pressure-coring tools becoming more common, there are opportunities to collect deep cores, return them to the surface while maintaining *in situ* pressure, and then transfer these samples into analysis systems without decompression. Still, only limited work has been done with such equipment (Parkes et al. 2009); however, along with other methods of restricting pressure (Bowles et al. 2011), the chance to examine microbial activities as they may actually occur *in situ* has been expanded.

Despite our ability to obtain samples over extended periods, it is important to note the bias associated with such samples. This concern was first considered in the late 1980s with observations of altered microbial communities following the coring of holes (Hirsch and Rades-Rohlkohl 1988). Lehman summarized the limitations of samplers incubated *in situ* with a guarded appraisal of the technologies (Lehman 2007). It is important to bear these cautions

in mind when analyzing the chemistry and microbiology of subsurface samples so obtained. Direct and rapid characterization of freshly acquired core and porewaters appears to be the best means by which to investigate native microbes in rocks.

In some cases, direct field methods of analysis or manipulation of microbial communities can provide essential data to understand the fundamental ecology of the subsurface or to determine the outcome of engineered processes in the subsurface. The new discipline of biogeophysics, an expansion from several geophysical methods of examining geological strata, keys on microbially-induced alterations of geological materials and how these alterations influence underground electrical signals in order to image the presence or activity of microbial communities (Allen et al. 2007; Revil et al. 2010). At this stage of development, these remote sensing strategies are mainly able to detect profound changes in such properties as electrical conductivity or specific conductance and are used mostly in places where purposeful stimulation of the microbiological characteristics of formations has occurred.

Microbiologists and hydrologists have also experimented with a miniaturized version of the macro-ecologist's "mark and recapture" experiment. In such investigations, microbes from the environment are collected and then cultivated in the presence of a ^{13}C -labelled substrate. The result is a population that is uniquely tagged with the heavy isotope and that can then be released in a well-field and hopefully collected down gradient to determine the degree to which microbes may be transported in the aquifer (Holben and Ostrom 2000; DeFlaun et al. 2001). This method is similar to stable isotope probing that uses a heavy isotope label to determine the most active communities in a sample based on their ability to take up the label and deposit the label in their DNA (Radajewski et al. 2003). Push-pull tests are another ingenious method of examining *in situ* activities and, if done with discretion, can provide useful information about the real metabolic capabilities of microbes in Earth (Haggerty et al. 1998; Pombo et al. 2005; Urmann et al. 2005).

Scientists also now possess a range of tools that can be used for biological characterization once samples have been recovered from the subsurface and are returned to the lab. Significant advances have been made in the ability to characterize the molecular capabilities of cells (i.e., genomics, transcriptome, proteome, metabolome, lipidome) that allow many of the same diagnostic tools as used in modern medicine. The Richmond Mine site has provided a wealth of information related to the function of highly constrained microbial communities of limited diversity. In a classic application of metagenomic information, the researchers investigating this site gained enough genomic data of the simple communities present that they were able to devise appropriate cultivation methods of uncultured members of the microbial assemblage (Tyson et al. 2005). The same communities have assisted the understanding of ecological divergence of similar but slightly distinct microbes according to the genomic and proteomic patterns detected in these cells (Denef et al. 2010). These relatively simple systems are teaching us how best to transfer this skill to environments that are intrinsically more biologically complex or where a limited number of samples may be available.

It is acknowledged that few microbes can be nurtured using traditional cultivation-based approaches. However, startling progress has been made in culturing pelagic marine microorganisms by taking advantage of high-throughput robotic systems to handle samples, parsing them into numerous novel media formulations, and then recognizing that populations of these cells simply do not achieve high density (Connon and Giovannoni 2002; Stevenson et al. 2004). Along with the aforementioned metagenomic approach applied in the Richmond Mine, the approach used for pelagic microbes could be applied to microbes from Earth's subsurface. Matching these cultivation approaches with single-cell manipulation techniques (Stepanuskas and Sieracki 2007) may tease geologically inclined microbes into culture and also help to minimize the influence that PCR-based amplification has on our view of microbial diversity. When these approaches are combined with imaging techniques that hinge on fluorescence

microscopy (Amann et al. 2001), synchrotron-based characterization (Holman et al. 1998), interferometry (Davis and Lutge 2005), atomic force microscopy (Warren et al. 2001), and different electron microscopy techniques, we gain new resolution of the relationships between these microbes and the minerals on which they depend.

To complete our ability to comprehend life underground, there are new opportunities in the sciences that are not directly associated with the field and lab. Collectively, computational and simulation sciences have made advances that enable rapid and detailed modeling of porous media, where microbes participate in the alteration and dissolution of mineral species. Reactive transport models (Steefel et al. 2005) are now merged with *in-silico* models and bioinformatics approaches (King et al. 2009) to yield simulations that explain or predict the active microbial taxa in a given subsurface setting under a given set of environmental conditions (Scheibe et al. 2009; Li et al. 2010; Zhuang et al. 2011). Thermodynamic modeling has been pacing these studies and also can provide essential insight into what microbial processes are likely to be active and when they are active (Spear et al. 2005) and, when combined with kinetic models, how rapidly the activities may occur (Jin and Bethke 2005).

Finally, advances in data science and in visualization are poised to help scientists who study the subsurface with their task of communicating the results of their findings. Web science and its new ways of using the internet (Fox and Hendler 2011) will allow a binding together of disparate disciplines such that the data acquired by scientists in different fields can begin to sketch relationships between the living and non-living in the subsurface, as is happening in other fields. We can now take advantage of work done to explore interwoven features of the oceans (Amaral-Zettler et al. 2010; Tittensor et al. 2010), or of the cosmos (Szalay and Gray 2001) and proteins (Askenazi et al. 2011), to do the same for places inside Earth. This visualization capability should lead us to the point of greater interaction with the public where engagement through websites and museums, and possibly problem-solving through crowdsourcing will draw our knowledge and questions about the subsurface into the vernacular of non-scientists.

THERE'S NO PLACE LIKE HOME

As already noted, the earliest intensive investigations of underground life often targeted locations of known resource prospects for fossil energy or metals (e.g., Taylorsville Basin, South African Gold mines), sites where groundwater or soils were contaminated (e.g., numerous U.S. Department of Energy [DOE] and other sites), planned nuclear waste repositories (e.g., Äspö, Yucca Mountain, Waste Isolation Pilot Plant), or simply where it was convenient to collect subsurface material (e.g., ODP sites). This pragmatic approach has surely biased our understanding of subsurface life. Many of the more recent sites that have been investigated were selected in order to test hypotheses grounded in prior knowledge of life in the subsurface and what appear to be the limits of the biosphere. The *Early studies and comprehensive reviews* section identifies papers (especially, Fredrickson and Balkwill 2006; Onstott et al. 2009a; Schrenk et al. 2010, 2013; Edwards et al. 2012) that highlight different locations where subsurface research has been conducted. We will not reiterate the reports of these papers but rather point to some selected studies that have helped us to understand the range of underground locations already investigated. Within the next few years, the newly initiated Census of Deep Life is expected to provide a catalog of subsurface life and the disparate geological settings where it has been detected.

The DOE is responsible for waste released into a range of subsurface environments over 40 years following World War II (Riley et al. 1992). Despite this very practical concern, the DOE Office of Science supported research paths to understand the basic properties of life underground. Initially, DOE-funded scientists were unfettered with the need to examine actual waste sites; rather, they “cut their teeth” at pristine locations where the limits of subterranean

life could be explored. This freedom led to reports of microbial life in deep and shallow coastal plain sediments and rocks on the southeastern coastal plain of the U.S. (Fredrickson et al. 1991), in Oyster, Virginia (Zhang et al. 1997), and in the Taylorsville Triassic Basin (Onstott et al. 1998); thick sedimentary zones in arid regions (McKinley et al. 1997) and in thick flood deposits (Brockman et al. 1992; Kieft et al. 1998) in southwestern Washington state; unsaturated and saturated fractured basalts of the Snake River Plain Aquifer (Colwell and Lehman 1997; Lehman et al. 2004) and the Columbia River Basalt Group (Stevens et al. 1993; Stevens and McKinley 1995); hot methane-charged fractured sandstones in the Piceance Basin (Colwell et al. 1997); volcanic tuffs in the Great Basin (Amy et al. 1992; Russell et al. 1994); and ancient marine sediments into which volcanic dikes impinged in New Mexico (Fredrickson et al. 1997; Krumholz et al. 1997).

Studies of deep seafloor locations for the presence of microbes have now reached well beyond the original skin of sediment examined by the first marine microbiologists. The ODP and later the Integrated Ocean Drilling Program (IODP), using the drilling platforms *JOIDES Resolution* and the *Chikyu*, catalyzed the first seagoing coring expeditions aimed at answering questions about the deep marine biosphere. Microbiologists participated in earlier drilling legs and continue to be opportunistically involved in expeditions.

Through targeted drilling and coring studies of different seafloor environments, we are slowly accumulating information about subsurface communities in a broader range of environments. The ODP Leg 201 was the first dedicated microbiology drilling leg. Leg 201 scientists used procedures to control contamination (Smith et al. 2000) that helped to reach out to a new scientific community. This sampling cruise occurred off the west coast of South America and explored deep sediment in near shore and pelagic sites (D'Hondt et al. 2004; Inagaki et al. 2006). More recent dedicated microbiology investigations sponsored by the IODP have explored the eastern flank of the Juan de Fuca Ridge, a hydrologically active basalt aquifer (Fisher et al. 2011); the South Pacific Gyre, in which sediment and basalt underlie waters of extremely low productivity (D'Hondt et al. 2011); a hydrothermal field in the Okinawa Trough (Takai et al. 2011); and basalt and sediment of North Pond, a sediment-filled basin off the main axis of the Mid-Atlantic Ridge (Expedition 336 Scientists 2012).

One of the curiosities of subsurface microbiology studies is that only rarely do these subsurface environments lack measureable life. Teams of scientists are typically able to tease evidence of life out of most subsurface rock or sediment that is within a temperature regime that embraces the known limits of life and that has enough connected pore space. Even environments contaminated with high levels of radioactive elements contain microbes (Fredrickson et al. 2004). Exceptions include the dry and thick unsaturated zone in the Eastern Snake River Plain (Colwell et al. 1992) and some sections of the deep, massive sandstones of the Piceance Basin (Colwell et al. 1997). It is possible that where life appears to be missing from moderate temperature regimes at moderate depths, the problem is either detection limit (with life present but not detectable) or inability to survey large enough samples due to the limited amount of material collected by coring.

Of course, our inquiries of the subsurface for microbes remain inadequate to survey the life there. Much of the field research has an exploratory element. We rarely acquire true replicate samples because drilling identical holes in the same location and sampling the same depths is both difficult and expensive. Furthermore, innumerable subsurface environments have thus far been ignored or simply been too difficult or expensive to reach. More sampling along lengthy, confined horizontal flow paths, similar to past studies on the southeastern coastal plain of the U.S. (Murphy et al. 1992), would provide excellent data about microbes limited by geological constraints. Notably static environments (Vreeland et al. 2000) will offer insight into how long cells can last and possibly the adaptations that they require in order to last on timescales of thousands to millions of years (Lomstein et al. 2012; Røy et al. 2012). Arctic and Antarctic

deep-Earth environments remain under-sampled, though we are gradually accumulating examples (Mikucki and Priscu 2007; D'Elia et al. 2008; Onstott et al. 2009b; Pham et al. 2009; Colwell et al. 2011). Deep samples from thick vadose zones are lacking in general as are studies that examine microbes present near faults in seismically active areas.

IS DIVERSITY THE SPICE OF SUBSURFACE LIFE?

Biologists, other scientists, and even non-scientists are justifiably attracted to the incredible diversity of life at Earth's surface and the essential aspect of surface life's diversity to the health of ecosystems is well recognized (Daily and Matson 2008; Rockstrom et al. 2009; Stein and Nicol 2011). Certainly, the amazing diversity of life at the surface is almost harrowing to those who must reach into the subsurface to get samples where biomass is typically low. We have devised ways of keeping our precious deep samples isolated from surface samples and also ways of determining whether surface contamination has occurred (Lehman et al. 1995; Masui et al. 2008). By relying on new methods of molecular characterization and accumulating enough information from the relatively rare sampling events, the nature of life's diversity underground is becoming clearer.

Bacteria and archaea are the common targets of investigations that aim to study subsurface diversity. Eukarya are rarely targeted. Subsurface studies often show that bacteria are more abundant in some subsurface environments than archaea (Schippers et al. 2005; Lin et al. 2006; Rastogi et al. 2009; Briggs et al. 2012), but others show that archaea predominate (Biddle et al. 2006) and some indicate more equal representation of the two domains (Pham et al. 2009). Microbiologists are now familiar with subsurface habitats like the sulfate-methane transition zone in the subseafloor, where the supply of sulfate from seawater and methane from deeper sediments dictate that both bacteria and archaea will be present. But this environment is now well defined and frequently sampled. While bacterial (vs. archaeal) abundance seems to be a common theme in the subsurface, more work is required to determine the relative abundance of these groups.

Before we can describe the patterns of abundance of these two domains in the subsurface, some technical hurdles associated with the methods of analysis must be overcome. Extraction of DNA from archaeal cells in deep sediments is difficult (Lipp et al. 2008) and, relative to bacteria, archaea typically have poorer representation in the gene databases that are required for construction of the amplification primers. Polymerase chain reaction (PCR) based amplification depends on these primers and some version of PCR is often used to assess microbial diversity. As usual, a robust approach using multiple methods of analysis (e.g., fluorescence *in situ* hybridization, DNA sequencing, and intact polar lipid characterization) is the best way to present a detailed description of an environmental microbial community. And new methods like single-cell sorting followed by whole genome amplification can help to explain where partiality associated with typical primer-based amplification has occurred.

Clone library-based investigations of the 16S rRNA and other genes obtained from DNA extracted from a number of samples find evidence of many taxa in the subsurface (cf., Biddle et al. (2006) and Inagaki et al. (2006), as well as examples as reported in Fredrickson and Balkwill (2006)). It is not unusual to detect "new" microbes in these surveys based on the presence of unique genes in the libraries. The results of diversity studies can be influenced by the manner of sampling and the types of sample used. That attached subsurface communities are different than free-living subsurface communities has been understood for some time (Hazen et al. 1991) but surveys of large volumes of subsurface space (e.g., pumping and filtering or concentrating aquifer samples for free-living microbial cells) may yield different findings than surveys based on a few grams of solid material. Examinations of subsurface microbial diversity rarely seem to address the question of how much of the subsurface was queried; however, adhering to

guidelines for how diversity is reported relative to the amount of material and the type material sampled (i.e., water, solids, or both) would help to develop our view of subsurface variations in communities (cf., Lehman et al. 2001; Lehman 2007).

That microbial diversity in the subsurface is typically lower than in surface systems is not necessarily surprising. Unique niches occur in the subsurface though certainly not as many as in surface systems. It seems that microbes that survive in the subsurface often need to overcome certain barriers that may not be unheard of at the surface (e.g., pressure, temperature, confinement) but may be more severe and persistent in the subsurface. Thus, species that can last at depth have been winnowed by the chemical and physical realities of their habitat once they arrive in the subsurface. Investigations of a low-pH system in the Richmond Mine at Iron Mountain in California attest to the stable presence of five dominant microbes making up most of the community (Tyson et al. 2004). Microbial communities obtained from deeply occurring fractures in the Mponeng Mine in South Africa provide another such example. There, a single sulfate-reducing microbe represents over 99.9% of the community (Lin et al. 2006); microbial ecosystems where diversity is so low are not commonly reported. Given the conditions of that environment and the genomic characteristics of this solo microbe, it seems to have the functional attributes needed in order to survive there, and indeed appears to be well distributed in the deep environment of the Witwatersrand Basin (Chivian et al. 2008). While both sites mentioned here are mine environments that are subject to unnatural forces during their creation, it seems plausible that similar low diversity communities can be found in utterly pristine locations of the subsurface.

Relatively low biomass and relatively low diversity may be the norm in much of the subsurface; however, some locations may not be so biologically depleted. Recent studies in the ocean crust at the East Pacific Rise reveal notable diversity (Santelli et al. 2008). These samples were not from the subsurface but might contain microbes representative of deeper crustal materials. While not as diverse as such well-studied surface systems as farm soils, these basaltic communities are considerably more complex than pelagic marine microbial communities (Santelli et al. 2008). Similar places that exhibit much milder geothermal gradients and probably milder flux through the system may foster simpler communities (Edwards et al. 2011). More open subsurface environments like those crustal basalts that benefit from a porous and fractured architecture and fluid circulation driven by geothermal processes (Delaney et al. 1998; Edwards et al. 2005) could be as abundant in the subsurface as the distribution of large igneous provinces around the planet (Saunders 2005) and volumetrically may be a significant source of the planet's underground diversity.

The search for eukarya was often a part of early investigations of aquifers (Sinclair and Ghiorse 1989; Sinclair et al. 1993; Novarino et al. 1997); however, many surveys of life underground may not even look for eukarya. The obvious spatial constraints in porous media prevent the occurrence of larger cells or multicellular organisms. However, we now have an example of a nematode that lives in deep fractures, apparently managing to subsist through grazing on microbes (Borgonie et al. 2011), and evidence of Collembola that exist in deep limestone caves (Jordana et al. 2012). These studies promote the idea that any deep system with cavities large enough (several microns?) such as the aforementioned crustal systems or fractured rocks may also be open enough to support eukarya if adequate unicellular biomass can be generated to provide food for the higher organisms. However, it should be noted that sterols, a key structural component of the membranes of eukarya, are difficult to make anaerobically and this may limit the extent to which these cells might penetrate anoxic zones.

Early considerations of the presence of viruses in the subsurface focused largely on shallow systems with considerable contact with surface environments (Gerba and Bales 1990; Matthes 1990). Certainly, in the subsurface viruses confront several factors such as low host biomass (Wiggins 1985), generally patchy, disconnected microbial communities (Brockman and Murray

1997), and limited fluid exchange between communities. These factors could minimize the effectiveness by which phages can infect bacteria or archaea. Some subsurface environments might be more likely to contain viruses (i.e., where microbes are abundant) and sand columns have been used to examine their distribution (Yates et al. 1997).

In recent years, studies in different environmental settings with new methods have revealed huge numbers of viruses (Anderson et al. 2013). These small packages of genetic information are now believed to provide significant paths for moving molecular information among microbial hosts and represent a massive means of turnover for living biomass (cf., Suttle 2005; Anderson et al. 2011a, 2013). Accordingly, there have been more studies of viruses in the subsurface. A metagenomic study conducted on marine sediments found highly diverse, largely unrecognized phage populations and identified marine sediments as a massive “reservoir of sequence space” (Breitbart et al. 2004). Finding more temperate than lytic phage suggests that an important infection strategy for subsurface viruses might be incorporation into the host genome rather than destroying the host. Shallow sediments worldwide contain large numbers of phage and collectively these have profound impact on biogeochemical cycles, at least as pelagic microbes are buried in the uppermost sediment layers (Danovaro et al. 2008). Microbial cell death due to phage in benthic systems of deep waters may significantly haze the viable microbes that are buried and increase the amount of organic detritus in the uppermost centimeters and alter the disposition of buried organic matter.

Recent identification of a “microbial immune system,” the so-called clustered regularly interspaced short palindromic repeat or CRISPRs, found within host genomes offers a new means of detecting phage populations and connecting them to their respective hosts (Banfield and Young 2009). The CRISPR approach was applied to hydrothermal vent samples and the results indicate that large numbers of microbial hosts are infected with viruses and that the hosts represent a diverse range of microbes (Anderson et al. 2011b). These vent fluids speak to the processes and populations in the subsurface and naturally lead to considering how phage may play a role in the transfer of genetic information in subsurface environments other than vents. That phages dictate the genetic diversity and evolution of microbial communities in vent systems of the shallow subsurface (Anderson et al. 2011a) suggests that microbial communities in other subsurface locations where fluids are actively moving may benefit from the enhanced genetic fitness and functional capacities that are conferred by prophages and that may assist subsurface survival.

BIOMASS OF SUBSURFACE LIFE

The pioneering study of global biomass by Whitman and colleagues (Whitman et al. 1998) proposed that subsurface bacteria and archaea comprise 35 to 47% of Earth’s total biomass, nearly equal to plants in their total carbon content. Microbes in subseafloor sediment comprised nearly 1/3 of their global biomass estimate. Microbes in terrestrial subsurface sediment comprised between 1/50 and 1/5 of their global biomass estimate. Their study was a great starting point for estimates of global microbial biomass. However, it was based on the relatively sparse data that were available in the mid-1990s. Estimates of subsurface biomass are changing as more data become available.

Cell abundance data for terrestrial subsurface sediment have not significantly improved since 1998. However, data for subseafloor sediment have improved greatly. Subsequent studies have generally yielded lower estimates than Whitman et al. (1998) for subseafloor sedimentary biomass (Parkes et al. 2000; Lipp et al. 2008; Kallmeyer et al. 2012). The most recent studies show that cell concentrations in the broad expanses of open-ocean sediment beneath the Pacific gyres are orders of magnitude lower than earlier counts from subseafloor sediment, which were largely limited to organic-rich sediment that underlies oceanic upwelling zones (D’Hondt et al.

2009; Kallmeyer et al. 2012). Consequently, total microbial abundance varies between sites by five orders of magnitude (Kallmeyer et al. 2012). This variation strongly co-varies with mean sedimentation rate and distance from shore. Based on these correlations, total cell abundance in subseafloor sediment is $\sim 3 \times 10^{29}$ cells, corresponding to ~ 4 petagram C and $\sim 0.6\%$ of Earth's total biomass (Kallmeyer et al. 2012).

Most estimates of subseafloor sedimentary biomass are based on visual cell counts, which do not include spores (the fluorescent dyes used for cell counts generally do not penetrate spores). However, a recent study of dipicolinic acid and muramic acid concentrations indicate that bacterial endospores are approximately as abundant as counted cells in deep subseafloor sediment of the Peru Margin (Lomstein et al. 2012; dipicolinic acid is limited to endospores and muramic acid is much more abundant in endospores than in vegetative cells). Because endospore abundance is not yet known for other subsurface habitats (or even for most sediment of the world ocean), this will be an intriguing avenue of research in the near future.

A further complication for global estimates of subsurface biomass is that the biomass resident in large subsurface habitats is not yet known. For example, biomass in the vast volume of fractured igneous basement in continents and oceans cannot yet be quantified, because the data do not yet exist.

Discussions of subsurface biomass to date have relied on counts of stained cells (e.g., (Thierstein and Störrlein 1991; Parkes et al. 1994, 2000; D'Hondt et al. 2004, 2009) or abundance of intact biomarkers (Lipp et al. 2008). These are powerful techniques that census overlapping, but non-identical subsets of a microbial community. Counts of cells stained with non-specific DNA-binding compounds (e.g., acridine orange or SYBR-Green) include intact vegetative bacteria and intact archaea, but do not include bacterial endospores. Molecular probes (fluorescence *in situ* hybridization [FISH] probes) specific to RNA in bacteria, archaea or more narrowly defined phylogenetic groups have also been used, albeit much less frequently (e.g., Mauclaire et al. 2004; Schippers et al. 2005). Intact biomarker assays have focused on archaeal biomarkers and consequently estimate subsurface archaeal biomass, not total subsurface biomass.

The results of these techniques (cell counts and biomarker assays) beg discussion of the distinction between "intact" and living cells. It is not yet certain how long cell membranes, their included nucleic acids, or their diagnostic phospholipids remain intact after cell death in deep subsurface environments. This said, RNA-based FISH counts are generally interpreted to suggest that a large fraction of counted subseafloor sedimentary cells are living or at least recently alive; RNA is widely recognized to degrade far more readily than DNA and RNA-based FISH counts constitute several percent to several tens of percent of DNA-based counts in subsurface environments (Mauclaire et al. 2004; Schippers et al. 2005). Compelling independent evidence that the majority of counted subseafloor cells in individual samples are alive was recently provided by experiments with isotopically-labeled organic substrates and sediment from hundreds of meters beneath the seafloor in the Japan Sea (Morono and al. 2011); in these experiments, as many as 76% of the counted cells assimilated the isotope-labeled substrates.

PHYSIOLOGICAL PROCESSES OF SUBSURFACE LIFE

Subsurface microorganisms include both heterotrophs (which consume organic matter) and lithoautotrophs (which consume inorganic compounds). Electron donors in subsurface environments include buried organic matter, reduced chemicals (such as reduced iron and reduced sulfur), and reduced compounds created by water-rock interactions; examples include H_2 from radioactive splitting of water (Pedersen 1997; Lin et al. 2006) and H_2 and CH_4 from serpentinization reactions (Kelley et al. 2005; Nealson et al. 2005). All of these electron donors occur in

a broad range of subsurface environments. For example, organic matter that was photosynthesized in the overlying ocean is the principal electron donor for microbes in subseafloor sediment (D'Hondt et al. 2004) and also circulates in dissolved form with seawater through oceanic basalt. The primary electron donors in subseafloor basaltic aquifers include reduced chemicals in mineral phases (e.g., Bach and Edwards 2003), which also commonly occur in both terrestrial and marine sediment. Hydrogen produced by natural radioactive splitting of water appears to sustain microbial life in deep continental aquifers (Lin et al. 2006) and may also be a significant electron donor in very organic-poor marine sediment (Blair et al. 2007).

Rates of subsurface microbial respiration have been most commonly quantified for subsurface sedimentary communities. Calculations based on concentration profiles of dissolved electron acceptors and products of microbial respiration indicate that the subsurface microbes of both terrestrial sedimentary aquifers (Chapelle and Lovley 1990; Phelps et al. 1994) and subseafloor sediment (D'Hondt et al. 2002a, 2004; Røy et al. 2012) respire orders of magnitude more slowly than microbes in the surface world (Onstott et al. 1999; Price and Sowers 2004).

Given the extraordinarily low rates of microbial respiration in many subsurface environments, subsurface microbes are generally assumed to reproduce very slowly, if at all. D'Hondt et al. (2002a) speculated that most subseafloor sedimentary microbes are either inactive (dormant) or adapted for extraordinarily low metabolic activity. Price and Sowers (2004) suggested that subsurface sedimentary microbes exhibit survival metabolism (sufficient to repair macromolecular damage but insufficient to sustain growth or motility). Whether they are actually growing or merely repairing macromolecular damage, amino acid racemization ratios indicate that subseafloor sedimentary biomass turns over very slowly, on timescales of hundreds to thousands of years (Lomstein et al. 2012). We do not yet know whether the microbes of these subsurface environments reproduce at these slow rates of biomass turnover or live without dividing for millions to tens of millions of years.

These extraordinarily slow rates of respiration and biomass turnover beg consideration of the factor(s) that control(s) rates of microbial activity in subsurface ecosystems. Where electron acceptors are present, areal or volumetric rates of subsurface microbial activities (e.g., activity in a square-meter sediment column or in a cubic meter of sediment or rock) are broadly related to electron donor availability. For example, areal rates of microbial respiration are orders of magnitude higher in subseafloor sediment rich in organic matter (D'Hondt et al. 2004) than in subseafloor sediment where organic matter is extremely dilute (D'Hondt et al. 2009). However, where electron acceptors are vanishingly rare, electron donors can build to extraordinarily high concentrations. For example, in some fractures intersected by deep South African gold mines, dissolved hydrogen from water radiolysis is present in millimolar concentrations but electron acceptors are scarce, indicating that microbial activity in those fractures is far too low to keep up with very low rates of hydrogen production on timescales of tens to hundreds of millions of years (Lin et al. 2006).

Such examples suggest that, in a broad sense, subsurface rates of bulk microbial activities are controlled by energy availability. However, the situation is much more problematic at closer inspection. For example, why don't sedimentary microbial communities oxidize all available organic matter within the first few centimeters of the seafloor? In other words, how does buried organic matter survive microbial activity to sustain slow rates of activity for millions to hundreds of millions of years? Why don't rapidly respiring cells outcompete the slowly respiring cells by oxidizing all available organic matter over a much shorter interval of geologic time?

The situation is also perplexing on a per-cell basis. For example, aerobic microbial communities of subseafloor sediment in the North Pacific Gyre exhibit per-cell rates of microbial activity (Røy et al. 2012) that are not vastly different from per-cell rates in anaerobic communities of subseafloor sediment in the Peru Margin and the equatorial Pacific Ocean

(D'Hondt et al. 2002a; 2004) although areal rates of activity differ by orders of magnitude between the oxic gyre sediment and the anoxic Peru Margin sediments. In both environments, mean per-cell rates of respiration are orders of magnitude lower than per-cell rates in surface sediment or laboratory cultures. What are the limits to survival that allow microbial hunger artists to eke out a living at such extraordinarily slow rates in both environments?

Finally, the extent to which subsurface organisms are (i) microbial zombies, incapable of being revived to a normal state, or (ii) capable of metabolism, growth, and reproduction at rates typical of the surface world is not yet known for many subsurface ecosystems. Isolation of many microbial strains from deep subsurface environments (e.g., (Balkwill 1989; Takai et al. 2001; D'Hondt et al. 2004; Batzke et al. 2007) has demonstrated that at least some deep subsurface microbes can emerge into the surface world, grow, and multiply. However, these few hundreds of laboratory isolates may not represent the majority of subsurface microbes. A recent study by Morono et al. (2011) sheds light on this issue. In short, Morono and colleagues demonstrated that many microbes from sediment hundreds of meters beneath the seafloor take up measurable quantities of isotopically-labeled substrates. In doing so, they effectively showed that many deeply-buried organisms maintain the potential to metabolize and grow, regardless of what they are doing deep beneath the seafloor (Jørgensen 2011). This result effectively demonstrates that the metabolic potential of long-buried microbes can be activated at much higher rates when they emerge into a moderate environment.

WHERE AND WHEN DOES LIFE IN THE SUBSURFACE REALLY MATTER TO US?

It is fair to ask when and where deep life matters to the life and processes at Earth's surface. Can we identify ecosystem services that are provided by life underground? The question might be considered for both naturally occurring subsurface microbes and those that are a part of a human-engineered process. A number of engineered systems that utilize or involve microbes and their activities are the result of stimulation of subsurface life. Microbes underground are responsible for numerous variations on the general theme of bioremediation. Where wastes have been carelessly released into aquifers, we now depend upon microbial communities to decontaminate these freshwater resources. The processes can take decades to be complete; however, it is usually far more economical to track these *in situ* reactors over time as they eliminate contaminants than it is to dig the waste out of the ground. Perhaps this situation is similar to how we depend upon subsurface microbes and the biogeochemical processes that they carry out to purify tainted water that enters the subsurface prior to our use of the water when it is collected down gradient. The mingled biological, chemical, and physical processes that are inherent to deep Earth can eliminate the human pathogens that are simply not able to survive.

Similar processes have been conceived for conducting *in situ* mining or biohydrometallurgy, where low-grade ores may be attacked by well-understood microbial processes under controlled conditions to extract the metals within (Das et al. 2011). This approach is conducted worldwide in managed "heap leach" operations, where the rubblized rock is piled onto a large impermeable pad, irrigation networks trickle the "lixiviant" fluid through the system, and the metal-rich liquid that results is collected (Rawlings 2002). This biologically driven process is responsible for the recovery of most of the world's copper (Rawlings 2002), as well as uranium and gold, and is contemplated for manganese extraction (Das et al. 2011).

Microbes and their astounding metabolic activities have been considered also for processes that would convert hydrocarbons deep in Earth into products that can more readily be extracted. Studies of anaerobic modification of hydrocarbons are relatively new as many aliphatic and aromatic structures were long considered to be inert (Heider et al. 1998). A better understanding

of the distribution of microbes in hydrocarbon-rich geological formations and the constraints under which they survive and modify the organic matter therein (Head et al. 2003) has also led to considering ways by which microbes might alter hydrocarbons in place where oxygen is absent. Oxygen-free reactions, including hydroxylation, methylation, fumarate addition, and reverse methanogenesis (anaerobic methane oxidation), allow microbes metabolic access to complex hydrocarbons and broaden our view of how organic matter can be converted in the subsurface (Heider 2007).

Many subsurface environments are used not for their resources but rather for their remoteness, stability, or controllability. Such subsurface settings are ideal repositories for nuclear waste, carbon dioxide, or as artificial reservoirs for natural gas. In each case, microbial activities may play a role in the security of the materials deposited therein. The microbiology of nuclear waste storage locations has been investigated to determine the degree to which biological activity may alter the waste in a range of geological environments designated as candidate underground repositories (Stroes-Gascoyne and West 1997; Pedersen 1999; Pitonzo et al. 1999; Jolley et al. 2003; Horn et al. 2004; Nazina et al. 2004). Locations close to the waste canisters shortly after enclosure may create conditions that are outside the range of microbial survival due to high-temperature or high-radiation fields of the newly deposited waste. But at some distance away from the waste, and over time as the radioactivity decays, these extreme conditions will moderate and microbes may recolonize the geologic niches.

Deep-Earth storage of carbon dioxide as a means to remove it from the atmosphere has received considerable attention. Most studies have focused on the physical or chemical controls on carbon dioxide stability in the subsurface (Benson and Cook 2005); however, microbes can survive in many environments suitable for CO₂ storage and for these settings we must also consider biogeochemical aspects of stability. To date, few studies have examined microbial communities where CO₂ would be secured or the conditions to which these cells would be exposed. These investigations make it clear that some microbes—likely as spores—can survive in the presence of supercritical CO₂ (Mitchell et al. 2008, 2009; Dupraz et al. 2009). Native communities in several geological habitats may resist the solvent properties of the supercritical CO₂ or survive proximal to the highest concentrations of the solvent (Morozova et al. 2010). Geochemical modeling suggests that subsurface microbes in some environments where CO₂ could be disposed (e.g., basalts) might be able to alter the disposition of the carbon (Onstott 2004). Assurance of the stability of deeply sequestered CO₂ is important and so there should be an effort to understand the biogeochemistry where life can survive.

Even though these microbial reactions occur only on the dimensions of single microbial cells or microcolonies or minerals or dissolved compounds, the large size of the biomass, its ability to permeate living space, and the relentless nature of this metabolism mean that the effects can translate to scales of hundreds of kilometers over millennia. An example that displays the cumulative effect of pervasive, sustained microbial activity is the accumulation of biogenic methane in continental shelf sediments where conditions are met for methanogenesis and capture of the methane in the form of hydrates (Hazen et al. 2013). The release of methane from this “large, dynamic microbially-mediated gas hydrate capacitor” (Dickens 2003) is one explanation for how massive quantities of isotopically-light carbon were injected into the Earth system at the Paleocene-Eocene thermal maximum and possibly at other times in Earth’s history. Computational modeling of how microbially-generated methane accumulates in sediments as hydrates, free-gas, or dissolved gas; how it is oxidized by microbes under normal conditions of leakage; and how it may escape from sediment and enter the overlying water or atmosphere and act as a greenhouse gas have matched the observed $\delta^{13}\text{C}$ excursions in the sediment records (Dickens 2003; Gu et al. 2011). Although Earth-system models indicate that the current phase of planetary warming is unlikely to cause large-scale release of methane present as hydrates (Archer 2007), modeling efforts that focus on high-latitude sediment suggest that more

immediate release of methane from hydrates is possible (Reagan and Moridis 2009). It seems that sediment and deep permafrost containing accumulations of biologically-produced methane are perhaps especially precarious (Westbrook et al. 2009; Ruppel 2011). That methane plumes can transition from the seafloor, through the water column, and then to the atmosphere is notable (Solomon et al. 2009). Polar field sites may be excellent places to observe how subsurface biota and their processes respond to the surface system (and how humans are changing it) and may be responsible for accelerating (i.e., by making methane) or quenching (i.e., by consuming the methane) the changes that are underway on the planet's skin.

Another example of microbes in their native state that may contribute significantly to processes of concern at the surface are those present in deep aquifers covered by the oceans and contained within large igneous provinces or proximal to spreading centers or seamounts (Schrenk et al. 2010; 2013). By virtue of their activity, these cells likely play an important role in planetary elemental cycling. These regions of considerable fluid movement are driven by advection of seawater into the crustal materials at the seafloor and by thermal convection cells generated when geothermally-heated waters circulate through the porous geological structure (Delaney et al. 1998; Edwards et al. 2005). Life in the crust consists of microbes that form complete ecosystems with lithoautotrophy and heterotrophy present (cf., Cowen et al. 2003; Santelli et al. 2008; Mason et al. 2010; Smith et al. 2011). It has been estimated that ca. 1×10^{12} g C/yr of primary biomass may accumulate based on the volume of accessible crustal material; the amount of water cycling through these sponge-like materials; and the iron-, sulfur- and hydrogen-based metabolisms upon which these ecosystems rely (Bach and Edwards 2003). Thus, crustal communities mediate the flux of crucial elements from the mantle to the overlying water, where chemical energy is converted into microbial cells (Menez et al. 2012). The surface (seafloor) exposures of these deep aquifers are windows through which the geochemical and microbiological fluxes may shine into the overlying water. These surface windows exhibit diverse and complex accumulations of life based on interaction of the released fluids and the microbes with the seawater into which they emerge (Bernardino et al. 2012; Thurber et al. 2012). And because of their distance from us, we have not yet completely seen deeply into these windows where there may be complex ecosystems projecting into the crustal materials.

Also of some importance is the concept that the subsurface was once a refuge for life when the surface was too harsh to allow survival (Stevens 1997). Early in the planet's history, perhaps after life started, but still when surface conditions were austere, the subsurface might have been relatively stable, perhaps even much as it is today. Bolide impacts might have routinely sterilized the surface proximal to the impact; however, at some distance the resultant fractures and fluid movement (Cockell et al. 2012) might have provided conditions that would enhance survival. The same might have been true in some regions that sustained ice cover. Here, at some depth (as is the case in present high-latitude locations), the balance between low surface temperatures and high subsurface temperatures would offer thermally optimal conditions for long-term stability of microbial communities. It is sobering to think that billions of years from now, as the Sun sears the surface of Earth, life may make its final stand in the refugial depths of the planet.

PROJECTIONS AND PRIORITIES FOR FUTURE STUDIES

The future of subsurface microbiology research is rich with opportunities to understand the peculiarities of this environment and how these characteristics define it as an important component of the biosphere. The priorities for future studies can be divided into topics that deal with how the subsurface is sampled and envisioned and also into topics that relate to traits of subsurface microbes or the ecology, diversity, biomass, activity, and constraints of microbial communities.

Imagining how we might sample and visualize deep life

Numerous research tools are available to those who would study microbes in the subsurface; however, new tools would enable even more information to be gleaned from our deep investigations. New experiments could help to address the bias that is expected to occur associated with sampling. Although there are few examples of such experiments (cf., Hirsch and Rades-Rohlkohl 1988), we can expect that the mere act of coring, pumping, or excavating in an underground environment may stimulate microbes in an otherwise quiescent setting. The creation of open boreholes that serve as vertical conduits for fluids where none existed before, or of hydrologic gradients associated with the strenuous pumping of aquifers, are certain stimuli to microflora that are used to stasis. Crustal systems are notoriously difficult geological environments to sample because of their inherent porosity and the tendency for fluid imbibition that can carry drilling materials into the rocks.

Equally delicate conditions are those required for studies that would manipulate the subsurface and determine the biogeochemical responses. Numerous examples reveal how microbes respond to purposeful changes in their environment in shallow or accessible systems derived from bioremediation research, but fewer studies have been conducted in deep or hard to access habitats and experimental alterations in such places are more complicated. Observatories such as CORKS and SCIMPIs in the seafloor have yielded exciting results, but even these devices are not always straightforward to deploy. As an example, settings where methane concentration, pressures, and temperatures are conducive to gas hydrate formation are not yet amenable to the observatory approach because hydrate formation prevents easy recovery of the samplers. Elsewhere, in methane-rich sediments with increased leakage of the gas, where fractures are important in distributing microbes, where shale gas is extracted, or in formations designated as CO₂ repositories, we can envision that sampling systems such as CORKS, osmosamplers, or FTISR will be essential for puzzling out the implications of microbial activity.

Advances are pending in the ways that we see or imagine the subsurface. From a computational perspective, new modeling approaches will strengthen the relationships of scientists dwelling on the disciplinary boundaries and drive new studies that can be approached using experimental work or field collections. The coupling of genome-scale models to reactive transport models has been accomplished in soils systems (cf., O'Donnell et al. 2007) and even in boutique subsurface settings like the Rifle uranium mill tailing cleanup site in western Colorado (Scheibe et al. 2009). The Rifle research sets a splendid example for other researchers, who must explain the biogeochemistry of their target environment underground. We eagerly anticipate the application of the same visualization and engagement tools that have been used for developing visual observatories in astronomy (Szalay and Gray 2001), mapping human proteins (Askenazi et al. 2011), detecting new evolutionary relationships in a spatiotemporal context (Kidd and Liu 2008; Sidlauskas et al. 2009), seeing the biogeographical distribution of large species (Kidd 2010), and observing how disease propagates through time (Janies et al. 2007). By using data science and visualization approaches, we can move towards conceptual models of subsurface life that will help us to realize principles of that life comparable to how progress has been made in Earth system models, simulations of deep astronomical time, and human behavior (Wright and Wang 2011). Perhaps just beyond the computational models of subsurface life will be the physical models, the holograms and GeoWalls, that allow us to see in our museums or auditoriums the inside of a living Earth.

Unexplored adaptations of subsurface microbes

Because the subsurface presents such extreme conditions for microbial life compared to life at the surface where most of our studies of life occur, it is a challenge to understand how processes that we recognize as essential for life, or even routine for life, can occur in the subsurface. We have already noted the records for long-term survival that life underground

appears to sustain; and subsurface cells may depend substantially on the formation of inert endospores (Lomstein et al. 2012). However, if cells remain marginally vegetative in some sense their protracted temporal survival evokes new questions:

- Do such vegetative cells grow at all or do they exercise some sort of contact inhibition when they lie in such intimate space with minerals?
- Can these living cells evolve—and if so, at what rate—without undergoing the cell division that normally seems so essential to introducing genetic change to a population? Might new evolutionary mechanisms be discovered such as CRISPRs (Banfield and Young 2009) or the apparent importance of viruses (Anderson et al. 2011a, 2013) in these deep dwelling cells?
- Are there specific adaptations (e.g., efficient nucleic acid repair) associated with survival proximal to minerals that may be undergoing radioactive decay (Arrage et al. 1993)?
- For cells that enter into extended dormancy, how do they muster the bare metabolic activity to repair damaged (i.e., oxidized or racemized) molecules that are associated with survival in aqueous media?
- Have these cells found new ways to resist high temperatures, as would appear to be true for some that live under the dual stress of high temperatures and pressures (Takai et al. 2008)?
- As some may be fixed in place for millennia, have these cells devised structures such as “nanowires” or pili to explore neighboring pores and fractures for the thermodynamic disequilibria that are essential for gaining energy (Nielsen et al. 2010)?

Unstudied physiologies and genotypes for the subsurface

The demands of life in Earth's deep reaches appear to invoke as-yet unstudied strategies by which microbes achieve the necessary energy. These strategies may not be new to life, but certainly may be new to science simply because they are discreet compared to the physiological capabilities that blare in surface systems that have been well investigated. What evocative new approaches to survival are awaiting discovery?

- What is the relative importance of “latent” redox systems in subsurface environments where microbes cling to life (Valentine 2011)? An example is the radiolytic splitting of molecules to generate transient reactive species. Such metastable oxidants (e.g., peroxides, oxidized Mn, Fe, or S species) could allow incremental metabolic activity in systems that are otherwise deprived of oxidants (Chivian et al. 2008; D'Hondt et al. 2009). Another example is dehalorespiration, whereby chlorinated organics that are buried in the sediments may serve as oxidants (Futagami et al. 2009). The means by which these halogenated species (of human origin) serve as electron acceptors have been understood for years (Lee et al. 1998); however, we have not fully explored the process by which low concentrations of naturally occurring versions of these organics may be accessed by microbial communities in seafloor settings. Is the anoxic production of oxygen at the expense of methane and nitrite (Ettwig et al. 2010) common in subsurface systems given the correct chemistry and might this be another reasonable oxidant source for cells that do not require much? Collectively, are these just metabolic eddies and mere curios or bona fide survival strategies by which microbes can and do survive the subsurface in broad measure?
- Are there strategies that would appear to be thermodynamically insurmountable that have been solved by subsurface microbes? These capabilities might be analogues to the still incompletely understood anaerobic oxidation of methane, where life

manages to exist at the extreme edge of thermodynamic probability. For example, methanogens appear to exist, albeit not in high numbers, in sediment where methane levels are high enough to make additional methane production exceedingly difficult. Is it possible that hydrates may serve as a sink for additional methane production for proximal methanogens? Could questions like these be solved by cultivation studies that creatively determined how to cultivate seawater microbes that resisted normal laboratory media (Connon and Giovannoni 2002)?

- Does the subsurface have a “rare” biosphere just as was found in the surface oceans (Sogin et al. 2006)? If there are numerous rare taxa in samples from the subsurface then what does this say about functional resilience of the subsurface? Do keystone taxa exist in the subsurface; that is, microbes that are hallmarks of the subsurface and therefore playing some fundamental roles underground? The Census of Deep Life, currently underway and a part of the Deep Carbon Observatory, may inform us.

Subsurface coupling of the living and the non-living

We now understand new ways that abiotic and biotic systems of our planet are inextricably linked to each other and to human systems (Liu et al. 2007; Watkins and Freeman 2008; Stafford 2010). The subsurface is no different. As we learn more about the ways in which life survives underground, the activities and identities of these cells, questions arise related to how deep Earth changes life and, in reciprocity, is changed as a result of the life therein. What are the various connections between large-scale Earth processes and subsurface microbes that require thermodynamic disequilibria to conserve energy for metabolic activity, however slim that activity might be (Fig. 1)?

- Does the fluid movement associated with tidal forces (Tolstoy et al. 2002) influence subsurface microbial communities? And to include another large-scale phenomenon, if earthquakes influence tidal activity (Glasby and Kasahara 2001), how then are these seismic events tied to microbial community activity?
- We understand that hydrogen may be key to microbial survival in the subsurface (Morita 2000; Sleep et al. 2004). Does hydrogen production from seismic activity represent yet another way that microbes in the subsurface can be kept alive by Earth movements (Hirose et al. 2011)?
- Besides hydrogen, what other microbial provisions might be generated by seismic energy release? Can new space in the subsurface in the form of fractures and porosity, as well as access to oxidants from newly cracked surfaces, be supplied to deep microbes that are otherwise so limited in this regard?
- Are continental margins, where tectonics dictate the adjustment of plates and promote fluid movement (Torres et al. 2002; Wood et al. 2002) by opening fractures and by changing the stability of gas hydrates within the sediments, also places where blooms of subsurface microbial activity can be expected?
- How do annual planetary cycles determine what may occur in the subsurface? For example, does the seasonal accumulation and then melting of snow, which loads and unloads Earth’s surface in places like northeastern Japan and causes deformation at the land surface (Heki 2001), also simulate a subterranean bellows opening and closing on an annual cycle that microbes might take advantage of?
- Do global events with longer cyclic periods, such as the current change in climate at Earth’s surface, cause changes realized by life underground? In high latitudes, where warming processes appear to have provoked the incipient thaw of permafrost and degeneration of methane hydrates (cf., Shakhova et al. 2010; Ruppel 2011; Walter

Anthony et al. 2012), will changes eventually translate into the subsurface sediments and prompt microbial activity as a result of renewed fluid movement in long-frozen materials? How will the microbes in these systems, as they become more active as they can in more surficial Arctic settings (Mackelprang et al. 2011), impose themselves on the fluxes of greenhouse gases that we are so concerned with?

SUMMARY

As investigators of life underground, we anticipate the chance to learn more about the unseen world of small life deep in Earth, but only if we continue to engage in essential collaborations with all those who possess complementary knowledge of the planet's history and systems. Soon, we hope, there will be more complete synthetic models of how the living and non-living aspects of Earth's uppermost layers function and integrate with one another. These models will guide our search for new deep life, its niches, capabilities, and adaptations. We look forward to new explanations for how the subsurface biosphere is sustained and how its expression matters to life at the surface.

ACKNOWLEDGMENTS

We thank the Deep Carbon Observatory funded by the Alfred P. Sloan Foundation and the Department of Energy, Office of Science, Office of Biological and Environmental Research, Subsurface Biogeochemical Research Program (Award Number DE-SC0001533) for support. We greatly appreciate the editorial support provided by Sachi and Tomi Nakama.

REFERENCES

- Allen JP, Atekwana EA, Atekwana EA, Duris JW, Werkema DD, Rossbach S (2007) The microbial community structure in petroleum-contaminated sediments corresponds to geophysical signatures. *Appl Environ Microbiol* 73:2860-2870
- Amann R, Fuchs BM, Behrens S (2001) The identification of microorganisms by fluorescence *in situ* hybridisation. *Curr Opin Biotech* 12:231-236
- Amaral-Zettler L, Artigas LF, Baross J, Bharathi PAL, Boetius A, Chandramohan D, Herndl G, Kogure K, Neal P, Pedros-Alio C, Ramette A, Schouten S, Stal L, Thessen A, de Leeuw J, Sogin M (2010) A global census of marine microbes. *In: Life in the World's Oceans: Diversity, Distribution and Abundance*. McIntyre AD (ed) Wiley-Blackwell, Oxford, p 223-245
- Amy P, Haldeman D, Ringelberg D, Hall D, Russell C (1992) Comparison of identification systems for classification of bacteria isolated from water and endolithic habitats within the deep subsurface. *Appl Environ Microbiol* 58:3367-3373
- Amy PS, Haldeman DL (eds) (1997) *The Microbiology of the Terrestrial Deep Subsurface*. CRC Press, Boca Raton Florida
- Anderson RE, Brazelton WJ, Baross J (2011a) Is the genetic landscape of the deep subsurface biosphere affected by viruses? *Front Microbiol* 2:doi: 10.3389/fmicb.2011.00219
- Anderson RE, Brazelton WJ, Baross JA (2011b) Using CRISPRs as a metagenomic tool to identify microbial hosts of a diffuse flow hydrothermal vent viral assemblage. *FEMS Microbiol Lett* 77:120-133
- Anderson RE, Brazelton WJ, Baross JA (2013) The deep virosphere: assessing the viral impact on microbial community dynamics in the deep subsurface. *Rev Mineral Geochem* 75:649-675
- Archer D (2007) Methane hydrate stability and anthropogenic climate change. *Biogeosciences* 4:521-544
- Arrage AA, Phelps TJ, Benoit RE, White DC (1993) Survival of subsurface microorganisms exposed to UV radiation and hydrogen peroxide. *Appl Environ Microbiol* 59:3545-3550
- Askenazi M, Webber JT, Marto JA (2011) mzServer: web-based programmatic access for mass spectrometry data analysis. *Molecular Cell Proteomics* 10, doi: 10.1074/mcp.M110.003988
- Bach W, Edwards K (2003) Iron and sulfide oxidation within the basaltic ocean crust: implications for chemolithoautotrophic microbial biomass production. *Geochim Cosmochim Acta* 67:3871-3887
- Bakermans C, Tsapin AI, Souza-Egipsy V, Gilichinsky DA, Nealson KH (2003) Reproduction and metabolism at -10 °C of bacteria isolated from Siberian permafrost. *Environ Microbiol.* 5:321-326

- Balkwill D (1989) Numbers, diversity, and morphological characteristics of aerobic, chemoheterotrophic bacteria in deep subsurface sediments from a site in South Carolina. *Geomicrobiol J* 7:33-52
- Balkwill DL, Leach FR, Wilson JT, McNabb JF, White DC (1988) Equivalence of microbial biomass measures based on membrane lipid and cell wall components, adenosine triphosphate, and direct counts in subsurface aquifer sediments. *Microb Ecol* 16:73-84
- Banfield JF, Young M (2009) Variety - the splice of life - in microbial communities. *Science* 326:1198-1199
- Bastin ES, Greer FE, Merritt CA, Moulton G (1926) Bacteria in oil field waters. *Science* 63:21-24
- Batzke A, Engelen B, Sass H, Cypionka H (2007) Phylogenetic and physiological diversity of cultured deep-biosphere bacteria from Equatorial Pacific Ocean and Peru Margin sediments. *Geomicrobiol J* 24:261-273
- Beatty JT, Overmann J, Lince MT, Manske AK, Lang AS, Blankenship RE, Van Dover CL, Martinson TA, Plumley FG (2005) An obligately photosynthetic bacterial anaerobe from a deep-sea hydrothermal vent. *Proc Natl Acad Sci USA* 102:9306-9310
- Benson SM, Cook P (2005) Chapter 5: Underground geological storage. *In: IPCC Special Report on Carbon Dioxide Capture and Storage, Intergovernmental Panel on Climate Change, Interlachen, Switzerland, p 5-1 to 5-134*
- Bernardino AF, Levin LA, Thurber AR, Smith CR (2012) Comparative composition, diversity and trophic ecology of sediment macrofauna at vents, seeps and organic falls. *PLoS One* 7, doi: 10.1371/journal.pone.0033515
- Biddle JF, Lipp JS, Lever MA, Lloyd KG, Sorensen KB, Anderson R, Fredricks HF, Elvert M, Kelly TJ, Schrag DP, Sogin ML, Brenchley JE, Teske A, House CH, Hinrichs K-U (2006) Heterotrophic Archaea dominate sedimentary subsurface ecosystems off Peru. *Proc Natl Acad Sci USA* 103:3846-3851
- Blair CC, D'Hondt S, Spivack AJ, Kingsley RH (2007) Potential of radiolytic hydrogen for microbial respiration in subseafloor sediments. *Astrobiology* 7:951-970
- Borgonie G, Garcia-Moyano A, Lithauer D, Bert W, Bester A, van Heerden E, Moller C, Erasmus M, Onstott TC (2011) Nematoda from the terrestrial deep subsurface of South Africa. *Nature* 474:79-82
- Bowles MW, Samarkin VA, Joye SB (2011) Improved measurement of microbial activity in deep-sea sediments at *in situ* pressure and methane concentration. *Limnol Oceanogr Meth* 9:499-506
- Breitbart M, Felts B, Kelley S, Mahaffy JM, Nulton J, Salamon P, Rohwer F (2004) Diversity and population structure of a near-shore marine-sediment viral community. *Proc Royal Soc London Ser B Biol Sci* 271:565-574
- Briggs BR, Inagaki F, Morono Y, Futagami T, Huguet C, Rosell-Mele A, Lorenson TD, Colwell FS (2012) Bacterial dominance in subseafloor sediments characterized by methane hydrates. *FEMS Microbiol Ecol* 81:88-98
- Brockman FJ, Kieft TL, Fredrickson JK, Bjornstad BN, Li SW, Spangenburg W, Long PE (1992) Microbiology of vadose zone paleosols in south-central Washington State. *Microb Ecol* 23:279-301
- Brockman FJ, Murray CJ (1997) Microbiological heterogeneity in the terrestrial subsurface and approaches for its description. *In: The Microbiology of the Terrestrial Deep Subsurface*. Amy PS, Haldeman DL (eds) CRC Press, Boca Raton Florida, p 75-102
- Cano RJ, Borucki MK (1995) Revival and identification of bacterial spores in 25- to 40-million- year-old Dominican amber. *Science* 268:1060-1064
- Chapelle F, Lovley D (1990) Rates of microbial metabolism in deep coastal plain aquifers. *Appl Environ Microbiol* 56:1865-1874
- Chivian D, Brodie EL, Alm EJ, Culley DE, Dehal PS, DeSantis TZ, Gihring TM, Lapidus A, Lin L-H, Lowry SR, Moser DP, Richardson PM, Southam G, Wanger G, Pratt LM, Andersen GL, Hazen TC, Brockman FJ, Arkin AP, Onstott TC (2008) Environmental genomics reveals a single-species ecosystem deep within earth. *Science* 322:275-278
- Cockell CS, Voytek MA, Gronstal AL, Finster K, Kirshtein JD, Howard K, Reitner J, Gohn GS, Sanford WE, J.W. HJ, Kallmeyer J, Kelly L, Powars DS (2012) Impact disruption and recovery of the deep subsurface biosphere. *Astrobiology* 12:231-246
- Colwell F, G. Stormberg, T. Phelps, S. Birnbaum, J. McKinley, S. Rawson, C. Veverka, S. Goodwin, P. Long, B. Russell, T. Garland, D. Thompson, Skinner P, Grover S (1992) Innovative techniques for collection of saturated and unsaturated subsurface basalts and sediments for microbiological characterization. *J Microbiol Meth* 15:279-292
- Colwell F, Schwartz A, Briggs B (2011) Microbial community distribution in sediments from the Mount Elbert Gas Hydrate Stratigraphic Test Well, Alaska North Slope. *Mar Petrol Geol* 28:404-410
- Colwell FS, Lehman RM (1997) Carbon source utilization profiles for microbial communities from hydrologically distinct zones in a basalt aquifer. *Microb Ecol* 33:240-251
- Colwell FS, Onstott TC, Delwiche ME, Chandler D, Fredrickson JK, Yao Q-J, McKinley JP, Boone DR, Griffiths R, Phelps TJ, Ringelberg D, White DC, LaFreniere L, Balkwill D, Lehman RM, Konisky J, Long PE (1997) Microorganisms from deep, high temperature sandstones: constraints on microbial colonization. *FEMS Microbiol Rev* 20:425-435

- Connon SA, Giovannoni SJ (2002) High-throughput methods for culturing microorganisms in very-low-nutrient media yield diverse new marine isolates. *Appl Environ Microbiol* 68:3878-3885
- Cowen JP, Giovannoni SJ, Kenig F, Johnson HP, Butterfield D, Rappe MS, Hutnak M, Lam P (2003) Fluids from aging ocean crust that support microbial life. *Science* 299:120-123
- D'Hondt S, Inagaki F, Alvarez Zariqian CA, Expedition 329 Scientists (2011) Proceedings of the Integrated Ocean Drilling Program Management International, Inc. doi: 10.2204/iodp.proc.329.2011
- D'Hondt S, Jorgensen BB, Miller DJ, Batzke A, Blake R, Cragg BA, Cypionka H, Dickens GR, Ferdelman T, Hinrichs K-U, Holm NG, Mitterer R, Spivack A, Wang G, Bekins B, Engelen B, Ford K, Gettemy G, Rutherford SD, Sass H, Skilbeck CG, Aiello IW, Guerin G, House CH, Inagaki F, Meister P, Naehr T, Niituma S, Parkes RJ, Schippers A, Smith DC, Teske A, Wiegel J, Padilla CN, Acosta JLS (2004) Distributions of microbial activities in deep seafloor sediments. *Science* 306:2216-2221
- D'Hondt S, Rutherford S, Spivak AJ (2002a) Metabolic activity of subsurface life in deep-sea sediments. *Science* 295:2067-2070
- D'Hondt S, Smith DC, Spivak AJ (2002b) Exploration of the marine subsurface biosphere. *JOIDES J* 28:51-54
- D'Hondt S, Spivack AJ, Pockalny R, Ferdelman TG, Fischer JP, Kallmeyer J, Abrams LJ, Smith DC, Graham D, Hasiuk F, Schrum H, A.M. S (2009) Subseafloor sedimentary life in the South Pacific Gyre. *Proc Natl Acad Sci USA* 106:11651-11656
- Daily GC, Matson PA (2008) Ecosystem services: from theory to implementation. *Proc Natl Acad Sci USA* 105:9455-9456
- Danovaro R, Dell'Anno A, Corinaldesi C, Magagnini M, Noble R, Tamburini C, Weinbauer M (2008) Major viral impact on the functioning of benthic deep-sea ecosystems. *Nature* 454:1084-1087
- Das AP, Sukla LB, Pradhan N, Nayak S (2011) Manganese biomining: a review. *Bioresour Technol* 102:7381-7387
- Davis EE, Becker K, Pettigrew T, Carson B, MacDonald BR (1992) CORK: a hydrological seal and downhole observatory for deep-ocean boreholes. *Proc Ocean Drill Program Initial Rep* 139:43-53
- Davis KJ, Lutge A (2005) Quantifying the relationship between microbial attachment and mineral surface dynamics using vertical scanning interferometry (VSI). *Am J Sci* 305:727-751
- DeFlaun MF, Fuller ME, Zhang P, Johnson WP, Mailloux BJ, Holben WE, Kovacic WP, Balkwill DL, Onstott TC (2001) Comparison of methods for monitoring bacterial transport in the subsurface. *J Microbiol Meth* 47:219-231
- Delaney JR, Kelley DS, Lilley MD, Butterfield DA, Baross JA, Wilcock WSD, Embley RW, Summit M (1998) The quantum event of oceanic crustal accretion: impacts of diking at mid-ocean ridges. *Science* 281:222-230
- D'Elia T, Veerapaneni R, Rogers SO (2008) Isolation of microbes from Lake Vostok accretion ice. *Appl Environ Microbiol* 74:4962-4965
- Denef VJ, Kalnejais LH, Mueller RS, Wilmes P, Baker BJ, Thomas BC, VerBerkmoes NC, Hettich RL, Banfield JF (2010) Proteogenomic basis for ecological divergence of closely related bacteria in natural acidophilic microbial communities. *Proc Natl Acad Sci USA* 107:2383-2390
- Dickens GR (2003) Rethinking the global carbon cycle with a large, dynamic and microbially mediated gas hydrate capacitor. *Earth Planet Sci Lett* 213:169-183
- Dupraz S, Parmentier M, Ménez B, Guyot F (2009) Experimental and numerical modeling of bacterially induced pH increase and calcite precipitation in saline aquifers. *Chem Geol* 265:44-53
- Edwards KJ, Bach W, McCollom TM (2005) Geomicrobiology in oceanography: microbe - mineral interactions at and below the seafloor. *Trends Microbiol* 13:449-456
- Edwards KT, Becker K, Colwell F (2012) The deep, dark energy biosphere: intraterrestrial life on earth. *Annu Rev Earth Planet Sci* 40:551-568
- Edwards KT, Glazer BT, Rouxel OJ, Bach W, Emerson D, Davis RE, Toner BM, Chan CS, Tebo BM, Staudigel H, Moyer CL (2011) Ultra-diffuse hydrothermal venting supports Fe-oxidizing bacteria and massive umber deposition at 5000 m off Hawaii. *ISME J* 5:1748-1758
- Edwards RA, Rodriguez-Brito B, Wegley L, Haynes M, Breitbart M, Peterson DM, Saar MO, Alexander S, Alexander EC, Rohwer F (2006) Using pyrosequencing to shed light on deep mine microbial ecology under extreme hydrogeologic conditions. *BMC Genomics* 7:57
- Ehrlich HL (1990) *Geomicrobiology*. Marcel Dekker, New York
- Ehrlich HL (2012) Reminiscences from a career in geomicrobiology. *Ann Rev Earth Planet Sci* 40:1-21
- Ettwig KF, Butler MK, Le Paslier D, Pelletier E, Mangenot S, Kuypers MMM, Schreiber F, Dutilh BE, Zedelius J, de Beer D, Gloerich J, Wessels HJCT, van Alen T, Luesken F, Wu ML, van de Pas-Schoonen KT, Op den Camp HJM, Janssen-Megens EM, Francois K-J, Stunnenberg H, Weissenbach J, Jetten MSM, Strous M (2010) Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* 464:543-548
- Expedition 336 Scientists (2012) Mid-Atlantic Ridge microbiology: initiation of long-term coupled microbiological, geochemical, and hydrological experimentation within the seafloor at North Pond, western flank of the Mid-Atlantic Ridge. *IODP Preliminary Report 336*. doi: 10.2204/iodp.pr.336.2012

- Fisher AT, Tsuji T, Petronotis K, Expedition 327 Scientists (2011) Proceedings of the Integrated Ocean Drilling Program Management International, Inc. doi: 10.2204/iodp.proc.327.2011
- Fox P, Hendler J (2011) Changing the equation on scientific data visualization. *Science* 331:705-708
- Fredrickson JK, Balkwill D, Zachara J, Li S, Brockman F, Simmons M (1991) Physiological diversity and distributions of heterotrophic bacteria in deep Cretaceous sediments of the Atlantic coastal plain. *Appl Environ Microbiol* 57:402-411
- Fredrickson JK, Balkwill DL (2006) Geomicrobial processes and biodiversity in the deep terrestrial subsurface. *Geomicrobiol J* 23:345-356
- Fredrickson JK, Fletcher M (eds) (2001) *Subsurface Microbiology and Biogeochemistry*. Wiley-Liss, New York
- Fredrickson JK, McKinley JP, Bjornstad BN, Long PE, Ringelberg DB, White DC, Suflita JM, Krumholz L, Colwell FS, Lehman RM, Phelps TJ (1997) Pore-size constraints on the activity and survival of subsurface bacteria in a Late Cretaceous shale-sandstone sequence, northwestern, New Mexico. *Geomicrobiol J* 14:183-202
- Fredrickson JK, Onstott TC (1996) Microbes deep inside the Earth. *Sci Am* 275:68-73
- Fredrickson JK, Zachara JM, Balkwill DL, Kennedy D, Li SMW, Kostandarithes HM, Daly MJ, Romine MF, Brockman FJ (2004) Geomicrobiology of high-level nuclear waste-contaminated vadose sediments at the Hanford Site, Washington State. *Appl Environ Microbiol* 70:4230-4241
- Freifeld BM, Trautz RC, Kharaka YK, Phelps TJ, Myer LR, Hovorka SD, Collins DJ (2005) The U-tube: a novel system for acquiring borehole fluid samples from a deep geologic CO₂ sequestration experiment. *J Geophys Res-Solid Earth* 110, doi: 10.1029/2005JB003735
- Futagami T, Morono Y, Terada T, Kaksonen AH, Inagaki F (2009) Dehalogenation activities and distribution of reductive dehalogenase homologous genes in marine subsurface sediments. *Appl Environ Microbiol* 75:6905-6909
- Gaidos EJ, Neelson KH, Kirschvink JL (1999) Life in ice-covered oceans. *Science* 284:1631-1632
- Gerba CP, Bales RC (1990) Virus transport in the subsurface. First International Symposium on Microbiology of the Deep Subsurface, WSRC Information Services Section Publications Group, p 7-23 to 7-31
- Glasby GP, Kasahara J (2001) Influence of tidal effects on the periodicity of earthquake activity in diverse geological settings with particular emphasis on submarine hydrothermal systems. *Earth-Sci Rev* 52:261-297
- Gu G, Dickens GR, Bhatnagar G, Colwell FS, Hirasaki GJ, Chapman WG (2011) Abundant Early Palaeogene marine gas hydrates despite warm deep-ocean temperatures. *Nature Geosci* 4: 848-851
- Haggerty R, Schroth MH, Istok JD (1998) Simplified method of "push-pull" test data analysis for determining *in situ* reaction rate coefficients. *Ground Water* 36:314-324
- Hazen RM, Downs RT, Jones AP, Kah L (2013) Carbon mineralogy and crystal chemistry. *Rev Mineral Geochem* 75:7-46
- Hazen TC (1997) Bioremediation. *In: The Microbiology of the Terrestrial Deep Subsurface*. Amy PS, Haldeman DL (eds) CRC Press, Boca Raton Florida, p 247-266
- Hazen TC, Jimenez L, Lopez de Victoria G (1991) Comparison of bacteria from deep subsurface sediment and adjacent groundwater. *Microb Ecol* 22:293-304
- Head IM, Jones DM, Larter SR (2003) Biological activity in the deep subsurface and the origin of heavy oil. *Nature* 426:344-352
- Heider J (2007) Adding handles to unhandy substrates: anaerobic hydrocarbon activation mechanisms. *Curr Opin Chem Biol* 11:188-194
- Heider J, Spormann AM, Beller HR, Widdel F (1998) Anaerobic bacterial metabolism of hydrocarbons. *FEMS Microbiol Rev* 22:459-473
- Heki K (2001) Seasonal modulation of interseismic strain buildup in northeastern Japan driven by snow loads. *Science* 293:89-92
- Hirose T, Kawagucci S, Suzuki K (2011) Mechanoradical H₂ generation during simulated faulting: implications for an earthquake-driven subsurface biosphere. *Geophys Res Lett* 38, doi: 10.1029/2011GL048850
- Hirsch P, Rades-Rohlkohl E (1988) Some special problems in the determination of viable counts of groundwater microorganisms. *Microb Ecol* 16:99-113
- Holben WE, Ostrom PH (2000) Monitoring bacterial transport by stable isotope enrichment of cells. *Appl Environ Microbiol* 66:4935-4939
- Holman H-YN, Perry DL, Hunter-Cevera JC (1998) Surface-enhanced infrared absorption-reflectance (SEIRA) microspectroscopy for bacteria localization on geologic material surfaces. *J Microbiol Meth* 34:59-71
- Horn JM, Masterson BA, Rivera A, Miranda A, Davis MA, Martin S (2004) Bacterial growth dynamics, limiting factors, and community diversity in a proposed geological nuclear waste repository environment. *Geomicrobiol J* 21:273-286
- Inagaki F, Nunoura T, Nakagawa S, Teske A, Lever M, Lauer A, Suzuki M, Takai K, Delwiche M, Colwell FS, Neelson KH, Horikoshi K, D'Hondt S, Jørgensen BB (2006) Biogeographical distribution and diversity of microbes in methane hydrate-bearing deep marine sediments on the Pacific Ocean Margin. *Proc Natl Acad Sci USA* 103:2815-2820

- Janies D, Hill AW, Guralnick R, Habib F, Waltari E, Wheeler WC (2007) Genomic analysis and geographic visualization of the spread of avian influenza (H5N1). *Syst Biol* 56:321-329
- Jin Q, Bethke CM (2005) Predicting the rate of microbial respiration in geochemical environments. *Geochim Cosmochim Acta* 69:1133-1143
- Jolley DM, Ehrhorn TF, Horn J (2003) Microbial impacts to the near-field environment geochemistry: a model for estimating microbial communities in repository drifts at Yucca Mountain. *J Contam Hydrol* 62-3:553-575
- Jordana R, Baquero E, Reboleira S, Sendra A (2012) Reviews of the genera *Schaefferia* Absolon, 1900, *Deuteraphorura* Absolon, 1901, *Plutomurus* Yosii, 1956 and the *Anurida* Laboulbène, 1865 species group without eyes, with the description of four new species of cave springtails (Collembola) from Krubera-Voronya cave, Arabika Massif, Abkhazia. *Terrest Arth Rev* 5:35-85
- Jørgensen BB (2011) Deep seafloor microbial cells on physiological standby. *Proc Natl Acad Sci USA* 108:18193-18194
- Kallmeyer J, Pockalny R, Adhikari RR, Smith DC, D'Hondt S (2012) Global distribution of microbial abundance and biomass in seafloor sediment. *Proc Natl Acad Sci USA* doi: 10.1073/pnas.1203849109
- Kappler A, Emerson D, Edwards KJ, Amend JP, Galnick JA, Grathwohl P, Hoehler TM, Straub KL (2005) Microbial activity in biogeochemical gradients - new aspects of research. *Geobiol* 3:229-233
- Kelley DS, Karson JA, Früh-Green GL, Yoerger DA, Butterfield DA, Hayes J, Shank T, Schrenk MO, Olson EJ, Proskurowski G, Jakuba M, Bradley A, Larson B, Ludwig K, Glickson D, Buckman K, Bradley AS, Brazelton WJ, Roe K, Elend MJ, Delacour A, Bernasconi SM, Lilley MD, Baross JA, Summons RE, Sylva SP (2005) A serpentinite-hosted submarine ecosystem: the Lost City Hydrothermal Field. *Science* 307:1428-1434
- Kidd DM (2010) Point of view: geophylogenies and the map of life. *Syst Biol* 59:741-752
- Kidd DM, Liu X (2008) GEOPHYLOBUILDER 1.0: an ARCGIS extension for creating 'geophylogenies'. *Molec Ecol Resour* 8:88-91
- Kieft TL, Murphy EM, Haldeman DL, Amy PS, Bjornstad BN, McDonald EV, Ringelberg DB, White DC, Stair J, Griffiths RP, Gsell TC, Holben WE, Boone DR (1998) Microbial transport, survival, and succession in a sequence of buried sediments. *Microb Ecol* 36:336-348
- Kieft TL, Phelps TJ, Fredrickson JK (2007) Drilling, coring, and sampling subsurface environments. *In: Manual of Environmental Microbiology*. Vol 3. Hurst CJ, Crawford RL, Garland JL, Lipson DA, Mills AL, Stetzenbach LD (eds) ASM Press, Washington, DC, p 799-817
- King EL, Tuncay K, Ortoleva P, Meile C (2009) In silico *Geobacter sulfurreducens* metabolism and its representation in reactive transport models. *Appl Environ Microbiol* 75:83-92
- Krumholz LR, McKinley JP, Ulrich FA, Suffita JM (1997) Confined subsurface microbial communities in Cretaceous rock. *Nature* 386:64-66
- Lee MD, Odom JM, Buchanan RJ (1998) New perspectives on microbial dehalogenation of chlorinated solvents: insights from the field. *Ann Rev Microbiol* 52:423-452
- Lehman RM (2007) Understanding of aquifer microbiology is tightly linked to sampling approaches. *Geomicrobiol J* 24:331-341
- Lehman RM, Colwell FS, Bala GA (2001) Attached and unattached microbial communities in a simulated basalt aquifer under fracture- and porous-flow conditions. *Appl Environ Microbiol* 67:2799-2809
- Lehman RM, Colwell FS, Ringelberg D, White DC (1995) Combined microbial community-level analyses for quality assurance of terrestrial subsurface cores. *J Microbiol Meth* 22:263-281
- Lehman RM, O'Connell SP, Banta A, Fredrickson JK, Reysenbach A-L, Kieft TL, Colwell FS (2004) Microbiological comparison of core and groundwater samples collected from a fractured basalt aquifer with that of dialysis chambers incubated *in situ*. *Geomicrobiology J* 21:169-182
- Li L, Steefel CI, Kowalsky MB, Englert A, Hubbard SS (2010) Effects of physical and geochemical heterogeneities on mineral transformation and biomass accumulation during biostimulation experiments at Rifle, Colorado. *J Contam Hydrol* 112:45-63
- Lin L-H, Wang P-L, Rumble D, Lippmann-Pipke J, Boice E, Pratt LM, Lollar BS, Brodie EL, Hazen TC, Andersen GL, DeSantis TZ, Moser DP, Kershaw D, Onstott TC (2006) Long-term sustainability of a high-energy, low-diversity crustal biome. *Science* 314:479-482
- Lipp JS, Morono Y, Inagaki F, Hinrichs K-U (2008) Significant contribution of Archaea to extant biomass in marine subsurface sediments. *Nature* 454:991-994
- Liu J, Dietz T, Carpenter SR, Alberti M, Folke C, Moran E, Pell AN, Deadman P, Kratz T, Lubchenco J, Ostrom E, Ouyang Z, Provencher W, Redman CL, Schneider SH, Taylor WW (2007) Complexity of coupled human and natural systems. *Science* 317:1513-1516
- Lomstein BA, Langerhuus AT, D'Hondt S, Jørgensen BB, Spivack AJ (2012) Endospore abundance, microbial growth and necromass turnover in deep sub-seafloor sediment. *Nature* 484:101-104
- Mackelprang R, Waldrop MP, DeAngelis KM, David MM, Chavarria KL, Blazewicz SJ, Rubin EM, Jansson JK (2011) Metagenomic analysis of a permafrost microbial community reveals a rapid response to thaw. *Nature* 480:368-371

- Mason OU, Nakagawa T, Rosner M, Van Nostrand JD, Zhou J, Maruyama A, Fisk MR, Giovannoni SJ (2010) First investigation of the microbiology of the deepest layer of ocean crust. *PLoS ONE* 5, doi: 15310.11371/journal.pone.0015399
- Masui N, Morono Y, Inagaki F (2008) Microbiological assessment of circulation mud fluids during the first operation of riser drilling by the deep-earth research vessel Chikyu. *Geomicrobiol J* 25:274-282
- Matthess G (1990) Hydrogeological controls of bacterial and virus migration in subsurface environments. First International Symposium on Microbiology of the Deep Subsurface, WSRC Information Services Section Publications Group, p 7-33 to 7-46
- Maucalire L, Zepp K, Meister P, McKenzie J (2004) Direct *in situ* detection of cells in deep-sea sediment cores from the Peru Margin (ODP Leg 201, Site 1229). *Geobiology* 2:217-223
- McKinley JP, Stevens TO, Fredrickson JK, Zachara JM, Colwell FS, Wagnon KB, Smith SC, Rawson SA, Bjornstad BN (1997) Biogeochemistry of anaerobic lacustrine and paleosol sediments within an aerobic unconfined aquifer. *Geomicrobiol J* 14:23-39
- Meersman F, Daniel I, Bartlett DH, Winter R, Hazael R, McMillain PF (2013) High-pressure biochemistry and biophysics. *Rev Mineral Geochem* 75:607-648
- Menez B, Pasini V, Brnelli D (2012) Life in the hydrated suboceanic mantle. *Nat Geosci* 5:133-137
- Mikucki JA, Priscu JC (2007) Bacterial diversity associated with Blood Falls, a subglacial outflow from the Taylor Glacier, Antarctica. *Appl Environ Microbiol* 73:4029-4039
- Mitchell AC, Phillips AJ, Hamilton MA, Gerlach R, Hollis WK, Kaszuba JP, Cunningham AB (2008) Resilience of planktonic and biofilm cultures to supercritical CO₂. *J Supercrit Fluids* 47:318-325
- Mitchell AC, Phillips AJ, Hiebert R, Gerlach R, Spangler LH, Cunningham A (2009) Biofilm enhanced geologic sequestration of supercritical CO₂. *Int J Greenhouse Gas Control* 3:90-99
- Moran K, Farrington S, Massion E, Paull C, Stephen R, Trehu A, Ussler III W (2006) SCIMPI: A New Seafloor Observatory System. *OCEANS 2006*, doi: 10.1109/OCEANS.2006.307103
- Morita RY (2000) Is H₂ the universal energy source for long-term survival? *Microb Ecol* 38:307-320
- Morono Y, Takeshi T, Nishizawa M, Ito M, Hillion F, Takahata N, Sano Y, Inagaki F (2011) Carbon and nitrogen assimilation in deep seafloor microbial cells. *Proc Natl Acad Sci USA* 108:18295-18300
- Morozova D, Wandrey M, Alawi M, Zimmer M, Vieth A, Zettlitzer M, Wurdemann H (2010) Monitoring of the microbial community composition in saline aquifers during CO₂ storage by fluorescence *in situ* hybridisation. *Int J Greenhouse Gas Control* 4:981-989
- Murphy E, Schramke J, Fredrickson J, Bledsoe H, Francis A, Sklarew D, Linehan J (1992) The influence of microbial activity and sedimentary organic carbon on the isotope geochemistry of the Middendorf Aquifer. *Water Resour Res* 28:723-740
- Nazina TN, Kosareva IM, Petrunyaka VV, Savushkina MK, Kudriavtsev EG, Lebedev VA, Ahunov VD, Revenko YA, Khafizov RR, Osipov GA, Belyaev SS, Ivanov MV (2004) Microbiology of formation waters from the deep repository of liquid radioactive wastes Severnyi. *FEMS Microbiology Ecology* 49:97-107
- Nealson KH, Inagaki F, Takai K (2005) Hydrogen-driven subsurface lithoautotrophic microbial ecosystems (SLiMEs): do they exist and why should we care? *Trends Microbiol* 13:405-410
- Nielsen LP, Risgaard-Petersen N, Fossing H, Christensen PB, Sayama M (2010) Electric currents couple spatially separated biogeochemical processes in marine sediment. *Nature* 463:1071-1074
- Nielsen ME, Fisk MR, Istok JD, Pedersen K (2006) Microbial nitrate respiration of lactate at *in situ* conditions in ground water from a granitic aquifer situated 450 m underground. *Geobiology* 4:43-52
- Novarino G, Warren A, Butler H, Lambourne G, Boxshall A, Bateman J, Kinner NE, Harvey RW, Mosse RA, Teltch B (1997) Protistan communities in aquifers: a review. *FEMS Microbiol Rev* 20:261-275
- O'Donnell AG, Young IM, Rushton SP, Shirley MD, Crawford JW (2007) Visualization, modelling and prediction in soil microbiology. *Nat Rev Microbiol* 5:689-699
- Onstott TC (2004) Impact of CO₂ injections on deep subsurface microbial ecosystems and potential ramifications for the surface biosphere. *In: The CO₂ Capture and Storage Project (CCP)* Thomas DC, Benson SM (eds) Lawrence Berkeley National Laboratory, Berkeley, CA, p 1207-1239
- Onstott TC, Colwell FS, Kieft TL, Murdoch L, Phelps TJ (2009a) New horizons for deep subsurface microbiology. *Microbe* 4:499-505
- Onstott TC, McGown DJ, Bakermans C, Ruskeeniemi T, Ahonen L, Telling J, Soffientino B, Pffiffer SM, Sherwood Lollar B, Frappe S, Stotler R, Johnson EJ, Vishnivetskaya TA, Rothmel R, Pratt LM (2009b) Microbial communities in subpermafrost saline fracture water at the Lupin Au Mine, Nunavut, Canada. *Microb Ecol* 58:786-807
- Onstott TC, Phelps TJ, Colwell FS, Ringelberg D, White DC, Boone DR, McKinley JP, Stevens TO, Long PE, Balkwill DL, Griffin WT, Kieft T (1998) Observations pertaining to the origin and ecology of microorganisms recovered from the deep subsurface of Taylorsville Basin, Virginia. *Geomicrobiol J* 15:353-385
- Onstott TC, Phelps TJ, Kieft T, Colwell FS, Balkwill DL, Fredrickson JK, Brockman F (1999) A global perspective on the microbial abundance and activity in the deep subsurface. *In: Enigmatic Microorganisms and Life in Extreme Environments*, Seckbach J (ed) Kluwer Academic Publishers, Netherlands, p 487-500

- Onstott TC, Tobin K, Dong H, DeFlaun M, Fredrickson J, Bailey T, Brockman F, Kieft T, Peacock A, White DC, Balkwill D, Phelps TJ, Boone DR (1997) The deep gold mines of South Africa: windows into the subsurface biosphere. *Proc. SPIE, International Society for Optical Engineering* 3111:344–357
- Orcutt BN, Wheat CG, Edwards KJ (2010) Subseafloor ocean crust microbial observatories: development of FLOCS (Flow-through Osmo Colonization System) and evaluation of borehole construction materials. *Geomicrobiol J* 27:143–157
- Parkes RJ, Cragg BA, Bale SJ, Getliff JM, Goodman K, Rochelle PA, Fry JC, Weightman AJ, Harvey SM (1994) Deep bacterial biosphere in Pacific Ocean sediments. *Nature* 371:410–413
- Parkes RJ, Cragg BA, Wellsbury P (2000) Recent studies on bacterial populations and processes in subseafloor sediments: a review. *Hydrogeol J* 8:11–28
- Parkes RJ, Sellek G, Webster G, Martin D, Anders E, Weightman AJ, Sass H (2009) Culturable prokaryotic diversity of deep, gas hydrate sediments: first use of a continuous high-pressure, anaerobic, enrichment and isolation system for subseafloor sediments (DeepIsoBUG). *Environ Microbiol* 11:3140–3153
- Pedersen K (1997) Microbial life in deep granitic rock. *FEMS Microbiol Rev* 20:399–414
- Pedersen K (1999) Subterranean microorganisms and radioactive waste disposal in Sweden. *Eng Geol* 52:163–176
- Pham VD, Hnatow LL, Zhang S, Fallon RD, Jackson SC, Tomb J-F, DeLong EF, Keeler SJ (2009) Characterizing microbial diversity in production water from an Alaskan mesothermic petroleum reservoir with two independent methods. *Environ Microbiol* 11:176–187
- Phelps TJ, Murphy EM, Pflfner SM, White DC (1994) Comparison between geochemical and biological estimates of subsurface microbial activities. *Microb Ecol* 28:335–349
- Pitonzo BJ, Amy PS, Rudin M (1999) Effect of gamma radiation on native endolithic microorganisms from a radioactive waste deposit site. *Radiation Res* 152:64–70
- Pombo SA, Kleikemper J, Schroth MH, Zeyer J (2005) Field-scale isotopic labeling of phospholipid fatty acids from acetate-degrading sulfate-reducing bacteria. *FEMS Microbiol Ecol* 51:197–207
- Price PB, Sowers T (2004) Temperature dependence of metabolic rates for microbial growth, maintenance, and survival. *Proc Natl Acad Sci USA* 101:4631–4636
- Radajewski S, McDonald IR, Murrell JC (2003) Stable-isotope probing of nucleic acids: a window to the function of uncultured microorganisms. *Curr Opin Biotechnol* 14:296–302
- Rastogi G, Stetler LD, Peyton BM, Sani RK (2009) Molecular analysis of prokaryotic diversity in the deep subsurface of the former Homestake gold mine, South Dakota, USA. *J Microbiol* 47:371–384
- Rawlings DE (2002) Heavy metal mining using microbes. *Annu Rev Microbiol* 56:65–91
- Reagan MT, Moridis GJ (2009) Large-scale simulation of methane hydrate dissociation along the West Spitsbergen Margin. *Geophys Res Lett* 36, doi: 10.1029/2009gl041332
- Revil A, Mendonça CA, Atekwan EA, Kulesa B, Hubbard SS, Bohlen KJ (2010) Understanding biogeobatteries: where geophysics meets microbiology. *J Geophys Res* 115, doi: 10.1029/2009JG001065
- Riley RG, Zachara JM, Wobber FJ (1992) Chemical Contaminants on DOE Lands and Selection of Contaminant Mixtures for Subsurface Science Research. US Department of Energy, Office of Energy Research, Subsurface Science Program, Washington, DC
- Rockstrom J, Steffen W, Noone K, Persson A, Chapin FS, Lambin EF, Lenton TM, Scheffer M, Folke C, Schellnhuber HJ, Nykvist B, de Wit CA, Hughes T, van der Leeuw S, Rodhe H, Sorlin S, Snyder PK, Costanza R, Svedin U, Falkenmark M, Karlberg L, Corell RW, Fabry VJ, Hansen J, Walker B, Liverman D, Richardson K, Crutzen P, Foley JA (2009) A safe operating space for humanity. *Nature* 461:472–475
- Røy H, Kallmeyer J, Adhikari RR, Pockalny R, Jorgensen BB, D'Hondt S (2012) Aerobic microbial respiration in 86-million-year-old deep-sea red clay. *Science* 336:922–925
- Ruppel CD (2011) Methane hydrates and contemporary climate change. *Nature Education Knowledge* 3(10): 29
- Russell CE, Jacobson R, Haldeman DL, Amy PS (1994) Heterogeneity of deep subsurface microorganisms and correlations to hydrogeological and geochemical parameters. *Geomicrobiol J* 12:37–51
- Santelli CM, Orcutt BN, Banning E, Bach W, Moyer CL, Sogin ML, Staudigel H, Edwards KJ (2008) Abundance and diversity of microbial life in ocean crust. *Nature* 453:653–656
- Satterfield CL, Lowenstein TK, Vreeland RH, Rosenzweig WD, Powers DW (2005) New evidence for 250 Ma age of halotolerant bacterium from a Permian salt crystal. *Geology* 33:265–268
- Saunders AD (2005) Large igneous provinces: origin and environmental consequences. *Elements* 1:259–263
- Scheibe TD, Mahadevan R, Fang Y, Garg S, Long PE, Lovley DR (2009) Coupling a genome-scale metabolic model with a reactive transport model to describe *in situ* uranium bioremediation. *Microb Biotechnol* 2:274–286
- Schippers A, Neretin LN, Kallmeyer J, Ferdelman TG, Cragg BA, John Parkes R, Jorgensen BB (2005) Prokaryotic cells of the deep sub-seafloor biosphere identified as living bacteria. *Nature* 433:861–864
- Schrenk MO, Brazelton WJ, Lang SQ (2013) Serpentinization, carbon, and deep life. *Rev Mineral Geochem* 75:575–606
- Schrenk MO, Huber JA, Edwards KJ (2010) Microbial provinces in the subseafloor. *Ann Rev Mar Sci* 2:279–304

- Shakhova N, Semiletov I, Salyuk A, Yusupov V, Kosmach D, Gustafsson O (2010) Extensive methane venting to the atmosphere from sediments of the East Siberian Arctic Shelf. *Science* 327:1246-1250
- Sidlauskas B, Ganapathy G, Hazkani-Covo E, Jenkins KP, Lapp H, McCall LW, Price S, Scherle R, Spaeth PA, Kidd DM (2009) Linking the big: the continuing promise of evolutionary synthesis. *Evolution*, doi: 10.1111/j.1558-5646.2009.00892.x
- Sinclair J, Kampbell D, Cook M, Wilson J (1993) Protozoa in subsurface sediments from sites contaminated with aviation gasoline or jet fuel. *Appl Environ Microbiol* 59:467-472
- Sinclair JL, Ghiorse WC (1989) Distribution of aerobic bacteria, protozoa, algae, and fungi in deep subsurface sediments. *Geomicrobiol J* 7:15-31
- Sleep NH, Meibom A, Fridriksson T, Coleman RG, Bird DK (2004) H₂-rich fluids from serpentinization: geochemical and biotic implications. *Proc Natl Acad Sci USA* 101:12818-12823
- Smith A, Popa R, Fisk MR, Nielsen M, Wheat CG, Jannasch HW, Fisher AT, Becker K, Sievert M, Flores G (2011) *In situ* enrichment of ocean crust microbes on igneous minerals and glasses using an osmotic flow-through device. *Geochem Geophys Geosyst* 12, doi: 10.1029/2010GC003424
- Smith DC, D'Hondt S (2006) Exploration of life in deep seafloor sediments. *Oceanography* 19:58-70
- Smith DC, Spivak AJ, Fisk MR, Haveman SA, Staudigel H (2000) Tracer-based estimates of drilling-induced microbial contamination of deep sea crust. *Geomicrobiol J* 17:207-219
- Smith RL, Harvey RW, LeBlanc DR (1991) Importance of closely spaced vertical sampling in delineating chemical and microbiological gradients in groundwater studies. *J Contam Hydrol* 7:285-300
- Sogin ML, Morrison HG, Huber JA, Welch DM, Huse SM, Neal PR, Arrieta JM, Herndl GJ (2006) Microbial diversity in the deep sea and the underexplored "rare biosphere". *Proc Natl Acad Sci USA* 103:12115-12120
- Solomon EA, Kastner M, MacDonald IR, Leifer I (2009) Considerable methane fluxes to the atmosphere from hydrocarbon seeps in the Gulf of Mexico. *Nature Geosci* 2, doi: 10.1038/ngeo574
- Spalding BP, Watson DB (2006) Measurement of dissolved H₂, O₂, and CO₂ in groundwater using passive samplers for gas chromatographic analyses. *Environ Sci Technol* 40:7861-7867
- Spear JR, Walker JJ, McCollom TM, Pace NR (2005) Hydrogen and bioenergetics in the Yellowstone geothermal ecosystem. *Proc Natl Acad Sci USA* 102:2555-2560
- Stafford SG (2010) Environmental science at the tipping point. *BioScience* 60:94-95
- Steeffel CI, DePaolo DJ, Lichtner PC (2005) Reactive transport modeling: an essential tool and a new research approach for the Earth sciences. *Earth Planet Sci Lett* 240:539-558
- Stein LY, Nicol GW (2011) Grand challenges in terrestrial microbiology. *Front Microbiol* 2, doi: 10.3389/fmicb.2011.00006
- Stepanauskas R, Sieracki ME (2007) Matching phylogeny and metabolism in the uncultured marine bacteria, one cell at a time. *Proc Natl Acad Sci USA* 104:9052-9057
- Stevens TO (1997) Subsurface microbiology and the evolution of the biosphere. *In: The Microbiology of the Terrestrial Deep Subsurface*. Amy PS, Haldeman DL (eds) CRC Press, Boca Raton Florida, p 205-223
- Stevens TO, McKinley JP (1995) Lithoautotrophic microbial ecosystems in deep basalt aquifers. *Science* 270:450-454
- Stevens TO, McKinley JP, Fredrickson JK (1993) Bacteria associated with deep, alkaline, anaerobic groundwaters in southeast Washington. *Microb Ecol* 25:35-50
- Stevenson BS, Eichorst SA, Wertz JT, Schmidt TM, Breznak JA (2004) New strategies for cultivation and detection of previously uncultured microbes. *Appl Environ Microbiol* 70:4748-4755
- Stroes-Gascoyne S, West JM (1997) Microbial studies in the Canadian nuclear fuel waste management program. *FEMS Microbiol Rev* 20:573-590
- Suttle CA (2005) Viruses in the sea. *Nature* 437:356-361
- Szalay A, Gray J (2001) The world-wide telescope. *Science* 293:2037-2040
- Takai K, Moser DP, Onstott TC, Spoelstra N, Pfiffner SM, Dohnalkova A, Fredrickson JK (2001) *Alkaliphilus transvaalensis* gen. nov., sp. nov., an extremely alkaliphilic bacterium isolated from a deep South African gold mine. *Int J Syst Evol Microbiol* 51:1245-1256
- Takai K, Mottl MJ, Nielsen SH, Expedition 331 Scientists (2011) Proceedings of the Integrated Ocean Drilling Program Management International, Inc. doi: 10.2204/iodp.proc.331.2011
- Takai K, Nakamura K, Toki T, Tsunogai U, Miyazaki M, Miyazaki J, Hirayama H, Nakagawa S, Nunoura T, Horikoshi K (2008) Cell proliferation at 122 °C and isotopically heavy CH₄ production by a hyperthermophilic methanogen under high-pressure cultivation. *Proc Natl Acad Sci USA* 105:10949-10954
- Thierstein HR, Störrlein U (1991) Living bacteria in Antarctic sediments from Leg 119. *In: Scientific Results, Ocean Drilling Program, Leg 119*. Barron J, Larsen B (eds) Ocean Drilling Program, College Station, TX, p 687-692
- Thurber AR, Levin LA, Orphan VJ, Marlow JJ (2012) Archaea in metazoan diets: implications for food webs and biogeochemical cycling. *ISME J* 6:602-612

- Tittensor DP, Mora C, Jetz W, Lotze HK, Ricard D, Vanden Berghe E, Worm B (2010) Global patterns and predictors of marine biodiversity across taxa. *Nature* 466:1098-1101
- Tolstoy M, Vernon FL, Orcutt JA, Wyatt FK (2002) Breathing of the seafloor: tidal correlations of seismicity at Axial volcano. *Geology* 30:503-506
- Torres ME, McManus J, Hammond DE, de Angelis MA, Heeschen KU, Colbert SL, Tryon MD, Brown KM, Suess E (2002) Fluid and chemical fluxes in and out of sediments hosting methane hydrate deposits on Hydrate Ridge, OR, I: hydrological provinces. *Earth Planet Sci Lett* 201:525-540
- Tyson GW, Chapman J, Hugenholtz P, Allen EE, Ram RJ, Richardson PM, Solovyev VV, Rubin EM, Rokhsar DS, Banfield JF (2004) Community structure and metabolism through reconstruction of microbial genomes from the environment. *Nature* 428:37-43
- Tyson GW, Lo I, Baker BJ, Allen EE, Hugenholtz P, Banfield JF (2005) Genome-Directed isolation of the key nitrogen fixer *Leptospirillum ferrodiazotrophum* sp. nov. from an acidophilic microbial community. *Appl Environ Microbiol* 71:6319-6324
- Urmann K, Gonzalez-Gil G, Schroth MH, Hofer M, Zeyer J (2005) New field method: gas push-pull test for the *in situ* quantification of microbial activities in the vadose zone. *Environ Sci Technol* 39:304-310
- Valentine DL (2011) Emerging topics in marine methane biogeochemistry. *Annu Rev Mar Sci* 3:147-171
- Vorobyova E, Soina V, Gorlenko M, Minkovskaya N, Zalinova N, Mamukelashvili A, Gilichinsky D, Rivkina E, Vishnivetskaya T (1997) The deep cold biosphere: facts and hypotheses. *FEMS Microbiol Rev* 20:277-290
- Vreeland RH, Rosenzweig WD, Powers DW (2000) Isolation of a 250 million-year-old halotolerant bacterium from a primary salt crystal. *Nature* 407:897-900
- Walter Anthony KM, Anthony P, Grosse G, Chanton J (2012) Geologic methane seeps along boundaries of Arctic permafrost thaw and melting glaciers. *Nature Geosci* 5:419-426
- Warren LA, Maurice PA, Parmar N, Ferris FG (2001) Microbially mediated calcium carbonate precipitation: implications for interpreting calcite precipitation and for solid-phase capture of inorganic contaminants. *Geomicrobiol J* 18:93-115
- Watkins NW, Freeman MP (2008) GEOSCIENCE: natural complexity. *Science* 320:323-324
- Westbrook GA, Thatcher KE, Rohling EJ, Piotrowski AM, Palike H, Osborne AH, Nisbet EG, Minshull TA, Lanoiselle M, Huhnerbach V, Green D, Fisher RE, Crocker AJ, Chabert A, Bolton C, Beszczynska-Moller A, Berndt C, Aquilina A (2009) Escape of methane gas from the seabed along the West Spitsbergen continental margin. *Geophys Res Lett* 36, doi: 10.1029/2009GL039191
- Whelan JK, Oremland R, Tarafa M, Smith R, Howarth R, Lee C (1986) Evidence for sulfate-reducing and methane-producing microorganisms in sediments from Sites 618, 619, 622. *In: Scientific Results, Ocean Drilling Program, Leg 119*. Bouma AH, Coleman JM, Meyer AW (eds) Ocean Drilling Program, College Station, TX, doi: 10.2973/dsdp.proc.96.147.1986
- Whitman WB, Coleman DC, Wiebe WJ (1998) Prokaryotes: the unseen majority. *Proc Natl Acad Sci USA* 95:6578-6583
- Wiggins B, Alexander, M. (1985) Minimum bacterial density for bacteriophage replication: implications for significance of bacteriophages in natural ecosystems. *Appl Environ Microbiol* 49:19-23
- Wood WT, Gettrust JF, Chapman NR, Spence GD, Hyndman RD (2002) Decreased stability of methane hydrates in marine sediments owing to phase-boundary roughness. *Nature* 420:656-660
- Wright DJ, Wang S (2011) The emergence of spatial cyberinfrastructure. *Proc Natl Acad Sci USA* 108:5488-5491
- Yates M, Thompson S, Jury W (1997) Sorption of viruses during flow through saturated sand columns. *Environ Sci Technol* 31:548-555
- Zhang CL, Lehman RM, Pfiffner SM, Scarborough SP, Palumbo AV, Phelps TJ, Beauchamp JJ, Colwell FS (1997) Spatial and temporal variations of microbial properties at different scales in shallow subsurface sediments. *Appl Biochem Biotechnol* 63-5:797-808
- Zhuang K, Izallalen M, Mouser P, Richter H, Risso C, Mahadevan R, Lovley DR (2011) Genome-scale dynamic modeling of the competition between *Rhodospirillum rubrum* and *Geobacter* in anoxic subsurface environments. *ISME J* 5:305-316
- Zobell CE, Anderson DQ (1936) Vertical distribution of bacteria in marine sediments. *Bull Am Assoc Petrol Geol* 20:258-269