

nblS, a Gene Involved in Controlling Photosynthesis-Related Gene Expression during High Light and Nutrient Stress in *Synechococcus elongatus* PCC 7942

Lorraine G. van Waasbergen,^{1*} Nadia Dolganov,² and Arthur R. Grossman³

Department of Biology, University of Texas at Arlington, Arlington, Texas 76019,¹ and Department of Microbiology and Immunology, Stanford Medical School, Stanford University,² and Department of Plant Biology, The Carnegie Institution of Washington,³ Stanford, California 94305

Received 29 October 2001/Accepted 3 February 2002

The HliA protein of the cyanobacterium *Synechococcus elongatus* PCC 7942 is a small, thylakoid-associated protein that appears to play a role in photoprotection; its transcript rapidly accumulates in response to high-intensity light (HL) and the *hli* gene family is required for survival of cells in high light. In order to discover regulatory factors involved in HL acclimation in cyanobacteria, a screen was performed for chemically generated mutants unable to properly control expression of the *hliA* gene in response to HL. One such mutant was identified, and complementation analysis led to the identification of the affected gene, designated *nblS*. Based on its deduced protein sequence, NblS appears to be a membrane-bound, PAS-domain-bearing, sensor histidine kinase of two-component regulatory systems in bacteria. The *nblS* mutant was unable to properly control light intensity-mediated expression of several other photosynthesis-related genes, including all three *psbA* genes and the *cpcBA* genes. The mutant was also unable to control expression of the *hliA* and *psbA* genes in response to low-intensity blue/UV-A light, a response that may be related to the HL-mediated regulation of the genes. Additionally, in response to nutrient deprivation, the *nblS* mutant was unable to properly control accumulation of the *nblA* transcript and associated degradation of the light-harvesting phycobilisomes. The *nblS* mutant dies more rapidly than wild-type cells following exposure to HL or nutrient deprivation, likely due to its inability to properly acclimate to these stress conditions. Thus, the NblS protein is involved in the control of a number of processes critical for altering the photosynthetic apparatus in response to both HL and nutrient stress conditions.

The photosynthetic apparatus is a dynamic assemblage of activities that are strongly controlled by the environment. Environmental parameters such as light quality, light intensity, temperature, water availability, and nutrient status play critical roles in determining the activities of photosynthetic complexes and the levels of pigments and proteins associated with those complexes (17, 22). It is vital that organisms tune photosynthesis to balance the energetic requirements of the cell with the light energy absorbed by antenna pigments and the transfer of that energy to photosynthetic reaction centers. Such coordination helps prevent oxidative damage caused by overreduction of electron carriers and the accumulation of excited pigment molecules. In the short term, photosynthetic activity in response to excess absorbed light can be modulated by redistribution of excitation energy utilization between the photosystems and the quenching of excess excitation energy within antenna complexes (7, 10, 17, 22, 35). Long-term modulations involve alterations in the composition of the photosynthetic apparatus or changes in processes that modulate the assembly or disassembly of specific protein complexes that function in photosynthesis (7, 10, 17, 35). Long-term acclimation processes may reflect altered patterns of gene expression, which may be triggered by changes in light quality (through the activity of

specific photoreceptors) or changes in light intensity (through the activity of redox-sensitive regulators) (2, 15, 20, 28, 31, 32, 39, 49).

Cyanobacteria possess several mechanisms for modifying the composition of the photosynthetic machinery with respect to environmental conditions. One dramatic example of this modulation is the degradation of the light-harvesting pigment-protein complex, the phycobilisome (PBS), during starvation for sulfur and nitrogen (13, 14, 52). The loss of PBS and a reduction in the level of chlorophyll per cell during nutrient-limited growth changes the appearance of cells from a deep blue-green to a chlorotic yellow, a process termed bleaching. Degradation of the PBS may provide some of the limiting nutrient and serve to reduce the light absorbed by the photosynthetic apparatus during starvation conditions, when the use of excitation energy for anabolic processes would markedly decline. One protein that appears to trigger PBS degradation during nutrient deprivation is NblA (14). The level of *nblA* transcript, which increases dramatically upon starvation of *Synechococcus elongatus* PCC 7942 for nitrogen or sulfur (14), is controlled by NblR (43). This response regulator controls activities that are critical for cell viability under a variety of stress conditions, including nutrient deprivation and high-intensity light (HL) stress.

A well-characterized response of both plants and cyanobacteria to HL is the turnover and exchange of the D1 subunits of photosystem II, the primary site of damage during photoinhibition (10). In *S. elongatus* PCC 7942, there are two forms of D1 that are encoded by three genes (21). The

* Corresponding author. Mailing address: Department of Biology, Box 19498, University of Texas at Arlington, Arlington, TX 76019. Phone: (817) 272-5239. FAX: (817) 272-2855. E-mail: Lorivw@uta.edu.

psbAI gene encodes D1, form I, which is most abundant under low light (LL) conditions. The *psbAII* and *psbAIII* genes each encode D1, form II, which becomes abundant when the cells are shifted from LL to HL, a consequence of the induction of the *psbAII* and *psbAIII* genes and a decrease in *psbAI* transcript levels (9). D1, form II, may provide cells with greater resistance to photooxidative damage (12). Light quality may provide part of the signal for light intensity regulation; blue light causes an increase in the accumulation of *psbAII* and *psbAIII* transcripts and a decrease in the level of the *psbAI* transcript (50).

Like the *psbAII* and *psbAIII* genes, expression of the *hliA* gene of *S. elongatus* PCC 7942 increases during exposure of the organism to HL or blue/UV-A light (19). The HliA protein of *S. elongatus* PCC 7942 has similarity to the extended chlorophyll *a/b*-binding family of proteins, especially to the early light-inducible proteins (ELIPs) of vascular plants, a carotenoid-binding protein (Cbr) of the alga *Dunaliella*, and analogous, single membrane-spanning helix proteins of other cyanobacteria (including a family of four *hli* genes present in *Synechocystis* sp. strain PCC 6803 [23]). Similar to the ELIPs, Hli polypeptides are localized in the thylakoid membrane and accumulate following exposure to HL or blue/UV-A light (19, 23). The Hli polypeptides are critical for survival in HL (23) and may be involved in the dissipation of excess absorbed excitation energy (M. Havaux, G. Guedeny, Q. He, and A. R. Grossman, unpublished data).

This paper describes the identification of an apparent sensor histidine kinase, NblS, that is involved in controlling *hliA* expression in response to HL. NblS was also found to be involved in the expression of *nblA* and degradation of the PBS during nutrient deprivation. In addition, it was found to be involved in the control of a number of other photosynthesis-related genes during exposure to HL and blue/UV-A light. Our results suggest that NblS functions as an important regulator controlling the acclimation of the photosynthetic apparatus to stress conditions.

MATERIALS AND METHODS

Culture conditions. Culturing of *Synechococcus elongatus* PCC 7942 (formerly known as *Anacystis nidulans* R2 or *Synechococcus* sp. strain PCC 7942 [24]) was as previously reported (30). Antibiotics were added when appropriate (see reference 18). Cells were grown at 50 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ incandescent light. The level of HL was 800 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ unless otherwise indicated. The level of UV-A light from black-light blue bulbs (see reference 19) was 27 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$. Prior to HL or UV-A treatment, cultures were grown to an optical density at 750 nm (OD_{750}) of approximately 1.0, diluted to an OD_{750} of 0.2 with fresh medium, and incubated for 18 h in LL at 10 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ (unless otherwise indicated). Starvation for sulfur or nitrogen was performed as previously reported (43).

Mutagenesis and screening for *hliA* regulatory mutants. The parental strain for mutagenesis was *S. elongatus* PCC 7942 hliNG (18). This strain carries the *hliA* promoter fused to both a promoterless *nblA* gene (14) (on an autonomously replicating plasmid) and a promoterless *uidA* gene (β -glucuronidase [GUS]) (incorporated into the chromosome at a neutral site). Since the presence of the NblA polypeptide in cyanobacterial cells triggers the bleaching process, a strain harboring the *nblA* gene fused to the *hliA* promoter will turn from a normal blue-green appearance to yellow upon exposure to conditions that activate the *hliA* promoter. Thus, the *nblA* reporter gene in the hliNG strain serves as a direct visual screen for *hliA* promoter activity. *S. elongatus* PCC 7942 hliNG was mutagenized with *N*-methyl-*N'*-nitrosoguanidine (18), and the mutagenized cells were screened on plates for reduced bleaching following a 12-h exposure to 500 $\mu\text{mol photon m}^{-2} \text{s}^{-1}$. Putative mutants that exhibited reduced bleaching in HL were quantified for *hliA* activity during HL exposure by assaying GUS activity as

previously reported (19). From a series of screens involving approximately 50,000 colonies, 11 isolates showed reduced bleaching and GUS activity in HL but exhibited normal bleaching under nutrient deprivation conditions and appeared to have lesions in the *hliA* promoter-*nblA* fusion. Four isolates had reduced bleaching but normal GUS activity under HL, also had reduced bleaching under nutrient deprivation conditions, and were complemented by a DNA fragment bearing the *nblB* gene, which encodes a polypeptide necessary for degradation of the PBS (18). A single mutant, designated *nblS-1*, was identified that exhibited both reduced bleaching and lowered GUS activity in HL. This mutant also was found to be nonbleaching under nutrient deprivation conditions (see below).

Complementation of the *nblS-1* mutant and characterization of the *nblS* gene. Molecular techniques were performed using standard protocols (42). The *nblS-1* mutant was transformed (30) with a library of *Sau3AI* partially digested chromosomal DNA of *S. elongatus* PCC 7942 in pUC118 (43), and transformants were selected for ampicillin resistance, which results when a plasmid integrates into the chromosome by single homologous recombination. Because *hliA* promoter-driven *nblA* expression was reduced but not completely absent in the *nblS-1* mutant, a significant level of bleaching occurred when the mutant was incubated overnight in HL. To more easily identify complemented strains, LL-adapted transformant colonies were screened for GUS activity after 3 h in HL by overspraying with the highly sensitive fluorogenic GUS substrate 4-methylumbelliferyl β -D-glucuronide (10 mg/ml dissolved in dimethyl sulfoxide) and visualizing fluorescence under long-wavelength UV light. Chromosomal DNA was extracted from one of the complemented strains, and the genomic regions flanking the inserted plasmid DNA were rescued as previously reported (43). One rescued plasmid contained a 4.5-kbp *PstI* segment of genomic DNA that was able to complement the *nblS-1* mutant. Subclones of this fragment were used in complementation tests to define the complementing gene. The 4.5-kbp *PstI* fragment was sequenced by using the Applied Biosystems PRISM system (Perkin-Elmer).

RNA isolation and RNA blot hybridizations. The wild-type and *nblS-1* strains used for RNA blot hybridizations and viability assays were the hliNG strain and the *nblS-1* mutant, respectively, after being cured of the *hliA-nblA* fusion-bearing plasmid by several passages through liquid medium devoid of antibiotics. The *nblR* mutant used for RNA blot analysis was the *nblRQ* strain, in which the *nblR* gene was disrupted by insertion of a kanamycin resistance gene (43). Following the various treatments, RNA was isolated from cells (6). For all RNA blot hybridizations, equal amounts of RNA (determined spectrophotocally) were resolved by electrophoresis in formaldehyde gels; ethidium bromide included in the loading buffer allowed visualization of the rRNA bands and confirmed equal loading of the RNA samples. Moreover, on one blot, equal loading of RNA samples based on ethidium bromide staining was confirmed by equal signals obtained by probing the RNA blot with an RNase P gene-specific probe as a control (see reference 41). Gene-specific probes were prepared as described for *hliA* (19), *nblA* (14), and *psbAI*, *psbAII*, and *psbAIII* (8). A 321-bp PCR-amplified fragment of the *cpcBA* operon (starting 27 bp upstream of the initial ATG of *cpcB* and extending into the coding region) was cloned into pGEM-T Easy (Promega). This plasmid served as the template for the preparation of an antisense RNA probe specific for transcripts bearing the *cpcB* gene. A 287-bp PCR-amplified fragment of *glnA* (starting 5 bp downstream of the initial ATG and extending into the coding region) was used to generate a DNA probe specific for that gene.

Nucleotide sequence accession number. The sequence of the 4.5-kbp *PstI* fragment identified in this study was deposited in the GenBank database (accession number AF299076).

RESULTS

Characterization of a mutant with aberrant regulation of *hliA* and discovery of its aberrant response to nutrient deprivation. To identify regulatory factors involved in light intensity control of gene expression in cyanobacteria, we screened for chemically generated mutants unable to properly activate transcription from the *hliA* promoter in HL. One such mutant identified was *nblS-1*. Figure 1A shows activity from the *hliA* promoter in HL quantified in strains bearing the *hliA* promoter fused to the GUS gene. Wild-type cells developed high levels of GUS activity following exposure to HL, showing a rapid increase in activity during the initial 2 h of exposure, followed by a slower rise; the *nblS-1* mutant exhibited a significantly




FIG. 1. Characterization of expression of the *hliA* gene in the *nblS-1* mutant. (A) Measurement of *hliA* promoter-driven GUS activity (phliA::GUS) during HL exposure of the hliNG wild-type (WT) strain and the *nblS-1* mutant. (B) RNA blot hybridization of an *hliA*-specific probe to RNA isolated from the wild-type or *nblS-1* strain grown for various times in HL or blue/UV-A light. For comparison, total RNA was stained with ethidium bromide (panel B, bottom).

lower level of activity, with the initial rapid rise in activity being absent but with the slower rise still apparent.

The kinetics of HL-triggered accumulation of *hliA* mRNA in the *nblS-1* mutant and wild-type strains agreed with the results observed using the GUS assay (Fig. 1B). A rapid increase in the level of *hliA* mRNA in wild-type cells in HL was followed by a gradual decrease. Similar kinetics of *hliA* transcript accumulation in the wild-type strain occurred in UV-A light (Fig. 1B); however, the peak at 30 min was followed by a more rapid decrease than that seen in HL, such that after 3.5 h of UV-A light there was almost no detectable *hliA* transcript. (This rapid decrease was not due to a general decrease in transcription during UV-A exposure, since a transcript for the *glnA* gene was actually more abundant following 3.5 h of UV-A light than at 30 min [data not shown].) In contrast, accumulation of *hliA* mRNA in the *nblS-1* mutant following exposure to either HL or UV-A light was much lower than that in wild-type cells; in the mutant, accumulation of *hliA* mRNA was apparent only following 3.5 h of exposure to HL and no accumulation was observed in UV-A light.

In addition to exhibiting aberrant regulation of *hliA*, the *nblS-1* mutant showed an abnormal response to nutrient deprivation. The mutant exhibited very little of the loss of pigmentation (bleaching) that is typically found in wild-type cells during nutrient deprivation (Fig. 2A, compare lanes +, -, S, and




FIG. 2. Characterization of the response to nutrient deprivation in the *nblS-1* mutant. (A) Liquid medium cultures of wild-type cells (WT) and the *nblS-1* mutant after 72 h in complete medium (+) or medium lacking sulfur (-S) or nitrogen (-N). (B) RNA blot hybridization of a riboprobe specific for *nblA* to RNA isolated from the wild-type or *nblS-1* strain after 24 h in complete medium or medium lacking sulfur or nitrogen. (C) RNA blot hybridization of a probe specific for *glnA* to RNA isolated from the wild-type or *nblS-1* strain grown in complete medium or in medium lacking nitrogen for 30 min. For comparison within each RNA blot, total RNA was stained with ethidium bromide (panels B and C, bottom).

-N); the name given to the mutant, *nblS* (nonbleaching sensor), reflects the fact that a mutation strongly affects the pathway that controls the bleaching process in *S. elongatus* PCC 7942 and that the gene altered in the strain encodes a putative sensor protein (see below). The finding that the *nblS-1* mutant did not bleach during nutrient deprivation suggested that this strain might be deficient in the expression of *nblA*, since this gene is induced during nutrient limitation and is required for bleaching to occur (14). This was confirmed by RNA blot hybridizations (Fig. 2B). On the other hand, genes critical for the uptake and assimilation of specific nutrients following nutrient deprivation were still properly regulated in the *nblS-1* mutant. For example, as seen in the wild-type cells, nitrogen deprivation of the *nblS-1* mutant resulted in an increased accumulation of the *glnA* transcript (Fig. 2C).

Since a mutant in the NblR response regulator that controls




FIG. 3. Viability of the *nblS-1* strain during stress conditions. (A) Growth, as measured by OD₇₅₀, of wild-type (WT) and *nblS-1* strains in HL. (B) Wild-type (WT) and *nblS-1* strains were spotted onto nutrient-replete solid medium following growth in liquid medium lacking sulfur (-S) or nitrogen (-N). Each spot consisted of 10 μ l from a twofold dilution of the original culture.

expression of *nblA* is unable to properly acclimate to stress conditions and dies upon exposure to HL or nutrient stress, we examined the *nblS-1* mutant for its viability under stress conditions. Similar to the *nblR* mutant, the *nblS-1* strain exhibited a marked decline in growth rate relative to that of wild-type cells in HL in liquid medium (Fig. 3A), with more cell death apparent in the mutant than in the wild-type strain. (After 6 h in HL, the *nblS-1* mutant culture contained 3.75×10^7 CFU/ml per OD₇₅₀ of 1.0, while the wild-type culture contained 1.17×10^8 CFU/ml per OD₇₅₀ of 1.0; prior to HL, the cultures contained 1.60×10^8 CFU/ml per OD₇₅₀ of 1.0 and 1.64×10^8 CFU/ml per OD₇₅₀ of 1.0 for the mutant and wild-type strains, respectively.) Cells from the mutant strain also died more rapidly than wild-type cells during nutrient deprivation (Fig. 3B).

Expression of other HL-regulated genes in the *nblS-1* mutant. We examined expression of a number of HL-regulated genes in the *nblS-1* mutant (Fig. 4). The *nblS-1* mutant was aberrant with regard to the control of all three *psbA* genes. The mutant strain exhibited little increase in *psbAII* and *psbAIII* transcript levels and no decrease in *psbAI* transcript level following exposure to HL or UV-A irradiation. In fact, the levels of *psbAI* transcript in the mutant increased in response to UV-A exposure. Furthermore, we found in wild-type *S. elongatus* PCC 7942 that the *cpcBA* transcripts (i.e., those encoding the phycocyanin subunits, which constitute the major polypeptides of the light-harvesting PBS) declined to very low levels in HL. Such a decrease in phycocyanin levels would reduce the excitation energy absorbed by the cells during HL. However, in the *nblS-1* strain there was little such decrease, with *cpcBA* transcript levels somewhat lower under LL and higher under HL relative to those of the wild-type strain (Fig. 4). Similarities in the light regulation of *hliA* and these other HL-regulated genes make it extremely likely that the altered expression of

these genes in the mutant was the result of the same lesion that affects *hliA* expression.

The *nblS* gene encodes a PAS domain-bearing putative sensor kinase. To identify the gene altered in *nblS-1*, the mutant was transformed with a wild-type recombinant library, and transformants were screened for complementation of the defect in *hliA* expression in HL as described in Materials and Methods. A 4.5-kbp *PstI* fragment isolated from one of the complemented strains restored *hliA* expression in HL. The entire fragment was sequenced, and the complementing region was localized (by using subcloning and complementation analyses) to a 664-amino acid open reading frame (ORF) that we designated NblS (Fig. 5). Another complete ORF (Orf 1) and a partial ORF (Orf 3) were found flanking NblS on the 4.5-kbp *PstI* fragment. Orf 1 has high sequence similarity to the family of phosphoribosylglycinamide formyltransferases (an enzyme involved in de novo purine biosynthesis), while Orf 3 shows high sequence similarity to the family of 5'-methylthioadenosine and purine-nucleoside phosphorylases (involved in purine salvage). It is not known whether these two ORFs share any functional relationship with NblS.

Analysis of the predicted NblS protein sequence showed that it shares strong sequence similarity with the superfamily of sensor histidine kinases of two-component regulatory systems (36, 44). It can be divided into several domains based on sequence similarity to other proteins in the databases and to characterized functional motifs (Fig. 6). The C terminus of the protein contains a typical histidine kinase transmitter module with N, G1, F, and G2 domains (37). The N-terminal putative sensory input region contains two predicted membrane-spanning regions (TM1 and TM2) followed by a region, recently termed the HAMP domain (3), with similarity to the linker region of methyl-accepting chemotaxis proteins and certain histidine kinases. HAMP domains may




FIG. 4. RNA blot analyses of light intensity-regulated genes in the wild-type and *nblS-1* strains. LL-acclimated cells were exposed for 30 min to either HL or blue/UV-A light as described in Materials and Methods. Under the usual LL conditions ($10 \mu\text{mol photon m}^{-2} \text{s}^{-1}$), *psbAI* was not transcribed significantly. Therefore, LL conditions of $50 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ were used for RNA blot analyses of *psbAI* expression. For comparison within each RNA blot, total RNA was stained with ethidium bromide (bottom panels).

be involved in intramolecular interactions that control activation of the polypeptides in which they are found (3). The HAMP domain of NblS is followed by a region that bears strong similarity to PAS domains. PAS domains are involved

in protein-protein interactions and the binding of redox-active cofactors that modulate protein output activity (48, 54).

The *nblS* gene from the *nblS-1* mutant was PCR amplified




FIG. 5. A physical map of the 4.5-kbp region carried by the integrative plasmid that complemented the *nblS-1* mutant and fragments of this plasmid used in subcloning analysis. Also indicated are the relative locations of the two complete ORFs and one partial ORF that were discovered upon DNA sequence analysis. Indicated by asterisks on the NblS ORF are the locations of the two amino acid changes generated by the two point mutations present in the *nblS-1* allele. To the right of each DNA fragment is listed the percentage of transformants bearing the integrated plasmid carrying the particular fragment that complemented *hliA* activity in the *nblS-1* mutant. This percentage is not always 100% for those fragments that did not bear a complete *nblS* gene because, depending upon the location of crossover relative to the mutations in the *nblS-1* allele, a complete, wild-type *nblS* gene was not always generated in the chromosome. The subcloning analyses localize the complementing region to the *nblS* gene. (The *ClaI* sites shown in parentheses are those sites at which the DNA was not cleaved during subcloning analyses due to methylation in the *Dam*⁺ *Escherichia coli*.)




FIG. 6. Domain structure of NblS. (A) Diagrammatic representation of the NblS domain structure; (B) sequence of the NblS protein annotated with the domain structure. The two predicted transmembrane regions (TM1 and TM2), the HAMP region, and the PAS domain are shown. Also indicated are motifs characteristic of a C-terminal kinase transmitter domain (from residue 400 to the end of the protein), which are highly conserved among histidine kinases (H, N, G1, F, and G2). Indicated in boldface italic and with asterisks in panel B are the two amino acid changes (G-379 to D and I-620 to V) present in the *nblS-1* mutant allele. (C) Alignment of the PAS domain with homologous regions from other proteins. Shown are PAS domains from the DspA protein (sll0698) from *Synechocystis* sp. strain PCC 6803 (4); the YCF26 protein from a red algal chloroplast (40); the ResE (45) and the PhoR (27) proteins, both from *Bacillus subtilis*; the FixL protein from *Rhizobium meliloti* (16); the ArcB protein from *E. coli* (26); and the photoactive-yellow protein (PYP) from *Halochromatium salexigens* (29). The consensus sequence from a published alignment of PAS domains (54) is also shown (U, bulky hydrophobic: FILMVWY; h, hydrophobic: ILMV). Residues conserved among 70% or more of the sequences are shaded in black; residues conserved among 50% or more are shaded in gray.

and sequenced. Although mutagenesis using nitrosoguanidine was performed at a rate that should generate a single mutation per cell, the gene was found to contain two mutations, both of which caused amino acid substitutions (described in the legend to Fig. 6B). This could be due to the fact that nitrosoguanidine

mutagenesis commonly acts at replication forks (34), resulting in multiple mutations within a cell that are close to one another.

Several characterized histidine kinase-like proteins have high overall sequence similarity to NblS. The highest sequence

similarities are between NblS and DspA of *Synechocystis* sp. strain PCC 6803 (58% identity overall, 54% identity within the input domain) and putative proteins encoded by the chloroplast genome of some red algae (49% identity overall, 28% identity within the input domain) (4, 40). The *dspA* gene product of *Synechocystis* sp. strain PCC 6803 was identified as a factor that confers resistance to a variety of herbicides and calmodulin antagonists (and was hypothesized to play a role in cell permeability) (4) and was later identified in a study as being involved in the response to chilling in *Synechocystis* sp. strain PCC 6803 (therein called Hik33) (46, 47). NblS also has high sequence similarity to ResE (45) and PhoR (27) of *Bacillus subtilis* (25% identity and 23% identity, respectively, overall; 19% identity and 14% identity, respectively, within the input domain). ResE is a global regulator controlling the expression of genes involved in aerobic and anaerobic respiratory growth (apparently by monitoring an intracellular signal, perhaps the reduced state of menaquinone, in controlling gene expression [33]), while PhoR is involved in controlling gene expression in response to phosphorus deprivation. (NblS also shares high sequence identity with an ResE homolog, SrrB, in *Staphylococcus aureus* that is involved in the global control of virulence factors in response to oxygen conditions [53].) Like NblS, all of the aforementioned proteins have TM, HAMP, and PAS domains. Figure 6C shows the PAS domain from NblS aligned with PAS domains from closely related, putative sensor proteins as well as with some of the more thoroughly characterized PAS domains.

We have attempted to generate a strain in which the *nblS* gene has been inactivated by insertion of an antibiotic cassette (data not shown). Despite repeated subculturing in LL, PCR analysis demonstrated that wild-type copies of the chromosome persist in this strain. The same result was seen for the *Synechocystis* sp. strain PCC 6803 *hik33* mutant strain (47). It may be that some amount of NblS activity is essential for cell viability. Nevertheless, the 4.5-kbp *PstI* fragment bearing *nblS* was able to complement the *nblS-1* mutant for the defects in HL-, UV-A-, and nutrient stress-mediated gene expression as well as in stress-related growth (data not shown). This indicates that the defects are due to the mutations in *nblS* and not to mutations elsewhere in the chromosome.

Are nutrient stress- and HL-regulated genes controlled through the same or different signal transduction pathways? The NblR response regulator controls transcription of *nblA* and the bleaching of the cyanobacterial cells during nutrient stress (43). To determine if NblR is also important for controlling expression of the light-responsive genes, we examined the effect of light intensity on the levels of *hliA*, *psbAI*, *psbAII*, *psbAIII*, and *cpcBA* transcripts following exposure of an *nblR* mutant to HL. As shown in Fig. 7, the *nblR* mutant still exhibits HL induction of *hliA*, *psbAII*, and *psbAIII* transcripts and HL depression of *psbAI* and *cpcBA* transcripts. These results suggest that the light-regulated expression of these photosynthetic genes is not controlled through NblR. Surprisingly, in the *nblR* mutant, transcripts from *hliA*, *psbA*, and *cpcBA* genes consistently accumulate to slightly higher levels than in wild-type cells under inducing conditions, and detectable levels of *psbAIII* and *psbAI* mRNA exist in the *nblR* mutant under noninducing conditions (Fig. 7).

DISCUSSION

Bacteria have evolved diverse ways for sensing and responding to changes in environmental conditions. One common mechanism for the perception and transduction of signals in bacteria is the two-component regulatory system (36, 44). This type of system typically involves a pair of proteins, namely, a histidine kinase that acts as a sensor of a signal and an associated response regulator that often is involved in binding of DNA and the regulation of gene expression. The study described here has identified, in a cyanobacterium, a putative sensor kinase, NblS, that is involved in controlling the expression of a variety of genes involved in photosystem function during HL and nutrient deprivation. NblS contains the appropriate motifs to function as a histidine kinase (including the conserved histidine residue), but it remains to be shown whether it actually functions as such at the biochemical level. We have not positively identified any response regulator(s) that interact with NblS. No gene with similarity to response-regulatory molecules was found contiguous to *nblS* on the genome (Fig. 5). One obvious candidate response regulator that may interact with NblS during nutrient limitation is NblR, since both NblS and NblR are involved in controlling *nblA* expression and PBS degradation. However, our data indicate that NblR is not directly involved in the regulation of the light-responsive genes apparently controlled by NblS. Higher-than-normal transcript levels from the light-responsive genes were observed in the *nblR* mutant under inducing conditions (and, in some cases, detectable levels were observed under noninducing conditions) (Fig. 7). One way this could be explained is to propose that, during changes in light intensity, NblS interacts with NblR (but not to control the light intensity-regulated genes analyzed here) as well as with one or more other response regulators. Thus, in the absence of NblR, more NblS would be free to interact with another response regulator(s), which would lead to the observed augmentation of NblS-dependent (but not NblR-dependent) light intensity-mediated responses. Of course, this hypothesis remains to be tested.

We have found that NblS is involved in controlling the expression of a number of different genes whose products are associated with photosynthesis during HL and nutrient stress. Cyanobacteria experience a variety of changes in the photosynthetic apparatus and its activities in response to HL and generalized nutrient deprivation (5, 13, 14, 25). *nblS-1* mutant cells were found to die more rapidly than wild-type cells during exposure to HL and nutrient limitation. The loss in viability is likely a consequence of the inability of the cells to properly control expression of genes that are necessary for acclimation to these stress conditions (including genes whose products are involved in photosynthesis such as those identified here). Interestingly, although NblR did not appear to control the light-regulated genes analyzed here (and, indeed, is not known to control any genes other than *nblA* during nutrient stress), a similar loss in viability during both nutrient deprivation and HL has been observed for an *nblR* mutant (43). This suggests that, in addition to modulating expression of *nblA* under nutrient stress, in HL NblR is likely to control the expression of genes other than those studied here that are important to survival in HL.




FIG. 7. RNA blot analysis of light-intensity controlled genes in the wild-type strain and a strain in which the *nblR* gene was inactivated. RNA was extracted from cells grown in LL and exposed to HL (as described in the Fig. 5 legend) for 30 min (*psbAII* and *psbAI*) or 2 h (*hliA*, *psbAIII*, and *cpcBA*). For comparisons within each RNA blot, total RNA was stained with ethidium bromide (bottom panels).

Input signals that control potential sensor kinases such as NblS are often difficult to specifically define. At this time, we do not know what signals modulate NblS action and how such signals specifically alter the HL and nutrient stress responses. Examination of the sequence of the potential sensory input region of NblS may give clues as to how NblS could be acting as a sensor. Since PAS domains in a number of proteins appear to be involved in redox sensing (48), it is possible that the PAS domain of NblS is involved in sensing changes in photosynthetic or cellular redox during HL and nutrient stress. Absorption of light in excess of that which can be used in photosynthesis can lead to overexcitation of the photosynthetic apparatus and the generation of damaging oxygen radicals. Such an excess of light can occur during periods of HL exposure or during periods of lower light when the organism is experiencing stresses, such as nutrient limitation, that reduce anabolic processes in the cell (17). The presence of two predicted TM domains in NblS makes it likely that the protein is anchored in the cytoplasmic or thylakoid membrane. In this location it could interact with photosynthesis and/or respiration. Another potential signal input region of NblS is the area between the two TM domains, which may be located either in the periplasm or in the thylakoid lumen. Indeed, HAMP domains, such as that found in NblS, have been proposed to be linkers involved in signal transduction, transmitting conforma-

tional changes from the periplasmic region (caused by the binding of specific ligands) to the cytoplasmic regions of the molecules (3, 51). It remains to be demonstrated what role each of the domains of NblS plays in the activity of the molecule.

Our results demonstrate that NblS is in some manner involved in blue/UV-A light-mediated gene expression. Several genes in plants and cyanobacteria are responsive both to HL and to relatively low levels of blue/UV-A light (levels that would not dramatically alter cellular redox conditions) (e.g., genes encoding ELIPs [1], the *hliA* gene [19], and the *psbA* genes [50]). This suggests that a blue/UV-A photoreceptor is involved in controlling gene expression during HL acclimation of photosynthetic organisms. It is not clear what role, if any, NblS plays in actual blue/UV-A photosensing. Since some PAS domains are involved in sensing of blue light (11, 38), it is possible that NblS itself functions as a blue light photoreceptor; however, it is also possible that NblS is only indirectly involved in integrating the blue light signal and that some other system is actually responsible for perception of blue/UV-A light.

Recently, the apparent *Synechocystis* sp. strain PCC 6803 NblS homolog, DspA (4) (Hik33 [47]), has been shown to be involved in controlling gene expression (including expression of photosynthesis-related genes) in response to low tempera-

ture conditions (46, 47). Chilling is another condition that, like nutrient deprivation, reduces the use of photosynthate by the cell, which could result in hyperreduction of the photosynthetic electron transport chain and the accumulation of reactive oxygen species (17). This adds support to the hypothesized role of NblS as a global regulator in these cyanobacteria that integrates redox and light signals and suggests that NblS may influence other signaling pathways involved in acclimation responses.

ACKNOWLEDGMENTS

The authors thank R. Schwarz, J. Christie, and W. Briggs for their helpful suggestions during the course of this work.

This work was supported in part by National Science Foundation Grant MCB 9727836 and U.S. Department of Agriculture grants 97-35301-4575 and 98-35301-6445 awarded to A.R.G., U.S.-Israel Binational Science Foundation Grant 9800146 awarded to A.R.G. and R. Schwarz, and National Science Foundation Postdoctoral Fellowship DBI 9750329 awarded to L.G.V.W.

REFERENCES

- Adamska, I., K. Kloppstech, and I. Ohad. 1992. UV light stress induces the synthesis of the early light-inducible protein and prevents its degradation. *J. Biol. Chem.* **267**:24732–24737.
- Anderson, J. M. 1986. Photoregulation of the composition, function, and structure of thylakoid membranes. *Annu. Rev. Plant Physiol.* **37**:93–136.
- Aravind, L., and C. P. Ponting. 1999. The cytoplasmic helical linker domain of receptor histidine kinase and methyl-accepting proteins is common to many prokaryotic signalling proteins. *FEMS Microbiol. Lett.* **176**:111–116.
- Bartsevich, V. V., and S. V. Shestakov. 1995. The *dspA* gene product of the cyanobacterium *Synechocystis* sp. strain PCC 6803 influences sensitivity to chemically different growth inhibitors and has amino acid similarity to histidine protein kinases. *Microbiology* **141**:2915–2920.
- Bhaya, D., R. Schwarz, and A. R. Grossman. 2000. Molecular responses to environmental stress, p. 397–442. In B. A. Whitton and M. Potts (ed.), *Ecology of cyanobacteria: diversity in time and space*. Kluwer Publishers, Dordrecht, The Netherlands.
- Bhaya, D., N. Watanabe, T. Ogawa, and A. R. Grossman. 1999. The role of an alternative sigma factor in motility and pilus formation in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Proc. Natl. Acad. Sci. USA* **96**:3188–3193.
- Bjorkman, O., and B. Demmig-Adams. 1994. Regulation of photosynthetic light energy capture, conversion, and dissipation in leaves of higher plants, p. 17–47. In E.-D. Schulze and M. M. Calwell (ed.), *Ecophysiology of photosynthesis*. Springer, Berlin, Germany.
- Brusslan, J., and R. Haselkorn. 1989. Resistance to the photosystem II herbicide diuron is dominant to sensitivity in the cyanobacterium *Synechococcus* sp. PCC7942. *EMBO J.* **8**:1237–1245.
- Bustos, S. A., M. R. Schaefer, and S. S. Golden. 1990. Different and rapid responses of four cyanobacterial *psbA* transcripts to changes in light intensity. *J. Bacteriol.* **172**:1998–2004.
- Chow, W. S. 1994. Photoprotection and photoinhibitory damage, p. 151–196. In J. Barber (ed.), *Advances in molecular and cell biology*, vol. 10. JAI Press, Greenwich, Conn.
- Christie, J. M., M. Salomon, K. Nozue, M. Wada, and W. R. Briggs. 1999. LOV (light, oxygen, or voltage) domains of the blue-light photoreceptor phototropin (nph1): binding sites for the chromophore flavin mononucleotide. *Proc. Natl. Acad. Sci. USA* **96**:8779–8783.
- Clarke, A. K., V. M. Hurry, P. Gustafsson, and G. Oquist. 1993. Rapid interchange between two distinct forms of cyanobacterial photosystem II reaction-center protein D1 in response to photoinhibition. *Proc. Natl. Acad. Sci. USA* **90**:9973–9977.
- Collier, J. L., and A. R. Grossman. 1992. Chlorosis induced by nutrient deprivation in *Synechococcus* sp. strain PCC 7942: not all bleaching is the same. *J. Bacteriol.* **174**:4718–4726.
- Collier, J. L., and A. R. Grossman. 1994. A small polypeptide triggers complete degradation of light-harvesting phycobiliproteins in nutrient-deprived cyanobacteria. *EMBO J.* **13**:1039–1047.
- Danon, A., and S. P. Mayfield. 1994. Light-regulated translation of chloroplast messenger RNAs through redox potential. *Science* **266**:1717–1719.
- David, M., M. L. Daveran, J. Batut, A. Dedieu, O. Domergue, J. Ghai, C. Hertig, P. Boistard, and D. Kahn. 1988. Cascade regulation of *nif* gene expression in *Rhizobium meliloti*. *Cell* **54**:671–683.
- Demmig-Adams, B., and W. W. Adams III. 1992. Photoprotection and other responses of plants to high light stress. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **43**:599–626.
- Dolganov, N., and A. R. Grossman. 1999. A polypeptide with similarity to phycocyanin α -subunit phycocyanobilin lyase involved in degradation of phycobilisomes. *J. Bacteriol.* **181**:610–617.
- Dolganov, N. A. M., D. Bhaya, and A. R. Grossman. 1995. Cyanobacterial protein with similarity to the chlorophyll *a/b* binding proteins of higher plants: evolution and regulation. *Proc. Natl. Acad. Sci. USA* **92**:636–640.
- Escoubas, J.-M., M. Lomas, J. LaRoche, and P. G. Falkowski. 1995. Light intensity regulation of *cab* gene transcription is signaled by the redox state of the plastoquinone pool. *Proc. Natl. Acad. Sci. USA* **92**:10237–10241.
- Golden, S. S., J. Brusslan, and R. Haselkorn. 1986. Expression of a family of *psbA* genes encoding a photosystem II polypeptide in the cyanobacterium *Anacystis nidulans* R2. *EMBO J.* **5**:2789–2798.
- Grossman, A. R., D. Bhaya, K. E. Apt, and D. M. Kehoe. 1995. Light-harvesting complexes in oxygenic photosynthesis: diversity, control, and evolution. *Annu. Rev. Genet.* **29**:231–288.
- He, Q., N. Dolganov, O. Björkman, and A. R. Grossman. 2001. The high light-inducible polypeptides in *Synechocystis* PCC 6803. Expression and function in high light. *J. Biol. Chem.* **276**:306–314.
- Herdman, M., R. W. Castenholz, I. Iteman, J. B. Waterbury, and R. Rippka. 2001. Order Chroococcales, Wettstein 1924, emend. Rippka, Deruelles, Waterbury, Herdman and Stanier 1979, p. 776. In G. M. Garrity (ed.), *Bergey's manual of systematic bacteriology: the archaea and the deeply branching and phototropic bacteria*, 2nd ed., vol. 1. Springer-Verlag, New York, N.Y.
- Hihara, Y., A. Kamei, M. Kanehisa, A. Kaplan, and M. Ikeuchi. 2001. DNA microarray analysis of cyanobacterial gene expression during acclimation to high light. *Plant Cell* **13**:793–806.
- Iuchi, S., Z. Matsuda, T. Fujiwara, and E. C. Lin. 1990. The *arcB* gene of *Escherichia coli* encodes a sensor-regulator protein for anaerobic repression of the *arc* regulon. *Mol. Microbiol.* **4**:715–727.
- Jensen, K. K., E. Sharkova, M. F. Duggan, Y. Qi, A. Koide, J. A. Hoch, and F. M. Hulett. 1993. *Bacillus subtilis* transcription regulator, SpoOA, decreases alkaline phosphatase levels induced by phosphate starvation. *J. Bacteriol.* **175**:3749–3756.
- Kehoe, D. M., and A. R. Grossman. 1996. Similarity of a chromatic adaptation sensor to phytochrome and ethylene receptors. *Science* **273**:1409–1412.
- Koh, M., G. van Driessche, B. Samyn, W. D. Hoff, T. E. Meyer, M. A. Cusanovich, and J. J. van Beeumen. 1996. Sequence evidence for strong conservation of the photoactive yellow proteins from the halophilic phototrophic bacteria *Chromatium salexigens* and *Rhodospirillum salexigens*. *Biochemistry* **35**:2526–2534.
- Laudenbach, D. E., and A. R. Grossman. 1991. Characterization and mutagenesis of sulfur-regulated genes in a cyanobacterium: evidence for function in sulfate transport. *J. Bacteriol.* **173**:2739–2750.
- Li, H., and L. A. Sherman. 2000. A redox-responsive regulator of photosynthesis gene expression in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **182**:4268–4277.
- Maxwell, D. P., D. E. Laudénbach, and N. P. A. Hunter. 1995. Redox regulation of light-harvesting complex II and *cab* mRNA abundance in *Dunaliella salina*. *Plant Physiol.* **109**:787–795.
- Nakano, M. M., and P. Zuber. 1998. Anaerobic growth of a "strict aerobe" (*Bacillus subtilis*). *Annu. Rev. Microbiol.* **52**:165–190.
- Nestmann, E. R. 1975. Mutagenesis by nitrosoguanidine, ethyl methanesulfonate, and mutator gene *mutH* in continuous cultures of *Escherichia coli*. *Mutat. Res.* **28**:323–330.
- Niyogi, K. K. 1999. Photoprotection revisited: genetic and molecular approaches. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **50**:333–359.
- Parkinson, J. S. 1993. Signal transduction schemes of bacteria. *Cell* **73**:857–871.
- Parkinson, J. S., and E. C. Kofoid. 1992. Communication modules in bacterial signaling proteins. *Annu. Rev. Genet.* **26**:71–112.
- Pellequer, J. L., K. A. Wager-Smith, S. A. Kay, and E. D. Getzoff. 1998. Photoactive yellow protein: a structural prototype for the three-dimensional fold of the PAS domain superfamily. *Proc. Natl. Acad. Sci. USA* **95**:5884–5890.
- Pfanschmidt, T., A. Nilsson, and J. F. Allen. 1999. Photosynthetic control of chloroplast gene expression. *Nature* **397**:625–628.
- Reith, M. E., and J. Munholland. 1995. Complete nucleotide sequence of the *Porphyra purpurea* chloroplast genome. *Plant Mol. Biol. Rep.* **13**:333–335.
- Reyes, J. C., and F. J. Florencio. 1995. Electron transport controls transcription of the glutamine synthetase gene (*glnA*) from the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Mol. Biol.* **27**:789–799.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Schwarz, R., and A. R. Grossman. 1998. A response regulator of cyanobacteria integrates diverse environmental signals and is critical for survival under extreme conditions. *Proc. Natl. Acad. Sci. USA* **95**:11008–11013.
- Stock, J. B., A. M. Stock, and J. M. Mottonen. 1990. Signal transduction in bacteria. *Nature* **344**:395–400.
- Sun, G., E. Sharkova, R. Chesnut, S. Birkey, M. F. Duggan, A. Sorokin, P.

- Pujic, D. Ehrlich, and F. M. Hulett.** 1996. Regulators of aerobic and anaerobic respiration in *Bacillus subtilis*. *J. Bacteriol.* **178**:1374–1385.
46. **Suzuki, I., Y. Kanesaki, K. Mikami, M. Kanehisa, and N. Murata.** 2001. Cold-regulated genes under control of the cold sensor Hik33 in *Synechocystis*. *Mol. Microbiol.* **40**:235–244.
47. **Suzuki, I., D. A. Los, Y. Kanesaki, K. Mikami, and N. Murata.** 2000. The pathway for perception and transduction of low-temperature signals in *Synechocystis*. *EMBO J.* **19**:1327–1334.
48. **Taylor, B. L., and I. B. Zhulin.** 1999. PAS domains: internal sensors of oxygen, redox potential, and light. *Microbiol. Mol. Biol. Rev.* **63**:479–506.
49. **Thompson, W. F., and M. J. White.** 1991. Physiological and molecular studies of light-regulated nuclear genes in higher plants. *Annu. Rev. Plant Physiol.* **42**:423–466.
50. **Tsinoremas, N. F., M. R. Schaefer, and S. S. Golden.** 1994. Blue and red light reversibly control *psbA* expression in the cyanobacterium *Synechococcus* sp. strain PCC 7942. *J. Biol. Chem.* **269**:16143–16147.
51. **Williams, S. B., and V. Stewart.** 1999. Functional similarities among two-component sensors and methyl-accepting chemotaxis proteins suggest a role for linker region amphipathic helices in transmembrane signal transduction. *Mol. Microbiol.* **33**:1093–1102.
52. **Yamanaka, G., and A. N. Glazer.** 1980. Dynamic aspects of phycobilisome structure. Phycobilisome turnover during nitrogen starvation in *Synechococcus* spp. *Arch. Microbiol.* **124**:39–47.
53. **Yarwood, J. M., J. K. McCormick, and P. M. Schlievert.** 2001. Identification of a novel two-component regulatory system that acts in global regulation of virulence factors of *Staphylococcus aureus*. *J. Bacteriol.* **183**:1113–1123.
54. **Zhulin, I. B., B. L. Taylor, and R. Dixon.** 1997. PAS domain S-boxes in Archaea, Bacteria and sensors for oxygen and redox. *Trends Biochem. Sci.* **22**:331–333.