

1 Reconstructing Neandertal behavior, diet, and disease using ancient DNA

2

3 Authors: Laura S Weyrich<sup>1</sup>, Sebastian Duchene<sup>2</sup>, Julien Soubrier<sup>1</sup>, Luis Arriola<sup>1</sup>,  
4 Bastien Llamas<sup>1</sup>, James Breen<sup>1</sup>, Alan G Morris<sup>3</sup>, Kurt W Alt<sup>4</sup>, David Caramelli<sup>5</sup>, Veit  
5 Dresely<sup>6</sup>, Milly Farrell<sup>7</sup>, Andrew G Farrer<sup>1</sup>, Michael Francken<sup>8</sup>, Neville Gully<sup>9</sup>,  
6 Wolfgang Haak<sup>1</sup>, Karen Hardy<sup>10</sup>, Katerina Harvati<sup>8</sup>, Petra Held<sup>11</sup>, Edward Holmes<sup>2</sup>,  
7 John Kaidonis<sup>9</sup>, Johannes Krause<sup>12</sup>, Carles Lalueza-Fox<sup>13</sup>, Marco de la Rasilla<sup>14</sup>,  
8 Antonio Rosas<sup>15</sup>, Patrick Semal<sup>16</sup>, Arkadiusz Soltysiak<sup>17</sup>, Grant Townsend<sup>9</sup>, Donatella  
9 Usai<sup>18</sup>, Joachim Wahl<sup>19</sup>, Daniel H. Huson<sup>20</sup>, Keith Dobney<sup>21</sup>, and Alan Cooper<sup>1</sup>

10

11 **Affiliations:**

12 <sup>1</sup> Australian Centre for Ancient DNA, School of Biological Sciences, University of  
13 Adelaide, Adelaide, South Australia, Australia

14 <sup>2</sup> Charles Perkins Centre, University of Sydney, Sydney, Australia

15 <sup>3</sup> Department of Human Biology, University of Cape Town, Cape Town, South Africa

16 <sup>4</sup> Danube Private University, Krems, Austria; State Office for Heritage Management  
17 and Archaeology, Saxony-Anhalt, Germany; Heritage Museum, Halle, Germany; and  
18 Institute for Prehistory and Archaeological Science, Basel University, Switzerland

19 <sup>5</sup> Department of Biology, University of Florence, Florence, Italy

20 <sup>6</sup> Archaeology Saxony-Anhalt and Heritage Museum, Halle, Germany

21 <sup>7</sup> Royal College of Surgeons, London, United Kingdom

22 <sup>8</sup> Institute for Archaeological Sciences, University of Tübingen, Tübingen, Germany

23 <sup>9</sup> School of Dentistry, The University of Adelaide, Adelaide, Australia

24 <sup>10</sup> ICREA (Catalan Institution for Research and Advanced Studies), Department of  
25 Prehistory Universitat Autònoma de Barcelona, Barcelona, Spain

26 <sup>11</sup> Institute of Anthropology, University of Mainz

27 <sup>12</sup> Institute for Archaeological Sciences, University of Tübingen, Tübingen, Germany,  
28 and Max Planck Institute for the Science of Human History, Jena, Germany

29 <sup>13</sup> Institute of Evolutionary Biology, CSIC-Universitat Pompeu Fabra, Barcelona,  
30 Spain

31 <sup>14</sup> Área de Prehistoria, Departamento de Historia, Universidad de Oviedo, Oviedo,  
32 Spain

33 <sup>15</sup> Paleoanthropology Group, Department of Paleobiology, Museo Nacional de  
34 Ciencias Naturales, CSIC, Madrid, Spain

35 <sup>16</sup> Scientific Service Heritage, Royal Belgian Institute of Natural Sciences, Brussels,  
36 Belgium

37 <sup>17</sup> Department of Bioarchaeology, Institute of Archaeology, University of Warsaw,  
38 Warsaw, Poland.

39 <sup>18</sup> Istituto Italiano per l’Africa e l’Oriente (IsIAO), Rome, Italy

40 <sup>19</sup> State Office for Cultural Heritage Management Baden-Württemberg, Esslingen  
41 Germany

42 <sup>20</sup> Department of Algorithms in Bioinformatics, University of Tübingen, Tübingen,  
43 Germany

44 <sup>21</sup> Department of Archaeology, Classics and Egyptology, School of Histories,  
45 Languages and Cultures, University of Liverpool, Liverpool, United Kingdom

46 **Abstract**

47           Recent genomic data has revealed multiple interactions between Neandertals  
48 and humans, but there is currently little genetic evidence about Neandertal behavior,  
49 diet, or health. We shotgun sequenced ancient DNA from five Neandertal dental  
50 calculus specimens to characterize regional differences in Neandertal ecology. At  
51 Spy, Belgium, Neandertal diet was heavily meat based, and included woolly  
52 rhinoceros and wild sheep - animals characteristic of a steppe environment. In El  
53 Sidrón, Spain, no meat was detected in the dental calculus, but dietary components  
54 including mushrooms, pine nuts, and moss reflected forest gathering. Differences in  
55 diet were also linked to an overall shift in the oral bacterial community (microbiota)  
56 in Neandertals, suggesting that meat consumption contributed to significant variation  
57 between Neandertal microbiota. Evidence for self-medication was identified in one  
58 El Sidrón Neandertal with a dental abscess, who also likely suffered from a chronic  
59 gastrointestinal pathogen (*Enterocytozoon bieneusi*). Lastly, we characterized a nearly  
60 complete genome of the archaeal commensal *Methanobrevibacter oralis* in  
61 Neandertals – the oldest draft microbial genome generated to date at ~48,000 years  
62 old (10.2 depth). DNA preserved within dental calculus represents an important new  
63 resource of behavioral and health information for ancient hominid specimens, as well  
64 as a unique long-term study system for microbial evolution.

65

66 **Main Text:**

67 Neandertals remain one of our closest extinct hominid relatives, and have been  
68 shown to have co-existed and occasionally interbred with anatomically modern  
69 humans (AMHs) across Eurasia in the Late Pleistocene. Neandertals went extinct in  
70 Europe around 42,000 years ago, while the extinction process across the rest of  
71 Eurasia is less clear (1, 2). Isotopic and archaeological data through the last glacial  
72 cycle (~120-12,000 years ago) suggests that Neandertals were as carnivorous as polar  
73 bears or wolves (3, 4), with a diet heavily based on large terrestrial herbivores, such  
74 as reindeer, woolly mammoth, and woolly rhinoceros (4, 5). In contrast, microwear  
75 analysis of tooth surfaces from Neandertals in different ecological settings, suggests  
76 diets were likely guided by local food availability (6). Furthermore, phytoliths and  
77 proteins preserved in calcified dental plaque (calculus) indicate Neandertal diets  
78 included many plants, including some potentially cooked prior to consumption or  
79 used for medicinal purposes (7, 8). As a result, Neandertal diet and behavior remains  
80 a topic of considerable debate, with important questions remaining about the specific  
81 animals and plants actually consumed, or potential impacts on Neandertal health or  
82 disease.

83 While genomic studies continue to reveal evidence of interbreeding between  
84 AMHs and Neandertals across Eurasia (9), little is known about the health  
85 consequences of these interactions. The genetic analysis of Neandertal dental calculus  
86 represents an opportunity to examine this issue, and to reconstruct Neandertal diet,  
87 behavior, and health (10, 11, 12). Here, we report the first genetic analysis of dental  
88 calculus from five Neandertals from El Sidrón cave in Spain (n=2), Spy cave in  
89 Belgium (n=2) and Breuil Grotta in Italy (n=1), alongside a historic chimpanzee (n=1)  
90 and a modern human (n=1) for comparison (Table S1). To provide increased

91 resolution of Neandertal diseases, we also deeply sequenced dental calculus from the  
92 best preserved specimen, El Sidrón 1 (>147 million reads), which was also suffering  
93 from an dental abscess (13).

94         Recent reports have identified significant size-based amplification biases  
95 when amplicon approaches are applied to ancient microbial DNA (*i.e.* when using  
96 16S ribosomal RNA (rRNA) amplicon sequencing of ancient bacterial taxa) (14).  
97 Hence, we examined the Neandertal dental calculus specimens using both Illumina  
98 16S rRNA amplicon (V4 region) (15) and metagenomic shotgun sequencing  
99 approaches. Blank control samples (EBCs) were sequenced to monitor contamination,  
100 and a stringent filtering strategy was applied to remove contaminant sequences in  
101 both datasets using QIIME (amplicon) or MEGAN5 (shotgun) (16, 17). The  
102 Neandertal sample from Breuil Grotta failed to produce amplicons, and was removed  
103 from downstream analysis. As expected, the amplicon datasets were not  
104 representative of the biodiversity revealed by shotgun sequencing (Figures S3-S4,  
105 S12, and S15-S16; Tables S2 and S7), and clustered together phylogenetically  
106 irrespective of sample age, while containing disproportionately large proportions of  
107 non-oral and environmental contaminant microorganisms (Figure 1, Table S2 and  
108 S7). Clearly, amplicon data from ancient specimens can be biased by DNA damage  
109 leading to preferential amplification of contaminating modern DNA and/or  
110 prokaryotic sequences with short ribosomal sequences (14). As a result, the  
111 metagenomic shotgun sequencing approach was used to analyze the Neandertal  
112 specimens.

113         The shotgun datasets consisted of short DNA fragments (<70 bp) which  
114 complicated accurate bacterial species identification using standard software, such as  
115 MG-RAST and DIAMOND (Figure S14) (17, 18). To circumvent this problem, we

116 used a novel metagenomic alignment tool that rapidly identifies species from shorter  
117 fragment lengths using a BLASTX like algorithm (MEGAN Alignment Tool with  
118 BLASTX-like alignments; MALT-X) (19). After benchmarking the tool (Figure S8-  
119 S10 and S14) (17), we applied MALT-X to the shotgun datasets and checked that the  
120 bacterial diversity accurately matched the subset of 16S rRNA fragments they  
121 contained (which are typically utilized to identify bacterial taxa); we also successfully  
122 repeated this test for two previously published ancient calculus datasets (Figure 1 and  
123 S13; Table S6) (20). Bioinformatic filtering was applied to remove laboratory and  
124 environmental contaminants from the samples, revealing that the Spy Neandertals  
125 were the most heavily impacted by environmental contamination (Figure S15-S17;  
126 Table S7) (17). Indeed, Spy I clustered within the biodiversity observed in the modern  
127 sample (Figure 1), contained high diversity analogous to environmental samples  
128 (Figure S20), and presented DNA damage patterns characteristic of modern  
129 contamination (Figure S22), so was therefore discarded from further analyses (17).  
130 The three robust Neandertal datasets (El Sidrón 1, El Sidrón 2, and Spy II) contained  
131 on average 93.76% bacterial, 5.91% archaeal, 0.27% eukaryotic, and 0.06% viral  
132 identifiable sequences, similar to previously published biodiversity of human dental  
133 calculus (Figure 2A and S17) (12). In addition, the six dominant bacterial phyla in the  
134 modern human mouth (Actinobacteria, Firmicutes, Bacteroidetes, Fusobacteria,  
135 Proteobacteria, and Spirochaetes) were also dominant in each of the Neandertals, with  
136 an average of 222 bacterial species per individual (Figure 2A and S17) (17).

137 We first examined Neandertal diets using the eukaryotic diversity preserved  
138 within dental calculus, after filtering spurious results (17). Calculus from the Spy II  
139 individual contained high numbers of reads mapping to rhinoceros (*Ceratotherium*  
140 *simum*) and sheep (*Ovis aries*), as well as the edible ‘grey shag’ mushroom

141 (*Coprinopsis cinerea*) (Table 1). Bones of woolly rhino and mouflon sheep were  
142 present in Spy Cave along with other mammalian fauna (21, 22), and both species are  
143 thought to have been part of the Neandertal diet at Spy and the nearby site of La  
144 Naulette (23). The dietary profile for El Sidrón Neandertals was markedly different,  
145 and contained no sequences matching large herbivores – no animal bone bearing  
146 contexts were associated with the recovery of the El Sidron Neanderthal remains.  
147 However, reads mapping to edible mushrooms (‘split gill’; *Schizophyllum commune*),  
148 pine nuts (*Pinus koraiensis*), forest moss (*Physcomitrella patens*), and poplar  
149 (*Populus trichocarpa*) were identified (Table 1). Sequences mapping to plant fungal  
150 pathogens were also observed (*Zymoseptoria tritici*, *Phaeosphaeria nodorum*,  
151 *Penicillium rubens*, and *Myceliophthora thermophile*), suggesting the El Sidrón  
152 Neandertals may have been consuming molded herbaceous material. The intracellular  
153 eukaryotic pathogen, *Enterocytozoon bieneusi*, which causes severe diarrhea (24),  
154 was also observed in El Sidrón 1, a young male that also had a dental abscess and  
155 may have used bitter medicinal compounds for self-medication (25). Our findings  
156 support earlier suggestions of self-medication, as he was the only individual whose  
157 calculus contained the natural antibiotic producing *Penicillium*, and poplar, whose  
158 bark, roots, and leaves contains the natural pain killer, salicylic acid (i.e. the active  
159 ingredient in aspirin) (26).

160 To examine how the oral microbial patterns in Neandertals were linked to  
161 dietary composition, we compared the data to a wide range of dental calculus  
162 specimens from ancient humans with varying diets, including ancient Later Stone Age  
163 (LSA) African gatherers; African Pastoralist Period individuals with high meat  
164 consumption (27); European hunter-gatherers with diets that included a wide range of  
165 proteins; and early European farmers with diets largely based around carbohydrates

166 and milk consumption (see SI for archaeological descriptions of dietary information)  
167 (17). We used UPGMA to cluster Bray Curtis distances obtained by comparing  
168 shotgun sequenced oral microbiota at an equal depth, and revealed four distinct  
169 groups: foragers with limited meat consumption (El Sidrón Neandertals, chimpanzee,  
170 and LSA African gatherers); hunter-gatherers or pastoralists with a frequent meat diet  
171 (Spy Neandertal, African pastoralists, and European hunter-gatherers); ancient  
172 agriculturalists (European farming individuals); and modern humans (Figure 2B).  
173 This analysis identifies a split between hunter-gatherers and agriculturalists, as  
174 previously observed (11), but also reveals two distinct hunter-gatherer clades,  
175 differentiated by the amount of meat consumed in their diet. Meat consumption  
176 appears to have impacted early hominid microbiota and health, similar to differences  
177 observed between carnivorous and herbivorous mammals (28). This finding also  
178 indicates that dental calculus may be used to directly infer the dietary behavior of  
179 ancient humans or hominids.

180 We then examined the Neandertal microbial diversity for signs of disease.  
181 Neandertal microbiota were more similar to the historic chimpanzee sample than  
182 modern humans, and contained less pathogenic Gram-negative species (18.9% Gram-  
183 negatives in Neandertals, compared to 77.6% in the modern human; Figure S21),  
184 which are associated with secondary enamel colonization, increased plaque formation,  
185 and periodontal disease (29). The lower levels of these immunostimulatory Gram-  
186 negative taxa in Neandertals may be related to the reduced presence of Fusobacteria  
187 taxa (Figure S17), as this keystone group facilitates the binding of Gram-negative  
188 microorganisms to the primary colonizers that bind to tooth enamel (e.g.  
189 *Streptococcus*, *Actinomyces* and *Methanobrevibacter* species).



190 Several oral pathogens could be identified within the shotgun data, although  
191 the short ancient sequences and diverse metagenomic background complicated  
192 identification. We established a number of exclusion criteria to verify the authenticity  
193 of short sequences, including the assessment of ancient DNA damage, phylogenetic  
194 position, and bioinformatic comparisons to differentiate close relatives (17).  
195 Pathogens that passed the exclusion criteria included the caries-associated pathogen  
196 *Streptococcus mutans*, which was identified in all Neandertals (0.08% to 0.18%)  
197 (Table S10-S11). All three members of the ‘red complex’ pathogens associated with  
198 modern periodontal disease were identified in at least one Neandertal  
199 (*Porphyromonas gingivalis*: 0-0.52%; *Tannerella forsythia* 0.05-2.4%; and  
200 *Treponema denticola* 0-1.87%), although all three pathogens were not identified in  
201 any single individual. These oral pathogens support the isolated evidence of  
202 Neandertal dental caries, periodontal disease, dental calculus, and associated tooth-  
203 picking to relieve dental pain observed from specimens at Krapina (Croatia), Shanidar  
204 (Iraq) and Cova Forada (Spain) (30–32). The microorganisms known to cause  
205 exacerbated whooping cough infections (*Bordetella parapertussis* and *Pasteurella*  
206 *multocida*) (Table S9) were also detected in El Sidrón 1; however, only a limited  
207 number of *B. parapertussis*-specific reads were identified (i.e. only 212 reads, mostly  
208 in a region containing hypothetical proteins, mapped more efficiently to *B.*  
209 *parapertussis* than *Bordetella petrii*, an environmental isolate). Similarly, pathogens  
210 closely related to common oral flora (*Neisseria gonorrhoeae*, *Streptococcus*  
211 *pyogenes*, and *Corynebacterium diphtheriae*) were identified but could not be  
212 unambiguously distinguished from closely related commensal oral taxa (Figure S23).  
213 These findings highlight the need for rigorous criteria when identifying pathogenic  
214 strains from ancient metagenomic data (17).

215 We also examined the commensal microorganisms in Neandertals in greater  
216 detail. Within the deeply sequenced El Sidrón 1 oral microbiome, we were able to  
217 recover eight draft ancient microbial genomes with >1-fold depth, corresponding to  
218 the most prevalent microbial taxa (Table 2). We were particularly interested in an  
219 archaeal species that dominated the oral metagenome of El Sidrón 1 (14.7%; Figure  
220 S17), but was present in lower proportions in the other Neandertal specimens (1.4%  
221 and 1.2% in El Sidrón 2, and Spy II, respectively). The large differences in G/C  
222 content between bacteria and archaea facilitated efficient read mapping of the  
223 archaeal sequences (Table 2), which mapped closest to the modern human-associated  
224 *Methanobrevibacter oralis* JMR01 strain. We were able to produce the first draft  
225 ancient archaeal genome, *Methanobrevibacter oralis neandertalensis* (10.3-fold  
226 depth; 44.7% of 2.1 Mbps; Figure 3), which is also the oldest draft microbial genome  
227 to date at ~48,000 years old (33). The DNA damage (33% C-T; 36% G-A) and  
228 fragment length distribution (average 58.67 bp) of *M. oralis neandertalensis* were  
229 consistent with an ancient microbial genome (Table 2).

230 Within the *M. oralis neandertalensis* genome, 1,929 coding sequences could  
231 be matched to modern human *M. oralis*, while 136 in the latter appeared to be absent  
232 (6.5% of all coding sequences in *M. oralis*) (Table S16). The absent loci included  
233 analogs of genes encoding antiseptic resistance (*qacE*) and those required for  
234 regulation of maltose metabolism (*sfsA*), likely reflecting the impacts of modern oral  
235 hygiene and relative dearth of carbohydrates in ancient Neandertal diets, respectively.  
236 As expected, bacterial immunity loci were also variable, as regions encoding for  
237 CRISPR Cas2 and Cas6 in modern *M. oralis* were missing from *M. oralis*  
238 *neandertalensis*, and the Cas1 CRISPR system in *M. oralis neandertalensis* could  
239 only be partially assembled (Table S16). The ratio of non-synonymous to

240 synonymous mutations ( $d_N/d_S$ ) between translatable *M. oralis neandertalensis* protein  
241 coding sequences and modern human *M. oralis* suggested that 58% were under  
242 purifying selection ( $d_N/d_S < 0.1$ ) (Table 3) (17). Only 4% appeared to be under positive  
243 selection ( $d_N/d_S > 1.2$ ), and included the conjugal transfer gene, *traB*, which aids in the  
244 uptake of foreign DNA (i.e. plasmid transfer) (34) and *mutT*, which is involved in  
245 DNA mismatch repair (35). These findings suggest that much of the genome is under  
246 purifying selective pressure, potentially due to the conserved environmental  
247 conditions in the hominid mouth over time (36). Overall, it appears that modern  
248 human *M. oralis* genome adaptation has occurred primarily through the uptake of new  
249 DNA sequences, rather than adaptive gene mutation.

250 Phylogenetic analysis of seven modern *Methanobrevibacter* genomes revealed  
251 that *M. oralis neandertalensis* is sister taxa to the modern human *M. oralis* strain  
252 (JMR01), and together, this clade is sister to *Methanobrevibacter smithii* (a  
253 commensal found in the modern gut) (Figure 3B). Molecular dating using a strict  
254 clock model indicates the divergence between *M. oralis neandertalensis* and modern  
255 human *M. oralis* strains occurred around 126,000 years ago (95% highest posterior  
256 density interval of 112-143K years) (Figure 3B), long after the genomic divergence of  
257 Neandertals and modern humans (450-750 kyr) (37). This suggests that microbial  
258 species were transferred during subsequent interactions, potentially during early  
259 human-Neandertal interbreeding in Africa (38).

260 Preserved dental calculus represents an important new resource of behavioral  
261 and health information for ancient hominid specimens, as well as a unique long-term  
262 study system for microbial evolution. A prime example is the potential link between  
263 the elevated levels of Fusobacteria taxa in modern humans and the increased diversity  
264 of Gram-negative immunostimulatory taxa. The latter are strongly associated with a

265 wide range of modern Western diseases, revealing the potential power of ancient

266 DNA to inform modern medical research.

267

268 **References**

- 269 1. M. Meyer *et al.*, A High-Coverage Genome Sequence from an Archaic  
270 Denisovan Individual. *Science*. **338**, 222–226 (2012).
- 271 2. T. Higham *et al.*, The timing and spatiotemporal patterning of Neanderthal  
272 disappearance. *Nature*. **512**, 306–309 (2014).
- 273 3. M. P. Richards *et al.*, Neanderthal diet at Vindija and Neanderthal predation:  
274 The evidence from stable isotopes. *PNAS*. **97**, 7663–7666 (2000).
- 275 4. M. P. Richards, E. Trinkaus, Isotopic evidence for the diets of European  
276 Neanderthals and early modern humans. *PNAS*. **106**, 16034–16039 (2009).
- 277 5. H. Bocherens, D. G. Drucker, D. Billiou, M. Patou-Mathis, B. Vandermeersch,  
278 Isotopic evidence for diet and subsistence pattern of the Saint-Césaire I  
279 Neanderthal: review and use of a multi-source mixing model. *Journal of Human  
280 Evolution*. **49**, 71–87 (2005).
- 281 6. S. El Zaatari, F. E. Grine, P. S. Ungar, J.-J. Hublin, Ecogeographic variation in  
282 Neanderthal dietary habits: Evidence from occlusal molar microwear texture  
283 analysis. *Journal of Human Evolution*. **61**, 411–424 (2011).
- 284 7. A. G. Henry, A. S. Brooks, D. R. Piperno, Microfossils in calculus demonstrate  
285 consumption of plants and cooked foods in Neanderthal diets (Shanidar III, Iraq;  
286 Spy I and II, Belgium). *PNAS* (2010), doi:10.1073/pnas.1016868108.
- 287 8. K. Hardy *et al.*, Neanderthal medics? Evidence for food, cooking, and medicinal  
288 plants entrapped in dental calculus, 617–626 (2012).
- 289 9. Q. Fu *et al.*, An early modern human from Romania with a recent Neanderthal  
290 ancestor. *Nature*. **524**, 216–219 (2015).
- 291 10. L. S. Weyrich, K. Dobney, A. Cooper, Ancient DNA analysis of dental calculus.  
292 *Journal of Human Evolution*. **79**, 119–124 (2015).
- 293 11. C. J. Adler *et al.*, Sequencing ancient calcified dental plaque shows changes in  
294 oral microbiota with dietary shifts of the Neolithic and Industrial revolutions.  
295 *Nat Genet*. **45**, 450–455 (2013).
- 296 12. C. Warinner *et al.*, Pathogens and host immunity in the ancient human oral  
297 cavity. *Nat Genet*. **46**, 336–344 (2014).
- 298 13. A. Rosas *et al.*, Paleobiology and comparative morphology of a late Neanderthal  
299 sample from El Sidrón, Asturias, Spain. *PNAS*. **103**, 19266–19271 (2006).
- 300 14. K. A. Ziesemer *et al.*, Intrinsic challenges in ancient microbiome reconstruction  
301 using 16S rRNA gene amplification. *Scientific Reports*. **5**, 16498 (2015).
- 302 15. J. G. Caporaso *et al.*, Ultra-high-throughput microbial community analysis on  
303 the Illumina HiSeq and MiSeq platforms. *ISME J*. **6**, 1621–1624 (2012).

- 304 16. D. H. Huson, S. Mitra, H.-J. Ruscheweyh, N. Weber, S. C. Schuster, Integrative  
305 analysis of environmental sequences using MEGAN4, 1552–1560 (2011).
- 306 17. Materials and methods are available as supplementary materials on Science  
307 Online.
- 308 18. F. Meyer *et al.*, The metagenomics RAST server – a public resource for the  
309 automatic phylogenetic and functional analysis of metagenomes, 386 (2008).
- 310 19. A. Herbig *et al.*, MALT: Fast alignment and analysis of metagenomic DNA  
311 sequence data applied to the Tyrolean Iceman. *bioRxiv*, 50559 (2016).
- 312 20. C. Warinner *et al.*, Pathogens and host immunity in the ancient human oral  
313 cavity. *Nat Genet.* **46**, 336–344 (2014).
- 314 21. M. Germonpré, M. Udrescu, E. Fiers, The fossil mammals of Spy.  
315 *Anthropologica et Præhistorica.* **123/2012**, 298–327 (2013).
- 316 22. M. Germonpré, M. Udrescu, E. Fiers, Possible evidence of mammoth hunting at  
317 the Neanderthal site of Spy (Belgium). *Quaternary International.* **337**, 28–42  
318 (2014).
- 319 M. Germonpré, M. Udrescu, E. Fiers, The fossil mammals of Spy. *Anthropologica et*  
320 *Præhistorica.* **123/2012**, 298–327 (2013).
- 321 22. M. Germonpré, M. Udrescu, E. Fiers, Possible evidence of mammoth hunting at  
322 the Neanderthal site of Spy (Belgium). *Quaternary International.* **337**, 28–42  
323 (2014).
- 324 23. M. Patou-Mathis, Neanderthal subsistence behaviours in Europe. *International*  
325 *Journal of Osteoarchaeology.* **10**, 379–395 (2000).
- 326 24. J. K. Tumwine *et al.*, Enterocytozoon bienewisi among children with diarrhea  
327 attending Mulago Hospital in Uganda. *Am. J. Trop. Med. Hyg.* **67**, 299–303  
328 (2002).
- 329 25. K. Hardy *et al.*, Neanderthal medics? Evidence for food, cooking, and medicinal  
330 plants entrapped in dental calculus. *Naturwissenschaften.* **99**, 617–626 (2012).
- 331 26. C. J. Tsai *et al.*, Salicylate metabolism in Populus. *BMC Proceedings.* **5**, 1–2  
332 (2011).
- 333 27. W. R. Leonard, M. H. Crawford, *The Human Biology of Pastoral Populations*  
334 (Cambridge University Press, 2002).
- 335 28. B. D. Muegge *et al.*, Diet Drives Convergence in Gut Microbiome Functions  
336 Across Mammalian Phylogeny and Within Humans. *Science.* **332**, 970–974  
337 (2011).
- 338 29. D. P. D. Marsh, D. J. Bradshaw, Dental plaque as a biofilm. *Journal of*  
339 *Industrial Microbiology.* **15**, 169–175 (1995).

- 340 30. M. Lozano, M. E. lia SubirÃ, J. Aparicio, C. Lorenzo, G. GÃmez-Merino,  
341 Toothpicking and Periodontal Disease in a Neanderthal Specimen from Cova  
342 ForadÃ Site (Valencia, Spain). *PLoS ONE*. **8**, e76852 (2013).
- 343 31. E. Trinkaus, Dental remains from the Shanidar adult Neanderthals. *Journal of*  
344 *Human Evolution*. **7**, 369–382 (1978).
- 345 32. B. Topić, H. Rašćić-Konjhodžić, M. Čížek Sajko, Periodontal disease and dental  
346 caries from Krapina Neanderthal to contemporary man - skeletal studies. *Acta*  
347 *Med Acad*. **41**, 119–130 (2012).
- 348 33. R. E. Wood *et al.*, A New Date for the Neanderthals from El Sidrón Cave  
349 (asturias, Northern Spain)\*. *Archaeometry*. **55**, 148–158 (2013).
- 350 34. S. Kosoño, M. Kataoka, T. Seki, T. Yoshida, The TraB protein, which mediates  
351 the intermycelial transfer of the *Streptomyces* plasmid pSN22, has functional  
352 NTP-binding motifs and is localized to the cytoplasmic membrane. *Mol.*  
353 *Microbiol*. **19**, 397–405 (1996).
- 354 35. R. G. Fowler, R. M. Schaaper, The role of the mutT gene of *Escherichia coli* in  
355 maintaining replication fidelity. *FEMS Microbiol. Rev.* **21**, 43–54 (1997).
- 356 36. L. Dethlefsen, M. McFall-Ngai, D. A. Relman, An ecological and evolutionary  
357 perspective on human–microbe mutualism and disease. *Nature*. **449**, 811–818  
358 (2007).
- 359 37. C. Stringer, The origin and evolution of *Homo sapiens*. *Phil. Trans. R. Soc. B.*  
360 **371**, 20150237 (2016).
- 361 38. S. Sankararaman, N. Patterson, H. Li, S. Paabo, D. Reich, The date of  
362 interbreeding between Neandertals and modern humans. *PLoS genetics*. **8**,  
363 e1002947 (2012).
- 364

365 **Figures and Legends**

366 **Figure 1: Comparison of 16S amplicon and shotgun datasets obtained from**  
367 **ancient, historic, and modern dental calculus samples.** Filtered and unfiltered 16S  
368 rRNA amplicon and shotgun datasets, as well as 16S rRNA shotgun sequences  
369 identified using GraftM, were compared using UPGMA clustering of Bray Curtis  
370 distances from a chimpanzee (red), Neandertals (El Sidrón 1 (green), El Sidrón 2  
371 (light green), Spy I (grey), Spy II (blue)), and a modern human (orange).



**Figure 2: Bacterial community composition at the phyla level of oral microbiota from chimpanzee, Neandertal, and modern human samples.** Oral microbiota from shotgun datasets of a wild-caught chimpanzee (A), Neandertals (n=3; B), and a modern human (C) are presented at the phyla level. Phyla names were simplified for clarity, and unidentified reads were excluded. Gram-positive (blue) and Gram-negative (red) phyla are differentiated by color. (D) UPGMA clustering of Bray Curtis values obtained from 22 oral rarefied metagenomes is displayed. Definitions for abbreviations can be found in the SI.

**Figure 3: 48,000 yBP archeal draft genome and phylogeny of *Methanobrevibacter oralis neandertalensis*.**

(A) Ancient sequences mapping to *Methanobrevibacter oralis* JMR01 are displayed in a Circos plot (black), alongside the depth of coverage obtained (red). The reference sequence is displayed (grey) with the GC content of the reference sequence calculated in 2500 bp bins (green). (B) A *Methanobrevibacter* phylogeny was constructed from whole genome alignments in RAxML with 100 bootstraps. The date for the split between *M. oralis* strains and *M. smithii* strains was calculated from whole genomes in BEAST using a strict clock model.

**Table 1: Dietary information preserved in calculus.**

DNA sequences mapping to eukaryotic species are shown as a proportion of the total eukaryotic reads identified within each sample. Eukaryotic sequencing identified in the extraction blank controls and the Spy I Neandertal, which is heavily contaminated with modern DNA, are shown to the right.

**Table 2: Draft microbial genomes present in El Sidrón 1.**

Eight draft microbial genomes were obtained from the deeply sequenced El Sidrón 1 specimen by mapping with ancient DNA parameters using bwa aln. The sequences coverage, GC content, sequencing depth, and damage profile (average fragment length and base pair modifications calculated from MapDamage v2) are displayed for each genome.

**Table 3: Purifying and positive selection in *M. oralis neandertalensis*.**

The ratio of nonsynonymous to synonymous ( $d_N/d_S$ ) mutations was calculated for coding regions with sufficient coverage and that were conserved between *M. oralis* and *M. oralis neandertalensis*. Genes that have undergone definite purifying ( $d_N/d_S = 0$ ) or positive selection ( $d_N/d_S > 1.2$ ; grey) are displayed if the function of the gene was annotated. Hypothetical proteins and those not matching to the *M. oralis* genome during BLAST searches are not shown.

## **Acknowledgements**

We thank Giorgio Manzi (University of Rome), the Odontological Collection of the Royal College of Surgeons, Royal Belgian Institute of Natural Sciences, Museo Nacional de Ciencias Naturales, and Adelaide Universities for access to dental calculus material. We thank Adam Croxford for DNA sequencing and Alan Walker, Alan Morris, and Alexander Herbig for critical feedback.

### **Author Contributions**

AGM, KWA, DC, VD, MF, MF, NG, WH, KH, KH, PH, JK, CLF, MR, AR, PS, AS, DU, and JW provided samples and interpretations of associated archaeological goods; LSW, KD, and AC designed experiments; LSW performed experiments; LSW, SD, EH, JS, BL, JB, LA, and AGF performed bioinformatics analysis and interpretation on the data; DHH developed bioinformatics tools; NG, JK, and GT analyzed medical relevance of data; LSW and AC wrote the paper; and all authors contributed to editing the manuscript.

## **Author Information**

Raw and analyzed datasets and the scripts utilized for this analysis are available in the Online Ancient Genome Repository (OAGR) (currently available at:

<https://www.oagr.org.au/experiment/view/16/?token=9LCF0GKSL7DHO3FPR4YBSZCGYD0ASJ>). The Australian Research Council supported this work, and the authors

declare no competing financial interests. Requests for materials should be addressed

to [laura.weyrich@adelaide.edu.au](mailto:laura.weyrich@adelaide.edu.au).