

 Open access • Journal Article • DOI:10.1645/16-34

Neospora caninum in Wild Waterfowl: Occurrence of Parasite DNA and Low Antibody Titers. — [Source link](#)

[Guido Rocchigiani](#), [Alessandro Poli](#), [Simona Nardoni](#), [Roberto Amerigo Papini](#) ...+1 more authors

Institutions: [University of Pisa](#)

Published on: 10 Mar 2017 - [Journal of Parasitology](#) (American Society of Parasitologists)

Topics: [Neospora caninum](#), [Neospora](#) and [Anas crecca](#)

Related papers:

- [Toxoplasma gondii and Neospora caninum in sparrows \(Passer domesticus\) in the Northeast of Brazil.](#)
- [Detection of Neospora Caninum DNA in Wild Birds from Italy.](#)
- [High seroprevalence of Toxoplasma gondii and Neospora caninum in the Common raven \(Corvus corax\) in the Northeast of Spain.](#)
- [Presence of Toxoplasma gondii and Neospora caninum DNA in the brain of wild birds](#)
- [Survey for natural Neospora caninum infection in wild and captive birds.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/neospora-caninum-in-wild-waterfowl-occurrence-of-parasite-4n6llzp5rn>

PubMed

Format: Abstract ▾

J Parasitol. 2016 Nov 2. [Epub ahead of print]

Neospora caninum in Wild Waterfowl: Occurrence of Parasite DNA and Low antibody titers.

Rocchigiani G¹, Poli A², Nardoni S³, Roberto P⁴, Mancianti F⁵.

⊕ Author information

Abstract

Thirty-five adult waterfowl (14 males and 21 females) representing various orders and species were sampled during the hunting season from 2015 to 2016. Antibodies to *Neospora caninum* were detected by IFAT on blood samples, while heart and brain were subjected to molecular analysis for the detection of *Neospora caninum* DNA. Twelve birds (34.3%) (6 *Anas crecca*, 3 *Anas platyrhynchos*, 2 *Anas penelope*, 1 *Anas acuta*) showed antibodies versus *N. caninum*, while 10 animals out of 35 (4 *Anas crecca*, 2 *Anas platyrhynchos*, 2 *Anas penelope*, 1 *Anas acuta*, and 1 *Vanellus vanellus*) scored positive for at least 1 sample, with an overall prevalence of 28.6%. The present report shows for the first time the occurrence of antibodies and *N. caninum* DNA in waterfowl. The avian species investigated in the present report, which feed directly from the soil and/or water, would be able to ingest oocysts excreted by final canid hosts and could contribute to parasite transmission in the sylvatic cycle. To achieve a definitive result about the role of these avian species in the epidemiology of this protozoan, the presence of viable parasites should be demonstrated by bioassay and/or culture, as well as histological evidence of *N. caninum* cysts in avian tissues.

PMID: 27805842 DOI: [10.1645/16-34](https://doi.org/10.1645/16-34)

[PubMed - as supplied by publisher]



LinkOut - more resources



PubMed Commons

[PubMed Commons home](#)

0 comments

[How to join PubMed Commons](#)