



Nerve Growth Factor and Pathogenesis of Leprosy: Review and Update

Tinara Leila de Souza Aarão¹, Jorge Rodrigues de Sousa², Aline Semblano Carreira Falcão³, Luiz Fábio Magno Falcão¹ and Juarez Antônio Simões Quaresma^{1,2,3*}

¹Center of Health and Biological Sciences, State University of Para, Belem, Brazil, ²Tropical Medicine Center, Federal University of Para, Belem, Brazil, ³Evandro Chagas Institute, Ministry of Health, Ananindeua, Brazil

OPEN ACCESS

Edited by:

Vânia Nieto Brito De Souza,
Instituto Lauro de Souza Lima, Brazil

Reviewed by:

Roberta Olmo Pinheiro,
Fundação Oswaldo Cruz (Fiocruz),
Brazil

Esmail Mortaz,
National Research Institute
of Tuberculosis and
Lung Diseases, Iran

*Correspondence:

Juarez Antônio Simões Quaresma
juarez.quaresma@gmail.com

Specialty section:

This article was submitted
to Microbial Immunology,
a section of the journal
Frontiers in Immunology

Received: 09 October 2017

Accepted: 16 April 2018

Published: 07 May 2018

Citation:



Aarão TLdS, de Sousa JR,
Falcão ASC, Falcão LFM and
Quaresma JAS (2018) Nerve
Growth Factor and Pathogenesis
of Leprosy: Review and Update.
Front. Immunol. 9:939.
doi: 10.3389/fimmu.2018.00939

Neurotrophins are a family of proteins that regulate different aspects of biological development and neural function and are of great importance in neuroplasticity. This group of proteins has multiple functions in neuronal cells, as well as in other cellular populations. Nerve growth factor (NGF) is a neurotrophin that is endogenously produced during development and maturation by multiple cell types, including neurons, Schwann cells, oligodendrocytes, lymphocytes, mast cells, macrophages, keratinocytes, and fibroblasts. These cells produce proNGF, which is transformed by proteolytic cleavage into the biologically active NGF in the endoplasmic reticulum. The present review describes the role of NGF in the pathogenesis of leprosy and its correlations with different clinical forms of the disease and with the phenomena of regeneration and neural injury observed during infection. We discuss the involvement of NGF in the induction of neural damage and the pathophysiology of pain associated with peripheral neuropathy in leprosy. We also discuss the roles of immune factors in the evolution of this pathological process. Finally, we highlight avenues of investigation for future research to broaden our understanding of the role of NGF in the pathogenesis of leprosy. Our analysis of the literature indicates that NGF plays an important role in the evolution and outcome of *Mycobacterium leprae* infection. The findings described here highlight an important area of investigation, as leprosy is one of the main causes of infection in the peripheral nervous system.

Keywords: nerve growth factor, leprosy, *Mycobacterium leprae*, pathogenesis, immunology

INTRODUCTION

The most extensively studied neurotrophin is nerve growth factor (NGF). This growth factor consists of 118 amino acids and has a molecular weight of 130 kDa. NGF was discovered as a growth factor that participates in the responses of sympathetic and sensory neurons regulating differentiation, neuronal regeneration, and the perception of pain (**Figure 1**) (1–5). In sensory neurons, NGF specifically binds to the TrkA receptor with high affinity and to receptor p75 with low affinity. The binding of NGF to receptor p75 seems to optimize the activity of TrkA (**Figure 2**) (6, 7). In skin and immune cells, NGF is produced by the proteolytic cleavage of its precursor, proNGF. proNGF is translocated to the lumen of the endoplasmic reticulum, transported through the exocytic pathway, and converted to its active mature biological form, NGF (3). The roles of NGF in physiological and pathological processes have been studied in several systems. Specifically, the significance of this molecule in responses to infectious and inflammatory diseases has recently been investigated in experimental



models of Chagas disease, respiratory syncytial virus infection, herpes simplex virus infection, myalgic encephalomyelitis, and other diseases (8–15).

DEFINITION AND BIOLOGICAL ASPECTS

Nerve growth factor was first described by Rita Levi-Montalcini (16), who showed its importance in the development, differentiation, maturation, and preservation of the integrity of sympathetic and sensory neurons (17). NGF is involved in modulating the sensitivity of peripheral nerve fibers to heat and pain in physiological and pathological events, such as genetic, metabolic (diabetes

mellitus), and infectious neuropathies (18). Further supporting a relationship between NGF and leprosy, Scully and Otten and others by previous studies reported the involvement of NGF in sympathetic and sensory neuron apoptosis (Figures 1 and 2) (19–21).

Nerve growth factor is endogenously produced as proNGF during development and into maturity by immune and nerve cells, as well as peripheral effector cells, such as keratinocytes, melanocytes, smooth muscle cells, fibroblasts, and Schwann cells (5, 17, 18). It is also synthesized in other organs, such as the gonads, thyroid, parathyroid, and exocrine glands (e.g., salivary glands) (5, 21, 22). The expression and receptor binding affinity of NGF, as well as the duration and intensity of cellular events

triggered by proNGF activation, determine its specific activity in effector cells or neurons (21, 23, 24).

Following its synthesis, NGF enters nerve endings by endocytosis, a process that permits NGF transport to the nerve cell body through contact with NGF receptors (NGF-R) located on the plasma membrane (25, 26). Several studies have described the roles of NGF in the central nervous system (CNS). In adults, NGF is involved in plasticity. However, the promotion of neuronal survival by NGF has only been observed in cholinergic neurons. NGF regulates the size of the cell body, dendritic branching, and neuronal connectivity (27, 28). In the peripheral nervous system, NGF acts on sympathetic and sensory neurons derived from the neural crest (29).

Nerve growth factor has been suggested to be involved in apoptosis of sympathetic and sensory neurons (20, 21). The binding of NGF to glial cells (i.e., oligodendrocytes and Schwann cells) through receptor p75 induces proapoptotic effects. However, these effects depend on NGF binding intensity and Trk receptor interactions (1, 30). Receptor p75 is a low-affinity accessory receptor for the TrkA receptor family to which all neurotrophins can bind. Receptor p75 seems to modulate the activity of TrkA and its intracellular signaling cascade. These receptors are present not only in nerve cells and CNS tumors, but also in immune cells such as macrophages, lymphocytes, and mast cells. These cells synthesize, store, and secrete proNGF, which suggests a possible role for NGF in the regulation of immune responses during inflammatory, infectious, and autoimmune processes (31, 32).

Thus, in addition to its primary functions in the nervous system, NGF participates in inflammatory processes and immune responses. NGF concentrations are increased during tissue inflammatory processes. Increased NGF levels can produce hyperalgesia through direct activation of nociceptors, which leads to CNS activation and neurogenic inflammation. Moreover, this process leads to local events comprising activation-induced release of histamine and increased numbers of mast cells and other immune system cells (7, 19).

Labouyrie et al. described the relationship between NGF and cells in normal and pathologic human lymphoid tissues and showed that NGF is involved in inflammatory or lymphoproliferative disorders (33). Immunoreactivity to the NGF-R (TrkA) has been observed in tissues, such as the thymic epithelium, cryptic tonsillar epithelium, epithelioid cells, multinucleated macrophages, and follicular dendritic cells. These findings demonstrate a relationship between NGF and the immune system. Owing to its production and activity in immune cells, NGF is thought by some to be a cytokine that participates in immune system events controlling the immune response in inflammatory and infectious processes (32, 34).

Different inflammatory and autoimmune diseases lead to altered expression of NGF. Increased anti-NGF antibody levels have been detected in patients with rheumatoid arthritis, systemic lupus erythematosus, and thyroiditis and are thought to contribute to the immune dysfunction and nerve damage observed in these diseases (21, 35, 36). However, the relationship between NGF autoantibodies and NGF expression remains unclear. Further studies are needed to investigate the effects of NGF autoantibodies on the expression of NGF and its receptors in autoimmune disease.

Other studies have shown an association between NGF and immune response mechanisms (37–42). For instance, Lambiase et al. have investigated the expression of NGF and the TrkA receptor in CD4+ T cells (38). Santambrogio et al. have shown that B cells express and secrete NGF, which in turn regulates the expression and secretion of calcitonin gene-related peptide by these cells *via* the TrkA receptor. This process has previously been described for sensory neurons (39). The immune response components interferon- β and interleukin (IL)-1 β increase NGF-R expression and activate its signaling pathways in neurons and participate in the control of apoptosis in these cells (40). T and B lymphocytes, dendritic cells, and macrophages express NGF, TrkA, and the p75 neurotrophin receptor (**Figure 2**). In the context of innate immunity, NGF is related to the activation of IL-1 β , nod-like receptor protein 1, NLRP3, and caspase-1 and contributes to inflammasome activation. TNF- α can induce differentiation and neuronal maturation through its interaction with NGF, which is in turn involved in the neuronal survival process (41, 42). Nevertheless, further investigation is needed to better elucidate the role of NGF in controlling the immune response in other cell types (41–43).

NGF, IMMUNE RESPONSE, AND LEPROSY

Leprosy is a chronic disease caused by *Mycobacterium leprae*, an intracellular bacillus that leads to loss of sensibility, innervation, intra-epidermal damage, and lesions associated with the loss of myelin in Schwann cells (44). Clinically, the different forms of the disease are characterized in part by the immune response patterns of the host (19, 44). According to the classification by Ridley and Jopling, leprosy has five clinical forms based on clinical, histopathological, immunological, and bacilloscopic criteria: borderline-tuberculoid, borderline-lepromatous, borderline-borderline, tuberculoid, and lepromatous (45). The initial stage of infection is referred to as the indeterminate form. The World Health Organization (WHO) classifies patients with leprosy as multibacillary or paucibacillary for treatment purposes. These classifications are made based on the identification of the bacilli and the number of lesions. Bacilloscopy is not always possible. Therefore, according to the WHO, the number of lesions can be used to classify patients into three groups: paucibacillary single-lesion leprosy (one skin lesion), paucibacillary leprosy (two to five skin lesions), and multibacillary leprosy (more than five skin lesions) (46).

In the tuberculoid form of the disease, the lesions are granulomatous and the individual displays an intense cell-mediated immune response (Th1) that prevents the proliferation of the bacillus. In the lepromatous form, the cell-mediated immune response is characterized by an anti-inflammatory cytokine profile (Th2) that contribute to multiplication of the bacillus in macrophage phagosomes. In the borderline form, the patients exhibit immunological and histopathological characteristics that vary between those of the tuberculoid and lepromatous forms (47).

Nerve injury-associated tissue damage is arguably the most important clinical consequence of leprosy (47). In the process of leprosy-associated neuropathy, the presence of bacilli in nerve endings and Schwann cells induces a response mediated by macrophages and other cells that eventually leads to the appearance

of immune-mediated lesions. In this process, cytokines, such as TNF- α , IL-6, and IL-17, may contribute to the evolution of neural lesions and deformities that are characteristic of some forms of the disease. The immune response and inflammation are not only defined by the presence of chemical mediators, such as cytokines and chemokines. Rather, inflammatory edema is also very important in the neuropathy associated with leprosy, which can induce degeneration of neural fibers. There are strong positive correlations among the levels of NGF, NGF-R, and TGF- β in patients with leprosy. This indicates that the above factors have synergistic actions that reduce tissue damage resulting from nerve injury (48). Study by Antunes et al. (48) in patients with the neuritic form of leprosy (pure neural form or primary neural form), it was observed that the NGF-R immunoreexpression was lower in nerve fibers and Schwann cells when compared to normal controls. In this study, hypoesthesia was correlated with decreased expression of NGF-R and protein gene product (PGP) 9.5, and electroencephalographic changes were observed in patients with altered immunolabeling for neurofilaments and PGP 9.5. These data point to a key role of NGF in the pathogenesis of neural lesions in leprosy.

Leprosy is the most common cause of non-traumatic neuropathy and is a classic example of an infectious neurodegenerative disease of the peripheral nervous system. It is estimated that more than one-fourth of patients with leprosy have some degree of disability and that about half of these patients have grade 2 disability corresponding to permanent neurological damage (44–59).

Studies have shown that different levels of NGF are associated with lepromatous and tuberculoid leprosy such that higher levels of NGF are associated with lepromatous forms and low levels of NGF are associated with tuberculoid forms of the disease (60). Specifically, Facer et al. have demonstrated the importance of NGF in leprosy (44). In patients with leprosy with and without lesions, TrkA receptors were shown to be present in subepidermal fibers and TrkA receptor messenger RNA was produced in the skin. The authors of the above study also evaluated the integrity of nerve endings and found that the presence of NGF in keratinocytes was correlated with deficient thermal sensation. Anand et al. detected low NGF levels in nerve and skin lesions of patients with leprosy and demonstrated that these low NGF levels contributed to the loss of NGF-dependent nociceptive fibers in damaged skin (19). Another study by Anand demonstrated that the loss of interaction between keratinocytes and nerves in affected skin drastically reduced the flow of NGF (61). Schwann cells produce NGF in response to axonal degeneration. However, while the levels of NGF are sharply reduced in the affected nerve trunks in patients with neuropathic lesions, there is a local increase in NGF levels in patients with chronic cutaneous hyperalgesia (61). The use of anti-NGF antibodies may be effective in treating hyperalgesia in patients with neuropathy and compromised nerve endings. In addition, physiological combinations of NGF, NT-3, and glial cell line-derived neurotrophic factor may assist in the reestablishment of homeostasis, and may thus be used in the treatment of neuropathic pain (61).

Other studies have suggested that NGF may restore pain sensitivity. Thus, NGF could play an important role in the prevention of ulcerations resulting from nociception loss, as observed

in leprosy and other peripheral neuropathies (50, 61). Some reports indicate that anti-NGF treatment may promote analgesia in patients with hyperalgesia, which suggests a modulatory role for NGF in nociception (61).

Immunostaining of damaged tissues using anti-NGF and anti-NGF-R antibodies revealed higher expression of NGF in patients with lepromatous leprosy and lower expression of NGF in those with the indeterminate form of the disease (49). In lepromatous forms of leprosy, higher mean expression levels of NGF and its receptor are associated with larger and more diffuse lesion patterns and greater nerve involvement. The presence of NGF is generally more apparent in patients with the highest probability of nerve damage (i.e., those with lepromatous leprosy and young patients) (Figure 3) (52, 53).


Several important findings have elucidated the role of NGF as an immune response mediator (38). In addition to its well-documented involvement in the differentiation and growth of neurons, there is growing evidence that NGF exerts a broad spectrum of effects on immune cells. As a result, NGF is considered a pleiotropic molecule involved in different functions (i.e., neuro-peptide modulation and tissue healing) that influence neural development, immune function, and injury responses (54).

Imbalance in the proNGF/NGF ratio, increased expression levels of TNF- α and p75 neurotrophin receptor, and impaired TrkA phosphorylation have been reported in microvascular preparations from Cre-proNGF transgenic mice when compared to normal control animals (43). TNF- α can induce differentiation and neuronal maturation through its interaction with NGF, which is in turn involved in the neuronal survival process (62). Endogenous TNF- α expression induced by NGF leads to a positive feedback loop comprising Akt activation through TNF receptor 2. This pathway promotes the survival of normal neural cells. Some studies suggest that TNF- α and IL-1 β control the activity of NGF in human synovial fibroblasts modulating the physiopathology of osteoarthritis (41, 62).

The relationship between NGF and TGF- β in glial cells has been described in a study on mice and rats with spinal injury (37). Following injury, TGF- β 1 levels increased, which in turn led to NGF messenger RNA and protein expression in the glial cells of these animals. In leprosy, the main target of *M. leprae* is the Schwann cell, which is the predominant cell type in the peripheral nervous system. Degradation of Schwann cells by *M. leprae* favors the development of peripheral neuropathy, which is the leading cause of morbidity in patients with leprosy. NGF may act as a protective factor for Schwann cells, and low levels of NGF may contribute to the development of neuropathy (38, 39).

Studies of systemic diseases (i.e., diabetes mellitus and osteoarthritis) have revealed an interaction between NGF and TGF- β (62, 63). Neurotrophins play a crucial role in the differentiation and survival of nerve cells. The characteristic positive correlations among NGF, NGF-R, and TGF- β in the clinical forms of leprosy highlight the interdependence of these factors (49).

Studies of lesions of patients with borderline leprosy have revealed strong correlations between TGF- β and NGF, and TGF- β and NGF-R ($r = 0.8722$ and $r = 0.7257$, respectively), with highly significant p -values for the two correlations ($p < 0.0001$ and



$p = 0.0015$, respectively) (48, 49). The borderline form of leprosy is immunologically dynamic and oscillates between the two polar forms. In patients with borderline-tuberculoid, borderline, and borderline-lepromatous leprosy, the progressive reduction in the cell-mediated response from the borderline-tuberculoid to the borderline-lepromatous form is accompanied by more extensive skin lesions and nerve damage, as well as increased bacillary burden and antibody levels (63, 64). In borderline forms of the disease, neurological manifestations resulting from immunological instability are characterized by nerve trunk impairment and frequent reactional episodes, which lead to early and asymmetrical nerve injuries and physical disability. This process is caused by increased numbers of bacilli in nerve branches close to Schwann cells (49). Cunha analyzed the relationships between the clinical forms of leprosy and episodes of neuritis. They found that patients with the borderline form of leprosy are 2.69 times more likely to progress to neuritis than those with the lepromatous form of the disease (65, 66).

During healing, NGF activates processes involved in the restoration of innervation (65, 67). In addition to inducing fibroblast migration, NGF exerts proliferative and antiapoptotic effects on keratinocytes and endothelial cells (68, 69). Moreover, NGF may be important for potentiating injury-specific responses through proinflammatory effects on neutrophils, eosinophils, mast cells, and T lymphocytes (67). The interactions of NGF in the tissue microenvironment are complex, and its relation to TNF- α , which can induce apoptosis in Schwann cells by binding to specific death receptors, may lead to antagonistic effects. This is because NGF can activate survival signals in the target cell. In fact, the same cytokine can have antagonistic effects depending on its interactions with specific receptors and the intracellular cascade activated after the activation of these receptors (Figure 3) (41–43, 62).

Cell-mediated immune responses may be beneficial against bacterial infections; however, inflammation can lead to irreversible

tissue damage. Nerve injury occurs in approximately 10% of patients with paucibacillary leprosy and 40% of patients with multibacillary leprosy and is particularly acute in patients with reverse reactions (69, 70).

TGF- β participates in the tissue repair process as an anti-inflammatory agent during intense inflammation by inducing nerve and tissue regeneration (67, 71). Higher TGF- β expression in patients with the lepromatous form of the disease is associated with a higher frequency of apoptosis in lesions, especially within Schwann cells (72). The positive correlation between TGF- β expression and NGF expression may be associated with the protection and regeneration of nerve endings (60).

In lepromatous leprosy, neurological manifestations progress slowly over many years and involve small nerve branches and multiple nerve trunks within the skin (73, 74). Peripheral nerve injury, or any pathological condition that causes an interruption between the target organ and the nerve cell body, acts as a signal that induces non-neural cell populations (e.g., fibroblasts) to produce NGF. The induction of NGF synthesis in these cells is also modulated by cytokines, which invade the site of nerve injury where nerve regeneration is initiated (74).

The expression of p75 receptor in peripheral nerve cells is induced by loss of contact between the target organ and the axon (60). Some studies suggest that the p75 receptor is involved in axonal NGF uptake, reflecting the importance of NGF in efficient nerve regeneration. Patients with lepromatous leprosy have nerve trunk lesions, as well as multiple mononeuropathies and polyneuropathies (33, 71–76). The relationship between NGF and TGF- β is a key determinant of the actions of NGF in patients with the lepromatous form of the disease. Coordination between NGF and TGF- β responses in inflammatory processes following tissue damage is thus fundamentally important in remodeling and tissue repair (Figure 3) (60).

NEED FOR FUTURE RESEARCH

Prospective cohort studies and intervention studies evaluating patients with different clinical forms of the disease and different reactive states (erythema nodosum and reverse reaction types) may help better elucidate the relationship between NGF and the immune response and the factors contributing to the protection and regeneration of nerves affected during infection. Further analysis of the relationship among tissue levels of NGF and a large panel of pro- and anti-inflammatory cytokines, blood levels of NGF, and the immune response is important for a better understanding of the involvement of NGF in the pathophysiology of chronic granulomatous peripheral nerve infections, especially leprosy.

CONCLUSION

This review demonstrates that, within lesions associated with leprosy, NGF and TGF- β respond to inflammatory processes and

REFERENCES

- Khan N, Smith MT. Neurotrophins and neuropathic pain: role in pathobiology. *Molecules* (2015) 2:10657–88. doi:10.3390/molecules200610657
- Hallbook F. Evolution of the vertebrate neurotrophin and Trk receptor gene families. *Curr Opin Neurobiol* (1999) 9:616–21. doi:10.1016/S0959-4388(99)00011-2
- Ibanez CF. Emerging themes in structural biology of neurotrophic factors. *Trends Neurosci* (1998) 21:438–44. doi:10.1016/S0166-2236(98)01266-1
- Terenghi G. Peripheral nerve regeneration and neurotrophic factors. *J Anat* (1999) 194:1–14. doi:10.1046/j.1469-7580.1999.19410001.x
- Sofroniew MV, Howe CL, Mobley WC. Nerve growth factor signaling, neuroprotection, and neural repair. *Annu Rev Neurosci* (2001) 24:1217–81. doi:10.1146/annurev.neuro.24.1.1217
- Hetman M, Xia Z. Signaling pathways mediating anti-apoptotic action of neurotrophins. *Acta Neurobiol Exp* (2000) 60:531–45.
- Wang H, Wang R, Thrimawithana T, Little PJ, Xu J, Feng ZP, et al. The nerve growth factor signaling and its potential as therapeutic target for glaucoma. *Biomed Res Int* (2014) 2014:759473. doi:10.1155/2014/759473
- Oliveira NK, Ferreira RN, Lopes SDN, Chiari E, Camargos ERDS, Martinelli PM. Cardiac autonomic denervation and expression of neurotrophins (NGF and BDNF) and their receptors during experimental Chagas disease. *Growth Factors* (2017) 35:161–70. doi:10.1080/08977194.2017.1395420
- Aridgides D, Salvador R, Pereira Perrin M. *Trypanosoma cruzi* coaxes cardiac fibroblasts into preventing cardiomyocyte death by activating nerve growth factor receptor TrkA. *PLoS One* (2013) 8:e57450. doi:10.1371/journal.pone.0057450
- Martinelli PM, Camargos ER, Azevedo AA, Chiari E, Morel G, Machado CR. Cardiac NGF and GDNF expression during *Trypanosoma cruzi* infection in rats. *Auton Neurosci* (2006) 130:32–40. doi:10.1016/j.autneu.2006.05.004
- Wu X, Zhou X, Hu Y, Liu C, Wang J. Neutralization of nerve growth factor (NGF) inhibits the Th2 response and protects against the respiratory syncytial virus (RSV) infection. *Immunol Res* (2017) 65:721–8. doi:10.1007/s12026-017-8909-z
- Yanez AA, Harrell T, Sriranganathan HJ, Ives AM, Bertke AS. Neurotrophic factors NGF, GDNF and NTN selectively modulate HSV1 and HSV2 lytic infection and reactivation in primary adult sensory and autonomic neurons. *Pathogens* (2017) 6:E5. doi:10.3390/pathogens6010005
- Chen ZW, Wang HP, Yuan FM, Zhang X, Dong XJ, Xie RS, et al. Releasing of herpes simplex virus carrying NGF in subarachnoid space promotes the functional repair in spinal cord injured rats. *Curr Gene Ther* (2016) 16:263–70. doi:10.2174/1566523217666161121105717
- Cabrera JR, Viejo-Borbolla A, Alcamí A, Wandosell F. Secreted herpes simplex virus-2 glycoprotein G alters thermal pain sensitivity by modifying NGF effects on TRPV1. *J Neuroinflammation* (2016) 13:210. doi:10.1186/s12974-016-0677-5
- Montoya JG, Holmes TH, Anderson JN, Maecker HT, Rosenberg-Hasson Y, Valencia IJ, et al. Cytokine signature associated with disease severity in chronic fatigue syndrome patients. *Proc Natl Acad Sci U S A* (2017) 114:E7150–8. doi:10.1073/pnas.1710519114
- Levi-Montalcini R, Hamburger VA. A diffusible agent of mouse sarcoma, producing hyperplasia of sympathetic ganglia and hyperneurotization of viscera in the chick embryo. *J Exp Zoolology* (1953) 123:233–87. doi:10.1002/jez.1401230203
- Levi-Montalcini R, Stephen SD, Toso RD, Petrelli L, Leon A. Nerve growth factor: from neurotrophin to neurokine. *Trends Neurosci* (1996) 19:514–20. doi:10.1016/S0166-2236(96)10058-8
- Lewin GR, Barde YA. Physiology of neurotrophins. *Annu Rev Neurosci* (1996) 19:289–317. doi:10.1146/annurev.ne.19.030196.001445
- Anand P, Pandya S, Ladiwala U, Singhal B, Sinicropi DV, Williams-Chestnut RE. Depletion of nerve growth factor in leprosy. *Lancet* (1994) 344:129–30. doi:10.1016/S0140-6736(94)91316-1
- Scully JL, Otten U. NGF. *Cell Biol Int* (1995) 19:459–69. doi:10.1006/cbir.1995.1089
- Ioannou MS, Fahnestock M. ProNGF, but not NGF, switches from neurotrophic to apoptotic activity in response to reductions in TrkA receptor levels. *Int J Mol Sci* (2017) 18:599. doi:10.3390/ijms18030599
- Vega JA, García-Suárez O, Hannestad J, Pérez-Pérez M, Germanà A. Neurotrophins and the immune system. *J Anat* (2003) 203:1–19. doi:10.1046/j.1469-7580.2003.00203.x
- Patapoutian A, Reichardt LF. Trk receptors: mediators of neurotrophin action. *Curr Opin Neurobiol* (2001) 11:272–80. doi:10.1016/S0959-4388(00)00208-7
- Aloe L, Rocco ML, Balzamino BO, Micera A. Nerve growth factor: a focus on neuroscience and therapy. *Curr Neuropharmacol* (2015) 13:294–303. doi:10.2174/1570159X13666150403231920
- Harper GP, Thoenen H. Target cells, biological effects, and mechanism of action of nerve growth factor and its antibodies. *Annu Rev Pharmacol Toxicol* (1981) 21:205–29. doi:10.1146/annurev.pa.21.040181.001225
- Wang W, Chen J, Guo X. The role of nerve growth factor and its receptors in tumorigenesis and cancer pain. *Biosci Trends* (2014) 8:68–74. doi:10.5582/bst.8.68
- Rush RA, Chie E, Liu D, Tafreshi A, Zetler C, Zouh XF. Neurotrophic factors are required by mature sympathetic neurons for survival, transmission and connectivity. *Clin Exp Pharmacol Physiol* (1997) 24:549–55. doi:10.1111/j.1440-1681.1997.tb02089.x
- Mendell LM, Munson JB, Arvanian VL. Neurotrophins and synaptic plasticity in the mammalian spinal cord. *J Physiol* (2001) 533:91–7. doi:10.1111/j.1469-7793.2001.0091b.x
- Aloe L, Rocco ML, Balzamino BO, Micera A. Nerve growth factor, differentiation and controlling cancer cell development. *J Exp Clin Cancer Res* (2016) 35:116. doi:10.1186/s13046-016-0395-y

tissue damage while triggering tissue remodeling. Further studies are needed to elucidate the broad role of NGF in the pathogenesis of leprosy. Although our understanding of the effects of NGF on nerve damage has increased, further insights into the functional roles of NGF (and other neurotrophins) in normal skin and during disease progression are needed.

AUTHOR CONTRIBUTIONS

TA, JS, AF, LF, and JQ conceived and wrote the manuscript.

FUNDING

JQ is a Research Productivity Fellow and Senior Postdoctoral Fellow at the Brazilian National Council for Scientific and Technological Development—CNPq/Brazil (Grants number 302553/2015-0 and 116427/2016-7). Program of support for qualified production—PAPQ/UFGA.

30. Berry A, Bindocci E, Alleva E. NGF, brain and behavioral plasticity. *Neural Plast* (2012) 2012:784040. doi:10.1155/2012/784040
31. Bracci-Laudiero L, De Stefano ME. NGF in early embryogenesis, differentiation, and pathology in the nervous and immune systems. *Curr Top Behav Neurosci* (2016) 29:125–52. doi:10.1007/7854_2015_420
32. Niederhauser O, Mangold M, Schubel R, Kuszniir EA, Schmidt D, Hertel C. NGF ligand alters NGF signaling via p75(NTR) and TrkA. *J Neurosci Res* (2000) 1:263–72. doi:10.1002/1097-4547(20000801)61:3<263::AID-JNR4>3.0.CO;2-M
33. Labouyrie E, Parrens M, de Mascarel A, Bloch B, Merlio JP. Distribution of NGF receptors in normal and pathologic human lymphoid tissues. *J Neuroimmunol* (1997) 77:161–73. doi:10.1016/S0165-5728(97)00055-6
34. Rook G, Lightman SL, Heijnen CJ. Can nerve damage disrupt neuroendocrine immune homeostasis? Leprosy as a case in point. *Trends Immunol* (2002) 23:18–22. doi:10.1016/S1471-4906(01)02090-7
35. Dicou E, Hurez D, Nerriere V. Natural antibodies against the nerve growth factor in autoimmune diseases. *J Neuroimmunol* (1993) 47:159–68. doi:10.1016/0165-5728(93)90026-U
36. Gartner A, Shostak Y, Hackel N, Ethell IM, Thoenen H. Ultrastructural identification of storage compartments and localization of activity-dependent secretion of neurotrophin 6 in hippocampal neurons. *Mol Cell Neurosci* (2000) 15:215–34. doi:10.1006/mcne.1999.0825
37. Yu G, Fahnestock M. Differential expression of nerve growth factor transcripts in glia and neurons and their regulation by transforming growth factor-beta1. *Brain Res Mol Brain Res* (2002) 30:115–25. doi:10.1016/S0169-328X(02)00399-6
38. Lambiase A, Bracci-Laudiero L, Bonini S, Bonini S, Starace G, D'Elia MM, et al. Human CD4+ T cell clones produce and release nerve growth factor and express high-affinity nerve growth factor receptors. *J Allergy Clin Immunol* (1997) 100:408–14. doi:10.1016/S0091-6749(97)70256-2
39. Santambrogio L, Benedetti M, Chao MV, Muzaffar R, Kulig K, Gabellini N, et al. Nerve growth factor production by lymphocytes. *J Immunol* (1994) 153:4488–92.
40. Dedoni S, Olianias MC, Ingiani A, Onali P. Type I interferons up-regulate the expression and signalling of p75 NTR/TrkA receptor complex in differentiated human SH-SY5Y neuroblastoma cells. *Neuropharmacology* (2014) 79:321–34. doi:10.1016/j.neuropharm.2013.12.002
41. Choi S, Friedman WJ. Interleukin-1 β enhances neuronal vulnerability to proNGF-mediated apoptosis by increasing surface expression of p75(NTR) and sortilin. *Neurosci* (2014) 257:11–9. doi:10.1016/j.neuroscience.2013.10.058
42. Datta-Mitra A, Kundu-Raychaudhuri S, Mitra A, Raychaudhuri SP. Cross talk between neuroregulatory molecule and monocyte: nerve growth factor activates the inflammasome. *PLoS One* (2015) 10:e0121626. doi:10.1371/journal.pone.0121626
43. Mohamed R, Coucha M, Elshaer SL, Artham S, Lemtali T, El-Remessy AB. Inducible overexpression of endothelial proNGF as a mouse model to study microvascular dysfunction. *Biochim Biophys Acta* (2017) 1864:746–57. doi:10.1016/j.bbdis.2017.12.023
44. Facer P, Mann D, Mathur R, Pandya S, Ladiwala U, Singhal B, et al. Do nerve growth factor-related mechanisms contribute to loss of cutaneous nociception in leprosy? *Pain* (2000) 85:231–8. doi:10.1016/S0304-3959(99)00273-0
45. Ridley DS, Jopling WH. Classification of leprosy according to immunity. A five-group system. *Int J Lepr Other Mycobact Dis* (1966) 34:255–73.
46. Pardillo FE, Fajardo TT, Abalos RM, Scollard D, Gelber RH. Methods for the classification of leprosy for treatment purposes. *Clin Infect Dis* (2007) 44:1096–9. doi:10.1086/512809
47. de Sousa JR, Sotto MN, Simões Quaresma JA. Leprosy as a complex infection: breakdown of the Th1 and Th2 immune paradigm in the immunopathogenesis of the disease. *Front Immunol* (2017) 8:1635. doi:10.3389/fimmu.2017.01635
48. Antunes SL, Chimelli LM, Rabello ET, Valentim VC, Corte-Real S, Sarno EN, et al. An immunohistochemical, clinical and electroneuro-myographic correlative study of the neural markers in the neuritic form of leprosy. *Braz J Med Biol Res* (2006) 39:1071–81. doi:10.1590/S0100-879X2006000800010
49. Aarão TL, Esteves NR, Esteves N, Soares LP, Pinto Dda S, Fuzii HT, et al. Relationship between growth factors and its implication in the pathogenesis of leprosy. *Microb Pathog* (2014) 77:66–72. doi:10.1016/j.micpath.2014.10.005
50. Chang DS, Hsu E, Hottinger DG, Cohen SP. Anti-nerve growth factor in pain management: current evidence. *J Pain Res* (2016) 9:373–83. doi:10.2147/JPR.S89061
51. Aloe L, Rocco ML, Bianchi P, Manni L. Nerve growth factor: from the early discoveries to the potential clinical use. *J Transl Med* (2012) 10:239. doi:10.1186/1479-5876-10-239
52. Schreuder PA. The occurrence of reactions and impairments in leprosy: experience in the leprosy control program of three provinces in northeastern Thailand, 1987–1995 [correction of 1978–1995]. II. Reactions. *Int J Lepr Other Mycobact Dis* (1998) 66:159–69.
53. Saunderson P. The epidemiology of reactions and nerve damage. *Leprosy Rev* (2000) 71:106–10. doi:10.5935/0305-7518.20000079
54. Richardus JH, Nicholls PG, Croft RP, Withington SG, Smith WC. Incidence of acute nerve function impairment and reactions in leprosy: a prospective cohort analysis after 5 years of follow-up. *Int J Epidemiol* (2004) 33:337–43. doi:10.1093/ije/dyg225
55. Nico B, Mangieri D, Benagiano V, Crivellato E, Ribatti D. Nerve growth factor as an angiogenic factor. *Microvasc Res* (2008) 75:135–41. doi:10.1016/j.mvr.2007.07.004
56. Job CK. Nerve damage in leprosy. *Int J Lepr Other Mycobact Dis* (1989) 57:532–9.
57. Bryceson A, Pfaltzgraff RE. Complications due to nerve damage. In: Hastings RC, editor. *Medicine in the Tropics: Leprosy*. 3rd ed. Edinburgh: Churchill Livingstone (1990). p. 133–51.
58. Chimelli L, Freitas M, Nascimento O. Value of nerve biopsy in the diagnosis and follow-up of leprosy: the role of vascular lesions and usefulness of nerve studies in the detection of persistent bacilli. *J Neurol* (1997) 244:318–23. doi:10.1007/s004150050094
59. Rose P, Water MF. Reversal reactions in leprosy and their management. *Leprosy Rev* (1991) 62:113–21. doi:10.5935/0305-7518.19910013
60. Ooi WW, Srinivasan J. Leprosy and peripheral nervous system: basic and clinical aspects. *Muscle Nerve* (2004) 30:393–409. doi:10.1002/mus.20113
61. Anand P. Neurotrophic factors and their receptors in human sensory neuropathies. *Prog Brain Res* (2004) 146:477–92. doi:10.1016/S0079-6123(03)46030-5
62. Takei Y, Laskey R. Intracellular and intercellular cross talk between NGF and TNF. *Adv Exp Med Biol* (2011) 691:559–65. doi:10.1007/978-1-4419-6612-4_58
63. Verge VM, Andreassen CS, Arnason TG, Andersen H. Mechanisms of disease: role of neurotrophins in diabetes and diabetic neuropathy. *Handb Clin Neurol* (2014) 126:443–60. doi:10.1016/B978-0-444-53480-4.00032-1
64. Takano S, Uchida K, Inoue G, Miyagi M, Aikawa J, Iwase D, et al. Nerve growth factor regulation and production by macrophages in osteoarthritic synovium. *Clin Exp Immunol* (2017) 190:235–43. doi:10.1111/cei.13007
65. Cunha AZS. Leprosy: evolution aspects of its diagnosis, treatment and control. *Cien Saude Colet* (2002) 2:235–42. doi:10.1590/S1413-81232002000200004
66. Britton WJ, Lockwood DNJ. Leprosy. *Lancet* (2004) 363:1209–19. doi:10.1016/S0140-6736(04)15952-7
67. Harsum S, Clarke JD, Martin P. A reciprocal relationship between cutaneous nerves and repairing skin wounds in the developing chick embryo. *Dev Biol* (2001) 238:27–39. doi:10.1006/dbio.2001.0395
68. Ching Y, Sutton TL, Pierpont YN, Robson MC, Payne WG. The use of growth factors and other humoral agents to accelerate and enhance burn wound healing. *Eplasty* (2011) 11:e41.
69. Pincelli C. Nerve growth factor and keratinocytes: a role in psoriasis. *Eur J Dermatol* (2000) 10:85–90.
70. Raychaudhuri SK, Raychaudhuri SP, Weltman H, Farber EM. Effect of nerve growth factor on endothelial cell biology: proliferation and adherence molecule expression on human dermal microvascular endothelial cells. *Arch Dermatol Res* (2001) 293:291–5. doi:10.1007/s004030100224
71. Spierings E, Boer T, Zulianello L, Ottenhoff TH. The role of Schwann cells, T cells and *Mycobacterium leprae* in the immunopathogenesis of nerve damage in leprosy. *Leprosy Rev* (2000) 71:121–9. doi:10.5935/0305-7518.20000083

72. Oliveira RB, Sampaio EP, Aarestrup F, Teles RM, Silva TP, Oliveira AL, et al. Cytokines and *Mycobacterium leprae* induce apoptosis in human Schwann cells. *J Neuropathol Exp Neurol* (2005) 64:882–90. doi:10.1097/01.jnen.0000182982.09978.66
73. Spierings E, De Boer T, Zulianello L, Ottenhoff TH. Novel mechanisms in the immunopathogenesis of leprosy nerve damage: the role of Schwann cells, T cells and *Mycobacterium leprae*. *Immunol Cell Biol* (2000) 78:349–55. doi:10.1046/j.1440-1711.2000.00939.x
74. Bedi A, Maak T, Walsh C, Rodeo SA, Grande D, Dines DM, et al. Cytokines in rotator cuff degeneration and repair. *J Shoulder Elbow Surg* (2012) 21:218–27. doi:10.1016/j.jse.2011.09.020
75. Araujo MG. Leprosy in Brazil. *Rev Soc Bras Med Trop* (2003) 36:373–82. doi:10.1590/S0037-86822003000300010
76. Singh N, Birdi TJ, Antia NH. Nerve growth factor production and expression of p75 by Schwann cells and neurofibroblasts in response to *M. leprae* infection

and macrophage secretory products. *Neuropathol Appl Neurobiol* (1997) 23:59–67. doi:10.1111/j.1365-2990.1997.tb01186.x

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Aarão, de Sousa, Falcão, Falcão and Quaresma. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.