



# Neurological manifestations of COVID-19: a systematic review and meta-analysis of proportions

T. T. Fava<sup>1</sup> · Priya Dev<sup>1</sup> · Rameshwar Nath Chaurasia<sup>1</sup> · Kamlesh Chakravarty<sup>2</sup> · Rahul Mishra<sup>3</sup> · Deepika Joshi<sup>1</sup> · Vijay Nath Mishra<sup>1</sup> · Anand Kumar<sup>1</sup> · Varun Kumar Singh<sup>1</sup> · Manoj Pandey<sup>4</sup> · Abhishek Pathak<sup>1</sup>

Received: 9 August 2020 / Accepted: 5 October 2020 / Published online: 21 October 2020  
© Fondazione Società Italiana di Neurologia 2020

## Abstract

**Background** Coronaviruses mainly affect the respiratory system; however, there are reports of SARS-CoV and MERS-CoV causing neurological manifestations. We aimed at discussing the various neurological manifestations of SARS-CoV-2 infection and to estimate the prevalence of each of them.

**Methods** We searched the following electronic databases; PubMed, MEDLINE, Scopus, EMBASE, Google Scholar, EBSCO, Web of Science, Cochrane Library, WHO database, and ClinicalTrials.gov. Relevant MeSH terms for COVID-19 and neurological manifestations were used. Randomized controlled trials, non-randomized controlled trials, case-control studies, cohort studies, cross-sectional studies, case series, and case reports were included in the study. To estimate the overall proportion of each neurological manifestations, the study employed meta-analysis of proportions using a random-effects model.

**Results** Pooled prevalence of each neurological manifestations are, smell disturbances (35.8%; 95% CI 21.4–50.2), taste disturbances (38.5%; 95% CI 24.0–53.0), myalgia (19.3%; 95% CI 15.1–23.6), headache (14.7%; 95% CI 10.4–18.9), dizziness (6.1%; 95% CI 3.1–9.2), and syncope (1.8%; 95% CI 0.9–4.6). Pooled prevalence of acute cerebrovascular disease was (2.3%; 95% CI 1.0–3.6), of which majority were ischaemic stroke (2.1%; 95% CI 0.9–3.3), followed by haemorrhagic stroke (0.4%; 95% CI 0.2–0.6), and cerebral venous thrombosis (0.3%; 95% CI 0.1–0.6).

**Conclusions** Neurological symptoms are common in SARS-CoV-2 infection, and from the large number of cases reported from all over the world daily, the prevalence of neurological features might increase again. Identifying some neurological manifestations like smell and taste disturbances can be used to screen patients with COVID-19 so that early identification and isolation is possible.

**Keywords** COVID-19 neurological manifestations · Acute cerebrovascular disease · SARS-CoV-2 infection · Meningoencephalitis · Guillain-Barré syndrome · Smell and taste disturbances

## Background

Coronaviruses are enveloped, positive-stranded RNA viruses that mainly cause respiratory and gastrointestinal tract infections

✉ Abhishek Pathak  
abhishekpathakaiims@gmail.com

<sup>1</sup> Department of Neurology, Institute of Medical Sciences, Banaras Hindu University, Varanasi, India

<sup>2</sup> Department of Neurology, PGIMER, Chandigarh, India

<sup>3</sup> Department of Statistics, Institute of science, Banaras Hindu University, Varanasi, India

<sup>4</sup> Department of Surgical Oncology, Institute of Medical Sciences, Banaras Hindu University, Varanasi, India

[1]. They are divided into four genera: alpha, beta, delta, and gamma. Alphacoronavirus and betacoronavirus cause human infections [1]. Betacoronaviruses are further divided into 4 clades, a–d [2]. SARS-CoV and MERS-CoV are betacoronaviruses which caused outbreaks in 2002 and 2012 respectively [3]. The likely reservoirs of SARS-CoV and MERS-CoV viruses were identified as bats [2]. SARS-CoV-2 is a coronavirus and is classified into the betacoronavirus 2b lineage; however, a distinct clade from the SARS-CoV and MERS-CoV [4, 5]. It has been postulated that reservoir of SARS-CoV-2 is also bats; however, more evidence is needed for proving the assumption [6]. The disease caused by SARS-CoV-2 is termed as COVID-19. COVID-19 outbreak started as a cluster of respiratory illnesses and the first case was reported from Wuhan, Hubei Province, China on 8th December [7, 8]. It was declared as a pandemic by WHO on March 11, 2020 [9].

The most common symptoms of COVID-19 are similar to other coronaviruses which include fever, fatigue, dry cough, anorexia, shortness of breath, myalgia, and headache [10–12]. Old age and co-morbidities are associated with higher mortality and morbidity as compared with younger patients and those without any co-morbidities [10, 12, 13].

The neuroinvasive and neurotropic potential of coronaviruses like SARS-CoV and MERS-CoV has been demonstrated in many previous studies [14, 15]. A similar mechanism is suggested for the SARS-CoV-2 also [16]. Neurological manifestations reported of SARS-CoV, MERS-CoV, and other coronaviruses include peripheral neuropathy [17], myopathies with elevated creatinine kinase [17], large vessel stroke [18], olfactory neuropathy/anosmia [19], meningoencephalitis [20, 21], post-infectious acute disseminated encephalomyelitis [22, 23], Bickerstaff's encephalitis overlapping with Guillain-Barré syndrome [24], and Guillain-Barré syndrome [24]. This review is aimed at discussing various neurological manifestations in COVID-19, including the frequency of neurological symptoms, morbidity, mortality, laboratory parameters, and imaging findings associated with patients with neurological symptoms. In the meta-analysis, we estimated the proportion of COVID-19 patients developing neurological manifestations.

## Methods

### Selection criteria and search strategy

We searched the following electronic databases for articles published between 1st December 2019 to 25th June 2020; PubMed, MEDLINE, Scopus, EMBASE, Google Scholar, EBSCO, Web of Science, Cochrane Library, WHO database, and [ClinicalTrials.gov](#). The MeSH terms and keywords used include: “COVID-19” OR “COVID 19” OR “SARS-CoV-2” OR “2019 novel coronavirus” OR “2019 nCoV” AND “Neurological” OR “Brain” OR “CNS features” OR “central nervous system features” OR “peripheral nervous system features” OR “neuropathy” OR “skeletal muscle” OR “myositis” OR “neuromuscular junction” OR “headache” OR “anosmia” OR “olfactory” OR “ageusia” OR “cranial neuropathy” OR “seizures” OR “encephalitis” OR “meningitis” OR “stroke” OR “cerebrovascular disease” OR “cerebral hemorrhage” OR “intracerebral hemorrhage” OR “cerebral infarct” OR “cortical venous thrombosis” OR “deep cerebral venous thrombosis” OR “impaired consciousness” OR “confusion” OR “weakness” OR “Guillain-Barre’ syndrome” OR “Miller Fisher syndrome” OR “ataxia” OR “myopathy” OR “myelitis” OR “myelopathy” with an additional filter of “studies in human subjects”. The search was done between 31st March 2020 and 25th June 2020. To ensure literature saturation, we inspected

the references of all studies included in this review. The protocol of this review was registered at PROSPERO (ID-CRD42020185593) prospectively in May 2020.

### Inclusion and exclusion criteria

All published randomized controlled trials, non-randomized controlled trials, case-control studies, cohort studies, cross-sectional studies, case series, and case reports, if they had sufficient data on neurological features, laboratory parameters, imaging findings were included in this review. Only those studies were included in which subjects were diagnosed with SARS-CoV-2 infection by real-time RT-PCR or high throughput sequencing analysis of swab specimens or serology or culture. We also included pre-prints and letters if they included data on neurological manifestations in COVID-19. Editorials, systematic reviews, meta-analysis, narrative reviews, conference abstracts, commentaries, animal studies, post-mortem studies, and where translation into English was not possible were excluded. The authors were contacted twice by email if any missing data in the articles.

### Data extraction and study quality assessment

Databases selected were searched independently by two members (TF and AP) in the team, and, following duplicate removal, reviewed all the articles and selected articles based on inclusion and exclusion criteria. Reporting was done according to the recommendations of the PRISMA statement [25]. Quality of the non-randomized studies was evaluated using the Newcastle-Ottawa Scale [26, 27] and the quality of one randomized controlled trial was assessed using the CONSORT criteria [28]. Any disagreements between two main reviewers were discussed with a third evaluator. Data about the author’s name, publication date, study setting and design, time and duration of the study, follow-up, the total number of patients evaluated, study population, age, gender, co-morbidities, neurological features, laboratory parameters, imaging findings, morbidity, and mortality were extracted.

### Outcome measures

Primary outcomes assessed were neurological manifestations in COVID-19 patients and its prevalence. For the categorical variables, simple and relative frequency and proportions were used. For continuous variables, central tendency (mean or median) and dispersion measures (standard error, standard deviation) were used. To measure association, risk ratios, odds ratios, and hazard ratios were used and 95% confidence intervals calculated. We also assessed secondary outcomes like the association of neurological manifestations with age, co-morbidities, lab parameters including CSF study, imaging features, length of hospital stay, ICU admission, time from onset

of typical COVID-19 symptoms to neurological manifestations, and morbidity/mortality.

### Strategy for data synthesis, statistical analysis for meta-analysis

Data synthesis and illustration were done in tables and figures. For the categorical variables, simple and relative frequency and proportions were used. For continuous variables, measures of central tendency (mean or median) and dispersion (standard error, standard deviation) were calculated. The primary aim of our study was to synthesize the findings from multiple studies that investigated the issues related to neurological manifestations in COVID-19 and thus provide a quantitative summary, to better direct future work. The data are available in the form of proportions, defined as the number of cases of interest in a sample with a particular characteristic divided by the size of the sample. To achieve the objective of obtaining a more precise estimate of the overall proportion for a certain event (neurological manifestations) related to COVID-19, the study employed meta-analysis of proportions using a random-effects model and by the DerSimonian-Laird method [29, 30]. We performed data analysis using meta-packages in R (version 3.5.0). Heterogeneity was assessed using the  $I^2$  value.  $I^2$  can take values from 0% to 100% and it is assumed that an  $I^2$  of 25%, 50%, and 75% indicate low, medium, and large heterogeneity respectively [31]. Forest plot was used to visualize the point estimates of study effects and their confidence intervals. Publication bias was evaluated using the funnel plot.

## Results

Among the 6789 articles identified, 212 studies were included in the systematic review and 74 studies in the meta-analysis (PRISMA flow diagram (Fig. 1)). Out of them, most were retrospective studies, 18 were cohort studies, 11 were prospective studies, nine were cross-sectional studies, one was a randomized controlled trial, one was a case-control study and the rest were all case series and case reports. Among these studies, we found only 19 studies, which investigated specifically neurological features in COVID-19 patients. Other studies, evaluated parameters in general. Table 1 shows a summary of all the observational studies included.

### Neurological manifestations

Neurological manifestations have been reported in patients with COVID-19 from all over the world. A multicentre, retrospective study by Mao et al. [32] was the first study to evaluate the neurological manifestations in COVID-19 and found that neurological manifestations were present in 36.4% of total

214 patients, out of which most common was CNS manifestations(24.8%) followed by peripheral nervous system manifestations(8.9%). Other large retrospective observational studies reported the incidence of neurological manifestations as 4.3% [45], 15% [47], and 57.4% [49]. The most common neurological manifestations reported in COVID-19 were smell disturbances, taste disturbances, headache, myalgia, and disturbances in consciousness/altered mental status. The prevalence of all the neurological manifestations assessed is given in Table 2. A summary estimate of pooled prevalence and heterogeneity of each neurological manifestation are given in Table 3. Forest plot and funnel plot is given in Figs. 2 and 3 respectively.

### Smell and taste disturbances

The overall incidence of smell disturbances in the studies ranged from 4.9–85.6% [49, 54] and the most common type of smell disturbance was anosmia. Other smell disturbances noticed were hyposmia, phantosmia, and parosmia [54]. Similarly, the incidence of taste disturbances reported was 0.3–88.8% [47, 54] and the most commonly reported were dysgeusia and ageusia. In the meta-analysis, we found 17 and 14 studies, which assessed the prevalence of smell and taste disturbances respectively and disturbances of smell (35.8%; 95%CI 21.4–50.2) and taste (38.5%; 95%CI 24.0–53.0) sensation were the most common neurological manifestation followed by non-specific neurological manifestations.

A case-control study of 79 COVID-19 patients and 40 historical controls of influenza patients from Spain [52] revealed that new-onset smell and taste disorders were significantly higher in the COVID-19 group. Patients in COVID-19 were significantly younger. Another study reported olfactory and taste disturbances occur more frequently in females than males [53]. Lechien et al. [54], Gilani S et al. [140], and Rachel Kaye et al. [141] reported that anosmia can be the initial and early manifestations of COVID-19. Population surveys on new-onset olfactory dysfunction from Iran [142] and UK [143] have reported an increase in olfactory dysfunction during the COVID-19 pandemic.

### Non-specific symptoms

The most common non-specific neurological symptoms reported in SARS-CoV-2 infection were myalgia, headache, syncope, and dizziness. The overall pooled prevalence estimate of the proportion of cases are given in Table 3. Incidence of myalgia reported in various studies ranged from 1.8–62.5% [47, 111], headache from 0.6–70.3% [90, 111], and dizziness from 0.6–21% [47, 113]. In children, myalgia and dizziness were less common and rarely reported. In health care workers, the incidence of myalgia, headache, and dizziness was higher compared with the general

**Table 1** Characteristics of studies included and neurological manifestations

First author	Article type	Study setting	Type of study	Enrolment date	Follow-up duration	Total number of patients (N)
Ling Mao [32]	Published	3 centers of Union Hospital of Huazhong University of Science and Technology, Wuhan, China	Retrospective, observational case series	January 16, 2020, to February 19, 2020	NA	214
Yanan Li [33]	Published	Single centre, Union Hospital of Huazhong University of Science and Technology, Wuhan, China	Retrospective, observational case series	16 January 2020, to 29 February 2020	NA	221
Lu Lu [34]	Published	Multicentre study from Hubei province, Sichuan province, Chongqing municipality, China	Retrospective study	January 18 to February 18, 2020	NA	304
F.A. Klok [35]	Published	Multicentre, Netherlands	Prospective study	March 7th 2020, to April 222,020	14 days	184
Corrado Lodigiani [36]	Published	Single centre, Humanitas Clinical and Research Hospital, Milan, Italy	Retrospective cohort study	13 February–10 April 2020	NA	388
Megan Fraisse [37]	Published	Single centre, France	Retrospective study	March 6 to April 22., 2020	NA	92 (1 lost to follow-up)
Siddhant Dogra [38]	Published	NYU Langone Health system, New York, USA	Retrospective cohort study	March 1st and April 27th, 2020	NA	3824
Julie Helms [39]	Published	Strasbourg, France	Observational Prospective case series	March 3 and April 3, 2020	NA	58
Julie Helms [40]	Published	Two centers of a French tertiary hospital France	Prospective cohort study	March 3rd and 31st 2020	April 7th	150
Sedat G Kandemirli [41]	Published	Multicentre (8 centers), Turkey	Retrospective study	March 1 to April 20,2020	NA	235
Silvia Garazzino [42]	Published	Italian Society of Paediatric Infectious Diseases, Multicentre, Italy	Retrospective study	25 March 2020, to 10 April 2020	At least 2 weeks	168
Rajan Jain [43]	Published	Multicentre(3 centers), New York	Retrospective cohort	March 1, 2020, and April 13, 2020	NA	3218
Alberto Benussi [44]	Published	ASST Spedali Civili Hospital, Lombardy, Italy	Retrospective, cohort study	February 21, 2020, to April v5, 2020	NA	56
Weixi Xiong [45]	Published	56 hospitals in Wuhan, Chongqing municipality, Sichuan province, China	Retrospective cohort study	18 January and 20 March 2020	NA	917 (1 asymptomatic patient excluded) (so total 918)
Tyler Scullen [46]	Published	Single center, New Orleans, Louisiana	Retrospective cross-sectional analysis	April 22, 2020	NA	27
Abdelkader Mahammed [47]	Published	Multicentre, Italy	Retrospective observational Study	Feb 29 to April 4	NA	725
Alireza Radmanesh [48]	Published	New York University Langone Medical Center, USA	Retrospective observational case series	March 1 and 31, 2020	2 weeks	3661
Carlos Manuel Romero-Sánchez [49]	Published	Two centers, Albacete, Spain	Retrospective observational	March 1st to April 1st, 2020	NA	841
Stephane Kremer [50]	Published	French Society of Neuroradiology, 16 hospitals, France	Retrospective cohort study	March 23th, 2020, to April 27th, 2020	NA	37
Pranusha Pirna [51]	Published	Rush University Medical Center, Chicago, Illinois, USA	Retrospective observational case series	March 1, 2020, to April 30, 2020	NA	50
	Published		Pilot multicentre case-control study	23rd to 25th March 2020	NA	79

**Table 1** (continued)

Álvaro Beltrán-Corbellini [52] Andrea Giacomelli [53]	Published	Multicentre (2 centres) Madrid, Spain L. Sacco Hospital in Milan, Italy	Cross-sectional study, verbal in-interview	19 March 2020	NA	59
Jerome R. Lechien [54] Giacomo Spinato [55]	Published	COVID-19 Task Force of YO-IFOS, Multicentre, Europe Treviso Regional Hospital, Italy	Prospective survey observational case series	NA	NA	417
Luigi Angelo Vaira [56]	Published	University Hospital of Sassari, Italy	Cross sectional telephone survey	March 19 and March 22, 2020	NA	202
Luigi Angelo Vaira [57] Luigi Angelo Vaira [58]	Published	Multicentre, Italy Multicentre, Italy	Prospective case series observational	March 31 and April 6, 2020	NA	72
Yonghyun Lee [59]	Published	The Daegu Medical Association, South Korea	Prospective study Multicentre prospective cohort study	April 9th and 10th 2020 NA	NA	33
Marlene M. Speth [60]	Published	Kantonsspital Aarau, Aarau, Switzerland	Prospective telephone interview	March 8, 2020 - March 31, 2020	NA	345
T. Klopfenstein [61]	Published	NFC (Nord Franche-Comté) Hospital, France	Prospective cross-sectional telephone questionnaire study	March 3, 2020, to April 17, 2020	NA	3191
Dawei Wang [10]	Published	Single centre,Zhongnan Hospital of Wuhan University in Wuhan, China	Retrospective, case series	March 1st to March 17th, 2020	March 24th, 2020	103
Wei-jie Guan [11] Nanshan Chen [12]	Published	Multicentre, 30 provinces in China	January 1 to January 28, 2020	January 1 to January 28, 2020	Till Feb. 3rd	138
Chaolin Huang [62]	Published	Jinyintan Hospital, Wuhan, China	Retrospective study	December 11, 2019, to January 31, 2020	NA	1099
Chaomin Wu [63]	Published	Jin Yintan Hospital, Wuhan, China	Retrospective study	Jan 1 to Jan 20, 2020	Till Jan 25,2020	99
Xiaobo Yang [64]	Published	Jinyintan Hospital Wuhan, China Jin Yin-tan Hospital, Wuhan, China	Prospective cohort Retrospective, observational study	Dec 16, 2019, to Jan 2, 2020 December 25, 2019, and January 26, 2020 Dec 24, 2019, to Jan 26, 2020	NA February 13, 2020 Feb 9, 2020	41
Tao Chen [65] Yingzhen Du [66]	Published	Tongji Hospital, Wuhan, China 2 centres, Hannan Hospital and Wuhan Union Hospital Wuhan, China	Retrospective case series Retrospective, observational study	13 January- 12 February 2020 January 9 to February 15, 2020	February 15, 2020 February 15, 2020	274 85
Yongli Zheng [67]	Published	Chengdu Public Health Clinical Medical Center, Chengdu, China	Retrospective case series	January 16 to February 20, 2020	February 23, 2020	99
Alfonso J. Rodriguez-Morales [68]	Published	Chile	Cross sectional	March 3, 2020, to March 23, 2020	NA	922
Feng Wang [69]	Published	Tongji Hospital Wuhan, China	Retrospective study	January 29, 2020, to February 10, 2020	February 22, 2020	28
Suxin Wan [70]	Published	Chongqing University Three Gorges Hospital, Chongqing, China	Retrospective case series	23 January - 8 February 2020	8 February 2020	135
Zhonghang Wang [71]	Published	Union hospital, Wuhan, China	Retrospective case series	January 16 to January 29, 2020	February 4, 2020	69
Dan Sun [72]	Published	Wuhan Children's Hospital, Wuhan, China	Case series	January 24 to February 24	February 24, 2020	8
Sijia Tian [73] Fei Zhou [74]	Published	Multicentre, 57 hospitals, Beijing, China 2 centers, Jinnyintan Hospital and Wuhan Pulmonary Hospital, Wuhan, China	Retrospective study Retrospective cohort	Jan 20 to Feb 10, 2020 Dec 29, 2019, to Jan 31, 2020	Feb 10, 2020 NA	262 191
Na Du [75]	Published	First Affiliated Hospital of Jilin University, Jilin, China	Case series	23 January 2020, to 11	NA	12

**Table 1** (continued)

Kui Liu [76] Alma Tostmann [77]	Published Published	9 tertiary hospitals, Hubei province, China Netherlands	Retrospective study Online anonymous questionnaire	December 30, 2019, to January 24, 2020 10 March to 29 March 2020	NA NA	137 90
Yongli Yan [78]	Published	Tongji Hospital, Wuhan, China	Retrospective, observational questionnaire	January 10, 2020, to February 24, 2020	NA	193
Xiao-Wei Xu [79]	Published	Multicentre, Zhejiang province, China	Retrospective case series	10 January 2020, to January 2020	NA	62
Jiang-shan Lian [80]	Published	Health Commission of Zhejiang Province Multicentre, Zhejiang province, China	Retrospective study	Jan 17 to Feb 7, 2020	Feb. 12, 2020	788
Nitesh Gupta [81]	Published	Safdarjung Hospital, India	Retrospective observational case series	Feb 1 st to 19th march 2020	19th March 2020	21
Xiaoli Zhang [82]	Published	Health Commission of Zhejiang Multicentre, Zhejiang, China	Retrospective study	January 17 to February 8	NA	645
Jie Li [83]	Published	Dazhou Central Hospital, Dazhou, China	Retrospective case series	22 January 2020, to 10 February 2020	11 February 2020	17
Ivan Fan-Ngai Hung [84]	Published	Multicentre, Hong Kong, China	Prospective, open-label, randomised, phase 2 trial	Feb 10 to March 20, 2020	NA	127
Huan Wu [85]	Published	Wuhan Children's Hospital, Wuhan, China	Retrospective case series	January 25 to April 18, 2020	April 18, 2020	148
Michael G Argenziano [86]	Published	NewYork-Presbyterian/Columbia University Irving Medical Center, New York, USA	Retrospective review	1 March to 5 April 2020	30 April	1000
Simone Bastrup Israelsen [87]	Published	Hvidovre Hospital, Denmark	Retrospective case series	10 March to 23 April 2020	NA	175
Matthew J Cummings [88]	Published	New York-Presbyterian hospitals affiliated with Columbia University Irving Medical Center, New York, USA	Prospective observational cohort	March 2 to April 1, 2020	April 28, 2020	257
Marjolein F. Q. Khuymans-van den Bergh [89]	Published	2 teaching Hospitals, Netherlands	Cross sectional	March 12, 2020, and March 16, 2020 (interview dates)	March 16, 2020	86
Blazej Nowak [90]	Published	Central Clinical Hospital, Warsaw, Poland	Retrospective study	March 16, 2020, to April 7, 2020	April 7, 2020	169
Xiaoquan Lai [91] X. Wang [92]	Published Published	Tongji Hospital Wuhan Dongxihu Fangcang Hospital, Wuhan, China	Retrospective case series Retrospective study	January 1 to February 9, 2020 7 February to 12 February 2020	NA 22 February	110 1012
Zhe Liu [93]	Published	Multicentre Xi'an, Shaanxi province, China	Retrospective study	January 16 to February 13, 2020	NA	72
Qiong Huang [94] Kyung Soo Hong [95]	Published Published	Multicentre, Human, China Yeungnam University Medical Center in Daegu, South Korea	Retrospective case series Retrospective study	January 17 to February 10, 2020 Up to March 29, 2020	NA March 29, 2020	54 98
Rui Huang [96] Mengyao Ji [97]	Published	Renmin Hospital of Wuhan University Wuhan, China	Retrospective study Retrospective study	January 22, 2020, to February 10, 2020 2nd January to 28 January 2020	February 10, 2020 8 February 2020	202 101
Dawei Wang [98]	Published	Zhongnan Hospital of Wuhan University in Wuhan and Xishui Hospital, Hubei Province, China	Retrospective study	Up to February 10, 2020	NA	107
Saurabh Aggarwal [99] Xin-Ying Zhao [100]	Published Published	Unity Point Clinic, USA Jingzhou Central Hospital Jingzhou, China	Retrospective study Retrospective study	March 1 to April 4, 2020 January 16, 2020, to February 10, 2020	NA February 10, 2020	16 91

**Table 1** (continued)

Yifan Meng [101] Qingchun Yao [102]	Published Published	Tongji Hospital, Wuhan, China Dabieshan Medical Center, Huanggang city, Hubei Province, China	Retrospective study Retrospective cohort	January 16th to February 4th, 2020 January 30, 2020 -February 11, 2020	March 24th, 2020 March 3	168 108 (1 pregnant patient excluded as information incomplete)
Li Zhu [103] Eu Suk Kim [104]	Published Published	Multicentre, Jiangsu province, China. Korea National Committee for Clinical Management of COVID-19, South Korea	Retrospective case series Nationwide multicentre retrospective study	January 24, 2020, to February 22, 2020 January 19th, 2020, to February 17th, 2020	February 25, 2020 February 17th, 2020	10 28
Pavan K. Bhattraj [105] Haiyan Qiu [106] Guang Chen [107]	Published Published Published	Multicentre(9), Seattle, USA Multicentre (3), Zhejiang, China Tongji Hospital, Wuhan, China	Retrospective study Retrospective cohort Retrospective study	February 24 to March 9, 2020 Jan 17 to March 1, 2020 Late December 2019 to January 27, 2020	March 23, 2020 Feb 28, 2020 February 2, 2020	24 36 21 (available data of symptoms in 20 only)
Wenjie Yang [108] Yu-Huan Xu [109]	Published Published	Multicentre(3 centers), Wenzhou city, Zhejiang, China Single centre, Beijing, China	Retrospective cohort Retrospective study	January 17th to February 10th, 2020 January to February 2020	Feb 15th, 2020 NA	149 50
Xi Xu [110] Jerome R. Lechien [111]	Published Published	Guangzhou Eighth People's Hospital, Guangzhou, China Multicentre, Europe	Retrospective study Observational, cross-sectional study	January 23, 2020, and February 4, 2020 March 22 to April 10, 2020	NA NA	90 1420
Sherry L. Burner [112]	Published	CDC COVID-19 Response Team, United states, USA	Retrospective study	February 12 to April 9, 2020	NA	9282 (symptom data for 4707) (age data for 8945) (sex data for 9667)
Ruth Levinson [113]	Published	Tel Aviv Medical Center, Israel	Retrospective with questionnaire via mobile and email	March 10 to 23, 2020	25th of March	42 (total 45 admitted, only 42 completed questionnaire)
Xu Zhu [114] Dan Wang [115] Chunning Chen [116]	Preprint Preprint Preprint	Renmin Hospital of Wuhan University, Wuhan, China Zhongshan Hospital, Wuhan, China Shenzhen Third People's Hospital, Guangdong, China	Retrospective study Cross-sectional study Prospective study	January 20 to February 15, 2020 January 15, 2020-February 28, 2020 Jan 16, 2020, to Feb 19, 2020	February 20, 2020 NA NA	114 143 31
Pingzheng Mo [117] Gu-qin Zhang [118]	Published Published	Zhongnan Hospital of Wuhan University, Wuhan, China Zhongnan Hospital of Wuhan University, Wuhan, China	Retrospective study Retrospective case series	January 1st to February 5th January 2, 2020, to February 10, 2020	NA NA	155 221
Jennifer Tomlins [119] Zonghao Zhao [120]	Published Preprint	North Bristol NHS Trust, UK First Affiliated Hospital of USTC Hefei, China	Retrospective study Retrospective study	March 10th to March 30th, 2020 Jan 21 to Feb 16, 2020	Feb 15, 2020 NA	95 75
Ying Huang [121]	Preprint	Fifth Hospital of Wuhan, Wuhan, China	Retrospective study	Jan 21 - Feb 10, 2020	Feb 14, 2020	36
Carol H. Yan [122]	Published	University of California San Diego Health, La Jolla, California, USA	Cross-sectional internet- and email-based platform	March 3, 2020, and March 29, 2020	NA	59

**Table 1** (continued)

First author	Study population	Age (years), mean $\pm$ SD or median(range) or median (IQR)	Sex (male) $n$ (%) Neurological features $n$ (%)	Remarks (groups compared)	Outcome $n$ (%)
Ling Mao [32]	Consecutive hospitalized patients	52.7 $\pm$ 15.5	87 (40.7)	Any—78 (36.4) CNS—53 (24.8) Dizziness—36 (16.8) Headache—28 (13.1) Impaired consciousness—16 (7.5) Acute cerebrovascular disease—6 (2.8)	Severe vs non-severe 5 ischaemic stroke, 1 hemorrhagic stroke NA
Yan Deng [123] Jiaojiao Chu [124] Håkon Ihle-Hansen [125]	Published Published Published	2 centers, Wuhan, China Tongji Hospital, Wuhan, China Berum Hospital, Norway	Retrospective study Retrospective study Observational qualitative study	January 1, 2020, to February 21, 2020 7 January to 11 February 2020 9–31 March 2020	NA NA 31 March 2020
Parag Goyal [126] Jianlei Cao [127]	Published Published	2 centres, New York, USA Wuhan University Zhongnan Hospital, Wuhan, China	Retrospective case series Retrospective cohort	March 3 to March 27, 2020 3 January to 1 February 2020	April 10th 15 February 2020
De Chang [128]	Published	Multicentre (3 centers), Beijing, China	Case series	January 16, 2020, to January 29, 2020	February 4, 2020
Huijun Chen [129]	Published	Zhongnan Hospital of Wuhan University, Wuhan, China	Retrospective case series	Jan 20 to Jan 31, 2020	Feb 4, 2020
Lang Wang [130]	Published	Renmin Hospital of Wuhan University, China	Retrospective study	Jan 1 to Feb 6, 2020	March 5
Gianfranco Spiteri [131]	Published	WHO European Region(except UK), Europe	Cross-sectional study	24 January to 21 February 2020	21 February 2020
Yingxia Liu [132]	Published	Shenzhen Third People's Hospital, China	Case series	Jan 11 to Jan 20, 2020	NA
Tianmin Xu [133]	Published	Third Hospital of Changzhou, Changzhou city, Jiangsu province, China	Retrospective cohort	Jan 23 to February 18, 2020	February 27, 2020
Michael Chung [134]	Published	Multicentre (3 centers), 3 provinces, China	Retrospective case series	January 18, 2020, to January 27, 2020	NA
Heshui Shi [135]	Published	Wuhan Jiniyintan hospital or Union Hospital of Tongji Medical College, China	Retrospective study	Dec 20, 2019, to Jan 23, 2020	Feb 8th, 2020
Luhuan Yang [136]	Published	Yichang Central People's Hospital, Yichang, Hubei Province, China	Retrospective study	Jan 30 to Feb 8, 2020	Feb 26, 2020
Wei Zhao [137] Ya-nan Han [138] Yang Wang [139]	Published Published Published	Multicentre (4 centers), Hunan, China Xian eighth hospital Shaanxi, China Tongji Hospital, China	Retrospective study Retrospective study Cohort	NA 31st January–16th February 2020 January 25, 2020, to February 25, 2020	NA NA 28 days follow-up

**Table 1** (continued)

Yanan Li [33]	Consecutive hospitalized patients	$53.3 \pm 15.9$	131 (59.3)	Ataxia—1 (0.5) Seizure—1 (0.5) PNS—19 (8.9) Taste disturbances—12 (5.6) Smell disturbances—11 (5.1) Vision impairment—3 (1.4) Nerve pain—5 (2.3) Skeletal muscle injury—23 (10.7) Acute cerebrovascular disease—13 (5.9)	Severe vs non-severe, with cerebrovascular disease vs without cerebrovascular disease	4 ischaemic stroke and 1 hemorrhagic stroke patients expired
Lu Lu [34]	Consecutive discharged or died patients from multiple centers	44 (33–59.25)	182 (59.9)	Ischaemic stroke—11 (84.6) Cerebral venous sinus thrombosis—1 (7.7) Cerebral haemorrhage—1 (7.7) Acute cerebrovascular disease—3 (1)	Mild, moderate vs severe, critical	NA
F.A. Klok [35]	Only ICU patients	NA	NA	Acute cerebrovascular disease—5 (2.8) (all ischaemic stroke) Acute ischaemic stroke—9 (2.5)	All patients received thrombo prophylaxis	41(22%) died and 78 (43%) discharged alive
Corrado Lodigiani [36]	Consecutive adult symptomatic patients admitted, 61 ICU patients	66 (55–75)	264 (68)	ICU vs general ward, survivors vs non-survivors, thromboprophylaxis in 100% ICU and 75% ward patients	ICU vs general ward, survivors vs non-survivors, thromboprophylaxis in 100% ICU and 75% ward patients	2 stroke patients died, 4 patients discharged
Megan Fraissé [37]	Only ICU patients	61 (55–70)	73 (79)	Acute cerebrovascular disease—4 (4.3) Ischaemic—2 (2.2) Hemorrhagic—2 (2.2) Acute hemorrhagic stroke—3 (3)	All received thromboprophylaxis	NA
Siddhant Dogra [38]	All hospitalized patients	62 (37–83) (among 33 patients)	26/33 (78.8)	(0.9) (only in 755 neuroimaging done)	37 had hemorrhage, but 4 excluded as hemorrhage secondary to trauma, bleeding in brain metastases, after tumor resection	NA
Julie Helms [39]	Consecutive hospitalized ICU patients	63	NA	Agitation—40/58 (69) Corticospinal tract signs—39/58 (67) Dysexecutive syndrome—14/39 (36) MRI—leptomeningeal enhancement—8/13 (62)	Perfusion abnormalities—11/11 (100) Cerebral ischaemic stroke—3/13 (23) Cerebral ischaemic attack—2 (1.3) (population after matching—0)	Historical prospective cohort of “non-COVID-19 ARDS” patients vs COVID-19 ARDS
Julie Helms [40]	All consecutive patients referred to ICU for ARDS	63 (53–71)	122 (81.3)	Neurological symptoms—50 (21) Cortical signal abnormalities on FLAIR images—10/27 (37) Acute transverse sinus thrombosis—1 (0.4)	Cerebral ischaemic stroke—3/13 (23) Cerebral ischaemic attack—2 (1.3) Historical prospective cohort of “non-COVID-19 ARDS” patients vs COVID-19 ARDS	Discharged—36 ICU admission—101 Died—13
Sedat G Kandemirli [41]	Patients admitted to ICU	63 (34–87)	21 (78)	Brain MRI done in 27/50 (54%) patients with neurological symptoms	Brain MRI done in 27/50 (54%) patients with neurological symptoms	NA

**Table 1** (continued)

Silvia Garazzino [42]	Pediatric patients under 18 years	2.3 (0.3–9.6)	94 (55.9)	Acute infarction in right middle cerebral artery territory—1 (0.4) Non-febrile seizures—3 (1.8) Febrile seizures—2 (1.2)	NA	Recovered—168
Rajan Jain [43]	All patients admitted	NA	NA	Imaging (32/18) Acute cerebrovascular disease—35 (1.1) Ischaemic—26 Hemorrhagic—9 Hypoxic anoxic brain injury—2	Neuro imaging done—454 (14.1%) Imaging Positive—38 (8.4) Stroke—35 (92.5) Ischaemic stroke—26 (68.5) Large vessel—17 (44.5) Lacunar—9 (24) Hemorrhagic stroke—9 (24) Hypoxic anoxic brain injury—2 (5) Encephalitis—1 (2.5)	NA
Alberto Benussi [44]	All adult ( $\geq 18$ years old) patients admitted for neurological disease and had a definite outcome	77.0 (67.0–83.8)	28 (50.0)	Ataxia (1.4%) Cerebrovascular disease—43 (76.8) TIA—5 (11.6) Ischaemic stroke—35 (81.4) Hemorrhagic stroke—3 (7.0) Epilepsy—4 (7.1) Delirium—15 (26.8)	COVID-19 vs non-COVID-19 Critical vs non-critical neurological events	Mortality—21 (37.5) Discharged—742 Hospitalized—145 Died—30
Weixi Xiong [45]	All consecutive symptomatic patients	48.7 ± 17.1	504 (55)	New-onset neurological events—39 (4.3) Disturbance of consciousness/delirium—21 (2.3) Syncope—3 (0.3) Traumatic brain injury—1 Acute Cerebrovascular accident—10 (early onset—2) Occipital neuralgia—1 Unexplained severe headache—2 Non-specific headache—8 Functional or? Tic/tremor—2	NA	
Tyler Scullen [46]	Severe cases with neurological features	59.8 (35–91)	14 (52)	Muscle cramp—2 Altered mental status—26 (96.3) Dysgeusia—1 (3.7) Generalized weakness—1 (3.7) Headache—2 (7.4) Focal Deficit—10 (37.0) Decerebrate posturing—1 (3.7) Facial droop—1 (3.7) Fixed pupils—1 (3.7) Gaze deviation—3 (11.1) Hemineglect—2 (7.4) Hemiparesis or hemiplegia—4 (14.9)	Imaging and EEG Encephalopathy—20 (74) Acute necrotizing encephalopathy—2 (7) Vasculopathy—5 (19) Subacute ischaemic stroke—4 (14.8) NCSE—1 (3.7) Large vessel occlusion—PCA P2B—1 (3.7)	NA

**Table 1** (continued)

			Quadruplegia 1 (3.7)	Focal stenosis ICA terminus—3 (11.1)
Abdelkader Mahammed [47]	Consecutive hospitalized patients	NA	Acute neurological symptoms—108 (15) Altered mental status—64(8.8) Ischaemic stroke—33(total was 34, but 1 is hypoxic encephalopathy added here) Headache—13 (1.8) Myalgia—13 (1.8) Seizures—10 Dizziness—4(0.6) Neuralgia—3 Ataxia—2 (0.3) Hypoxmia—2 (0.3) ICH-6 Hypoxic ischaemic encephalopathy—1 Cerebral venous thrombosis—2 GBS—2	119 patients had neurological symptoms, however, only 108 received neuroimaging evaluation NA
Alireza Radmanesh [48]	All patients diagnosed	NA	MFS—1 PRES—1 Acute encephalopathy—1 Non-specific encephalopathy—2 MS plaque exacerbation—2 Acute/subacute infarct—13 Haemorrhage—7 (excluding previous) Altered mental status—102 (2.9%) Syncope/fall (79 patients	242 underwent imaging (CT or MRI) NA
Carlos Manuel Romero-Sánchez [49]	All patients admitted	66.42 ± 14.96	473 (56.2) Focal neurologic deficit—30 Neurological manifestations—483 (57.4) Myalgias—145 (17.2) Headache—119 (14.1) Dizziness—51 (6.1) Syncope—5 (0.6) Anosmia—41 (4.9) Dysgeusia—52 (6.2) Disorders of consciousness—165 (19.6) Seizures—6 (0.7) Dysautonomia—21 (2.5) ALDP—1	Non-severe vs severe Mortality—197 (23.42) NA
			HyperCKemia—73 (9.2) Rhabdomyolysis—9 (1.1) Myopathy—26 (3.1) Ischaemic stroke—11 (1.3) Intracranial hemorrhage—3 (0.4) Movement disorders—6 (0.7) Encephalitis—1 (0.1) Optic neuritis—1 (0.1)	

**Table 1** (continued)

Stephane Kremmer [50]	Severe patients with abnormal MRI Only	61 (8–78)	30 (81)	Neuropsychiatric symptoms—167 (19.9) Headache—4 (11) Seizures—5 (14) Clinical signs of corticospinal tract involvement—4(11) Disturbances of consciousness—27 (73) Confusion—12 (32) Agitation—7(19)	Non-hemorrhagic vs hemorrhagic forms CSF—1 patient's CSF SARS-CoV-2 RT-PCR positive	Died—5 (14)
Pranusha Pirna [51]	Only 50 patients admitted to neurology ward or referred to neurology is studied	NA	NA	Pathological wakefulness in intensive care units—15(41) CNS Altered mental status—30 Seizures—13 Headache—12 Short-term memory loss—12 Acute cerebrovascular accident—19 Acute ischaemic stroke—10 Hypoxic ischaemic brain injury—7 ICH—4 Non-aneurysmal SAH—4 PRES—2 TIA—1 PNS Dyautonomia—6 Muscle injury with elevated CK—6 Hypogeausia/dygeusia—5 Hyposmia—3 Extraocular muscle abnormalities—5 Isolated unilateral facial palsy—3 Paresthesias—1 Ataxia—1 Smell and/or taste disorder—31 (39.2) Smell disorder—25 (31.65) (Most common—anosmia—14/31 (45.7))	Neurological manifestations—7.7% (total patients in the hospital were 650; however, not all evaluated for neurological symptoms, mentioned in the limitations of the study)	NA
Álvaro Beltrán-Corbellini [52]	Consecutive patients hospitalized, > 18 years	61.6 ± 17.4	48 (60.8)	Case—COVID-19 patients Control—40 historical group of 2019/2020 season influenza patients	Case—COVID-19 patients Control—40 historical group of 2019/2020 season influenza patients	NA
Andrea Giacomelli [53]	All hospitalized patients who were able to be interviewed	60 (50–74)	40 (67.8)	Taste disorder—28 (35.44) Most common—ageusia 14/31 (45.2) Headache—2 (3.4) Olfactory and/or taste disorders—20 (33.9) Olfactory disorders—14 Taste disorder—17 Olfactory dysfunction—357 (85.6) Anosmia—284 (79.6) Hyposmia—73 (20.4) Phantosmia—12.6%	NA	NA
Jerome R. Lechien [54]	Adult > 18 years, mild to moderate cases (ICU cases excluded) hospitalized and home patients	36.9 ± 11.4	154 (36.9)			

Table 1 (continued)

Giacomo Spinato [55]	Adults ( $\geq 18$ years) consecutively assessed and mildly symptomatic (only home managed patients)	56 (45–67)	97 (48.0)	Headache—86 (42.6) Muscle or joint pains—90 (44.6) Dizziness—28 (13.9) Altered sense of smell or taste 130—(64.4%) Headache—30 (41.6) Olfactory and taste disorders—53 (73.6) Olfactory disorder—44 (61.1) Taste disorder—39 (54.2) Olfactory and taste disorders—21 (63.6) Olfactory disorder—17 (51.5) Taste disorder—17 (51.5) Olfactory and/or taste disorders—256(74.2) Olfactory disorder—225 Taste disorder—234 Anosmia and/or ageusia—488 (15.3) Anosmia—389	NA	NA
Luigi Angelo Vaira [56]	Adults over 18 years of age (excluded assisted ventilation patients)	49.2 ± 13.7	27 (37.5)	Olfactory and taste disorders—53 (73.6) Olfactory disorder—39 (54.2) Olfactory and taste disorders—21 (63.6) Olfactory disorder—17 (51.5) Taste disorder—17 (51.5) Olfactory and/or taste disorders—256(74.2) Olfactory disorder—225 Taste disorder—234 Anosmia and/or ageusia—488 (15.3) Anosmia—389	NA	NA
Luigi Angelo Vaira [57]	Health care staff, home quarantined, age $> 18$ years	47.2 ± 10	11 (33.3)	Olfactory and taste disorders—21 (63.6) Olfactory disorder—17 (51.5) Taste disorder—17 (51.5) Olfactory and/or taste disorders—256(74.2) Olfactory disorder—225 Taste disorder—234 Anosmia and/or ageusia—488 (15.3) Anosmia—389	NA	Validation of a self-administered olfactory and gustatory test done
Luigi Angelo Vaira [58]	Both hospitalized and home quarantined patients, $\geq 18$ years (excluded assisted ventilation patients)	48.5 ± 12.8	146 (42.3)	Olfactory and taste disorders—21 (63.6) Olfactory disorder—17 (51.5) Taste disorder—17 (51.5) Olfactory and/or taste disorders—256(74.2) Olfactory disorder—225 Taste disorder—234 Anosmia and/or ageusia—488 (15.3) Anosmia—389	NA	Objective assessment done
Konghyun Lee [59]	COVID-19 patients awaiting hospitalization or facility isolation	44.0(25.0–58.0)	1161(36.4)	Olfactory and taste disorders—21 (63.6) Olfactory disorder—17 (51.5) Taste disorder—17 (51.5) Olfactory and/or taste disorders—256(74.2) Olfactory disorder—225 Taste disorder—234 Anosmia and/or ageusia—488 (15.3) Anosmia—389	NA	Presence vs absence of anosmia or ageusia
Marlene M. Speth [60]	All positive (ICU and deceased excluded)	NA	50 (48.5)	Olfactory dysfunction—63 (61.2) Decreased smell—14.6% Anosmia—46.6%	NA	NA
F. Klopferstein [61]	All admitted adults	NA	NA	Gustatory dysfunction—67 (65.0) Decreased taste—25.2% Ageusia—39.8% Anosmia—54 (47) Dysgeusia—46/54 (85) Myalgia—40/54 (74) Headache—44/54 (82) Dizziness—13 (9.4) Headache—9 (6.5) Headache—150 (13.6) Myalgia or arthralgia—164 (14.9) Rhabdomyolysis—2 (0.2)	NA	Death—2/54(4)
Dawei Wang [10]	Consecutive patients admitted	56 (42–68)	75 (54.3)	ICU vs non-ICU	NA	NA
Wei-je Guan [11]	All patients with data available	47.0 (35.0–58.0)	637/1096 (58.1)	All Severe vs non-severe	NA	Death—15 (1.4) Discharged—55 (5.0) Hospitalization—1029 (93.6) Recovery—9 (0.8) Remained in hospital—57 (58) Discharged—31 (31) Died—11 (11)
Nanshan Chen [12]	All hospitalized patients	55.5 ± 13.1	67 (68)	Muscle ache—11 (11) Headache—8 (8) Confusion—9 (9)	NA	Hospitalization—7 (17) Discharge—28 (68)
Chaolin Huang [62]	Hospitalized	49.0 (41.0–58.0)	30 (73)	ICU vs non-ICU	NA	Myalgia or fatigue—18 (44) Headache—3/38 (8)

**Table 1** (continued)

Chaomin Wu [63]	All hospitalized patients	51 (43–60)	128 (63.7)	Fatigue or myalgia—65 (32.3)	ARDS vs non-ARDS
Xiaobo Yang [64]	Only critically ill patient admitted in ICU	59·7 (13·3)	35 (67)	Myalgia—6 (11·5) Headache—3 (6)	Survivors vs non-survivors
Tao Chen [65]	113 died and 161 fully recovered and discharged patients	62·0 (44·0–70·0)	171 (62)	Myalgia—60 (22) Headache—31 (11) Dizziness—21 (8)	Deaths vs recovered
				Hypoxic encephalopathy—24 (9)	
Yingzhen Du [66]	Consecutive severe patients	65·8 ± 14·2	62 (72·9)	Myalgia—14 (16·5) Headache—4 (4·7)	Died—85
Yongli Zheng [67]	Consecutively hospitalized All ages First notified cases of COVID-19	49·40 ± 18·45 NA	51(52) NA	Muscle ache and headache—12 (12) Headache—59·7 (64·8) Myalgia—32 (3·5)	NA
Alfonso J. Rodriguez-Morales [68]				Headache—3 (10·7)	
Feng Wang [69]	Diabetic, hospitalized patients	68·6 ± 9·0	21(75)	Headache—3 (10·7)	ICU vs non-ICU
Suxin Wan [70]	Hospitalized patients	47 (36–55)	72 (53·3)	Myalgia or fatigue—44 (32·5) Headache—24 (17·7)	Mild vs severe
Zhongliang Wang [71]	Hospitalized patients	42·0(35·0–62·0)	32(46)	SpO2 < 90 vs SpO2 > 90	
Dan Sun [72]	Pediatric ICU (severe and critically ill only)	2 months to 15 years	6	Myalgia—21 (30) Headache—10 (14) Dizziness—5 (7)	Hospitalization—44(65·7)
Sijia Tian [73]	Hospitalized, all age groups	47·5 (1–94)	127 (48·5)	Myalgia or fatigue—1 Headache—1—1 Headache—17 (6·5)	Discharge—18 (26·9)
Fei Zhou [74]	All adult ≥ 18 hospitalized and either dead or discharged patients	56·0 (46·0–67·0)	119 (62)	Myalgia—29 (15)	Death—5 (7·5)
Na Du [75]	Consecutive hospitalized patients	45·25(23–79)	7(54·3)	Headache—3 (20)	Hospitalized—3 (42·9)
Kui Liu [76]	Hospitalized patients	57 (20–83)	61 (44·5)	Myalgia or fatigue—44(32·1) Headache—13(9·5)	Discharged—44 (32·1)
Alma Tostmann [77]	Only health care workers	NA	19 (21·1)	Anosmia—37/79 (46·8) Muscle ache—57/90 (63·3) Headache—64/90 (71·1) Headache—21 (10·9)	Hospitalized—77 (56·2)
Yongli Yan [78]	Adults over 18 years, hospitalized, severe (all hospitalized admitted there included)	64 (49–73)	114 (59·1)	negative 48 diabetic vs 148 non-diabetic, survivors	Death—16 (11·7)
Xiao-Wei Xu [79]	Adult hospitalized patients	41 (32–52)	35 (56)	Myalgia or fatigue—32 (52) Headache—21 (34)	Mortality—108 (56·0)
				Symptom onset > 10 days vs < 10 days	Hospital admission—61 (98) Discharge—1 (2) Death—0

**Table 1** (continued)

Jiang-shan Lian [80]	All confirmed cases	NA	407(51.65)	Muscle ache—91(11.54) Headache—75(9.52)	With Wuhan exposure vs without	Discharged—322 (40.86)
Nitesh Gupta [81]	First 21 hospitalized patients in the centre	40.3 (16–73)	14 (66.7)	Headache—3 (13.6)	Death—0	Death—0
Xiaoli Zhang [82]	All hospitalized patients	NA	328(50.85)	Muscle ache—71(11.01) Headache—67(10.39)	Normal imaging vs abnormal imaging	Discharged—5
Jie Li [83]	All hospitalized patients	45.1 ± 12.8 45 (22–65)	9 (52.9)	Myalgia—4 (23.5) Dizziness—2 (11.8)	Discharged vs non-discharged	Hospitalized—12
Ivan Fan-Ngai Hung [84]	Adult at least 18 years, admitted	NA	68 (53.4)	Myalgia—18 (14.17) Headache—6 (4.72)	Combination triple antiviral drug vs control	Death—0
Huan Wu [85]	Pediatric mild and moderate cases only	84 (18–123)months	60 (40.5)	Anosmia—5 (3.93) Headache—5 (3.4)	group (lopinavir–ritonavir)	Discharged—148 (100)
Michael G Argenziano [86]	First 1000 consecutive patients presented to centre	63.0 (50.0–75.0)	596 (59.6)	Myalgia—268 (26.8) Headache—101 (10.1)	NA	Died—0
Simone Bastrop Israelson [87]	Consecutive patients, adult ≥ 18, hospitalized	71 (55–81)	85 (48.6)	Syncope—48 (4.8) Myalgia—46 (26.3) Headache—32 (18.3)	Emergency vs ward vs ICU	Discharged—699
Matthew J Cummings [88]	Only critically ill adults aged ≥ 18 years	62 (51–72)	171 (67)	Altered sense of taste—5 (2.9) Headache—10 (4)	General Ward vs ICU	Died—211
Marjolein F. Q. Kluytmans-van den Bergh [89]	Only health care workers infected	49 (22–66)	15 (17)	Severe myalgia—54 (63) Headache—49 (57) Altered or lost sense of taste—6 (7)	Interview within 7 d of the onset of Symptoms vs >7d	Hospitalized—98 (38)
Błażej Nowak [90]	Consecutive patients hospitalized	63.7 ± 19.6	87 (51.5)	Headache—1 Anosmia and ageusia—3 (1.7)	Recovered—19 (22) Hospital admission—2 (2)	Recovered—19 (22) Hospital admission—2 (2)
Xiaoquan Lai [91]	Only health care workers	36.5 (30.0–47.0)	31 (28.2)	Myalgia or fatigue—66 (60.0) Muscle ache—50 (45.5) Headache—33 (30.0)	Hcw with COVID-19 vs without	Hospitalized—80 (45.7)
X. Wang [92]	Only non-critically ill (however, all patient admitted in that hospital included)	50 (39–58)	524 (51.8)	Dizziness—24 (21.8) Headache—152 (15.0) Myalgia—170 (16.8)	With and without aggravation during follow up	Discharged—58 (23)
Zhe Liu [93]	All hospitalized	46.2 ± 15.9	39 (54.2)	Muscle soreness—7 (9.7) Headache—4 (5.6)	Uncomplicated vs mild vs severe	Died—46 (26.3)
Qiong Huang [94]	All hospitalized patients	41 (31–51)	28 (51.9)	Muscle soreness—9 (16.7) Headache—3 (5.6)	Common vs severe	Died—1 (0.9)
Kyung Soo Hong [95]	Consecutive hospitalized patients	55.4 ± 17.1	38 (38.8)	Dizziness—3 (5.6) Myalgia—37 (37.8)	ICU vs non-ICU	Remains in hospital—57 (58.2)
						Discharged—30 (30.6)
						Died—5 (5.1)

**Table 1** (continued)


Rui Huang [96]	All hospitalised	44.0 (33.0–54.0)	116 (57.4)	Muscle ache—21 (10.4) Headache—12 (5.9)	Severe vs non-severe	Transferred—6 (6.1) Remained in hospital—165 (81.7) Hospital discharge—37 (18.3)
Mengyao Ji [97]	Random selection of confirmed patients	51.0 (37.0–61.0)	48 (48)	Myalgia—16 (16) Vertigo—4 (4) Headache—6 (6)	Medica staff vs non-medical	Death—0 (0) Death—11 (11) Hospitalization—53 (52)
Dawei Wang [98]	All the discharged (alive at home and dead) patients with confirmed COVID-19	51.0 (36.0–65.0)	57 (53.3)	Myalgia—33 (30.8) Headache—7 (6.5) Dizziness—7 (6.5)	Survivors vs non-survivors	Cured—37 (37) Died—19 Survived—88
Saurabh Aggarwal [99]	88 patients overlap with Wang D [100] All admitted patients	67 (38–95)	12 (75)	Lightheadedness—3 (19) Headache—4 (25) Anosmia—3 (19) Dysgeusia—3 (19) Myalgia—15 (16.5) Dizziness—3 (3.3)	ICU, shock, death vs no	Died—3 (19) Discharged—11 Admitted—2
Xin-Ying Zhao [100]	All hospitalized patients	46.00	49 (53.8)	Severe vs mild	Remained in hospital—75 (82.4) Discharged—14 (15.4)	
Yifan Meng [101]	All consecutive admitted all were severe or critically ill patients)	56.7 ± 15.1	86	Disturbance of consciousness—3 (3.3) Male vs female	Died—2 (2.2) Died—17 (8.9) Discharge—136 Hospital—15	
Qingchun Yao [102]	Consecutive adult patients admitted 1–18 years, children	52 (37–58)	43 (39.8)	Myalgia or fatigue—28 (25.9) Headache—1 (0.9)	Died—12 Discharged—96	
Li Zhu [103]	First 28 patients in Republic of Korea, hospitalized	NA	5 (50.0)	Headache—2 (20.0)	Discharged—5 (50.0) Hospitalized—5 (50.0)	
Eu Sul Kim [104]	Only critically ill ICU patients	42.6 ± 13.4	15 (53.6)	Myalgia—7 (25.0) Headache—7 (25.0) Headache—2 (8)	Discharged—10 Hospitalized—18 Died—12 (50) Discharged—5 (21) Hospitalized—7 (30)	
Pavan K. Bhattraij [105]	All pediatric 0–16 years All hospitalized patients	64 ± 18 (23–97)	15 (63)	Myalgia or fatigue—28 (25.9) Headache—1 (0.9)	All cured Died—4 Recovered—2	
Haiyan Qiu [106]	Consecutive hospitalized patients	8.3 ± 3.5 56.0 (50.0–65.0)	23 (64) 17 (81.0)	Headache—3 (8) Myalgia—8/20 (40.0%) Headache—2/20 (10.0%) Muscle pain—5 (3.36%) Headache—13/8.72%	Mild vs moderate Severe vs moderate	
Guang Chen [107]					NA	
Wenjie Yang [108]					Remained in hospital—76 (51.01) Discharged—73 (48.99) Died—0 (0.0)	
Yu-Huan Xu [109]	All hospitalized patients	43.9 ± 16.8	29 (58)	Headache—5 (10) Muscle ache—8 (16) Myalgia—25 (28) Headache—4 (4)	Mild vs moderate vs severe vs critically severe	
Xi Xu [110]	All hospitalized patients	50 (18–86)	39 (43)	Headache—99/8 (70.3) Loss of smell—99/7 (70.2) Reduction of smell—201 (14.2) Myalgia—887 (62.5)	NA	
Jerome R. Lechien [111]	Mild to moderate (but all reported)	39.17 ± 12.09	458 (32.3)	Based on age	NA	

**Table 1** (continued)

Sherry L. Burner [112]	Cases reported to CDC, only health care personal	42 (32–54)	246(27)	Taste dysfunction—770 (54.2) Muscle ache—3122(66) Headache—3048(65) Loss of smell or taste—750(16)	NA	Data of 8945 Not hospitalized—6760 (90%) Hospitalized—723 (8–10%) ICU admission—184 (2–5%) Died—27 (0.3–0.6%) NA
Ruth Levinson [113]	Hospitalized adults and adolescents (age $\geq$ 15 years), and mild symptoms (all admitted were mild)	34 (15–82)	23	Myalgia or arthralgia—24 (57) Headache—20 (48) Anosmia—14 (33) Dysgeusia—15 (36) Dizziness—9 (21) Myalgia—4 (3.5)	NA	
Xu Zhu [114]	Only elderly(> 70) patients	76 (72–82)	67 (58.8)	Myalgia—49(34.3) Headache—7(4.9) Headache—1 (3.2)	Severe vs non-severe Mild/moderate vs severe/critical NA	Alive—87 (76.3) Dead—27(23.7) NA
Dan Wang [115]	All consecutive admitted patients	58(39–67)	73(51.0)	Myalgia or arthralgia—50 (61.0) Headache—8 (9.8) Dizziness—2 (2.4) Headache—17(7.7)	General vs refractory NA	Died—0 NA
Chuning Chen [116]	Only pediatric, < 18 years, hospitalized patients	7.33 $\pm$ 4.35	13 (41.9)	Myalgia—13 (14) Confusion—20 (21) Seizure—1 (1.1) Headache—9 (9.5)	Severe vs non-severe NA	Hospitalization—168 (76.0) Discharge—42 (19.0) Death—12 (5.4) Died—21 (21) Discharged—44(43) Hospitalized—30 (29)
Pingzheng Mo [117]	All Consecutive admitted patients	54 (42–66)	86 (55.5)	Myalgia—13 (14) Confusion—20 (21) Seizure—1 (1.1) Headache—9 (9.5)	NA	
Gu-qin Zhang [118]	All hospitalized patients	55.0 (39.0–66.5)	108(48.9)	Myalgia—13 (14) Confusion—20 (21) Seizure—1 (1.1) Headache—9 (9.5)	NA	
Jennifer Tomlins [119]	All sequential hospitalized patients	75 (59–82)	60 (63)	Myalgia—13 (14) Confusion—20 (21) Seizure—1 (1.1) Headache—9 (9.5)	NA	
Zonghao Zhao [120]	All positive cases	47 (34–55)	42 (56)	Myalgia—13 (14) Confusion—20 (21) Seizure—1 (1.1) Headache—9 (9.5)	NA	
Ying Huang [121]	Non survivors only	69.22 (9.64)	25 (69.4)	Myalgia—13 (14) Confusion—20 (21) Seizure—1 (1.1) Headache—9 (9.5)	NA	Died—36
Carol H. Yan [122]	All positive COVID-19 who completed survey(most are mild cases)	NA	29 (49.2)	Disturbance of consciousness—8 (22.22) Headache—39 (66.1) Myalgia/arthritis—37 (62.7) Ageusia—42 (71.2) Anosmia—40 (67.8)	With subjective olfaction score COVID-19 vs non-COVID-19 Death group vs recovered group	NA
Yan Deng [123]	Only dead and recovered patients admitted	NA	124	Myalgia or fatigue—57 Headache—13 (11.5)	NA	Died—109 Recovered—116
Jiaojiao Chu [124]	Only medical staff(54 tested, but only 38positive for nucleic acid tests)	39 (26–66)	24 (63.2)	Muscle ache—2 (5.3)	Common vs severe, positive RT-PCR vs negative	NA
Håkon Ihl-Hansen [125]	All consecutive admitted	72.5 (30–95)	28 (67)	New-onset confusion—8 (19)	Severe vs critical	NA
Parag Goyal [126]	First consecutive patients hospitalized, adults $\geq$ 18 years	62.2 (48.6–73.7)	238 (60.6)	Myalgia—107 (27.2)	Invasive mechanical ventilation vs no invasive mechanical ventilation	Died—40 (10.2) Discharged—260 (66.2)

**Table 1** (continued)

				Outcome data incomplete—93 (23.7) Discharge—85 (83.3) Died—17 (16.7) All recovered (12 still quarantined) All nine live birth Died—0 Discharged—91 (26.8) Hospitalized—183 (54.0) Died—65 (19.2) Died—1
Jianlei Cao [127]	All patients admitted	54 (37–67)	53	Muscle ache—35 (34.3) Non survivors vs survivors
De Chang [128]	All hospitalized patients	34 (34–48)	10 (77)	Myalgia—3 (23.1) Headache—3 (23.1) NA
Huijun Chen [129]	Only pregnant patients	26–40 years	NA	Myalgia—3 (33%) NA
Lang Wang [130]	Consecutive cases over 60 years old	69 (65–76)	166(49)	Myalgia—16 (4.7) Dizziness—13 (3.8) Headache—12 (3.5) Headache—6 Myalgia—1 (3.22) Infected in Europe vs china
Gianfranco Spiteri [131]	First cases in the WHO European region except UK	42(2–81)	25	NA NA Imported vs secondary vs tertiary (1 patient diagnosed with anal swab)
Yingxia Liu [132] Tianmin Xu [133]	Patients admitted Patients admitted	10–72 years NA	8 25	Myalgia—4 (33.3) Myalgia—8 (15.7) NA NA
Michael Chung [134]	Admitted patients who underwent chest CT Admitted and had CT chest done	51±14	13 (62)	Headache—3 (14) Muscle soreness—3 (14) NA
Heshui Shi [135]	All admitted patients	49.5±11.0	42 (52)	Headache—5 (6) Dizziness—2 (2) NA
Luhuan Yang [136]		55±17.1	98 (49.0)	Myalgia or malaise—44 (22.0) Headache 27—(13.5) NA
Wei Zhao [137]	Consecutive laboratory confirmed COVID-19 who underwent CT	44.44(17–75)	56 (55.4)	Myalgia or fatigue—17 (16.8) Emergency vs non-emergency group Only 30/32 (93.8%) lab confirmed (2 included based on clinical and epidemiological evidence)
Ya-nan Han [138]	All admitted patients	NA	16	Myalgia or fatigue-13(all adults) Only 30/32 (93.8%) lab confirmed (2 included based on clinical and epidemiological evidence)
Yang Wang [139]	Severely and critically ill (ICU)	64 (52–72)	179 (52.0)	Rhabdomyolysis—9 (2.6) Pediatrics vs adults Survivors vs non-survivors Died—133 (38.7) Discharged—185 (87.7) Hospitalized—26



**Fig. 1** Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram

population. Syncope was reported in three studies with incidence of 0.3% [45], 0.6% [49], and 4.8% [86]. Few studies showed an increase in creatine kinase, LDH, and myoglobin in COVID-19 patients [12, 62, 66].

### Acute cerebrovascular disease

Acute cerebrovascular disease (CVD) was reported in 0.5–5.9% [33, 48] of COVID-19 patients. Out of them, the most common type was acute ischaemic stroke and severe COVID-19 patients were more at risk of developing the acute CVD [33]. From these studies, the incidence of acute CVD in severe/ICU patients reported were 0.8–9.8% [33,

41]. The incidence of ischaemic stroke, hemorrhagic stroke, and cerebral venous thrombosis reported from various studies ranged from 0.4–4.9% [33, 48], 0.2–0.9% [38, 48], and 0.3–0.5% [33, 47] respectively. A study by Mao et al. [32] reported that two patients presented with hemiplegia without any typical COVID-19 symptoms. The median time to onset of cerebrovascular disease was 9 days. Another study by Li Y et.al [33] showed that acute CVD was more likely to be present with severe COVID-19; however, they were older, and had cardiovascular risk factors. These findings were similar to the above study by Mao et al. [32]. In both these studies, the laboratory parameters in patients with CNS symptoms were different from the other COVID-19

**Table 2** Prevalence of neurological manifestations reported from systematic assessment

	Studies (N)	Sample size (N)	Cases (n)	Prevalence (95% CI)
Smell disturbances	17	7919	2488	31.4% (30.4–32.4)
Taste disturbances	14	7033	1979	28.1% (27.1–29.2)
Headache	54	13,623	2751	20.2% (19.5–20.9)
Myalgia	38	11,169	2288	20.5% (19.7–21.2)
Disturbances in consciousness/altered mental status	9	6687	408	6.1% (5.5–6.7)
Syncope	3	1000	56	5.6% (4.3–7.2)
Dizziness	12	2595	137	5.3% (4.5–6.2)
Acute cerebrovascular disease	8	10,186	148	1.4% (1.2–1.7)
Ischaemic stroke	7	9268	108	1.2% (1.0–1.4)
Hemorrhagic stroke	7	12,704	60	0.5% (0.4–0.6)
Cerebral venous thrombosis	2	946	3	0.3% (0.1–0.9)
Seizures	5	2043	23	1.1% (0.7–1.7)
Ataxia	2	939	3	0.3% (0.1–0.9)

patients, with a higher white cell and neutrophil counts, reduced lymphocyte and platelet counts, elevated CRP and D-dimer levels [32, 33].

We found two studies that specifically studied the thrombotic complications in COVID-19 patients and found acute ischaemic stroke in COVID-19 patients receiving thromboprophylaxis [35, 36]. A retrospective observational case series in COVID-19 patients from Italy [144] reported six cases of stroke, four were ischaemic and two were hemorrhagic. Five of them had pre-existing vascular risk factors. Three patients with ischaemic stroke and one patient with hemorrhagic stroke showed hypercoagulable blood parameters [144]. Two studies reported six cases of stroke in young(< 50 years) COVID-19 patients, out of which three patients did not have any risk factors [145, 146].


Also there are multiple case reports and case series of ischaemic stroke including large artery [147], aneurysmal [148, 149] and non-aneurysmal SAH [51], deep cerebral venous thrombosis [150–157], hemorrhagic stroke [38, 158, 159] and CNS vasculitis [160] from all over the world in COVID-19 patients [38, 51, 147–173].

### Meningoencephalitis, encephalopathy, disturbances in consciousness

Several cases of meningoencephalitis and encephalopathy were reported in COVID-19 patients [39, 43, 49, 174–183]. The incidence of encephalitis reported in two retrospective studies was 0.03% [43] and 0.1% [49]. Only in four of the 15 reported cases of encephalitis, CSF RT-PCR test was

**Table 3** Meta-analysis, summary estimate of pooled prevalence and heterogeneity of each neurological manifestations

	Number of studies (N)	Summary estimate (%)	95% CI	$\chi^2$
Smell disturbances	17	35.8	(21.4, 50.2)	99.87
Taste disturbances	14	38.5	(24.0, 53.0)	99.65
Headache	54	14.7	(10.4, 18.9)	99.09
Myalgia	38	19.3	(15.1, 23.6)	98.98
Disturbances in consciousness/altered mental status	9	9.6	(4.9, 14.3)	98.26
Dizziness	12	6.1	(3.1, 9.2)	93.44
Acute cerebrovascular disease	8	2.3	(1.0, 3.6)	96.61
Ischaemic stroke	7	2.1	(0.9, 3.3)	96.67
Hemorrhagic stroke	7	0.4	(0.2, 0.6)	62.36
Cerebral venous thrombosis	2	0.3	(0.1, 0.6)	0.00
Syncope	3	1.8	(0.9, 4.6)	98.48
Ataxia	2	0.3	(0.1, 0.7)	0.00
Seizure	5	0.9	(0.5, 1.3)	9.03




**Fig. 2** Forest plot of each neurological manifestations

positive for SARS-CoV-2 RNA, and surprisingly two cases among them had negative nasopharyngeal swab [50, 174–176]. Two reports showed elevated levels of cytokines like IL-6, IL-8, TNF- $\alpha$ ,  $\beta$ 2-microglobulin, IP-10, MCP-1 in CSF [177, 181]. Interestingly, fluid from the surgical evacuation of subdural hematoma was positive for SARS-CoV-2 RT-PCR in a COVID-19 patient [184]. Isolated meningoencephalitis without any respiratory involvement has also been reported [175, 185]. Another case of rhombencephalitis as a rare complication of COVID-19 patient has been reported [186]. Few retrospective studies [32, 47, 49] reported seizures with the incidence ranging from 0.5–1.4% [32, 47]. Cases of all types of seizures like febrile seizures [42], focal seizures [180, 187–189], generalized tonic-clonic seizures [183, 190–192], myoclonic status epilepticus [193], status epilepticus [188, 194] and non-convulsive status epilepticus [46] were reported in COVID-19 patients.

Generally, the SARS-CoV-2 virus causes mild disease in children. However, a study from Italy showed a total five patients with seizures, and out of them, two had febrile seizures (three children had a known history of epilepsy, one child had a history of febrile seizures, one child had a first

episode of febrile seizures) [42]. Also, a case of a 6-week-old infant with SARS-CoV-2 in addition to rhinovirus, presenting with brief 10–15-s episodes of upward gaze and bilateral leg stiffening was reported with normal EEG and MRI brain [195]. Another case of an 11-year-old child with COVID-19 viral encephalitis has been reported, with CSF showing viral encephalitis picture [194].

PRES syndrome has also been reported in studies [47, 51]. Transient cortical blindness like presentation of PRES syndrome with MRI brain at admission revealing bilateral T2/FLAIR hyperintensities, especially left occipital, frontal cortical white matter and splenium of the corpus callosum and diffusion restriction in DWI revealing vasogenic edema has been reported [196]. Repeat MRI after 2 weeks showed a complete resolution of findings. Cases of acute necrotizing hemorrhagic encephalopathy [191, 197], hypoxic brain injury with encephalopathy [43, 47, 51, 65], delayed post-hypoxic leukoencephalopathy [198], mild encephalitis/encephalopathy with a reversible splenial lesion(MERS) [199], ADEM in elderly females [200, 201], MS plaque exacerbation [47] and CIS [176] were reported in SARS-CoV-2 infected patients.



**Fig. 3** Funnel plot for assessing publication bias of each neurological manifestations studied

Incidence of disturbances of consciousness/delirium ranged from 3.3–19.6% [49, 100] in retrospective studies. S.R. Beach, et al. [202] reported four cases of elderly COVID-19 patients, who presented to the hospital with altered mental status without any respiratory complaints, and only one among them developed respiratory complaints during the hospital stay. Similar cases have been reported in elderly patients from Saudi Arabia [203], Norway [204] and China [205]. An observational case series from France [39] in 58 COVID-19 patients with ARDS admitted in ICU reported agitation in 40(69%) patients, confusion in 26 of 40 patients, diffuse corticospinal tract signs in 39 patients (67%) and out of the 45 patients discharged, 15(33%) had a dysexecutive syndrome. MRI Brain showed enhancement of leptomeningeal spaces in eight patients, bilateral frontotemporal hypoperfusion in 11 patients who underwent perfusion imaging, two asymptomatic patients with small acute ischaemic stroke and one patient with subacute ischaemic stroke.

### Guillain-Barré syndrome

There are multiple reports of GBS in patients with confirmed COVID-19. GBS has also been reported to be a presenting feature in one case report by Zhao H et al. [206] where the patient, later on, developed fever and other symptoms of COVID-19. All the variants of GBS like AIDP, AMAN, AMSAN has been reported in COVID-19 patients [47, 206–219] including both para [206–212, 220–223] and post-infectious pattern [210, 211, 214–219, 224–226]. Toscano et al. [227] reported a series of five patients of COVID-19 with GBS, with the interval between the onset of fever, cough and symptoms of GBS ranging from 5 to 10 days. Cases of MFS were also reported [47, 226, 228, 229]. One case of MFS was associated with a positive serum GD1b-IgG antibody [228]. Other rare variants reported were GBS/MF overlap syndrome [219], AMSAN variants with severe autonomic neuropathy [219], facial diplegia [222, 227] and post-infectious pattern of the demyelinating type of GBS with

brainstem and cervical leptomeningeal enhancement [225]. Cranial neuropathies with abnormal perineural or cranial nerve findings [230], multiple cranial neuropathies [211, 219], peripheral motor neuropathy [231] and ataxia [32, 43, 51] are all reported as presentations of COVID-19.

### Other neurological manifestations

The incidence of rhabdomyolysis has been reported between 0.2–2.6% in different studies [11, 49, 139]. A report illustrates a 38 year-old COVID-19 patient presenting with fever, dyspnea, and severe myalgia, with high creatine kinase (>42,670 U/L) and LDH (4301 U/La) and was diagnosed as viral myositis [232]. Another two cases of adult COVID-19 patients with lower extremity pain and weakness with rhabdomyolysis with high creatine kinase and LDH were reported [233, 234]. First case developed rhabdomyolysis on the 9th day of admission [233] and 2nd case presented to the hospital with rhabdomyolysis [234]. An isolated case of post-infectious myelitis has been reported from Germany in a COVID-19 patient [235].

Three cases of generalized brainstem type of myoclonus were reported from Spain, with normal CSF study in one patient (others not done) and normal imaging findings. However, nasopharyngeal RT-PCR for SARS-CoV-2 was positive in only one patient. In all these patients, EEG was showing mild diffuse slowing without any epileptic activity [236]. Paresthesias [51] and cutaneous hyperesthesia [237] were reported as a presentation in COVID-19 patients. A case of COVID-19 patient with oropharyngeal dysphagia followed by aspiration pneumonia, taste impairment, impaired pharyngolaryngeal sensation, and nasopharyngeal contractile dysfunction with absent gag reflex was reported from Japan [238]. Visual symptoms were also reported in a few studies. Mao L et al. [32] reported visual impairment in 1.4% of the COVID-19 patients. Cases of optic neuritis [49], isolated central retinal artery occlusion [239], non-arteritic type of posterior ischaemic optic neuropathy (PION) [240] as a COVID-19

manifestation were also reported. The summary of all the neurological manifestations reported in COVID-19 is given in Table 4.

## Heterogeneity

The heterogeneity was high in most of the neurological manifestations studied except for hemorrhagic stroke (medium), cerebral venous thrombosis (low), seizure (low), and ataxia (low). The funnel plots were symmetric in hemorrhagic stroke, ataxia, seizures, cerebral venous thrombosis and myalgia, which is pointing towards no bias in the selection of publications that are included in the study. However, the funnel plots were asymmetric in other neurological manifestations studied, which pointed towards the heterogeneity in the

studies undertaken or bias in the selection of publications included in the study.

## Discussion

In this systematic review and meta-analysis, we assessed the neurological manifestations, risk factors, mortality, laboratory parameters, and imaging findings in those patients with neurological features. Involving 30,159 patients, our meta-analysis is the first and most comprehensive study about the neurological manifestations of COVID-19.

The most common neurological manifestations reported were smell and taste disturbances. Another interesting finding is the geographical variations in the frequency of smell and

**Table 4** Summary of all the neurological manifestations of COVID-19

Non-specific	CNS manifestations	Peripheral nervous system manifestations
Myalgia	Disturbances in consciousness	Taste disturbances (ageusia, reduced taste, distorted taste)
Headache	Agitation	Smell disturbances (anosmia, phantosmia, parosmia)
Dizziness	Pathological wakefulness	Vision impairment
Vertigo	Encephalitis	Nerve pain/neuralgia
Lightheadedness	Encephalopathy	Skeletal muscle injury
	Acute necrotizing hemorrhagic encephalopathy	Rhabdomyolysis
	Post-hypoxic encephalopathy/hypoxic ischaemic brain injury	Myositis
	Mild encephalitis/encephalopathy with a reversible splenial lesion (MERS)	Occipital neuralgia
	Rhombencephalitis/myelitis	Dysautonomia
	Seizure (focal, GTCS, NCSE, status epilepticus, febrile seizures)	Extraocular muscle abnormalities
	Acute cerebrovascular disease	Isolated unilateral facial palsy
	Ischaemic stroke/TIA	GBS (AIDP/AMAN/AMSAN)
	Hemorrhagic stroke	DP (facial diplegia) variant of GBS
	SAH (aneurysmal and non-aneurysmal)	Miller Fisher syndrome
	Cerebral venous sinus thrombosis	Cranial neuropathy
	Ataxia	Oropharyngeal dysphagia
	Dysexecutive syndrome	Optic neuritis
	Corticospinal tract signs	Posterior ischaemic optic neuropathy (non-arteritic) (PION)
	Syncope	Central retinal artery occlusion
	Short term memory loss	Cutaneous hyperesthesia
	Movement disorders	Parasthesias
	Neuropsychiatric symptoms	
	PRES syndrome	
	MS plaque exacerbation	
	Clinically isolated syndrome (CIS)	
	ADEM	
	Post-infectious myelitis	
	Generalized brainstem type of myoclonus	
	CNS vasculitis	

taste disturbances. High incidence of smell and taste disturbances were noted in studies from most of the European countries [54] while studies from Asian countries showed a lower incidence [32]. However, most of the studies which reported a higher incidence of smell and taste disturbances evaluated mainly olfactory and taste symptoms only and studied mild to moderate cases and excluded severe/ICU patients compared with studies with lower incidence. This bias might have caused under-reporting of smell and taste disturbances in severe/ICU patients or could also be because of decreased awareness of investigator about these symptoms at the beginning of the pandemic. Supporting our assumption, a study from Spain which evaluated 841 COVID-19 patients with neurological manifestations reported only 4.9% of cases of smell disturbances and 6.2% cases of taste disturbances [49]. Other possibilities for these variations are, the difference in affinity of SARS-CoV-2 to tissues between populations, a different strain of mutated virus circulating in Europe compared with Asian countries. However, more studies are required to confirm these assumptions. Interestingly a study by Wan Y et al. [241] predicted that binding affinity between 2019-nCoV and human ACE2 may be enhanced by a single N501T mutation. Also, ACE2 receptors are highly expressed by sustentacular cells of the olfactory epithelium. Olfactory and taste disorders were more common in younger patients [52, 140] most occurs in the early stages as initial manifestations of the disease and even as the only manifestation of COVID-19. Hence, olfactory and gustatory disorders can be the initial and early manifestations of COVID-19 and early identification of these symptoms might lead to early diagnosis and disease containment.

Non-specific neurological manifestations could be just systemic features of a viral infection. Similar to olfactory disturbances, the incidence of myalgia, headache, and dizziness also shows geographical variations with the highest incidence reported from Europe, the USA, and Chile. The incidence of non-specific symptoms was lower in children. We noticed that non-specific symptoms were higher among the studies conducted in health care workers. This may be due to increased knowledge and awareness of the symptoms and disease.

The most common type of acute CVD reported was an ischaemic stroke. Hemorrhagic stroke, deep cerebral venous thrombosis, SAH (both non-aneurysmal and aneurysmal), and TIA were also reported; however, with much lesser prevalence. Severe infection or ICU requirement, older age, cardiovascular risk factors, prior co-morbidities, and hypercoagulable lab parameters were found to be a risk factor for developing acute CVD [32, 33]. The apparent association of COVID-19 and stroke is likely due to the sharing of similar risk factors. The severity of COVID-19 has been proved to be directly related to the presence of co-morbidities like hypertension and DM. An earlier meta-analysis by Yang J et al. [242] comprising [46, 243] COVID-19 patients reported the prevalence of risk

factors, hypertension in 21.1%, DM in 9.7%, and cardiovascular diseases in 8.4%. Also, hypercoagulable blood parameters as shown by Li Y et.al [33], can lead to ischaemic stroke and cerebral venous thrombosis. Nervous system involvement in SARS-CoV-2 infection can be due to direct invasion of neural tissues, inflammatory response, or immune dysregulation. The SARS-CoV-2 virus uses the ACE2 and TMPRSS2 for entry to the host cell and it is one of the main determinants of infectivity [241, 244]. Susceptibility to infection correlated with ACE2 expression in previous studies [245].

Very few retrospective studies showed meningoencephalitis as a presentation of COVID-19; however, there are multiple case reports from all over the world. The probable mechanism can again be direct invasion via the hematogenous route or retrograde pathway via peripheral nerve terminals. Two studies even showed higher levels of inflammatory cytokines in the CSF analysis of these patients [177, 181]. SARS-CoV-2 could trigger a seizure in predisposing patients through neurotropic mechanisms as explained earlier [188]. However, more evaluation is required in this field to find a temporal factor. All types of seizures were reported like febrile seizures, focal seizures, generalized tonic-clonic seizures, status epilepticus and myoclonic status epilepticus, NCSE and also brainstem type of myoclonus. Demyelinating disorders like ADEM, exacerbation of MS plaque, and the clinically isolated syndrome were all reported in COVID-19 patients.

Cases of GBS and its variants were also reported in COVID-19. Both post-infectious and pre-infectious pattern of GBS were reported. The most common type of GBS reported was AIDP. Other variants like AMAN, AMSAN, Miller Fisher syndrome, and facial diplegic variant were also reported. Patients presenting as GBS without any other typical symptoms of COVID-19 were also reported. Possible pathogenesis of GBS in COVID-19 includes immune dysregulation secondary to systemic hyper inflammation and cytokines produced as described by McGonagle et al. [246] and Quin et al. [247]. Hence, it is important to suspect and test for COVID-19 in those patients presenting with GBS and MFS. However, more studies are required to conclude that these cases were not just coincidental and COVID-19 itself is a trigger for GBS and MFS. GBS was also reported in other recent important viral infections like MERS-CoV [248] and Zika virus [243].

Change in laboratory parameters was also reported in COVID-19 patients with neurological manifestations like higher white cell and neutrophil counts, reduced lymphocyte and platelet counts, elevated CRP and D-dimer levels, and higher levels of creatine kinase, LDH, and myoglobin [12, 32, 33, 62].

High heterogeneity in our study could be because of differences in the selection of patients and ethnicity, the severity of the disease, co-morbidities, only a few studies evaluated neurological symptoms specifically, variation in the number of patients in different studies, or due to publication bias and differences in the methodology among the studies.

## Comparison with previous systematic reviews

Earlier meta-analyses addressing general clinical features in COVID-19 were published. One such study showed myalgia in (28.5%; 95%CI 21.2–36.2), headache (14.0%; 95%CI 9.9–18.6), and dizziness (7.6%; 95%CI 0.0–23.5) [249]. Our results also found similar results for myalgia, headache, and dizziness, i.e. (19.3%; 95%CI 15.1–23.6), (14.7%; 95%CI 10.4–18.9), and (6.1%; 95%CI 3.1–9.2) respectively. Another similar meta-analysis also showed myalgia in (21.9%; 95%CI 17.7–26.4) and headache in (11.3%; 95%CI 8.9–14.0) [250]. One more study reported the prevalence of headache as (8.0%; 95%CI 5.7–10.2) [251]. However, no meta-analyses are published on the specific neurological manifestations till now.

## Strengths and limitations

The strength of our study is that we did a comprehensive search in all the electronic databases. Study limitations include high heterogeneity in the estimation of the prevalence of some neurological manifestations, the inclusion of studies with very small sample size, and lack of meta-regression analysis. We excluded studies in languages other than English where translation was not possible. Most of the included studies were of moderate quality. More good-quality prospective cohort studies are required to establish that the neurological manifestations reported in the studies were not just coincidental.

## Conclusions

In conclusion, our study showed neurological manifestations are common in COVID-19 and are even present as the only symptom without any other manifestation of the respiratory system involvement. Hence it is important to suspect every COVID-19 patient with neurological manifestations. In this pandemic, a neurologist needs to take necessary precautions while examining the patients presenting to them. Also, some symptoms like smell and taste disturbance can be used as a screening tool for SARS-CoV-2 infection and can help isolate suspected patients earlier to avoid the spread of the disease.

**Author contributions** TF and AP conceptualized the study and searched and screened the literature. PD and RNC extracted and analysed the data. RNC contributed to figures, tables, and interpretation of images. KC was involved in extraction of data. DJ, VNM, TF, and AP were involved in study design, data interpretation, and data analysis. RM and MP did the statistical analysis. AK drafted the manuscript, data collection, figures. VKS did literature search, drafted the manuscript, and contributed to study design. TF wrote the first draft of the manuscript with input from AP.

**Data availability** All data available on request.

## Compliance with ethical standards

**Conflict of interest** The authors declare that they have no competing interests.

**Ethics approval** Not applicable.

**Consent to participate** Not applicable.

**Consent for publication** Not applicable.

## References

1. de Wilde AH, Snijder EJ, Kikkert M, van Hemert MJ (2018) Host factors in coronavirus replication. *Curr Top Microbiol Immunol* 419:1–42. [https://doi.org/10.1007/82\\_2017\\_25](https://doi.org/10.1007/82_2017_25)
2. Drexler JF, Corman VM, Drosten C (2014) Ecology, evolution and classification of bat coronaviruses in the aftermath of SARS. *Antivir Res* 101:45–56. <https://doi.org/10.1016/j.antiviral.2013.10.013>
3. de Wit E, van Doremalen N, Falzarano D, Munster VJ (2016) SARS and MERS: recent insights into emerging coronaviruses. *Nat Rev Microbiol* 14(8):523–534. <https://doi.org/10.1038/nrmicro.2016.81>
4. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, Niu P, Zhan F, Ma X, Wang D, Xu W, Wu G, Gao GF, Tan W, China Novel Coronavirus Investigating and Research Team (2020) A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 382(8):727–733. <https://doi.org/10.1056/NEJMoa2001017>
5. Letko M, Marzi A, Munster V (2020) Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol* 5(4):562–569. <https://doi.org/10.1038/s41564-020-0688-y>
6. Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, Chen HD, Chen J, Luo Y, Guo H, Jiang RD, Liu MQ, Chen Y, Shen XR, Wang X, Zheng XS, Zhao K, Chen QJ, Deng F, Liu LL, Yan B, Zhan FX, Wang YY, Xiao GF, Shi ZL (2020) A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 579(7798):270–273. <https://doi.org/10.1038/s41586-020-2012-7>
7. Novel Coronavirus – China. World Health Organization. <https://www.who.int/csr/don/12-january-2020-novel-coronavirus-china/en/>. Published January 12, 2020. Accessed June 19, 2020
8. Chen X, Yu B (2020) First two months of the 2019 coronavirus disease (COVID-19) epidemic in China: real-time surveillance and evaluation with a second derivative model. *Glob Health Res Policy* 5(1):7. <https://doi.org/10.1186/s41256-020-00137-4>
9. Coronavirus Disease (COVID-19) - events as they happen. World Health Organization. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen>. Published January 12, 2020. Accessed June 19, 2020
10. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y, Zhao Y, Li Y, Wang X, Peng Z (2020) Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 323(11):1061–1069. <https://doi.org/10.1001/jama.2020.1585>
11. Guan W, Ni Z, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC, du B, Li LJ, Zeng G, Yuen KY, Chen RC, Tang CL, Wang T, Chen PY, Xiang J, Li SY, Wang JL, Liang ZJ, Peng YX, Wei L, Liu Y, Hu YH, Peng P, Wang JM, Liu JY, Chen Z, Li G, Zheng ZJ, Qiu SQ, Luo J, Ye CJ, Zhu SY, Zhong NS, China

- Medical Treatment Expert Group for Covid-19 (2020) Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 382(18):1708–1720. <https://doi.org/10.1056/NEJMoa2002032>
- 12. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Xia J', Yu T, Zhang X, Zhang L (2020) Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 395(10223):507–513. [https://doi.org/10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7)
  - 13. Wu C, Chen X, Cai Y, Xia J', Zhou X, Xu S, Huang H, Zhang L, Zhou X, du C, Zhang Y, Song J, Wang S, Chao Y, Yang Z, Xu J, Zhou X, Chen D, Xiong W, Xu L, Zhou F, Jiang J, Bai C, Zheng J, Song Y (2020) Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med* 180(7): 934–943. <https://doi.org/10.1001/jamainternmed.2020.0994>
  - 14. Netland J, Meyerholz DK, Moore S, Cassell M, Perlman S (2008) Severe acute respiratory syndrome coronavirus infection causes neuronal death in the absence of encephalitis in mice transgenic for human ACE2. *J Virol* 82(15):7264–7275. <https://doi.org/10.1128/JVI.00737-08>
  - 15. Li K, Wohlford-Lenane C, Perlman S, Zhao J, Jewell AK, Reznikov LR, Gibson-Corley KN, Meyerholz DK, McCray PB Jr (2016) Middle East respiratory syndrome coronavirus causes multiple organ damage and lethal disease in mice transgenic for human dipeptidyl peptidase 4. *J Infect Dis* 213(5):712–722. <https://doi.org/10.1093/infdis/jiv499>
  - 16. Baig AM, Khaleeq A, Ali U, Syeda H (2020) Evidence of the COVID-19 virus targeting the CNS: tissue distribution, host-virus interaction, and proposed neurotropic mechanisms. *ACS Chem Neurosci* 11(7):995–998. <https://doi.org/10.1021/acscchemneuro.0c00122>
  - 17. Tsai L, Hsieh S, Chao C et al (2004) Neuromuscular disorders in severe acute respiratory syndrome. *Arch Neurol* 61(11):1669–1673. <https://doi.org/10.1001/archneur.61.11.1669>
  - 18. Umapathi T, Kor AC, Venkatasubramanian N, Lim CCT, Pang BC, Yeo TT, Lee CC, Lim PL, Ponnudurai K, Chuah KL, Tan PH, Tai DYH, Ang SPB (2004) Large artery ischaemic stroke in severe acute respiratory syndrome (SARS). *J Neurol* 251(10):1227–1231. <https://doi.org/10.1007/s00415-004-0519-8>
  - 19. Hwang CS (2006) Olfactory neuropathy in severe acute respiratory syndrome: report of a case. *Acta Neurol Taiwanica* 15(1):26–28
  - 20. Li Y, Li H, Fan R, Wen B, Zhang J, Cao X, Wang C, Song Z, Li S, Li X, Lv X, Qu X, Huang R, Liu W (2016) Coronavirus infections in the central nervous system and respiratory tract show distinct features in hospitalized children. *Intervirology*. 59(3):163–169. <https://doi.org/10.1159/000453066>
  - 21. Morfopoulou S, Brown JR, Davies EG, Anderson G, Virasami A, Qasim W, Chong WK, Hubank M, Plagnol V, Desforges M, Jacques TS, Talbot PJ, Breuer J (2016) Human coronavirus OC43 associated with fatal encephalitis. *N Engl J Med* 375(5): 497–498. <https://doi.org/10.1056/NEJMc1509458>
  - 22. Arabi YM, Harthi A, Hussein J, Bouchama A, Johani S, Hajer AH, Saeed BT, Wahbi A, Saedy A, AlDabbagh T, Okaili R, Sadat M, Balkhy H (2015) Severe neurologic syndrome associated with Middle East respiratory syndrome corona virus (MERS-CoV). *Infection*. 43(4):495–501. <https://doi.org/10.1007/s15010-015-0720-y>
  - 23. Ann Yeh E, Collins A, Cohen ME, Duffner PK, Faden H (2004) Detection of coronavirus in the central nervous system of a child with acute disseminated encephalomyelitis. *PEDIATRICS*. 113(1):e73–e76. <https://doi.org/10.1542/peds.113.1.e73>
  - 24. Kim J-E, Heo J-H, Kim H, Song SH, Park SS, Park TH, Ahn JY, Kim MK, Choi JP (2017) Neurological complications during treatment of Middle East Respiratory Syndrome. *J Clin Neurol* 13(3):227–233. <https://doi.org/10.3988/jcn.2017.13.3.227>
  - 25. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 6(7):e1000097. <https://doi.org/10.1371/journal.pmed.1000097>
  - 26. Stang A (2010) Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 25(9):603–605. <https://doi.org/10.1007/s10654-010-9491-z>
  - 27. Wells GA, Shea B, O'Connell D, et al. The Ottawa Hospital Research Institute. [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp). Published October 19, 2009. Accessed June 19, 2020
  - 28. Schulz KF, Altman DG, Moher D, for the CONSORT Group (2010) CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. *BMJ* 340(mar23 1): c332–c332. <https://doi.org/10.1136/bmj.c332>
  - 29. Viechtbauer W. Conducting meta-analyses in R with the metafor Package. *J Stat Softw*. 2010;36(3). <https://doi.org/10.18637/jss.v036.i03>
  - 30. Kontopantelis E, Reeves D (2012) Performance of statistical methods for meta-analysis when true study effects are non-normally distributed: a comparison between DerSimonian–Laird and restricted maximum likelihood. *Stat Methods Med Res* 21(6): 657–659. <https://doi.org/10.1177/0962280211413451>
  - 31. Higgins JPT (2003) Measuring inconsistency in meta-analyses. *BMJ*. 327(7414):557–560. <https://doi.org/10.1136/bmj.327.7414.557>
  - 32. Mao L, Jin H, Wang M, Hu Y, Chen S, He Q, Chang J, Hong C, Zhou Y, Wang D, Miao X, Li Y, Hu B (2020) Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol* 77(6):683–690. <https://doi.org/10.1001/jamaneurol.2020.1127>
  - 33. Li Y, Li M, Wang M, et al. Acute cerebrovascular disease following COVID-19: a single center, retrospective, observational study. *Stroke Vasc Neurol*. Published online July 2, 2020:svn-2020-000431. <https://doi.org/10.1136/svn-2020-000431>
  - 34. Lu L, Xiong W, Liu D, Liu J, Yang D, Li N, Mu J, Guo J, Li W, Wang G, Gao H, Zhang Y, Lin M, Chen L, Shen S, Zhang H, Sander JW, Luo J, Chen S, Zhou D (2020) New onset acute symptomatic seizure and risk factors in coronavirus disease 2019: a retrospective multicenter study. *Epilepsia*. 61(6):e49–e53. <https://doi.org/10.1111/epi.16524>
  - 35. Klok FA, Kruip MJHA, van der Meer NJM, Arbous MS, Gomers D, Kant KM, Kaptein FHJ, van Paassen J, Stals MAM, Huisman MV, Endeman H (2020) Confirmation of the high cumulative incidence of thrombotic complications in critically ill ICU patients with COVID-19: An updated analysis. *Thromb Res* 191:148–150. <https://doi.org/10.1016/j.thromres.2020.04.041>
  - 36. Lodigiani C, Iapichino G, Carenzo L, Cecconi M, Ferrazzi P, Sebastian T, Kucher N, Studt JD, Sacco C, Alexia B, Sandri MT, Barco S, Humanitas COVID-19 Task Force (2020) Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. *Thromb Res* 191: 9–14. <https://doi.org/10.1016/j.thromres.2020.04.024>
  - 37. Fraisse M, Logre E, Pajot O, Mentec H, Plantefève G, Contou D (2020) Thrombotic and hemorrhagic events in critically ill COVID-19 patients: a French monocenter retrospective study. *Crit Care* 24(1):275. <https://doi.org/10.1186/s13054-020-03025-y>
  - 38. Dogra S, Jain R, Cao M, Bilaloglu S, Zagzag D, Hochman S, Lewis A, Melmed K, Hochman K, Horwitz L, Galetta S, Berger J (2020) Hemorrhagic stroke and anticoagulation in COVID-19. *J Stroke Cerebrovasc Dis* 29(8):104984. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2020.104984>

39. Helms J, Kremer S, Merdji H, Clere-Jehl R, Schenck M, Kummerlen C, Collange O, Boulay C, Fafi-Kremer S, Ohana M, Anheim M, Meziani F (2020) Neurologic features in severe SARS-CoV-2 infection. *N Engl J Med* 382(23):2268–2270. <https://doi.org/10.1056/NEJMc2008597>
40. Helms J, Tacquard C, Severac F et al (2020) High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med* 46(6):1089–1098. <https://doi.org/10.1007/s00134-020-06062-x>
41. Kandemirli SG, Dogan L, Sarikaya ZT, et al. (2020) Brain MRI findings in patients in the intensive care unit with COVID-19 infection [published online ahead of print, 2020 May 8]. *Radiology*. 201697. <https://doi.org/10.1148/radiol.2020201697>
42. Garazzino S, Montagnani C, Donà D, Meini A, Felici E, Vergine G, Bernardi S, Giaccherio R, Lo Vecchio A, Marchisio P, Nicolini G, Pierantoni L, Rabbone I, Banderoli G, Denina M, Venturini E, Krzysztofiak A, Badolato R, Bianchini S, Galli L, Villani A, Castelli-Gattinara G, Italian SITIP-SIP Pediatric Infection Study Group, Italian SITIP-SIP SARS-CoV-2 paediatric infection study group\* (2020) Multicentre Italian study of SARS-CoV-2 infection in children and adolescents, preliminary data as at 10 April 2020. *Euro Surveill* 25(18):2000600. <https://doi.org/10.2807/1560-7917.ES.2020.25.18.2000600>
43. Jain R, Young M, Dogra S, Kennedy H, Nguyen V, Jones S, Bilaloglu S, Hochman K, Raz E, Galetta S, Horwitz L (2020) COVID-19 related neuroimaging findings: a signal of thromboembolic complications and a strong prognostic marker of poor patient outcome. *J Neurol Sci* 414:116923. <https://doi.org/10.1016/j.jns.2020.116923>
44. Benussi A, Pilotto A, Premi E, et al. Clinical characteristics and outcomes of inpatients with neurologic disease and COVID-19 in Brescia, Lombardy, Italy [published online ahead of print, 2020 May 22]. *Neurology*. 2020; <https://doi.org/10.1212/WNL.0000000000009848>
45. Xiong W, Mu J, Guo J, et al. New onset neurologic events in people with COVID-19 infection in three regions in China [published online ahead of print, 2020 Jun 17]. *Neurology*. 2020; <https://doi.org/10.1212/WNL.00000000000010034>
46. Scullen T, Keen J, Mathkour M, Dumont AS, Kahn L (2020) Coronavirus 2019 (COVID-19)-associated encephalopathies and cerebrovascular disease: the New Orleans experience [published online ahead of print, 2020 May 28]. *World Neurosurg*. S1878-8750(20)31163-3. <https://doi.org/10.1016/j.wneu.2020.05.192>
47. Mahammed A, Saba L, Vagal A, et al. (2020) Imaging in neurological disease of hospitalized COVID-19 patients: an Italian multicenter retrospective observational study [published online ahead of print, 2020 May 21]. *Radiology*. 201933. <https://doi.org/10.1148/radiol.2020201933>
48. Radmanesh A, Raz E, Zan E, Derman A, Kaminetzky M (2020) Brain imaging use and findings in COVID-19: a single academic center experience in the epicenter of disease in the United States. *AJNR Am J Neuroradiol* 41(7):1179–1183. <https://doi.org/10.3174/ajnr.A6610>
49. Romero-Sánchez CM, Díaz-Maroto I, Fernández-Díaz E, et al. (2020) Neurologic manifestations in hospitalized patients with COVID-19: The ALBACOVID registry [published online ahead of print, 2020 Jun 1]. *Neurology*. <https://doi.org/10.1212/WNL.0000000000009937>
50. Kremer S, Lersy F, de Sèze J, et al. (2020) Brain MRI findings in severe COVID-19: a retrospective observational study [published online ahead of print, 2020 Jun 16]. *Radiology* ;202222. <https://doi.org/10.1148/radiol.2020202222>
51. Pinna P, Grewal P, Hall JP, Tavarez T, Dafer RM, Garg R, Osteraas ND, Pellack DR, Asthana A, Fegan K, Patel V, Conners JJ, John S, Silva ID (2020) Neurological manifestations and COVID-19: experiences from a tertiary care center at the frontline. *J Neurol Sci* 415:116969. <https://doi.org/10.1016/j.jns.2020.116969>
52. Beltrán-Corbellini Á, Chico-García JL, Martínez-Poles J, et al. (2020) Acute-onset smell and taste disorders in the context of COVID-19: a pilot multicentre polymerase chain reaction based case-control study [published online ahead of print, 2020 Apr 22]. *Eur J Neurol*. <https://doi.org/10.1111/ene.14273>
53. Giacomelli A, Pezzati L, Conti F, et al. (2020) Self-reported olfactory and taste disorders in SARS-CoV-2 patients: a cross-sectional study [published online ahead of print, 2020 Mar 26]. *Clin Infect Dis*. ciaa330. <https://doi.org/10.1093/cid/ciaa330>
54. Lechien JR, Chiesa-Estomba CM, De Sisti DR, et al (2020) Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Otorhinolaryngol* 277(8): 2251–2261. <https://doi.org/10.1007/s00405-020-05965-1>
55. Spinato G, Fabbris C, Polesel J, Cazzadori D, Borsetto D, Hopkins C, Boscolo-Rizzo P (2020) Alterations in smell or taste in mildly symptomatic outpatients with SARS-CoV-2 infection. *JAMA* 323(20):2089–2090. <https://doi.org/10.1001/jama.2020.6771>
56. Vaira LA, Deiana G, Fois AG, Pirina P, Madeddu G, de Vito A, Babudieri S, Petrocelli M, Serra A, Bussu F, Ligas E, Salzano G, de Riu G (2020) Objective evaluation of anosmia and ageusia in COVID-19 patients: single-center experience on 72 cases. *Head Neck* 42(6):1252–1258. <https://doi.org/10.1002/hed.26204>
57. Vaira LA, Salzano G, Petrocelli M, Deiana G, Salzano FA, De Riu G (2020) Validation of a self-administered olfactory and gustatory test for the remotely evaluation of COVID-19 patients in home quarantine. *Head Neck* 42(7):1570–1576. <https://doi.org/10.1002/hed.26228>
58. Vaira LA, Hopkins C, Salzano G, Petrocelli M, Melis A, Cucurullo M, Ferrari M, Gagliardini L, Pipolo C, Deiana G, Fiore V, de Vito A, Turra N, Canu S, Maglio A, Serra A, Bussu F, Madeddu G, Babudieri S, Giuseppe Fois A, Pirina P, Salzano FA, de Riu P, Biglioli F, de Riu G (2020) Olfactory and gustatory function impairment in COVID-19 patients: Italian objective multicenter-study. *Head Neck* 42(7):1560–1569. <https://doi.org/10.1002/hed.26269>
59. Lee Y, Min P, Lee S, Kim S-W (2020) Prevalence and duration of acute loss of smell or taste in COVID-19 patients. *J Korean Med Sci* 35(18):e174. <https://doi.org/10.3346/jkms.2020.35.e174>
60. Speth MM, Singer-Cornelius T, Oberle M, Gengler I, Brockmeier SJ, Sedaghat AR (2020) Olfactory dysfunction and sinonasal symptomatology in COVID-19: prevalence, severity, timing, and associated characteristics. *Otolaryngol Head Neck Surg* 163(1): 114–120. <https://doi.org/10.1177/0194599820929185>
61. Klopfenstein T, Kadiane-Oussou NJ, Toko L, et al. (2020) Features of anosmia in COVID-19 [published online ahead of print, 2020 Apr 17]. *Med Mal Infect*. S0399-077X(20)30110-4. <https://doi.org/10.1016/j.medmal.2020.04.006>
62. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B (2020) Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395(10223):497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)
63. Wu C, Chen X, Cai Y, Xia J, Zhou X, Xu S, Huang H, Zhang L, Zhou X, du C, Zhang Y, Song J, Wang S, Chao Y, Yang Z, Xu J, Zhou X, Chen D, Xiong W, Xu L, Zhou F, Jiang J, Bai C, Zheng J, Song Y (2020) Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med* 180(7): 934–943. <https://doi.org/10.1001/jamainternmed.2020.0994>
64. Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, Wu Y, Zhang L, Yu Z, Fang M, Yu T, Wang Y, Pan S, Zou X, Yuan S, Shang Y (2020)

- Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 8(5):475–481. [https://doi.org/10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5)
65. Chen T, Wu D, Chen H, Yan W, Yang D, Chen G, Ma K, Xu D, Yu H, Wang H, Wang T, Guo W, Chen J, Ding C, Zhang X, Huang J, Han M, Li S, Luo X, Zhao J, Ning Q (2020) Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study [published correction appears in BMJ. 2020 Mar 31;368:m1295]. *BMJ* 368:m1091 Published 2020 Mar 26. <https://doi.org/10.1136/bmj.m1091>
66. Du Y, Tu L, Zhu P et al (2020) Clinical features of 85 fatal cases of COVID-19 from Wuhan. A retrospective observational study. *Am J Respir Crit Care Med* 201(11):1372–1379. <https://doi.org/10.1164/rccm.202003-0543OC>
67. Zheng Y, Xu H, Yang M, Zeng Y, Chen H, Liu R, Li Q, Zhang N, Wang D (2020) Epidemiological characteristics and clinical features of 32 critical and 67 noncritical cases of COVID-19 in Chengdu. *J Clin Virol* 127:104366. <https://doi.org/10.1016/j.jcv.2020.104366>
68. Rodriguez-Morales AJ, Rodriguez-Morales AG, Méndez CA, Hernández-Botero S. Tracing new clinical manifestations in patients with COVID-19 in Chile and its potential relationship with the SARS-CoV-2 divergence [published online ahead of print, 2020 Apr 18]. *Curr Trop Med Rep* 2020;1–4. <https://doi.org/10.1007/s40475-020-00205-2>
69. Wang F, Yang Y, Dong K, Yan Y, Zhang S, Ren H, Yu X, Shi X (2020) Clinical characteristics of 28 patients with diabetes and COVID-19 in Wuhan, China. *Endocr Pract* 26(6):668–674. <https://doi.org/10.4158/EP-2020-0108>
70. Wan S, Xiang Y, Fang W, Zheng Y, Li B, Hu Y, Lang C, Huang D, Sun Q, Xiong Y, Huang X, Lv J, Luo Y, Shen L, Yang H, Huang G, Yang R (2020) Clinical features and treatment of COVID-19 patients in northeast Chongqing. *J Med Virol* 92(7):797–806. <https://doi.org/10.1002/jmv.25783>
71. Wang Z, Yang B, Li Q, Wen L, Zhang R (2020) Clinical features of 69 cases with coronavirus disease 2019 in Wuhan, China [published online ahead of print, 2020 Mar 16]. *Clin Infect Dis*. ciaa272. <https://doi.org/10.1093/cid/ciaa272>
72. Sun D, Li H, Lu X-X, Xiao H, Ren J, Zhang FR, Liu ZS (2020) Clinical features of severe pediatric patients with coronavirus disease 2019 in Wuhan: a single center's observational study. *World J Pediatr* 16(3):251–259. <https://doi.org/10.1007/s12519-020-00354-4>
73. Tian S, Hu N, Lou J, Chen K, Kang X, Xiang Z, Chen H, Wang D, Liu N, Liu D, Chen G, Zhang Y, Li D, Li J, Lian H, Niu S, Zhang L, Zhang J (2020) Characteristics of COVID-19 infection in Beijing. *J Inf Secur* 80(4):401–406. <https://doi.org/10.1016/j.jinf.2020.02.018>
74. Zhou F, Yu T, Du R et al (2020) Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 395(10229):1054–1062. [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3)
75. Du N, Chen H, Zhang Q et al (2020) A case series describing the epidemiology and clinical characteristics of COVID-19 infection in Jilin Province. *Virulence* 11(1):482–485. <https://doi.org/10.1080/21505594.2020.1767357>
76. Liu K, Fang Y-Y, Deng Y et al (2020) Clinical characteristics of novel coronavirus cases in tertiary hospitals in Hubei Province. *Chin Med J* 133(9):1025–1031. <https://doi.org/10.1097/CM9.0000000000000744>
77. Tostmann A, Bradley J, Bousema T, et al. (2020) Strong associations and moderate predictive value of early symptoms for SARS-CoV-2 test positivity among healthcare workers, the Netherlands, March 2020. *Eurosurveillance*. 25(16). <https://doi.org/10.2807/1560-7917.EU.2020.25.16.2000508>
78. Yan Y, Yang Y, Wang F, Ren H, Zhang S, Shi X, Yu X, Dong K (2020) Clinical characteristics and outcomes of patients with severe covid-19 with diabetes. *BMJ Open Diabetes Res Care* 8(1):e001343. <https://doi.org/10.1136/bmjdrc-2020-001343>
79. Xu XW, Wu XX, Jiang XG, Xu KJ, Ying LJ, Ma CL, Li SB, Wang HY, Zhang S, Gao HN, Sheng JF, Cai HL, Qiu YQ, Li LJ (2020) Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-CoV-2) outside of Wuhan, China: retrospective case series [published correction appears in BMJ. 2020 Feb 27;368:m792]. *BMJ* 368:m606 Published 2020 Feb 19. <https://doi.org/10.1136/bmj.m606>
80. Lian J, Cai H, Hao S, Jin X, Zhang XL, Zheng L, Jia HY, Hu JH, Zhang SY, Yu GD, Gu JQ, Ye CY, Jin CL, Lu YF, Sheng JF, Yang YD (2020) Comparison of epidemiological and clinical characteristics of COVID-19 patients with and without Wuhan exposure. *J Zhejiang Univ-Sci B* 21(5):369–377. <https://doi.org/10.1631/jzus.B2000112>
81. Gupta N, Agrawal S, Ish P, et al. (2020) Clinical and epidemiologic profile of the initial COVID-19 patients at a tertiary care centre in India. *Monaldi Arch Chest Dis*. 90(1). <https://doi.org/10.4081/monaldi.2020.1294>
82. Zhang X, Cai H, Hu J, Lian J, Gu J, Zhang S, Ye C, Lu Y, Jin C, Yu G, Jia H, Zhang Y, Sheng J, Li L, Yang Y (2020) Epidemiological, clinical characteristics of cases of SARS-CoV-2 infection with abnormal imaging findings. *Int J Infect Dis* 94:81–87. <https://doi.org/10.1016/j.ijid.2020.03.040>
83. Li J, Li S, Cai Y et al (2020) Epidemiological and clinical characteristics of 17 hospitalized patients with 2019 novel coronavirus infections outside Wuhan, China. *Epidemiology*. <https://doi.org/10.1101/2020.02.11.20022053>
84. Hung IF-N, Lung K-C, Tso EY-K, Liu R, Chung TWH, Chu MY, Ng YY, Lo J, Chan J, Tam AR, Shum HP, Chan V, Wu AKL, Sin KM, Leung WS, Law WL, Lung DC, Sin S, Yeung P, Yip CCY, Zhang RR, Fung AYF, Yam EYW, Leung KH, Ip JD, Chu AWH, Chan WM, Ng ACK, Lee R, Fung K, Yeung A, Wu TC, Chan JWM, Yan WW, Chan WM, Chan JFW, Lie AKW, Tsang OTY, Cheng VCC, Que TL, Lau CS, Chan KH, To KK, Yuen KY (2020) Triple combination of interferon beta-1b, lopinavir–ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomised, phase 2 trial. *Lancet* 395(10238):1695–1704. [https://doi.org/10.1016/S0140-6736\(20\)31042-4](https://doi.org/10.1016/S0140-6736(20)31042-4)
85. Wu H, Zhu H, Yuan C, Yao C, Luo W, Shen X, Wang J, Shao J, Xiang Y (2020) Clinical and immune features of hospitalized pediatric patients with coronavirus disease 2019 (COVID-19) in Wuhan, China. *JAMA Netw Open* 3(6):e2010895. <https://doi.org/10.1001/jamanetworkopen.2020.10895>
86. Argenziano MG, Bruce SL, Slater CL et al (2020) Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 369:m1996. Published 2020 May 29. <https://doi.org/10.1136/bmj.m1996>
87. Israelsen SB, Kristiansen KT, Hindsberger B et al (2020) Characteristics of patients with COVID-19 pneumonia at Hvidovre Hospital, March–April 2020. *Dan Med J* 67(6):A05200313 Published 2020 May 15
88. Cummings MJ, Baldwin MR, Abrams D, Jacobson SD, Meyer BJ, Balough EM, Aaron JG, Claassen J, Rabbani LRE, Hastie J, Hochman BR, Salazar-Schicchi J, Yip NH, Brodie D, O'Donnell MR (2020) Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *Lancet* 395(10239):1763–1770. [https://doi.org/10.1016/S0140-6736\(20\)31189-2](https://doi.org/10.1016/S0140-6736(20)31189-2)
89. Kluytmans-van den Bergh MFQ, AGM B, Pas SD et al (2020) Prevalence and clinical presentation of health care workers with symptoms of coronavirus disease 2019 in 2 Dutch hospitals during

- an early phase of the pandemic. *JAMA Netw Open* 3(5):e209673. <https://doi.org/10.1001/jamanetworkopen.2020.9673>
90. Nowak B, Szymański P, Pańkowski I, et al. Clinical characteristics and short-term outcomes of coronavirus disease 2019: retrospective, single-center experience of designated hospital in Poland. *Pol Arch Intern Med.* Published online May 18, 2020. <https://doi.org/10.20452/pamw.15361>
91. Lai X, Wang M, Qin C, Tan L, Ran L, Chen D, Zhang H, Shang K, Xia C, Wang S, Xu S, Wang W (2020) Coronavirus disease 2019 (COVID-2019) infection among health care workers and implications for prevention measures in a tertiary Hospital in Wuhan, China. *JAMA Netw Open* 3(5):e209666. <https://doi.org/10.1001/jamanetworkopen.2020.9666>
92. Wang X, Fang J, Zhu Y, et al. Clinical characteristics of non-critically ill patients with novel coronavirus infection (COVID-19) in a Fangcang Hospital. *Clin Microbiol Infect.* Published online April 2020:S1198743X20301774. <https://doi.org/10.1016/j.cmi.2020.03.032>
93. Liu Z, Jin C, Wu CC, Liang T, Zhao H, Wang Y, Wang Z, Li F, Zhou J, Cai S, Zeng L, Yang J (2020) Association between initial chest CT or clinical features and clinical course in patients with coronavirus disease 2019 pneumonia. *Korean J Radiol* 21(6):736–745. <https://doi.org/10.3348/kjr.2020.0171>
94. Huang Q, Deng X, Li Y, Sun X, Chen Q, Xie M, Liu S, Qu H, Liu S, Wang L, He G, Gong Z (2020) Clinical characteristics and drug therapies in patients with the common-type coronavirus disease 2019 in Hunan, China. *Int J Clin Pharm* 42(3):837–845. <https://doi.org/10.1007/s11096-020-01031-2>
95. Hong KS, Lee KH, Chung JH, Shin KC, Choi EY, Jin HJ, Jang JG, Lee W, Ahn JH (2020) Clinical features and outcomes of 98 patients hospitalized with SARS-CoV-2 infection in Daegu, South Korea: a brief descriptive study. *Yonsei Med J* 61(5):431–437. <https://doi.org/10.3349/ymj.2020.61.5.431>
96. Huang R, Zhu L, Xue L et al (2020) Clinical findings of patients with coronavirus disease 2019 in Jiangsu province, China: a retrospective, multi-center study. Santiago H da C, ed. *PLoS Negl Trop Dis* 14(5):e0008280. <https://doi.org/10.1371/journal.pntd.0008280>
97. Ji M, Yuan L, Shen W et al (2020) Characteristics of disease progress in patients with coronavirus disease 2019 in Wuhan, China. *Epidemiol Infect* 148:e94. Published 2020 May 6. <https://doi.org/10.1017/S0950268820000977>
98. Wang D, Yin Y, Hu C, Liu X, Zhang X, Zhou S, Jian M, Xu H, Prowle J, Hu B, Li Y, Peng Z (2020) Clinical course and outcome of 107 patients infected with the novel coronavirus, SARS-CoV-2, discharged from two hospitals in Wuhan, China. *Crit Care* 24(1):188. <https://doi.org/10.1186/s13054-020-02895-6>
99. Aggarwal S, Garcia-Telles N, Aggarwal G, Lavie C, Lippi G, Henry BM (2020) Clinical features, laboratory characteristics, and outcomes of patients hospitalized with coronavirus disease 2019 (COVID-19): early report from the United States. *Diagnosis* 7(2):91–96. <https://doi.org/10.1515/dx-2020-0046>
100. Zhao X-Y, Xu X-X, Yin H-S, Hu QM, Xiong T, Tang YY, Yang AY, Yu BP, Huang ZP (2020) Clinical characteristics of patients with 2019 coronavirus disease in a non-Wuhan area of Hubei Province, China: a retrospective study. *BMC Infect Dis* 20(1):311. <https://doi.org/10.1186/s12879-020-05010-w>
101. Meng Y, Wu P, Lu W et al (2020) Sex-specific clinical characteristics and prognosis of coronavirus disease-19 infection in Wuhan, China: A retrospective study of 168 severe patients. Perlman S, ed. *PLoS Pathog* 16(4):e1008520. <https://doi.org/10.1371/journal.ppat.1008520>
102. Yao Q, Wang P, Wang X, et al. Retrospective study of risk factors for severe SARS-CoV-2 infections in hospitalized adult patients. *Pol Arch Intern Med.* Published online April 24, 2020. <https://doi.org/10.20452/pamw.15312>
103. Zhu L, Wang J, Huang R, Liu L, Zhao H, Wu C, Zhu C (2020) Clinical characteristics of a case series of children with coronavirus disease 2019. *Pediatr Pulmonol* 55(6):1430–1432. <https://doi.org/10.1002/ppul.24767>
104. Kim ES, Chin BS, Kang CK, Kim NJ, Kang YM, Choi JP, Oh DH, Kim JH, Koh B, Kim SE, Yun NR, Lee JH, Kim JY, Kim Y, Bang JH, Song KH, Kim HB, Chung KH, Oh MD, on behalf of the Korea National Committee for Clinical Management of COVID-19 (2020) Clinical course and outcomes of patients with severe acute respiratory syndrome coronavirus 2 infection: a preliminary report of the first 28 patients from the Korean cohort study on COVID-19. *J Korean Med Sci* 35(13):e142. <https://doi.org/10.3346/jkms.2020.35.e142>
105. Bhatraju PK, Ghassemieh BJ, Nichols M, Kim R, Jerome KR, Nalla AK, Greninger AL, Pipavath S, Wurfel MM, Evans L, Kritek PA, West TE, Luks A, Gerbino A, Dale CR, Goldman JD, O'Mahony S, Mikacenic C (2020) Covid-19 in critically ill patients in the Seattle region — case series. *N Engl J Med* 382(21):2012–2022. <https://doi.org/10.1056/NEJMoa2004500>
106. Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D (2020) Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet Infect Dis* 20(6):689–696. [https://doi.org/10.1016/S1473-3099\(20\)30198-5](https://doi.org/10.1016/S1473-3099(20)30198-5)
107. Chen G, Wu D, Guo W, Cao Y, Huang D, Wang H, Wang T, Zhang X, Chen H, Yu H, Zhang X, Zhang M, Wu S, Song J, Chen T, Han M, Li S, Luo X, Zhao J, Ning Q (2020) Clinical and immunological features of severe and moderate coronavirus disease 2019. *J Clin Invest* 130(5):2620–2629. <https://doi.org/10.1172/JCI137244>
108. Yang W, Cao Q, Qin L, Wang X, Cheng Z, Pan A, Dai J, Sun Q, Zhao F, Qu J, Yan F (2020) Clinical characteristics and imaging manifestations of the 2019 novel coronavirus disease (COVID-19): a multi-center study in Wenzhou city, Zhejiang, China. *J Inf Secur* 80(4):388–393. <https://doi.org/10.1016/j.jinf.2020.02.016>
109. Xu Y-H, Dong J-H, An W-M, Lv XY, Yin XP, Zhang JZ, Dong L, Ma X, Zhang HJ, Gao BL (2020) Clinical and computed tomographic imaging features of novel coronavirus pneumonia caused by SARS-CoV-2. *J Inf Secur* 80(4):394–400. <https://doi.org/10.1016/j.jinf.2020.02.017>
110. Xu X, Yu C, Qu J, Zhang L, Jiang S, Huang D, Chen B, Zhang Z, Guan W, Ling Z, Jiang R, Hu T, Ding Y, Lin L, Gan Q, Luo L, Tang X, Liu J (2020) Imaging and clinical features of patients with 2019 novel coronavirus SARS-CoV-2. *Eur J Nucl Med Mol Imaging* 47(5):1275–1280. <https://doi.org/10.1007/s00259-020-04735-9>
111. Lechien JR, Chiesa-Estomba CM, Place S, et al. Clinical and epidemiological characteristics of 1420 European patients with mild-to-moderate coronavirus disease 2019. *J Intern Med.* Published online June 17, 2020:joim.13089. <https://doi.org/10.1111/joim.13089>
112. CDC COVID-19 response team. Characteristics of health care personnel with COVID-19 - United States, February 12–April 9, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(15):477–481. Published 2020 Apr 17. <https://doi.org/10.15585/mmwr.mm6915e6>
113. Levinson R, Elbaz M, Ben-Ami R, Shasha D, Levinson T, Choshen G, Petrov K, Gadoth A, Paran Y (2020) Time course of anosmia and dysgeusia in patients with mild SARS-CoV-2 infection. *Infect Dis (Lond)* 52(8):600–602. <https://doi.org/10.1080/23744235.2020.1772992>
114. Zhu X, Yuan W, Huang K, et al. Clinical features and short-term outcomes of 114 elderly patients with COVID-19 in Wuhan, China: a single-center, retrospective, observational study. *SSRN Electron J.* Published online 2020. <https://doi.org/10.2139/ssrn.3548774>

115. Wang D, Li R, Wang J, et al. Correlation analysis between disease severity and clinical and biochemical characteristics of 143 cases of COVID-19 in Wuhan, China: a descriptive study. *BMC Infect Dis.* 2020;20(1):519. Published 2020 Jul 16. <https://doi.org/10.1186/s12879-020-05242-w>
116. Chen C, Cao M, Peng L, et al. Coronavirus disease-19 among children outside Wuhan, China. *SSRN Electron J.* Published online 2020. <https://doi.org/10.2139/ssrn.3546071>
117. Mo P, Xing Y, Xiao Y, et al. Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China. *Clin Infect Dis.* Published online March 16, 2020:ciaa270. <https://doi.org/10.1093/cid/ciaa270>
118. Zhang G, Hu C, Luo L, Fang F, Chen Y, Li J, Peng Z, Pan H (2020) Clinical features and short-term outcomes of 221 patients with COVID-19 in Wuhan, China. *J Clin Virol* 127:104364. <https://doi.org/10.1016/j.jcv.2020.104364>
119. Tomlins J, Hamilton F, Gunning S, Sheehy C, Moran E, MacGowan A (2020) Clinical features of 95 sequential hospitalised patients with novel coronavirus 2019 disease (COVID-19), the first UK cohort. *J Infect* Published online April S0163445320302322:e59–e61. <https://doi.org/10.1016/j.jinf.2020.04.020>
120. Zhao Z, Xie J, Yin M, et al. Clinical and laboratory profiles of 75 hospitalized patients with novel coronavirus disease 2019 in Hefei, China. *Infectious Diseases (except HIV/AIDS);* 2020. <https://doi.org/10.1101/2020.03.01.20029785>
121. Huang Y, Yang R, Xu Y, Gong P. Clinical characteristics of 36 non-survivors with COVID-19 in Wuhan, China. *Infectious Diseases (except HIV/AIDS);* 2020. <https://doi.org/10.1101/2020.02.27.20029009>
122. Yan CH, Faraji F, Prajapati DP, Boone CE, DeConde AS (2020) Association of chemosensory dysfunction and COVID-19 in patients presenting with influenza-like symptoms. *Int Forum Allergy Rhinol* 10(7):806–813. <https://doi.org/10.1002/alr.22579>
123. Deng Y, Liu W, Liu K, Fang YY, Shang J, Zhou L, Wang K, Leng F, Wei S, Chen L, Liu HG (2020) Clinical characteristics of fatal and recovered cases of coronavirus disease 2019 in Wuhan, China: a retrospective study. *Chin Med J* 133(11):1261–1267. <https://doi.org/10.1097/CM9.0000000000000824>
124. Chu J, Yang N, Wei Y, Yue H, Zhang F, Zhao J, He L, Sheng G, Chen P, Li G, Wu S, Zhang B, Zhang S, Wang C, Miao X, Li J, Liu W, Zhang H (2020) Clinical characteristics of 54 medical staff with COVID-19: a retrospective study in a single center in Wuhan, China. *J Med Virol* 92(7):807–813. <https://doi.org/10.1002/jmv.25793>
125. Ihle-Hansen H, Berge T, Tveita A, et al. (2020) COVID-19: Symptoms, course of illness and use of clinical scoring systems for the first 42 patients admitted to a Norwegian local hospital. *Covid-19: Symptomer, forlop og bruk av kliniske skåningsverktøy hos de 42 første pasientene innlagt på et norsk lokalsykehus. Tidsskr Nor Laegeforen.* 140(7):<https://doi.org/10.4045/tidsskr.20.0301>. Published 2020 Apr 10. <https://doi.org/10.4045/tidsskr.20.0301>
126. Goyal P, Choi JJ, Pinheiro LC, Schenck EJ, Chen R, Jabri A, Satlin MJ, Campion TR Jr, Nahid M, Ringel JB, Hoffman KL, Alshak MN, Li HA, Wehmeyer GT, Rajan M, Reshetnyak E, Hupert N, Horn EM, Martinez FJ, Gulick RM, Safford MM (2020) Clinical characteristics of Covid-19 in New York City. *N Engl J Med* 382(24):2372–2374. <https://doi.org/10.1056/NEJMcp2010419>
127. Cao J, Tu W-J, Cheng W, et al. Clinical features and short-term outcomes of 102 patients with coronavirus disease 2019 in Wuhan, China. *Clin Infect Dis.* Published online March 13, 2020:ciaa243. <https://doi.org/10.1093/cid/ciaa243>
128. Chang D, Lin M, Wei L, Xie L, Zhu G, dela Cruz CS, Sharma L (2020) Epidemiologic and clinical characteristics of novel coronavirus infections involving 13 patients outside Wuhan, China. *JAMA* 323(11):1092–1093. <https://doi.org/10.1001/jama.2020.1623>
129. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, Li J, Zhao D, Xu D, Gong Q, Liao J, Yang H, Hou W, Zhang Y (2020) Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet* 395(10226):809–815. [https://doi.org/10.1016/S0140-6736\(20\)30360-3](https://doi.org/10.1016/S0140-6736(20)30360-3)
130. Wang L, He W, Yu X, Hu D, Bao M, Liu H, Zhou J, Jiang H (2020) Coronavirus disease 2019 in elderly patients: characteristics and prognostic factors based on 4-week follow-up. *J Inf Secur* 80(6):639–645. <https://doi.org/10.1016/j.jinf.2020.03.019>
131. Spiteri G, Fielding J, Diercke M, et al. (2020) First cases of coronavirus disease 2019 (COVID-19) in the WHO European Region, 24 January to 21 February 2020. *Eurosurveillance*, 25(9). <https://doi.org/10.2807/1560-7917.ES.2020.25.9.2000178>
132. Liu Y, Yang Y, Zhang C, Huang F, Wang F, Yuan J, Wang Z, Li J, Li J, Feng C, Zhang Z, Wang L, Peng L, Chen L, Qin Y, Zhao D, Tan S, Yin L, Xu J, Zhou C, Jiang C, Liu L (2020) Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Sci China Life Sci* 63(3):364–374. <https://doi.org/10.1007/s11427-020-1643-8>
133. Xu T, Chen C, Zhu Z, Cui M, Chen C, Dai H, Xue Y (2020) Clinical features and dynamics of viral load in imported and non-imported patients with COVID-19. *Int J Infect Dis* 94:68–71. <https://doi.org/10.1016/j.ijid.2020.03.022>
134. Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, Cui J, Xu W, Yang Y, Fayad ZA, Jacobi A, Li K, Li S, Shan H (2020) CT imaging features of 2019 novel coronavirus (2019-nCoV). *Radiology* 295(1):202–207. <https://doi.org/10.1148/radiol.2020200230>
135. Shi H, Han X, Jiang N, Cao Y, Alwailid O, Gu J, Fan Y, Zheng C (2020) Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet Infect Dis* 20(4):425–434. [https://doi.org/10.1016/S1473-3099\(20\)30086-4](https://doi.org/10.1016/S1473-3099(20)30086-4)
136. Yang L, Liu J, Zhang R, Li M, Li Z, Zhou X, Hu C, Tian F, Zhou F, Lei Y (2020) Epidemiological and clinical features of 200 hospitalized patients with corona virus disease 2019 outside Wuhan, China: a descriptive study. *J Clin Virol* 129:104475. <https://doi.org/10.1016/j.jcv.2020.104475>
137. Zhao W, Zhong Z, Xie X, Yu Q, Liu J (2020) Relation between chest CT findings and clinical conditions of coronavirus disease (COVID-19) pneumonia: a multicenter study. *Am J Roentgenol* 214(5):1072–1077. <https://doi.org/10.2214/AJR.20.22976>
138. Han Y, Feng Z, Sun L, et al. A comparative-descriptive analysis of clinical characteristics in 2019-coronavirus-infected children and adults. *J Med Virol.* Published online April 17, 2020:jmv.25835. <https://doi.org/10.1002/jmv.25835>
139. Wang Y, Lu X, Li Y, Chen H, Chen T, Su N, Huang F, Zhou J, Zhang B, Yan F, Wang J (2020) Clinical course and outcomes of 344 intensive care patients with COVID-19. *Am J Respir Crit Care Med* 201(11):1430–1434. <https://doi.org/10.1164/rccm.202003-0736LE>
140. Gilani S, Roditi R, Naraghi M (2020) COVID-19 and anosmia in Tehran, Iran. *Med Hypotheses* 141:109757. <https://doi.org/10.1016/j.mehy.2020.109757>
141. Kaye R, Chang CWD, Kazahaya K, Brereton J, Denneny JC (2020) COVID-19 anosmia reporting tool: initial findings. *Otolaryngol Neck Surg* 163(1):132–134. <https://doi.org/10.1177/0194599820922992>
142. Bagheri SHR, Asghari AM, Farhadi M et al (2020) Coincidence of COVID-19 epidemic and olfactory dysfunction outbreak. *Otolaryngology*. <https://doi.org/10.1101/2020.03.23.20041889>

143. Hopkins C, Surda P, Kumar N (2020) Presentation of new onset anosmia during the COVID-19 pandemic. *Rhinol J* 58(3):295–298. <https://doi.org/10.4193/Rhin20.116>
144. Morassi M, Bagatto D, Cobelli M, D'Agostini S, Gigli GL, Bnà C, Vogrig A (2020) Stroke in patients with SARS-CoV-2 infection: case series. *J Neurol* 267(8):2185–2192. <https://doi.org/10.1007/s00415-020-09885-2>
145. Oxley TJ, Mocco J, Majidi S, Kellner CP, Shoirlah H, Singh IP, de Leacy RA, Shigematsu T, Ladner TR, Yaeger KA, Skluit M, Weinberger J, Dangayach NS, Bederson JB, Tuhrim S, Fifi JT (2020) Large-vessel stroke as a presenting feature of Covid-19 in the young. *N Engl J Med* 382(20):e60. <https://doi.org/10.1056/NEJMc2009787>
146. Deliwala S, Abdulhamid S, Abusalihi MF, Al-Qasmi MM, Bachuwa G. Encephalopathy as the sentinel sign of a cortical stroke in a patient infected with coronavirus disease-19 (COVID-19). *Cureus*. Published online May 14, 2020. <https://doi.org/10.7759/cureus.8121>
147. Moshayed P, Ryan TE, Mejia LLP, Nour M, Liebeskind DS (2020) Triage of acute ischemic stroke in confirmed COVID-19: large vessel occlusion associated with coronavirus infection. *Front Neurol* 11:353. <https://doi.org/10.3389/fneur.2020.00353>
148. Muhammad S, Petridis A, Cornelius JF, Hägg D (2020) Letter to editor: severe brain haemorrhage and concomitant COVID-19 infection: a neurovascular complication of COVID-19. *Brain Behav Immun* 87:150–151. <https://doi.org/10.1016/j.bbi.2020.05.015>
149. Al Saiegh F, Ghosh R, Leibold A, et al. (2020) Status of SARS-CoV-2 in cerebrospinal fluid of patients with COVID-19 and stroke [published online ahead of print, 2020 Apr 30]. *J Neurol Neurosurg Psychiatry*. jnnp-2020-323522. <https://doi.org/10.1136/jnnp-2020-323522>
150. Klein DE, Libman R, Kirsch C, Arora R (2020) Cerebral venous thrombosis: a typical presentation of COVID-19 in the young. *J Stroke Cerebrovasc Dis* 29(8):104989. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2020.104989>
151. Garaci F, Di Giuliano F, Picchi E, Da Ros V, Floris R (2020) Venous cerebral thrombosis in COVID-19 patient. *J Neurol Sci* 414:116871. <https://doi.org/10.1016/j.jns.2020.116871>
152. Chougar L, Mathon B, Weiss N, Degos V, Shor N. Atypical deep cerebral vein thrombosis with hemorrhagic venous infarction in a patient positive for COVID-19. *Am J Neuroradiol*. Published online June 18, 2020:ajnr;ajnr.A6642v1. <https://doi.org/10.3174/ajnr.A6642>
153. Cavalcanti DD, Raz E, Shapiro M, et al. Cerebral venous thrombosis associated with COVID-19. *Am J Neuroradiol*. Published online June 18, 2020:ajnr;ajnr.A6644v1. <https://doi.org/10.3174/ajnr.A6644>
154. Hemasian H, Ansari B (2020) First case of Covid-19 presented with cerebral venous thrombosis: a rare and dreaded case. *Rev Neurol (Paris)* 176(6):521–523. <https://doi.org/10.1016/j.neurol.2020.04.013>
155. Sweid A, Hammoud B, Weinberg JH, et al. Letter: thrombotic neurovascular disease in COVID-19 patients. *Neurosurgery*. Published online June 4, 2020:nyaa254. <https://doi.org/10.1093/neurology/nyaa254>
156. Hughes C, Nichols T, Pike M, Subbe C, Elghenai S. Cerebral venous sinus thrombosis as a presentation of COVID-19. *Eur J Case Rep Intern Med*. Published online 2020:1–4. [https://doi.org/10.12890/2020\\_001691](https://doi.org/10.12890/2020_001691)
157. Zhou B, She J, Wang Y, Ma X (2020) A case of coronavirus disease 2019 with concomitant acute cerebral infarction and deep vein thrombosis. *Front Neurol* 11:296. <https://doi.org/10.3389/fneur.2020.00296>
158. Muhammad S, Petridis A, Cornelius JF, Hägg D (2020) Letter to editor: severe brain haemorrhage and concomitant COVID-19 infection: a neurovascular complication of COVID-19. *Brain Behav Immun* 87:150–151. <https://doi.org/10.1016/j.bbi.2020.05.015>
159. Sharifi-Razavi A, Karimi N, Rouhani N (2020) COVID-19 and intracerebral haemorrhage: causative or coincidental? *New Microbes New Infect* 35:100669. <https://doi.org/10.1016/j.nmni.2020.100669>
160. Hanafi R, Roger PA, Perin B, et al. (2020) COVID-19 neurologic complication with CNS vasculitis-like pattern [published online ahead of print, 2020 Jun 18]. *AJNR Am J Neuroradiol*. <https://doi.org/10.3174/ajnr.A6651>
161. Fara MG, Stein LK, Skluit M, Morgello S, Fifi JT, Dhamoon MS. Macrothrombosis and stroke in patients with mild Covid-19 infection. *J Thromb Haemost*. Published online June 25, 2020: jth.14938. <https://doi.org/10.1111/jth.14938>
162. Brüggemann R, Gietema H, Jallah B, ten Cate H, Stehouwer C, Spaetgens B (2020) Arterial and venous thromboembolic disease in a patient with COVID-19: a case report. *Thromb Res* 191:153–155. <https://doi.org/10.1016/j.thromres.2020.04.046>
163. Guillam M, Villacíeros-Alvarez J, Bellido S, Perez-Jorge Peremarch C, Suarez-Vega VM, Aragones-Garcia M, Cabrera-Rojo C, Fernandez-Ferro J (2020) Unusual simultaneous cerebral infarcts in multiple arterial territories in a COVID-19 patient. *Thromb Res* 193:107–109. <https://doi.org/10.1016/j.thromres.2020.06.015>
164. Escalard S, Maier B, Redjem H, Delvoye F, Hébert S, Smajda S, Ciccio G, Desilles JP, Mazighi M, Blanc R, Piotin M. Treatment of acute ischemic stroke due to large vessel occlusion with COVID-19: experience from Paris. *Stroke*. Published online May 29, 2020. <https://doi.org/10.1161/STROKEAHA.120.030574>, 2540, 2543
165. Griffin DO, Jensen A, Khan M, et al. (2020) Arterial thromboembolic complications in COVID-19 in low-risk patients despite prophylaxis. *Br J Haematol*. 190(1). <https://doi.org/10.1111/bjh.16792>
166. Beyrouti R, Adams ME, Benjamin L, et al. Characteristics of ischaemic stroke associated with COVID-19. *J Neurol Neurosurg Psychiatry*. Published online April 30, 2020:jnnp-2020-323586. <https://doi.org/10.1136/jnnp-2020-323586>
167. Nepal P, Batchala PP, Songmen S, Parashar K, Sapire J. An unresponsive COVID-19 patient. *Emerg Radiol*. Published online May 28, 2020. <https://doi.org/10.1007/s10140-020-01799-w>
168. Rudilosso S, Esteller D, Urra X, Chamorro Á (2020) Thalamic perforating artery stroke on computed tomography perfusion in a patient with coronavirus disease 2019. *J Stroke Cerebrovasc Dis* 29(8):104974. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2020.104974>
169. Viguer A, Delamarre L, Duplantier J, Olivot J-M, Bonneville F. Acute ischemic stroke complicating common carotid artery thrombosis during a severe COVID-19 infection. *J Neuroradiol*. Published online May 2020:S0150986120301590. <https://doi.org/10.1016/j.neurad.2020.04.003>
170. Zhang Y, Xiao M, Zhang S, Xia P, Cao W, Jiang W, Chen H, Ding X, Zhao H, Zhang H, Wang C, Zhao J, Sun X, Tian R, Wu W, Wu D, Ma J, Chen Y, Zhang D, Xie J, Yan X, Zhou X, Liu Z, Wang J, du B, Qin Y, Gao P, Qin X, Xu Y, Zhang W, Li T, Zhang F, Zhao Y, Li Y, Zhang S (2020) Coagulopathy and antiphospholipid antibodies in patients with Covid-19. *N Engl J Med* 382(17):e38. <https://doi.org/10.1056/NEJMc2007575>
171. Valderrama EV, Humbert K, Lord A, Frontera J, Yaghi S (2020) Severe acute respiratory syndrome coronavirus 2 infection and ischemic stroke. *Stroke*. 51(7). <https://doi.org/10.1161/STROKEAHA.120.030153>
172. Goldberg MF, Goldberg MF, Cerejo R, Tayal AH (2020) Cerebrovascular disease in COVID-19. *Am J Neuroradiol* 41(7): 1170–1172. <https://doi.org/10.3174/ajnr.A6588>

173. Sharifi-Razavi A, Karimi N, Zarvani A, Cheraghmakan H, Baghbanian SM. Ischemic stroke associated with novel coronavirus 2019: a report of three cases. *Int J Neurosci* Published online June 17, 2020:1–5. <https://doi.org/10.1080/00207454.2020.1782902>
174. Moriguchi T, Harii N, Goto J, Harada D, Sugawara H, Takamino J, Ueno M, Sakata H, Kondo K, Myose N, Nakao A, Takeda M, Haro H, Inoue O, Suzuki-Inoue K, Kubokawa K, Ogihara S, Sasaki T, Kinouchi H, Kojin H, Ito M, Onishi H, Shimizu T, Sasaki Y, Enomoto N, Ishihara H, Furuya S, Yamamoto T, Shimada S (2020) A first case of meningitis/encephalitis associated with SARS-Coronavirus-2. *Int J Infect Dis* 94:55–58. <https://doi.org/10.1016/j.ijid.2020.03.062>
175. Huang YH, Jiang D, Huang JT (2020) SARS-CoV-2 detected in cerebrospinal fluid by PCR in a case of COVID-19 encephalitis. *Brain Behav Immun* 87:149. <https://doi.org/10.1016/j.bbi.2020.05.012>
176. Domingues RB, Mendes-Correa MC, de Moura Leite FBV, Sabino EC, Salarini DZ, Claro I, Santos DW, de Jesus JG, Ferreira NE, Romano CM, Soares CAS First case of SARS-CoV-2 sequencing in cerebrospinal fluid of a patient with suspected demyelinating disease. *J Neurol* Published online June 20, 2020. <https://doi.org/10.1007/s00415-020-09996-w>
177. Farhadian S, Glick LR, Vogels CBF, Thomas J, Chiarella J, Casanovas-Massana A, Zhou J, Odio C, Vijayakumar P, Geng B, Fournier J, Bermejo S, Fauver JR, Alpert T, Wyllie AL, Turcotte C, Steinle M, Paczkowski P, dela Cruz C, Wilen C, Ko AI, MacKay S, Grubaugh ND, Spudich S, Barakat LA (2020) Acute encephalopathy with elevated CSF inflammatory markers as the initial presentation of COVID-19. *BMC Neurol* 20(1):248. <https://doi.org/10.1186/s12883-020-01812-2>
178. Filatov A, Sharma P, Hindi F, Espinosa PS. Neurological complications of coronavirus disease (COVID-19): encephalopathy. *Cureus* Published online March 21, 2020. <https://doi.org/10.7759/cureus.7352>
179. Espinosa PS, Rizvi Z, Sharma P, Hindi F, Filatov A. Neurological complications of coronavirus disease (COVID-19): encephalopathy, MRI brain and cerebrospinal fluid findings: case 2. *Cureus*. Published online May 2, 2020. <https://doi.org/10.7759/cureus.7930>
180. Hepburn M, Mullaguri N, George P, Hantus S, Punia V, Bhimraj A, Newey CR. Acute symptomatic seizures in critically ill patients with COVID-19: is there an association? *Neurocrit Care* Published online May 28, 2020. <https://doi.org/10.1007/s12028-020-01006-1>
181. Pilotto A, Odolini S, Masciocchi S, et al. Steroid-responsive encephalitis in coronavirus disease 2019. *Ann Neurol*. Published online June 9, 2020:ana.25783. <https://doi.org/10.1002/ana.25783>
182. Yin R, Feng W, Wang T, et al. Concomitant neurological symptoms observed in a patient diagnosed with coronavirus disease 2019. *J Med Virol*. Published online June 29, 2020:jmv.25888. <https://doi.org/10.1002/jmv.25888>
183. Afshar H, Yassin Z, Kalantari S, Aloosh O, Lotfi T, Moghaddasi M, Sadeghipour A, Emamikahah M Evolution and resolution of brain involvement associated with SARS-CoV2 infection: a close clinical – paraclinical follow up study of a case. *Mult Scler Relat Disord* 2020;43:102216. <https://doi.org/10.1016/j.msard.2020.102216>
184. Al-olama M, Rashid A, Garozzo D (2020) COVID-19-associated meningoencephalitis complicated with intracranial hemorrhage: a case report. *Acta Neurochir* 162(7):1495–1499. <https://doi.org/10.1007/s00701-020-04402-w>
185. Duong L, Xu P, Liu A (2020) Meningoencephalitis without respiratory failure in a young female patient with COVID-19 infection in Downtown Los Angeles, early April 2020. *Brain Behav Immun* 87:33. <https://doi.org/10.1016/j.bbi.2020.04.024>
186. Wong PF, Craik S, Newman P, Makan A, Srinivasan K, Crawford E, Dev D, Moudgil H, Ahmad N (2020) Lessons of the month 1: a case of rhombencephalitis as a rare complication of acute COVID-19 infection. *Clin Med* 20(3):293–294. <https://doi.org/10.7861/clinmed.2020-0182>
187. Zanin L, Saraceno G, Panciani PP, Renisi G, Signorini L, Migliorati K, Fontanella MM (2020) SARS-CoV-2 can induce brain and spine demyelinating lesions. *Acta Neurochir* 162(7):1491–1494. <https://doi.org/10.1007/s00701-020-04374-x>
188. Vollono C, Rollo E, Romozzi M, Frisullo G, Servidei S, Borghetti A, Calabresi P (2020) Focal status epilepticus as unique clinical feature of COVID-19: a case report. *Seizure* 78:109–112. <https://doi.org/10.1016/j.seizure.2020.04.009>
189. Kadono Y, Nakamura Y, Ogawa Y, Yamamoto S, Kajikawa R, Nakajima Y, Matsumoto M, Kishima H (2020) A case of COVID-19 infection presenting with a seizure following severe brain edema. *Seizure* 80:53–55. <https://doi.org/10.1016/j.seizure.2020.06.015>
190. Sohal S, Mansur M (2020) COVID-19 presenting with seizures. *IDCases* 20:e00782. <https://doi.org/10.1016/j.idcr.2020.e00782>
191. Dixon L, Varley J, Gontsarová A, Mallon D, Tona F, Muir D, Luqmani A, Jenkins IH, Nicholas R, Jones B, Everitt A (2020) COVID-19-related acute necrotizing encephalopathy with brain stem involvement in a patient with aplastic anemia. *Neurol - Neuroimmunol Neuroinflammation* 7(5):e789. <https://doi.org/10.1212/NXI.0000000000000789>
192. Karimi N, Sharifi Razavi A, Rouhani N (2020) Frequent convulsive seizures in an adult patient with COVID-19: a case report. *Iran Red Crescent Med J* 22(3). <https://doi.org/10.5812/ircmj.102828>
193. Somani S, Pati S, Gaston T, Chitlangia A, Agnihotri S (2020) De novo status epilepticus in patients with COVID-19. *Ann Clin Transl Neurol* 7(7):1240–1244. <https://doi.org/10.1002/acn3.51071>
194. McAbee GN, Brosgol Y, Pavlakis S, Agha R, Gaffoor M. Encephalitis associated with COVID-19 infection in an 11-year-old child. *Pediatr Neurol*. Published online April 2020: S0887899420301430. <https://doi.org/10.1016/j.pediatrneurool.2020.04.013>
195. Dugue R, Cay-Martínez KC, Thakur KT, Garcia JA, Chauhan LV, Williams SH, Briese T, Jain K, Foca M, McBrian DK, Bain JM, Lipkin WI, Mishra N (2020) Neurologic manifestations in an infant with COVID-19. *Neurology* 94(24):1100–1102. <https://doi.org/10.1212/WNL.0000000000009653>
196. Kaya Y, Kara S, Akinci C, Kocaman AS (2020) Transient cortical blindness in COVID-19 pneumonia; a PRES-like syndrome: case report. *J Neurol Sci* 413:116858. <https://doi.org/10.1016/j.jns.2020.116858>
197. Poyiadji N, Shahin G, Noujaim D, Stone M, Patel S, Griffith B. COVID-19-associated acute hemorrhagic necrotizing encephalopathy: CT and MRI features. *Radiology*. Published online March 31, 2020:201187. <https://doi.org/10.1148/radiol.2020201187>
198. Radmanesh A, Derman A, Ishida K (2020) COVID-19-associated delayed posthypoxic necrotizing leukoencephalopathy. *J Neurol Sci* 415:116945. <https://doi.org/10.1016/j.jns.2020.116945>
199. Hayashi M, Sahashi Y, Baba Y, Okura H, Shimohata T (2020) COVID-19-associated mild encephalitis/encephalopathy with a reversible splenial lesion. *J Neurol Sci* 415:116941. <https://doi.org/10.1016/j.jns.2020.116941>
200. Parsons T, Banks S, Bae C, Gelber J, Alahmadi H, Tichauer M. COVID-19-associated acute disseminated encephalomyelitis (ADEM). *J Neurol* Published online May 30, 2020. <https://doi.org/10.1007/s00415-020-09951-9>
201. Novi G, Rossi T, Pedemonte E, Saitta L, Rolla C, Roccatagliata L, Inglese M, Farinini D (2020) Acute disseminated

- encephalomyelitis after SARS-CoV-2 infection. *Neurol - Neuroimmunol Neuroinflammation* 7(5):e797. <https://doi.org/10.1212/NXI.00000000000000797>
202. Beach SR, Praschan NC, Hogan C, Dotson S, Merideth F, Kontos N, Fricchione GL, Smith FA (2020) Delirium in COVID-19: a case series and exploration of potential mechanisms for central nervous system involvement. *Gen Hosp Psychiatry* 65:47–53. <https://doi.org/10.1016/j.genhosppsych.2020.05.008>
203. Alkeridy WA, Almaghlouth I, Alrashed R, Alayed K, Binkhamis K, Alsharidi A, Liu-Ambrose T (2020) A unique presentation of delirium in a patient with otherwise asymptomatic COVID -19. *J Am Geriatr Soc* 68(7):1382–1384. <https://doi.org/10.1111/jgs.16536>
204. Neerland BE, Dobloug A, Nore KG, Mikaelson EE, Halsen A, Ahmed MV. COVID-19 in an elderly woman with acute functional decline. Covid-19 påvist hos eldre kvinne med akutt funksjonssvikt. *Tidsskr Nor Laegeforen*. 2020;140(7):<https://doi.org/10.4045/tidsskr.20.0307>. Published 2020 Apr 9.
205. Yin R, Feng W, Wang T, et al. Concomitant neurological symptoms observed in a patient diagnosed with coronavirus disease 2019. *J Med Virol*. Published online June 29, 2020:jmv.25888. <https://doi.org/10.1002/jmv.25888>
206. Zhao H, Shen D, Zhou H, Liu J, Chen S (2020) Guillain-Barré syndrome associated with SARS-CoV-2 infection: causality or coincidence? *Lancet Neurol* 19(5):383–384. [https://doi.org/10.1016/S1474-4422\(20\)30109-5](https://doi.org/10.1016/S1474-4422(20)30109-5)
207. Camdessanche J-P, Morel J, Pozzetto B, Paul S, Tholance Y, Botelho-Nevers E (2020) COVID-19 may induce Guillain–Barré syndrome. *Rev Neurol (Paris)* 176(6):516–518. <https://doi.org/10.1016/j.neurol.2020.04.003>
208. Coen M, Jeanson G, Culebras Almeida LA, Hübers A, Stierlin F, Najjar I, Ongaro M, Moulin K, Makrygianni M, Leemann B, Kronig I, Bertrand J, Reny JL, Schibler M, Serratrice J (2020) Guillain-Barré syndrome as a complication of SARS-CoV-2 infection. *Brain Behav Immun* 87:111–112. <https://doi.org/10.1016/j.bbi.2020.04.074>
209. El Otmani H, El Moutawakil B, Rafai M-A et al (2020) Covid-19 and Guillain-Barré syndrome: more than a coincidence! *Rev Neurol (Paris)* 176(6):518–519. <https://doi.org/10.1016/j.neurol.2020.04.007>
210. Lascano AM, Epiney J, Coen M, et al. SARS-CoV-2 and Guillain-Barré syndrome: AIDP variant with favorable outcome. *Eur J Neurol*. Published online June 2020:ene.14368. <https://doi.org/10.1111/ene.14368>
211. Bigaut K, Mallaret M, Baloglu S, Nemoz B, Morand P, Baicry F, Godon A, Vouleminot P, Kremer L, Chanson JB, de Seze J (2020) Guillain-Barré syndrome related to SARS-CoV-2 infection. *Neurol - Neuroimmunol Neuroinflammation* 7(5):e785. <https://doi.org/10.1212/NXI.0000000000000785>
212. Alberti P, Beretta S, Piatti M, Karantzoulis A, Piatti ML, Santoro P, Viganò M, Giovannelli G, Pirro F, Montisano DA, Appollonio I, Ferrarese C (2020) Guillain-Barré syndrome related to COVID-19 infection. *Neurol - Neuroimmunol Neuroinflammation* 7(4): e741. <https://doi.org/10.1212/NXI.0000000000000741>
213. Padroni M, Mastrangelo V, Asioli GM, Pavolucci L, Abu-Rumeileh S, Piscaglia MG, Querzani P, Callegarini C, Foschi M (2020) Guillain-Barré syndrome following COVID-19: new infection, old complication? *J Neurol* 267(7):1877–1879. <https://doi.org/10.1007/s00415-020-09849-6>
214. Helbok R, Beer R, Löscher W, et al. Guillain-Barré syndrome in a patient with antibodies against SARS-COV-2. *Eur J Neurol*. Published online June 12, 2020:ene.14388. <https://doi.org/10.1111/ene.14388>
215. Scheidl E, Canseco DD, Hadji-Naumov A, Bereznai B (2020) GUILLAIN-BARRÉ syndrome during SARS-CoV-2 pandemic: a case report and review of recent literature. *J Peripher Nerv Syst* 25(2):204–207. <https://doi.org/10.1111/jns.12382>
216. Kilinc D, van de Pasch S, Doets AY, Jacobs BC, van Vliet J, Garssen MPJ. Guillain-Barré syndrome after SARS-CoV-2 infection. *Eur J Neurol*. Published online June 13, 2020:ene.14398. <https://doi.org/10.1111/ene.14398>
217. Sedaghat Z, Karimi N (2020) Guillain Barre syndrome associated with COVID-19 infection: a case report. *J Clin Neurosci* 76:233–235. <https://doi.org/10.1016/j.jocn.2020.04.062>
218. Rana S, Lima AA, Chandra R, Valeriano J, Desai T, Freiberg W, Small G (2020) Novel coronavirus (COVID-19)-associated Guillain-Barré syndrome: case report. *J Clin Neuromuscul Dis* 21(4):240–242. <https://doi.org/10.1097/CND.0000000000000309>
219. Assini A, Benedetti L, Di Maio S, Schirinzi E, Del Sette M (2020) New clinical manifestation of COVID-19 related Guillain-Barré syndrome highly responsive to intravenous immunoglobulins: two Italian cases. *Neurol Sci* 41(7):1657–1658. <https://doi.org/10.1007/s10072-020-04484-5>
220. Virani A, Rabold E, Hanson T, Haag A, Elrufay R, Cheema T, Balaam M, Bhanot N (2020) Guillain-Barré syndrome associated with SARS-CoV-2 infection. *IDCases* 20:e00771. <https://doi.org/10.1016/j.idcr.2020.e00771>
221. Ottaviani D, Bosco F, Tranquilli E, Gapani I, Pedrotti G, Cozzio S, Guerrera GM, Giometto B (2020) Early Guillain-Barré syndrome in coronavirus disease 2019 (COVID-19): a case report from an Italian COVID-hospital. *Neurol Sci* 41(6):1351–1354. <https://doi.org/10.1007/s10072-020-04449-8>
222. Juliao Caamaño DS, Alonso BR (2020) Facial diplegia, a possible atypical variant of Guillain-Barré syndrome as a rare neurological complication of SARS-CoV-2. *J Clin Neurosci* 77:230–232. <https://doi.org/10.1016/j.jocn.2020.05.016>
223. Webb S, Wallace VC, Martin-Lopez D, Yogarajah M (2020) Guillain-Barré syndrome following COVID-19: a newly emerging post-infectious complication. *BMJ Case Rep* 13(6):e236182. <https://doi.org/10.1136/bcr-2020-236182>
224. Arnaud S, Budowski C, Wing N, Tin S, Degos B (2020) Post SARS-CoV-2 Guillain-Barré syndrome. *Clin Neurophysiol* 131(7):1652–1654. <https://doi.org/10.1016/j.clinph.2020.05.003>
225. Sancho-Saldaña A, Lambea-Gil Á, Liesa JLC, Caballo MRB, Garay MH, Celada DR, Serrano-Ponz M (2020) Guillain–Barré syndrome associated with leptomeningeal enhancement following SARS-CoV-2 infection. *Clin Med* 20(4):e93–e94. <https://doi.org/10.7861/clinmed.2020-0213>
226. Reyes-Bueno JA, García-Trujillo L, Urbaneja P, et al. Miller-Fisher syndrome after SARS-CoV-2 infection. *Eur J Neurol*. Published online June 5, 2020:ene.14383. <https://doi.org/10.1111/ene.14383>
227. Toscano G, Palmerini F, Ravaglià S, Ruiz L, Invernizzi P, Cuzzoni MG, Franciotta D, Baldanti F, Daturi R, Postorino P, Cavallini A, Micieli G (2020) Guillain–Barré syndrome associated with SARS-CoV-2. *N Engl J Med* 382(26):2574–2576. <https://doi.org/10.1056/NEJMc2009191>
228. Gutiérrez-Ortiz C, Méndez A, Rodrigo-Rey S, et al. Miller Fisher syndrome and polyneuritis cranialis in COVID-19. *Neurology*. Published online April 17, 2020:<https://doi.org/10.1212/WNL.0000000000009619>.
229. Manganotti P, Pesavento V, Buote Stella A, Bonzi L, Campagnolo E, Bellavita G, Fabris B, Luzzati R. Miller Fisher syndrome diagnosis and treatment in a patient with SARS-CoV-2. *J Neurovirol*. Published online June 11, 2020. <https://doi.org/10.1007/s13365-020-00858-9>
230. Dinkin M, Gao V, Kahan J, et al. COVID-19 presenting with ophthalmoparesis from cranial nerve palsy. *Neurology*. Published online May 1, 2020:<https://doi.org/10.1212/WNL.0000000000009619>.

00000000000009700. <https://doi.org/10.1212/WNL.00000000000009700>
231. Abdelnour L, Eltahir Abdalla M, Babiker S (2020) COVID 19 infection presenting as motor peripheral neuropathy. *J Formos Med Assoc* 119(6):1119–1120. <https://doi.org/10.1016/j.jfma.2020.04.024>
232. Zhang Q, Shan KS, Minalyan A, O'Sullivan C, Nace T. A rare presentation of coronavirus disease 2019 (COVID-19) induced viral myositis with subsequent rhabdomyolysis. *Cureus* Published online May 12, 2020. <https://doi.org/10.7759/cureus.8074>
233. Jin M, Tong Q (2020) Rhabdomyolysis as potential late complication associated with COVID-19. *Emerg Infect Dis* 26(7):1618–1620. <https://doi.org/10.3201/eid2607.200445>
234. Suwanwongse K, Shabarek N. Rhabdomyolysis as a presentation of 2019 novel coronavirus disease. *Cureus* Published online April 6, 2020. <https://doi.org/10.7759/cureus.7561>
235. Munz M, Wessendorf S, Koretsis G, Tewald F, Baegi R, Krämer S, Geissler M, Reinhard M (2020) Acute transverse myelitis after COVID-19 pneumonia. *J Neurol* 267(8):2196–2197. <https://doi.org/10.1007/s00415-020-09934-w>
236. Rábano-Suárez P, Bermejo-Guerrero L, Méndez-Guerrero A, et al. Generalized myoclonus in COVID-19. *Neurology*. Published online May 21, 2020: <https://doi.org/10.1212/WNL.0000000000009829>.
237. Krajewski PK, Szepietowski JC, Maj J (2020) Cutaneous hyperesthesia: a novel manifestation of COVID-19. *Brain Behav Immun* 87:188. <https://doi.org/10.1016/j.bbi.2020.05.064>
238. Aoyagi Y, Ohashi M, Funahashi R, Otaka Y, Saitoh E (2020) Oropharyngeal dysphagia and aspiration pneumonia following coronavirus disease 2019: a case report. *Dysphagia* 35(4):545–548. <https://doi.org/10.1007/s00455-020-10140-z>
239. Acharya S, Diamond M, Anwar S, Glaser A, Tyagi P (2020) Unique case of central retinal artery occlusion secondary to COVID-19 disease. *IDCases* 21:e00867. <https://doi.org/10.1016/j.idcr.2020.e00867>
240. Selvaraj V, Sacchetti D, Finn A, Dapaah-Afriyie K (2020) Acute vision loss in a patient with COVID-19. *R I Med J* (2013) 103(6):37–38 Published 2020 Jun 10
241. Wan Y, Shang J, Graham R, Baric RS, Li F (2020) Receptor recognition by the novel coronavirus from Wuhan: an analysis based on decade-long structural studies of SARS coronavirus. Gallagher T, ed. *J Virol* 94(7):e00127–e00120, /jvi/94/7/JVI.00127–20.atom. <https://doi.org/10.1128/JVI.00127-20>
242. Yang J, Zheng Y, Gou X, Pu K, Chen Z, Guo Q, Ji R, Wang H, Wang Y, Zhou Y (2020) Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis. *Int J Infect Dis* 94:91–95. <https://doi.org/10.1016/j.ijid.2020.03.017>
243. Krauer F, Riesen M, Reveiz L et al (2017) Zika virus infection as a cause of congenital brain abnormalities and Guillain–Barré syndrome: systematic review. von Seidlein L, ed. *PLoS Med* 14(1):e1002203. <https://doi.org/10.1371/journal.pmed.1002203>
244. Hoffmann M, Kleine-Weber H, Schroeder S et al (2020) SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 181(2):271–280.e8. <https://doi.org/10.1016/j.cell.2020.02.052>
245. Hofmann H, Geier M, Marzi A, Krumbiegel M, Peipp M, Fey GH, Gramberg T, Pöhlmann S (2004) Susceptibility to SARS coronavirus S protein-driven infection correlates with expression of angiotensin converting enzyme 2 and infection can be blocked by soluble receptor. *Biochem Biophys Res Commun* 319(4):1216–1221. <https://doi.org/10.1016/j.bbrc.2004.05.114>
246. McGonagle D, Sharif K, O'Regan A, Bridgewood C (2020) The role of cytokines including interleukin-6 in COVID-19 induced pneumonia and macrophage activation syndrome-like disease. *Autoimmun Rev* 19(6):102537. <https://doi.org/10.1016/j.autrev.2020.102537>
247. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, Xie C, Ma K, Shang K, Wang W, Tian DS (2020) Dysregulation of immune response in patients with coronavirus 2019 (COVID-19) in Wuhan, China. *Clin Infect Dis* 71(15):762–768. <https://doi.org/10.1093/cid/ciaa248>
248. Kim J-E, Heo J-H, Kim H, Song SH, Park SS, Park TH, Ahn JY, Kim MK, Choi JP (2017) Neurological complications during treatment of Middle East respiratory syndrome. *J Clin Neurol* 13(3):227. <https://doi.org/10.3988/jcn.2017.13.3.227>
249. Fu L, Wang B, Yuan T, Chen X, Ao Y, Fitzpatrick T, Li P, Zhou Y, Lin YF, Duan Q, Luo G, Fan S, Lu Y, Feng A, Zhan Y, Liang B, Cai W, Zhang L, du X, Li L, Shu Y, Zou H (2020) Clinical characteristics of coronavirus disease 2019 (COVID-19) in China: a systematic review and meta-analysis. *J Inf Secur* 80(6):656–665. <https://doi.org/10.1016/j.jinf.2020.03.041>
250. Zhu J, Zhong Z, Ji P et al (2020) Clinicopathological characteristics of 8697 patients with COVID-19 in China: a meta-analysis [published correction appears in Fam Med Community Health. 2020 May;8(2)]. *Fam Med Community Health* 8(2):e000406. <https://doi.org/10.1136/fmch-2020-000406>
251. Rodriguez-Morales AJ, Cardona-Ospina JA, Gutiérrez-Ocampo E, Villamizar-Peña R, Holguin-Rivera Y, Escalera-Antezana JP, Alvarado-Arnez LE, Bonilla-Aldana DK, Franco-Paredes C, Henao-Martinez AF, Paniz-Mondolfi A, Lagos-Grisales GJ, Ramírez-Vallejo E, Suárez JA, Zambrano LI, Villamil-Gómez WE, Balbin-Ramon GJ, Rabaan AA, Harapan H, Dhama K, Nishiura H, Kataoka H, Ahmad T, Sah R, Latin American Network of Coronavirus Disease 2019-COVID-19 Research (LANCOVID-19). Electronic address: <https://www.lancovid.org> (2020) Clinical, laboratory and imaging features of COVID-19: a systematic review and meta-analysis. *Travel Med Infect Dis* 34:101623. <https://doi.org/10.1016/j.tmaid.2020.101623>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.