

Neuronal Activity Patterns During Hippocampal Network Oscillations In Vitro

Tengis Gloveli, Nancy Kopell, and Tamar Dugladze

Overview

Neurons form transient, functionally specialized assemblies by coordinating their activity within networks. Assembly activity is important for coding and information processing in the brain; oscillations are assumed to entrain and provide temporal structure to this. Recent work from different laboratories has uncovered cell type-specific activity patterns during network oscillations, indicating that the cells may differentially contribute to the generation of oscillation and thereby the coordination of cell assemblies. The purpose of this chapter is to summarize recent findings from these works in in vitro preparations highlighting the importance of different neuronal activity patterns of hippocampal principal cells and different subtypes of interneurons. Special attention will be paid to the role of the firing properties of hippocampal interneurons on the network oscillatory activity at the theta and gamma frequency range. Models based on these ideas are found in “Gamma and Theta Rhythms in Biophysical Models of Hippocampal Circuits” by Kopell et al., this book.

In Vitro Models of Network Oscillations

Hippocampal Population Activity Patterns In Vivo and In Vitro

Hippocampal networks show rhythmic oscillations in various frequency ranges in a behavior-dependent manner (Singer, 1999; Buzsaki and Draguhn, 2004). In the freely moving rat, three types of hippocampal oscillatory activities have been observed (Leung et al., 1982). Theta (5–10 Hz) and gamma (30–100 Hz) frequency rhythms are observed in the rat during exploration and rapid eye movement sleep (Fig. 1A). The frequency range of both rhythms is described differently in different

T. Gloveli (✉)

Institute of Neurophysiology, Charité – Universitätsmedizin Berlin, 10117 Berlin, Germany
e-mail: tengis.gloveli@charite.de

studies. These two rhythms often coexist but can also occur separately (Fig. 1A, for review see Whittington and Traub, 2003). Gamma and theta rhythms also occur throughout the neocortex *in vivo* and have been proposed to constitute a fundamental mechanism underlying cognitive tasks such as feature recognition, associative learning, and content-sensitive and context-sensitive processing of sensory information. In addition, intermittent population bursts, sharp wave-associated field ripples (100–300 Hz), are present in the CA3–CA1–subiculum–entorhinal cortex axis during awake immobility, consummatory behaviors, and slow-wave sleep (Fig. 1B, Vanderwolf, 1969; Buzsaki et al., 1983; Bland, 1986; Chrobak and Buzsaki, 1996; Csicsvari et al., 1999).




Fig. 1 Hippocampal network oscillations *in vivo* **A**, **B** and *in vitro* **C–E**. **A** Theta-related and gamma-related modulation of the field in the dentate gyrus (hilar region) during exploratory waking. **B** Sharp wave-associated field ripples in CA1 area during slow-wave sleep. *Upper traces*, wide band recording, *lower traces*, band bath (40–150 Hz, **A**; 150–250 Hz, **B**) filtered gamma and ripple activity. **C** Metabotropic glutamate receptor activation under conditions of reduced AMPA receptor activation generates in CA1 area theta population activity. **D** Kainate receptor activation induces network oscillations at the gamma frequency range in CA3 area. **E** Spontaneously occurring sharp wave-associated ripple oscillation in CA1 area *in vitro*. *Upper trace*, wide band recording, *lower trace*, ripple band-pass (140–320 Hz) filtered activity. Panels are adapted from **A**, Bragin et al. (1995); **B**, Csicsvari et al. (1999); **C**, Gillies et al. (2002); **D**, Gloveli et al. (2005b); **E**, Both et al. (2008)

Various *in vitro* models have been developed to gain insight into the cellular and synaptic mechanisms of theta, gamma, and ripple oscillations (Fig. 1C–E). *In vitro* models of network oscillations, such as the carbachol (Fisahn et al., 1998; Buhl et al., 1998), the kainate (Buhl et al., 1998), the metabotropic glutamate receptor activation (Gillies et al., 2002), and the tetanically induced (Whittington et al., 1997) gamma activity models, reproduce salient features of oscillatory activity in slice preparations maintained in “interface” slice chamber. To determine activity

pattern of individual neurons, sharp microelectrode or blind whole-cell patch-clamp recordings have been obtained from principal cells or putative interneurons. In addition, an in vivo model, the juxtacellular recording technique, was developed to conjointly record action potential series from single neurons and the extracellular field potential during different forms of network activity in anesthetized animals (Pinault, 1996; Klausberger et al., 2003, 2004). These in vitro and in vivo methods have some clear advantages in studying network activity. However, the sparse distribution of interneurons makes them unlikely targets for these blind approaches. Therefore, these investigations are very inefficient in mapping neuronal activity patterns. Whole-cell patch-clamp recordings using infrared differential contrast videomicroscopy (Dodt and Zieglängsberger, 1994) have greatly facilitated selection and recordings from interneuron. However, this approach has been hampered by the difficulty of generating population activity in the submerged-type slice chambers. Recently, technical modification of the pharmacological paradigms, brief pressure ejection of kainate (Gloveli et al., 2005a, b) or bath application of carbachol (Hájos et al., 2004), permitted the reproduction of the network oscillatory activity in submerged slices. Using these approaches, it was possible to record from visually identified pyramidal cells and interneurons during gamma and theta frequency network oscillation in vitro.

Cell Types Involved in Rhythms

Morphological properties discriminate hippocampal pyramidal cells (PCs) from inhibitory interneurons. In addition, further distinctions exist within both PCs and interneurons. It is reasonable to postulate that hippocampal neurons with different structural features are also likely to have different functions in the network.

Pyramidal cells. Despite the morphological similarities (pyramid-shaped somata, apical and basal dendritic trees), PCs in CA1 and CA3 areas display some important differences such as the existence of excitatory recurrent collaterals. The latter is considered to be the hallmark of the CA3 but not the CA1 area. The pyramidal cells of the CA3 area themselves are not homogeneous. Whereas most axon collaterals of the CA3a and CA3b neurons give rise to extensive recurrent collaterals that are confined to the CA3 region, pyramidal cells in CA3c subregion are mostly projection cells, with most of their axon collaterals terminating in the CA1 region (Li et al., 1994; Wittner et al., 2007). It was hypothesized (Csicsvari et al., 2003) that intrahippocampal gamma oscillations emerge in the recurrent collateral-rich CA3a,b subregions; their activity recruits CA3c subregion, which, in turn, entrains CA1 cells.

Interneuron types. In contrast to glutamatergic principal cells, GABAergic interneurons of the hippocampus exhibit substantial diversity. In the CA1 area, for instance, at least 21 classes of interneurons were described (for review see Klausberger and Somogyi, 2008 [see “Morphology of Hippocampal Neurons” by Vida, this book]). In contrast to principal cells, the vast majority of interneurons have

locally restricted axons and lack spines. Interneurons can be broadly classified into several classes on the basis of different criteria, such as action potential firing properties, somato-dendritic architecture and axonal ramification pattern, neurochemical content, voltage- and ligand-gated conductances as well as plastic changes in excitatory synaptic transmission (for reviews see Freund and Buzsáki, 1996; McBain and Fisahn, 2001; Whittington and Traub, 2003). Functionally, at least three main GABAergic cell types coexist in hippocampal networks: perisomatic inhibitory neurons, dendritic inhibitory interneurons, and GABAergic cells specifically innervating other inhibitory interneurons (Miles et al., 1996). The most striking morpho-functional dichotomy in the population of cortical interneurons is the targeting of the dendritic versus the perisomatic domain of principal cells. Dendritic inhibition is likely to control the efficacy and plasticity of excitatory synaptic inputs of principal cells, whereas perisomatic inhibition is ideally suited to control output, synchronizing the action potential firing of large groups of principal cells (Freund and Buzsáki, 1996). Further distinctions exist within the same class of interneurons. Thus, different types of *perisomatic targeting* parvalbumin (PV)-expressing interneurons innervate distinct subcellular domains of principal cells. Axo-axonic cells (AACs) innervate exclusively the axon initial segment of PCs; in contrast, basket cells (BCs) innervate the somata and proximal apical dendrites. In addition, two distinct populations of basket cells – PV-expressing and cholecystokinin (CCK)-expressing interneurons – could be defined on the basis of their neurochemical content (see “Morphology of Hippocampal Neurons” by Vida, this book). Dendrite-targeting interneurons could also be subdivided into the distal (such as oriens lacunosum-moleculare, O-LM, and radiatum lacunosum-moleculare – R-LM cells) and proximal (such as trilaminar, bistratified, and radiatum cells) dendrite-targeting cells. Interneurons belonging to distinct classes defined by their axonal target domain on the pyramidal cell have clearly different intrinsic, synaptic, and firing properties.

As an example, Fig. 2 illustrates the morphology and the physiological properties of two types of interneurons: non-fast spiking distal dendrite-targeting O-LM cells, which present one of the best studied interneuron classes in the hippocampus, and fast-spiking proximal dendrite-targeting trilaminar cells. These cells differ in their morphology, neurochemical marker contents, and intrinsic membrane properties (see Fig. 2A, B). Clear differences were also detected in spontaneous EPSC properties between these two subtypes of interneurons – with slower kinetics in O-LM than those in trilaminar interneurons (Fig. 2B). Furthermore, while the excitatory input displayed a late-persistent firing in O-LM cells, fast-spiking trilaminar interneurons displayed an onset-transient firing in response to stimulation of CA1 axons in the alveus (Fig. 2B; Pouille and Scanziani, 2004).

These differences in morphological and electrophysiological properties of interneurons indicate that they are likely to have specific roles in the network. In fact, analysis of their spike timing during the oscillations suggests that a division of labor exist among interneuron subtypes involved in hippocampal network oscillations (see Sections “Firing Patterns in Gamma Oscillations” and “Firing Patterns in Theta Oscillations”).




Fig. 2 Properties of distal (O-LM) and proximal (trilaminar) dendrite-targeting interneurons. **A** Morphology of O-LM and trilaminar cells. Somata and dendrites are drawn in *red*, axons are in *black*. The somata of O-LM cells are located in stratum oriens and have mainly horizontally running dendrites. The main axon of these cells crosses strata pyramidale and radiatum and branches in stratum lacunosum-moleculare. O-LM cells innervate the distal dendrites of pyramidal cells which are co-aligned with the entorhinal input (Sik et al., 1995; Maccaferri et al., 2000). **A** inset, O-LM cells are immunopositive for the metabotropic glutamate receptor (mGluR1 α) and the neuropeptide somatostatin (SOM, Tukker et al., 2007) and express low levels of calcium binding protein PV (Maccaferri et al., 2000; Klausberger et al., 2003). The trilaminar cells have similar horizontally distributed dendrites in stratum oriens, but are clearly different from O-LM cells in respect of axonal arborization (Sik et al., 1995). Sub – subiculum; str. or. – stratum oriens; str. pyr. – stratum pyramidale; str. rad. – stratum radiatum; str. l.-m. – stratum lacunosum-moleculare. **B** *Intrinsic*, intrinsic membrane (i) and firing properties (ii) of O-LM and trilaminar cells during hyperpolarizing and depolarizing current injection. O-LM cells demonstrate clear “sag” potential and non-fast spiking pattern in marked contrast to trilaminar cells showing no “sag” and fast-spiking character upon hyperpolarizing and depolarizing pulses. **B** *Synaptic* (i), spontaneous EPSC (sEPSC) in O-LM and trilaminar interneurons. Forty individual traces are *black* and superimposed averaged currents are *red*. **B** *Synaptic* (ii), cell-attached responses from an O-LM and trilaminar cells. Current deflections indicate action potential firing in response to the stimulation of CA1 axons in the alveus. Panels **A** and **B**, *Intrinsic* (i, ii) and *Synaptic* (i) are from Gloveli et al. (2005a); panel **B** *Synaptic* (ii) is from Pouille and Scanziani (2004); inset **A**, (O-LM) is from Tukker et al. (2007)

Gamma Oscillations

Two forms of local network gamma frequency oscillations can be induced in vitro in hippocampal slices (Table 1). Transient forms of gamma frequency oscillations (lasting for a few seconds or minutes) can be evoked in vitro by titanic stimulation (Whitington et al., 1997) or through pressure ejection of glutamate

Table 1 Properties of in vitro models of theta and gamma oscillations

Oscillation type	Activated by	Blocked by	Region	Mean freq. (rec. temp.)	Main references
Persistent θ	mGlutR	NMDAR, GABA _A R	CA1	7 Hz (35°C)	Gillies et al. (2002)
	mAChR	NMDAR, GABA _A R	CA1	9 Hz (35°C)	Gillies et al. (2002)
	mGlutR, mAChR	AMPA	CA3	8 Hz (33°C)	Cobb et al. (2000)
	KAR, long. slice	N.T.	CA3	8 Hz (29°C)	Konopacki et al. (1992) Gloveli et al. (2005b)
Transient θ	PuffKA, long. slice	GABA _A R	CA3	8 Hz (29°C)	Gloveli et al. (2005b)
Persistent γ	KAR	GABA _A R	CA3	35 Hz (35°C) 37 Hz (34°C)	Fisahn et al. (2004) Gloveli et al. (2005b)
	mAChR	AMPA, GABA _A R	CA3	32 Hz (30°C)	Pálhalmi et al. (2004)
			CA3	39 Hz (34°C)	Fisahn et al. (1998) Hájos et al. (2004)
	mGluR	AMPA, GABA _A R	CA3	41 Hz (30°C)	Pálhalmi et al. (2004)
Transient γ	Puff Glut.	GABA _A R	CA1/DG	42/64 Hz (36°C)	Pöschel et al. (2002)
	Puff KA	GABA _A R	CA3	33 Hz (29°C)	Gloveli et al. (2005a)
	Puff K ⁺	GABA _A R	DG	67 Hz (34°C)	Towers et al. (2002)
		AMPA, GABA _A R	CA1	63 Hz (34°C)	LeBeau et al. (2002)

Abbreviations: R, receptor; m, metabotropic; KAR, kainate R; long., longitudinal; Puff, pressure ejection; N.T., not tested.

(Pöschel et al., 2002), high molarities of kainate (Gloveli et al., 2005a, b) or potassium (LeBeau et al., 2002; Towers et al., 2002) (Table 1).

Another model of gamma frequency oscillations is known as “persistent gamma” (lasting for hours). This kind of oscillation can be induced in the hippocampal CA3 area in vitro by bath application of agonists of muscarinic acetylcholine (mAChR) (Fisahn et al., 1998; Fellous and Sejnowski, 2000; Shimono et al., 2000; Fisahn et al., 2002) and kainate receptors (KAR) (Fisahn et al., 2004; Gloveli et al., 2005a, b) (Table 1). mAChR agonist (carbachol)-induced and kainate-induced fast network oscillations provide a useful model to explore the mechanisms underlying physiological gamma frequency oscillations for the following reasons. The hippocampus receives a dense cholinergic projection from the medial septum/diagonal band of Broca, which plays an important role in the generation of hippocampal network activity (Leung, 1985). In addition, kainate receptors are expressed by both principal cells and interneurons of the hippocampus (Cossart et al., 1998; Frerking et al., 1998; for review see Lerma, 2003). Moreover, these oscillations in vitro share many of the features of intrahippocampal gamma oscillations in vivo, including the

firing of pyramidal neurons at low frequencies (<5 Hz) phase-locked to the oscillation, and the generation of oscillations in CA3 then propagating to CA1 (Fisahn et al., 1998; Gloveli et al., 2005a). Finally, both in vivo and in vitro cholinergically induced oscillations have similar current source density profiles, and the gamma phase relationship between pyramidal cells and perisomatic innervating interneurons is comparable (Csicsvari et al., 2003; Mann et al., 2005; Oren et al., 2006).

Both persistent and transient gamma oscillations can be evoked in different hippocampal areas, including CA3, CA1, and DG (Table 1, Towers et al., 2002; Pöschel et al., 2002; Gloveli et al., 2005a). However, there are regional differences in frequency and power of the oscillations, suggesting the existence of different rhythm-generating networks in the hippocampus. In line with this suggestion, both persistent and transient forms of kainate-induced gamma oscillations demonstrate faster gamma frequency oscillations in isolated CA1 area than those in CA3 area (N. Maziashvili and T. Gloveli, unpublished observation, Middleton et al., 2008). However, gamma oscillations in the same area (CA3 area) induced by different pharmacological drugs (carbachol and DHPG) also show significant differences in their properties (the peak frequencies, maximal power, and spectral width, Table 1, Pálhalmi et al., 2004), suggesting involvement of different network mechanisms, such as the recruitment of distinct types of interneurons. In addition, the gamma oscillations evoked under different conditions differ in their dependence on excitation and inhibition (Table 1). Thus, one form of transient oscillations, “Interneuronal network gamma” (ING) (Whittington et al., 1995), are based on mutual inhibition between the interneurons (for computational models see Wang and Buzsáki, 1996; White et al., 1998; Vida et al., 2006) whereas “pyramidal-interneuronal network gamma” (PING) (Whittington et al., 1997) is based on reciprocal interneuron–pyramidal cell interaction (see “Gamma and Theta Rhythms in Biophysical Models of Hippocampal Circuits” by Kopell et al., this book). It seems likely that all of these forms are relevant in vivo, possibly reflecting region and state dependence of mechanisms underlying hippocampal gamma oscillations.

Firing Patterns in Gamma Oscillations

A key requirement for the generation of network oscillations is rhythmic and synchronized activity of large sets of neurons. An important step in understanding the role of hippocampal neurons in network oscillations is to examine their spike patterns during these oscillations.

Principal cells. Analysis of firing properties of electrophysiologically and morphologically identified pyramidal cells in CA3 area has recently been performed in vitro for KAR (Gloveli et al., 2005a, b) and mAChR (Fisahn et al., 1998; Hájos et al., 2004) agonist-induced gamma frequency oscillations. Both KAR and mAChR activation (by kainate and carbachol, respectively) revealed low-frequency, <5 Hz, firing of pyramidal cells (Table 2, Fisahn et al., 1998; Hájos et al., 2004; Gloveli et al., 2005a). These results are in agreement with in vivo observations demonstrating similar low-frequency firing of PCs (Csicsvari et al., 2003). Moreover,

PC firing is phase-locked to the field oscillations (Table 2). In carbachol-induced gamma oscillations, PCs fired action potentials around the negative peak of the field recorded in the pyramidal cell layer (Fig. 3a, d, Hájos et al., 2004). Both in vivo and in vitro observations suggest that during gamma frequency oscillations, PCs of CA3 area drive local interneurons in a feedback manner (Fisahn et al., 1998; Csicsvari et al., 2003; Pálhalmi et al., 2004; Hájos et al., 2004). If PC–interneuron interactions generate gamma oscillations, the firing of PCs should precede interneuron discharge so that PC can recruit interneuron activity in the next gamma cycle (Oren et al., 2006). Consistent with this suggestion, interneuron responses were indeed preceded by PC firing (Fig. 3d, Hájos et al., 2004).

Table 2 Firing properties of some hippocampal neurons during gamma frequency oscillations in vitro

Neuron type	Activated by	Mean firing frequency (Hz)	Spikes/gamma cycle	Angle of spikes relative to the field	Main references
Pyramidal	KAR	3.5 ± 0.6	0.18 ± 0.05	N.T.	Gloveli et al. (2005a)
	mAChR	2.82 ± 0.7	0.09 ± 0.02	58.1 ± 5.3^0	Hájos et al. (2004) Fisahn et al. (1998)
O-LM	KAR	8.3 ± 2.1	0.26 ± 0.04	N.T.	Gloveli et al. (2005a)
	mAChR	12.9 ± 1.8	0.4 ± 0.07	88.1 ± 6.1^0	Hájos et al. (2004)
Trilaminar	KAR	32.1 ± 2.8	1.82 ± 0.07	N.T.	Gloveli et al. (2005a)
	mAChR	18.2 ± 2.7	0.6 ± 0.09	96.8 ± 2.2^0	Hájos et al. (2004)
Bistratified	KAR	35.0 ± 2.5	1.04 ± 0.08	N.T.	Gloveli et al. (2005a)
Basket	KAR	33.6 ± 2.6	1.28 ± 0.06	N.T.	Gloveli et al. (2005a)
	mAChR	18.1 ± 2.7	0.62 ± 0.09	93 ± 2.1^0	Hájos et al. (2004)
R-LM	mAChR	13.2 ± 3.9	N.T.		Hájos et al. (2004)
Radiatum	mAChR	2.3 ± 0.6	0.07 ± 0.02	128.4 ± 12.4^0	Hájos et al. (2004)

Abbreviations: O-LM, oriens lacunosum-moleculare; R-LM, radiatum lacunosum-moleculare; N.T., not tested.

During in vitro gamma frequency oscillations induced by kainate, the interneurons receive a high-frequency barrage of compound EPSPs, modulated at gamma frequency, which are temporally correlated with extracellular population activity (Gloveli et al., 2005a). Since the slice is deafferented, it is likely that the action potential-dependent excitatory events are mediated by local excitatory input from neighboring pyramidal neurons. Given the relatively low PC somatic spike rate with

respect to the frequency of EPSPs invading interneurons, the question remains as to how PCs generate these rhythmic burst of events and reliably discharge interneurons. Interneurons may receive a rhythmic barrage of gamma frequency EPSPs for the following reasons. First, in the active network, multiple pyramidal cells are likely to fire on any given oscillatory cycle. Due to the convergence of numerous pyramidal cell axons onto a single postsynaptic interneuron it follows that each interneuron is also likely to receive multiple unitary excitatory inputs on each successive gamma wave. Second, there are suggestions that activity in pyramidal cell axons may orthodromically excite interneurons, without pyramidal cell somata necessarily firing (Traub et al., 2003). Computational models of carbachol (Traub et al., 2000) and kainate-induced gamma oscillations (Fisahn et al., 2004) emphasize the importance of ectopic axonal action potentials for the generation of hippocampal gamma oscillations. The coexistence of phasic, high-frequency oscillations in principal cell axon populations and field potential gamma frequency oscillations was demonstrated in kainate model (Traub et al., 2003).

Interneuron types. During gamma frequency oscillation in vivo and in vitro, the different classes of interneurons fire action potentials at different times and inhibit distinct subcellular domains of PCs (Fig. 2–4). During pharmacologically induced gamma frequency oscillations in vitro, perisomatic-targeting *PV-expressing basket* cells generate a predominantly gamma frequency output (Fig. 3B, Gloveli et al., 2005a; Hájos et al., 2004). Moreover, the firing of perisomatic basket cells is tightly coupled to the oscillation (Fig. 3B, D). The anatomical and physiological properties make these neurons ideally suited for generating local gamma rhythms. In contrast, spiking of other *PV-expressing perisomatic-targeting interneurons, axo-axonic* cells, was found to be only moderately coupled to the field gamma in anesthetized animals (Tukker et al., 2007).

There are no in vitro data available on the activity of *CCK-expressing basket* cells. In vivo results indicate that, in contrast to *PV-expressing BCs*, these interneurons fire earlier than pyramidal cells and out of phase with *PV-expressing interneurons* during the gamma oscillations in anesthetized animals (Tukker et al., 2007). Therefore, *CCK-expressing basket* cells are likely to interfere with gamma synchronicity (Freund and Katona, 2007; Galarreta et al., 2008).

While *PV-expressing perisomatic inhibitory interneurons* are thought to play a major role in gamma oscillations (Hájos et al., 2004; Gloveli et al., 2005a), other classes of fast-spiking interneurons, such as *bistratified* and *trilaminar* cells, may also be important for this rhythm (Gloveli et al., 2005a). *Bistratified* cells were so named because the axonal arbor is found in two strata: oriens and radiatum (Buhl et al., 1994). In addition to pyramidal cells they also innervate interneurons including basket cells (Halasy et al., 1996). During the gamma oscillations in vitro *bistratified* cells discharge at high frequency, phase-locked to the field gamma (Gloveli et al., 2005a; Hájos et al., 2004; Tukker et al., 2007). Therefore, they are also likely to be involved in the generation of the gamma oscillatory activity. Interestingly, the most prominent interneuronal output seen during pharmacologically induced gamma oscillations in vitro was associated with *trilaminar* interneurons (Gloveli et al., 2005a; Hájos et al., 2004). These fast-spiking cells project to three layers




Fig. 3 Morphological and firing properties of hippocampal neurons during pharmacologically induced gamma oscillations. Reconstructions of representative biocytin-filled pyramidal (**Ai**), PV-positive basket (**Bi**), and trilaminar (**Ci**) cells. The soma and dendrites are drawn in red, whereas the axons are in black. CA3, CA3 area; str. or., stratum oriens; str. pyr., stratum pyramidale; str. rad., stratum radiatum; str. l.-m., stratum lacunosum-moleculare. During kainate-induced field oscillatory activity (**Aii**; **Bii**) pyramidal cells fire sporadically (**Aiii**), whereas basket cells discharged with single spikes interrupted by irregularly occurring doublets of action potentials, phase-locked to the field gamma activity (**Biii**). Trilaminar cells produced spike doublets (**Ciii**) on every gamma cycle (**Cii**). (**D, left**) Time sequence of firing of different neuron types during carbachol-induced oscillatory cycle (top trace). Pyramidal cells fired at the negative peak of the oscillation followed by the interneurons. Gaussian functions were fitted to the spike time distribution for each type of neuron, and the average mean and SD were used to represent each cell class as a Gaussian function. (**D, right**) Schematic diagram of the connectivity among phase-coupled neuron types in the CA3 hippocampal circuitry taking part in the gamma oscillation. Panel **Aii–C**, adapted from Gloveli et al. (2005a); **D**, adapted from Hájos et al. (2004)

of CA3 area, the strata oriens, pyramidale, and radiatum. Additionally, axon collaterals of trilaminar cells were seen projecting to area CA1 and into the subiculum, and possibly to other brain areas as well (Somogyi and Klausberger, 2005). These cells generated highly regular, short latency spike doublets (Fig. 3C, Gloveli et al., 2005a). Their axonal arborization indicates that these interneurons innervate somatic and dendritic compartments of pyramidal cells, locally as well as in distant regions. Thus, via these cells, gamma rhythms generated locally in area CA3 could be efficiently transmitted to distal sites “downstream” in the hippocampal processing pathway.

Interneurons located in the stratum radiatum (with both the dendrites and axonal arborization localized in the stratum radiatum) have the lowest firing rate among all dendrite-targeting interneurons with weak coupling to the gamma oscillations in vitro (Table 2, Hájos et al., 2004). Although *R-LM* cells (with dendritic tree in stratum radiatum and axon restricted to stratum lacunosum-moleculare) fire at higher frequency than other radiatum cells, they also do not show significant phase-related firing (Hájos et al., 2004).

Theta Oscillations

A prominent network pattern in the hippocampus of all mammals studied to date, including humans (Arnolds et al., 1980; Tesche and Karhu, 2000), is a slow oscillation in the theta frequency band (5–10 Hz). Theta oscillations are most consistently present during various types of locomotor activities (Vanderwolf, 1969) and rapid eye movement (REM) sleep (Jouvet, 1969). In general, theta waves are absent in the immobile animal (Bland, 1986; for review see Buzsáki, 2002). To explain the generation of these oscillations, various external pacemakers have been proposed (for review see Buzsáki, 2002). One classical hypothesis is that cholinergic excitation from the septum and the diagonal band of Broca activates inhibitory interneurons, which in turn induce rhythmic IPSPs on the soma of pyramidal cells (Petsche et al., 1962). Alternatively, the entorhinal cortex may entrain hippocampal areas at theta frequency. In rodents, hippocampal theta activity has maximal power in the CA1 region and the synaptic currents underlying these oscillations are mainly generated by the entorhinal input (for review see Buzsáki, 2002). However, recent in vitro experimental data and computational analysis indicate that theta activity can be generated intrinsically in the CA1 (Gillies et al., 2002; Rotstein et al., 2005) and CA3 (Gloveli et al., 2005b) areas of the hippocampus. In fact, Cobb et al. (1995) demonstrated that individual GABAergic interneurons can effectively phase subthreshold membrane potential oscillations and spontaneous firing in pyramidal cells at theta frequencies. Alternating inhibition and post-inhibitory “rebound” activation underlies the entrainment of pyramidal cells (Cobb et al., 1995). Intrinsic GABAergic mechanisms are thus sufficient to generate theta activity in cortical networks.

Various in vitro models of the theta oscillatory activity have been developed based on either mAChR (Konopacki et al., 1992; Fisahn et al., 1998), metabotropic glutamatergic-receptor (mGluR) (Gillies et al., 2002) or kainate receptor activation (Gloveli et al., 2005a, b). Coactivation of mGluRs and metabotropic cholinergic receptors has also been reported to generate robust theta frequency oscillations in the hippocampus in vitro (Cobb et al., 2000).

Metabotropic GluR activation generates prominent, inhibition-based, atropine-resistant theta population oscillations under conditions of reduced AMPA receptor activation in the hippocampal CA1 area (Gillies et al., 2002). This field oscillation was independent of muscarinic cholinergic receptor drive, but strongly dependent on NMDA receptor and GABA_A receptor activity (Table 1, Gillies et al., 2002). The mechanism of generation of theta frequency population activity in this model appeared to involve intrinsic theta frequency membrane potential oscillations in a subset of stratum oriens interneurons. The blockade of AMPA receptors was a critical requirement of the experimental conditions needed to see this population theta activity. Many of the properties of theta frequency oscillations in this reduced model match those seen in area CA1 in vivo (Gillies et al., 2002). In particular, the resulting population theta rhythm resembled atropine-resistant theta oscillations recorded in vivo (Buzsáki et al., 1986) and may be generated by a subset of stratum oriens interneurons displaying intrinsic membrane potential oscillations at theta frequency

(Gillies et al., 2002). In addition, the coherent theta oscillations may come from the interaction of other GABAergic interneurons with the O-LM cells (see “Gamma and Theta Rhythms in Biophysical Models of Hippocampal Circuits” by Kopell et al., this book).


In the CA3 area, theta oscillations can be induced by application of kainate. A necessary prerequisite to ensure precisely synchronized theta activity was specific orientation of the slices: theta frequency population activity was detected predominantly in longitudinal hippocampal slice preparation (Gloveli et al., 2005b). These data demonstrate that theta activity can be generated intrinsically both in the CA1 and in the CA3 areas of the hippocampus. While there are several differences between these models a common feature is their dependence on GABAergic inhibition (Table 1).

Firing Patterns in Theta Oscillations

Principal cells. In a model of atropine-resistant theta oscillations following mGluR activation, with AMPA receptor activation blocked, pyramidal cell somatic firing was seen in only few cells recorded, but could be elicited with injection of tonic depolarizing current (0.1–0.2 nA). In these conditions, pyramidal cells fired one spike per field theta (~ 7 Hz) cycle during the trough of the field oscillation (Gillies et al., 2002). Consistent with this finding, pyramidal neurons in the CA1 region showed subthreshold resonance and firing preference at theta frequencies (range 2–7 Hz) (Pike et al., 2000).

Interneuron types. Ample evidence supports the critical involvement of hippocampal interneurons in theta oscillations. The best documented is the involvement of stratum oriens distal dendrite-targeting *O-LM* interneurons (Fig. 2, Fig. 4A) in generation of theta rhythm. This cell type was found to participate in hippocampal theta activity both in vivo (Buzsáki, 2002; Klausberger et al., 2003) and in vitro (Pike et al., 2000; Gillies et al., 2002; Hájos et al., 2004; Gloveli et al., 2005a, b). In particular, involvement of O-LM cells was investigated in vitro in kainate and mAChR-mediated network oscillatory activity. In kainate-induced oscillations O-LM cells fired at the theta frequency range during both theta and gamma population activity (Fig. 4B, Gloveli et al., 2005b). O-LM cells show prominent membrane potential oscillations in the theta frequency range (Maccaferri and McBain, 1996). In contrast, hippocampal fast-spiking cells preferentially resonate in the gamma range (Pike et al., 2000). Furthermore, O-LM cells have longer membrane time constants than the gamma-preferring interneurons and a considerably longer afterhyperpolarization (AHP). Changes in AHP profiles in interneurons have been shown to have dramatic effects on firing patterns (e.g., see Savić et al., 2001). Thus O-LM cells and gamma-preferring interneurons discharge at different frequencies and participate preferentially in theta or gamma activity, respectively (Gloveli et al., 2005a). The theta frequency discharge of O-LM interneurons (Fig. 4B, C, Gloveli et al., 2005b) will provide a robust theta frequency rhythmic inhibitory output to the apical dendrites of PCs.

In vitro



In vivo




Fig. 4 Morphological and firing properties of O-LM interneurons. **A** Neurolucida reconstruction of biocytin-filled O-LM cell in area CA3 from transverse, longitudinal, and coronal slices. The soma and dendrites are drawn in *red*, whereas the axon is in *green*. Note different axonal ramification pattern in stratum lacunosum-moleculare in different slice preparations. Hippocampal layers are depicted schematically. CA3, CA3 area; str. or., stratum oriens; str. pyr., stratum pyramidale; str. rad., stratum radiatum; str. l-m., stratum lacunosum-moleculare. **B** Typical example of extracellular field potential (fp) and concomitant current clamp (−60 mV) recordings in an O-LM cell after induction of oscillatory activity with kainate in different slices. **C** Corresponding power spectra (60-s epoch) from field (*black*) and current clamp (*red*) recordings. **D–F** O-LM firing in vivo is specifically associated with different types of brain state and network activity. Filtered extracellular network oscillations (*top*) and extracellularly recorded action potentials (*bottom*). Note that the O-LM cell firing is not phase-coupled to gamma cycle, but fired rhythmically on the trough of theta oscillations and was silent during sharp-wave-associated ripples. Calibrations: **D** 0.1 mV (*upper trace*); 0.2 mV (*lower trace*) and 0.1 s; **E, F**, 0.3 mV (*lower traces*), 0.2 mV (*upper theta trace*), 0.05 mV (*upper sharp-wave-ripple trace*) and 300 ms (theta), 50 ms (sharp-wave-ripple). Adapted: Panels **A–C** from Gloveli et al. (2005b), Dugladze et al. (2007), Tort et al. (2007); Panel **D** from Tukker et al. (2007); Panels **E** and **F** from Klausberger et al. (2003)

In addition to distal dendrite inhibiting O-LM cells, the dendrite domains of principal cells are innervated by other types of interneurons, whose involvement in the hippocampal oscillations has not been addressed *in vitro*. This includes, for example, recently described interneuron type in CA1 area in anesthetized animal, *Ivy* cells, expressing neuropeptide Y (NPY), and the neuronal nitric oxide (NO). The soma of these cells is located in stratum pyramidale and axonal collaterals innervate two strata: oriens and radiatum. *Ivy* cells discharged at low frequency during theta as well as gamma and ripple oscillations in anesthetized animals (Fuentetaja et al., 2008). Another GABAergic interneuron type, *neurogliaform* cells that share some similarity with *Ivy* cells, such as dense axonal fields, low-frequency discharge, and slow synaptic transmission (Vida et al., 1998; Price et al., 2005; Szabadics et al., 2007), are located in stratum lacunosum-moleculare and innervate the apical dendrite tuft of CA1 pyramidal cells co-aligned with the entorhinal input (Price et al., 2005). This cell type provides both fast GABA_A receptor-mediated and slow GABA_B receptor-mediated (Price et al., 2005; Szabadics et al., 2007) inhibition and therefore represents a potential candidate to be involved in both theta and gamma frequency oscillations. However, there is very little information about the activity pattern of this interneuron type and their role in network oscillations.

Nested Theta and Gamma Oscillations

Theta and gamma oscillations often occur simultaneously and show interaction. Amplitude of gamma oscillations is modulated with the phase of the theta rhythm. In addition, the frequencies of the two oscillations are also correlated, providing additional evidence of their interrelated function (Bragin et al., 1995). The coordinated nature of the two rhythms, and the observation that gamma power is stronger during theta-associated behavior (Leung et al., 1982; Bragin et al., 1995), implies that the neuronal generators of the two rhythms interact (and may be also overlap). This nested activity pattern is hypothesized to play a critical role in memory encoding and retrieval (Lisman and Idiart, 1995; Lisman, 2005).

Recently, combined anatomical and physiological studies have provided evidence that *in vitro* gamma and theta rhythms are supported by neuronal circuits arranged orthogonally along the transverse and longitudinal axes, respectively (Table 1, Gloveli et al., 2005b). In hippocampal coronal slice preparation with intermediate orientations (between the transverse and longitudinal axis), both theta and gamma population rhythms were manifest (Fig. 4B, C, Gloveli et al., 2005b). The reason for that is a differential preservation of rhythm-generating microcircuits in transverse, longitudinal, and coronal slice preparation. Analysis of the three-dimensional axonal arborization patterns of different hippocampal CA3 interneurons recorded in transverse slices show that PV-expressing perisomatic-targeting interneurons, along with trilaminar and bistratified cells, show a clear tendency to arborize widely within the transverse plane (Gloveli et al., 2005a, b). In contrast,

distal dendrite-targeting O-LM cells arborized most extensively in the longitudinal plane forming two or three clusters in this direction (see Fig. 4A, Gloveli et al., 2005b). In longitudinal slices, the preservation of these projections facilitated the generation of theta rhythms (Fig. 4B, C) with robust coherence over large distances (Gloveli et al., 2005b; Tort et al., 2007). Thus, orthogonal arrangement of rhythm-generating microcircuits alongside the longitudinal and transverse axis, and distinct firing patterns of certain classes of interneurons during the theta and gamma frequency oscillations (Gloveli et al., 2005b; Tort et al., 2007), enables the hippocampus to produce different (solely or combined) population activity (for related simulation see “Gamma and Theta Rhythms in Biophysical Models of Hippocampal Circuits” by Kopell et al., this book).

Sharp-Waves Ripple Activity

The high-frequency oscillations termed ripples (100–300 Hz) are typically associated with sharp-wave activity (Buzsaki et al., 1992; Wilson and McNaughton, 1994; O’Neill et al., 2006). The hippocampal sharp-wave-ripple (SPW-R) complex is thought to play an important role in synaptic plasticity and the transfer of new memory trace from the hippocampus to the neocortex (Buzsáki, 1989).

The mechanisms of these fast oscillatory patterns in the hippocampus and neocortex are not fully understood (Buzsaki et al., 1992; Ylinen et al., 1995; Draguhn et al., 1998). Both in vivo and in vitro studies suggest that SWP-Rs arise in the recurrent collateral system of the CA3 area (similar to gamma oscillations), propagate toward CA1, and leave the hippocampal formation via the subiculum and the EC (Chrobak and Buzsaki, 1996; Csicsvari et al., 2000; Maier et al., 2003; Both et al., 2008). During this state, the hippocampus seems to be less controlled by input from the EC; rather, it generates output signals itself (Chrobak and Buzsaki, 1996).

SPW-R complex can be induced in vitro by electrical stimulation, pharmacologically, or can occur spontaneously with properties similar to the events seen in vivo (Maier et al., 2003; Nimmrich et al., 2005; Behrens et al., 2005). Several local network mechanisms underlying these patterns have been identified within CA1, including strong inhibition of non-participating pyramidal cells during SPW-R (Ylinen et al., 1995; Maier et al., 2003) and electrical coupling of CA1 pyramidal cells (Draguhn et al., 1998; Schmitz et al., 2001; Nimmrich et al., 2005). Concomitant extracellular and intracellular recordings of SPW-R complexes show that pyramidal cells display EPSP-IPSP sequences, IPSP-EPSP sequences, and prominent IPSPs, but never isolated EPSPs (Behrens et al., 2005). These results suggest that inhibitory inputs are strong during the development of ripple complexes. Consistent with this finding, fast-spiking basket and bistratified interneurons strongly increase their firing rate during ripple oscillations in vivo (Ylinen et al., 1995; Klausberger et al., 2004). Another fast-spiking cell type, axo-axonic interneurons fire before the ripple episode but are silenced during and after it (Klausberger

et al., 2003). In contrast, non-fast-spiking O-LM cell firing is suppressed during ripples (Klausberger et al., 2003). Gap junctions also seem to be important for ripples, since the blockade of gap junctions with carbenoxolone attenuated ripple occurrence (Behrens et al., 2005; LeBeau et al., 2003). Interestingly, SPW-R can be induced with stimulation protocols known to induce LTP, a model of learning and memory, suggesting that this pattern is associated with changes in functional connectivity (Behrens et al., 2005).

Cellular and Synaptic Mechanisms Involved in Oscillations

Intrinsic properties. The voltage-gated ion channels strongly contribute to *pyramidal cell* excitability. These channels influence intrinsic properties of the neuron, such as the action potential threshold, spike AHP and afterdepolarization (ADP), and action potential firing mode. Na^+ and A-type K^+ channels are expressed in both CA1 and CA3 PCs whereas hyperpolarization-activated cation channels (HCN channels) are expressed in CA1 PCs, but are almost absent from CA3 PCs (for review see Spruston, 2008). The HCN channels in PCs have important influences also on synaptic integration. Deactivation of these channels reduces EPSP duration and results in a slight hyperpolarization following EPSPs (Magee, 1999). Conversely, activation of HCN channels reduces IPSP duration and produces a slight depolarization following the IPSP (Williams and Stuart, 2003, for review see Spruston, 2008). This interaction of HCN channels and synaptic conductances may represent elementary mechanisms for rhythmogenesis at the cellular and subcellular levels.

Another feature relevant for the firing pattern of pyramidal cells in the hippocampus is their ability to generate subthreshold membrane potential oscillations (MPOs) in the theta frequency range and their resonance properties (Leung and Yu, 1998; Pike et al., 2000). These properties of hippocampal pyramidal cells are likely to contribute to theta activity (Leung and Yu, 1998). In hippocampal pyramidal cells the electrical resonance at theta frequencies is generated by M-current, h-current, and persistent Na^+ current (Hu et al., 2002).

How the different firing patterns of certain *GABAergic interneurons* are generated remains largely unknown. Intrinsic membrane properties of these cells may be important for hippocampal network oscillations. For instance, O-LM cells have a longer membrane time constant and a considerably longer (5-fold to 10-fold slower) AHP than the gamma-preferring interneurons (Gloveli et al., 2005a), restricting their firing to low, theta frequencies (see Savić et al., 2001). In addition, O-LM cells show prominent slow subthreshold membrane potential oscillations and the resonance properties in the theta frequency range (Maccaferri and McBain, 1996; Pike et al., 2000).

Hyperpolarization-activated cationic currents (I_h) and I_A currents which have been detected in hippocampal interneurons may influence not only the intrinsic and firing properties but also their synchronization. I_h currents are activated at voltages close to rest (Gu et al., 2005). Different subunit composition (HCN1–4) that is

co-expressed in hippocampal GABAergic interneurons (Notomi and Shigemoto, 2004) influences not only the kinetics but also the voltage dependency of I_h activation (see Chen et al., 2002). Besides O-LM and other types of non-fast-spiking interneurons, I_h channels are expressed in the somato-dendritic region, axon, and presynaptic elements of fast-spiking basket cell in the hippocampus (Aponte et al., 2006). In contrast, hippocampal lacunosum-moleculare and radiatum interneurons display subthreshold MPOs generated by an interplay of Na^+ and 4-AP-sensitive A-type K^+ currents, independent of I_h currents and muscarine-sensitive K^+ currents, I_M (Bourdeau et al., 2007).

Synaptic properties. The properties of excitatory events discriminate hippocampal principal cells from inhibitory neurons (Miles, 1990; Jonas et al., 1993; Geiger et al., 1997; Toth et al., 2000). It appears that excitatory synapses onto interneurons not only tend to have a larger number of AMPA receptors (Nusser et al., 1998), thereby increasing the quantal amplitude, but the postsynaptic receptors also appear to have a different molecular composition (Geiger et al., 1995), which, in turn, endows them with faster kinetics (Geiger et al., 1997). Further discrimination in the properties of excitatory input was found between different interneuron types. Different classes of hippocampal interneurons with distinct axonal ramification patterns and efferent target profiles show clear differences in both amplitude and kinetics of EPSCs/Ps during gamma frequency network oscillations (Gloveli et al., 2005a). For instance, the amplitudes of excitatory drive are considerably larger in fast-spiking BCs and trilaminar cells than in O-LM cells (Fig. 2B), suggesting that the intensity of synaptic drive may play a role in generating their different outputs (Gloveli et al., 2005a).

Similar to the kinetics of excitatory postsynaptic currents at PC–BC, unitary inhibitory postsynaptic currents at BC–BC synapses demonstrated very fast kinetics (Bartos et al., 2001, 2002). In addition to IPSCs with fast kinetic properties ($\text{GABA}_{A,\text{fast}}$) mediated by perisomatic synapses, IPSCs with slowly rising and decaying kinetic ($\text{GABA}_{A,\text{slow}}$) mediated by dendritic synapses were also detected in CA1 area (Banks et al., 2000). Interplay of CA1 interneurons, mediating $\text{GABA}_{A,\text{slow}}$ and $\text{GABA}_{A,\text{fast}}$, may contribute to theta and gamma rhythms occurring separately or as a nested gamma/theta rhythm (Banks et al., 2000).

Thus, different intrinsic membrane properties together with different kinetics of excitatory and inhibitory inputs govern the specific roles of hippocampal cells in shaping distinct network oscillatory activity.

Neuromodulatory Control of In Vitro Oscillations

Sources of neuromodulators in the brain are the four aminergic systems: the dopaminergic, histaminergic, serotonergic, and noradrenergic systems. All four of the associated modulators (dopamine, histamine, serotonin, and noradrenalin) are released from small groups of neurons, which have projection patterns to most of the brain, including the hippocampus. Effects of these neuromodulators have been tested in vitro on theta and gamma oscillatory activity.

The hippocampus receives dopaminergic input from the ventral tegmental area. Activation of D1-like *dopamine* receptors strongly depresses cholinergic gamma oscillations in area CA3 of rat hippocampus, and this effect is most likely mediated via impairment of interneurons involved in generation and maintenance of the carbachol-induced network rhythm (Weiss et al., 2003).

Histamine 3 (H3) receptors seem likely to play an important role in regulation of hippocampal theta oscillation. Systemic administration of the H3 receptor antagonists (ciproxifan and thioperamide) enhances the power of spontaneous theta in anesthetized rats. Since H3 receptors are located at axon terminals of histamine-containing neurons and function as autoreceptors (Arrang et al., 1983), their blockade could enhance histamine release and subsequently promote hippocampal theta oscillation. Regulation of hippocampal theta oscillations by H3 receptors may represent one of the probable mechanisms involved in histamine-induced modulation of higher brain functions, such as attention and learning (Hajós et al., 2008).

Serotonergic neurons of the midbrain raphe have been implicated in the control of affective and cognitive functions and in modulating the neural activities of networks across the sleep–wake cycle. The midbrain raphe nuclei form a strong serotonergic projection to the hippocampus. Recent in vivo findings suggest that a subpopulation of raphe neurons discharged action potentials that were phase-locked to the hippocampal theta rhythm (Kocsis et al., 2006). Hippocampal PCs and interneurons show different expression of metabotropic 5-HT receptor subtypes, such as 5-HT_{1A}, 5-HT_{1B}, 5-HT₂ (Ropert and Guy, 1991; Schmitz et al., 1995; Shen and Andrade, 1998), which may result in a differential modulation of intrinsic and synaptic properties of these cells in response to serotonin (for review see Schmitz et al., 1998). Serotonin input may influence the hippocampal network also via ionotropic 5-HT₃ receptors, which are expressed by several classes of GABAergic interneurons (Tecott et al., 1993; Ropert and Guy, 1991; Morales et al., 1998). These cells include CCK-containing basket cells, an interneuron type that has been proposed to hamper the gamma rhythm (see Section “Firing Patterns in Gamma Oscillations”, Freund and Katona, 2007; Galarreta et al., 2008), as well as calbindin-containing and calretinin-containing GABAergic cells (Morales and Bloom, 1997). In contrast, serotonergic fibers do not contact the PV-containing GABAergic basket cells, which are responsible for some gamma frequency oscillations (see Section “Firing Patterns in Gamma Oscillations”). Therefore, the rhythmic serotonergic input may modulate, but not drive, hippocampal network oscillations at gamma frequency range.

The brain *noradrenergic* (NE) neurons, located in the pontine nucleus of locus coeruleus (LC), are presumed to play a role in regulation of the circadian sleep–wake cycle and alertness (Aston-Jones and Cohen, 2005). Several experimental findings suggest involvement of these neuromodulators on the hippocampal network activity. Local injection of glutamate in the LC results in multiple actions on the hippocampus, which include an increase in theta rhythm (Brown et al., 2005). Activation of LC–NE neurons by local application of a cholinergic agonist (bethanechol) induces theta oscillation of MS/DB neurons and theta-wave oscillation of hippocampal EEG in anesthetized rats (Berridge and Foote, 1991). Furthermore, the selective NE

reuptake inhibitor reboxetine modulates hippocampal theta activity in a state-dependent manner, i.e., can either increase or decrease theta amplitude depending on the behavioral state of the animal (Kocsis et al., 2007).

Pharmacological agents which are used with in vitro models of oscillation, such as KAR, mGluR, and mAChR agonist, also have a direct modulatory effect on hippocampal neurons in a manner that is remarkably cell specific. Various types of interneurons express KARs (Cossart et al., 1998, 2002; Frerking et al., 1998; Mulle et al., 2000; Lerma, 2003). Kinetics of KA-mediated EPSCs is slower than that of AMPAR (Frerking et al., 1998; Cossart et al., 2002), which could enable these two receptor types to generate oscillations with different dominating frequencies (Frerking and Ohliger-Frerking, 2002). Consistent with this, O-LM interneurons, which receive a large input mediated by KARs (Cossart et al., 2002), show post-synaptic KAR-mediated action potential firing at 10 Hz during theta stimulations, in contrast to perisomatic, bistratified, or septum/back-projecting cells (Goldin et al., 2007). Activation of *mGluRs* in the hippocampus has a range of effects (Anwyl, 1999) which include decreases in I_M and I_{AHP} currents. Therefore, activation of these receptors increases the excitability of hippocampal cells. mGluR subtypes are differentially expressed in specific hippocampal interneurons resulting in their different responsiveness to agonists. Thus, O-LM cells express a large number of group I mGluRs and are very sensitive to agonists, in marked contrast to other stratum oriens interneurons, including basket cells that express only a small number of this receptor subtype and are less sensitive (van Hooft et al., 2000). Also *mAChR* agonists may act on different GABAergic inhibitory interneurons that possess the muscarinic receptors (Pitler and Alger, 1992). Activation of these receptors may increase the excitability of interneuronal activity. In particular, muscarinic receptor agonist-carbachol blocks several potassium conductances, including I_{AHP} and I_M in a concentration-dependent manner (Madison et al., 1987). This depolarizes the pyramidal cells, unmasking subthreshold membrane potential oscillations in the theta frequency range (Leung and Yim, 1986; Fellous and Sejnowski, 2000). In addition, muscarinic receptor activation consistently enhanced firing frequency and produced large, sustained ADPs of O-LM but not other stratum oriens interneurons (Lawrence et al., 2006).

In summary, the effects of state-dependent activation of different neuromodulators can be markedly different on hippocampal network activity and depend on the expression and distribution of receptors across the cellular components of the network.

Oscillations in Disease

Schizophrenia

A number of studies have shown changes in gamma frequency EEG activity in schizophrenia. Reduced gamma activity was found in stimulus-dependent responses

in the auditory and visual cortices of schizophrenic patients (for review see Kehrer et al., 2008), and there is also evidence for a change in neuronal synchrony during high-frequency oscillations (Spencer et al., 2004). Interestingly, the amounts of RNA and immunoreactivity for PV are reduced in postmortem tissue from the frontal cortex and the hippocampus, pointing to a reduction in perisomatic inhibitory interneuron population (Zhang and Reynolds, 2002). The loss of these interneurons could directly explain the observed changes in gamma oscillations (Lewis et al., 2005; Vierling-Claassen et al., 2008). Although most clinical studies have found reductions in gamma band activity in schizophrenics (e.g., Slewa-Younan et al., 2001), there appears to be a symptom-specific pattern in the alterations in gamma activity indicating that increases in amplitude and power are associated with positive symptoms, particularly hallucinations and reality distortions, whereas negative symptoms, such as psychomotoric deficits, are linked to decreased gamma activity (Baldeweg et al., 1998; Bucci et al., 2007). Similar to clinical observations, in *in vitro* NMDA-hypofunction models of schizophrenia, both increased (Kehrer et al., 2007; Pinault, 2008) and decreased gamma activities (Cunningham et al., 2006) have been demonstrated (for review see Kehrer et al., 2008; Roopun et al., 2008).

Alterations of cortical interneurons in schizophrenia, especially parvalbumin and somatostatin-containing interneurons, are well documented. These alterations are likely to have significant effects on the network oscillatory activity and therefore on cognitive processes (Gonzalez-Burgos and Lewis, 2008; Morris et al., 2008). The axo-axonic subclass of GABAergic interneurons containing the calcium binding protein parvalbumin has attracted the most scrutiny in studies of schizophrenia (Behrens et al., 2007; Sakai et al., 2008; Wang et al., 2008). Although the altered network activities were recently reported *in vivo* and *in vitro* (see e.g., Cunningham et al., 2006; Behrens et al., 2007; Braun et al., 2007; Gonzalez-Burgos and Lewis, 2008; Spencer, 2008), further investigation needs to be undertaken to address possible model-specific and region-specific alterations in the gamma network oscillatory activity in different animal models of schizophrenia. Establishing the contingencies of increased versus decreased gamma band activity is of high importance since aberrant network oscillatory activity may underlie the cognitive decline observed in schizophrenic patients and offer vital clues to the relationship between positive and negative symptoms in schizophrenia at a network level (Cho et al., 2006; Bucci et al., 2007; Ford et al., 2007).

Mesial Temporal Lobe Epilepsy (mTLE)

Epileptic seizures are less frequent in conditions during which theta frequency occurs (e.g., wakefulness or REM sleep, Montplaisir et al., 1987) and thus the theta rhythm appears to indicate a hippocampal functional state in which generation of seizures is hindered (Colom et al., 2006).

In epileptic mice, the power of low-frequency theta oscillations has been reported to be reduced (Arabadzisz et al., 2005; Dugladze et al., 2007). Apart from

suppression of the theta oscillatory activity, as a potential anticonvulsant factor (Colom et al., 2006), a strong enhancement of the gamma activity (Dugladze et al., 2007) may underlie emergence of epileptiform activity. Observations in humans support this scenario: spatially localized increase in the power of gamma frequency oscillations has been observed preceding seizures in human TLE patients (Fisher et al., 1992). In addition, gamma oscillatory activity and increases in firing rate of the interneuronal network have been suggested as mechanisms of seizure occurrence in patients with drug-resistant TLE (Bragin et al., 2007). Consistent with this suggestion, intracellular recordings in kainate model of mTLE reveal an increased firing frequency of both pyramidal cells and dendrite-inhibiting interneurons in the ventral hippocampal CA3 area of epileptic mice (Dugladze et al., 2007). The involvement of the perisomatic targeting interneurons remains, however, to be investigated.

The epileptic tissue may also generate a specific rhythm, transient high-frequency oscillations (HFOs) in the 200–500 Hz frequency band known as fast ripple (Bragin et al., 1999). In fact, fast ripples may represent a specific marker for the area of the brain in which seizures begin (Bragin et al., 2002). The frequency of these oscillations is about twice as fast as the maximum rate at which most neurons in the hippocampus can fire action potentials. This fact raises the question of how these oscillations are generated. Analysis of the firing properties of hippocampal neurons during HFOs in vitro low-Mg²⁺ model of epileptiform activity showed that the pyramidal cells fired at the rising phase of the highest frequency portion of the field oscillation. In addition, distal dendrite-targeting interneurons (R-LM cells) fired at the start of the epileptiform bursts (on average 140 Hz) but stopped firing before its end (Spampanato and Mody, 2007). However, neither the principal cells nor the distal dendrite-targeting interneurons (R-LM and O-LM cell) fired action potentials at high frequencies (200–600 Hz) seen in the field oscillations (Spampanato and Mody, 2007). Another study (Foffani et al., 2007) suggested that these synchronous population oscillations could be generated as a consequence of the out-of-phase activities of two independent oscillators, each operating at half the frequency of the ensemble. In line with this suggestion, it was found that hippocampal pyramidal cells fired short bursts of action potentials at frequencies up to 300 Hz (Kandel and Spencer, 1961) and some interneurons could sustain frequencies of 400 Hz (Foffani et al., 2007).

Thus, numerous studies suggest that alterations in inhibition-based network oscillations may underlie the pathophysiology in schizophrenia and mTLE, which are associated with impaired information processes.

Perspectives

The spatiotemporal patterns of activity during network oscillations would ideally be explored in vivo (Bragin et al., 1995; Penttonen et al., 1998; Csicsvari et al., 2003; Buzsáki et al., 2003). However, it is reasonable to use relevant in vitro models to test hypotheses for the basic mechanisms involved. The replication of an endogenous

brain pattern in vitro allows for the investigation of a number of important cellular and synaptic mechanisms that are difficult or even impossible to explore in vivo. In addition, using transgenic, fluorescent EGFP-expressing mice under the control of different gene promoters (Oliva et al., 2000; Meyer et al., 2002) enables the identification and selection of different interneurons in the acute slice preparation. The firing properties of single cells in the active network and the contribution of excitation and inhibition to the generation of network oscillatory activity can be systematically examined in different models of in vitro oscillations (Table 1), which may reflect region and state dependence of mechanisms underlying network oscillations in vivo. Although there are some differences between the data obtained from in vitro and in vivo observations, these observations show substantial homology. The development of new transgenic methods for activating, inactivating, and labeling neurons and synapses (Marek and Davis, 2003; Polleux, 2005) will certainly facilitate progress in this area. Furthermore, several recently developed methods in vitro may help to overcome the limitations in vivo. These include the following: simultaneous patch-clamp recording from several cells in a brain slice (Miles and Poncer, 1996; Markram et al., 1997); recording of sufficiently large numbers of cells at once using optical methods; stimulation/suppression of different cellular compartments using uncaging of different substances or optically activated channels (Callaway, 2002; Boyden et al., 2005; Deisseroth et al., 2006; Zhang et al., 2007).

Further Reading

- Anwyl R (1999) Metabotropic glutamate receptors: electrophysiological properties and role in plasticity. *Brain Res Rev* 29:83–120.
- Aponte Y, Lien CC, Reisinger E, Jonas P (2006) Hyperpolarization-activated cation channels in fast-spiking interneurons of rat hippocampus. *J Physiol* 574:229–243.
- Arabadzisz D, Antal K, Parpan F, Emri Z, Fritschy JM (2005) Epileptogenesis and chronic seizures in a mouse model of temporal lobe epilepsy are associated with distinct EEG patterns and selective neurochemical alterations in the contralateral hippocampus. *Exp Neurol* 194:76–90.
- Arnolds DE, Lopes da Silva FH, Aitink JW, Kamp A, Boeijinga, P (1980) The spectral properties of hippocampal EEG related to behaviour in man. *Electroencephalogr Clin Neurophysiol* 50:324–328.
- Arrang JM, Garbarg M, Schwartz JC (1983) Auto-inhibition of brain histamine release mediated by a novel class (H3) of histamine receptor. *Nature (Lond)* 302:832–837.
- Aston-Jones G, Cohen JD (2005) An integrative theory of locus coeruleus-norepinephrine function: adaptive gain and optimal performance. *Annu Rev Neurosci* 28:403–450.
- Baldeweg T, Spence S, Hirsch SR, Gruzeliér J (1998) Gamma-band electroencephalographic oscillations in a patient with somatic hallucinations. *Lancet* 352:620–621.
- Banks MI, White JA, Pearce RA (2000) Interactions between distinct GABA(A) circuits in hippocampus. *Neuron* 25:449–457.
- Bartos M, Vida I, Frotscher M, Geiger JR, Jonas P (2001) Rapid signaling at inhibitory synapses in a dentate gyrus interneuron network. *J Neurosci* 21:2687–2698.
- Bartos M, Vida I, Frotscher M, Meyer A, Monyer H, Geiger JR, Jonas P (2002) Fast synaptic inhibition promotes synchronized gamma oscillations in hippocampal interneuron networks. *Proc Natl Acad Sci USA* 99:13222–13227.
- Behrens CJ, van den Boom LP, de Hoz L, Friedman A, Heinemann U (2005) Induction of sharp wave-ripple complexes in vitro and reorganization of hippocampal networks. *Nat Neurosci* 8:1560–1567.

- Behrens MM, Ali SS, Dao DN, Lucero J, Shekhtman G, Quick KL, Dugan LL (2007) Ketamine-induced loss of phenotype of fast-spiking interneurons is mediated by NADPH-oxidase. *Science* 318:1645–1647.
- Berridge CW, Foote SL (1991) Effects of locus coeruleus activation on electroencephalographic activity in neocortex and hippocampus. *J Neurosci* 11:3135–3145.
- Bland BH (1986) Physiology and pharmacology of hippocampal formation theta rhythms. *Prog Neurobiol* 26:1–54.
- Both M, Böhner F, von Bohlen und Halbach O, Draguhn A (2008) Propagation of specific network patterns through the mouse hippocampus. *Hippocampus* 18:899–908.
- Bourdeau ML, Morin F, Laurent CE, Azzi M, Lacaille JC (2007) Kv4.3-mediated A-type K⁺ currents underlie rhythmic activity in hippocampal interneurons. *J Neurosci* 27:1942–1953.
- Boyden ES, Zhang F, Bamberg E, Nagel G, Deisseroth K (2005) Millisecond-timescale, genetically targeted optical control of neural activity. *Nat Neurosci* 8:1263–1268.
- Bragin A, Jandó G, Nádasdy Z, Hetke J, Wise K, Buzsáki G (1995) Gamma (40–100 Hz) oscillation in the hippocampus of the behaving rat. *J Neurosci* 15:47–60.
- Bragin A, Engel J Jr, Wilson CL, Fried I, Buzsáki G (1999) High-frequency oscillations in human brain. *Hippocampus* 9:137–142.
- Bragin A, Mody I, Wilson CL, Engel J Jr (2002) Local generation of fast ripples in epileptic brain. *J Neurosci* 22:2012–2021.
- Bragin A, Claeys P, Vonck K, Van Roost D, Wilson C, Boon P, Engel J Jr (2007) Analysis of initial slow waves (ISWs) at the seizure onset in patients with drug resistant temporal lobe epilepsy. *Epilepsia* 48:1883–1894.
- Braun I, Genius J, Grunze H, Bender A, Möller HJ, Rujescu D (2007) Alterations of hippocampal and prefrontal GABAergic interneurons in an animal model of psychosis induced by NMDA receptor antagonism. *Schizophr Res* 97:254–263.
- Brown RA, Walling SG, Milway JS, Harley CW (2005) Locus coeruleus activation suppresses feedforward interneurons and reduces β - γ electroencephalogram frequencies while it enhances theta frequencies in rat dentate gyrus. *J Neurosci* 25:1985–1991.
- Bucci P, Mucci A, Merlotti E, Volpe U, Galderisi S (2007) Induced gamma activity and event-related coherence in schizophrenia. *Clin EEG Neurosci* 38:96–104.
- Buhl EH, Halasy K, Somogyi P (1994) Diverse sources of hippocampal unitary inhibitory postsynaptic potentials and the number of synaptic release sites. *Nature* 368:823–828.
- Buhl EH, Tamás G, Fisahn A (1998) Cholinergic activation and tonic excitation induce persistent gamma oscillations in mouse somatosensory cortex in vitro. *J Physiol* 513:117–126.
- Buzsáki, G (1989) Two-stage model of memory trace formation: a role for “noisy” brain states. *Neuroscience* 31:3551–3570.
- Buzsáki G (2002) Theta oscillations in the hippocampus. *Neuron* 33:325–340.
- Buzsáki G, Buhl DL, Harris KD, Csicsvari J, Czéh B, Morozov A (2003) Hippocampal network patterns of activity in the mouse. *Neuroscience* 116:201–211.
- Buzsáki G, Czopf J, Kondakor J, Kelenyi L (1986) Laminar distribution of hippocampal rhythmic slow activity (RSA) in the behaving rat; current source density analysis, effects of urethane and atropine. *Brain Res* 365:125–137.
- Buzsaki G, Draguhn A (2004) Neuronal oscillations in cortical networks. *Science* 304:1926–1929.
- Buzsaki G, Horvath Z, Urioste R, Hetke J, Wise K (1992) High-frequency network oscillation in the hippocampus. *Science* 256:1025–1027.
- Buzsaki G, Leung LW, Vanderwolf CH (1983) Cellular bases of hippocampal EEG in the behaving rat. *Brain Res* 287:139–171.
- Callaway EM (2002) Cell type specificity of local cortical connections. *J Neurocytol* 31:231–237.
- Chen K, Aradi I, Santhakumar V, Soltesz I (2002) H-channels in epilepsy: new targets for seizure control? *Trends Pharmacol Sci* 23:552–557.
- Cho RY, Konecny RO, Carter CS (2006) Impairments in frontal cortical gamma synchrony and cognitive control in schizophrenia. *Proc Natl Acad Sci USA* 103:19878–19883.

- Chrobak JJ, Buzsáki G (1996) High-frequency oscillations in the output networks of the hippocampal-entorhinal axis of the freely behaving rat. *J Neurosci* 16:3056–3066.
- Cobb SR, Buhl EH, Halasy K, Paulsen O, Somogyi P (1995) Synchronisation of neuronal activity in hippocampus by individual GABAergic interneurons. *Nature* 378:75–78.
- Cobb SR, Butlers DO, Davies CH (2000) Coincident activation of mGluRs and MACHRs imposes theta frequency patterning on synchronised network activity in the hippocampal CA3 region. *Neuropharmacology* 23:1933–1942.
- Colom LV, Garcia-Hernandez A, Castaneda MT, Perez-Cordova MG, Garrido-Sanabria ER (2006) Septo-hippocampal networks in chronically epileptic rats: potential antiepileptic effects of theta rhythm generation. *J Neurophysiol* 95:3645–3653.
- Cossart R, Esclapez M, Hirsch JC, Bernard C, Ben-Ari Y (1998) GluR5 kainate receptor activation in interneurons increases tonic inhibition of pyramidal cells. *Nature Neurosci* 1:470–478.
- Cossart R, Epsztein J, Tyzio R, Becq H, Hirsch J, Ben-Ari Y, Crépel V (2002) Quantal release of glutamate generates pure kainate and mixed AMPA/kainate EPSCs in hippocampal neurons. *Neuron* 35:147–159.
- Csicsvari J, Hirase H, Czurko A, Mamiya A, Buzsáki G (1999) Oscillatory coupling of hippocampal pyramidal cells and interneurons in the behaving rat. *J Neurosci* 19:274–287.
- Csicsvari J, Hirase H, Mamiya A, Buzsáki G (2000) Ensemble patterns of hippocampal CA3-CA1 neurons during sharp wave-associated population events. *Neuron* 28:585–594.
- Csicsvari J, Jamieson B, Wise KD, Buzsáki G (2003) Mechanisms of gamma oscillations in the hippocampus of the behaving rat. *Neuron* 37:311–322.
- Cunningham MO, Hunt J, Middleton S, LeBeau FE, Gillies MJ, Davies CH, Maycox PR, Whittington MA, Racca C (2006) Region-specific reduction in entorhinal gamma oscillations and parvalbumin-immunoreactive neurons in animal models of psychiatric illness. *J Neurosci* 26:2767–2776.
- Deisseroth K, Feng G, Majewska AK, Miesenböck G, Ting A, Schnitzer MJ (2006) Next-generation optical technologies for illuminating genetically targeted brain circuits. *J Neurosci* 26:10380–10386.
- Dodt HU, Ziegglänsberger W (1994) Infrared videomicroscopy: a new look at neuronal structure and function. *Trends Neurosci* 17:453–458.
- Draguhn A, Traub RD, Schmitz D, Jefferys JG (1998) Electrical coupling underlies high-frequency oscillations in the hippocampus in vitro. *Nature* 394:189–192.
- Dugladze T, Vida I, Tort AB, Gross A, Otahal J, Heinemann U, Kopell NJ, Gloveli T (2007) Impaired hippocampal rhythmogenesis in a mouse model of mesial temporal lobe epilepsy. *Proc Natl Acad Sci USA* 104:17530–17535.
- Fellous JM, Sejnowski TJ (2000) Cholinergic induction of oscillations in the hippocampal slice in the slow bands (0.5–2 Hz), theta (5–12 Hz), and gamma (35–70 Hz). *Hippocampus* 10:187–197.
- Fisahn A, Pike FG, Buhl EH, Paulsen O (1998) Cholinergic induction of network oscillations at 40 Hz in the hippocampus in vitro. *Nature* 394:186–189.
- Fisahn A, Yamada M, Duttaroy A, Gan JW, Deng CX, McBain CJ, Wess J (2002) Muscarinic induction of hippocampal gamma oscillations requires coupling of the M1 receptor to two mixed cation currents. *Neuron* 33:615–624.
- Fisahn A, Contractor A, Traub RD, Buhl EH, Heinemann SF, McBain CJ (2004) Distinct roles for the kainate receptor subunits GluR5 and GluR6 in kainate-induced hippocampal gamma oscillations. *J Neurosci* 24:9658–9668.
- Fisher RS, Webber WR, Lesser RP, Arroyo S, Uematsu S (1992) High-frequency EEG activity at the start of seizures. *J Clin Neurophysiol* 9:441–448.
- Ford JM, Krystal JH, Mathalon DH (2007) Neural synchrony in schizophrenia: from networks to new treatments. *Schizophr Bull* 33:848–852.
- Frerking M, Malenka RC, Nicoll RA (1998) Synaptic activation of kainate receptors on hippocampal interneurons. *Nat Neurosci* 1:479–486.

- Frerking M, Ohliger-Frerking P (2002) AMPA receptors and kainate receptors encode different features of afferent activity. *J Neurosci* 22:7434–7443.
- Freund TF, Buzsáki G (1996) Interneurons of the hippocampus. *Hippocampus* 6:347–470.
- Freund TF, Katona I (2007) Perisomatic inhibition. *Neuron* 56:33–42.
- Foffani G, Uzcategui YG, Gal B, Menendez de la Prida L (2007) Reduced spike-timing reliability correlates with the emergence of fast ripples in the rat epileptic hippocampus. *Neuron* 55:930–941.
- Fuentealba P, Begum R, Capogna M, Jinno S, Márton LF, Csicsvari J, Thomson A, Somogyi P, Klausberger T (2008) Ivy cells: a population of nitric-oxide-producing, slow-spiking GABAergic neurons and their involvement in hippocampal network activity. *Neuron* 57:917–929.
- Galarreta M, Erdélyi F, Szabó G, Hestrin S (2008) Cannabinoid sensitivity and synaptic properties of 2 GABAergic networks in the neocortex. *Cereb Cortex* 18:2296–22305.
- Geiger JR, Melcher T, Koh DS, Sakmann B, Seeburg PH, Jonas P, Monyer H (1995) Relative abundance of subunit mRNAs determines gating and Ca²⁺ permeability of AMPA receptors in principal neurons and interneurons in rat CNS. *Neuron* 15:193–204.
- Geiger JR, Lübke J, Roth A, Frotscher M, Jonas P (1997) Submillisecond AMPA receptor-mediated signaling at a principal neuron-interneuron synapse. *Neuron* 18:1009–1023.
- Gillies MJ, Traub RD, LeBeau FE, Davies CH, Gloveli T, Buhl EH, Whittington MA (2002) A model of atropine-resistant theta oscillations in rat hippocampal area CA1. *J Physiol (Lond)* 543:779–793.
- Gloveli T, Dugladze T, Saha S, Monyer H, Heinemann U, Traub RD, Whittington MA, Buhl EH (2005a) Differential involvement of oriens/pyramidal interneurons in hippocampal network oscillations in vitro. *J Physiol* 562:131–147.
- Gloveli T, Dugladze T, Rotstein HG, Traub RD, Monyer H, Heinemann U, Whittington MA, Kopell NJ (2005b) Orthogonal arrangement of rhythm-generating microcircuits in the hippocampus. *Proc Natl Acad Sci USA* 102:13295–13300.
- Goldin M, Epsztein J, Jorquera I, Represa A, Ben-Ari Y, Crépel V, Cossart R (2007) Synaptic kainate receptors tune oriens-lacunosum moleculare interneurons to operate at theta frequency. *J Neurosci* 27:9560–9572.
- Gonzalez-Burgos G, Lewis DA (2008) GABA Neurons and the mechanisms of network oscillations: implications for understanding cortical dysfunction in schizophrenia. *Schizophr Bull* 34:944–961.
- Gu N, Vervaeke K, Hu H, Storm JF (2005) Kv7/KCNQ/M and HCN/h, but not KCa2/SK channels, contribute to the somatic medium after-hyperpolarization and excitability control in CA1 hippocampal pyramidal cells. *J Physiol* 566:689–715.
- Hajós N, Pálhalmi J, Mann EO, Németh B, Paulsen O, Freund TF (2004) Spike timing of distinct types of GABAergic interneuron during hippocampal gamma oscillations in vitro. *J Neurosci* 24:9127–9137.
- Hajós M, Siok CJ, Hoffmann WE, Li S, Kocsis B (2008) Modulation of hippocampal theta oscillation by histamine H3 receptors. *J Pharmacol Exp Ther* 324:391–398.
- Halasy K, Buhl EH, Lörinczi Z, Tamás G, Somogyi P (1996) Synaptic target selectivity and input of GABAergic basket and bistratified interneurons in the CA1 area of the rat hippocampus. *Hippocampus* 6:306–329.
- Hu H, Vervaeke K, Storm JF (2002) Two forms of electrical resonance at theta frequencies, generated by M-current, h-current and persistent Na⁺ current in rat hippocampal pyramidal cells. *J Physiol* 545:783–805.
- Jonas P, Major G, Sakmann B (1993) Quantal components of unitary EPSCs at the mossy fibre synapse on CA3 pyramidal cells of rat hippocampus. *J Physiol* 472:615–663.
- Jouvet M (1969) Biogenic amines and the states of sleep. *Science* 163:32–41.
- Kandel ER, Spencer WA (1961) Electrophysiology of hippocampal neurons. II. After-potentials and repetitive firing. *J Neurophysiol* 24:243–259.
- Kehrer C, Dugladze T, Maziashvili N, Wójtowicz A, Schmitz D, Heinemann U, Gloveli T (2007) Increased inhibitory input to CA1 pyramidal cells alters hippocampal gamma frequency oscillations in the MK-801 model of acute psychosis. *Neurobiol Dis* 25:545–552.

- Kehrer C, Maziashvili N, Dugladze T, Gloveli T (2008) Altered excitatory-inhibitory balance in the NMDA-hypofunction model of schizophrenia. *Front Mol Neurosci* 1:6.
- Klausberger T, Somogyi P (2008) Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science* 321:53–57.
- Klausberger T, Magill PJ, Márton LF, Roberts JD, Cobden PM, Buzsáki G, Somogyi P (2003) Brain-state- and cell-type-specific firing of hippocampal interneurons in vivo. *Nature* 421: 844–848.
- Klausberger T, Márton LF, Baude A, Roberts JD, Magill PJ, Somogyi P (2004) Spike timing of dendrite-targeting bistratified cells during hippocampal network oscillations in vivo. *Nat Neurosci* 7:41–47.
- Kocsis B, Varga V, Dahan L, Sik A (2006) Serotonergic neuron diversity: identification of raphe neurons with discharges time-locked to the hippocampal theta rhythm. *Proc Natl Acad Sci USA* 103:1059–1064.
- Kocsis B, Li S, Hajos M (2007) Behavior-dependent modulation of hippocampal EEG activity by the selective norepinephrine reuptake inhibitor reboxetine in rats. *Hippocampus* 17:627–633.
- Konopacki J, Gołębiewski H, Eckersdorf B (1992) Carbachol-induced rhythmic slow activity (theta) in cat hippocampal formation slices. *Brain Res* 578:13–16.
- Lawrence JJ, Statland JM, Grinspan ZM, McBain CJ (2006) Cell type-specific dependence of muscarinic signalling in mouse hippocampal stratum oriens interneurons. *J Physiol* 570: 595–610.
- LeBeau FEN, Towers SK, Traub RD, Whittington MA, Buhl EH (2002) Fast network oscillations induced by potassium transients in the rat hippocampus in vitro. *J Physiol (Lond)* 542:167–179.
- LeBeau FE, Traub RD, Monyer H, Whittington MA, Buhl EH (2003) The role of electrical signalling via gap junctions in the generation of fast network oscillations. *Brain Res Bull* 62:3–13.
- Lerma J (2003) Roles and rules of kainate receptors in synaptic transmission. *Nat Rev Neurosci* 4:481–495.
- Leung LW (1985) Spectral analysis of hippocampal EEG in the freely moving rat effects of centrally active drugs and relations to evoked potentials. *Electroencephalogr Clin Neurophysiol* 60:65–77.
- Leung LS, Yim CY (1986) Intracellular records of theta rhythm in hippocampal CA1 cells of the rat. *Brain Res* 367:323–327.
- Leung LS, Yu HW (1998) Theta-frequency resonance in hippocampal CA1 neurons in vitro demonstrated by sinusoidal current injection. *J Neurophysiol* 79:1592–1596.
- Leung LS, Lopes da Silva F, Wadman WJ (1982) Spectral characteristics of the hippocampal EEG in the freely moving rat. *Clin Neurophysiol* 54:203–219.
- Lewis DA, Hashimoto T, Volk DW (2005) Cortical inhibitory neurons and schizophrenia. *Nat Rev Neurosci* 6:312–324.
- Li XG, Somogyi P, Ylinen A, Buzsáki G (1994) The hippocampal CA3 network: an in vivo intracellular labeling study. *J Comp Neurol* 339:181–208.
- Lisman JE (2005) The theta/gamma discrete phase code occurring during the hippocampal phase precession may be a more general brain coding scheme. *Hippocampus* 15:913–922.
- Lisman JE, Idiart MAP (1995) Storage of 7+– 2 short term memories in oscillatory subcycles. *Science* 267:1512–1515.
- Maccaferri G, McBain CJ (1996) The hyperpolarization-activated current (I_h) and its contribution to pacemaker activity in rat CA1 hippocampal stratum oriens-alveus interneurons. *J Physiol* 497:119–130.
- Maccaferri G, Roberts JD, Szucs P, Cottingham CA, Somogyi P (2000) Cell surface domain specific postsynaptic currents evoked by identified GABAergic neurones in rat hippocampus in vitro. *J Physiol* 524:91–116.
- Madison DV, Lancaster B, Nicoll RA (1987) Voltage clamp analysis of cholinergic action in the hippocampus. *J Neurosci* 7:733–741.
- Magee JC (1999) Dendritic hyperpolarization-activated currents modify the integrative properties of hippocampal CA1 pyramidal neurons. *J Neurosci* 18:7613–7624.

- Maier N, Nimmrich V, Draguhn A (2003) Cellular and network mechanisms underlying spontaneous sharp wave-ripple complexes in mouse hippocampal slices. *J Physiol* 550:873–887.
- Mann EO, Suckling JM, Hajos N, Greenfield SA, Paulsen O (2005) Perisomatic feedback inhibition underlies cholinergically induced fast network oscillations in the rat hippocampus in vitro. *Neuron* 45:105–117.
- Marek KW, Davis GW (2003) Controlling the active properties of excitable cells. *Curr Opin Neurobiol* 13:607–611.
- Markram H, Lubke J, Frotscher M, Roth A, Sakmann B (1997) Physiology and anatomy of synaptic connections between thick tufted pyramidal neurones in the developing rat neocortex. *J Physiol* 500:409–440.
- McBain CJ, Fisahn A (2001) Interneurons unbound. *Nat Rev Neurosci* 2:11–23.
- Meyer AH, Katona I, Bлатow M, Rozov A, Monyer H (2002) In vivo labeling of parvalbumin-positive interneurons and analysis of electrical coupling in identified neurons. *J Neurosci* 22:7055–7064.
- Middleton S, Jalics J, Kispersky T, Lebeau FE, Roopun AK, Kopell NJ, Whittington MA, Cunningham MO (2008) NMDA receptor-dependent switching between different gamma rhythm-generating microcircuits in entorhinal cortex. *Proc Natl Acad Sci USA* 105:18572–18577.
- Miles R (1990) Synaptic excitation of inhibitory cells by single CA3 hippocampal pyramidal cells of the guinea-pig in vitro. *J Physiol* 428:61–77.
- Miles R, Poncer JC (1996) Paired recordings from neurones. *Curr Opin Neurobiol* 6:387–394.
- Miles R, Tóth K, Gulyás AI, Hájos N, Freund TF (1996) Differences between somatic and dendritic inhibition in the hippocampus. *Neuron* 16:815–823.
- Montplaisir J, Laverdiere M, Saint-Hilaire JM, Rouleau I (1987) Nocturnal sleep recording in partial epilepsy: a study with depth electrodes. *J Clin Neurophysiol* 4:383–388.
- Morales M, Bloom FE (1997) The 5-HT₃ receptor is present in different subpopulations of GABAergic neurons in the rat telencephalon. *J Neurosci* 17:3157–3167.
- Morales M, Battenberg E, Bloom F (1998) Distribution of neurons expressing immunoreactivity for the 5HT₃ receptor subtype in the rat brain and spinal cord. *J Comp Neurol* 402:385–401.
- Morris HM, Hashimoto T, Lewis DA (2008) Alterations in somatostatin mRNA expression in the dorsolateral prefrontal cortex of subjects with schizophrenia or schizoaffective disorder. *Cereb Cortex* 18:1575–1587.
- Mulle C, Sailer A, Swanson GT, Brana C, O’Gorman S, Bettler B, Heinemann SF (2000) Subunit composition of kainate receptors in hippocampal interneurons. *Neuron* 28:475–484.
- Nimmrich V, Maier N, Schmitz D, Draguhn A (2005) Induced sharp wave-ripple complexes in the absence of synaptic inhibition in mouse hippocampal slices. *J Physiol* 563:663–670.
- Notomi T, Shigemoto R (2004) Immunohistochemical localization of Ih channel subunits, HCN1–4, in the rat brain. *J Comp Neurol* 471:241–276.
- Nusser Z, Lujan R, Laube G, Roberts JD, Molnar E, Somogyi P (1998) Cell type and pathway dependence of synaptic AMPA receptor number and variability in the hippocampus. *Neuron* 21:545–559.
- Oliva AA Jr, Jiang M, Lam T, Smith KL, Swann JW (2000) Novel hippocampal interneuronal subtypes identified using transgenic mice that express green fluorescent protein in GABAergic interneurons. *J Neurosci* 20:3354–3368.
- O’Neill J, Senior T, Csicsvari J (2006) Place-selective firing of CA1 pyramidal cells during sharp wave/ripple network patterns in exploratory behavior. *Neuron* 49:143–155.
- Oren I, Mann EO, Paulsen O, Hajos N (2006) Synaptic currents in anatomically identified CA3 neurons during hippocampal gamma oscillations in vitro. *J Neurosci* 26:9923–9934.
- Pálhalmi J, Paulsen O, Freund TF, Hájos N (2004) Distinct properties of carbachol- and DHPG-induced network oscillations in hippocampal slices. *Neuropharmacology* 47:381–289.
- Penttonen M, Kamondi A, Acsády L, Buzsáki G (1998) Gamma frequency oscillation in the hippocampus of the rat: intracellular analysis in vivo. *Eur J Neurosci* 10:718–728.

- Petsche H, Stumpf C, Gogolák G (1962) The significance of the rabbit's septum as a relay station between midbrain and the hippocampus: I. The control of hippocampus arousal activity by the septum cells. *Electroencephalogr Clin Neurophysiol* 14:202–211.
- Pike FG, Goddard RS, Suckling JM, Ganter P, Kasthuri N, Paulsen O (2000) Distinct frequency preferences of different types of rat hippocampal neurones in response to oscillatory input currents. *J Physiol* 529:205–213.
- Pinault D (1996) A novel single-cell staining procedure performed in vivo under electrophysiological control: morpho-functional features of juxtacellularly labeled thalamic cells and other central neurons with biocytin or neurobiotin. *J Neurosci Methods* 65:113–136.
- Pinault D (2008) N-methyl d-aspartate receptor antagonists ketamine and MK-801 induce wake-related aberrant gamma oscillations in the rat neocortex. *Biol Psychiatry* 63:730–735.
- Pitler TA, Alger BE (1992) Cholinergic excitation of GABAergic interneurons in the rat hippocampal slice. *J Physiol* 450:127–142.
- Polleux F (2005) Genetic mechanisms specifying cortical connectivity: let's make some projections together. *Neuron* 46:395–400.
- Pöschel B, Draguhn A, Heinemann U (2002) Glutamate-induced gamma oscillations in the dentate gyrus of rat hippocampal slices. *Brain Res* 938:22–28.
- Pouille F, Scanziani M (2004) Routing of spike series by dynamic circuits in the hippocampus. *Nature* 429:717–723.
- Price CJ, Cauli B, Kovacs ER, Kulik A, Lambolez B, Shigemoto R, Capogna M (2005) Neurogliaform neurons form a novel inhibitory network in the hippocampal CA1 area. *J Neurosci* 25:6775–6786.
- Roopun AK, Cunningham MO, Racca C, Alter K, Traub RD, Whittington MA (2008) Region-specific changes in gamma and beta2 rhythms in NMDA receptor dysfunction models of schizophrenia. *Schizophr Bull* 34:962–973.
- Ropert N, Guy N (1991) Serotonin facilitates GABAergic transmission in the CA1 region of rat hippocampus in vitro. *J Physiol (Lond)* 441:121–136.
- Rotstein HG, Pervouchine DD, Acker CD, Gillies MJ, White JA, Buhl EH, Whittington MA, Kopell N (2005) Slow and fast inhibition and an H-current interact to create a theta rhythm in a model of CA1 interneuron network. *J Neurophysiol* 94:1509–1518.
- Sakai T, Oshima A, Nozaki Y, Ida I, Haga C, Akiyama H, Nakazato Y, Mikuni M (2008) Changes in density of calcium-binding-protein-immunoreactive GABAergic neurons in prefrontal cortex in schizophrenia and bipolar disorder. *Neuropathology* 28:143–150.
- Savić N, Pedarzani P, Sciancalepore M (2001) Medium afterhyperpolarization and firing pattern modulation in interneurons of stratum radiatum in the CA3 hippocampal region. *J Neurophysiol* 85:1986–1997.
- Schmitz D, Empson RM, Heinemann U (1995) Serotonin reduces inhibition via 5-HT_{1A} receptors in area CA1 of rat hippocampal slices in vitro. *J Neurosci* 15:7217–7225.
- Schmitz D, Gloveli T, Empson RM, Heinemann U (1998) Comparison of the effects of serotonin in the hippocampus and the entorhinal cortex. *Mol Neurobiol* 17:59–72.
- Schmitz D, Schuchmann S, Fisahn A, Draguhn A, Buhl EH, Petrasch-Parwez E, Dermietzel R, Heinemann U, Traub RD (2001) Axi-axonal coupling. A novel mechanism for ultrafast neuronal communication. *Neuron* 31:831–840.
- Shen RY, Andrade R (1998) 5-Hydroxytryptamine₂ receptor facilitates GABAergic neurotransmission in rat hippocampus. *J Pharmacol Exp Ther* 285:805–12.
- Shimono K, Brucher F, Granger R, Lynch G, Taketani M (2000) Origins and distribution of cholinergically induced beta rhythms in hippocampal slices. *J Neurosci* 20:8462–8473.
- Sik A, Penttonen M, Ylinen A, Buzsáki G (1995) Hippocampal CA1 interneurons: an in vivo intracellular labeling study. *J Neurosci* 15:6651–6665.
- Singer W (1999) Neuronal synchrony: a versatile code for the definition of relations? *Neuron* 24:49–65.
- Slawa-Younan S, Gordon E, Williams L, Haig AR, Goldberg E (2001) Sex differences, gamma activity and schizophrenia. *Int J Neurosci* 107:131–144.

- Somogyi P, Klausberger T (2005) Defined types of cortical interneurone structure space and spike timing in the hippocampus. *J Physiol (Lond)* 562:9–26.
- Spampanato J, Mody I (2007) Spike timing of lacunosum-molecular targeting interneurons and CA3 pyramidal cells during high-frequency network oscillations in vitro. *J Neurophysiol* 98:96–104.
- Spencer KM, Nestor PG, Perlmuter R, Niznikiewicz MA, Klump MC, Frumin M, Shenton ME, McCarley RW (2004) Neural synchrony indexes disordered perception and cognition in schizophrenia. *Proc Natl Acad Sci USA* 101:17288–17293.
- Spencer KM (2008) Visual gamma oscillations in schizophrenia: implications for understanding neural circuitry abnormalities. *Clin EEG Neurosci* 39:65–68.
- Spruston N (2008) Pyramidal neurons: dendritic structure and synaptic integration. *Nat Rev Neurosci* 9:206–221.
- Szabadics J, Tamas G, Soltesz I (2007) Different transmitter transients underlie presynaptic cell type specificity of GABAA, slow and GABAA, fast. *Proc Natl Acad Sci USA* 104:14831–14836.
- Tecott LH, Maricq AV, Julius D (1993) Nervous system distribution of the serotonin 5-HT₃ receptor mRNA. *Proc Natl Acad Sci USA* 90:1430–1434.
- Tesche CD, Karhu J (2000) Theta oscillations index human hippocampal activation during a working memory task. *Proc Natl Acad Sci USA* 97:919–924.
- Tort AB, Rotstein HG, Dugladze T, Gloveli T, Kopell NJ (2007) On the formation of gamma-coherent cell assemblies by oriens lacunosum-molecular interneurons in the hippocampus. *Proc Natl Acad Sci USA* 104:13490–13495.
- Toth K, Soares G, Lawrence JJ, Philips-Tansey E, McBain CJ (2000) Differential mechanisms of transmission at three types of mossy fiber synapse. *J Neurosci* 20:8279–8289.
- Towers SK, LeBeau FE, Gloveli T, Traub RD, Whittington MA, Buhl EH (2002) Fast network oscillations in the rat dentate gyrus in vitro. *J Neurophysiol* 87:1165–1168.
- Traub RD, Bibbig A, Fisahn A, LeBeau FE, Whittington MA, Buhl EH (2000) A model of gamma-frequency network oscillations induced in the rat CA3 region by carbachol in vitro. *Eur J Neurosci* 12:4093–4106.
- Traub RD, Cunningham MO, Gloveli T, LeBeau FE, Bibbig A, Buhl EH, Whittington MA (2003) GABA-enhanced collective behavior in neuronal axons underlies persistent gamma-frequency oscillations. *Proc Natl Acad Sci USA* 100:11047–11052.
- Tukker JJ, Fuentealba P, Hartwich K, Somogyi P, Klausberger T (2007) Cell type-specific tuning of hippocampal interneuron firing during gamma oscillations in vivo. *J Neurosci* 27:8184–8189.
- van Hooft JA, Giuffrida R, Blatow M, Monyer H (2000) Differential expression of group I metabotropic glutamate receptors in functionally distinct hippocampal interneurons. *J Neurosci* 20:3544–3551.
- Vanderwolf CH (1969) Hippocampal electrical activity and voluntary movement in the rat. *Electroencephalogr Clin Neurophysiol* 26:407–418.
- Vida I, Halasy K, Szinyei C, Somogyi P, Buhl EH (1998) Unitary IPSPs evoked by interneurons at the stratum radiatum-stratum lacunosum-molecular border in the CA1 area of the rat hippocampus in vitro. *J Physiol* 506:755–773.
- Vida I, Bartos M, Jonas P (2006) Shunting inhibition improves robustness of gamma oscillations in hippocampal interneuron networks by homogenizing firing rates. *Neuron* 49:107–117.
- Vierling-Claassen D, Siekmeier P, Stufflebeam S, Kopell N (2008) Modeling GABA alterations in schizophrenia: a link between impaired inhibition and altered gamma and beta range auditory entrainment. *J Neurophysiol* 99:2656–2671.
- Wang X-J, Buzsáki G (1996) Gamma oscillation by synaptic inhibition in a hippocampal interneuronal network model. *J Neurosci* 16:6402–6413.
- Wang CZ, Yang SF, Xia Y, Johnson KM (2008) Postnatal phencyclidine administration selectively reduces adult cortical parvalbumin-containing interneurons. *Neuropsychopharmacology* 33:2442–2455.
- Weiss T, Veh RW, Heinemann U (2003) Dopamine depresses cholinergic oscillatory network activity in rat hippocampus. *Eur J Neurosci* 18:2573–2580.

- White JA, Chow CC, Ritt J, Soto-Treviño C, Kopell N (1998) Synchronization and oscillatory dynamics in heterogeneous, mutually inhibited neurons. *J Comput Neurosci* 5:5–16.
- Whittington MA, Traub RD (2003) Interneuron diversity series: inhibitory interneurons and network oscillations in vitro. *Trends Neurosci* 26:676–682.
- Whittington MA, Stanford IM, Colling SB, Jefferys JG, Traub RD (1997) Spatiotemporal patterns of gamma frequency oscillations tetanically induced in the rat hippocampal slice. *J Physiol* 502:591–607.
- Whittington MA, Traub RD, Jefferys JGR (1995) Synchronized oscillations in interneuron networks driven by metabotropic glutamate receptor activation. *Nature* 373:612–615.
- Williams SR, Stuart GJ (2003) Voltage- and site-dependent control of the somatic impact of dendritic IPSPs. *J Neurosci* 23:7358–7367.
- Wilson MA, McNaughton BL (1994) Reactivation of hippocampal ensemble memories during sleep. *Science* 265:676–679.
- Wittner L, Henze DA, Záborszky L, Buzsáki G (2007) Three-dimensional reconstruction of the axon arbor of a CA3 pyramidal cell recorded and filled in vivo. *Brain Struct Funct* 212:75–83.
- Ylinen A, Bragin A, Nadasdy Z, Jando G, Szabo I, Sik A, Buzsaki G (1995) Sharp wave-associated high-frequency oscillation (200 Hz) in the intact hippocampus: network and intracellular mechanisms. *J Neurosci* 15:30–46.
- Zhang ZJ, Reynolds GP (2002) A selective decrease in the relative density of parvalbumin-immunoreactive neurons in the hippocampus in schizophrenia. *Schizophr Res* 55:1–10.
- Zhang F, Wang LP, Brauner M, Liewald JF, Kay K, Watzke N, Wood PG, Bamberg E, Nagel G, Gottschalk A, Deisseroth K (2007) Multimodal fast optical interrogation of neural circuitry. *Nature* 446:633–639.