

Poster presentation

Open Access

Neuronal correlates of emotions in human-machine interaction

Benjamin Blankertz*^{1,2}, Klaus-Robert Müller¹ and Gabriel Curio³

Address: ¹Berlin Institute of Technology, Machine Learning Laboratory, Germany, ²Fraunhofer FIRST, Berlin, Germany and ³Dept. of Neurology, Campus Benjamin Franklin, Charité University Medicine Berlin, Germany

Email: Benjamin Blankertz* - blanker@cs.tu-berlin.de

* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS*2009
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, **10**(Suppl 1):P80 doi:10.1186/1471-2202-10-S1-P80

This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/P80>

© 2009 Blankertz et al; licensee BioMed Central Ltd.

Introduction and method

Previous neurophysiological studies of emotions have focused on the affective response in the emotional valence of a situation in which the reaction is rooted in perception or memories [1]. Furthermore emotions have been investigated with regard to the trait of a subject, e.g. *anger-out* vs. *anger control* [2] and regarding motivational direction, e.g. *approach* vs. *withdrawal* [3]. Aiming at an enhancement of human-computer interaction by incorporating the emotional state of the user, a novel type of investigation is required. Neuronal correlates of emotional reactions related to interaction (e.g. annoyance due to one's own failure or an error of the machine; joy of success) have to be analyzed and methods for their detection in real-time need to be developed. In the present study we have acquired multi-channel EEG in four subjects while they were interacting with computer applications that have been specifically designed in order to provoke – in alternating phases – neural, positive or negative (stress, annoyance) emotions. In particular, a two-player variant of a two-alternative forced-choice task had to be performed while in alternating periods either one or the other player was given "unfair" preferential treatment by providing the task stimulus slightly in advance. This bias could not be noticed by the players.

Results and discussion

The behavioral data are consistent between subjects and indicate that the participants had a temporary feeling of inferiority and adapted their strategy (accepting higher

error rates in order to achieve faster reaction times to cope with the competitor). As neuronal correlates, intra-individual significant differences between periods of negative and positive emotions were found in the theta-, alpha-, or beta-band with widely distributed and spatially coherent topographies, see Figure 1 for the result from one subject. Notably, the frequency band as well as its spatial focus varied between subjects. The variety of EEG correlates found among the four subjects already demonstrates the need for adaptive methods in order to enhance human-machine interfaces by emotional decoding. Experimental studies with a larger number of subjects will show whether some pattern clusters could be identified that potentially correlate with individual attitudes towards the particular emotion-provoking environmental situation.

Acknowledgements

This work was supported in parts by grants of the BMBF (01IB001A/B, 01GQ0850) and the EU (ICT-216886).

References

1. Krause CM, Viemerö V, Rosenqvist A, Sillanmäki L, Aström T: **Relative electroencephalographic desynchronization and synchronization in humans to emotional film content: an analysis of the 4–6, 6–8, 8–10 and 10–12 Hz frequency bands.** *Neurosci Lett* 2000, **286**:9-12.
2. Hewig J, Hagemann D, Seifert J, Naumann E, Bartusek D: **On the selective relation of frontal cortical asymmetry and anger-out versus anger-control.** *J Personal Social Psychol* 2004, **87**:926-939.
3. Harmon-Jones E: **Contributions from research on anger and cognitive dissonance to understanding the motivational functions of asymmetrical frontal brain activity.** *Biol Psychol* 2004, **67**:51-76.




Figure 1
Topographical maps of alpha band-power (9–13 Hz) for the emotional states "stress" and "annoyance" (minus baseline) in one subject. For comparison a map of alpha band-power is shown for the condition "eyes closed" which has a focus clearly different from the other states.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

