

# Neuroprotective Effects of Lithium: Implications for the Treatment of Alzheimer's Disease and Related Neurodegenerative Disorders


O. V. Forlenza,<sup>\*,†</sup> V. J. R. De-Paula,<sup>\*,†</sup> and B. S. O. Diniz<sup>‡</sup>

<sup>†</sup>Laboratory of Neuroscience (LIM-27), Department and Institute of Psychiatry, Faculty of Medicine, University of Sao Paulo, SP, Brazil

<sup>‡</sup>Department of Mental Health and National Institute of Science and Technology, Molecular Medicine, Faculty of Medicine, Federal University of Minas Gerais, Belo Horizonte, MG, Brazil

**ABSTRACT:** Lithium is a well-established therapeutic option for the acute and long-term management of bipolar disorder and major depression. More recently, based on findings from translational research, lithium has also been regarded as a neuroprotective agent and a candidate drug for disease-modification in certain neurodegenerative disorders, namely, Alzheimer's disease (AD), amyotrophic lateral sclerosis (ALS), and, more recently, Parkinson's disease (PD). The putative neuroprotective effects of lithium rely on the fact that it modulates several homeostatic mechanisms involved in neurotrophic response, autophagy, oxidative stress, inflammation, and mitochondrial function. Such a wide range of intracellular responses may be secondary to two key effects, that is, the inhibition of glycogen synthase kinase-3 beta (GSK-3 $\beta$ ) and inositol monophosphatase (IMP) by lithium. In the present review, we revisit the neurobiological properties of lithium in light of the available evidence of its neurotrophic and neuroprotective properties, and discuss the rationale for its use in the treatment and prevention of neurodegenerative diseases.

**KEYWORDS:** Lithium, neuroprotection, GSK-3 $\beta$ , autophagy, bipolar disorder, Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis



Lithium salts have long been used in psychiatry for the treatment of severe mental disorders.<sup>1</sup> Currently, the main medical indications of lithium are for the acute and long-term treatment of bipolar disorder (BD) and for the adjunctive treatment of major depression, given its well-established mood stabilizing properties.<sup>2</sup> More recently, there has been a growing body of evidence indicating that the neurobiological benefits of lithium may go beyond mood stabilization. In experimental and clinical models, lithium treatment has been associated with neuroprotection, due to its effects on several mechanisms of neuronal homeostasis involved in the activation of neurotrophic responses, modulation of oxidative stress, inflammatory cascades, up-regulation of mitochondrial function, and other specific biological effects implicated in the pathogenesis of neurodegenerative diseases such as Alzheimer's disease (AD) and amyotrophic lateral sclerosis (ALS).<sup>3</sup> This article aims to review the mechanisms by which lithium may exert its neuroprotective effects, and how these mechanisms may help delay the progression of neurobiological changes in mood and neurocognitive disorders. Additionally, we address the potential of lithium as a disease-modifying agent for certain neurodegenerative and dementing conditions.

## NEUROBIOLOGICAL PROPERTIES OF LITHIUM

The pharmacological mechanisms of lithium are not completely understood, but current evidence suggests the direct involve-

ment of classic pharmacological targets affecting neurotransmission and signal transduction. These include the modulation of cell-surface receptors, the release of second-messengers and downstream signaling molecules, and the subsequent effect on the activity of important regulatory systems, with an impact on the release of transcription factors and gene expression.<sup>4</sup> Monovalent lithium (Li<sup>+</sup>) competes with bivalent magnesium (Mg<sup>2+</sup>) to the similar ionic radii of these cations (0.60 and 0.65 Å respectively), rendering the ability of lithium to bind to Mg<sup>2+</sup> substrate sites. Therefore, lithium can inhibit a wide range of enzymes that depend on Mg<sup>2+</sup> as a cofactor.<sup>5,6</sup> The competition between lithium and Mg<sup>2+</sup> on these substrate sites has a significant influence on the activity of several enzymes and therefore the release of their metabolic products; in particular, glycogen synthase kinase-3 beta (GSK-3 $\beta$ ), inositol monophosphatase (IMP), and Akt/ $\beta$ -arrestin2 (Akt) are important lithium targets. Therefore, the modification of these intracellular pathways through enzymatic inhibition is

**Special Issue:** Current Hypotheses of Lithium's Mechanism of Action as a Neuropsychiatric Medication

**Received:** February 17, 2014

**Revised:** April 22, 2014

**Published:** April 25, 2014

relevant to the understanding of the pathogenesis of certain neuropsychiatric and neurodegenerative disorders.

GSK-3 has two isoforms, alpha and beta, with distinct patterns of distribution and homeostatic roles. GSK-3 $\beta$  is more abundant in the brain, and is implicated in cytoskeletal organization and remodeling.<sup>7</sup> Conversely, cerebral GSK-3 $\alpha$  is involved in neurodevelopment<sup>8</sup> and its inhibition by lithium has also been associated with disease-modification in a transgenic mouse model of AD.<sup>9</sup> The inhibition of GSK-3 $\beta$  is one of the most relevant mechanisms of action of lithium, and substantiates its putative role as a candidate for a disease-modifying drug in the treatment or prevention of AD.<sup>10,11</sup> Lithium inhibits GSK-3 $\beta$  activity by two distinct and interrelated mechanisms: directly by preventing the binding of Mg<sup>2+</sup> to the catalytic core of GSK-3 $\beta$ , and indirectly through inducing the phosphorylation of the serine-9 residue of GSK-3 $\beta$ , leading to conformational changes and inactivation, which is required for enzymatic activity. Therefore, by the competitive dislocation of Mg<sup>2+</sup>, lithium reversibly inhibits the enzyme.<sup>12,13</sup> The indirect mechanism is followed by the activation of intracellular kinases (e.g., Akt) or the inhibition of intracellular phosphatases (e.g., PP2A) by lithium.<sup>14–16</sup> Finally, lithium can also reduce the availability of GSK-3 $\beta$  at the transcriptional level, therefore reducing its protein expression as a consequence of the inhibition of mRNA transcription.<sup>17</sup>

Another relevant mechanism of action of lithium is the inhibition of inositol monophosphatase (IMP) and inositol polyphosphate-1 (IPP). As with GSK-3 $\beta$ , lithium directly inhibits IMP and IPP activity by the competitive displacement of Mg<sup>2+</sup> from the catalytic site of the enzyme.<sup>18</sup> The inhibition of IMP and IPP prevents the reuptake of inositol, leading to depletion of intracellular levels and subsequent inhibition of the phosphoinositol cycle. Another important consequence of IMP inhibition is the suppression of the formation of its metabolic product inositol triphosphate (IP3); IP3 is an intracellular messenger implicated in the regulation of many intracellular pathways relevant to neuropsychiatric disorders, including autophagy, an important homeostatic mechanism based on the degradation of cytoplasmic proteins and organelles.<sup>19</sup> Sarkar et al.<sup>20</sup> found in mammalian cell cultures that lithium induced autophagy as a downstream effect of the inhibition of IMP.

Lithium, in a dose-dependent manner, modulates autophagy through both the GSK-3 $\beta$  and IMPase pathways, with opposite effects. Lithium-induced IMPase inhibition at lower doses ( $\approx 0.8$  mM), up-regulating autophagy,<sup>20</sup> while the inhibition of GSK-3 $\beta$  by higher doses of lithium ( $\approx 2$  mM) down-regulates autophagy via activation of the inhibitory regulator mTOR.<sup>21,22</sup> The overall ability of lithium to induce autophagy is due to the prevailing inhibition of IMPase.<sup>20</sup> Finally, lithium has been shown to act on other homeostatic pathways as well, such as extracellular signal-regulated kinase (ERK), PI3k/Akt, and phospholipase C (PLC), which are proteins with further impact on the regulation of autophagy.<sup>23,24</sup>

### ■ PRECLINICAL EVIDENCE OF THE NEUROPROTECTIVE EFFECT OF LITHIUM

As stated above, lithium has specific properties that may attenuate the effect of critical pathological changes that occur in AD, namely, through the inhibition of the GSK-3 $\beta$ . In fact, this is the cornerstone to support the “GSK hypothesis of AD”,<sup>25</sup> according to which the inhibition of GSK-3 $\beta$  activity by lithium is associated with the down-regulation of two central processes in the pathogenesis of AD, namely, the reduction of the

hyperphosphorylation of microtubule-associated tau protein,<sup>26–28</sup> and the induction of neuronal death via overproduction of the A $\beta$  peptide.<sup>29–31</sup> In transgenic mice overexpressing GSK-3 $\beta$  and in other animal models of AD, chronic lithium treatment significantly reduced tau phosphorylation.<sup>32–34</sup> Likewise, chronic lithium treatment reduced A $\beta$ <sub>42</sub> production by a direct modulation of APP processing and by inhibition of GSK-3 $\beta$  activity.<sup>35,36</sup> It is noteworthy that the attenuation or reversal of AD-related neuropathology was accompanied by a significant improvement in memory deficits in these animal models.<sup>37–39</sup>

In addition to these disease-specific mechanisms, lithium may also protect neurons against the neurotoxic effects of A $\beta$ <sub>42</sub> by favoring other neurotrophic and/or neuroprotective responses.<sup>40,41</sup> Chen and Chuang<sup>42</sup> showed that lithium increases the expression of p53 and Bcl-2, favoring neuronal survival. Chen et al.<sup>43</sup> showed the chronic administration of two structurally dissimilar mood stabilizing agents, lithium and valproate, increases Bcl-2 levels in the cortex, with beneficial neuroprotective effects. Also, lithium significantly stimulates the proliferation of progenitor cells in neuronal cell cultures,<sup>44,45</sup> and increases the expression of antiapoptotic proteins (e.g., Bcl-2).<sup>46–48</sup> Recent evidence suggests that lithium treatment enhances the mitochondrial respiratory rate, reduces oxidative stress, protects DNA against damage from oxidative stress, and modulates calcium influx in the mitochondria.<sup>49–54</sup> Lithium treatment also stimulates autophagic processes due to its inhibition of IMP/IPP activity and reduction of IP3 formation, in spite of GSK-3 $\beta$  inhibition.<sup>55,56</sup>

Another important neuroprotective effect of lithium is the stimulation of synthesis and release of neurotrophic factors, in particular brain-derived neurotrophic factor (BDNF) and vascular endothelial growth factor (VEGF). Increased availability of these factors protects neurons against neurotoxic insults, stimulates hippocampal neurogenesis, increases synaptic plasticity and long-term potentiation (LTP), and positively regulates cell survival.<sup>3</sup> Finally, lithium can regulate inflammatory processes by lessening the pro-inflammatory response. Lithium has also been shown to reduce the production of interleukin-1 beta (IL-1 $\beta$ ) and tumor necrosis factor alpha (TNF- $\alpha$ ), inducers of lipopolysaccharide (LPS)-induced inflammation in glial cells,<sup>57</sup> and to reduce microglial activation secondary to ischemic insult in mice.<sup>58</sup> Chronic lithium treatment can attenuate arachidonic acid production, an essential feature of the innate inflammatory response.<sup>59</sup> Therefore, the modulation of inflammatory processes by lithium is relevant in light of the prominent role of inflammation in neurodegenerative and mood disorders.<sup>60</sup>

Complex interactions between genetic and environmental factors are believed to play a critical role in the pathophysiology of neuropsychiatric disorders. The epigenetic status is affected by environmental stimuli and insults, leading to DNA methylation and histone modifications. Therefore, a better understanding of the epigenetic mechanisms affecting neuronal cells will provide important insights into the pathophysiology of cognitive and mood disorders and clues to new treatment approaches.<sup>61,62</sup> Epigenetic studies conducted in the post-mortem brains of patients with major depression found evidence of hypermethylation of several genes involved in neuronal response, such as *BDNF*, *DBN1*, *SLC6A4*, and *PRIMA1* in the prefrontal cortex.<sup>63–65</sup> Studies further provided evidence implicating of GSK-3 $\beta$  in the regulation of DNA methylation in mouse embryonic stem cells (ESCs).<sup>66</sup> The de

novo DNA methyltransferase gene (*Dnmt3a2*) is down-regulated in GSK-3 $\beta$  double knockout ESCs, decreasing DNA methylation. The inhibition of GSK-3 $\beta$  activity by lithium mimics the effects of reducing DNA methylation in both wild-type ESCs and wild-type neural stem cells. In addition, the inactivation of GSK-3 $\beta$  via components of the insulin signaling pathway also results in reduced DNA methylation.<sup>67</sup>

### ■ CLINICAL AND IMAGING FINDINGS SUPPORTING THE NEUROTROPHIC AND NEUROPROTECTIVE PROPERTIES OF LITHIUM IN BIPOLAR DISORDER

In addition to the cumulative evidence derived from experimental models, clinical and neuroimaging studies with patients with bipolar and other mood disorders further corroborate the neuroprotective properties of lithium. Most of the current clinical evidence derives from studies using subjects with BD. Large case registry studies found that BD patients continuously treated with lithium had a significantly lower risk of dementia, compared to those on other mood stabilizers or without treatment.<sup>68,69</sup> In a retrospective study, Terao and colleagues<sup>70</sup> found that patients on chronic lithium treatment had lower rates of cognitive decline as measured by the Mini-Mental State Examination (MMSE). In a cross-sectional study from our group, we found that older bipolar patients on chronic lithium treatment had a significantly lower incidence of AD (3%) compared to those with no or minimal lifetime lithium exposure (19%).<sup>71</sup> In this study, the incidence rates of AD in the group treated with lithium was comparable to those observed in the general population,<sup>72</sup> suggesting that chronic lithium treatment can be protective against the development of dementia (particularly AD) in the long term outcome of BD.

Acute and chronic lithium treatment of BD patients has been associated with the up-regulation of certain biological cascades related to neuroprotection. Lithium treatment significantly increased plasma concentrations of BDNF, with influence on response to treatment.<sup>73,74</sup> In a clinical trial with patients in acute mania, a significant increase in plasma concentrations of BDNF was observed after 4 weeks of treatment with lithium; however, increased BDNF levels were not associated with treatment response.<sup>75</sup> In addition, maintenance treatment with lithium was associated with persistently higher levels of BDNF and reduced risk of relapse after a major affective episode.<sup>76</sup>

Lithium can also modulate other important biological processes related to inflammation and oxidative stress. Lithium treatment of acute mania episodes was associated with a reduction of pro-oxidative stress markers, namely TBARS.<sup>77</sup> In addition, lithium treatment increased antioxidative, and reduced pro-oxidative, markers in healthy subjects.<sup>78</sup> The reduction in pro-oxidative stress markers was associated with significant clinical improvement in depressive symptoms after lithium treatment.<sup>79</sup> Finally, a recent study demonstrated that patients with BD who had a good response to lithium also had a significant reduction in plasma concentrations of TNF- $\alpha$ ; in contrast, patients who did not respond well to lithium showed a significant increase in TNF- $\alpha$  levels.<sup>80</sup> Lithium can restore the balance between the production of IL-1 $\beta$  and IL-6 in monocytes of bipolar patients *in vitro*; this effect is similar to those observed *in vivo*.<sup>81</sup>

Neuroimaging studies have provided further support for the neuroprotective effects of lithium. Structural neuroimaging studies showed that short and long-term lithium treatment was associated with a volumetric increase in the hippocampus and

amygdala, in addition to increased cortical thickness.<sup>82–85</sup> In a recent multicenter, observational study, BD patients on continuous lithium treatment had significantly larger hippocampi compared to those with no or minimal lifetime exposure to lithium.<sup>86</sup> Finally, lithium treatment was associated with increased *N*-acetyl aspartate (NAA) and myo-inositol levels as shown by magnetic resonance spectroscopy.<sup>87,88</sup> These imaging and neurochemical findings suggest that long-term lithium treatment may have a significant effect on synaptic density, neuronal vitality, and mitochondrial function in BD patients.

### ■ CLINICAL EVIDENCE OF NEUROPROTECTIVE EFFECTS OF LITHIUM IN NEURODEGENERATIVE DISORDERS

**Alzheimer's Disease (AD).** The rationale for the clinical use of lithium as a neuroprotective therapy derives from preclinical models of AD, indicating that lithium can preclude or attenuate A $\beta$  and tau pathology, and improve cognitive function in transgenic mice. These results encouraged the conduction of clinical studies in patients with AD; however, few studies have been presented thus far. In a small open label trial with 25 patients with mild to moderate AD conducted in the United Kingdom, MacDonald et al.<sup>89</sup> found no significant effects of lithium on cognitive function after 1 year of treatment. Nonetheless, the authors concluded that lithium was a safe drug in older adults as most of the side effects and dropouts from the trial were due to mild and reversible side effects. In a single-blind, multicenter clinical trial with 71 patients with mild to moderate AD conducted in Europe, Hampel et al.<sup>90</sup> did not find any significant benefits of lithium on cognitive performance associated with 10 weeks of treatment at therapeutic levels (0.5–0.8 mmol/L). In this study, the authors evaluated the effect of lithium on biomarkers related to AD, and did not find any significant changes in cerebrospinal fluid (CSF) concentrations of A $\beta$ <sub>42</sub> and phosphorylated tau, nor in phosphorylated GSK-3 $\beta$  (i.e., the inactive form of this enzyme) levels in leukocytes. The authors hypothesize that their negative results, in light of the short duration of treatment, were insufficient for lithium to exert its neuroprotective effects, or at least for these effects to be represented by changes in biomarker levels. Secondary analysis of this trial showed that lithium treatment was associated with increased plasma concentrations of BDNF. In this subset of AD patients, lithium treatment restored low baseline BDNF to levels similar to controls, and patients who displayed an increase in BDNF also had significant cognitive improvement.<sup>91</sup> The effect of lithium was specific to the BDNF response, as no significant changes were observed in levels of glial cell-derived neurotrophic factor (GDNF) both in the CSF and serum of AD patients after 10 weeks of lithium treatment.<sup>92</sup>

More recently, we conducted a double-blind, placebo controlled, clinical trial to evaluate whether long-term treatment with lithium at subtherapeutic levels (serum levels of 0.2–0.5 mmol/L) could delay the progression from amnesic mild cognitive impairment (MCI) to dementia, and to evaluate the disease-modifying properties as illustrated by the modification of clinical and biological markers of AD in patients with MCI.<sup>11</sup> This study recruited 45 older adults with amnesic MCI, and the preliminary analysis after 1 year of follow-up showed that amnesic MCI subjects receiving lithium presented stable cognitive performance and lower conversion rates to AD compared to subjects on placebo, although the latter difference was not statistically significant. However, significant differences

in favor of the lithium group were observed on multiple cognitive parameters, namely memory, attention and global cognitive function. In addition, lithium use was associated with a significant reduction in CSF concentrations of phosphorylated tau as compared to subjects in the placebo group. Additional analyses revealed that the effect size of lithium on phosphorylated tau levels was even greater in MCI subjects who did not progress to AD upon follow-up. Overall, these results suggest that long-term lithium treatment may have disease-modifying properties on the core pathophysiologic features of AD and deliver a marginal clinical benefit, mostly if started at the earlier stages of the disease process. In another recent clinical trial conducted by a different Brazilian group, Nunes et al.<sup>71</sup> demonstrated a significant improvement in global cognitive performance (as shown by the MMSE) using continuous microdoses of lithium (300  $\mu\text{g}$  daily) for 18 months. The authors state that these benefits started after 6 months of treatment and persisted until the end of the trial.

As stated earlier, autophagy is a key intracellular pathway dedicated to the degradation of mutant proteins, some of which are associated with neurodegeneration.<sup>93</sup> Lithium ultimately induces autophagy via its effect on the dominant regulatory mechanism, which is dependent on the inhibition of IMP.<sup>20</sup> Autophagy is also induced by active GSK-3 $\beta$ ;<sup>93</sup> therefore, the inhibition of GSK-3 $\beta$  by lithium leads to the attenuation of autophagy. This effect has been shown to occur via the activation of mTOR.<sup>21</sup> Therefore, there is a clear interplay between distinct regulatory mechanisms that may be differentially affected by lithium depending on the prevailing pathological process of the neurodegenerative disease. The overall effect of lithium on these mechanisms and their clinical implications still need to be clarified by future controlled studies.

**Amyotrophic Lateral Sclerosis (ALS).** The potential neuroprotective effects of lithium were also evaluated in ALS, a severe progressive neurodegenerative disorder that affects motor neurons leading to premature disability and death.<sup>94</sup>

Dill et al.<sup>95</sup> suggested the potential neuroprotective effect of lithium in ALS by demonstrating the ability of lithium to induce the sprouting of pyramidal neurons in the corticospinal tract following mechanical injury. In primary neuronal cultures obtained from the ventral spinal cord, Busceti et al.<sup>96</sup> suggested that the neurotrophic response and synaptogenesis induced by lithium could be relevant for the treatment of ALS, with a possible impact on disease progression. This effect was related to the inhibition of GSK-3 $\beta$  (and subsequent decrease in tau phosphorylation) and upregulation of autophagy, which may further increase the clearance of hyperphosphorylated tau. Therefore, it is likely that distinct pathways may contribute to the neuroprotective effects of lithium on neurodegenerative diseases associated with hyperphosphorylated tau, such as AD, ALS, and some forms of frontotemporal dementia.<sup>97</sup>

Preclinical studies have shown a significant improvement in motor function in animal models of ALS treated with lithium. The main hypothesized mechanism for such improvement was the stimulation of autophagy by lithium.<sup>98,99</sup> In an early clinical trial, lithium treatment for 15 months was shown to be safe and significantly associated with a slower rate of disease progression and death in these patients.<sup>98</sup> However, more recent and larger trials failed to show a significant benefit of lithium for this condition.<sup>100,101</sup>

In animal models of ALS, the cotreatment of lithium with valproate (another mood stabilizing drug) has been shown to

produce more beneficial effects than the treatment with either drug alone.<sup>102</sup> Similar findings were presented by other authors using animal models of Huntington's disease<sup>103</sup> and traumatic brain injury.<sup>104</sup>

Chiò et al.<sup>105</sup> recently conducted a phase 3 multicenter, double-blind, placebo-controlled trial of lithium versus placebo in ALS. Patients were randomly assigned into two groups to receive either lithium ( $n = 107$ ) or matched placebo tablets ( $n = 107$ ). Oral doses of lithium carbonate (mean serum levels ranging from 0.4 to 0.8 mmol/L) or placebo were continuously administered for 18 months. The primary end point was the rate of survival after 18 months, which was ascertained by intention to treat analysis. Unfortunately, the study results did not support any evidence of increased survival associated with lithium treatment.

**Other Neurodegenerative Diseases.** Lithium has been also studied in preclinical models of other neurodegenerative diseases, including Parkinson's<sup>106,107</sup> and Huntington's disease.<sup>108–110</sup> The pattern of cell death in Parkinson's disease is complex, having features of apoptosis and necrosis in addition to accumulations of autophagosome-like structures.<sup>111</sup> Using an *in vitro* model of Parkinson's disease, Chen et al.<sup>112</sup> demonstrated that 6-OHDA activates GSK-3 $\beta$  in cultured human neuroblastoma SH-SY5Y cells as well as in cultures of rat cerebellar granule neurons. Lithium and other specific GSK-3 $\beta$  inhibitors effectively protected against neuronal death after exposure to 6-OHDA, indicating that GSK-3 $\beta$  is involved in 6-OHDA-induced apoptosis of SH-SY5Y cells and cerebellar granule neurons. However, other studies in dopaminergic neurons have presented conflicting results: 6-OHDA treatment was not associated with GSK-3 $\beta$  activation, and 6-OHDA-induced degeneration was not inhibited by lithium.<sup>113</sup> These results suggest that GSK-3 $\beta$  activity may not be centrally involved in 6-OHDA-induced dopaminergic neurodegeneration in the substantia nigra (pars compacta) of rats. In a rat model of Huntington's disease, a protocol of chronic subcutaneous injections of lithium showed that lithium treatment may protect against brain damage caused by focal cerebral ischemia and suppresses excitotoxicity-induced striatal lesions.<sup>114</sup> Despite the promising neuroprotective potential against disease mechanisms described in these studies, no clinical trials have been conducted so far in human patients to test these findings.

## CONCLUSIONS

Converging lines of evidence derived from preclinical and clinical models support the rationale for the study of the protective effects of lithium in neuropsychiatric conditions associated with chronic degeneration of the central nervous system. This effect is probably due to the modulation of multiple biological cascades that are involved in cell survival, neuronal plasticity, transcriptional control, energetic metabolism, and resilience against neurotoxic insults. Some of these mechanisms may be represented as core pathological processes of mood and neurodegenerative disorders. The knowledge on the specific effects of lithium on distinct pathways critically relevant to neuronal homeostasis, and the broad understanding of their interactions, will guide the development of novel therapeutic strategies against neurodegeneration, aiming at both symptom reduction and attenuation of disease progression.

## ■ AUTHOR INFORMATION

## Corresponding Author

\*Mailing address: Laboratory of Neuroscience, Institute of Psychiatry, Rua Dr. Ovidio Pires de Campos 785, 3th floor CEP 05403-010 - São Paulo, SP, Brazil. Phone:+55 11 2661-7267. Fax: +55 11 2661-8010. E-mail: forlenza@usp.br.

## Author Contributions

O.V.F., V.J.R.P. and B.S.O.D. equally contributed to this review article (design, critical review of the literature, and manuscript writing).

## Funding

Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, 2009/52825-8), Associação Beneficente Alzira Denise Hertzog da Silva (ABADHS).

## Notes

The authors declare no competing financial interest.

## ■ REFERENCES

- (1) Cade, J. F. J. (1949) Lithium salts in the treatment of psychotic excitement. *Med. J. Aust.* 36, 349–352.
- (2) Nivoli, A. M., Colom, F., Murru, A., Pacchiarotti, I., Castro-Loli, P., González-Pinto, A., et al. (2011) New treatment guidelines for acute bipolar depression: a systematic review. *J. Affective Disord.* 129 (1–3), 14–26.
- (3) Diniz, B. S., and Teixeira, A. L. (2011) Brain-derived neurotrophic factor and Alzheimer's disease: physiopathology and beyond. *Neuromol. Med.* 13 (4), 217–222.
- (4) Pasquali, L., Busceti, C. L., Fulceri, F., Paparelli, A., and Fornai, F. (2010) Intracellular pathways underlying the effects of lithium. *Behav. Pharmacol.* 21 (5–6), 473–492.
- (5) Birch, N. J. (1974) Letter: lithium and magnesium-dependent enzymes. *Lancet* 2, 965–966.
- (6) Amari, L., Layden, B., Rong, Q., Gerald, C. F., and Mota de Freitas, D. (1999) Comparison of fluorescence,  $^{31}\text{P}$  NMR, and  $^7\text{Li}$  NMR spectroscopic methods for investigating  $\text{Li}^+/\text{Mg}^{2+}$  competition for biomolecules. *Anal. Biochem.* 272, 1–7.
- (7) Grimes, C. A., and Jope, R. S. (2001) The multifaceted roles of glycogen synthase kinase  $3\beta$  in cellular signaling. *Prog. Neurobiol.* 65 (4), 391–426.
- (8) Lee, F. H., Kaidanovich-Belil, O., Roder, J. C., Woodgett, J. R., and Wong, A. H. (2011) Genetic inactivation of GSK3 $\alpha$  rescues spine deficits in Disc1-L100P mutant mice. *Schizophr. Res.* 129 (1), 74–79.
- (9) Phiel, C. J., Wilson, C. A., Lee, V. M., and Klein, P. S. (2003) GSK-3 $\alpha$  regulates production of Alzheimer's disease amyloid- $\beta$  peptides. *Nature* 423 (6938), 435–439.
- (10) Hooper, C., Killick, R., and Lovestone, S. (2008) The GSK3 hypothesis of Alzheimer's disease. *J. Neurochem.* 104 (6), 1433–1439.
- (11) Forlenza, O. V., Diniz, B. S., Radanovic, M., Santos, F. S., Talib, L. L., and Gattaz, W. F. (2011) Disease-modifying properties of long-term lithium treatment for amnesic mild cognitive impairment: randomised controlled trial. *Br J. Psychiatry* 198 (5), 351–356.
- (12) Klein, P. S., and Melton, D. A. (1996) A molecular mechanism for the effect of lithium on development. *Proc. Natl. Acad. Sci. U.S.A.* 93 (16), 8455–8459.
- (13) Ryves, W. J., and Harwood, A. J. (2001) Lithium inhibits glycogen synthase kinase-3 by competition for magnesium. *Biochem. Biophys. Res. Commun.* 280, 720–725.
- (14) Chalecka-Franaszek, E., and Chuang, D. M. (1999) Lithium activates the serine/threonine kinase Akt-1 and suppresses glutamate-induced inhibition of Akt-1 activity in neurons. *Proc. Natl. Acad. Sci. U.S.A.* 96 (15), 8745–8750.
- (15) O'Brien, W. T., Huang, J., Buccafusca, R., Garskof, J., Valvezan, A. J., Berry, G. T., and Klein, P. S. (2011) Glycogen synthase kinase-3 is essential for  $\beta$ -arrestin-2 complex formation and lithium-sensitive behaviors in mice. *J. Clin. Invest.* 121 (9), 3756–3762.
- (16) Pan, J. Q., Lewis, M. C., Ketterman, J. K., Clore, E. L., Riley, M., Richards, K. R., Berry-Scott, E., Liu, X., Wagner, F. F., Holson, E. B., Neve, R. L., Biechele, T. L., Moon, R. T., Scolnick, E. M., Petryshen, T. L., and Haggarty, S. J. (2011) AKT kinase activity is required for lithium to modulate mood-related behaviors in mice. *Neuropsychopharmacology* 36 (7), 1397–1411.
- (17) Mendes, C. T., Mury, F. B., de Sá Moreira, E., Alberto, F. L., Forlenza, O. V., Dias-Neto, E., and Gattaz, W. F. (2009) Lithium reduces Gsk3b mRNA levels: implications for Alzheimer Disease. *Eur. Arch. Psychiatry Clin. Neurosci.* 259 (1), 16–22.
- (18) Patel, S., Yenush, L., Rodríguez, P. L., Serrano, R., and Blundell, T. L. (2002) Crystal structure of an enzyme displaying both inositol-polyphosphate-1-phosphatase and 3'-phosphoadenosine-5'-phosphate phosphatase activities: a novel target of lithium therapy. *J. Mol. Biol.* 315 (4), 677–685.
- (19) Garcia-Arencibia, M., Hochfeld, W., Toh, P., and Rubinsztein, D. C. (2010) Autophagy, a guardian against neurodegeneration. *Semin. Cell Dev. Biol.* 7, 691–698.
- (20) Sarkar, S., Floto, R. A., Berger, Z., Imarisio, S., Cordenier, A., Pasco, M., Cook, L. J., and Rubinsztein, D. C. (2005) Lithium induces autophagy by inhibiting inositol monophosphatase. *J. Cell Biol.* 170 (7), 1101–1111.
- (21) Sarkar, S., Perlstein, E. O., Imarisio, S., Pineau, S., Cordenier, A., Maglathlin, R. L., Webster, J. A., Lewis, T. A., O'Kane, C. J., Schreiber, S. L., and Rubinsztein, D. C. (2007) Small molecules enhance autophagy and reduce toxicity in Huntington's disease models. *Nat. Chem. Biol.* 3 (6), 331–338.
- (22) Chiu, C. T., and Chuang, D. M. (2010) Molecular actions and therapeutic potential of lithium in preclinical and clinical studies of CNS disorders. *Pharmacol. Ther.* 128 (2), 281–304.
- (23) Kang, H. J., Noh, J. S., Bae, Y. S., and Gwag, B. J. (2003) Calcium-dependent prevention of neuronal apoptosis by lithium ion: essential role of phosphoinositide 3-kinase and phospholipase *Cy*. *Mol. Pharmacol.* 64, 228–234.
- (24) Sasaki, T., Han, F., Shioda, N., Moriguchi, S., Kasahara, J., Ishiguro, K., and Fukunaga, K. (2006) Lithium-induced activation of Akt and CaM kinase II contributes to its neuroprotective action in a rat microsphere embolism model. *Brain Res.* 1108, 98–106.
- (25) Hooper, C., Killick, R., and Lovestone, S. (2008) The GSK3 hypothesis of Alzheimer's disease. *J. Neurochem.* 104 (6), 1433–1439.
- (26) Lovestone, S., Davis, D. R., Webster, M. T., Kaech, S., Brion, J. P., Matus, and Anderton, B. H. (1999) Lithium reduces tau phosphorylation: effects in living cells and in neurons at therapeutic concentrations. *Biol. Psychiatry* 45 (8), 995–1003.
- (27) Takahashi, M., Yasutake, K., and Tomizawa, K. (1999) Lithium inhibits neurite growth and tau protein kinase I/glycogen synthase kinase-3 $\beta$ -dependent phosphorylation of juvenile tau in cultured hippocampal neurons. *J. Neurochem.* 73 (5), 2073–2083.
- (28) Fu, Z. Q., Yang, Y., Song, J., Jiang, Q., Lin, Z. C., Wang, Q., and Tian, Q. (2010) LiCl attenuates thapsigargin-induced tau hyperphosphorylation by inhibiting GSK-3 $\beta$  in vivo and in vitro. *J. Alzheimer's Dis.* 21 (4), 1107–1117.
- (29) Esselmann, H., Maler, J. M., Kunz, N., Otto, M., Paul, S., Lewczuk, P., Rütger, E., Kornhuber, J., and Wiltfang, J. (2004) Lithium decreases secretion of A $\beta$ 1–42 and C-truncated species A $\beta$ 1–37/38/39/40 in chicken telencephalic cultures but specifically increases intracellular A $\beta$ 1–38. *Neurodegener. Dis.* 1 (4–5), 236–241.
- (30) Phiel, C. J., Wilson, C. A., Lee, V. M., and Klein, P. S. (2003) GSK-3 $\alpha$  regulates production of Alzheimer's disease amyloid- $\beta$  peptides. *Nature* 423 (6938), 435–439.
- (31) Wei, H., Leeds, P. R., Qian, Y., Wei, W., Chen, R., and Chuang, D. (2000)  $\beta$ -Amyloid peptide-induced death of PC 12 cells and cerebellar granule cell neurons is inhibited by long-term lithium treatment. *Eur. J. Pharmacol.* 392 (3), 117–123.
- (32) Engel, T., Goñi-Oliver, P., Lucas, J. J., Avila, J., and Hernández, F. (2006) Chronic lithium administration to FTDP-17 tau and GSK-3 $\beta$  overexpressing mice prevents tau hyperphosphorylation and neurofibrillary tangle formation, but pre-formed neurofibrillary tangles do not revert. *J. Neurochem.* 99 (6), 1445–1455.

- (33) Leroy, K., Ando, K., Héraud, C., Yilmaz, Z., Authélet, M., Boeynaems, J. M., Buée, L., De Decker, R., and Brion, J. P. (2010) Lithium treatment arrests the development of neurofibrillary tangles in mutant tau transgenic mice with advanced neurofibrillary pathology. *J. Alzheimer's Dis* 19 (2), 705–719.
- (34) Noble, W., Planel, E., Zehr, C., Olm, V., Meyerson, J., Suleman, F., Gaynor, K., Wang, L., LaFrancois, J., Feinstein, B., Burns, M., Krishnamurthy, P., Wen, Y., Bhat, R., Lewis, J., Dickson, D., and Duff, K. (2005) Inhibition of glycogen synthase kinase-3 by lithium correlates with reduced tauopathy and degeneration in vivo. *Proc. Natl. Acad. Sci. U.S.A.* 102 (19), 6990–6995.
- (35) Rockenstein, E., Torrance, M., Adame, A., Mante, M., Bar-on, P., Rose, J. B., Crews, L., and Masliah, E. (2007) Neuroprotective effects of regulators of the glycogen synthase kinase-3 $\beta$  signaling pathway in a transgenic model of Alzheimer's disease are associated with reduced amyloid precursor protein phosphorylation. *J. Neurosci.* 27 (8), 1981–1991.
- (36) Su, Y., Ryder, J., Li, B., Wu, X., Fox, N., Solenberg, P., Brune, K., Paul, S., Zhou, Y., Liu, F., and N. B. (2004) Lithium, a common drug for bipolar disorder treatment, regulates amyloid- $\beta$  precursor protein processing. *Biochemistry* 43, 6899–6908.
- (37) Yu, F., Zhang, Y., and Chuang, D. M. (2012) Lithium reduces BACE1 overexpression,  $\beta$  amyloid accumulation, and spatial learning deficits in mice with traumatic brain injury. *J. Neurotrauma* 29 (13), 2342–2351.
- (38) Zhang, X., Heng, X., Li, T., Li, L., Yang, D., Zhang, X., Du, Y., Doody, R. S., and Le, W. (2011) Long-term treatment with lithium alleviates memory deficits and reduces amyloid- $\beta$  production in an aged Alzheimer's disease transgenic mouse model. *J. Alzheimer's Dis* 24 (4), 739–749.
- (39) Fiorentini, A., Rosi, M. C., Grossi, C., Luccarini, I., and Casamenti, F. (2010) Lithium improves hippocampal neurogenesis, neuropathology and cognitive functions in APP mutant mice. *PLoS One* 5 (12), e14382.
- (40) Alvarez, G., Muñoz-Montano, J. R., Satrustegui, J., Avila, J., Bogónez, E., and Díaz-Nido, J. (1999) Lithium protects cultured neurons against  $\beta$ -amyloid-induced neurodegeneration. *FEBS Lett.* 453 (3), 260–264.
- (41) Alvarez, G., Muñoz-Montano, J. R., Satrustegui, J., Avila, J., Bogónez, E., and Díaz-Nido, J. (2002) Regulation of tau phosphorylation and protection against  $\beta$ -amyloid-induced neurodegeneration by lithium. Possible implications for Alzheimer's disease. *Bipolar Disord.* 4 (3), 153–165.
- (42) Chen, R. W., and Chuang, D. M. (1999) Long term lithium treatment suppresses p53 and Bax expression but increases Bcl-2 expression. A prominent role in neuroprotection against excitotoxicity. *J. Biol. Chem.* 274 (10), 6039–6042.
- (43) Chen, G., Zeng, W. Z., Yuan, P. X., Huang, L. D., Jiang, Y. M., Zhao, Z. H., and Manji, H. K. (1999) The mood-stabilizing agents lithium and valproate robustly increase the levels of the neuroprotective protein bcl-2 in the CNS. *J. Neurochem.* 72 (2), 879–882.
- (44) Hashimoto, R., Senatorov, V., Kanai, H., Leeds, P., and Chuang, D. M. (2003) Lithium stimulates progenitor proliferation in cultured brain neurons. *Neuroscience* 117 (1), 55–61.
- (45) Kim, J. S., Chang, M. Y., Yu, I. T., Kim, J. H., Lee, S. H., Lee, Y. S., and Son, H. (2004) Lithium selectively increases neuronal differentiation of hippocampal neural progenitor cells both in vitro and in vivo. *J. Neurochem.* 89 (2), 324–336.
- (46) Chen, C. L., Lin, C. F., Chiang, C. W., Jan, M. S., and Lin, Y. S. (2006) Lithium inhibits ceramide- and etoposide-induced protein phosphatase 2A methylation, Bcl-2 dephosphorylation, caspase-2 activation, and apoptosis. *Mol. Pharmacol.* 70 (2), 510–517.
- (47) Ghribi, O., Herman, M. M., Spaulding, N. K., and Savory, J. (2002) Lithium inhibits aluminum-induced apoptosis in rabbit hippocampus, by preventing cytochrome c translocation, Bcl-2 decrease, Bax elevation and caspase-3 activation. *J. Neurochem.* 82 (1), 137–145.
- (48) Chou, C. H., Chou, A. K., Lin, C. C., Chen, W. J., Wei, C. C., Yang, M. C., Hsu, C. M., Lung, F. W., Loh, J. K., Howng, S. L., and Hong, Y. R. (2012) GSK3 $\beta$  regulates Bcl2L12 and Bcl2L12A anti-apoptosis signaling in glioblastoma and is inhibited by LiCl. *Cell Cycle* 11 (3), 532–542.
- (49) Shalbuyeva, N., Brustovetsky, T., and Brustovetsky, N. (2007) Lithium desensitizes brain mitochondria to calcium, antagonizes permeability transition, and diminishes cytochrome C release. *J. Biol. Chem.* 282 (25), 18057–18068.
- (50) Bachmann, R. F., Wang, Y., Yuan, P., Zhou, R., Li, X., Alesci, S., Du, J., and Manji, H. K. (2009) Common effects of lithium and valproate on mitochondrial functions: protection against methamphetamine-induced mitochondrial damage. *Int. J. Neuropsychopharmacol.* 12 (6), 805–822.
- (51) Quiroz, J. A., Machado-Vieira, R., Zarate, C. A., Jr., and Manji, H. K. (2010) Novel insights into lithium's mechanism of action: neurotrophic and neuroprotective effects. *Neuropsychobiology* 62 (1), 50–60.
- (52) Bosche, B., Schäfer, M., Graf, R., Härtel, F. V., Schäfer, U., and Noll, T. (2013) Lithium prevents early cytosolic calcium increase and secondary injurious calcium overload in glycolytically inhibited endothelial cells. *Biochem. Biophys. Res. Commun.* 434 (2), 268–272.
- (53) Ngok-Ngam, P., Watcharavit, P., Thiantanawat, A., and Satayavivad, J. (2013) Pharmacological inhibition of GSK3 attenuates DNA damage-induced apoptosis via reduction of p53 mitochondrial translocation and Bax oligomerization in neuroblastoma SH-SY5Y cells. *Cell Mol. Biol. Lett.* 18 (1), 58–74.
- (54) Feier, G., Valvassori, S. S., Varela, R. B., Resende, W. R., Bavaresco, D. V., Morais, M. O., Scaini, G., Andersen, M. L., Streck, E. L., and Quevedo, J. (2013) Lithium and valproate modulate energy metabolism in an animal model of mania induced by methamphetamine. *Pharmacol., Biochem. Behav.* 103 (3), 589–596.
- (55) Li, Q., Li, H., Roughton, K., Wang, X., Kroemer, G., Blomgren, K., and Zhu, C. (2010) Lithium reduces apoptosis and autophagy after neonatal hypoxia-ischemia. *Cell Death Dis.* 1, e56.
- (56) Chang, J. W., Choi, H., Cotman, S. L., and Jung, Y. K. (2011) Lithium rescues the impaired autophagy process in CbCln3( $\Delta$ ex7/8/ $\Delta$ ex7/8) cerebellar cells and reduces neuronal vulnerability to cell death via IMPase inhibition. *J. Neurochem.* 116 (4), 659–668.
- (57) Nahman, S., Belmaker, R. H., and Azab, A. N. (2012) Effects of lithium on lipopolysaccharide-induced inflammation in rat primary glia cells. *Innate Immun.* 18 (3), 447458.
- (58) Li, H., Li, Q., Du, X., Sun, Y., Wang, X., Kroemer, G., Blomgren, K., and Zhu, C. (2011) Lithium-mediated long-term neuroprotection in neonatal rat hypoxia-ischemia is associated with antiinflammatory effects and enhanced proliferation and survival of neural stem/progenitor cells. *J. Cereb. Blood Flow Metab.* No. 10, 2106–2115.
- (59) Basselin, M., Villacreses, N. E., Lee, H. J., Bell, J. M., and Rapoport, S. I. (2007) Chronic lithium administration attenuates up-regulated brain arachidonic acid metabolism in a rat model of neuroinflammation. *J. Neurochem.* 102 (3), 761–772.
- (60) Schwartz, M., Kipnis, J., Rivest, S., and Prat, A. (2013) How do immune cells support and shape the brain in health, disease, and aging? *J. Neurosci.* 33 (45), 17587–17596.
- (61) Nivoli, A. M., Colom, F., Murru, A., Pacchiarotti, I., Castro-Loli, P., and González-Pinto, A. (2011) New treatment guidelines for acute bipolar depression: a systematic review. *J. Affective Disord.* 129 (1–3), 14–26.
- (62) Petronis, A. (2010) Epigenetics as a unifying principle in the aetiology of complex traits and diseases. *Nature.* Jun 10 465 (7299), 721–7.
- (63) Sugawara, H., Iwamoto, K., Bundo, M., Ueda, J., Miyauchi, T., Komori, A., Kazuno, A., Adati, N., Kusumi, I., Okazaki, Y., Ishigooka, J., Kojima, T., and Kato, T. (2011) Hypermethylation of serotonin transporter gene in bipolar disorder detected by epigenome analysis of discordant monozygotic twins. *Transl. Psychiatry* 1, e25.
- (64) Rao, J. S., Keleshian, V. L., Klein, S., and Rapoport, S. I. (2012) Epigenetic modifications in frontal cortex from Alzheimer's disease and bipolar disorder patients. *Transl. Psychiatry* 2, e132.
- (65) Sabuncian, S., Aryee, M. J., Irizarry, R. A., Rongione, M., Webster, M. J., Kaufman, W. E., Murakami, P., Lessard, A., Yolken, R.

- H., Feinberg, A. P., and Potash, J. B. (2012) Genome-wide DNA methylation scan in major depressive disorder. *PLoS One* 7, 34451.
- (66) Rao, J. S., Keleshian, V. L., Klein, S., and Rapoport, S. I. (2012) Epigenetic modifications in frontal cortex from Alzheimer's disease and bipolar disorder patients. *Transl. Psychiatry* 2, e132.
- (67) Popkie, A. P., Zeidner, L. C., Albrecht, A. M., D'Ippolito, A., Eckardt, S., Newsom, D. E., Groden, J., Doble, B. W., Aronow, B., McLaughlin, K. J., White, P., and Phiel, C. J. (2010) Phosphatidylinositol 3-kinase (PI3K) signaling via glycogen synthase kinase-3 (Gsk-3) regulates DNA methylation of imprinted loci. *J. Biol. Chem.* 285 (53), 41337–41347.
- (68) Kessing, L. V., Søndergård, L., Forman, J. L., and Andersen, P.K. (2008) Lithium treatment and risk of dementia. *Arch. Gen. Psychiatry*. 65 (11), 1331–1335.
- (69) Kessing, L. V., Forman, J. L., and Andersen, P. K. (2010) Does lithium protect against dementia? *Bipolar Disord.* 12 (1), 87–94.
- (70) Terao, T., Nakano, H., Inoue, Y., Okamoto, T., Nakamura, J., and Iwata, N. (2006) Lithium and dementia: a preliminary study. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 30 (6), 1125–1128.
- (71) Nunes, P. V., Forlenza, O. V., and Gattaz, W. F. (2007) Lithium and risk for Alzheimer's disease in elderly patients with bipolar disorder. *Br. J. Psychiatry*. 190, 359–360.
- (72) Nitrini, R., Caramelli, P., Herrera, E., Jr., Bahia, V. S., Caixeta, L. F., Radanovic, M., Anghinah, R., Charchat-Fichman, H., Porto, C. S., Carthery, M. T., Hartmann, A. P., Huang, N., Smid, J., Lima, E. P., Takada, L. T., and Takahashi, D. Y. (2004) Incidence of dementia in a community-dwelling Brazilian population. *Alzheimer Dis. Assoc. Disord.* 18 (4), 241–246.
- (73) Rybakowski, J. K., and Suwalska, A. (2010) Excellent lithium responders have normal cognitive functions and plasma BDNF levels. *Int. J. Neuropsychopharmacol.* 13 (5), 617–622.
- (74) Machado-Vieira, R., Manji, H. K., and Zarate, C. A., Jr. (2009) The role of lithium in the treatment of bipolar disorder: convergent evidence for neurotrophic effects as a unifying hypothesis. *Bipolar Disord.* 11 (2), 92–109.
- (75) de Sousa, R. T., van de Bilt, M. T., Diniz, B. S., Ladeira, R. B., Portela, L. V., Souza, D. O., Forlenza, O. V., Gattaz, W. F., and Machado-Vieira, R. (2011) Lithium increases plasma brain-derived neurotrophic factor in acute bipolar mania: a preliminary 4-week study. *Neurosci. Lett.* 494 (1), 54–56.
- (76) Suwalska, A., Sobieska, M., and Rybakowski, J. K. (2010) Serum brain-derived neurotrophic factor in euthymic bipolar patients on prophylactic lithium therapy. *Neuropsychobiology* 62 (4), 229–234.
- (77) Machado-Vieira, R., and Manji, H. K. (2007) The role of lithium in the treatment of bipolar disorder: convergent evidence for neurotrophic effects as a unifying hypothesis. *Bipolar Disord.* 11 (2), 92–109.
- (78) Khairova, R., Pawar, R., Salvatore, G., Juruena, M. F., de Sousa, R. T., Soeiro-de-Souza, M. G., Salvador, M., Zarate, C. A., Gattaz, W. F., and Machado-Vieira, R. (2012) Effects of lithium on oxidative stress parameters in healthy subjects. *Mol. Med. Rep.* 5 (3), 680–682.
- (79) de Sousa, R. T., Zarate, C. A., Jr., Zanetti, M. V., Costa, A. C., Talib, L. L., Gattaz, W. F., and Machado-Vieira, R. (2014) Oxidative stress in early stage Bipolar Disorder and the association with response to lithium. *J. Psychiatr. Res.* 50, 36–41.
- (80) Guloksuz, S., Altinbas, K., Aktas, C. E., Kenis, G., Bilgic, Gazioglu, S., Deniz, G., Oral, E. T., and van Os, J. (2012) Evidence for an association between tumor necrosis factor- $\alpha$  levels and lithium response. *J. Affective Disord.* 143 (1–3), 148–152.
- (81) Knijff, E. M., Breunis, M. N., Kupka, R. W., de Wit, H. J., Ruwhof, C., Akkerhuis, G. W., Nolen, W. A., and Drexhage, H. A. (2007) An imbalance in the production of IL-1 $\beta$  and IL-6 by monocytes of bipolar patients: restoration by lithium treatment. *Bipolar Disord.* 9 (7), 743–753.
- (82) Bearden, C. E., Thompson, P. M., Dalwani, M., Hayashi, K. M., Lee, A. D., Nicoletti, M., Trakhtenbroit, M., Glahn, D. C., Brambilla, P., Sassi, R. B., Mallinger, A. G., Frank, E., Kupfer, D. J., and Soares, J. C. (2007) Greater cortical gray matter density in lithium-treated patients with bipolar disorder. *Biol. Psychiatry* 62 (1), 7–16.
- (83) Moore, G. J., Cortese, B. M., Glitz, D. A., Zajac-Benitez, C., Quiroz, J. A., Uhlde, T. W., Drevets, W. C., and Manji, H. K. (2009) A longitudinal study of the effects of lithium treatment on prefrontal and subgenual prefrontal gray matter volume in treatment-responsive bipolar disorder patients. *J. Clin. Psychiatry* 70 (5), 699–705.
- (84) Lyoo, I. K., Dager, S. R., Kim, J. E., Yoon, S. J., Friedman, S. D., Dunner, D. L., and Renshaw, P. F. (2010) Lithium-induced gray matter volume increase as a neural correlate of treatment response in bipolar disorder: a longitudinal brain imaging study. *Neuropsychopharmacology* 35 (8), 1743–1750.
- (85) van Erp, T. G., Thompson, P. M., Kieseppä, T., Bearden, C. E., Marino, A. C., Hoftman, G. D., Haukka, J., Partonen, T., Huttunen, M., Kaprio, J., Lonnqvist, J., Poutanen, V. P., Toga, A. W., and Cannon, T. D. (2012) Hippocampal morphology in lithium and non-lithium-treated bipolar I disorder patients, non-bipolar co-twins, and control twins. *Hum. Brain Mapp.* 33 (3), 501–510.
- (86) Hajek, T., Calkin, C., Blagdon, R., Slaney, C., and Alda, M. (2013) Type 2 Diabetes Mellitus: A Potentially Modifiable Risk Factor for Neurochemical Brain Changes in Bipolar Disorders. *Biol. Psychiatry*, DOI: 10.1016/j.biopsych.2013.11.007.
- (87) Forester, B. P., Finn, C. T., Berlow, Y. A., Wardrop, M., Renshaw, P. F., and Moore, C. M. (2008) Brain lithium, N-acetyl aspartate and myo-inositol levels in older adults with bipolar disorder treated with lithium: a lithium-7 and proton magnetic resonance spectroscopy study. *Bipolar Disord.* 10 (6), 691–700.
- (88) Silverstone, P. H., Wu, R. H., O'Donnell, T., Ulrich, M., Asghar, S. J., and Hanstock, C. C. (2003) Chronic treatment with lithium, but not sodium valproate, increases cortical N-acetyl-aspartate concentrations in euthymic bipolar patients. *Int. Clin. Psychopharmacol.* 18 (2), 73–79.
- (89) Macdonald, A., Briggs, K., Poppe, M., Higgins, A., Velayudhan, L., and Lovestone, S. (2008) A feasibility and tolerability study of lithium in Alzheimer's disease. *Int. J. Geriatr. Psychiatry* 23 (7), 704–711.
- (90) Hampel, H., Ewers, M., Bürger, K., Annas, P., Mörtberg, A., Bogstedt, A., Frölich, L., Schröder, J., Schönknecht, P., Riepe, M. W., Kraft, I., Gasser, T., Leyhe, T., Möller, H. J., Kurz, A., and Basun, H. (2009) Lithium trial in Alzheimer's disease: a randomized, single-blind, placebo-controlled, multicenter 10-week study. *J. Clin. Psychiatry* 70 (6), 922–931.
- (91) Leyhe, T., Eschweiler, G. W., Stransky, E., Gasser, T., Annas, P., Basun, H., and Laske, C. (2009) Increase of BDNF serum concentration in lithium treated patients with early Alzheimer's disease. *J. Alzheimer's Dis.* 16 (3), 649–656.
- (92) Straten, G., Saur, R., Laske, C., Gasser, T., Annas, P., Basun, H., and Leyhe, T. (2011) Influence of lithium treatment on GDNF serum and CSF concentrations in patients with early Alzheimer's disease. *Curr. Alzheimer Res.* 8 (8), 853–859.
- (93) Yang, J., Takahashi, Y., Cheng, E., Liu, J., Terranova, P. F., Zhao, B., Thrasher, J. B., Wang, H. G., and Li, B. (2010) GSK-3 $\beta$  promotes cell survival by modulating Bif-1-dependent autophagy and cell death. *J. Cell Sci.* 123 (6), 861–870.
- (94) Gordon, P. H. (2011) Amyotrophic lateral sclerosis: pathophysiology, diagnosis and management. *CNS Drugs* 25 (1), 1–15.
- (95) Dill, J., Wang, H., Zhou, F., and Li, S. (2008) Inactivation of glycogen synthase kinase 3 promotes axonal growth and recovery in the CNS. *J. Neurosci.* 28, 8914–8928.
- (96) Busceti, C. L., Biagioni, F., Rizzo, B., Battaglia, G., Storto, M., Cinque, C., Molinaro, G., Gradini, R., Caricasole, A., Canudas, A. M., Bruno, V., Nicoletti, F., and Fornai, F. (2008) Enhanced tau phosphorylation in the hippocampus of mice treated with 3,4-methylenedioxymethamphetamine ("Ecstasy"). *J. Neurosci.* 28, 3234–3245.
- (97) Strong, M. J. (2008) The syndromes of frontotemporal dysfunction in amyotrophic lateral sclerosis. *Amyotrophic Lateral Scler.* 9, 323–338.
- (98) Fornai, F., Longone, P., Cafaro, L., Kastsuchenka, O., Ferrucci, M., Manca, M. L., Lazzeri, G., Spalloni, A., Bellio, N., Lenzi, P.,

Modugno, N., Siciliano, G., Isidoro, C., Murri, L., Ruggieri, S., and Paparelli, A. (2008) Lithium delays progression of amyotrophic lateral sclerosis. *Proc. Natl. Acad. Sci. U.S.A.* 105 (6), 2052–2057.

(99) Feng, H. L., Leng, Y., Ma, C. H., Zhang, J., Ren, M., and Chuang, D. M. (2008) Combined lithium and valproate treatment delays disease onset, reduces neurological deficits and prolongs survival in an amyotrophic lateral sclerosis mouse model. *Neuroscience* 155 (3), 567–572.

(100) Chiò, A., and Mora, G. (2013) The final chapter of the ALS lithium saga. *Lancet Neurol.* 12 (4), 324–325.

(101) Morrison, K. E., Dhariwal, S., Hornabrook, R., Savage, L., Burn, D. J., Khoo, T. K., Kelly, J., Murphy, C. L., Al-Chalabi, A., Dougherty, A., Leigh, P. N., Wijesekera, L., Thornhill, M., Ellis, C. M., O'Hanlon, K., Panicker, J., Pate, L., Ray, P., Wyatt, L., Young, C. A., Copeland, L., Ealing, J., Hamdalla, H., Leroi, I., Murphy, C., O'Keefe, F., Oughton, E., Partington, L., Paterson, P., Rog, D., Sathish, A., Sexton, D., Smith, J., Vanek, H., Dodds, S., Williams, T. L., Steen, I. N., Clarke, J., Eziefula, C., Howard, R., Orrell, R., Sidle, K., Sylvester, R., Barrett, W., Merritt, C., Talbot, K., Turner, M. R., Whatley, C., Williams, C., Williams, J., Cosby, C., Hanemann, C. O., Iman, I., Philips, C., Timings, L., Crawford, S. E., Hewamadduma, C., Hibberd, R., Hollinger, H., McDermott, C., Mils, G., Rafiq, M., Shaw, P. J., Taylor, A., Waines, E., Walsh, T., Addison-Jones, R., Birt, J., Hare, M., and Majid, T. (2013) Lithium in patients with amyotrophic lateral sclerosis (LiCALS): a phase 3 multicentre, randomised, double-blind, placebo-controlled trial. *Lancet Neurol.* 12 (4), 339–345.

(102) Feng, H. L., Leng, Y., Ma, C. H., Zhang, J., Ren, M., and Chuang, D. M. (2008) Combined lithium and valproate treatment delays disease onset, reduces neurological deficits and prolongs survival in an amyotrophic lateral sclerosis mouse model. *Neuroscience* 155 (3), 567–572.

(103) Chiu, C.T.1, Liu, G., Leeds, P., and Chuang, D. M. (2011) Combined treatment with the mood stabilizers lithium and valproate produces multiple beneficial effects in transgenic mouse models of Huntington's disease. *Neuropsychopharmacology* 36 (12), 2406–2421.

(104) Yu, F.1, Wang, Z., Tanaka, M., Chiu, C. T., Leeds, P., Zhang, Y., and Chuang, D. M. (2013) Posttrauma cotreatment with lithium and valproate: reduction of lesion volume, attenuation of blood-brain barrier disruption, and improvement in motor coordination in mice with traumatic brain injury. *J. Neurosurg.* 119 (3), 766–773.

(105) Chiò, A., and Mora, G. (2013) The final chapter of the ALS lithium saga. *Lancet Neurol.* 12 (4), 324–325.

(106) Youdim, M. B., and Arraf, Z. (2004) Prevention of MPTP (*N*-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) dopaminergic neurotoxicity in mice by chronic lithium: involvements of Bcl-2 and Bax. *Neuropharmacology* 46 (8), 1130–1140.

(107) Yong, Y., Ding, H., Fan, Z., Luo, J., and Ke, Z. J. (2011) Lithium fails to protect dopaminergic neurons in the 6-OHDA model of Parkinson's disease. *Neurochem. Res.* 36 (3), 367–374.

(108) Senatorov, V. V., Ren, M., Kanai, H., Wei, H., and Chuang, D. M. (2004) Short-term lithium treatment promotes neuronal survival and proliferation in rat striatum infused with quinolinic acid, an excitotoxic model of Huntington's disease. *Mol. Psychiatry* 9 (4), 371–385.

(109) Wei, H., Qin, Z. H., Senatorov, V. V., Wei, W., Wang, Y., Qian, Y., and Chuang, D. M. (2001) Lithium suppresses excitotoxicity-induced striatal lesions in a rat model of Huntington's disease. *Neuroscience* 106 (3), 603–612.

(110) Wood, N. I., and Morton, A. J. (2003) Chronic lithium chloride treatment has variable effects on motor behaviour and survival of mice transgenic for the Huntington's disease mutation. *Brain Res. Bull.* 61 (4), 375–383.

(111) Xilouri, M., Vogiatzi, T., Vekrellis, K., and Stefanis, L. (2008) Alphasynuclein degradation by autophagic pathways: a potential key to Parkinson's disease pathogenesis. *Autophagy* 4, 917–919.

(112) Chen, G., Bower, K. A., Ma, C., Fang, S., Thiele, C. J., and Luo, J. (2004) Glycogen synthase kinase 3 $\beta$  (GSK3 $\beta$ ) mediates 6-hydroxydopamine-induced neuronal death. *FASEB J.* 18, 1162–1164.

(113) Ge, X. H., Zhu, G. J., Geng, D. Q., Zhang, Z. J., and Liu, C. F. (2012) Erythropoietin attenuates 6-hydroxydopamine-induced apoptosis via glycogen synthase kinase 3 $\beta$ -mediated mitochondrial translocation of Bax in PC12 cells. *Neurol. Sci.* 33 (6), 1249–1256.

(114) Wei, H., Qin, Z. H., Senatorov, V. V., Wei, W., Wang, Y., Qian, Y., and Chuang, D. M. (2001) Lithium suppresses excitotoxicity-induced striatal lesions in a rat model of Huntington's disease. *Neuroscience* 106, 603–612.