

Research Article

New Estimates of q_1q_2 -Ostrowski-Type Inequalities within a Class of n -Polynomial Preconvexity of Functions

Humaira Kalsoom ¹, Muhammad Idrees,² Dumitru Baleanu,^{3,4,5} and Yu-Ming Chu ^{6,7}

¹School of Mathematical Sciences, Zhejiang University, Hangzhou 310027, China

²Zhejiang Province Key Laboratory of Quantum Technology and Device, Department of Physics, Zhejiang University, Hangzhou 310027, China

³Department of Mathematics, Çankaya University, 06530 Ankara, Turkey

⁴Institute of Space Sciences, Magurele, Bucharest, Romania

⁵Institute of Space Sciences, PO Box MG-23, Magurele (Ilfov) Bucharest 077123, Romania

⁶Department of Mathematics, Huzhou University, Huzhou 313000, China

⁷Hunan Provincial Key Laboratory of Mathematical Modeling and Analysis in Engineering, Changsha University of Science & Technology, Changsha 410114, China

Correspondence should be addressed to Yu-Ming Chu; chuyuming@zjhu.edu.cn

Received 24 March 2020; Accepted 19 June 2020; Published 24 July 2020

Academic Editor: Serkan Araci

Copyright © 2020 Humaira Kalsoom et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In this article, we develop a novel framework to study for a new class of preinvex functions depending on arbitrary nonnegative function, which is called n -polynomial preinvex functions. We use the n -polynomial preinvex functions to develop q_1q_2 -analogues of the Ostrowski-type integral inequalities on coordinates. Different features and properties of excitement for quantum calculus have been examined through a systematic way. We are discussing about the suggestions and different results of the quantum inequalities of the Ostrowski-type by inferring a new identity for q_1q_2 -differentiable function. However, the problem has been proven to utilize the obtained identity, we give q_1q_2 -analogues of the Ostrowski-type integrals inequalities which are connected with the n -polynomial preinvex functions on coordinates. Our results are the generalizations of the results in earlier papers.

1. Introduction

Calculus is an imperative study of the derivatives and integrals. The classical derivative was convoluted with the strength regulation kind kernel, and eventually, this gave upward thrust to new calculus referred to as the quantum calculus. In mathematics, quantum calculus (named q -calculus) is the study of calculus without limits. The interest in this subject has exploded, and the q -calculus has in the last twenty years served as a bridge between mathematics and physics. The q -calculus has numerous applications in various fields of mathematics, for example, dynamical systems, number theory, combinatorics, special functions, fractals, and also for scientific problems in some applied areas such as computer science, quantum mechanics, and quantum physics.

Jackson [1] defined the q -analogue of derivative and integral operator as well as provided some of their applications. It is imperative to mention that quantum integral inequalities are more practical and informative than their classical counterparts. It has been mainly due to the fact that quantum integral inequalities can describe the hereditary properties of the processes and phenomena under investigation. Historically, the subject of quantum calculus can be traced back to Euler and Jacobi, but in recent decades, it has experienced a rapid development. As a result, new generalizations of the classical concepts of quantum calculus have been initiated and reviewed in many literature. Tariboon and Ntouyas [2, 3] proposed the quantum calculus concepts on finite intervals and obtained several q -analogues of classical mathematical objects, which inspired many other researchers to study the

subject in depth, and as a consequence, numerous novel results concerning quantum analogues of classical mathematical results have been launched. Noor et al. [4] obtained new q -analogues of inequality utilizing the first-order q -differentiable convex function.

Inequality plays an irreplaceable role in the development of mathematics. Very recently, many new inequalities such as the Hermite-Hadamard-type inequality [5–9], Petrović-type inequality [10], Pólya-Szegő and Čebyšev-type inequalities [11], Ostrowski-type inequality [12], reverse Minkowski inequality [13], Jensen-type inequality [14–16], Bessel function inequality [17], trigonometric and hyperbolic functions inequalities [18], fractional integral inequality [19–22], complete and generalized elliptic integrals inequalities [23–28], generalized convex function inequality [29–31], and mean values inequality [32–34] have been discovered by many researchers. While the concept of classical convexity has been brought into a streamline by mathematical inequalities [35–50]. In fact, convex function and its connection with mathematical inequalities have wide applications in the estimation of some parameters in scientific observations and calculations [51–65]. In recent years, the classical concept of convexity has been extended and generalized in different directions, one of the important generalization of convexity is the invexity, which was studied by Hanson [66]; this work has greatly expanded the role of invexity in optimization. In [67, 68], the authors introduced a class of functions, which is called preinvexity as a generalization of convex functions.

Now, we recall the classical and well-known Hermite-Hadamard inequality [69], which can be stated as

$$\Phi\left(\frac{\xi_1 + \xi_2}{2}\right) \leq \frac{1}{\xi_2 - \xi_1} \int_{\xi_1}^{\xi_2} \Phi(\gamma) d\gamma \leq \frac{\Phi(\xi_1) + \Phi(\xi_2)}{2}, \quad (1)$$

for all $\xi_1, \xi_2 \in I$ if $\Phi : I \rightarrow \mathbb{R}$ is a convex function.

Ostrowski [70] established an integral inequality for continuous and differentiable function as follows.

Theorem 1 (See [70]). *Let $\Phi : [\xi_1, \xi_2] \rightarrow \mathbb{R}$ be continuous and differentiable on (ξ_1, ξ_2) such that $|\Phi'(\gamma)| \leq M$ for all $\gamma \in (\xi_1, \xi_2)$. Then, one has*

$$\left| \Phi(Q) - \frac{1}{\xi_2 - \xi_1} \int_{\xi_1}^{\xi_2} \Phi(\gamma) d\gamma \right| \leq \left[\frac{1}{4} + \frac{(Q - \xi_1 + \xi_2/2)^2}{(\xi_2 - \xi_1)^2} \right] (\xi_2 - \xi_1) M, \quad (2)$$

for all $Q \in [\xi_1, \xi_2]$ with the best possible constant $1/4$.

The inequality (2) can be described in an identical kind as

$$\left| \Phi(Q) - \frac{1}{\xi_2 - \xi_1} \int_{\xi_1}^{\xi_2} \Phi(\gamma) d\gamma \right| \leq \frac{M}{\xi_2 - \xi_1} \left[\frac{(Q - \xi_1)^2 + (Q)^2}{2} \right]. \quad (3)$$

Latif et al. [71] generalized the Ostrowski inequality (2) to the coordinated convex function by establishing an identity as follows.

Theorem 2 (See [71]). *Let $\xi_1 < \xi_2, \xi_3 < \xi_4$ and $\Phi : [\xi_1, \xi_2] \times [\xi_3, \xi_4] \rightarrow \mathbb{R}$ be continuous and differentiable on $(\xi_1, \xi_2) \times (\xi_3, \xi_4)$ such that $\partial^2 \Phi / \partial z \partial w \in L([\xi_1, \xi_2] \times [\xi_3, \xi_4])$.*

Then the identity

$$\begin{aligned} & \Phi(Q, \rho) + \frac{1}{(\xi_2 - \xi_1)(\xi_4 - \xi_3)} \int_{\xi_1}^{\xi_2} \int_{\xi_3}^{\xi_4} \Phi(u, v) dudv - \rho \\ &= \frac{(Q - \xi_1)^2 (\rho - \xi_3)^2}{(\xi_2 - \xi_1)(\xi_4 - \xi_3)} \int_0^1 \int_0^1 zw \frac{\partial^2}{\partial z \partial w} \Phi \\ & \cdot (zQ + (1-z)\xi_1, w\rho + (1-w)\xi_3) dzdw \\ & - \frac{(Q - \xi_1)^2 (\xi_4 - \rho)^2}{(\xi_2 - \xi_1)(\xi_4 - \xi_3)} \int_0^1 \int_0^1 zw \frac{\partial^2}{\partial z \partial w} \Phi \\ & \cdot (zQ + (1-z)\xi_1, w\rho + (1-w)\xi_4) dzdw \\ & - \frac{(\xi_2 - Q)^2 (\rho - \xi_3)^2}{(\xi_2 - \xi_1)(\xi_4 - \xi_3)} \int_0^1 \int_0^1 zw \frac{\partial^2}{\partial z \partial w} \Phi \\ & \cdot (zQ + (1-z)\xi_2, w\rho + (1-w)\xi_3) dzdw \\ & + \frac{(\xi_2 - Q)^2 (\xi_4 - \rho)^2}{(\xi_2 - \xi_1)(\xi_4 - \xi_3)} \int_0^1 \int_0^1 zw \frac{\partial^2}{\partial z \partial w} \Phi \\ & \cdot (zQ + (1-z)\xi_2, w\rho + (1-w)\xi_4) dzdw, \end{aligned} \quad (4)$$

holds for all $(Q, \rho) \in [\xi_1, \xi_2] \times [\xi_3, \xi_4]$, where

$$y = \frac{1}{\xi_4 - \xi_3} \int_{\xi_3}^{\xi_4} \Phi(Q, v) dv + \frac{1}{\xi_2 - \xi_1} \int_{\xi_1}^{\xi_2} \Phi(u, \rho) du. \quad (5)$$

Noor et al. [4] presented the Ostrowski-type inequality for quantum calculus.

Theorem 3 (See [4]). *Let $q \in (0, 1), \xi_1 < \xi_2$ and $\Phi[\xi_1, \xi_2] \rightarrow \mathbb{R}$ be continuous such that ${}_{\xi_1} D_q \Phi$ is integrable on (ξ_1, ξ_2) . Then*

$$\begin{aligned} & \Phi(Q) - \frac{1}{\xi_2 - \xi_1} \int_{\xi_1}^{\xi_2} \Phi(u) {}_{\xi_1} d_q u = \frac{q(Q - \xi_1)^2}{\xi_2 - \xi_1} \int_0^1 z {}_{\xi_1} D_q \Phi \\ & \cdot ((1-z)\xi_1 + zQ) {}_0 d_q z + \frac{q(\xi_2 - Q)^2}{\xi_2 - \xi_1} \int_0^1 z {}_{\xi_1} D_q \Phi \\ & \cdot ((1-z)\xi_2 + zQ) {}_0 d_q z. \end{aligned} \quad (6)$$

The following quantum integral version of the Hermite-Hadamard-type inequality for the coordinated convex function was proved by Alp and Sarikaya [72].

Theorem 4 (See [72]). *Let $q_1, q_2 \in (0, 1), \xi_1 < \xi_2, \xi_3 < \xi_4$ and $\Phi : \mathcal{N} = [\xi_1, \xi_2] \times [\xi_3, \xi_4] \rightarrow \mathbb{R}$ be a coordinated convex*

function on \mathcal{N} . Then one has

$$\begin{aligned} & \Phi\left(\frac{q_1\xi_1 + \xi_2}{1 + q_1}, \frac{q_2\xi_3 + \xi_4}{1 + q_2}\right) \\ & \leq \frac{1}{2} \left[\frac{1}{\xi_2 - \xi_1} \int_{\xi_1}^{\xi_2} \Phi\left(z, \frac{q_2\xi_3 + \xi_4}{1 + q_2}\right)_{\xi_1} d_{q_1} z \right. \\ & \quad \left. + \frac{1}{\xi_3 - \xi_4} \int_{\xi_3}^{\xi_4} \Phi\left(\frac{q_1\xi_1 + \xi_2}{1 + q_1}, w\right)_{\xi_3} d_{q_2} w \right] \\ & \leq \frac{1}{(\xi_1 - \xi_2)(\xi_3 - \xi_4)} \int_{\xi_1}^{\xi_2} \int_{\xi_3}^{\xi_4} \Phi(z, w)_{\xi_1} d_{q_1} z_{\xi_3} d_{q_2} w \\ & \leq \frac{1}{4} \left[\frac{1}{(1 + q_1)(\xi_2 - \xi_1)} \left(q_2 \int_{\xi_1}^{\xi_2} \Phi(z, \xi_3)_{\xi_1} d_{q_1} z + \int_{\xi_1}^{\xi_2} \Phi(z, \xi_4)_{\xi_1} d_{q_1} z \right) \right. \\ & \quad \left. + \frac{1}{(1 + q_2)(\xi_3 - \xi_4)} \left(q_1 \int_{\xi_3}^{\xi_4} \Phi(\xi_1, w)_{\xi_3} d_{q_2} w + \int_{\xi_3}^{\xi_4} \Phi(\xi_2, w)_{\xi_3} d_{q_2} w \right) \right] \\ & \leq \frac{q_1 q_2 \Phi(\xi_1, \xi_3) + q_2 \Phi(\xi_2, \xi_3) + q_1 \Phi(\xi_1, \xi_4) + \Phi(\xi_2, \xi_4)}{(1 + q_1)(1 + q_2)}. \end{aligned} \tag{7}$$

Kalsoom et al. [73] found the quantum integral inequality for two parameters function on the finite rectangle.

Next, we present the definitions of $q_1 q_2$ -derivative and integral, and their two known results.

Definition 5. Let $q_1, q_2 \in (0, 1)$, $\xi_1 < \xi_2, \xi_3 < \xi_4$ and $\Phi : [\xi_1, \xi_2] \times [\xi_3, \xi_4] \rightarrow \mathbb{R}$ be a continuous function. Then, the partially q_1 -derivative, q_2 -derivative, and $q_1 q_2$ -derivative at $(z, w) \in [\xi_1, \xi_2] \times [\xi_3, \xi_4]$ for the function Φ are defined by

$$\frac{\xi_1 \partial_{q_1} \Phi(z, w)}{\xi_1 \partial_{q_1} z} = \frac{\Phi(z, w) - \Phi(q_1 z + (1 - q_1)\xi_1, w)}{(1 - q_1)(z - \xi_1)} \quad (z \neq \xi_1),$$

$$\frac{\xi_3 \partial_{q_2} \Phi(z, w)}{\xi_3 \partial_{q_2} w} = \frac{\Phi(z, w) - \Phi(z, q_2 w + (1 - q_2)\xi_3)}{(1 - q_2)(w - \xi_3)} \quad (w \neq \xi_3),$$

$$\begin{aligned} \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(z, w)}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} &= \frac{1}{(1 - q_1)(1 - q_2)(z - \xi_1)(w - \xi_3)} \\ & \times [\Phi(q_1 z + (1 - q_1)\xi_1, q_2 w + (1 - q_2)\xi_3) \\ & - \Phi(q_1 z + (1 - q_1)\xi_1, w) \\ & - \Phi(z, q_2 w + (1 - q_2)\xi_3) + \Phi(z, w)] \\ & \quad (z \neq \xi_1, w \neq \xi_3), \end{aligned} \tag{8}$$

respectively. The function Φ is said to be partially q_1 -, q_2 -, and $q_1 q_2$ -differentiable on $[\xi_1, \xi_2] \times [\xi_3, \xi_4]$ if $\xi_1 \partial_{q_1} \Phi(z, w)/_{\xi_1} \partial_{q_1} z, \xi_3 \partial_{q_2} \Phi(z, w)/_{\xi_3} \partial_{q_2} w$, and $\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(z, w)/_{\xi_1} \partial_{q_1} z_{\xi_3} \partial_{q_2} w$ exist for all $(z, w) \in [\xi_1, \xi_2] \times [\xi_3, \xi_4]$.

Definition 6. Let $q_1, q_2 \in (0, 1)$, $\xi_1 < \xi_2, \xi_3 < \xi_4$ and $\Phi : [\xi_1, \xi_2] \times [\xi_3, \xi_4] \rightarrow \mathbb{R}$ be a continuous function. Then the $q_1 q_2$

-integral of the function Φ on $[\xi_1, \xi_2] \times [\xi_3, \xi_4]$ is defined by

$$\begin{aligned} & \int_{\xi_3}^t \int_{\xi_1}^s \Phi(z, w)_{\xi_1} d_{q_1} z_{\xi_3} d_{q_2} w = (1 - q_1)(1 - q_2)(s - \xi_1)(t - \xi_3) \\ & \quad \times \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \Phi(q_1^n s + (1 - q_1^n)\xi_1, q_2^m t + (1 - q_2^m)\xi_3) \\ & \quad \times \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \Phi(q_1^n s + (1 - q_1^n)\xi_1, q_2^m t + (1 - q_2^m)\xi_3), \end{aligned} \tag{9}$$

for $(s, t) \in [\xi_1, \xi_2] \times [\xi_3, \xi_4]$.

Theorem 7. Let $\xi_1 < \xi_2, \xi_3 < \xi_4$ and $\Phi : [\xi_1, \xi_2] \times [\xi_3, \xi_4] \rightarrow \mathbb{R}$ be a continuous function. Then, we have the identities

$$\frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \int_{\xi_4}^t \int_{\xi_1}^s \Phi(z, w)_{\xi_1} d_{q_1} z_{\xi_3} d_{q_2} w = \Phi(s, t),$$

$$\int_{\xi_3}^t \int_{\xi_1}^s \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(z, w)}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \xi_1 d_{q_1} z_{\xi_3} d_{q_2} w = \Phi(s, t),$$

$$\begin{aligned} & \int_{t_1}^t \int_{s_1}^s \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(z, w)}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \xi_1 d_{q_1} z_{\xi_3} d_{q_2} w = \Phi(s, t) - \Phi(s, t_1) \\ & \quad - \Phi(s_1, t) + \Phi(s_1, t_1), \end{aligned} \tag{10}$$

for $(s_1, t_1) \in (\xi_1, s) \times (\xi_4, t)$.

Theorem 8. Let $a \in \mathbb{R}, \xi_1 < \xi_2, \xi_3 < \xi_4$ and $\Phi_1, \Phi_2 : [\xi_1, \xi_2] \times [\xi_3, \xi_4] \rightarrow \mathbb{R}$ be continuous functions. Then, the identities

$$\begin{aligned} & \int_{\xi_3}^t \int_{\xi_1}^s [\Phi_1(z, w) + \Phi_2(z, w)]_{\xi_1} d_{q_1} z_{\xi_3} d_{q_2} w \\ & = \int_{\xi_3}^t \int_{\xi_1}^s \Phi_1(z, w)_{\xi_1} d_{q_1} z_{\xi_3} d_{q_2} w + \int_{\xi_3}^t \int_{\xi_1}^s \Phi_2(z, w)_{\xi_1} d_{q_1} z_{\xi_3} d_{q_2} w, \\ & \int_{\xi_3}^t \int_{\xi_1}^s a \Phi(z, w)_{\xi_1} d_{q_1} z_{\xi_3} d_{q_2} w = a \int_{\xi_3}^t \int_{\xi_1}^s \Phi(z, w)_{\xi_1} d_{q_1} z_{\xi_3} d_{q_2} w, \end{aligned} \tag{11}$$

holds for $(s, t) \in [\xi_1, \xi_2] \times [\xi_3, \xi_4]$.

Very recently, Toplu et al. [74] improved the Hermite-Hadamard inequality (1) by investigating the n -polynomial convexity. The main purpose of the article is to introduce the notion of n -polynomial preinvex function, provide a new generalized quantum integral identity, establish new quantum analogues of Ostrowski-type inequalities for the n -polynomial preinvex function on coordinates, and generalize and unify the previous known results.

2. Discussions and Main Results

In the beginning of this section, we introduce the definition of n -polynomial preinvexity.

Definition 9 (See [75]). Let $\eta(\cdot, \cdot) : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuous bi-function. Then, $\Omega_\eta \subset \mathbb{R}^n$ is said to be invex if

$$\xi_1 + \gamma\eta(\xi_2, \xi_1) \in \Omega_\eta, \tag{12}$$

for all $\xi_1, \xi_2 \in \Omega_\eta$ and $\gamma \in [0, 1]$.

Definition 10 (See [67]). The function $\Phi : \Omega_\eta \subset \mathbb{R}^n \rightarrow \mathbb{R}$ is said to be preinvex if

$$\Phi(\xi_1 + \gamma\eta(\xi_2, \xi_1)) \leq (1 - \gamma)\Phi(\xi_1) + \gamma\Phi(\xi_2), \tag{13}$$

for all $\xi_1, \xi_2 \in \Omega_\eta$ and $\gamma \in [0, 1]$.

Definition 11. Let $n \in \mathbb{N}$. Then, the nonnegative function $\Phi : \Omega_\eta \subset \mathbb{R}^n \rightarrow \mathbb{R}$ is said to be n -polynomial preinvex if

$$\begin{aligned} \Phi(\xi_1 + \gamma\eta(\xi_2, \xi_1)) &\leq \frac{1}{n} \sum_{p=1}^n [1 - (1 - \gamma)^p] \Phi(\xi_1) \\ &+ \frac{1}{n} \sum_{p=1}^n [1 - (1 - \gamma^p)] \Phi(\xi_2), \end{aligned} \tag{14}$$

for all $\xi_1, \xi_2 \in \Omega_\eta$ and $\gamma \in [0, 1]$.

Note that if $n = 1$, then the definition of n -polynomial preinvex function reduce to the definition of preinvex function.

If we take $n = 2$, then we have 2-polynomial preinvex function inequality

$$\Phi(\xi_1 + \gamma\eta(\xi_2, \xi_1)) \leq \frac{3\gamma - \gamma^2}{2} \Phi(\xi_1) + \frac{1 - \gamma - \gamma^2}{2} \Phi(\xi_2). \tag{15}$$

Proposition 12. Let $\xi_2 > 0$ and $\Phi_\alpha : [\xi_1, \xi_1 + \eta(\xi_2, \xi_1)] \rightarrow \mathbb{R}$ be an arbitrary family of n -polynomial preinvex functions and $\Phi(\xi) = \sup_\alpha \Phi_\alpha(\xi)$. If $J_\eta = \{u \in [\xi_1, \xi_1 + \eta(\xi_2, \xi_1)] : \Phi(u) < 1\}$ is nonempty, then J_η is an interval and Φ is an n -polynomial preinvex function on J_η .

Proof. Let $\gamma \in [0, 1]$ and $\xi_1, \xi_1 + \eta(\xi_2, \xi_1) \in J_\eta$. Then, we have

$$\begin{aligned} \Phi(\xi_1 + \gamma\eta(\xi_2, \xi_1)) &= \sup_\alpha \Phi_\alpha(\xi_1 + \gamma\eta(\xi_2, \xi_1)) \\ &\leq \sup_\alpha \left[\frac{1}{n} \sum_{p=1}^n [1 - (1 - \gamma)^p] \Phi_\alpha(\xi_1) + \frac{1}{n} \sum_{p=1}^n [1 - \gamma]^p \Phi_\alpha(\xi_2) \right] \\ &\leq \frac{1}{n} \sum_{p=1}^n [1 - (1 - \gamma)^p] \sup_\alpha \Phi_\alpha(\xi_1) + \frac{1}{n} \sum_{p=1}^n (1 - \gamma)^p \sup_\alpha \Phi_\alpha(\xi_2) \\ &\leq \frac{1}{n} \sum_{p=1}^n [1 - (1 - \gamma)^p] \Phi(\xi_1) + \frac{1}{n} \sum_{p=1}^n (1 - \gamma)^p \Phi(\xi_2) < \infty. \end{aligned} \tag{16}$$

This completes the proof.

In order to establish new quantum analogues of the Ostrowski-type inequalities on coordinates for the n -polynomial preinvex function, we need a key lemma, which we present in this section.

Lemma 13. Let $q_1, q_2 \in (0, 1)$, $\xi_1 < \xi_2$, $\xi_3 < \xi_4$, $\mathcal{N} = [\xi_1, \xi_2] \times [\xi_3, \xi_4]$, \mathcal{N}° , be the interior of \mathcal{N} , and $\Phi : \mathcal{N} \rightarrow \mathbb{R}$ be mixed partial q_1, q_2 -differentiable on \mathcal{N}° such that $|(\frac{\partial^2}{\partial_{q_1} \partial_{q_2}} \Phi)|_{(\xi_1, \xi_3)}$ is continuous and integrable on $[\xi_1, \xi_1 + \eta_1(\xi_2, \xi_1)] \times [\xi_3, \xi_3 + \eta_2(\xi_4, \xi_3)] \subset \mathcal{N}^\circ$ for $\eta_1(\xi_2, \xi_1), \eta_2(\xi_4, \xi_3) > 0$. Then, we have the identity

$$\begin{aligned} \Omega_{q_1, q_2}(\xi_1, \xi_2, \xi_3, \xi_4)(\Phi) &= -\frac{q_1 q_2 [\eta_1(Q, \xi_1)]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \\ &\times \int_0^1 \int_0^1 z w \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(Q, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\ &- \frac{q_1 q_2 [\eta_1(Q, \xi_1)]^2 [\eta_2(\xi_4, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \\ &\times \int_0^1 \int_0^1 z w \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(Q, \xi_1), \xi_4 + w\eta_2(\rho, \xi_4))_0 d_{q_1} z_0 d_{q_2} w \\ &- \frac{q_1 q_2 [\eta_1(\xi_2, Q)]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \\ &\times \int_0^1 \int_0^1 z w \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(Q, \xi_2), \xi_3 + w\eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\ &- \frac{q_1 q_2 [\eta_1(\xi_2, Q)]^2 [\eta_2(\xi_4, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \\ &\times \int_0^1 \int_0^1 z w \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(Q, \xi_2), \xi_4 + w\eta_2(\rho, \xi_4))_0 d_{q_1} z_0 d_{q_2} w, \end{aligned} \tag{17}$$

where

$$\begin{aligned}
 \Omega_{q_1 q_2}(\xi_1, \xi_2, \xi_3, \xi_4)(\Phi) &= \Phi(\mathbf{Q}, \rho) \\
 &+ \frac{1}{\eta_2(\xi_4, \xi_3)} \left[\int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathbf{Q}, v)_0 d_{q_2} v + \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(\mathbf{Q}, v)_0 d_{q_2} v \right] \\
 &+ \frac{1}{\eta_1(\xi_2, \xi_1)} \left[\int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \Phi(u, \rho)_0 d_{q_1} u + \int_{\xi_2}^{\xi_2 + \eta_1(\mathbf{Q}, \xi_2)} \Phi(u, \rho)_0 d_{q_1} v \right] - \mathcal{Q}, \\
 \mathcal{Q} &= \frac{1}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(u, v)_0 d_{q_1} u d_{q_2} v \\
 &+ \frac{1}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(u, v)_0 d_{q_1} u d_{q_2} v \\
 &+ \frac{1}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_2}^{\xi_2 + \eta_1(\mathbf{Q}, \xi_2)} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(u, v)_0 d_{q_1} u d_{q_2} v \\
 &+ \frac{1}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_2}^{\xi_2 + \eta_1(\mathbf{Q}, \xi_2)} \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(u, v)_0 d_{q_1} u d_{q_2} v.
 \end{aligned} \tag{18}$$

Proof. Considering

$$\begin{aligned}
 & - \int_0^1 \int_0^1 z w \frac{\xi_1 \xi_3 \partial_{q_1 q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_1 + z \eta_1(\mathbf{Q}, \xi_1), \xi_3 + w \eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\
 & - \int_0^1 \int_0^1 z w \frac{\xi_1 \xi_3 \partial_{q_1 q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_1 + z \eta_1(\mathbf{Q}, \xi_1), \xi_4 + w \eta_2(\rho, \xi_4))_0 d_{q_1} z_0 d_{q_2} w \\
 & - \int_0^1 \int_0^1 z w \frac{\xi_1 \xi_3 \partial_{q_1 q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_2 + z \eta_1(\mathbf{Q}, \xi_2), \xi_3 + w \eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\
 & - \int_0^1 \int_0^1 z w \frac{\xi_1 \xi_3 \partial_{q_1 q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_2 + z \eta_1(\mathbf{Q}, \xi_2), \xi_4 + w \eta_2(\rho, \xi_4))_0 d_{q_1} z_0 d_{q_2} w,
 \end{aligned} \tag{19}$$

it follows from the definitions of partial $q_1 q_2$ -derivative and $q_1 q_2$ -integral that

$$\begin{aligned}
 & \int_0^1 \int_0^1 z w \frac{\xi_1 \xi_3 \partial_{q_1 q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_1 + z \eta_1(\mathbf{Q}, \xi_1), \xi_3 + w \eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\
 &= \frac{1}{(1 - q_1)(1 - q_2) \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \times \left[\int_0^1 \int_0^1 \Phi(\xi_1 + z q_1 \eta_1(\mathbf{Q}, \xi_1), \xi_3 \right. \\
 &+ w q_2 \eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w - \int_0^1 \int_0^1 \Phi(\xi_1 + z q_1 \eta_1(\mathbf{Q}, \xi_1), \xi_3 \\
 &+ w \eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w - \int_0^1 \int_0^1 \Phi(\xi_1 + z \eta_1(\mathbf{Q}, \xi_1), \xi_3 + w q_2 \eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\
 &+ \left. \int_0^1 \int_0^1 \Phi(\xi_1 + z \eta_1(\mathbf{Q}, \xi_1), \xi_3 + w \eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \right] = \frac{1}{\eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \\
 &\times \left[\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi \left(\xi_1 + q_1^{n+1} \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^{m+1} \eta_2(\rho, \xi_3) \right) \right. \\
 &\left. - \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^{n+1} \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)) \right)
 \end{aligned}$$

$$\begin{aligned}
 & - \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^{m+1} \eta_2(\rho, \xi_3)) \\
 &+ \left. \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^{n+1} \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)) \right] \\
 &= \frac{1}{q_1 q_2 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)) \\
 &- \frac{1}{q_1 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)) \\
 &- \frac{1}{q_2 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)) \\
 &+ \frac{1}{\eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)). \tag{20}
 \end{aligned}$$

Note that

$$\begin{aligned}
 & \frac{1}{q_1 q_2 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)) \\
 &= - \frac{\Phi(\mathbf{Q}, \rho)}{q_1 q_2 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} - \frac{1}{q_1 q_2 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \sum_{n=0}^{\infty} q_1^n \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \rho) \\
 &- \frac{1}{q_1 q_2 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \sum_{m=0}^{\infty} q_2^m \Phi(\mathbf{Q}, \xi_3 + q_2^m \eta_2(\rho, \xi_3)) - \frac{1}{q_1 q_2 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \\
 &\times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)), \\
 & \tag{21} \\
 & - \frac{1}{q_1 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)) \\
 &= \frac{1}{q_1 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \sum_{m=0}^{\infty} q_2^m \Phi(\mathbf{Q}, \xi_3 + q_2^m \eta_2(\rho, \xi_3)) - \frac{1}{q_1 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} \\
 &\times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)),
 \end{aligned} \tag{22}$$

$$\begin{aligned}
& -\frac{1}{q_2\eta_1(\mathbf{Q}, \xi_1)\eta_2(\rho, \xi_3)} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)) \\
& = \frac{1}{q_2\eta_1(\mathbf{Q}, \xi_1)\eta_2(\rho, \xi_3)} \sum_{n=0}^{\infty} q_1^n \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \rho) - \frac{1}{q_2\eta_1(\mathbf{Q}, \xi_1)\eta_2(\rho, \xi_3)} \\
& \quad \times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 + q_2^m \eta_2(\rho, \xi_3)).
\end{aligned} \tag{23}$$

From (20)–(23) we get

$$\begin{aligned}
& \int_0^1 \int_0^1 zw \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\
& = -\frac{\Phi(\mathbf{Q}, \rho)}{q_2 q_2 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} - \frac{(1 - q_2) \eta_2(\rho, \xi_3)}{q_1 q_2 \eta_1(\mathbf{Q}, \xi_1) [\eta_2(\rho, \xi_3)]^2} \sum_{m=0}^{\infty} q_2^m(\mathbf{Q}, \xi_3) \\
& \quad + q_2^m \eta_2(\rho, \xi_3) - \frac{(1 - q_1) \eta_1(\mathbf{Q}, \xi_1)}{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 \eta_2(\rho, \xi_3)} \sum_{n=0}^{\infty} q_2^n \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \rho) \\
& \quad + \frac{(1 - q_1)(1 - q_2) \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)}{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 [\eta_2(\rho, \xi_3)]^2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Phi(\xi_1 + q_1^n \eta_1(\mathbf{Q}, \xi_1), \xi_3 \\
& \quad + q_2^m \eta_2(\rho, \xi_3)),
\end{aligned} \tag{24}$$

$$\begin{aligned}
& \int_0^1 \int_0^1 zw \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_2), \xi_3 + w\eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\
& = -\frac{\Phi(\mathbf{Q}, \rho)}{q_1 q_2 \eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)} - \frac{\Phi(\mathbf{Q}, \rho)}{q_1 q_2 \eta_1(\mathbf{Q}, \xi_1) [\eta_2(\rho, \xi_3)]^2} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathbf{Q}, v)_0 d_{q_2} v \\
& \quad - \frac{1}{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 \eta_2(\rho, \xi_3)} \int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \Phi(u, \rho)_0 d_{q_1} v \\
& \quad + \frac{1}{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 [\eta_2(\rho, \xi_3)]^2} \int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_2)} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(u, v)_0 d_{q_1} u_0 d_{q_2} v.
\end{aligned} \tag{25}$$

Multiplying both sides of equality (25) by

$$\frac{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)}, \tag{26}$$

leads to

$$\begin{aligned}
& \frac{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 [\eta_1(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_0^1 \int_0^1 zw \frac{\partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \times \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\
& = -\frac{\eta_1(\mathbf{Q}, \xi_1) \eta_2(\rho, \xi_3)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \Phi(\mathbf{Q}, \rho) - \frac{\eta_1(\mathbf{Q}, \xi_1)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathbf{Q}, v)_0 d_{q_2} v \\
& \quad - \frac{\eta_2(\rho, \xi_3)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \Phi(u, \rho)_0 d_{q_1} u + \frac{1}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(u, v)_0 d_{q_1} u_0 d_{q_2} v.
\end{aligned} \tag{27}$$

Similarly, we have

$$\begin{aligned}
& \frac{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 [\eta_2(\xi_4, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_0^1 \int_0^1 zw \frac{\partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \times \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_4 + z\eta_2(\rho, \xi_4))_0 d_{q_1} z_0 d_{q_2} w \\
& = -\frac{\eta_1(\mathbf{Q}, \xi_1) \eta_2(\xi_4, \rho)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \Phi(\mathbf{Q}, \rho) - \frac{\eta_1(\mathbf{Q}, \xi_1)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(\mathbf{Q}, v)_0 d_{q_2} v \\
& \quad + \frac{\eta_2(\xi_4, \rho)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \Phi(u, \rho)_0 d_{q_1} u + \frac{1}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(u, v)_0 d_{q_1} u_0 d_{q_2} v, \\
& \quad \times \frac{q_1 q_2 [\eta_1(\xi_2, \mathbf{Q})]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_0^1 \int_0^1 zw \frac{\partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \times \Phi(\xi_2 + z\eta_1(\mathbf{Q}, \xi_2) + w\eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\
& = -\frac{\eta_1(\xi_2, \mathbf{Q}) \eta_2(\rho, \xi_3)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \Phi(\mathbf{Q}, \rho) - \frac{\eta_1(\xi_2, \mathbf{Q})}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathbf{Q}, v)_0 d_{q_2} v \\
& \quad - \frac{\eta_2(\rho, \xi_3)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_2}^{\xi_2 + \eta_1(\rho, \xi_2)} \Phi(u, \rho)_0 d_{q_1} u + \frac{1}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_2}^{\xi_2 + \eta_1(\mathbf{Q}, \xi_2)} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(u, v)_0 d_{q_1} u_0 d_{q_2} v,
\end{aligned} \tag{28}$$

$$\begin{aligned}
 & \frac{q_1 q_2 [\eta_1(\xi_2, \mathcal{Q})]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_0^1 \int_0^1 zw \frac{\partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \times \Phi(\xi_2 + z\eta_1(\mathcal{Q}, \xi_2) + w\eta_2(\rho, \xi_3))_0 d_{q_1} z_0 d_{q_2} w \\
 &= -\frac{\eta_1(\xi_2, \mathcal{Q}) \eta_2(\rho, \xi_3)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \Phi(\mathcal{Q}, \rho) - \frac{\eta_1(\xi_2, \mathcal{Q})}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathcal{Q}, \nu)_0 d_{q_2} \nu \\
 & \quad - \frac{\eta_2(\rho, \xi_3)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_2}^{\xi_2 + \eta_1(\mathcal{Q}, \xi_2)} \Phi(u, \rho)_0 d_{q_1} u + \frac{1}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_2}^{\xi_2 + \eta_1(\mathcal{Q}, \xi_2)} \int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(u, \nu)_0 d_{q_1} u_0 d_{q_2} \nu,
 \end{aligned} \tag{29}$$

$$\begin{aligned}
 & \frac{q_1 q_2 [\eta_1(\xi_2, \mathcal{Q})]^2 [\eta_2(\xi_4, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_0^1 \int_0^1 zw \frac{\partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \times \Phi(\xi_2 + z\eta_1(\mathcal{Q}, \xi_2), \xi_4 + w\eta_2(\rho, \xi_4))_0 d_{q_1} z_0 d_{q_2} w \\
 &= -\frac{\eta_1(\xi_2, \mathcal{Q}) \eta_2(\xi_4, \rho)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \Phi(\mathcal{Q}, \rho) - \frac{\eta_1(\xi_2, \mathcal{Q})}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(\mathcal{Q}, \nu)_0 d_{q_2} \nu \\
 & \quad - \frac{\eta_2(\xi_4, \rho)}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_2}^{\xi_2 + \eta_1(\mathcal{Q}, \xi_2)} \Phi(u, \rho)_0 d_{q_1} u + \frac{1}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_{\xi_2}^{\xi_2 + \eta_1(\mathcal{Q}, \xi_2)} \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(u, \nu)_0 d_{q_1} u_0 d_{q_2} \nu.
 \end{aligned} \tag{30}$$

Therefore, Lemma 13 follows from (19), (27), (28), (29), and (30).

Theorem 14. Let $q_1, q_2 \in (0, 1)$, $\xi_1 < \xi_2$, $\xi_3 < \xi_4$, $\Phi : \mathcal{N} \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be mixed partial $q_1 q_2$ differentiable on \mathcal{N}° (the interior of \mathcal{N}) such that its mixed partial $q_1 q_2$ -derivatives is continuous and integrable on $[\xi_1, \xi_1 + \eta_1(\xi_2, \xi_1)] \times [\xi_3, \xi_3 + \eta_2(\xi_4, \xi_3)] \subset \mathcal{N}^\circ$ for $\eta_1(\xi_2, \xi_1), \eta_2(\xi_4, \xi_3) > 0$. If $|(\xi_1, \xi_3 \partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w) \Phi|$ is an n -polynomial preinvex function on the coordinates on $[\xi_1, \xi_1 + \eta_1(\xi_2, \xi_1)] \times [\xi_3, \xi_3 + \eta_2(\xi_4, \xi_3)]$ and $|(\xi_1, \xi_3 \partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w) \Phi(\mathcal{Q}, \rho)| \leq M$ for $\mathcal{Q}, \rho \in \mathcal{N}$, then we have

$$\begin{aligned}
 & \left| \Phi(\mathcal{Q}, \rho) + \frac{1}{\eta_2(\xi_4, \xi_3)} \left[\int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathcal{Q}, \nu)_0 d_{q_2} \nu + \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_3)} \Phi(\mathcal{Q}, \nu)_0 d_{q_2} \nu \right] \right. \\
 & \quad \left. + \frac{1}{\eta_1(\xi_2, \xi_1)} \left[\int_{\xi_1}^{\xi_1 + \eta_1(\mathcal{Q}, \xi_1)} \Phi(u, \rho)_0 d_{q_1} u + \int_{\xi_2}^{\xi_2 + \eta_1(\mathcal{Q}, \xi_1)} \Phi(u, \rho)_0 d_{q_1} u \right] - \mathcal{Q} \right| \\
 & \leq q_1 q_2 M (\mathcal{A}_{q_1} + \mathcal{B}_{q_1}) (\mathcal{A}_{q_2} + \mathcal{B}_{q_2}) \left[\frac{[\eta_1(\mathcal{Q}, \xi_1)]^2 + [\eta_1(\xi_2, \mathcal{Q})]^2}{\eta_1(\xi_2, \xi_1)} \right] \\
 & \quad \times \left[\frac{[\eta_2(\rho, \xi_3)]^2 + [\eta_2(\xi_4, \rho)]^2}{\eta_2(\xi_4, \xi_3)} \right],
 \end{aligned} \tag{31}$$

where \mathcal{Q} is defined in Lemma 13.

Proof. It follows from (19) and the properties of the n -polynomial preinvexity function of the function $|(\partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w) \Phi|$ on coordinates that

$$\begin{aligned}
 & \left| \Phi(\mathcal{Q}, \rho) + \frac{1}{\eta_2(\xi_4, \xi_3)} \left[\int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathcal{Q}, \rho)_0 d_{q_2} \nu + \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_3)} \Phi(\mathcal{Q}, \rho)_0 d_{q_2} \nu \right] + \frac{1}{\eta_1(\xi_2, \xi_1)} \left[\int_{\xi_1}^{\xi_1 + \eta_1(\mathcal{Q}, \xi_1)} \Phi(u, \rho)_0 d_{q_1} u + \int_{\xi_2}^{\xi_2 + \eta_1(\mathcal{Q}, \xi_1)} \Phi(u, \rho)_0 d_{q_1} u \right] - \mathcal{Q} \right| \\
 & \leq \frac{q_1 q_2 [\eta_1(\mathcal{Q}, \xi_1)]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(\mathcal{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_0 d_{q_1} z_0 d_{q_2} w \\
 & \quad + \frac{q_1 q_2 [\eta_1(\mathcal{Q}, \xi_1)]^2 [\eta_2(\xi_4, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(\mathcal{Q}, \xi_1), \xi_4 + w\eta_2(\rho, \xi_4)) \right|_0 d_{q_1} z_0 d_{q_2} w \\
 & \quad + \frac{q_1 q_2 [\eta_1(\xi_2, \mathcal{Q})]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(\mathcal{Q}, \xi_2), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_0 d_{q_1} z_0 d_{q_2} w \\
 & \quad + \frac{q_1 q_2 [\eta_1(\xi_2, \mathcal{Q})]^2 [\eta_2(\xi_4, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(\mathcal{Q}, \xi_2), \xi_4 + w\eta_2(\rho, \xi_4)) \right|_0 d_{q_1} z_0 d_{q_2} w.
 \end{aligned} \tag{32}$$

Considering the first integral

$$\begin{aligned} & \int_0^1 \int_0^1 zw \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(Q, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_0 d_{q_1} z_0 d_{q_2} w \\ & \leq \int_0^1 w \left\{ \int_0^1 z \left[\frac{1}{n} \sum_{p=1}^n [1 - (1-z)^p] \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(Q, \xi_3 + w\eta_2(\rho, \xi_3)) \right| \right. \right. \\ & \quad \left. \left. + \frac{1}{n} \sum_{p=1}^n [1 - z^p] \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1, \xi_3 + w\eta_2(\rho, \xi_3)) \right| \right] \right\}_0 d_{q_1} z \cdot_0 d_{q_2} w, \end{aligned} \tag{33}$$

in view of the Definition 6 for $k = 1, 2$, we get

$$\begin{aligned} \mathcal{A}_{qk} &= \frac{1}{n} \sum_{p=1}^n \int_0^1 z [1 - (1-z)^p]_0 d_{qk} z = \frac{1}{1+q_k} - \frac{(1-q_k)}{n} \sum_{p=1}^n \sum_{e=0}^{\infty} q_k^{2e} (1-q_k^e)^p, \\ \mathcal{B}_{qk} &= \frac{1}{n} \sum_{p=1}^n \int_0^1 z [1 - z^p]_0 d_{qk} z = \frac{1}{1+q_k} - \frac{1}{n} \sum_{p=1}^n \frac{1-q_k}{1-q_k^{p+2}}. \end{aligned} \tag{34}$$

From (33) and computing the q_1 -integral, we get

$$\begin{aligned} & \int_0^1 \int_0^1 zw \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(Q, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_0 d_{q_1} z_0 d_{q_2} w \\ & \leq \int_0^1 w \left[\begin{aligned} & \mathcal{A}_{q_1} \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(Q, \xi_3 + w\eta_2(\rho, \xi_3)) \right| \\ & + \mathcal{B}_{q_1} \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1, \xi_3 + w\eta_2(\rho, \xi_3)) \right| \end{aligned} \right]_0 d_{q_2} w. \end{aligned} \tag{35}$$

Computing the q_2 -integral and utilizing the fact $|\frac{\xi_1, \xi_3}{\partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z \partial_{q_2} w} \Phi(Q, \rho)| \leq M$ for $Q, \rho \in \mathcal{N}$, inequality (35) leads to the conclusion that

$$\begin{aligned} & \int_0^1 \int_0^1 zw \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(Q, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_0 d_{q_1} z_0 d_{q_2} w \\ & \leq M (\mathcal{A}_{q_1} + \mathcal{B}_{q_1}) (\mathcal{A}_{q_2} + \mathcal{B}_{q_2}). \end{aligned} \tag{36}$$

Analogously, we also have

$$\begin{aligned} & \int_0^1 \int_0^1 zw \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(Q, \xi_1), \xi_4 + w\eta_2(\rho, \xi_4)) \right|_0 d_{q_1} z_0 d_{q_2} w \\ & \leq M (\mathcal{A}_{q_1} + \mathcal{B}_{q_1}) (\mathcal{A}_{q_2} + \mathcal{B}_{q_2}), \end{aligned} \tag{37}$$

$$\begin{aligned} & \int_0^1 \int_0^1 zw \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(Q, \xi_2), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_0 d_{q_1} z_0 d_{q_2} w \\ & \leq M (\mathcal{A}_{q_1} + \mathcal{B}_{q_1}) (\mathcal{A}_{q_2} + \mathcal{B}_{q_2}), \end{aligned} \tag{38}$$

$$\begin{aligned} & \int_0^1 \int_0^1 zw \left| \frac{\xi_1, \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(Q, \xi_2), \xi_4 + w\eta_2(\rho, \xi_4)) \right|_0 d_{q_1} z_0 d_{q_2} w \\ & \leq M (\mathcal{A}_{q_1} + \mathcal{B}_{q_1}) (\mathcal{A}_{q_2} + \mathcal{B}_{q_2}). \end{aligned} \tag{39}$$

Making use of the inequalities (36), (37), (38), and (39) and the fact that

$$\begin{aligned} & [\eta_1(Q, \xi_1)]^2 [\eta_2(\rho, \xi_3)]^2 + [\eta_1(Q, \xi_1)]^2 [\eta_2(\xi_4, \rho)]^2 \\ & \quad + [\eta_1(\xi_2, Q)]^2 [\eta_2(\rho, \xi_3)]^2 + [\eta_1(\xi_2, Q)]^2 [\eta_2(\xi_4, \rho)]^2 \\ & = \left[[\eta_1(Q, \xi_1)]^2 + [\eta_1(\xi_2, Q)]^2 \right] \left[[\eta_2(\rho, \xi_3)]^2 + [\eta_2(\xi_4, \rho)]^2 \right], \end{aligned} \tag{40}$$

we get the desired inequality (31).

Theorem 15. Let $\gamma_1, \gamma_2 > 1$ with $(1/\gamma_2) + (1/\gamma_1) = 1$, $q_1, q_2 \in (0, 1)$, and $\Phi : \mathcal{N} \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be mixed partial $q_1 q_2$ -differentiable on \mathcal{N}° such that its mixed partial $q_1 q_2$ -derivative is continuous and integrable on $[\xi_1, \xi_1 + \eta_1(\xi_2, \xi_1)] \times [\xi_3, \xi_3 + \eta_2(\xi_4, \xi_3)] \subset \mathcal{N}^\circ$ for $\eta_1(\xi_2, \xi_1), \eta_2(\xi_4, \xi_3) > 0$. If $|\frac{\xi_1, \xi_3}{\partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi|^{\gamma_2}$ is an n -polynomial preinvex function on the coordinates on $[\xi_1, \xi_1 + \eta_1(\xi_2, \xi_1)] \times [\xi_3, \xi_3 + \eta_2(\xi_4, \xi_3)]$ and $|\frac{\xi_1, \xi_3}{\partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(Q, \rho)| \leq M$ for $Q, \rho \in \mathcal{N}$, then we have

$$\begin{aligned} & \left| \Phi(Q, \rho) + \frac{1}{\eta_2(\xi_4, \xi_3)} \left[\int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(Q, \rho)_0 d_{q_2} v + \int_{\xi_4}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(Q, \rho)_0 d_{q_2} v \right] \frac{1}{\eta_1(\xi_2, \xi_1)} \right. \\ & \quad \cdot \left[\int_{\xi_1}^{\xi_1 + \eta_1(Q, \xi_1)} \Phi(u, \rho)_0 d_{q_1} u + \int_{\xi_2}^{\xi_1 + \eta_1(Q, \xi_1)} \Phi(u, \rho)_0 d_{q_1} u \right] - Q \left| \leq \frac{q_1 q_2 M \sqrt{\gamma_2} \sqrt{(\mathcal{E}_{q_1} + \mathcal{D}_{q_1}) (\mathcal{E}_{q_2} + \mathcal{D}_{q_2})}}{\sqrt{\gamma_1} \sqrt{[\gamma_1 + 1]_{q_1} [\gamma_1 + 1]_{q_2}}} \right. \\ & \quad \cdot \left[\frac{[\eta_1(Q, \xi_1)]^2 + [\eta_1(\xi_2, Q)]^2}{\eta_1(\xi_2, \xi_1)} \right] \times \left[\frac{[\eta_2(\rho, \xi_3)]^2 + [\eta_2(\xi_4, \rho)]^2}{\eta_2(\xi_4, \xi_3)} \right], \end{aligned} \tag{41}$$

where \mathcal{Q} is defined in Lemma 13.

Proof. It follows from (19), the Hölder inequality and the property of n -polynomial preinvexity of the function $|\partial_{q_1, q_2}^2 \Phi / \xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w|^{\gamma_2}$ on coordinates that

$$\begin{aligned} & \left| \Phi(\mathbf{Q}, \rho) + \frac{1}{\eta_2(\xi_4, \xi_3)} \left[\int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathbf{Q}, v) {}_0 d_{q_2} v \right. \right. \\ & \quad \left. \left. + \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(\mathbf{Q}, v) {}_0 d_{q_2} v \right] \right. \\ & \quad \left. + \frac{1}{\eta_1(\xi_2, \xi_1)} \left[\int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \Phi(u, \rho) {}_0 d_{q_1} u \right. \right. \\ & \quad \left. \left. + \int_{\xi_2}^{\xi_2 + \eta_1(\mathbf{Q}, \xi_2)} \Phi(u, \rho) {}_0 d_{q_1} u \right] - \mathcal{Q} \right| \\ & \leq \left(\int_0^1 \int_0^1 z^{\gamma_1} w^{\gamma_1} {}_0 d_{q_1} z {}_0 d_{q_2} w \right)^{1/\gamma_1} \\ & \quad \times \left[\frac{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \right. \\ & \quad \cdot \left(\int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} {}_0 d_{q_1} z {}_0 d_{q_2} w \right)^{1/\gamma_2} \\ & \quad + \frac{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 [\eta_2(\xi_3, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \\ & \quad \cdot \left(\int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_4 + w\eta_2(\rho, \xi_4))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} {}_0 d_{q_1} z {}_0 d_{q_2} w \right)^{1/\gamma_2} \\ & \quad + \frac{q_1 q_2 [\eta_1(\xi_2, \mathbf{Q})]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \\ & \quad \cdot \left(\int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_2 + z\eta_1(\mathbf{Q}, \xi_2), \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} {}_0 d_{q_1} z {}_0 d_{q_2} w \right)^{1/\gamma_2} \\ & \quad + \frac{q_1 q_2 [\eta_1(\xi_2, \mathbf{Q})]^2 [\eta_2(\xi_4, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \\ & \quad \cdot \left. \left(\int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_2 + z\eta_1(\mathbf{Q}, \xi_2), \xi_4 + w\eta_2(\rho, \xi_4))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} {}_0 d_{q_1} z {}_0 d_{q_2} w \right)^{\frac{1}{\gamma_2}} \right], \end{aligned} \tag{42}$$

for all $\mathbf{Q}, \rho \in N$.

From the n -polynomial preinvexity of the function $|(\xi_1 \xi_3 \partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w) \Phi|^{\gamma_2}$, we get

$$\begin{aligned} & \int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} {}_0 d_{q_1} z {}_0 d_{q_2} w \\ & \leq \int_0^1 \left\{ \int_0^1 \left[\frac{1}{n} \sum_{p=1}^n [1 - (1-z)^p] \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\mathbf{Q}, \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} \right. \right. \\ & \quad \left. \left. + \frac{1}{n} \sum_{p=1}^n [1 - z^p] \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1, \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} \right] {}_0 d_{q_1} z \right\} {}_0 d_{q_2} w. \end{aligned} \tag{43}$$

Computing the q_1 -integral on the right-hand side of (43),

we have

$$\begin{aligned} & \int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} {}_0 d_{q_1} z {}_0 d_{q_2} w \\ & \leq \int_0^1 \left[\frac{1}{n} \sum_{p=1}^n [1 - (1-z)^p] \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\mathbf{Q}, \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} \right. \\ & \quad \left. + \frac{1}{n} \sum_{p=1}^n [1 - z^p] \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1, \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} \right] {}_0 d_{q_1} z. \end{aligned} \tag{44}$$

In view of the Definition 6 for $k = 1, 2$, we obtain

$$\begin{aligned} \mathcal{E}_{q_k} &= \frac{1}{n} \sum_{p=1}^n \int_0^1 [1 - (1-z)^p] {}_0 d_{q_k} z = 1 - \frac{(1-q_k)}{n} \sum_{p=1}^n \sum_{e=0}^{\infty} q_k^e (1-q_k^e)^p, \\ \mathcal{D}_{q_k} &= \frac{1}{n} \sum_{p=1}^n \int_0^1 [1 - z^p] {}_0 d_{q_k} z = 1 - \frac{1}{n} \sum_{p=1}^n \frac{1 - q_k}{1 - q_k^{p+2}}. \end{aligned} \tag{45}$$

Therefore, we get

$$\begin{aligned} & \int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} {}_0 d_{q_1} z {}_0 d_{q_2} w \\ & \leq \int_0^1 \left[\mathcal{E}_{q_1} \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\mathbf{Q}, \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} \right. \\ & \quad \left. + \mathcal{D}_{q_1} \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1, \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} \right] {}_0 d_{q_2} w. \end{aligned} \tag{46}$$

Similarly, computing the q_2 -integral and utilizing the fact $|(\xi_1 \xi_3 \partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z \partial_{q_2} w) \Phi(\mathbf{Q}, \rho)| \leq M$ for $\mathbf{Q}, \rho \in N$ on the right-hand side of (46), one has

$$\begin{aligned} & \int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} {}_0 d_{q_1} z {}_0 d_{q_2} w \\ & \leq M^{\gamma_2} (\mathcal{E}_{q_1} + \mathcal{D}_{q_1}) (\mathcal{E}_{q_2} + \mathcal{D}_{q_2}). \end{aligned} \tag{47}$$

Analogously, we also can get

$$\begin{aligned} & \int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2 \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_4 + w\eta_2(\rho, \xi_4))}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \right|^{\gamma_2} {}_0 d_{q_1} z {}_0 d_{q_2} w \\ & \leq M^{\gamma_2} (\mathcal{E}_{q_1} + \mathcal{D}_{q_1}) (\mathcal{E}_{q_2} + \mathcal{D}_{q_2}), \end{aligned} \tag{48}$$

$$\int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(\mathbf{Q}, \xi_2), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_{0d_{q_1} z_0 d_{q_2} w}^{\gamma_2} \leq M^{\gamma_2} (\mathcal{C}_{q_1} + \mathcal{D}_{q_1}) (\mathcal{C}_{q_2} + \mathcal{D}_{q_2}), \tag{49}$$

$$\int_0^1 \int_0^1 \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(\mathbf{Q}, \xi_2), \xi_4 + w\eta_2(\rho, \xi_4)) \right|_{0d_{q_1} z_0 d_{q_2} w}^{\gamma_2} \leq M^{\gamma_2} (\mathcal{C}_{q_1} + \mathcal{D}_{q_1}) (\mathcal{C}_{q_2} + \mathcal{D}_{q_2}). \tag{50}$$

Therefore, the desired inequality (41) follows from (47), (48), (49), and (50) and the fact that

$$\int_0^1 \int_0^1 z^{\gamma_1} w^{\gamma_1} {}_0d_{q_1} z_0 d_{q_2} w = \frac{1}{[\gamma_1 + 1]_{q_1} [\gamma_1 + 1]_{q_2}}, \tag{51}$$

where $[\gamma_1 + 1]_{q_1}$ and $[\gamma_1 + 1]_{q_2}$ are the q_1 - and q_2 -analogues of $\gamma_1 + 1$ and $\gamma_2 + 1$, respectively.

Theorem 16. Let $q_1, q_2 \in (0, 1)$, $\gamma > 1$ and $\Phi : \mathcal{N} \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be mixed partial q_1, q_2 -differentiable over \mathcal{N}° such that its partial q_1, q_2 -derivative is continuous and integrable on $[\xi_1, \xi_1 + \eta_1(\xi_2, \mathbf{Q})]$ for all $\mathbf{Q}, \rho \in \mathcal{N}$.

$\xi_1) \times [\xi_3, \xi_3 + \eta_2(\xi_4, \xi_3)] \subset \mathcal{N}^\circ$ for $\eta_1(\xi_2, \xi_1), \eta_2(\xi_4, \xi_3) > 0$. If $|(\xi_1, \xi_3 \partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w) \Phi|$ is an n -polynomial preinvex function on the coordinates on $[\xi_1, \xi_1 + \eta_1(\xi_2, \xi_1)] \times [\xi_3, \xi_3 + \eta_2(\xi_4, \xi_3)]$ and $|(\xi_1, \xi_3 \partial_{q_1, q_2}^2 / \xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w) \Phi(\mathbf{Q}, \rho)| \leq M$ for $\mathbf{Q}, \rho \in \mathcal{N}$, then we have the inequality

$$\left| \Phi(\mathbf{Q}, \rho) + \frac{1}{\eta_2(\xi_4, \xi_3)} \left[\int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathbf{Q}, v) {}_0d_{q_2} v + \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(\mathbf{Q}, v) {}_0d_{q_2} v \right] + \frac{1}{\eta_1(\xi_2, \xi_1)} \left[\int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \Phi(u, \rho) {}_0d_{q_1} u + \int_{\xi_2}^{\xi_2 + \eta_1(\mathbf{Q}, \xi_2)} \Phi(u, \rho) {}_0d_{q_1} u \right] - \mathcal{Q} \right| \leq \frac{q_1 q_2 M \sqrt{(\mathcal{A}_{q_1} + \mathcal{B}_{q_1})(\mathcal{A}_{q_2} + \mathcal{B}_{q_2})}}{[(1 + q_1)(1 + q_2)]^{1-1/\gamma}} \times \left[\frac{[\eta_1(\mathbf{Q}, \xi_1)]^2 + [\eta_1(\xi_2, \mathbf{Q})]^2}{\eta_1(\xi_2, \xi_1)} \right] \left[\frac{[\eta_2(\rho, \xi_3)]^2 + [\eta_2(\xi_4, \rho)]^2}{\eta_2(\xi_4, \xi_3)} \right], \tag{52}$$

where \mathcal{Q} is defined in Lemma 13.

Proof. It follows from (19), the power mean inequality and the property of n -polynomial preinvexity of the function $|(\xi_1, \xi_3 \partial_{q_1, q_2}^2 \Phi / \xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w)|$ on coordinates that

$$\begin{aligned} & \left| \Phi(\mathbf{Q}, \rho) + \frac{1}{\eta_2(\xi_4, \xi_3)} \left[\int_{\xi_3}^{\xi_3 + \eta_2(\rho, \xi_3)} \Phi(\mathbf{Q}, \rho) {}_0d_{q_2} v + \int_{\xi_4}^{\xi_4 + \eta_2(\rho, \xi_4)} \Phi(\mathbf{Q}, v) {}_0d_{q_2} v \right] + \frac{1}{\eta_1(\xi_2, \xi_1)} \left[\int_{\xi_1}^{\xi_1 + \eta_1(\mathbf{Q}, \xi_1)} \Phi(u, \rho) {}_0d_{q_1} u + \int_{\xi_2}^{\xi_2 + \eta_1(\mathbf{Q}, \xi_2)} \Phi(u, \rho) {}_0d_{q_1} u \right] - \mathcal{Q} \right| \\ & \leq \left(\int_0^1 \int_0^1 zw {}_0d_{q_1} z_0 d_{q_2} w \right)^{1-(1/\gamma)} \times \left[\frac{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \right. \\ & \quad \cdot \left. \left(\int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_{0d_{q_1} z_0 d_{q_2} w}^{\gamma} \right)^{1/\gamma} \right] \\ & \quad + \left[\frac{q_1 q_2 [\eta_1(\mathbf{Q}, \xi_1)]^2 [\eta_2(\xi_4, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \left(\int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(\mathbf{Q}, \xi_1), \xi_4 + w\eta_2(\rho, \xi_4)) \right|_{0d_{q_1} z_0 d_{q_2} w}^{\gamma} \right)^{1/\gamma} \right] \\ & \quad + \left[\frac{q_1 q_2 [\eta_1(\xi_2, \mathbf{Q})]^2 [\eta_2(\rho, \xi_3)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \left(\int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(\mathbf{Q}, \xi_2), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_{0d_{q_1} z_0 d_{q_2} w}^{\gamma} \right)^{1/\gamma} \right] \\ & \quad + \left[\frac{q_1 q_2 [\eta_1(\xi_2, \mathbf{Q})]^2 [\eta_2(\xi_4, \rho)]^2}{\eta_1(\xi_2, \xi_1) \eta_2(\xi_4, \xi_3)} \left(\int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z_{\xi_3} \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(\mathbf{Q}, \xi_2), \xi_4 + w\eta_2(\rho, \xi_4)) \right|_{0d_{q_1} z_0 d_{q_2} w}^{\gamma} \right)^{1/\gamma} \right], \end{aligned} \tag{53}$$

By similar argument as in Theorem 14, we can prove that

$$\int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(\mathcal{Q}, \xi_1), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_0^y d_{q_1} z_0 d_{q_2} w \leq M^y (\mathcal{A}_{q_1} + \mathcal{B}_{q_1}) (\mathcal{A}_{q_2} + \mathcal{B}_{q_2}), \quad (54)$$

$$\int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_1 + z\eta_1(\mathcal{Q}, \xi_1), \xi_4 + w\eta_2(\rho, \xi_4)) \right|_0^y d_{q_1} z_0 d_{q_2} w \leq M^y (\mathcal{A}_{q_1} + \mathcal{B}_{q_1}) (\mathcal{A}_{q_2} + \mathcal{B}_{q_2}), \quad (55)$$

$$\int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(\mathcal{Q}, \xi_2), \xi_3 + w\eta_2(\rho, \xi_3)) \right|_0^y d_{q_1} z_0 d_{q_2} w \leq M^y (\mathcal{A}_{q_1} + \mathcal{B}_{q_1}) (\mathcal{A}_{q_2} + \mathcal{B}_{q_2}), \quad (56)$$

$$\int_0^1 \int_0^1 zw \left| \frac{\xi_1 \xi_3 \partial_{q_1, q_2}^2}{\xi_1 \partial_{q_1} z \xi_3 \partial_{q_2} w} \Phi(\xi_2 + z\eta_1(\mathcal{Q}, \xi_2), \xi_4 + w\eta_2(\rho, \xi_4)) \right|_0^y d_{q_1} z_0 d_{q_2} w \leq M^y (\mathcal{A}_{q_1} + \mathcal{B}_{q_1}) (\mathcal{A}_{q_2} + \mathcal{B}_{q_2}). \quad (57)$$

Now by making use of the inequalities (54), (55), (56), and (57) and the fact that

$$\int_0^1 \int_0^1 zw_0 d_{q_1} z_0 d_{q_2} w = \frac{1}{(1+q_1)(1+q_2)}, \quad (58)$$

we get the desired inequality (52).

3. Conclusion

In the article, we have introduced a new class of preinvex functions which is named n -polynomial preinvex functions and discovered a new quantum integral identity involving second-order mixed partial differentiable function. By using the obtained quantum integral identity as an auxiliary result, we have established several $q_1 q_2$ -Ostrowski-type inequalities for the class of n -polynomial preinvex functions on coordinates, which have improved and unified many previously known results. Our given ideas and approaches may lead to a lot of follow-up research for the interested readers.

Data Availability

No data were used to support this study.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Authors' Contributions

H. Kalsoom introduced the Definition 11, carried out the proof of Proposition 12 and Lemma 13, and drafted the manuscript. M. Idrees carried out the proof of Theorem 14. D. Baleanu carried out the proof of Theorem 15. Y.-M. Chu provided the main idea of the manuscript, carried out the proof of Theorem 16, completed the final revision and submitted the article. All authors read and approved the final manuscript.

Acknowledgments

The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions. This work was supported by the Natural Science Foundation of China (Grant Nos. 61673169, 11971142, 11871202, 11301127, 11701176, 11626101, and 11601485).

References

- [1] F. H. Jackson, "On a q -definite integrals," *Quarterly Journal of Pure and Applied Mathematics*, vol. 41, pp. 193–203, 1910.
- [2] J. Tariboon and S. K. Ntouyas, "Quantum integral inequalities on finite intervals," *Journal of Inequalities and Applications*, vol. 2014, no. 1, 2014.
- [3] J. Tariboon and S. K. Ntouyas, "Quantum calculus on finite intervals and applications to impulsive difference equations," *Advances in Difference Equations*, vol. 2013, no. 1, 2013.
- [4] M. A. Noor, M. U. Awan, and K. I. Noor, "Quantum Ostrowski inequalities for q -differentiable convex functions," *Journal of Mathematical Inequalities*, vol. 10, no. 4, pp. 1013–1018, 2007.
- [5] M. A. Latif, S. Rashid, S. S. Dragomir, and Y.-M. Chu, "Hermite-Hadamard type inequalities for co-ordinated convex and qausi-convex functions and their applications," *Journal of Inequalities and Applications*, vol. 2019, no. 1, 2019.
- [6] M. U. Awan, N. Akhtar, S. Iftikhar, M. A. Noor, and Y.-M. Chu, "New Hermite-Hadamard type inequalities for n -polynomial harmonically convex functions," *Journal of Inequalities and Applications*, vol. 2020, no. 1, 2020.
- [7] M. Adil Khan, N. Mohammad, E. R. Nwaeze, and Y.-M. Chu, "Quantum Hermite Hadamard inequality by means of a Green function," *Advances in Difference Equations*, vol. 2020, no. 1, 2020.
- [8] A. Iqbal, M. Adil Khan, S. Ullah, and Y.-M. Chu, "Some new Hermite-Hadamard-type inequalities associated with conformable fractional integrals and their applications," *Journal of Function Spaces*, vol. 2020, Article ID 9845407, 18 pages, 2020.
- [9] M. U. Awan, S. Talib, Y.-M. Chu, M. A. Noor, and K. I. Noor, "Some new refinements of Hermite-Hadamard-type inequalities involving Ψ_k -Riemann-Liouville fractional integrals and applications," *Mathematical Problems in Engineering*, vol. 2020, Article ID 3051920, 10 pages, 2020.
- [10] I. Abbas Baloch and Y.-M. Chu, "Petrović-type inequalities for harmonic h -convex functions," *Journal of Function Spaces*, vol. 2020, Article ID 3075390, 7 pages, 2020.
- [11] S. Rashid, F. Jarad, H. Kalsoom, and Y.-M. Chu, "On Pólya-Szegő and Čebyšev type inequalities via generalized

- k -fractional integrals,” *Advances in Difference Equations*, vol. 2020, no. 1, 2020.
- [12] S. Rashid, M. A. Noor, K. I. Noor, and Y.-M. Chu, “Ostrowski type inequalities in the sense of generalized K -fractional integral operator for exponentially convex functions,” *AIMS Mathematics*, vol. 5, no. 3, pp. 2629–2645, 2020.
- [13] S. Rashid, F. Jarad, and Y.-M. Chu, “A note on reverse Minkowski inequality via generalized proportional fractional integral operator with respect to another function,” *Mathematical Problems in Engineering*, vol. 2020, Article ID 7630260, 12 pages, 2020.
- [14] S. Khan, M. Adil Khan, and Y.-M. Chu, “Converses of the Jensen inequality derived from the Green functions with applications in information theory,” *Mathematical Methods in the Applied Sciences*, vol. 43, no. 5, pp. 2577–2587, 2020.
- [15] M. Adil Khan, J. Pečarić, and Y. M. Chu, “Refinements of Jensen’s and McShane’s inequalities with applications,” *AIMS Mathematics*, vol. 5, no. 5, pp. 4931–4945, 2020.
- [16] T.-H. Zhao, L. Shi, and Y.-M. Chu, “Convexity and concavity of the modified Bessel functions of the first kind with respect to Hölder means,” *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales Serie A Matemáticas*, vol. 114, no. 2, p. 96, 2020.
- [17] M.-K. Wang, M.-Y. Hong, Y.-F. Xu, Z.-H. Shen, and Y.-M. Chu, “Inequalities for generalized trigonometric and hyperbolic functions with one parameter,” *Journal of Mathematical Inequalities*, vol. 14, no. 1, pp. 1–21, 2020.
- [18] Y.-M. Chu, M. Adil Khan, T. Ali, and S. S. Dragomir, “Inequalities for α -fractional differentiable functions,” *Journal of Inequalities and Applications*, vol. 2017, no. 1, 2017.
- [19] S.-S. Zhou, S. Rashid, F. Jarad, H. Kalsoom, and Y.-M. Chu, “New estimates considering the generalized proportional Hadamard fractional integral operators,” *Advances in Difference Equations*, vol. 2020, no. 1, 2020.
- [20] Y. Khurshid, M. Adil Khan, and Y.-M. Chu, “Conformable fractional integral inequalities for GG- and GA-convex functions,” *AIMS Mathematics*, vol. 5, no. 5, pp. 5012–5030, 2020.
- [21] S. Rafeeq, H. Kalsoom, S. Hussain, S. Rashid, and Y.-M. Chu, “Delay dynamic double integral inequalities on time scales with applications,” *Advances in Difference Equations*, vol. 2020, no. 1, 2020.
- [22] S. Rashid, İ. İşcan, D. Baleanu, and Y. M. Chu, “Generation of new fractional inequalities via n polynomials s -type convexity with applications,” *Advances in Difference Equations*, vol. 2020, no. 1, 2020.
- [23] M.-K. Wang, H.-H. Chu, and Y.-M. Chu, “Precise bounds for the weighted Hölder mean of the complete p -elliptic integrals,” *Journal of Mathematical Analysis and Applications*, vol. 480, no. 2, article 123388, 2019.
- [24] M.-K. Wang, Z.-Y. He, and Y.-M. Chu, “Sharp power mean inequalities for the generalized elliptic integral of the first kind,” *Computational Methods and Function Theory*, vol. 20, no. 1, pp. 111–124, 2020.
- [25] T.-H. Zhao, M.-K. Wang, and Y.-M. Chu, “A sharp double inequality involving generalized complete elliptic integral of the first kind,” *AIMS Mathematics*, vol. 5, no. 5, pp. 4512–4528, 2020.
- [26] M.-K. Wang, H.-H. Chu, Y.-M. Li, and Y.-M. Chu, “Answers to three conjectures on convexity of three functions involving complete elliptic integrals of the first kind,” *Applicable Analysis and Discrete Mathematics*, vol. 14, pp. 255–271, 2020.
- [27] W.-M. Qian, Z.-Y. He, and Y.-M. Chu, “Approximation for the complete elliptic integral of the first kind,” *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales Serie A Matemáticas*, vol. 114, no. 2, p. 57, 2020.
- [28] Z.-H. Yang, W.-M. Qian, W. Zhang, and Y.-M. Chu, “Notes on the complete elliptic integral of the first kind,” *Mathematical Inequalities & Applications*, vol. 23, no. 1, pp. 77–93, 2020.
- [29] S. Zaheer Ullah, M. Adil Khan, and Y.-M. Chu, “A note on generalized convex functions,” *Journal of Inequalities and Applications*, vol. 2019, no. 1, 2019.
- [30] M. U. Awan, N. Akhtar, A. Kashuri, M. A. Noor, and Y.-M. Chu, “2D approximately reciprocal ρ -convex functions and associated integral inequalities,” *AIMS Mathematics*, vol. 5, no. 5, pp. 4662–4680, 2020.
- [31] S. Rashid, R. Ashraf, M. A. Noor, K. I. Noor, and Y.-M. Chu, “New weighted generalizations for differentiable exponentially convex mapping with application,” *AIMS Mathematics*, vol. 5, no. 4, pp. 3525–3546, 2020.
- [32] W.-M. Qian, Y.-Y. Yang, H.-W. Zhang, and Y.-M. Chu, “Optimal two-parameter geometric and arithmetic mean bounds for the Sándor–Yang mean,” *Journal of Inequalities and Applications*, vol. 2019, no. 1, 2019.
- [33] W.-M. Qian, Z.-Y. He, H.-W. Zhang, and Y.-M. Chu, “Sharp bounds for Neuman means in terms of two-parameter contra-harmonic and arithmetic mean,” *Journal of Inequalities and Applications*, vol. 2019, no. 1, 2019.
- [34] B. Wang, C.-L. Luo, S.-H. Li, and Y.-M. Chu, “Sharp one-parameter geometric and quadratic means bounds for the Sándor–Yang means,” *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales Serie A Matemáticas*, vol. 114, no. 1, p. 7, 2020.
- [35] Z. Dai and F. Wen, “Another improved Wei-Yao-Liu nonlinear conjugate gradient method with sufficient descent property,” *Applied Mathematics and Computation*, vol. 218, no. 14, pp. 7421–7430, 2012.
- [36] Y. Jiang and J. Ma, “Spectral collocation methods for Volterra-integro differential equations with noncompact kernels,” *Journal of Computational and Applied Mathematics*, vol. 244, pp. 115–124, 2013.
- [37] Z.-F. Dai and F.-H. Wen, “Robust CVaR-based portfolio optimization under a genal affine data perturbation uncertainty set,” *Journal of Computational Analysis and Applications*, vol. 16, no. 1, pp. 93–103, 2014.
- [38] C.-X. Huang, S. Guo, and L.-Z. Liu, “Boundedness on Morrey space for Toeplitz type operator associated to singular integral operator with variable Calderón-Zygmund kernel,” *Journal of Mathematical Inequalities*, vol. 8, no. 3, pp. 453–464, 2007.
- [39] W. Zhou and F. Wang, “A PRP-based residual method for large-scale monotone nonlinear equations,” *Applied Mathematics and Computation*, vol. 261, pp. 1–7, 2015.
- [40] X. Fang, Y. Deng, and J. Li, “Plasmon resonance and heat generation in nanostructures,” *Mathematical Methods in the Applied Sciences*, vol. 38, no. 18, pp. 4663–4672, 2015.
- [41] J. Li, G. Sun, and R. Zhang, “The numerical solution of scattering by infinite rough interfaces based on the integral equation method,” *Computers & Mathematics with Applications*, vol. 71, no. 7, pp. 1491–1502, 2016.
- [42] L. Duan and C. Huang, “Existence and global attractivity of almost periodic solutions for a delayed differential neoclassical growth model,” *Mathematical Methods in the Applied Sciences*, vol. 40, no. 3, pp. 814–822, 2017.

- [43] W. Wang, "On A -stable one-leg methods for solving nonlinear Volterra functional differential equations," *Applied Mathematics and Computation*, vol. 314, pp. 380–390, 2017.
- [44] Y. Tan, C. Huang, B. Sun, and T. Wang, "Dynamics of a class of delayed reaction-diffusion systems with Neumann boundary condition," *Journal of Mathematical Analysis and Applications*, vol. 458, no. 2, pp. 1115–1130, 2018.
- [45] Z. Liu, N. Wu, X. Qin, and Y. Zhang, "Trigonometric transform splitting methods for real symmetric Toeplitz systems," *Computers & Mathematics with Applications*, vol. 75, no. 8, pp. 2782–2794, 2018.
- [46] J. Wang, X. Chen, and L. Huang, "The number and stability of limit cycles for planar piecewise linear systems of node-saddle type," *Journal of Mathematical Analysis and Applications*, vol. 469, no. 1, pp. 405–427, 2019.
- [47] C. Huang, H. Zhang, and L. Huang, "Almost periodicity analysis for a delayed Nicholson's blowflies model with nonlinear density-dependent mortality term," *Communications on Pure & Applied Analysis*, vol. 18, no. 6, pp. 3337–3349, 2019.
- [48] W. Zhou, "On the convergence of the modified Levenberg-Marquardt method with a nonmonotone second order Armijo type line search," *Journal of Computational and Applied Mathematics*, vol. 239, pp. 152–161, 2013.
- [49] W. Zhou and X. Chen, "On the convergence of a modified regularized Newton method for convex optimization with singular solutions," *Journal of Computational and Applied Mathematics*, vol. 239, pp. 179–188, 2013.
- [50] L. Zhang and S. Jian, "Further studies on the Wei-Yao-Liu nonlinear conjugate gradient method," *Applied Mathematics and Computation*, vol. 219, no. 14, pp. 7616–7621, 2013.
- [51] X.-F. Li, G. J. Tang, and B. Q. Tang, "Stress field around a strike-slip fault in orthotropic elastic layers via a hypersingular integral equation," *Computers & Mathematics with Applications*, vol. 66, no. 11, pp. 2317–2326, 2013.
- [52] G. Qin, C. Huang, Y. Xie, and F. Wen, "Asymptotic behavior for third-order quasi-linear differential equations," *Advances in Difference Equations*, vol. 2013, no. 1, 2013.
- [53] C. Huang, Z. Yang, T. Yi, and X. Zou, "On the basins of attraction for a class of delay differential equations with non-monotone bistable nonlinearities," *Journal of Differential Equations*, vol. 256, no. 7, pp. 2101–2114, 2014.
- [54] Y. Liu and J. Wu, "Fixed point theorems in piecewise continuous function spaces and applications to some nonlinear problems," *Mathematical Methods in the Applied Sciences*, vol. 37, no. 4, pp. 508–517, 2014.
- [55] W. Tang and Y. Sun, "Construction of Runge-Kutta type methods for solving ordinary differential equations," *Applied Mathematics and Computation*, vol. 234, pp. 179–191, 2014.
- [56] D. Xie and J. Li, "A new analysis of electrostatic free energy minimization and Poisson-Boltzmann equation for protein in ionic solvent," *Nonlinear Analysis: Real World Applications*, vol. 21, pp. 185–196, 2015.
- [57] X.-S. Zhou, "Weighted Sharp function estimate and boundedness for commutator associated with singular integral operator satisfying a variant of Hörmander's condition," *Journal of Mathematical Inequalities*, vol. 9, no. 2, pp. 587–596, 2007.
- [58] Z. Dai, X. Chen, and F. Wen, "A modified Perry's conjugate gradient method-based derivative-free method for solving large-scale nonlinear monotone equations," *Applied Mathematics and Computation*, vol. 270, pp. 378–386, 2015.
- [59] Z. Dai, "Comments on a new class of nonlinear conjugate gradient coefficients with global convergence properties," *Applied Mathematics and Computation*, vol. 276, pp. 297–300, 2016.
- [60] Z. Dai, D. Li, and F. Wen, "Worse-case conditional value-at-risk for asymmetrically distributed asset scenarios returns," *Journal of Computational Analysis and Applications*, vol. 20, no. 2, pp. 237–251, 2016.
- [61] Y. Tan and K. Jing, "Existence and global exponential stability of almost periodic solution for delayed competitive neural networks with discontinuous activations," *Mathematical Methods in the Applied Sciences*, vol. 39, no. 11, pp. 2821–2839, 2016.
- [62] L. Duan, L. Huang, Z. Guo, and X. Fang, "Periodic attractor for reaction-diffusion high-order Hopfield neural networks with time-varying delays," *Computers & Mathematics with Applications*, vol. 73, no. 2, pp. 233–245, 2017.
- [63] W. Wang and Y. Chen, "Fast numerical valuation of options with jump under Merton's model," *Journal of Computational and Applied Mathematics*, vol. 318, pp. 79–92, 2017.
- [64] C. Huang and L. Liu, "Boundedness of multilinear singular integral operator with non-smooth kernels and mean oscillation," *Quaestiones Mathematicae*, vol. 40, no. 3, pp. 295–312, 2017.
- [65] H. Hu and L. Liu, "Weighted inequalities for a general commutator associated to a singular integral operator satisfying a variant of Hörmander's condition," *Mathematical Notes*, vol. 101, no. 5-6, pp. 830–840, 2017.
- [66] M. A. Hanson, "On sufficiency of the Kuhn-Tucker conditions," *Journal of Mathematical Analysis and Applications*, vol. 80, no. 2, pp. 545–550, 1981.
- [67] T. Weir and B. Mond, "Pre-invex functions in multiple objective optimization," *Journal of Mathematical Analysis and Applications*, vol. 136, no. 1, pp. 29–38, 1988.
- [68] S. R. Mohan and S. K. Neogy, "On invex sets and preinvex functions," *Journal of Mathematical Analysis and Applications*, vol. 189, no. 3, pp. 901–908, 1995.
- [69] J. Hadamard, "Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann," *Journal de Mathématiques Pures et Appliquées*, vol. 58, pp. 171–215, 1893.
- [70] A. Ostrowski, "Über die Absolutabweichung einer differenzierbaren Funktion von ihrem Integralmittelwert," *Commentarii Mathematici Helvetici*, vol. 10, no. 1, pp. 226–227, 1937.
- [71] M. A. Latif, S. Hussain, and S. S. Dragomir, "New Ostrowski type inequalities for coordinated convex functions," *Transylvanian Journal of Mathematics and Mechanics*, vol. 4, no. 2, pp. 125–136, 2012.
- [72] N. Alp and M. Z. Sarikaya, "Quantum Hermite-Hadamard's type inequalities for functions on the co-ordinated convex functions," <http://www.researchgate.net/publication/329814922>.
- [73] H. Kalsoom, J.-D. Wu, S. Hussain, and M. A. Latif, "Simpson's type inequalities for coordinated convex functions on quantum calculus," *Symmetry*, vol. 11, no. 6, p. 768, 2019.
- [74] T. Toplu, M. Kadakal, and İ. İşcan, "On n -polynomial convexity and some related inequalities," *AIMS Mathematics*, vol. 5, no. 2, pp. 1304–1318, 2020.
- [75] H. Hudzik and L. Maligranda, "Some remarks on s -convex functions," *Aequationes Mathematicae*, vol. 48, no. 1, pp. 100–111, 1994.