

New Synthesis of Furochromenyl Imidazo [2a-1b] Thiazole Derivatives, Studies on Their Antitumor Activities.

Asmaa A. Magd-El-Din^{1*}, Amira S. Abd-El-All¹, Hanaa M. F. Roaiah¹ and Mashalla M.S. El-Baroudy¹,

¹Chemistry of Natural Products and Microbial Department, National Research Centre, Dokki, Cairo, Egypt.

*E-mail: asmaaaly2003@yahoo.com.

Abstract 4, 9-Dimethoxy-5-oxo-5H-furo [3, 2-g] benzopyran-6-carboxaldehyde 1 was condensed with 2-thiox-4-imidazolinone 2 to form 3. Treatment of 3 with α -chloroacetyl chloride gave 4. Cyclization of 4 with acetic anhydride took place by heating to give 5. Condensation of 5 with aromatic aldehydes gave the arylidene derivatives 6a-c. Coupling of 5 with diazonium salts gave azo derivatives 7a-c. The work was further extended to investigate the behavior of 3 with 1, 2-dichloroethane to give (4Z)-2-(2-chloroethylthio)-4-((4, 9-dimethoxy-5-oxo-5H-furo [3, 2-g] chromen-6-yl) methylene)-1H-imidazol-5(4H)-one 8. Then 8 was cyclized with acetic anhydride to give (6Z)-2, 3-dihydro-6-[(4, 9-dimethoxy-5-oxo-5H-furo-[3, 2-g] chromen-6-yl) methylene] imidazo [2, 1-b] thiazol-5-(6H)-one 9. [Journal of American Science 2010; 6(5):251-256]. (ISSN: 1545-1003).

Key words: Furochromon; arylidene derivatives; azo; antitumor activity

1. Introduction

2-Thioxo-4-imidazolinone derivatives possess mathematical correlation of plasma levels of anticonvulsant drugs in epileptic patients which introduced in 1978 by Abarbanel⁽¹⁾ and use of it as antiasthmatic drugs⁽²⁾. On the other hand, furochromen and flavones were known to possess coronary dilator activities⁽³⁾. Some derivatives of furochromen composite for treating chronic skin or eye diseases which used in ophthalmic drugs and in dermatological diseases⁽⁴⁾. Recently some furochromone derivatives showed potent antispasmodic, and antitumor activities⁽⁵⁻¹⁰⁾.

A compound having both imidazolinone and furochromone moieties could expect to possess marked biological activities. This prompted us to design and synthesis new furochromen imidazo [2, 1-b]thiazole derivatives to study their antitumor activities.

2. Material and Methods

1-Chemistry

Experimental

All melting points were uncorrected. IR spectra recorded on a Pye Unicam SP- 1100 spectrophotometer using KBr discs. The ¹HNMR spectra were recorded on a Varian EM-390-90 MHz spectrometer using DMSO-d₆ as a solvent and TMS as an internal standard. Chemical shifts expressed as δ ppm units. The micro analytical Centre at Cairo University performed the microanalysis. The antitumor activity of the newly compounds were tested at Cancer Biology Department, National Cancer Institute Cairo, Egypt.

General procedure for preparation of 4 and 8

A solution of (4Z)-2-mercapto-4-[(4,9-dimethoxy-5-oxo-5H-furo-[3,2-g]chromen-6-yl)methylene]-1-H-imidazol-5(4H) one (3) (0.01 mole) in a mixture of 2% potassium hydroxide (56 ml) and absolute ethanol(40 ml) was added α -chloroacetyl chloride (0.01 mole) The reaction mixture was reflux on a steam bath for 3hrs then left to cool at room temperature. It acidified with dilute hydrochloric acid. The solid obtain was filtered off and crystallized from ethanol as yellow crystals of 4. *S*-(4Z)-4, 5-dihydro-4-[(4, 9-dimethoxy-5-oxo-5H-furo [3, 2-g] chromen-6-yl) methylen]-5-oxo-1H-imidazol-2-yl-2-chloroethanethioate (4)

4 : m.p 261^oC yields 75%.

Analysis : C₁₉H₁₃ClN₂O₇S

Calculated :C, 50.84; H, 2.92; N, 6.24; S, 7.14; Cl, 7.90

Found : C, 50.61; H, 2.73; N, 6.11; S, 7.31; Cl, 8.01. IR (Cm⁻¹): 3380 (NH); 1720 (ring C=O); 1690 (γ -pyrone C=O); 1640 (C=N).

MS : m,z 448,450

¹H NMR (DMSO-d₆) (δ ppm): 3.8, 3.7 (2s, 6H, 2OCH₃); 4.5 (s, 2H, CH₂); 6.3-6.8 (2s, 2H, 2CH=C); 8.0, 7.2 (2d, 2H, H-2, H-3 furan); 9.2 (s, 1H, NH exchangeable with D₂O).

(4Z)-2-(2-Chloroethylthio)-4-[(4, 9-dimethoxy-5-oxo-5H-furo [3, 2-g]chromen-6-yl) methylen]-1H-imidazol-5(4H)-one (8) crystallized from ethanol as a yellowish green of 8.

8 : m.p. 243^oC yield 65%.

Analysis : C₁₉H₁₃ClN₂O₆S

Calculate: C, 52.46; H, 4.39; N, 6.44; S, 7.38; Cl, 8.16

Found : C, 52.23; H, 4.12; N, 6.23; S, 7.50; Cl, 8.45 IR (Cm⁻¹): 3440 (NH); 1740 (ring C=O); 1665 (γ -pyrone C=O); 1640 (C=N); 1243 (C-S).

MS : m/z 434, 436

¹H NMR (DMSO-d₆) (δ ppm) : 3.8, 3.9 (2s, 6H, 2OCH₃); 3.3 (t, CH₂-S); 4.1 (t, CH₂-Cl);

6.2, 7.1 (2s, 2H, 2CH=C); 8.1, 7.2 (2d, 2H, H-2, H-3 furan); 8.7 (s, 1H, NH exchangeable with D₂O).

General procedure for preparation of 5 and 9

A suspension of each of 4, 8 (0.01 mole) in acetic anhydride (30 ml) was refluxed for 4hrs. The reaction mixture allowed cooling, and then the mixture poured into cold water. The product obtained was filtered off and crystallized from ethanol as yellowish green, dark brown crystals for 5 and 9 respectively.

(6Z)-6-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylen]imidazo [2,1-b]thiazol-2,5(3H,6H)-dione (5) crystallized from ethanol as a yellowish green of 5.

5 : m.p. 272°C yield 70%.

Analysis : C₁₉H₁₂N₂O₇S

Calculate : C, 55.34; H, 2.93; N, 6.79; S, 7.78

Found : C, 55.60; H, 2.72; N, 6.54; S, 8.03

IR (Cm⁻¹): 1720-1710(two ring C=O); 1680 (γ - pyrone C=O); 1640 (C=N).

MS : m/z 434, 436

¹HNMR (DMSO-d₆)

(δppm):3.7,3.6(2s,6H,2OCH₃);4.1(s,2H,CH₂);6.4,6.7(2s,2H,2CH=C);7.1,7.9 (2d,2H,H-2,H-3furan)

(6Z)-2, 3-dihydro-6-[(4, 9-dimethoxy-5-oxo-5H-furo [3, 2-g]chromen-6-yl) methylen] imidazo[2,1-b]thiazol-5(6H)-one (9) crystallized from ethanol as dark brown of 9.

9 : m.p. 255°C yield 75%.

Analysis : C₁₉H₁₄N₂O₆S

Calculated : C, 62.86; H, 3.07; N, 6.11; S, 7.00

Found : C, 63.01; H, 2.93; N, 5.89; S, 7.40

IR (Cm⁻¹) : 1736 (ring C=O); 1655 (γ -pyroneC=O); 1648(C=N); 1253(C-S

MS : m/z 398

¹HNMR (DMSO-d₆)

(δppm):3.6,3.9(2s,6H,2OCH₃);3.2,3.7(2t,2CH₂);7.0,6.3 (2s, 2H, 2CH=C);8.1,7.2 (2d,2H,H-2,H-3 furan).

Reaction of (5) with aromatic aldehydes

A mixture of 5 (0.005 mole) fused sodium acetate (2.5g) and a slight excess of aromatic aldehyde (0.005 mole) benzaldehyde, chlorobenzaldehyde, bromobenzaldehyde in (25ml) glacial acetic acid were refluxed for 2 hrs. The reaction mixture poured over cooled water, and then separated solid filtered off washed with water and crystallized from acetic acid to give greenish yellow, olive green and dark brown for 6a-c respectively.

(3E,6Z)-3-benzylidene-6-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylen] imidazo[2,1-b]thiazol-2,5(3H,6H)-dione (6a) crystallized from acetic acid as greenish yellow of 6a.

6a : m.p. 281°C yield 65%.

Analysis : C₂₆H₁₆N₂O₇S

Calculated : C, 62.34; H, 3.19; N, 5.59; S, 6.39

Found : C, 62.50; H, 3.12; N, 5.82; S, 6.25

IR (Cm⁻¹) : 1715, 1700 (two ring C=O); 1680 (γ - pyrone C=O); 1635(C=N)

MS : m/z 500

¹HNMR(DMSO-d₆)(δppm):

3.8,3.6(2s,6H,2OCH₃);6.3,6.5,6.8(3s,3H,3CH=C);7.1,7.8(2d,2H,H- 2,H-3furan);7.3-7.6 (m,5H,aromatic protons).

(3E,6Z)-3-(7-chlorohepta-2,4,6-triynylidene)-6-[(4,9-dimethoxy-5-oxo-5H-furo [3,2-g]chromen-6-yl)methylen]imidazo[2,1-b]thiazol 2,5(3H,6H)-dione (6b) crystallized from acetic acid as olive green of 6b.

6b : m.p. 259°C yield 70 %.

Analysis : C₂₆H₁₅ClN₂O₇S

Calculated :C, 58.38; H, 2.83; N, 5.24; S, 5.99; Cl, 6.63

Found : C, 58.21; H, 3.10; N, 5.41; S, 6.23; Cl, 6.42

IR (Cm⁻¹) : 1720, 1710 (two ring C=O); 1690 (γ - pyrone C=O); 1640 (C=N).

MS : m/z 534,536

(3E,6Z)-3-(7-bromohepta-2,4,6 triynylidene)-6-[(4,9-dimethoxy-5-oxo-5H-furo [3,2-g]chromen-6-yl)methylen]imidazo[2,1-b]thiazol 2,5(3H,6H)-dione (6c) crystallized from acetic acid as dark brown of 6c.

6c : m.p. 247°C yield 75%

Analysis : C₂₆H₁₅BrN₂O₇S

Calculated :C, 53.90; H, 2.61; N, 4.84; S, 5.53; Br, 13.79

Found : C, 54.17; H, 2.86; N, 5.11; S, 5.81; Br, 13.53

IR (Cm⁻¹) : 1715, 1705 (two ring C=O); 1685 (γ - pyrone C=O); 1635(C=N).

MS : m/z 578,580

Reaction of 5 with diazotised aromatic amines:-

In a very cold condition, a solution of (0.01 mole) of the appropriate diazotised aromatic amines (prepared from the equivalent amounts of the amine, HCl and NaNO₂) was gradually added to a cold solution of 5 (0.01 mole) in aqueous sodium hydroxide solution (2%, 20 ml) in about 15 min. The reaction mixture kept in the ice-chest for 2hrs with constant stirring; the solid product collected by filtration, washed with water then crystallized from appropriate solvent to give reddish brown, dark brown and brown crystals of compounds 7a-c respectively.

(6Z)-3-(2-phenyldiazenyl)-6-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylen]imidazo[2,1-b]thiazol-2,5(3H,6H)-dione (7a) crystallized from acetone as reddish brown of 7a.

7a : m.p. 278°C yield 70%
 Analysis : C₂₅H₁₆N₄O₇S
 Calculated : C, 58.13; H, 3.12; N, 10.84; S, 6.21
 Found : C, 57.91; H, 3.32; N, 11.00; S, 6.13.
 IR (Cm⁻¹):3370 (NH); 1720,1700 (two rings C=O);1680 (γ - pyroneC=O);1635(C=N)
 MS : m/z 516
¹HNMR(DMSO-d₆)(δppm) : 3.7, 3.6(2s,6H,2OCH₃); 6.4, 6.6(2s,2H,2CH=C);7.2, 7.7(2d,2H,H-2,H-3 furan); 7.3-7.6 (m,5H,aromatic protons and 12.4(s,1H,NH).

(6Z)-3-(2-(6-chlorohexa-1,3,5-triynyl)diazenyl)-6-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylen]imidazo[2,1-b]thiazol-2,5(3H,6H)-dione (7b)crystallized from chloroform as dark brown of 7b.

7b : m.p. 263°C yield 70%
 Analysis: C₂₅H₁₅ClN₄O₇S
 Calculated: C, 54.49; H, 2.72; Cl, 6.44; N, 10.17; S, 5.82.
 Found : C, 54.27; H, 2.95; Cl, 6.2; N, 10.34; S, 6.11.
 IR (Cm⁻¹):3390 (NH); 1725,1710 (two rings C=O);1690 (γ pyroneC=O);(C=N)
 MS : m/z 551,553
 (6Z)-3-(2-(6-bromohexa-1,3,5-triynyl)diazenyl)-6-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylen]imidazo[2,1-b]thiazol-2,5(3H,6H)-dione (7c) crystallized from chloroform as brown of 7c.

7c : m.p. 262°C yield 75%
 Analysis : C₂₅H₁₅BrN₄O₇S
 Calculated: C, 50.41; H,2.54; Br,13.42; N,9.41; S,5.93
 Found : C, 50.67; H,2.76; Br,13.16; N,9.59; S,5.63.
 IR (Cm⁻¹) :3380 (NH); 1716, 1700 (two ring C=O); 1685 (γ -pyrone C=O); 1635 (C=N).
 MS : m/z 595,593

2-Antitumor

Different concentration of the tested compounds between 1-10 µg/ml were added to the cell monolayer using SRB ASSAY (Sulfrhodamine B stain), and compared with the standard drug Doxorubicin DXR⁽¹⁹⁾ using the method of Skehan et al⁽²⁰⁾.

The antitumor activity of the new formed compounds were tested at Cancer Biological Department, National Cancer Institute, Cairo, Egypt .

Results and Discussion

1-Chemistry

4,9-Dimethoxy-5-oxo-5H-furo[3,2-g]benzopyran-6-carboxaldehyde (1)⁽¹¹⁾ condensed with 2-thio-4-imidazolinone (2)⁽¹²⁾ to give (4Z)-2-mercapto-4-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylene]-1-H-imidazol-5-(4H)-one (3). The reaction product (3) was formed via the condensation of the formyl group of (1) with active methylene group of (2).

Treatment of (3) with α-chloroacetyl chloride gave S-(4Z)-4, 5-dihydro-4-[(4, 9-dimethoxy-5-oxo-5H-furo [3, 2-g] chromen-6-yl)methylene]-1H-imidazol-2-yl-2-chloro-ethanethioate (4). Compound (4) confirmed by elemental analysis and spectral data.


When compound (4) was heated with acetic anhydride, cyclization took place and (6Z)-6-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylene]imidazo[2,1-b]thiazole-2,5(3H,6H)-dione (5) was obtained via loss of HCl, the IR and ¹H NMR spectrum of (5) was characterized by the absence of NH proton. The cyclic structure proposed for compound (5) was the favor one.

Moreover, compound (5) having an active methylene group adjacent to carbonyl group was condensed with aromatic aldehydes (benzaldehyde chlorobenzaldehyde, bromobenzaldehyde in glacial acetic acid in presence of fused sodium acetate at 140°C to give (3E,6Z)-3-benzylidene-6-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylene]imidazo[2,1-b]thiazole-2,5-(3H,6H)-dione derivatives (6a-c). The arylidene derivatives (6a-c) showed the correct analytical values, their UV absorption spectra were studied (6a) absorbed at 358 nm (ε= 1553). This absorption is compatible with the benzene ring conjugated with C=C bond which in turn is conjugated with carbonyl group⁽¹³⁾ Exhibiting another property of active methylene groups. Compound (5) reacted in sodium hydroxide solution with aromatic diazonium compounds to give (3E,6Z)-3-(aryl-hydrazone)-6-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylene]imidazo[2,1-b]thiazole-2,5(3H,6H)-dione derivatives (7a-c). However, it is generally assumed that hydrazone is the stable form, whenever, coupling occurs at a methylene carbon atom (Scheme 1).

Wiley and Jarboe⁽¹⁴⁾ and Tanner⁽¹⁵⁾ have presented the IR absorption data which corroborated the above view, and in addition the presence of maximum band at 410 nm in the UV spectrum of (7a) which proved that (7a) exists in the hydrazone form rather than the azo form^(16,17) (Scheme 1).

The work was further extended to investigate the behavior of 3 with 1,2-dichloroethane to give (4Z)-2-(2-chloroethylthio)-4-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-yl)methylene]-1H-imidazol-5(4H)-one(8) which upon cyclization with acetic anhydride gave (6Z)-2,3-dihydro-6-[(4,9-dimethoxy-5-oxo-5H-furo[3,2-g]chromen-6-

yl)methylene]imidazo[2,1-b]thiazol-5(6H)-one(9) by elimination of HCl (Scheme 2). The structure assigned for the cyclized product based on the correct analytical and spectral data. The $^1\text{H-NMR}$ spectrum of the above compound revealed the absence of the NH group.




Azo-form

Hydrazo-form

a, Ar=C₆H₅; b, Ar=C₆H₄-Cl-p; c, Ar=C₆H₄-Br-p

Scheme-1-



Scheme 2

Antitumor activity

The cytotoxic activity:


All the newly synthesized compounds were tested for their cytotoxic activity using tumor cell Lines ⁽¹⁸⁾ ,[HEPG2 (Human Liver Carcinoma Cell Line) and MCF7 (Human Breast Carcinoma Cell Line)].

2-Antitumor

The cytotoxic activity of the tested compounds on HEPG2 and MCF7 were expressed as IC50, table (I), where IC50 (UM) is the dose of compound which reduces survival to 50%. The relation between the surviving fraction and drug concentration plotted to get the survival curve of the tumor cell line. The tested compound showed this activity only at the specified concentration and this cell lines.c.f.Table (I)

Table (I):

Compound No.	Cell lines	
	HEPG2 IC50	MCF7 IC50
1	-ve	0.769
2	-ve	0.694
3	-ve	0.731
4	-ve	0.769
5	-ve	0.694
6a	-ve	0.656
6b	-ve	0.806
7b	3.68	0.769
8	-ve	0.769
9	4.95	0.731



The standard curves for the most active compounds and the standard drugs Doxorubicin (DXR) are given below.

Conclusion:

All the tested compounds showed remarkable antitumor activity against human MCF7 cell line. Compound 6b was the most potent one comparing with the standard drug DXR. The following compounds 6_a<2, 5<3, 9<1,4,7_b,8 showed varying activity in an increasing order. When those compounds tested against human HEPG2 cell line, compounds 7_b, 9 showed moderate activity. Compounds 1, 2, 3, 4, 5, 6_{a,b}, 8 has no activity, comparing with the standard drug DXR.

References:

1. Abarbanel, J., Herishanu, Y., Rosenberg, P., and Elyath, U., : In vivo interaction of anticonvulsant drugs. The mathematical correlation of plasma levels of anticonvulsant drugs in epileptic patients. *J. Neurol.*218, 137(1978),C. A.89, 140641x. (1978).
2. Jamieson, W.B., Ross, W.J., Simmonds, R. G., and Verge, J.P., Hydantion compounds formulation. Containing them and their use as pharmaceuticals. *Eur. Pat Appl.*5,647 (Cl. Co7 D233/74);28 Nov.1979 Brit. Appl. , 78/21, 352/23 May. 1978; C.A.93,71772j(1980).
3. Mantaxd, I., Berger, T., and Silaghi, E., The synthesis of some chromones and flavones, coronary dilators. *Rev. Chim. (Bucharest)* 10, 69 (1959).
4. Bell, J. H., Clarke, C., Walter, F., Taylor, J. E., and Sullivan T. J. (*Fisons Ltd.*): Composition for treating chronic skin or eye diseases. Ger. Offen. 2, 634 (1) (CIA61K31/35); 14 Apr. 1977 Brit. Appl. .03/507, 75/40 Oct. 1975.
5. Atassi, G., Briet, P., Berthelon, J. J., Collonges, F., Synthesis and antitumor activity of some (8-substituted 4-oxo-4H-1-benzopyrans). *Eur. J. Med. Chem. Chim. Ther.* , 20, 393-402(1985).
6. Hudson J., Towers G.H.N., phytomedicines as antivirals. *drugs Future* 24, 295 (1999)
7. Galal, S.A., Abd El All, A.S., Abdallah, M.M., and El-Diwan, H.I., Synthesis of potent antitumor and antiviral benzofuran derivatives. *Bioorg. Med. Chem. Lett.* 19, 2420-2428 (2009).
8. Fawzy N.M., Swelam S.A., Batran S.A.,Synthesis and Pharmaceutical activity of new Series of Chromonyl derivatives *Boll.Chim.farm.*143, 24-33 (2004).
9. Fawzy N.M., Synthesis of Novel Dihydropyridine, Dihydropyrimidine, Dithoacetal and Chalcone derivatives from formylchromones *Boll.Chim.farm.*143, 70-77 (2004).
10. El-Sawy E. R., Mandour A.H., Islam I.E., Abo-Salem H. M., Synthesis and Antimicrobial Evaluation of Some New 6- Substituted Furobenzopyrone derivatives *Egypt. J Chem.*51, No.4,523-538 (2008).
11. Eidin F., Schlemann J., darstellung und reaktionen von 6-acylhellin-derivaten.Fritz Eiden,Jurgen Schunemann *Arch Pharm (Weinheim)* 201, 316 (1983).
12. Klason, P., *Chem. Ztg.* 14, 543 (1980) Cf. also Tiwari, S. S., and Swaroop, A., Search for new oral hypoglycemic agents. part V j. Ind. *J. Ind. Chem.. Soc.* 40, 693 (1963).
13. Mustafa, A., Asker, W., Harhash, A.H., and Fleifel, A.M., search for new oral hypoglycemic agents. part V, Reactivity of unsaturated centers in heterocycles and chalkones towards diazoalkanes. *Tetrahydron[London]*,21,2215 (1965).
14. Wiley, R.H., and Jarboe, C.H., 2-Pyrones..XII.γ-Keto-β-methylglutaconic anhydride. *J. Amer. Chem. Soc.* 77, 403(1955).
15. Tanner, E.M., the inferared absorption spectra of some α-phenylhrdrazone-ketones and alcohols(p20-26)v(15). *Spectrochim. Acta [oxford]* 15, 20-26(1959).
16. Yao, H.C., The Coupling of Diazonium ion with β-Diketones. *J.Org Chem.* 29, 2959 (1964).
17. Depuy, C.H., and Wells, P.P.R., Cyclopentene-3,5-dione.III.Reactions of the Methylene Group^{1,2} *J. Am. Chem. Soc.* 82, 2909 (1960).
18. Macleod, R., Skehan, et al., Widespread interspecies cross of human tumor cell lines *International Journal Cancer.* 83, 555-563 (1999).
19. Groopman, Jerome, E., *How Doctors Think. Boston: Houghton Mifflin. ISBN,* 49, 6018-61003 (2007).
20. Monks, A. ,Scudiero, D., Skehan, P., Shoemaker, R., Paull, K., Vistica, D., Hose, C., Langley, J., Cronise, P., Vaigro-wolff, A., Gray-Goodrich, M., Campbell, H., Mayo, J., and Boyd, M., Feasibility of a high-flur anticancer drug screen using a diverse panel of cultured human tumor cell lines. *J. Nat. Cancer Inst.* 83, 757-766(1991).

3/5/2010