

Published in final edited form as:


J Am Chem Soc. 2009 September 30; 131(38): 13606–13607. doi:10.1021/ja906472m.

A Nine-Step Enantioselective Total Synthesis of (+)-Minfiensine


Spencer B. Jones, Bryon Simmons, and David W. C. MacMillan*

Merck Center for Catalysis at Princeton University, Princeton, New Jersey 08544

Abstract






Diels–Alder/Amine Cyclization Cascade



An enantioselective total synthesis of the *Strychnos* alkaloid (+)-minfiensine has been accomplished. Prominent features of this synthesis include (i) a new enantioselective organocatalytic Diels–Alder/amine cyclization sequence to build the central tetracyclic pyrroloindoline framework in four steps from commercial materials and (ii) a 6-*exo*-dig radical cyclization to forge the final piperidinyl ring system. This total synthesis of (+)-minfiensine was completed in nine chemical steps and 21% overall yield.

Over the last 50 years, the *Strychnos* alkaloids have become recognized as a family of molecular benchmarks, used to calibrate the utility of new complexity-generating reactions or novel synthetic strategies.¹ In this context, minfiensine (**1**),² a structurally unique isolation product from *Strychnos minfiensis*, has received considerable attention from the chemical synthesis community,³ culminating recently in the highly elegant (and first) enantioselective total synthesis by Overman and coworkers.^{3a} Minfiensine is characterized by an embedded 1,2,3,4-tetrahydro-9a,4a-(iminoethano)-9*H*-carbazole moiety, a tetracyclic core that is also found among related akuammiline alkaloids vincorine (**2**) and echitamine (**3**).⁴ In this communication, we document a new organocatalytic Diels–Alder/amine cyclization sequence that allows rapid and enantioselective access to this tetracyclic carbazole framework using only an amine catalyst, propynal, and a simple tryptamine derivative. Moreover, we elaborate upon this new asymmetric protocol to achieve a nine-step total synthesis of minfiensine beginning from commercial materials.



dmacmill@princeton.edu.

Supporting Information Available. Experimental procedures and spectral data for all new compounds are provided (21 pages) (PDF).

Design Plan

As outlined in Scheme 1, we envisioned two key steps that would allow rapid access to the complete pentacyclic topography of minfiensine. From a disconnection approach, we assumed the fifth and final ring of the natural product might be forged via a 6-*exo*-dig cyclization between an allylic radical and a pendent alkyne **4**, a carbon-carbon bond union that would also create the requisite stereogenicity at C(15) and the exocyclic olefin at C(20).⁵ As for our second key step, we hypothesized that the structurally complex pyrroloindoline tetracycle **5** might arise directly from vinyl indole **6** via a cascade reaction that would incorporate an organocatalytic Diels–Alder cycloaddition, enamine to iminium isomerization, and an amine cyclization sequence.⁶ More specifically, we proposed that condensation of secondary amine catalyst **7** with propynal should generate an activated iminium ion with the acetylenic group being partitioned away from the bulky *t*-butyl substituent of the catalyst framework (Scheme 2, **TS-A**). In this conformation, the aryl ring would shield the top face of the reactive alkyne, facilitating an *endo*-selective⁷ Diels–Alder cycloaddition with 2-vinylindole **8** in a regioselective manner to produce the tricyclic diene **9**. Protonation of the enamine moiety would then give rise to an iminium ion **10**, facilitating a 5-*exo* amine heterocyclization to deliver the tetracyclic pyrroloindoline **11**. As a key design element, the dienyl substructure of indole **8** incorporates a 1-methyl sulfide substituent, a moiety that we hoped would accelerate the Diels–Alder cycloaddition while providing a latent handle for radical formation,⁸ as required in the final ring-forming step.

Synthesis of the minfiensine core began with production of the requisite [4+2] cycloaddition substrate **8** in three steps from commercial materials using the standard procedures outlined in Scheme 3. With this material in hand, we were able to examine the pivotal Diels–Alder-cyclization cascade in detail. As shown in Table 1, we were delighted to find that subjection of 2-vinylindole **8** to propynal in the presence of imidazolidinone catalyst **7** produced the desired tetracycle **15**, albeit with moderate selectivity (entry 1, 75% ee). A catalyst structure evaluation revealed that the 1-naphthyl substituted catalyst **14** in conjunction with tribromoacetic acid (TBA) co-catalyst provided superior levels of yield and enantioselectivity, presumably due to the extended shielding effect of the naphthyl ring in the [4+2] transition state (entry 3, 87% yield, 96% ee). It is important to note that catalyst loadings as low as 5 mol % were sufficient to effect the cascade while maintaining high levels of reaction efficiency (entries 4 and 5, 80% yield, 94% ee).

Subsequent conversion of the pyrroloindoline tetracycle **15** to minfiensine (**1**) was achieved in a five-step sequence as shown in Scheme 3. Simultaneous *N*-Boc deprotection and primary alcohol protection were performed by exposure of carbamate **15** to TESOTf in acetonitrile at 0 °C to afford silyl ether **16** in 84% yield. Reductive amination of secondary amine **16** with butynal *t*-butyl sulfide **17** was readily accomplished with NaBH(OAc)₃ in CH₂Cl₂ to render the requisite radical cyclization substrate **18** in 96% yield. At this stage, we were surprised to find that all attempts to forge the final piperidine ring of minfiensine via alkyne radical cyclization were unsuccessful using prototypical conditions (AIBN, Bu₃SnH).⁹ However, replacement of Bu₃SnH with the more bulky *t*-Bu₃SnH¹⁰ (with AIBN in refluxing toluene) cleanly afforded the allene **19** in 61% yield.^{11,12} Next, we envisioned that regio- and diastereoselective allene hydrogenation would provide the *trans*-ethylidene subunit that is commonly found throughout the *Strychnos* alkaloid family. In the event, we were pleased to find that the proposed reduction could be realized via subjection of allene **19** to 10% Pd/C and H₂ in THF at –15 °C to give the desired *trans*-ethylidene with greater than 20:1 E:Z selectivity.¹³ Finally, a global deprotection using neat TFA at room temperature delivered (+)-minfiensine (**1**) in 90% yield, a substance that was identical in all respects to the natural isolate.

In summary, the total synthesis of (+)-minfiensine (**1**) was completed in nine steps and 21% overall yield from commercial materials. Prominent features of this synthesis include (i) a new cascade organocatalysis sequence to build the central tetracyclic pyrroloindoline framework and (ii) a 6-*exo*-dig radical cyclization to forge the final piperidiny ring system.

Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

Acknowledgement



Financial support was provided by the NIHGM (R01 GM078201-01-01) and kind gifts from Merck and Amgen. B.S. and S.B.J. would like to thank Merck and Bristol-Myers Squibb respectively for graduate fellowships.

References


1. Bonjoch J, Solé D. Chem. Rev 2000;100:3455. [PubMed: 11777429]
2. Massiot G, Thépenier P, Jacquier M, Le Men-Olivier L, Delaude C. Heterocycles 1989;29:1435.
3. Total syntheses: (a)Dounay AB, Overman LE, Wrobleski AD. J. Am. Chem. Soc 2005;127:10186. [PubMed: 16028927] (b)Dounay AB, Humphreys PG, Overman LE, Wrobleski AD. J. Am. Chem. Soc 2008;130:5368. [PubMed: 18303837] (c)Shen L, Zhang M, Wu Y, Qin Y. Angew. Chem. Int. Ed 2008;47:3618. Partial synthesis: dBobeck DR, France S, Leverette CA, Sánchez-Cantalejo F, Padwa A. Tetrahedron. Lett 2009;50:3145.
4. (a)Ramírez A, García-Rubio S. Curr. Med. Chem 2003;10:1891. [PubMed: 12871110] (b) For a total synthesis of racemic vincorine, see: Zhang M, Hunag X, Shen L, Qin Y. J. Am. Chem. Soc 2009;131:6013. [PubMed: 19326910]
5. Sha CK, Zhan ZP, Wang FS. Org. Lett 2000;2:2011. [PubMed: 10891217]
6. (a) It should be noted that a racemic, thermally promoted Diels-Alder/amine cyclization sequence between a 2-vinylindole and dimethyl maleate has previously been reported: Lévy J, Sapi J, Laronze JY, Royer D, Toupet L. Synlett 1992;601 (b) For iminium catalyzed pyrroloindoline formation see: Austin JF, Kim S-G, Sinz CS, Xiao W-J, MacMillan DWC. Proc. Natl. Acad. Sci 2004;101:5482–5486. [PubMed: 15067109]
7. Acetylenic [4 + 2] cycloadditions are typically *exo*-selective: (a)Corey EJ, Lee TW. Tetrahedron. Lett 1997;38:5755. (b)Ishihara K, Kondo S, Kurihara H, Yamamoto H. J. Org. Chem 1997;62:3026. [PubMed: 11671675]
8. Gutierrez C, Summerhays LR. J. Org. Chem 1984;49:5206.
9. Only alkyne stannylation was observed under these conditions.
10. Pike PW, Gilliat V, Ridenour M, Hershberger JW. Organometallics 1988;7:2220.
11. Bachi MD, Bar-Ner N, Melman A. J. Org. Chem 1996;61:7116. [PubMed: 11667614]
12. Cyclization onto the analogous methyl alkyne gives a 1:1 *E*:*Z* ratio.
13. A related allene hydrogenation has been achieved in modest yield: Bonjoch J, Solé D, García-Rubio S, Bosch J. J. Am. Chem. Soc 1997;119:7230.



Scheme 1.
Retrosynthetic Approach to Minfiensine Pentacycle.



Scheme 2.
Enantioselective Catalytic Cascade Sequence to Core.




Scheme 3. Nine-Step Total Synthesis of Minfiensine (1)^a

^a Reagents and Conditions: (a) NaH, PMBCl, DMF, 0 °C. (b) *n*-BuLi, THF, −78 °C; then DMF, −78 °C to rt. (c) (EtO)₂P(O)CH₂SMe, NaH, THF, 0 °C to rt. (d) Table 1, entry 3. (e) TESOTf, MeCN, 0 °C. (f) 4-(*tert*-butylthio)but-2-ynal (17), NaBH(OAc)₃, CH₂Cl₂, rt. (g) 3 equiv. *t*-Bu₃SnH, 0.3 equiv AIBN, toluene, 110 °C. (h) Pd/C, H₂, THF, −15 °C; >20:1 E:Z selectivity. (i) PhSH, TFA, rt.

Table 1

Organocatalytic Diels–Alder–Cascade Cyclization Studies.



entry	catalyst•HA	mol%	time (h)	% yield ^a	% ee ^b
1	7 ·TFA	20	1	78	75
2	7 ·TBA	20	1	81	88
3 ^c	14 ·TBA	15	24	87 ^d	96
4	14 ·TBA	10	48	80	94
5	14 ·TBA	5	72	80	94

^aYield determined by ¹H NMR with internal standard.

^bEnantiomeric excess determined by chiral SFC analysis.

^cAt -50 °C.

^dIsolated yield.