



UPPSALA
UNIVERSITET

DiVA 

<http://uu.diva-portal.org>

This is a preprint version of a paper published in *Acta Anaesthesiologica Scandinavica*. The definitive version is available at www3.interscience.wiley.com

Citation for the published paper:

Miclescu, Adriana and Gordh, Torsten

" Nitric oxide and pain: "Something old, something new"

Acta Anaesthesiologica Scandinavica, 2009, 53 (9)

1107-1120 pp

URL: <http://dx.doi.org/10.1111/j.1399-6576.2009.02054.x>

Access to the published version may require subscription.





Nitric oxide and pain

Journal:	<i>Acta Anaesthesiologica Scandinavica</i>
Manuscript ID:	AAS-08-0284.R1
Manuscript Type:	Review Article
Date Submitted by the Author:	22-Dec-2008
Complete List of Authors:	Miclescu, Adriana; Uppsala University Hospital, Anesthesia and Intensive therapy Gordh, Torsten; Uppsals University, Anaestehsiology and Intensive care
Key Words:	nitric oxide, NOS (nitric oxide synthases), NOS cofactors, reactive oxidative species, NO-cGMP pathway, pain treatment, cyclo-oxygenase-inhibitor NO donors (CINODs), NO-donors, NOS inhibitors



Review

1
2
3
4
5
6
7
8 *Review article*
9

10
11 **Nitric oxide and pain**

12 *“Something old, something new“*
13
14
15
16
17
18
19
20
21
22
23
24
25

26 Adriana Miclescu^{1,2} MD, PhD, DEAA, Torsten Gordh^{1,2} MD, PhD, DEAA
27 Department of Surgical Sciences/Anesthesiology and Intensive Care Medicine¹,
28 Pain Clinic²
29

30
31 Uppsala University Hospital, SE-75185 Uppsala, Sweden
32
33
34
35
36
37
38
39
40
41
42
43

44 Key words: nitric oxide (NO), NOS (nitric oxide synthases), NOS cofactors,
45 reactive oxidative species, NO-cGMP pathway, pain treatment, cyclo-
46 oxygenase-inhibitor NO donors (CINODs), NO-donors, NOS inhibitors
47
48
49
50
51
52
53

54 Correspondence: Adriana Miclescu, as above
55 E-mail: adriana.miclescu@akademiska.se
56
57
58
59
60

Abstract

Challenges have emerged following the revival of nitric oxide (NO) from “something old”, a simple gas derived from nitrogen and oxygen with a role in the early stages of evolution, into “something new”, an endogenously formed biological mediator regulating a wide variety of physiological functions. Although pain is a common sensation, it encompasses multiple neurobiologic components of which NO is only one. In pain research, the study of NO is complicated by convoluted problems related mostly to the effects of NO, which are pro- or antinociceptive depending on the circumstances. This dual function reflects the two faces of the NO molecule described in physiology. This review covers current information about NO and its implications in pain mechanisms. In addition, it follows the pain pathways, demonstrating the role of NO in peripheral nociceptive transmission as well in central sensitization. This knowledge may provide the scientific basis for developing new drugs that are indicated for different types of pain, drugs that may be related to the chemical links of NO. A comprehensive approach to understanding the effects of NO will help clinicians identify novel agents that combine the pharmacological profile of native drugs with a controllable manner of NO release. Inhibitors of NO synthesis may have analgesic effects and would be of interest for treating inflammatory and neuropathic pain. Many specific NO inhibitors have been described and the most promising substances have been studied in animals. Unfortunately, only a few of these compounds have reached the stage of clinical pain trials.

Summary:

1. Introduction
2. Nitric oxide (NO) biosynthesis and pain
 - 2.1. NO biosynthesis- basic physiology
 - 2.2. The role of different NOS (nitric oxide synthases) isoforms in pain
 - 2.3. The potential role of tetrahydrobiopterin (BH4) in pain
3. Reactive oxidative (oxygen/nitrogen) species and pain
4. The NO-cGMP pathway and pain
 - 4.1. Basic physiology
 - 4.2. The NO-cGMP pathway and pain
5. Link between NO and NMDA
6. Peripheral effects of NO
7. Central nervous system (CNS) effects of NO
8. Cyclo-oxygenase (COX) and NO
9. Potential implications of NO in pain treatment
 - 9.1. NO-releasing drugs
 - 9.1.1. Cyclo-oxygenase-inhibiting nitric oxide donating drugs (CINODs)
 - 9.1.2. NO donating drugs (NODDs)
 - 9.1.3. NO-donors (nitroglycerine and isosorbid dinitrate)
 - 9.2. NO inhibitors
 - 9.2.1. NOS inhibitors
 - 9.2.2. Inhibitors of Soluble Guanylyl/Guanylate Cyclase
10. Concluding remarks
11. References

1 2 3 **I. Introduction** 4

5 The revival of nitric oxide (NO) indicates that “something simple and something old” may
6 now be “new” again. It should be remembered that NO is a simple gas (1) derived from
7 nitrogen and oxygen, and it played a crucial role in the early stages of evolution (2). NO may
8 have constituted a critical defence mechanism for primitive microorganisms by counteracting
9 oxidative destruction and giving them an evolutionary advantage (2). Not only is NO an
10 ancient and widely used regulator of the life history of different species of eukaryotes and
11 solitary ascidians (3), but the presence of NO was observed in plants much earlier than in
12 animals (4). Discovered in the 17th century by Jan B. Helmont (5) and studied under the name
13 “phlogisticated nitrous air” by Joseph Priestley (6), NO was regarded for many years as an
14 environmental pollutant (7). It became “something new” in 1992 when it received intense
15 media coverage due to its stature as a biological messenger (8) and became the molecule of
16 the year (9). The 1998 Nobel Prize in Physiology or Medicine was awarded for “the first
17 discovery that a gas can act as a signal molecule”, emphasising that NO is a biological
18 mediator produced by mammalian cells (10). The fascinating history of NO thereafter
19 continued, with extensive research showing that NO plays an important role in most human
20 organ systems, and in neurotransmission, immune defence, regulation of death cells and cell
21 motility (10).
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 It was reported that NO modulates spinal and sensory neuron excitability that contributes to
46 different pain states. However, investigating the implications of this molecule in nociception
47 remains a challenging task. The aim of the present review is to summarise the contributions of
48 NO to pain mechanisms, both in peripheral and central nociceptive transmission, and to
49 elucidate therapeutic implications of its action.
50
51
52
53
54
55
56

57 **2. Nitric oxide biosynthesis and pain**

58 **2.1. Basic physiology** 59 60

1
2
3 In biological systems, NO is a molecular gas generated from L-arginine (11) and molecular
4 oxygen by the activity of nitric oxide synthase (NOS). L-citrulline results from this reaction
5 and is shunted into a metabolic pathway that regenerates free L-arginine (Fig.1). Furchgott
6 first identified NO synthases implicated in NO generation in experiments on the aortas of
7 rabbits (12). The different forms of NO synthase have been identified and named based upon
8 the cells from which they were first isolated: neuronal NOS (nNOS, type 1 NOS) (13);
9 inducible NOS, induced in macrophages upon stimulation with bacterial endotoxines and
10 cytokines (iNOS, type 2 NOS) (14); endothelial NOS (eNOS, type 3 NOS) (15); and in some
11 tissues they may exist in mitochondria (mNOS) (16). The NOS enzymes differ not only in
12 their localization, but also in their function. The two calcium dependent constitutive NOSs-
13 eNOS and nNOS, generate small amounts of NO, while iNOS production that is independent
14 of intracellular calcium concentrations generates NO levels able to damage cells and
15 microorganisms (17). These enzymes are distinct proteins encoded by genes on disparate
16 chromosomes, but all of them have the same genomic structure and all of them facilitate the
17 addition of the guanidino nitrogen of the amino acid arginine to molecular oxygen, producing
18 NO and water (Fig.1) (15). The enzyme consists of an oxygenase domain and a reductase
19 domain, each with its own catalytic activity (15) (Fig.2).

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1 Production of nitric oxide. Nitric oxide synthases (NOS) catalyze an oxidation of one N° -atom of the guanidine group of L-arginine to form NO (nitric oxide) and L-citrulline, with the intermediate N° -hydroxy-L-arginine (NOHA). ASS, argininosuccinate synthase; ASL, argininosuccinate lyase; OTC, ornithine carbamoyltransferase.

2.2. Nitric oxide synthases (NOSs) in relation to pain

Knowledge about the role of NO in different types of pain has been derived primarily from animal experiments examining the expression of the NOS isoforms in relation to a nociceptive stimulus. The idea that nNOS contributes to the role of NO in nociception is supported by the observation that its expression is rapidly upregulated in the dorsal horn neurons of the spinal cord after a peripheral noxious stimulation (18, 19). Genetic knockout and pharmacologic inhibition of neuronal nNOS attenuate nerve injury-induced mechanical hypersensitivity in mice, suggesting that nNOS may participate in the development and maintenance of mechanical hypersensitivity after nerve injury (20).

In mice lacking iNOS, there were fewer and smaller regenerating myelinated fibres and slowed reinnervation of muscle endplates distal to the injury zone, demonstrating that iNOS is a critical factor in the repair of injured tissue (21). Inducible NOS is upregulated in the inflamed tissue (22) and is involved in the development of hypersensitivity to pain in inflammatory and neuropathic pain models (23). These findings lead to the conclusion that

administration of iNOS inhibitors may have potential in the treatment of inflammatory and neuropathic pain syndromes (24).

NO generated by eNOS may modulate leukocyte adherence, a key factor in acute tissue inflammation (25). It participates in the regulation of vascular tone, vascular remodelling, angiogenesis (26, 27) and neurogenesis (28).

2.3. Potential role of tetrahydrobiopterin (BH4) in pain

Various cofactors, including tetrahydrobiopterin (BH4), flavin mononucleotide (FMN), flavin adenine dinucleotide (FAD), reduced nicotinamide adenine dinucleotide phosphate (NADPH), and calmodulin/Ca²⁺, are necessary for the formation of NO if the synthase isoform is the constitutive NOS I or NOS III (29, 30) (Fig.2).




Figure 2 NOS cofactors. NOS has a reductase domain and an oxygenase domain. Electrons (e^-) are donated by NADPH (nikotinamid-adenin-dinukleotid) to the reductase domain of the enzyme and proceed via FAD (flavin-adenin-dinukleotid) and FMN (flavin mononucleotide) carriers to the oxygenase domain. There they interact with the heme iron and BH₄ at the active site to catalyze the reaction of oxygen with L-arginine, generating citrulline and NO. Electron flow through the reductase domain requires the presence of bound Ca²⁺/CaM (modified after Alderton) (30).

1
2
3 The presence of BH₄ is important in dopamine formation, and in several neurotransmitters
4 and NO formation, and any deficiency leads to the formation of other reactive oxygen species
5 (ROS) (15). The role of BH₄ as an intrinsic regulator in neuropathic and inflammatory pain
6 (ROS) (15). The role of BH₄ as an intrinsic regulator in neuropathic and inflammatory pain
7 was confirmed by increasing BH₄ synthesis in the rat sensory neurons in response to both
8 axonal injury and peripheral inflammation, and also by the observation that blocking the BH₄
9 synthesis reduced neuropathic and inflammatory pain (31). The enzyme implicated in BH₄
10 synthesis reduced neuropathic and inflammatory pain (31). The enzyme implicated in BH₄
11 synthesis, GTP cyclohydrolase (GCH1), is a key modulator of peripheral neuropathic and
12 inflammatory pain (32) (Fig.3). In animal models of both neuropathic pain and inflammatory
13 pain, injecting GTP cyclohydrolase inhibitor, called 2, 4-diamino-6-hydroxypyrimidine
14 (DAHP), alleviated hypersensitivity to pain and reduced the levels of NO (31, 32).
15 Furthermore, the animal models were correlated to clinical pain when a “pain-protective”
16 haplotype of the GCH1 gene was identified (31). This haplotype that reduces BH₄
17 upregulation was protective against persistent neuropathic pain, and it decreased pain
18 sensitivity in low back pain patients following herniated disc surgery and also in normal
19 volunteers in an experimental pain model (31).
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3 Implications of BH₄ and GTP cyclohydrolase in pain (modified after Pasternak)

(33). Noxious stimuli induce the upregulation of GTP cyclohydrolase (GCH1). BH₄ modulates several

1
2
3 *enzymes, including neuronal nitric oxide synthase (nNOS), which generates nitric oxide (NO) that is under*
4 *the control of NMDA (N-methyl-D-aspartate) receptors through their regulation of intracellular calcium.*

7 In the absence of substrate and cofactors, NOS no longer produces NO, but transfers the free
8 electrons and produces free oxygen radicals (34).

11 **3. Reactive oxidative species (oxygen/nitrogen) and pain**

14 Reactive oxidative species are highly reactive chemicals that contain only oxygen atoms
15 (reactive oxygen species-ROS) or both nitrogen and oxygen (reactive nitrogen species-RNS).
16 ROS react easily with other molecules, resulting in potentially damaging modifications (35).
17 The main members of reactive oxygen species are superoxide (O_2^-), H_2O_2 and the hydroxyl
18 radical (OH^-). RNS are a group of chemically reactive molecules derived from nitric oxide
19 (NO) that may react with other radicals to give species of greater reactivity and toxicity such
20 as peroxynitrite ($ONOO^-$), nitrogen dioxide (NO_2) and dinitrogen trioxide (N_2O_3) (36). These
21 reactions account for the **indirect effects** of NO (35) observed under pathophysiologic
22 circumstances when NO flux becomes enhanced, mainly as a consequence of iNOS
23 expression. The interaction between nitric oxide and superoxide is responsible for the
24 formation of peroxynitrite (37), which is a potent cytotoxic and pro-inflammatory molecule
25 (38). Both superoxide and peroxynitrite are mediators of pain that accompanies inflammation
26 (39, 40, 41). Furthermore, ROS contribute to hyperalgesia (clinically defined as an augmented
27 sensitivity to painful stimuli) due to nerve injury (42, 43, 44) and to delayed recovery of
28 injured nerves (45). Thus, peroxynitrite formed after partial nerve injury contributes to the
29 initiation of hyperalgesia and Wallerian degeneration (46), damage that is alleviated by ROS
30 scavengers (42). In spite of many published articles, investigation of the involvement of
31 ROS/RNS in pain is limited to animal studies describing the effects of antioxidants in
32 attenuation of neuropathic or inflammatory pain (47, 48). “Spin-trap reagents” are the most
33 potent ROS scavengers, with phenyl-N-tert-butyl nitron (PBN) being especially noteworthy
34 (47). PBN has been shown to possess a neuroprotective action in several animal models of

1
2
3 brain ischaemia (49) not only because of its capacity to scavenge free radicals, but also due to
4
5 the inhibition of inducible nitric oxide synthase (iNOS) expression, as well as to the inhibition
6
7 of cytokine production (50). In animal models PBN relieves mechanical allodynia and inhibits
8
9 development of neuropathic pain-like behavior (51). The site of action of PBN is considered
10
11 to be spinal rather than peripheral (42). The results suggest that ROS scavengers can be
12
13 developed as powerful analgesic drugs for neuropathic pain (42). *N-t*-butyl hydroxylamine
14
15 (NtBHA) is one of the breakdown products of PBN and is the only spin-trap reagent that has
16
17 been used in humans as a neuroprotective agent in acute ischaemic stroke (52). However, it is
18
19 still not clear whether antioxidant therapy is effective in human pain, but results obtained
20
21 from small clinical trials indicate that treatment with nutritional antioxidants (vitamin C,
22
23 vitamin E, and beta-caroten) may reduce pain in patients suffering from chronic pancreatitis
24
25 (53, 54) and fibromyalgia (55, 56).

31 **4. The NO-cyclic guanosine monophosphate (cGMP) pathway**

32 **4.1. Basic physiology**

33
34 The stimulation of cyclic guanosine monophosphate cGMP-dependent kinases is one of the
35
36 numerous **direct biological actions** of NO (57). NO activates the guanylyl cyclase and
37
38 increases the synthesis of cyclic guanosine monophosphate (58). The major target of NO is
39
40 NO-sensitive guanylyl cyclase (GC) or soluble guanylyl cyclase (sGC). Activation of
41
42 guanylyl cyclase results in conversion of guanosine triphosphate to the second messenger
43
44 cGMP (59). The NO/cGMP signalling cascade (Fig.4) is important in the cardiovascular and
45
46 nervous systems, where it controls smooth muscle relaxation and modulation of synaptic
47
48 transmission (60).
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4 NO/cGMP signaling cascade. *NO endogenously produced by NO synthases activates NO-sensitive GC and leads to increased synthesis of cGMP. This intracellular messenger in turn modulates the activity of cGMP-dependent kinases, cGMP-gated ion channels, and cGMP-regulated phosphodiesterases.*

The second messenger cGMP has several targets, including cGMP-dependent protein kinase I (PKG-I) and PKG-II, activation of cGMP-regulated phosphodiesterase and cGMP-regulated ion channels (60). PKG-I is expressed in the spinal cord and is involved in the facilitation of synaptic transmission of nociceptive stimuli in the spinal cord (61, 62) and also in substance P synthesis (32).

4.2. The NO-cGMP pathway in pain

The NO-cGMP signalling pathway is present in neurons of the spinal cord and contributes to the development of hyperalgesia in animal models of pain (63, 64, 65) through PKG-I activation (66). It was recently demonstrated that during spinal nociceptive processing, cGMP produced by NO-GC may activate signalling pathways different from those activated by cGMP-dependent protein kinase I (cGKI) (67). Different studies have demonstrated the implications of the NO-cGMP pathway in the analgesic effect of several drugs that are indicated for the treatment of neuropathic pain, such as tramadol (68), spinal-administered clonidine (69), gabapentin (70, 71), and also in the antinociceptive effect of anti-inflammatory

1
2
3 drugs such as indomethacin (72), and in the hyperalgesia evoked by intrathecal high dose
4 morphine (73,74). The NO-cGMP pathway interacts with other signalling pathways (75) such
5
6 as the cholinergic, adrenergic, purinergic and peptidergic pathways in the peripheral nerve
7
8 system, and also with endocannabinoids (76). In pain research, the significance of these
9
10 interactions of the NO-cGMP pathway with other signalling pathways is only speculative,
11
12 because they are still poorly understood.
13
14
15

16 17 **5. The link between NO and NMDA**

18
19 The *N*-methyl-D-aspartate (NMDA) receptor is activated via a voltage-gated ion channel that,
20
21 once activated, allows Ca^{2+} to enter the neuron. It is the increase of intracellular Ca^{2+} that
22
23 triggers the cascade of events that includes activation of a constitutive form of NOS, followed
24
25 by increased production of NO (77). Many of the effects of NMDA activation are mediated
26
27 via production of NO. The NO diffuses to adjacent neurons and glia, where it activates
28
29 soluble guanylate cyclase (sGC), which in turn increases the intracellular content of cyclic
30
31 guanylate mono phosphate (cGMP) (78). After tissue injury, sensitization of nociceptors and
32
33 changes in the excitability of spinal neurons occur, which could underlie development of
34
35 primary hyperalgesia and secondary hyperalgesia. The spinal NMDA receptor is associated
36
37 with secondary hyperalgesia (79).
38
39
40
41
42

43 **6. NO and peripheral nociception**

44
45 The role of nitric oxide and cyclic GMP in the periphery has been controversial. Several
46
47 studies have demonstrated that NO may promote both pro- and antinociceptive effects (80).
48
49 Intracutaneous and paravascular injections of NO in humans can directly evoke pain in a
50
51 dose-related manner (81, 82). NO-donors led to hypersensitivity in rats (83) and to
52
53 antinociception in humans (84, 85). Thus, transdermal NO-donor glyceryl trinitrate was found
54
55 to act as an antinociceptive when used to treat shoulder pain syndrome due to supraspinatus
56
57 tendinitis (84) and elbow tendinosis (85). Topical application of nitric oxide using
58
59
60

1
2
3 transdermal glyceryl trinitrate patches can increase blood supply to the region due to local
4
5 vasodilatation, which in turn increases the clearance of local inflammatory mediators or
6
7 bioactive proteins such as substance P (85). It may also stimulate wound fibroblasts and
8
9 increase fibroblast synthesis, as seen in the healing of tendons (84, 85). Furthermore, the
10
11 arginine/NO/cGMP pathway is antinociceptive in subcutaneous tissues, while it is
12
13 pronociceptive in intradermal tissues (86). In inflammatory pain, the action of NO depends on
14
15 the stages of inflammation (80); it is protective during the initial hours after inflammation
16
17 (80) and cytotoxic later (87). These studies suggest that NO and cGMP signalling are able to
18
19 induce pro- or antinociception depending of the different concentrations of NO produced
20
21 locally (87), and with a dose-response relationship (88), depending on the location (89) and
22
23 also on the stage of pathological processes (80).
24
25
26
27
28

29 **7. Effects of NO in central nervous system**

30
31 A peripheral nerve lesion gives rise to structural changes in the spinal cord, probably related
32
33 to the fact that NO acts as a modulator of dorsal horn spinal cord nociceptive pathways (90).
34
35 NO is suggested to act as a “retrograde transmitter” because it can easily pass through
36
37 neuronal membranes. Activation of a receptor results in the production of NO in a post-
38
39 synaptic neuron from which it rapidly diffuses to enter the presynaptic neuron, where it
40
41 modulates excitability and enhances synaptic connection (91). Our laboratory is engaged to
42
43 examine changes in NOS expressions using immunohistochemistry and to relate these
44
45 changes with morphological alterations in the spinal cord using light microscopy (92). Thus,
46
47 in our model of chronic neuropathic pain in rats, induced by lesion or ligation of peripheral
48
49 spinal nerves that induced hyperalgesia behaviour during 4 to 8 weeks after nerve lesion, we
50
51 reported the role of NO in nociceptive signalling based on the upregulation of neuronal NOS
52
53 in the superficial dorsal horn and intermediolateral cell column (92). This finding lead to the
54
55 notion that NO regulates sensory transduction at the spinal cord level. Furthermore, the
56
57
58
59
60

1
2
3 mechanisms underlying pain hypersensitivity have been associated with release of NO within
4 the spinal cord (93). The involvement of NO in nociception is supported by experiments in
5 which inhibitors of NOS (L-NAME) were used to reduce NO production, which in turn
6 reduced NOS upregulation and structural changes in the spinal cord (94). NO donors
7 administered intrathecally have been shown to elicit a nociceptive behavioural response (95),
8 whereas NOS inhibitors provide analgesia (96). Our group demonstrated that a peripheral
9 nerve lesion in a rat model of chronic neuropathic pain induces selective alterations in the
10 blood-spinal cord barrier (BSCB) permeability in a time-related manner and activates
11 microglia (97). We assessed the integrity of the BSCB using immunohistochemistry for
12 endogenous albumin that represents the breakdown of the BSCB following nerve lesion and
13 also the state of glial activation was examined using antibodies to glial fibrillary acidic protein
14 (GFAP), a specific marker of astrocytes (98). We hypothesise that neurochemical alteration
15 would lead to morphological changes in the spinal cord endothelial cells and the astrocytes
16 and contributes to the pathobiology of the neuropathic pain (90, 92, 99). There are reasons to
17 believe that nerve lesion is associated with upregulation of nitric oxide synthase (NOS) is
18 instrumental in morphological alterations in the spinal cord in animal models (90, 92, 100).
19 The brain expresses all three identified nitric oxide synthase isoforms, and NOS activity is
20 higher than in any other tissue (101). In the brain, NO has been proposed to be involved in
21 synaptic plasticity, or to act as a neurotoxin when produced in excess (102).

22 **8. Cyclo-oxygenase (COX) and NO**

23 Cyclo-oxygenase (COX) converts arachidonic acid into prostaglandins, prostacyclin and
24 thromboxane A₂. The link between COX and NO pathways was observed in 1993 by
25 Salvemini and co-workers when they demonstrated that the enhanced release of
26 prostaglandines, which follows inflammatory mechanisms, was nearly entirely driven by NO
27 (103). NO is implicated in the regulation of cyclo-oxygenase (104) activity and activates
28 cyclo-oxygenase, followed by an increase in prostaglandin synthesis (105). Nitric oxide
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 mediates several of its beneficial effects, such as maintenance of blood vessel tone, inhibition
4 of platelet aggregation and cytoprotection, through the activation of cyclo-oxygenase (106).
5
6 The discovery of the reciprocal interaction between NO/COX has opened up the possibility of
7 designing new drugs, with a better toxicological and safety profile, containing NO donors
8 conjugated with COX inhibitors.
9
10
11
12

13 14 15 **9. Potential applications of nitric oxide in pain treatment**

16
17 Pharmacological modulation of NO levels and NO biosynthesis may be a therapeutic strategy
18 for treating different conditions, including pain (107, 108).
19
20

21 22 **9.1. NO-releasing drugs**

23 24 **9.1.1. Cyclo-oxygenase-inhibiting nitric oxide donating drugs (CINODs)**

25
26 Cyclo-oxygenase-inhibiting nitric oxide donating drugs (CINODs) are also known under the
27 name nitric-oxide releasing non-steroidal anti-inflammatory drugs (NO-NSAID) because they
28 combine the pharmacological profile of NSAID drugs with a controllable manner of NO
29 release (107). The release of NO together with NSAID prevents the major side effects of
30 NSAIDs that appear after inhibition of cyclo-oxygenase and suppression of prostaglandin
31 synthesis (109). CINODs are synthesized by the ester linkage of an NO-releasing moiety to
32 conventional non-steroidal anti-inflammatory drugs such as aspirin (NO-aspirin), flurbiprofen
33 (NO-flurbiprofen), naproxen (NO-naproxen), diclofenac (NO-diclofenac) and ibuprofen (NO-
34 ibuprofen) (110). NO-naproxen is the first drug in the new class of analgesic and anti-
35 inflammatory CINOD compounds (111,112) that have been evaluated clinically in treating the
36 signs and symptoms of osteoarthritis of the knee (113,114).
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 52 *a. CINODs and the gastrointestinal tract*

53
54 Gastrointestinal disturbances are a side effect of NSAIDs, and have been attributed to
55 inhibition of gastric COX-1 activity leading to loss of cytoprotective prostaglandin (PGI₂ and
56 PGE₂) formation. CINODs exploit the functional role of NO in gastric protection, leading to
57 generation of anti-inflammatory drugs that are associated with less gastric toxicity than the
58
59
60

1
2
3 parent NSAID (Fig. 5), and with fewer erosions and a lower incidence of gastric and duodenal
4
5 ulcers (110, 112). The mechanism of gastrointestinal protection is related to NO-mediated
6
7 mucosal vasodilatation, which increases gastric mucosal blood flow and is also related to the
8
9 inhibition of leukocyte adhesion in the gastric microcirculation (112).
10
11

12 *b. CINODs exhibit enhanced anti-inflammatory activity (Fig.5)*

13 NSAIDs exhibit anti-inflammatory action, but the mechanisms of enhanced anti-inflammatory
14
15 activity of CINODs are probably related to the inhibition of caspase activity, thereby reducing
16
17 cytokine formation (115). In contrast to conventional NSAIDs, as well as selective COX-2
18
19 inhibitors, CINODs inhibit the generation of T helper 1 (Th1)-type cytokines *in vitro* and
20
21 interrupt Th1-type responses in animal models of inflammation (116). Caspases are a group of
22
23 proteases implicated in apoptosis, but also in cytokine maturation and cell growth and
24
25 differentiation (117). Among the caspases, those that most resemble caspase-1 are involved in
26
27 mediating cytokine release and play a major role in inflammation, and those resembling
28
29 members of the caspase-3 family are involved in apoptosis (117). Low levels of NO inactivate
30
31 caspase 1 and 3, and thus the enhanced anti-inflammatory effects of CINODs are NO-
32
33 dependent (110).
34
35
36
37
38
39

40 *c. CINODs in pain and hyperalgesia (Fig.5)*

41 It is well known that NSAIDs induce analgesia by inhibition of prostaglandin production.
42
43 Based on previous reports, it might be expected that NO donor drugs would promote pain
44
45 perception. On the contrary, however, NO donors acting at peripheral sites have been
46
47 demonstrated to reduce pain perception, although the rationale for the enhanced anti-
48
49 nociceptive activity of CINODSs is not clear (110).
50
51
52

53 *d. CINODs have increased anti-thrombotic potency (106, 110), which is related to the effects*
54
55 of NSAIDs, as well as to a greater degree of inhibition of platelet function and vasodilator
56
57
58
59
60

activity by NO that counteracts vasoconstriction caused by platelet-derived mediators such as TXA₂.




Figure 5. Cyclo-oxygenase inhibiting NO donating drugs (CINODs). *Column 1: the effects of CINODs. Column 2: interaction between nitric oxide (NO) and COX (cyclo-oxygenase) pathway. NSAID (non-steroidal anti-inflammatory drugs) inhibit COX1 and COX2 enzymes, followed by prostaglandin (PG) inhibition. Column 3: NO and NSAID effects.*

9.1.2. NO-donating drugs (NODD)

NO-donating drugs (NODD) represent second-generation compounds with an ester-linked NO moiety, unrelated to NSAIDs. They have reduced adverse effects and have an improved profile of pharmacological activity in terms of enhanced therapeutic efficacy (106). Several compounds have been developed that fit these criteria, examples of which include nitroparacetamol (NO-paracetamol), nitroprednisolone (NO-prednisolone) and nitromesalamine (NO-mesalamine) (106). These drugs have not been developed for clinical use, and further research is necessary to investigate these compounds in humans.

NO-prednisolone has been shown to be more potent than prednisolone alone in a rat model of cholesten induced arthritis (118).

1
2
3 *NO-paracetamol* was found to be devoid of the hepatotoxic effects of co-administered
4 paracetamol (119) and it is safe in mice with pre-existing liver disease. It even protects
5 against the hepatotoxic effect of co-administered paracetamol (120). One of the first
6 compounds developed was NCX-701, or nitroparacetamol. It has been shown to be effective
7 in acute nociception as well as in neuropathic pain, situations in which paracetamol and other
8 COX inhibitors have no effect. In addition, NCX-701 is more potent and, in some
9 circumstances, more effective than its parent compound in different models of inflammatory
10 pain (121). The augmented biological activity of this compound has been assumed to be due
11 to the NO release from nitroparacetamol (122). The NO donating derivate of gabapentin
12 alleviates neuropathic pain-like behaviour after spinal cord and peripheral nerve injury (123).

26 **9.1.3. NO-donors (nitroglycerine, isosorbid dinitrate)**

27
28
29 These are the classic nitric oxide donors which include *nitroglycerine (NTG)*, used for pain
30 relief in coronary artery disease. The physiological effect of nitroglycerine as a potent
31 vasodilator results after constitutive nitric oxide synthase (NOS) activation and synthesis of
32 NO, which is responsible for vasodilatation (124). It seems worthwhile to mention the effects
33 of transdermal nitroglycerine in analgesia, including enhancing the antinociception from
34 spinal administration of either sufentanil following orthopaedic surgery (125) or neostigmine
35 (126) in acute postoperative pain, and acting as coadjuvant in opiate therapy for the control of
36 cancer pain (127). Impaired neuronal NO generation in diabetic rats induces hyperalgesia
37 (128) and impaired endoneurial blood flow (129). Impaired NO generation may play a role in
38 diabetic neuropathic pain as it was found that *isosorbide dinitrate spray (ISDN)*, a NO donor
39 with local vasodilating properties, relieved pain in a small number of patients with diabetic
40 neuropathic pain (130). It was speculated that alleviation of pain with ISDN spray follows
41 NO-induced vasodilation, which improves microvascular blood flow and may induce
42 angiogenesis of the vasa nervorum (130).

9.2. NO- inhibitors and pain

Chronic pain patients show a significant increase in plasma levels of NO in comparison with healthy individuals (131). Consequently, development of nitric oxide synthase (NOS) inhibitors could constitute future therapy for patients with chronic pain syndromes.

In animals there are well documented studies where NOS inhibitors suppress experimental nociception resulting from intraplantar formalin injection (132), suppress carrageenan-induced mechanical and thermal hyperalgesia (133) and inhibit NMDA hyperalgesia (134).

Inhibition of NO could be achieved by targeting the differential cofactors (tetrahydrobiopterin) (31, 32), or the differential substrate requirements for expressing various isoforms of NOS (L-arginine uptake blockers or arginase) (135). An important measure in treating different types of pain would be to achieve selective inhibition of NOS isoforms, particularly iNOS and nNOS; in other words, to develop pharmacological inhibitors with isoform specificity (106).

Finally, a combination of NOS inhibitors and COX-inhibitors should be mentioned. A significant decrease of neuropathic-like behaviour was demonstrated in animal models that received co-administered cyclooxygenase-2 (COX-2) inhibitors (meloxicam and rofecoxib) and an inducible nitric oxide synthase (iNOS) inhibitor (aminoguanidine hydrochloride) as compared with the groups treated with an iNOS inhibitor or COX-2 inhibitors alone (136). These results are controversial, because the effects of COX-2 inhibitors are not considered efficacious in neuropathic pain. It was previously reported that iNOS and COX-2 pathways interact closely and that iNOS stimulates COX-2 activity (103), and thus it is speculated that this may be why the combination of NOS inhibitors and COX-2 inhibitors was effective in the treatment of neuropathic pain (136).

1
2
3 As described before with other combinations of drugs that modulate NO levels (CINODs), the
4 efficacy of the combination of iNOS and COX-2 inhibitors may be due to an added
5 advantage, such as potentiated action over and above that of each of the compounds alone.
6
7
8
9

10 **9.2.1. N^G-monomethyl-L-arginine hydrochlorides (L-NMMA) and methylene blue**
11 **(MB)** are the only inhibitors of NOS that have been thoroughly tested in humans. L-NMMA
12 inhibits all three types of NOS (endothelial NOS, neuronal NOS, and inducible NOS) (137).
13 L-NMMA is effective in the treatment of chronic tension-type headache (138, 139) and has
14 also been tested in the treatment of neuropathic pain in humans (140). The mechanism
15 responsible for headache has been studied using the NO-donor (glycerol trinitrate) model for
16 experimental headache in humans (138). The L-NMMA inhibitor of NOS has an analgesic
17 effect in different types of headache such as migraine and cluster-type headache (139).
18
19
20
21
22
23
24
25
26
27
28

29 **9.2.2. Inhibitors of Soluble Guanylyl/Guanylate Cyclase**

30 **Methylene blue** has direct inhibitory effects on nitric oxide synthases, both constitutive and
31 inducible (141, 142), and blocks accumulation of cyclic guanosine monophosphate (cGMP)
32 by inhibiting the enzyme guanylate cyclase (141). Data suggest that MB is a more specific
33 and potent inhibitor of NOS than guanylyl cyclase, because direct NO-donating compounds in
34 the presence of MB can still partially activate c-GMP signalling pathways (141, 142). It was
35 recently demonstrated that MB reduces chronic discogenic low back pain (143) as well as the
36 pain after lateral sphincterotomy (144). The underlying mechanisms of MB treatment in the
37 above clinical pain conditions are not completely understood. However, MB is known to
38 inhibit the formation of free oxygen radicals and superoxides (reactive oxygen species) by
39 competing with molecular oxygen for the transfer of electrons by xanthine oxidase (145). MB
40 acts as a neuroprotective agent after an experimental global ischemia-reperfusion incident in
41 experimental animal models (146). Furthermore, a number of observational studies and case
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

reports have evaluated the use of MB in both the prevention and treatment of ifosfamide-induced neurotoxicity (147).




Figure 6. NO-cGMP pathway and inhibitors. The scheme shows the blockers used and their target enzymes. Nitric oxide synthase catalyses the formation of NO from L-arginine, NO activates the soluble guanylyl cyclase which in turn catalyses the formation of cyclic GMP from GTP. At the end of the cascade, the cGMP specific phosphodiesterase type 5 hydrolyses cGMP to 5'-cGMP. Inhibitors: L-NAME (NG-nitro-L-arginine methyl ester, a blocker of the nitric oxide synthase); ODQ, 1H-[1, 2, 4]oxadiazolo[4,3-a]quinoxalin-1-one, a selective blocker of NO-sensitive guanylyl cyclase; Sildenafil, a selective blocker of the cGMP specific phosphodiesterase type 5 (leads to an increase in cGMP level). NO (nitric oxide); GMP, guanosine monophosphate; cGMP, cyclic guanosine monophosphate; GTP, guanosine triphosphate.

In conclusion, we believe that the inhibitors of the NO-cGMP pathway may be a tool for treatment of pain, and we hope that a specific therapeutic strategy will be developed in the near future.

10. Concluding remarks

This review has attempted to update the accumulated evidence concerning the mechanisms of NO action and the importance of NO in pain. The subject is controversial, in part because nitric oxide has an ambivalent character: it can have both harmful and beneficial effects in pain, with mechanisms that appear to contradict one another. Thus, the interpretation of data

1
2
3 requires caution and the beneficial and deleterious consequences of NO must be determined
4
5 in order to attain effective therapeutic interventions in the treatment of pain. Also, the NO
6
7 pathway interacts with other transmitter pathways as it is the reciprocal interaction between
8
9 NO/COX pathways. If COX-inhibiting NO donating drugs and donating NO drugs are
10
11 incorporated in the treatment of pain in humans, they will probably replace established drugs
12
13 currently used in treating pain. An effective NO inhibitor is likely to provide a new tool for
14
15 treating the pain associated with central sensitization. Unfortunately, only a few of these
16
17 compounds have reached the stage of clinical pain trials.
18
19
20
21
22
23
24

25 **II. References:**

- 26
27
28
29 **1** Snyder SH: Nitric oxide: first in a new class of neurotransmitters. *Science* 1992;
30
31 24;257:494-6
32
33
- 34
35 **2** Feelisch M, Martin FJ: The early role of nitric oxide in evolution. *TREE* 1995; 10:
36
37 496-499
38
- 39
40 **3** Bishop CD, Brandhorst BP: On nitric oxide signaling, metamorphosis, and the
41
42 evolution of biphasic life cycles. *Evolution & Development* 2003; 5: 542–550
43
- 44
45 **4** Klepper LA: Nitric oxide (NO) and nitrogen dioxide (NO₂) emissions from
46
47 herbicide-treated soybean plants. *Atmos Environ* 1979; 13: 537-542
48
- 49
50 **5** Encyclopedia Britannica online: Nitric oxide retrieved from URL:
51
52 <http://www.britannica.com/ebc/article-9373539/nitric-oxide>
53
- 54
55 **6** Priestley J: Experiments and Observations on Different Kinds of Air, Vol. II, and
56
57 section III, 1775, London, retrieved from
58
59 URL:<http://web.lemoyne.edu/~GIUNTA/EA/PRIESTLEYann.HTML>
60

- 1
2
3 7 Wright P: Nitric oxide: From menace to marvel of the decade. A briefing
4 document prepared for the Royal Society and Association of British Science
5 Writers 1996; retrieved from URL:
6
7
8
9
10 <http://www.absw.org.uk/briefings/nitric%20oxide.htm>
11
12 8 Lewis R, Palevitz B: Nobel honors pioneers of NO. *The Scientist* 1998; 12:1
13
14 9 Culotta E, Koshland DE: NO news is good news. *Science* 1992; 258:1862-5
15
16
17 10 The Nobel Assembly at Karolinska Insitutet Press Release/Nobelprize.org press
18 release The Nobel Prize in Physiology or Medicine 1998 retrieved from
19
20 URL:http://nobelprize.org/nobel_prizes/medicine/laureates/1998/press.html
21
22
23 11 Palmer RMJ, Ashton DS, Moncada S: Vascular endothelial cells synthesise nitric
24 oxide from L-arginine. *Nature* 1988; 333: 664-6
25
26
27 12 Furchgott R, Zawadzki J: The obligatory role of endothelial cells in the relaxation
28 of arterial smooth muscle by acetylcholine. *Nature* 1980; 288: 373-6
29
30
31 13 Bredt DS, Hwang PM, Snyder SH: Localization of nitric oxide synthase indicating
32 a neuronal role for nitric oxide. *Nature* 1990; 347:768-770
33
34
35 14 Knowles RG, Moncada S : Nitric oxide synthases in mammals. *Biochem J* 1994;
36 298:249-58
37
38
39 15 Bruckdorfer R: The basics about nitric oxide. *Mol Asp Med* 2005; 26: 3-31
40
41
42 16 Schild L, Reinheckel T, Reiser M, et al: Nitric oxide produced in rat liver
43 mitochondria causes oxidative stress and impairment of respiration after transient
44 hypoxia. *FASEB J* 2002; 17: 2194-2201
45
46
47 17 MacMicking J, Xie QW, Nathan C: Nitric oxide and macrophage function. *Annu*
48 *Rev Immunol* 1997; 15:323-350
49
50
51 18 Meller ST, Gebhart GF: Nitric oxide (NO) and nociceptive processing in the spinal
52 cord. *Pain* 1993; 52: 127-136
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 19 Lam HH, Hanley DF, Trapp BD, et al: Induction of spinal cord neuronal nitric oxide synthase (NOS) after formalin injection in the rat hind paw. *Neurosci Lett* 1996; 210: 201-204
- 20 Guan Y, Yaster M, Raja SN, et al: Genetic knockout and pharmacologic inhibition of neuronal nitric oxide synthase attenuate nerve injury-induced mechanical hypersensitivity in mice. *Mol Pain* 2007;3:29
- 21 Levy D, Kubes P, Zochodne DW: Delayed peripheral nerve degeneration, regeneration, and pain in mice lacking inducible nitric oxide synthase. *J Neuropathol Exp Neur* 2001; 60: 411-421
- 22 Gühring H, Tegeder I, Lötsch J, et al: Role of nitric oxide in zymosan induced paw inflammation and thermal hyperalgesia. *Inflamm Res* 2001; 50: 083-088
- 23 De Alba J, Clayton NM, Collins SD, et al: GW274150, a novel and highly selective inhibitor of the inducible isoform of nitric oxide synthase (iNOS), shows analgesic effects in rat models of inflammatory and neuropathic pain. *Pain* 2006; 120: 170-81
- 24 Sung CS, Wen ZH, Chang WK, et al: Intrathecal interleukin-1 β administration induces thermal hyperalgesia by activating inducible nitric oxide synthase expression in the rat spinal cord. *Brain Res* 2004; 1015: 145-153
- 25 Kubes P, Suzuki M, Granger DN: Nitric oxide: an endogenous modulator of leukocyte adhesion. *Proc Natl Acad Sci USA* 1991; 88:4651-4655
- 26 Rudic RD, Shesely EG, Maeda N, et al: Direct evidence for the importance of endothelium-derived nitric oxide in vascular remodeling. *J Clin Invest* 1998; 101:731-736
- 27 Murohara T, Asahara T, Silver M, et al: Nitric oxide synthase modulates angiogenesis in response to tissue ischemia. *J Clin Invest* 1998; 101:2567-2578

- 1
2
3 28 Chen J, Zacharek A, Zhang C, et al: Endothelial nitric oxide synthase regulates
4 brain-derived neurotrophic factor expression and neurogenesis after stroke in mice.
5
6 *J Neurosci* 2005; 25: 2366–2375
7
8
9
- 10 29 Moncada D, Palmer RMJ, Higgs EA: Nitric oxide: physiology, pathophysiology
11 and pharmacology. *Pharmacol Rev* 1991; 43: 109-142
12
13
14
- 15 30 Alderton WK, Cooper CE, Knowles RG: Nitric oxide synthase : structure, function
16 and inhibition. *Biochem J* 2001; 357: 593-615
17
18
19
- 20 31 Tegeder I, Costigan M, Griffin RS, et al: GTP cyclohydrolase and
21 tetrahydrobiopterin regulate pain sensitivity and persistence. *Nat Med* 2006; 12 :
22 1269-77
23
24
25
26
- 27 32 Tegeder I, Del Turco D, Schmidtko A, et al: Reduced inflammatory hyperalgesia
28 with preservation of acute thermal nociception in mice lacking cGMP-dependent
29 protein kinase I. *Proc Natl Acad Sci USA* 2004; 101:3253-7
30
31
32
33
- 34 33 Pasternak GW, Inturrisi CE: Feeling pain? Who's your daddy...*Nature Medicine*
35 2006; 12: 1243 - 1244
36
37
38
- 39 34 Schulz R, Kelm M, Heusch G: Nitric oxide in myocardial ischemia/reperfusion
40 injury. *Cardiovasc Research* 2004; 61: 402-413
41
42
43
- 44 35 Duncan AJ, Heales SJR: Nitric oxide and neurological disorders. *Mol Asp Med*
45 2005; 26; 67-96
46
47
48
- 49 36 Hughes M: Chemistry of Nitric Oxide and Related Species. *Methods in*
50 *Enzymology* 2008; 436:3-19
51
52
53
- 54 37 Beckman JS, Beckman TW, Chen J: Apparent hydroxyl radical production by
55 peroxynitrite: implications for endothelial injury from nitric oxide and superoxide.
56
57 *Proc Natl Acad Sci U S A* 1990; 87: 1620-1624
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 38 Salvemini D, Doyle TM, Cuzzocrea S: Superoxide, peroxynitrite and oxidative/nitrative stress in inflammation. *Biochem Soc Transactions* 2004; 34: 965-970
- 39 Wang ZQ, Porreca F, Cuzzocrea S, et al: A newly identified role for superoxide in inflammatory pain. *J Pharmacol Exp Ther* 2004; 309: 869–878
- 40 Liu T, Knight KR, Tracey DJ: Hyperalgesia due to nerve injury—role of peroxynitrite. *Neuroscience* 2000; 97:125-131
- 41 Ndengele M, Cuzzocrea S, Esposito E, et al: Cyclooxygenases 1 and 2 contribute to peroxynitrite-mediated inflammatory pain hypersensitivity. *The FASEB Journal* 2008;22: 3154-3164
- 42 Chung JM: The role of reactive oxygen species (ROS) in persistent pain. *Molecular interventions* 2004;4: 248-249
- 43 Khalil Z, Liu T, Helme RD: Free radicals contribute to the reduction in peripheral vascular responses and the maintenance of thermal hyperalgesia in rats with chronic constriction injury. *Pain* 1999; 79 :31-7
- 44 Wagner R, Heckman HM, Meyers RR: Wallerian degeneration and hyperalgesia after peripheral nerve injury are glutathione-dependent. *Pain* 1998; 77: 173-9
- 45 Khalil Z, Khodr B: A role for free radicals and nitric oxide in delayed recovery in aged rats with chronic constriction nerve injury. *Free Radic Biol Med* 2001; 31:430-9.
- 46 Ridger VC, Greenacre SA, Handy RL, et al: Effect of peroxynitrite on plasma extravasation, microvascular blood flow and nociception in the rat. *Br J Pharmacol* 1997;122:1083-8
- 47 Kim HK, Park SK, Zhou JL, et al: Reactive oxygen species (ROS) play an important role in a rat model of neuropathic pain. *Pain* 2004; 111: 116-124

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 48 Nishiyama T, Ogawa M: Intrathecal edaravone, a free radical scavenger, is effective on inflammatory-induced pain in rats. *Acta Anaesthesiol Scandinavica* 2005; 49: 147-151
- 49 Wiklund L, Sharma HS, Basu S : Circulatory arrest as a model for studies of global ischemic injury and neuroprotection *Ann N Y Acad Sci* 2005;1053:205-19
- 50 Dhainaut A, Tizot A, Eric Raimbaud E, et al: Synthesis, Structure, and Neuroprotective Properties of Novel Imidazolyl Nitrones. *J Med Chem* 2000; 43: 2165 -2175
- 51 Floyd RA: Protective Action of Nitronone-Based Free Radical Traps against oxidative Damage to the Central Nervous Systems. *Advances in Pharmacology*, Academic Press Inc: New York, 1997; 38: 361-378
- 52 Lees KR, Zivin JA, Ashwood T, et al: NXY-059 for acute ischemic stroke. *N Engl J Med* 2006; 354: 588–600
- 53 Kirk GR, White JS, Lloyd McKie L, et al: Combined antioxidant therapy reduces pain and improves quality of life in chronic pancreatitis. *J Gastroint Surg* 2006; 10: 499-503
- 54 Bhardwaj P, Grag PK, Saray A, et al: Antioxidant Supplementation for Pain Relief in Chronic Pancreatitis: A Randomized Placebo Controlled Double Blind Trial. *Gastroenterology* 2007; 132: A51
- 55 Bagis S, Tamer L, Sahin G, et al: Free radicals and antioxidants in primary fibromyalgia: an oxidative stress disorder? *Rheumatol Int* 2005; 25: 188-90
- 56 Salih O, Huseyin O, Sadik S, et al: Current concepts in the pathophysiology of fibromyalgia: the potential role of oxidative stress and nitric oxide *Rheumatol Int*. 2006; 26: 585-597

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 57 Miki N, Kawabe Y and Kuriyama K: Activation of cerebral guanylate cyclase by nitric oxide. *Biochem Biophys Res Commun* 1977; 75:851-856
- 58 Byers MR, Bonica J : Peripheral pain mechanisms pp26-72 in *Bonica`s Management of Pain* third edition Editor JDLoeser, Lippincott Williams&Wilkins 2001
- 59 Friebe A, Koesling D: Regulation of nitric-oxide-sensitive guanylyl cyclase. *Circ Res* 2003; 93:96-105
- 60 Garthwaite J, Charles SL, Chess-Williams R: Endothelium-derived relaxing factor release on activation of NMDA receptors suggests role as intercellular messenger in the brain. *Nature* 1988; 336: 385–388
- 61 Qian Y, Chao D, Santillano DR, et al: cGMP-dependent protein kinase in dorsal root ganglion: relationship with nitric oxide synthase and nociceptive neurons. *J Neurosci* 1996; 16: 3130-3138
- 62 Schmidt H, Werner M, Heppenstall PA, et al: cGMP-mediated signaling via cGKIalpha is required for the guidance and connectivity of sensory axons. *J Cell Biol* 2002; 159: 489-498
- 63 Xu L, Mabuchi T, Katano T, et al: Nitric oxide (NO) serves as a retrograde messenger to activate neuronal NO synthase in the spinal cord via NMDA receptors. *Nitric Oxide* 2007; 17 (1):18-24).
- 64 Meller ST, Dykstra C, Gebhart GF: Production of endogenous nitric oxide and activation of soluble guanylate cyclase are required for *N*-methyl-d-aspartate-produced facilitation of the nociceptive tail-flick reflex. *Eur J Pharmacol* 1992;214: 93–96

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 65 YX Tao, RA Johns: Activation of cGMP-dependent protein kinase I α is required for *N*-methyl-d-aspartate- or nitric oxide-produced spinal thermal hyperalgesia. *Eur J Pharmacol* 2000; 392: 141–145
- 66 A Schmidtko, P Ruth, G Geisslinger, et al: Inhibition of cyclic guanosine 5'-monophosphate-dependent protein kinase I (PKG-I) in lumbar spinal cord reduces formalin-induced hyperalgesia and PKG upregulation *Nitric Oxide* 2003; 8: 89–94
- 67 Schmidtko A, Gao W, König P, et al: cGMP produced by NO-sensitive guanylyl cyclase essentially contributes to inflammatory and neuropathic pain by using targets different from cGMP-dependent protein kinase I. *J Neurosci* 2008;28:8568-76
- 68 Dal D, Salman MA, Salman AE, et al: The involvement of nitric oxide on the analgesic effect of tramadol. *Eur J Anaesthesiol* 2006; 23: 175-177
- 69 Ge YX, Xin WJ, Hu NW, et al: Clonidine depresses LTP of C-fiber evoked potentials in spinal dorsal horn via NO-cGMP pathway. *Brain Research* 2006; 1118: 58-65
- 70 Mixcoatl-Zecuatl T, Flores-Murrieta F, Grannados-Soto V: The nitric oxide-cyclic GMP-protein kinase G-K⁺ channel pathway participates in the antiallodynic effect of spinal gabapentin. *Eur J Pharmacol* 2006; 531: 87-95
- 71 Ortiz MI, Medina-Tato DA, Sarmiento-Heredia D, et al: Possible activation of the NO-cyclic GMP-protein kinase G-K⁺ channels pathway by gabapentin on the formalin test. *Pharmacol Biochem Behav* 2006; 83:420-7
- 72 Ventura-Martinez R, Dèciga-Campos M, Diaz-Reval MI, et al: Peripheral involvement of the nitric oxide–cGMP pathway in the indomethacin-induced antinociception in rat. *Eur J Pharmacol* 2004; 503: 43– 48

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 73 Komatsu T, Sakurada C, Sasaki M, et al: The role of nitric oxide in the nociceptive behaviour evoked by intrathecal high-dose morphine. *J Pharmacol Sci* 2004; 94: 229
- 74 Watanabe C, Sakurada T, Okuda K, et al: The role of spinal nitric oxide and glutamate in nociceptive behaviour evoked by high-dose intrathecal morphine in rats. *Pain* 2003; 106: 269-283
- 75 Garthwaite J: Concepts of neural nitric oxide-mediated transmission. *Eur J Neurosci* 2008; 27: 2783-2802
- 76 Sergeeva OA, Doreulee N, Chepkova AN, et al: Long-term depression of corticostriatal synaptic transmission by DHPG depends on the endocannabinoid release and nitric oxide synthesis. *Eur J Neurosci* 2008; 26: 1889-1894
- 77 Meller ST, Gebhart GF: Nitric oxide (NO) and nociceptive processing in the spinal cord. *Pain* 1993; 52: 127-136
- 78 Morris R, Southam E, Braid DJ, et al: Nitric oxide may act as a messenger between dorsal root ganglia neurones and their satellite cells. *Neurosci Lett* 1992; 137: 29-32
- 79 Urban MO, Gebhart GF: Supraspinal contributions to hyperalgesia. *Proc Natl Acad Sci USA* 1999; 96: 7687-7692
- 80 Paul-Clark MJ, Gilroy DW, Willis D, et al: Nitric oxide synthase inhibitors have opposite effects on acute inflammation depending on their route of administration. *J Immunol* 2001; 166: 1169-1177
- 81 Holtusen H, Arndt JO: Nitric oxide evokes pain in humans on intracutaneous injection. *Neurosci Lett* 1994; 165: 71-74
- 82 Holtusen H, Arndt JO: Nitric oxide evokes pain at nociceptors of the paravascular tissue and veins in humans. *J Physiol* 1995; 253-258

- 1
2
3 **83** Aley KO, McCarter G, Levine JD: Nitric oxide signalling in pain and nociceptor
4 sensitization in the rat. *J Neurosci* 1998;18:7008-7014
5
6
7
8 **84** Berrazueta JR, Losada A, Poveda J, et al: Successful treatment of shoulder pain
9 syndrome due to supraspinatus tendinitis with transdermal nitroglycerin. A double
10 blind study. *Pain* 1995; 66: 63–67
11
12
13
14
15 **85** Paoloni JA, Appleyard RC, Nelson J, et al: Topical nitric oxide application in the
16 treatment of chronic extensor tendinosis at the elbow. A randomized, double-
17 blinded, placebo-controlled clinical trial. *Am J Sports Med* 2003; 31:915-920
18
19
20
21
22 **86** Vivancos GG, Parada CA, Ferreira SH: Opposite nociceptive effects of the
23 arginine/NO/cGMP pathway stimulation in dermal and subcutaneous tissues. *Br J*
24 *Pharmacol* 2003; 138: 1351-1357
25
26
27
28
29 **87** Iadecola C: Bright and dark sides of nitric oxide in ischemic brain injury. *Trends*
30 *Neurosci* 1997; 20:132-139
31
32
33
34 **88** Kawabata A, Manabe S, Manabe Y, et al: Effect of Topical Administration of L-
35 arginine on Formalin-induced Nociception in the Mouse: A Dual Role of
36 Peripherally Formed Nitric Oxide in Pain Modulation. *Br J Pharmacol*
37 1994;112:547-550
38
39
40
41
42
43 **89** Bhave G, Gereau RW: Postranslational mechanisms of peripheral sensitization. *J*
44 *Neurobiol* 2004; 61: 88-106
45
46
47
48 **90** Gordh T: The role of nitric oxide in neuropathic pain and neurodegeneration. *Acta*
49 *Anaesthesiol Scand* 1998; 42: 29-30
50
51
52
53 **91** Barinaga M: Is nitric oxide the `retrograde messenger`? *Science* 1991; 254: 1296-
54 1297
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 92** Gordh T, Sharma HS, Alm P, et al: Spinal nerve lesion induces upregulation of neuronal nitric oxide synthase in the spinal cord. An immunohistochemical investigation in the rat. *Amino Acids* 1998; 14; 105-112
- 93** Meller ST, Gebhart GF: Nitric oxide (NO) and nociceptive processing in the spinal cord. *Pain* 1993; 52:127–136
- 94** Inoue T, Mashimo T, Shibata M, et al: Rapid development of nitric oxide-induced hyperalgesia depends on an alternate to the cGMP-mediated pathway in the rat neuropathic pain model. *Brain Res* 1998;792:263-270
- 95** Roche AK, Cook M, Wilcox GL, et al: A nitric oxide synthesis inhibitor (L-NAME) reduces licking behavior and Fos-labeling in the spinal cord of rats during formalin-induced inflammation. *Pain* 1996; 66: 331-341
- 96** Bredt DS, Glatt CE, Hwang PM, et al: Nitric oxide synthase protein and mRNA are discretely localized in neuronal populations of the mammalian CNS together with NADPH diaphorase. *Neuron* 1991; 7: 615-624
- 97** Gordh T, Chu H, Sharma HS: Spinal nerve lesion alters blood–spinal cord barrier function and activates astrocytes in the rat. *Pain* 2006;124:211–221
- 98** Sharma HS, Zimmer C, Westman J, et al: Acute systemic heat stress increases glial fibrillary acidic protein immunoreactivity in brain: experimental observations in conscious normotensive young rats. *Neuroscience* 1992;48:889–901
- 99** Karlsten R, Gordh T. How do drugs relieve neurogenic pain? *Drugs Aging* 1997;11:398–412
- 100** Sharma HS, Alm P: Role of nitric oxide on the blood–brain and the spinal cord barriers. In: Sharma HS, Westman J, editors. *The blood–spinal cord and brain barriers in health and disease*. San Diego: Elsevier Academic Press; 2004. p. 191–230.

- 1
2
3 **101**Garthwaite J, Boulton CL Nitric oxide signalling in the central nervous system.
4
5 *Annu Rev Physiol* 1995; 57:683-706
6
7
8 **102**Vincent SR: Nitric oxide: A radical neurotransmitter in the central nervous system.
9
10 *Prog Neurobiol* 1994; 42: 129-160
11
12 **103**Salvemini D, Misko TP, Masferrer JL, et al: Nitric oxide activates cyclooxygenase
13 enzyme. *Proc Natl Acad Sci USA* 1993; 90: 7240-7244
14
15
16
17 **104**Mollace V, Muscoli C, Masini E, et al: Modulation of prostaglandin biosynthesis
18 by nitric oxide and nitric oxide donors. *Pharmacol Rev* 2005; 57:217
19
20
21
22 **105**McCormack K: Non-steroidal anti-inflammatory drugs and spinal nociceptive
23 processing. *Pain* 1994; 59: 9-43
24
25
26
27 **106**Salvemini D, Boyle TM, Cuzzocrea S: Superoxide, peroxynitrite and
28 oxidative/nitrative stress in inflammation. *Biochem Soc Transactions* 2006;
29 34:965-970
30
31
32
33
34 **107**Low SY: Application of pharmaceuticals to nitric oxide. *Mol Aspects Med* 2005;
35 26: 97-138
36
37
38
39 **108**Alcaraz MJ, Guillén MI: Nitric oxide related therapeutic phenomenon: A
40 challenging task. *Curr Pharmac Design* 2002; 8: 215-231
41
42
43
44 **109**Cashman JN: The mechanism of action of NSAIDs in analgesia. *Drugs* 1996; 52:
45 13-23
46
47
48
49 **110**Burgaud JL, Riffaud JP, Del Soldato P: Nitric-oxide releasing molecules: A new
50 class of drugs with several major indications. *Curr Pharmac Design* 2002; 8: 201-
51 213
52
53
54
55 **111**Keeble JE, Moore PK: Pharmacology and potential therapeutic applications of
56 nitric-oxide- releasing non-steroidal anti-inflammatory and related nitric oxide-
57 donating drugs. *Br J Pharmacol* 2002; 137: 295-310
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 112**Fiorucci S, Santucci L, Distrutti E: NSAIDs, coxibs, CINOD and H (2) S-releasing NSAIDs: What lies beyond the horizon. *Dig Liver Dis* 2007; 39: 1043-51
- 113**Schnitzer TJ, Kivitz AJ, Lipetz RS, et al: Comparison of the COX-inhibiting nitric oxide donator AZD3582 and rofecoxib in treating the signs and symptoms of osteoarthritis of the knee. *Arthritis Rheum* 2005; 53: 827-37
- 114**Lohmander LS, McKeith D, Svensson O, et al: STAR Multinational Study Group. *Ann Rheum Dis* 2005;64:449
- 115**Fiorucci: NO-releasing NSAIDs are caspase inhibitors. *Trends Immunol* 2001; 22: 232-235
- 116**Fiorucci S, Santucci L, Antonelli E, et al: NO-aspirin protects from T-cell mediated liver injury by inhibiting caspase- dependent processing of Th1-like cytokines. *Gastroenterology* 2000; 118: 404–421
- 117**Salvesen GS, Dixit VM: Caspases: intracellular signalling by proteolysis. *Cell* 1997; 91: 443–446
- 118**Paul-Clark MJ, Mancini L, Del Soldato P et al: Potent antiarthritic properties of a glucocorticoid derivative, NCX-1015, in an experimental model of arthritis. *Proc Natl Acad Sci USA* 2002; 99: 1677-82.
- 119**Futter LE, Al-Swayeh OA, Moore PK: A comparison of the effect of nitroparacetamol and paracetamol in liver injury. *Br J Pharmacol* 2001; 132: 10-12
- 120**Fiorucci S, Antonelli E, Mencarelli A, et al: A NO-releasing derivative of acetaminophen spares the liver by acting at several checkpoints in the Fas pathway. *Br J Pharmacol* 2002; 135: 589-599

- 1
2
3 **121**Romero-Sandoval EA, Curros-Criado MM, Gaitan G, et al: Nitroparacetamol
4
5 (NCX-701) and pain: first in a series of novel analgesics. *CNS Drug Rev* 2007; 13:
6
7 279-95
8
9
- 10 **122**al-Swayeh OA, Futter LE, Clifford RH, et al: Nitroparacetamol exhibits anti-
11
12 inflammatory and anti-nociceptive activity *Br J Pharmacol* 2000; 130: 1453-1456
13
14
- 15 **123**Wu WP, Hao JX, Ongini E, et al: A nitric oxide (NO)-releasing derivative of
16
17 gabapentin, NCX 8001, alleviates neuropathic pain-like behaviour after spinal cord
18
19 and peripheral nerve injury. *Br J Pharmacol* 2004; 141: 65-74
20
21
- 22 **124**Bonini MG, Stadler K, Silva SO, et al: Constitutive nitric oxide synthase
23
24 activation is a significant route for nitroglycerin-mediated vasodilation. *Proc Natl*
25
26 *Acad Sci USA* 2008 ;105 : 8569-74
27
28
- 29 **125**Lauretti GR, de Oliveira R, Reis MP, et al: Transdermal nitroglycerine enhances
30
31 spinal sufentanil postoperative analgesia following orthopedic surgery.
32
33 *Anesthesiology* 1999; 90:734–9
34
35
- 36 **126**Lauretti GR, Oliveira AP, Julião MC, et al: Transdermal nitroglycerine enhances
37
38 spinal neostigmine postoperative analgesia following gynecological surgery.
39
40 *Anesthesiology* 2000; 93: 943–946
41
42
- 43 **127**Lauretti GR, Lima ICPR, Reis MP, et al: Oral ketamine and transdermal
44
45 nitroglycerin as analgesic adjuvants to oral morphine therapy for cancer pain
46
47 management. *Anesthesiology* 1999;90:1528–1533
48
49
- 50 **128**Sasaki T, Yasuda H, Maeda K, et al: Hyperalgesia and decreased neuronal nitric
51
52 oxide synthase in diabetic rats. *Neuroreport* 1998; 9: 243-247
53
54
- 55 **129**Pitei DL, Watkins PJ, Edmonds ME: NO-dependent smooth muscle vasodilatation
56
57 is reduced in NIDDM patients with peripheral sensory neuropathy. *Diab Med*
58
59 1997; 14: 284-290
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 130**Yuen KCJ, Baker NR, Ratman G: Treatment of chronic painful diabetic neuropathy with isosorbide dinitrate spray: A double-blind placebo-controlled cross-over study. *Diabetes Care* 2002; 25: 1699-1703
- 131**Koch A, Zacharowski K, Boehm O, et al: Nitric oxide and pro-inflammatory cytokines correlate with pain intensity in chronic pain patients. *Inflamm Res* 2007; 56: 32-37
- 132**Moore PK, Oluyomi AO, Babbedge RC, et al: L-NG-nitro arginine methyl ester exhibits antinociceptive activity in the mouse. *Br J Pharmacol* 1991; 102: 198–202
- 133**Handy RLC, Moore PK: Effects of selective inhibitors of neuronal nitric oxide synthase on carrageenan-induced mechanical and thermal hyperalgesia. *Neuropharmacology* 1998; 37: 37-43
- 134**Kitto KF, Haley JE, Wilcox GL: Involvement of nitric oxide in spinally mediated hyperalgesia in the mouse. *Neurosci-Lett.* 1992; 148: 1-5
- 135**Southan GJ, Szabò C: Selective pharmacological inhibition of distinct nitric oxide synthase isoforms. *Biochem Pharmacol* 1996; 51: 384-394
- 136**Dudhgaonkar SP, Tandan SK, Kumar D, et al: Ameliorative effect of combined administration of inducible nitric oxide synthase inhibitor with cyclooxygenase-2 inhibitors in neuropathic pain in rats. *Eur J Pain* 2007; 11: 528-3
- 137**Hoheisel U, Unger T, Mense S: The possible role of the NO-cGMP pathway in nociception: Different spinal and supraspinal action of enzyme blockers on rat dorsal horn neurones. *Pain* 2005; 117: 358-36
- 138**Ashina M, Bendtsen L, Jensen R, et al: Nitric oxide-induced headache in patients with chronic tension-type headache. *J Brain* 2000; 123: 1830-1837

- 1
2
3 **139**Ashina M, Lassen LH, Bendtsen L, et al: Effect of inhibition of nitric oxide
4
5 synthase on chronic tension-type headache: a randomised crossover trial. *Lancet*
6
7 1999; 353: 287-289
8
9
- 10 **140**Gordh T, Blomberg H: The effects of intravenous L-NMMA, a NOS inhibitor, in
11
12 the treatment of patients with neuropathic pain. A double blind, placebo-
13
14 controlled, cross over study. *Svenska Läkaresällskapets Riksstämman* 2004; 82
15
16
- 17 **141**Mayer B, Brunner F, Schmidt K: Inhibition of nitric oxide synthesis by methylene
18
19 blue. *Biochem Pharmacol* 1993; 45: 367-74
20
21
- 22 **142**Mayer B, Brunner F, Schmidt K: Novel actions of methylene blue. *Eur Heart J*
23
24 1993; 14; S I: 22-6
25
26
- 27 **143**Peng B, Zhang Y, Hou S, et al: Intradiscal methylene blue injection for the
28
29 treatment of chronic discogenic low back pain. *Eur Spine J* 2007; 16:33-8
30
31
- 32 **144**Tan KY, Seow-Choen F: Methylene blue reduces pain after lateral anal
33
34 sphincterotomy. *Tech Coloproctol* 2007; 11: 68-9
35
36
- 37 **145**Salaris SC, Babs CF, Voorhees III WD: Methylene blue as an inhibitor of
38
39 superoxide generation by xanthine oxidase. A potential new drug for the
40
41 attenuation of ischemia/reperfusion injury. *Biochem Pharmacol* 1991; 42: 499-506
42
43
- 44 **146**Wiklund L, Basu S, Miculescu A, et al: Neuro- and cardioprotective effects of
45
46 blockade of nitric oxide action by administration of methylene blue. *Ann NY Acad*
47
48 *Sci* 2007;1122: 231–244
49
- 50 **147**Pelgrims J, De Vos F, Van den Brande J, et al: Methylene blue in the treatment
51
52 and prevention of ifosfamide-induced encephalopathy: report of 12 cases and a
53
54 review of the literature. *Br J Cancer* 2000; 82: 291-294
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review