



Published in final edited form as:

*Mov Disord.* 2009 September 15; 24(12): 1840–1842. doi:10.1002/mds.22701.

## No Evidence for Cognitive Dysfunction or Depression in Patients with Mild Restless Legs Syndrome

Erika Driver-Dunckley, MD<sup>1,\*</sup>, Donald Connor, PhD<sup>2</sup>, Joe Hentz, MS<sup>3</sup>, Marwan Sabbagh, MD<sup>2</sup>, Nina Silverberg, PhD<sup>2</sup>, Jose Hernandez, BA<sup>3</sup>, Linda Vedders<sup>2</sup>, Virgilio Gerald Evidente, MD<sup>1</sup>, Holly Shill, MD<sup>2</sup>, John Caviness, MD<sup>1</sup>, and Charles Adler, MD, PhD<sup>3</sup>

<sup>1</sup>Parkinson's Disease and Movement Disorders Center, Mayo Clinic, Scottsdale, Arizona, USA

<sup>2</sup>Sun Health Research Institute, Phoenix, Arizona, USA

<sup>3</sup>Biostatistics Section, Mayo Clinic, Scottsdale, Arizona, USA

### Abstract

Restless legs syndrome is a common disorder that may interrupt sleep and has been reported to produce daytime fatigue and/or mood changes. This study assessed whether patients with RLS have more cognitive dysfunction and depression than individuals of the same age and education who do not have RLS. The study showed that older individuals with mild RLS for at least 1 year do not have cognitive dysfunction and are not depressed compared with a control group of similar age and education.

### Keywords

restless legs syndrome; cognitive dysfunction

---

Restless legs syndrome (RLS) affects up to 25% of the adult population.<sup>1</sup> RLS is characterized by uncomfortable sensations in the legs, which are relieved by movement but commonly worsen at night. This often interrupts sleep and may cause daytime fatigue. It has been reported that patients with RLS suffer from increased rates of irritability, anxiety, and depression.<sup>2–5</sup> However, almost all previous studies (both population-based and clinic-based) have used

---

© 2009 Movement Disorder Society

\*Correspondence to: Dr. Erika Driver-Dunckley, Department of Neurology, Mayo Clinic Scottsdale, 13400 East Shea Blvd, Scottsdale, AZ 85259. driverdunckley.erika@mayo.edu.

Potential conflict of interest: None reported.

**Author Roles:** Erika Driver-Dunckley MD, conception and design, data acquisition or analysis, drafting, editing, or revising of the text. Donald Connor PhD, data acquisition or analysis. Joseph G Hentz MS, data acquisition or analysis. Marwan Sabbagh MD, drafting, editing, or revising of the text. Nina Silverberg PhD, data acquisition or analysis. Jose Hernandez BA, data acquisition or analysis. Linda Vedders data acquisition or analysis. Virgilio GH Evidente MD, drafting, editing, or revising of the text. Holly Shill MD, drafting, editing, or revising of the text. John N Caviness MD, drafting, editing, or revising of the text. Charles H Adler MD, PhD, conception and design, data acquisition or analysis, drafting, editing or revising of the text.

**Financial Disclosures:** Erika Driver-Dunckley MD = none; Donald Connor PhD = consulting and advisory board - MedAvante; Joseph G. Hentz MS = Research funding from Allergan, Astellas Pharma US, Dynatherm Medical, Genentech, and Millenium Pharmaceuticals. Marwan Sabbagh MD = on advisory boards for Elan, Wyeth, Lilly, and Amerisciences. Receives royalty income from Amerisciences and consults for them as well. Nina Silverberg PhD = none; Jose Hernandez BA = none; Linda Vedders = none; Virgilio GH Evidente MD = Consulting for: Teva Neurosciences. Speaking honorarium from: GSK. Research support from: Allergan, Merz Pharmaceuticals, Boehringer, UCB-Schwarz, Neurogen. Holly Shill MD = Consulting fees from Medtronics and Ipsen. Honoraria from Allergan and Teva. Research funding from Boehringer-Ingelheim, Chelsea, Schering-Plough, International Essential Tremor Foundation, Nexalin. John N. caviness MD = none; Charles H. Adler MD, PhD = Consulting fees for advisory boards from GlaxoSmithKline, Ipsen, MerckSerono. Contracts with Ipsen and MerckSerono. Research funding from Allergan and Novartis.

varying RLS diagnostic criteria and various depression scales and have not completely assessed the associated cognitive function.<sup>5</sup> This study assessed whether patients with RLS have more cognitive dysfunction and depression than individuals of the same age and education who do not have RLS.

## SUBJECTS AND METHODS

The Sun Health Research Institute Brain and Body Donation Program (BBDP) database was reviewed for subjects with and without RLS. All subjects had received annual movement disorder and neuropsychological evaluations. Subjects with Parkinson's disease, parkinsonism, essential tremor, other tremor disorders, progressive supranuclear palsy, dystonia, peripheral neuropathy, fibromyalgia, or dementia were excluded from this analysis. The diagnosis of RLS was made by a movement disorder specialist using the IRLSSG criteria.<sup>6</sup> Neuropsychological testing included Rey-AVLT, Trails A and B, stroop, controlled oral word association, animal fluency, judgment of line orientation, digit span, Folstein mini-mental status examination (MMSE), and clock drawing. The 30-item Geriatric Depression Scale (GDS) was used to assess depression.

Data were compiled from the subject's most recent visit that had both movement and cognitive testing. The mean level of each measure in the RLS group was compared with that of the control group and the statistical significance was calculated by using the two-sample *t* test.

## RESULTS

After exclusion criteria were applied there were 26 subjects with RLS, 208 without RLS (control group), and 79 subjects were excluded. There was no difference in mean age (RLS, 77 years; control, 78 years) (Table 1). The 95% CI for the difference in mean age (95% CI = -5.0 to 2.3) indicates that the mean age is equivalent within 5 years. The mean duration of education was 15 years in both groups. The mean duration of education was equivalent within 1 year (95% CI = -0.9 to 1.3).

The mean RLS rating scale score for the RLS group was 11.0 (SD = 7.6, N = 25) and the mean duration of RLS was 11.0 years (range, 1–51 years). Ten RLS subjects had a family history of RLS and one had a family history of Parkinson's disease.

Twelve of the 26 subjects had been or were being treated for RLS (6-gabapentin, 1-ropinirole, 1-pramipexole, 3-narcotics, 1-quinine), 9 had been or were on antidepressants, and 8 had been or were using benzodiazepines. Only 4 patients were using an antidepressant or benzodiazepine at the time of cognitive testing. There were no significant differences in neuropsychometric testing between these two groups of RLS patients. Therefore, they were combined into one group for comparison with controls. Also, no significant difference was found in cognitive testing between the RLS patients on medications for RLS and those who were not on treatment medications. Therefore, these subjects were not separated from the RLS group in the comparative analysis. Medications for the control subjects were not recorded.

None of the mean cognitive scores differed by more than half of the standard deviation of the group without RLS (Table 1). All of the mean cognitive scores were equivalent within one standard deviation of the group without RLS. The mean GDS scores differed by less than 1 point on a 30-point scale. The percentage of subjects with GDS  $\geq 10$  was 6% (n = 3/208) among controls and 4% (n = 1/26) among RLS cases ( $\Delta = -0.02$ , 95% CI = -0.08 to 0.13,  $P > 0.99$ ). The sample was too small to assess for a correlation between the GDS score and severity of RLS.

## DISCUSSION

These findings suggest that older individuals with mild RLS for at least 1 year do not have cognitive dysfunction and are not depressed compared with a control group of similar age and education. Previous epidemiological studies have reported that patients with RLS have more anxiety and depression.<sup>2–5</sup> Whether this is secondary to the actual RLS symptoms, insomnia or another sleep disorder, or other factors is unclear. These prior reports varied significantly in their methodology, including study design (population to random sampling), sample size (range, seven subjects to over 3,000), the RLS diagnostic criteria (the IRLSSG was not normally used), the exclusion or inclusion of patients with a known history of depression, and various assessment scales for evaluation of depression.<sup>2–5</sup> Furthermore, cognitive dysfunction in this population has not been systematically or extensively evaluated. In two previous conflicting reports, untreated RLS patients showed cognitive deficits similar to that seen in sleep-deprived patients, yet the follow-up study found untreated RLS patients to perform better than sleep-deprived controls.<sup>7,8</sup> These studies were limited by a small sample size ( $n = 16$ ), the RLS cases being withdrawn from treatment, and no correlation with RLS severity.<sup>7</sup> These factors make it difficult to assess for medication-withdrawal effects and RLS severity on the cognitive dysfunction found.

The current study used the International RLS Study Group criteria assessed by a movement disorder specialist, the RLS rating scale for quantifying the symptoms, and prospective neuropsychological testing to objectively assess cognitive function. Additionally, patients with depression were not prospectively excluded. Limitations of this study included the cases having relatively mild RLS, no treatment withdrawal for RLS medications ( $n = 12$ ), no controlling for the use of antidepressants or benzodiazepines ( $n = 4$ ), and no measurement for sleep disturbances in either group. Further prospectively designed studies of RLS and control populations are needed to determine if cognitive or neuropsychiatric symptoms occur in more severe cases of RLS and what factors may be correlated with these disorders.

## Acknowledgments

This work was funded by Federal Grants NS40669 and P30 AG019610, the Arizona Biomedical Research Commission (contracts 04-800, 4001, 05-901), the Michael J. Fox Foundation for Parkinson's Research (Prescott Family Initiative), Mayo Foundation for Medical Research, and Sun Health Research Institute.

## REFERENCES

1. Nichols DA, Allen RP, Grauke JH, et al. Restless legs syndrome symptoms in primary care: a prevalence study. *Arch Intern Med* 2003;163:2323–2329. [PubMed: 14581252]
2. Sevim S, Dogu O, Kaleagasi H, Aral M, Metin O, Camdeviren H. Correlation of anxiety and depression symptoms in patients with restless legs syndrome: a population based survey. *J Neurol Neurosurg Psychiatry* 2004;75:226–230. [PubMed: 14742594]
3. Ulfberg J, Nystrom B, Carter N, Edling C. Prevalence of restless legs syndrome among men aged 18 to 64 years: an association with somatic disease and neuropsychiatric symptoms. *Mov Disord* 2001;16:1159–1163. [PubMed: 11748753]
4. Winkelman J, Prager M, Lieb R, et al. *Anxietas tibiarium*: depression and anxiety disorders in patients with restless legs syndrome. *J Neurol* 2005;252:67–71. [PubMed: 15654556]
5. Picchietti D, Winkelman J. Restless legs syndrome, periodic limb movements in sleep and depression: review. *Sleep* 2005;28:891–898. [PubMed: 16124671]
6. Allen RP, Picchietti D, Hening WA, Trenkwalder C, Walters AS, Montplaisir J. A report from the restless legs syndrome diagnosis and epidemiology workshop at the national institutes of health. *Sleep Med* 2003;4:101–119. [PubMed: 14592341]
7. Pearson VE, Allen RP, Dean T, Gamaldo CE, Lesage SR, Earley CJ. Cognitive deficits associated with restless legs syndrome (RLS). *Sleep Med* 2006;7:25–30. [PubMed: 16198145]

8. Gamaldo CE, Benbrook AR, Allen RP, Oguntimein O, Earley CJ. A further evaluation of the cognitive deficits associated with restless legs syndrome (RLS). *Sleep Med* 2008;9:500–505. [PubMed: 17869573]

TABLE 1

Demographic and cognitive measures of subjects with and without restless legs syndrome

	With RLS: mean (SD), N	Without RLS: mean (SD), N	$\Delta$	95% CI	P*
Age (yr)	76.8 (9.2), 26	78.1 (9.0), 208	-1.4	-5.0 to 2.3	0.46
Duration of education (yr)	15.3 (2.8), 26	15.1 (2.6), 208	0.2	-0.9 to 1.3	0.71
Folstein MMSE	28.2 (1.7), 25	28.7 (1.6), 204	-0.5	-1.2 to 0.2	0.15
AVLT total learning	47.7 (9.1), 26	44.7 (11.3), 205	3.0	-1.6 to 7.5	0.19
AVLT Trial 1	5.7 (1.7), 26	5.5 (2.1), 205	0.2	-0.6 to 1.0	0.64
AVLT Trial 2	8.5 (1.9), 26	8.2 (2.6), 205	0.2	-0.8 to 1.3	0.67
AVLT Trial 3	10.0 (2.5), 26	9.6 (2.7), 205	0.4	-0.7 to 1.5	0.49
AVLT Trial 4	11.4 (2.2), 26	10.3 (2.7), 205	1.1	0.01 to 2.2	0.048
AVLT Trial 5	12.1 (2.1), 26	11.0 (2.7), 205	1.1	0.01 to 2.2	0.047
AVLT intrusions (1-5)	0.6 (1.2), 25	1.5 (2.3), 200	-0.9	-1.8 to 0.0	0.06
AVLT STM (A6)	9.6 (3.4), 26	9.0 (3.3), 205	0.5	-0.8 to 1.9	0.43
AVLT intrusions (STM)	0.36 (0.70), 25	0.50 (0.96), 200	-0.14	-0.52 to 0.25	0.50
AVLT LTM (A7)	9.0 (3.5), 26	8.9 (3.7), 205	0.1	-1.4 to 1.6	0.88
AVLT LT % recall	73 (24), 26	79 (26), 205	-6	-17 to 4	0.25
AVLT List B	5.2 (2.0), 26	4.8 (2.0), 204	0.4	-0.4 to 1.2	0.33
AVLT recognition TP	13.5 (1.7), 25	13.2 (2.3), 199	0.3	-0.7 to 1.2	0.56
AVLT recognition FP	0.20 (0.41), 25	0.64 (1.25), 199	-0.44	-0.94 to 0.05	0.08
WAIS digit span - forward	8.6 (1.5), 24	9.6 (2.0), 188	-1.0	-1.8 to -0.2	0.02
WAIS digit span - backward	6.2 (1.8), 24	6.4 (2.0), 187	-0.2	-1.1 to 0.6	0.60
WAIS digit span - total	14.8 (2.6), 24	16.0 (3.3), 187	-1.2	-2.6 to 0.2	0.09
WAIS digit span - backward span	4.6 (1.2), 23	4.7 (1.2), 182	-0.1	-0.6 to 0.4	0.74
WAIS digit span - forward span	5.7 (0.9), 23	6.3 (1.2), 183	-0.6	-1.1 to -0.1	0.03
Clock drawing	9.0 (1.6), 25	9.2 (1.1), 204	-0.3	-0.8 to 0.2	0.26
STROOP word (#)	87 (15), 24	89 (16), 199	-2	-8.9 to 5.0	0.58
STROOP color (#)	63 (12), 23	63 (12), 196	0	-5 to 5	0.98
STROOP word/color (#)	32.6 (7.0), 23	33.0 (10.1), 19	-0.4	-4.6 to 3.9	0.86
STROOP uncorrected W/C errors	0.4 (1.1), 22	0.4 (1.0), 191	0.1	-0.4 to 0.5	0.72
STROOP interference	-4.0 (5.4), 23	-4.0 (7.2), 195	0.0	-3.1 to 3.0	0.99
TRAILS A/B - time (3 min limit)	37 (16), 24	38 (22), 203	-1	-10 to 78	0.79

	With RLS: mean (SD), N	Without RLS: mean (SD), N	$\Delta$	95% CI	$P^*$
TRAILS A/B - A errors	0.13 (0.34), 24	0.12 (0.42), 203	0.01	-0.17 to 0.18	0.94
TRAILS A/B - B (5 min limit)	101 (43), 24	102 (53), 200	-1	-24 to 21	0.90
TRAILS A/B - B errors	0.46 (0.72), 24	0.57 (1.11), 201	-0.11	-0.57 to 0.35	0.64
COWA (CFL) total	38 (16), 26	38 (12), 205	1	-4 to 6	0.79
COWA (CFL) perseverations	2.8 (2.9), 25	2.8 (2.7), 200	0.0	-1.2 to 1.1	0.97
COWA (CFL) intrusions	0.08 (0.28), 25	0.29 (0.64), 200	-0.21	-0.46 to 0.05	0.12
Animal fluency	16.7 (5.2), 26	16.8 (5.0), 204	-0.1	-2.2 to 2.0	0.91
Animal fluency perseverations	1.0 (1.4), 25	1.0 (1.3), 199	0.0	-0.5 to 0.5	1.00
Judgment of line orientation	23.6 (3.9), 23	23.7 (4.3), 191	-0.1	-2.0 to 1.7	0.89
GDS	3.7 (3.2), 26	3.4 (3.7), 206	0.4	-1.1 to 1.8	0.63

\* Two sample *t* test.