

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

TITLE NOISE-SUSTAINED STRUCTURE, INTERMITTENCY, AND THE GINZBURG-LANDAU EQUATION

LA-UR--85-1055

AUTHOR(S) Robert J. Deissler

DE85 009626

SUBMITTED TO Proceedings of International Conference on Solitons and Coherent Structures In Santa Barbara, CA (Jan. 1985)

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U.S. Government and certain other persons have certain rights in the published form of this contribution, and it is to be used for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

WASTE

Los Alamos Los Alamos National Laboratory Los Alamos, New Mexico 87545



**Noise-Sustained Structure, Intermittency,  
and the Ginzburg-Landau Equation**

Robert J. Deissler \*

Center for Nonlinear Studies, MS B258, Los Alamos National Laboratory

Los Alamos, New Mexico 87545

\* Permanent Address: Physics Department, 307 Nat Sci II  
University of California at Santa Cruz, Santa Cruz, CA 95064

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

6/1/83

The time-dependent generalized Ginzburg-Landau equation is a partial differential equation that is related to many physical systems. In the stationary (ie. laboratory) frame of reference the equation is:

$$\frac{\partial \psi}{\partial t} = a \psi - v \frac{\partial \psi}{\partial x} + b \frac{\partial^2 \psi}{\partial x^2} - c |\psi|^2 \psi \quad (1)$$

where the dependent variable  $\psi$  is in general complex;  $a$ ,  $b$ , and  $c$  are constants which are in general complex; and  $v$  is the group velocity.

Consider a small initial localized perturbation about the equilibrium state  $\psi = 0$ . A linear stability analysis reveals that there are three types of behavior which this perturbation can undergo. 1) The perturbation will be damped in any frame of reference. This behavior corresponds to the system being *absolutely stable*. 2) The perturbation will grow and spread such that the edges of the perturbation move in opposite directions. This behavior corresponds to the system being *absolutely unstable*. 3) The perturbation will be damped at any given stationary point but a frame of reference may be found in which the perturbation is growing. In other words, even though the perturbation is growing and spreading, it is moving at a sufficiently large velocity such that both edges of the perturbation are moving in the same direction. Thus the system behind the perturbation returns to its undisturbed state. This behavior corresponds to the system being *spatially unstable* (ie. convectively unstable).

Under conditions when the system is absolutely unstable, an initial small (microscopic) perturbation will grow to macroscopic size, saturate (assuming  $c_p \neq 0$ ), and produce a structure for all time. In contrast, if the system is spatially unstable, the perturbation and resulting structure will move spatially such that the structure eventually leaves the boundaries of the system. Thus the system returns to the equilibrium state

A *single* perturbation therefore produces only a *temporary* structure. However if the the system is continuously perturbed by microscopic external noise it will be unable to return to the equilibrium state and a new state will be established which is sustained by the presence of the noise (ie. a *noise-sustained state*).

Under conditions in which the system is spatially unstable (ie.  $a_r = \frac{r^2 b_r}{4|b_r|^2} < 0$  and  $a_r > 0$ ), eq. (1) is numerically solved in the presence of low-level external noise. At the conference the time evolution of  $\psi$  plotted as a function of  $x$  was shown via a movie. Two striking features were the coherent structure forming as a result of the selective amplification of the external noise and the interspersion of this coherent structure with turbulent behavior.

Fig. 1 shows one frame of the movie --  $\psi$  plotted as a function of  $x$  for a particular  $t$  after the system has reached a statistically steady state. The microscopic noise near the left boundary grows spatially to macroscopic proportion resulting in the observed structure. This is a *noise-sustained structure*. If the external noise is removed the structure moves out through the right boundary and the system returns to the state  $\psi = 0$  everywhere except for some slight ( $\approx 10^{-100}$ ) fluctuations due to computer roundoff.

The intermittency is seen in fig. 2 which shows  $\psi$  plotted as a function of  $t$  at  $x = 150$ . This intermittent behavior may be qualitatively understood as follows. The microscopic noise near the left boundary is spatially and selectively amplified resulting in the formation of spatially growing waves. These waves are sufficiently regular to produce the coherent structure seen in fig. 1. However the coherent structure itself is *spatially unstable* (ie. a secondary spatial instability). This spatial instability causes the coherent structure to break up at some spatial point. The point at which it breaks

up depends on the degree of irregularity in the spatially growing waves. Since the irregularities in the spatially growing waves change with time (the source of the waves being random noise), the point at which the coherent structure breaks up changes with time resulting in intermittency.

A few other points are: 1) The external noise (or other external perturbation) was necessary for the formation of the structure -- no noise  $\rightarrow$  no structure. 2) The *microscopic* noise played an important role in the *macroscopic* dynamics of the system (eg. the intermittency). 3) The chaotic behavior is not associated with a strange attractor (ie. not deterministic chaos). 4) The system exhibits a laminar region followed spatially by a turbulent region. This type of behavior occurs in many fluid systems such as pipe flow, fluid flow over a flat plate, and smoke rising from a cigarette. 5) A few fluid systems from which eq. (1) (with nonzero group velocity  $v$ ) has been derived are plane Poiseuille flow and wind-induced water waves. A nonzero group velocity is necessary in order for the equation to exhibit a spatial instability.

For references and a more detailed account the reader is referred to ref. [1] and the references contained therein.

This work was partially supported by the Air Force Office of Scientific Research under AFOSR grant #ISSA-84-00017 and the U. S. Office of Naval Research under contract #N00014-80 C-0657.

### References

- 1) R. J. Deissler, "Noise-Sustained Structure, Intermittency, and the Ginzburg-Landau Equation", Los Alamos preprint #LA-UR-84-3629 and submitted to J. Stat. Phys.

### Figures

- 1) Plot of  $\psi_r$  as a function of  $x$  for a given  $t$  ( $t = 300$ ) after transients have settled down.  $a = 2$ ,  $v = 6$ ,  $b_r = 2.8$ ,  $b_i = -1$ ,  $c_r = .5$ ,  $c_i = 1$ . Noise level  $= r = 10^{-6}$ .  
The noise, a random number uniformly distributed between  $-r$  and  $r$ , is added to the grid point adjacent to the left boundary. The microscopic noise at this grid point grows spatially to macroscopic proportion resulting in the observed structure.
- 2) Plots of  $\psi_r$  as a function of  $t$ .  $a = 2$ ,  $v = 6$ ,  $b_r = 2.8$ ,  $b_i = -1$ ,  $c_r = .5$ ,  $c_i = 1$ ,  
 $x = 150$ . High and low frequencies are seen to be interspersed.





