

REVIEW

Non-coding RNAs: long non-coding RNAs and microRNAs in endocrine-related cancers

Carolyn M Klinge

Department of Biochemistry & Molecular Genetics, Center for Genetics and Molecular Medicine, University of Louisville School of Medicine, Louisville, Kentucky, USA

Correspondence should be addressed to C M Klinge: carolyn.klinge@louisville.edu

Abstract

The human genome is 'pervasively transcribed' leading to a complex array of non-coding RNAs (ncRNAs) that far outnumber coding mRNAs. ncRNAs have regulatory roles in transcription and post-transcriptional processes as well numerous cellular functions that remain to be fully described. Best characterized of the 'expanding universe' of ncRNAs are the ~22 nucleotide microRNAs (miRNAs) that base-pair to target mRNA's 3' untranslated region within the RNA-induced silencing complex (RISC) and block translation and may stimulate mRNA transcript degradation. Long non-coding RNAs (lncRNAs) are classified as >200 nucleotides in length, but range up to several kb and are heterogeneous in genomic origin and function. lncRNAs fold into structures that interact with DNA, RNA and proteins to regulate chromatin dynamics, protein complex assembly, transcription, telomere biology and splicing. Some lncRNAs act as sponges for miRNAs and decoys for proteins. Nuclear-encoded lncRNAs can be taken up by mitochondria and lncRNAs are transcribed from mtDNA. Both miRNAs and lncRNAs are dysregulated in endocrine cancers. This review provides an overview on the current understanding of the regulation and function of selected lncRNAs and miRNAs, and their interaction, in endocrine-related cancers: breast, prostate, endometrial and thyroid.

Key Words

- ▶ miRNA
- ▶ lncRNA
- ▶ ncRNA
- ▶ transcription
- ▶ translation

Endocrine-Related Cancer
(2018) **25**, R259–R282

Introduction

Cancer is a disease of the genome in which tumors have a constellation of genomic and epigenetic alterations that drive their clinical behavior and patient prognosis (Macconail & Garraway 2010). The Central Dogma proposed by Francis Crick in 1958 envisioned that information flowed from DNA to mRNA, which is translated into functional proteins with the assistance of tRNA and rRNA (reviewed in Morris & Mattick 2014, Jarroux *et al.* 2017). With time, additional classes of RNA that have important regulatory roles in cellular biology were discovered: short nuclear and small nucleolar RNAs (snRNAs and snoRNAs), followed by microRNAs (miRNAs) and long non-coding RNAs (lncRNAs). The Encyclopedia

of DNA Elements (ENCODE) Consortium was started in 2003 as an international collaboration funded by the National Human Genome Institute (NHGRI). Analysis of the human genome revealed that while ~85% is transcribed, only ~1% is protein-coding mRNA (ENCODE Project Consortium 2012, Djebali *et al.* 2012, Dykes & Emanueli 2017). The GENCODE consortium (<https://www.genencodegenes.org/>) is a subproject of ENCODE that produces high-quality reference gene annotation and experimental validation. The current GENCODE, version 27 of the human genome includes 58,288 genes; 200,401 total transcripts; 19,826 protein-coding genes; 15,778 lncRNAs; 14,694 pseudogenes; 7569 small non-coding

RNA (ncRNA) genes and 1881 miRNAs. Most (85%) small ncRNAs are snRNAs, snoRNAs, miRNAs and tRNAs (Djebali *et al.* 2012). In addition to GENCODE, the FANTOM (Functional Annotation of Mammalian Genomes) project is creating a comprehensive catalog of transcripts encoded in human, mouse, rat, dog, Rhesus monkey and chicken genomes (<http://fantom.gsc.riken.jp>) (Kawaji *et al.* 2017). Analysis of data from FANTOM5 identified 27,919 human lncRNA genes with high-confidence 5' ends and provided evidence that 69% (19,175) are functional (Hon *et al.* 2017). The history of the discovery of regulatory RNAs, notably the ncRNAs, has been reviewed (Morris & Mattick 2014).

The advent of next-generation sequencing (NGS) by RNA sequencing (RNA-seq), also called 'whole transcriptome shotgun sequencing' and 'deep RNA-seq', has allowed characterization of the cellular transcriptomes, i.e., the entire spectrum of RNAs produced in a cell type (Wolf 2013), resulting in a glut of data that can be mined by investigators. One database of transcriptomes of potential interest for readers of this review is the NURSA website (www.nursa.org). The transcriptome includes mRNA, rRNA and tRNA; and the ncRNAs: miRNAs, enhancer RNAs (eRNAs), endogenous small-interfering RNAs (siRNAs), Piwi-interacting RNAs (piRNAs), circular RNAs (circRNAs) and lncRNAs (Marrone *et al.* 2014). Table 1 highlights features of these ncRNAs and their roles in cellular processes. The function of piRNAs and snoRNAs in endocrine-related cancers was reviewed (Venkatesh *et al.* 2015). miRNAs are best characterized for their gene silencing of target mRNA by complementary base-pairing with the 3'UTR within the RNA-induced silencing complex (RISC) to repress translation and/or cause mRNA degradation (Muluhngwi & Klinge 2015). Like miRNAs, siRNAs and piRNAs bind argonaute (AGO) family members and base-pair with target RNA to cause RNA degradation and/or translation repression (Watanabe & Lin 2014). Studies have shown that RNA polymerase (pol) II-mediated transcriptional activity at gene enhancers is 'pervasive' and bidirectional, generally producing transcripts in both directions including short eRNAs that play roles in enhancer-promoter interactions (reviewed in Lam *et al.* 2014). Enhancers are relatively insensitive to position and distance of their target genes and about half are intragenic, which can result in inhibition or attenuation of nascent transcript elongation (Cinghu *et al.* 2017). circRNAs are usually formed by noncanonical splice reactions and reside in the cytoplasm where one circRNA, CDR1AS, regulates miR-7 stability and transport in neurons (Piwecka *et al.* 2017).

Both miRNAs and lncRNAs are epigenetic regulators of human cancers (Liz & Esteller 2016). Moreover, these RNAs are chemically modified, e.g., by post-transcriptional methylation on N6 of adenosine (m⁶A), which alters their activities, providing an additional layer of regulation termed 'epitranscriptomics' (Li *et al.* 2016c). Many more papers have been published on miRNAs than lncRNAs in endocrine cancers (Fig. 1). This review provides a brief overview of the identity, regulation and roles of miRNAs and lncRNAs in selected endocrine-related cancers. Because of the scope of published literature on these topics, this review is not a comprehensive analysis of lncRNAs and miRNAs in breast, prostate, endometrial and thyroid cancers. The reader is provided with citations to recent reviews and primary literature for further information.

lncRNAs: classification and functional activities

lncRNAs are defined as ncRNAs of >200 nucleotides that are transcribed by RNA pol II from regulatory regions including promoter upstream elements and technically include eRNAs (Andersson *et al.* 2014). However, eRNAs differ from lncRNAs by being shorter (<2 kbp), without being spliced and are not polyadenylated (Kim *et al.* 2015). By total cellular RNA mass, lncRNA constitutes only 0.03–0.2% while mRNA constitute 3–7% (Dykes & Emanuelli 2017). lncRNAs are classified based on their genomic organization: (1) intergenic lncRNAs (lincRNAs) are transcribed between two protein-coding genes; (2) intronic lncRNAs are transcribed from introns of protein-coding genes; (3) overlapping lncRNAs, as the name implies, constitute transcripts that overlap known protein-coding genes; and (4) antisense (as) lncRNAs are transcribed in a direction opposite that of the protein-coding gene (Peng *et al.* 2017b) (Fig. 2). Pseudogenes that arise from DNA duplication of coding genes followed by inactivating mutations that render the pseudogene non-coding are transcribed as lncRNAs and sequester some of the miRNAs that interact with conserved miRNA response elements (MREs) in the functional gene (Dykes & Emanuelli 2017). However, given the low levels of lncRNA, the functional role of lncRNAs as 'miRNA sponges' by base-pairing and thus blocking miRNA-MRE binding to target mRNA-3'UTRs, is debated. It seems likely that more highly expressed lncRNAs (with higher FKPM (Fragments Per Kilobase Million reads in RNA-seq)) would be more likely to act as competing endogenous RNA (ceRNA) to sequester miRNAs (Dykes & Emanuelli 2017). Additional classifications of lncRNAs are by (1) length,

Table 1 Summary of ncRNAs in humans, excluding rRNA and tRNA, with their size, cellular location, aspects of biogenesis, function, citations for reviews for roles in various endocrine-related cancers and websites.

ncRNA	Description, size, location, function and example of findings in endocrine cancers	Websites
miRNA	<p>~21 nt ssRNA</p> <ul style="list-style-type: none"> Nucleus and cytoplasm Transcribed as pri-RNAs, processed from pre-miRs bearing one or more inverted repeats or hairpins Form complimentary base-pairs with the 3' untranslated region of target mRNAs within RISC block translation and/or stimulate mRNA transcript degradation Expressed in a tissue-specific manner Most are considered highly stable IsomiRs are sequence variants of the canonical miRNA currently in the miRBase generated from a single miRNA locus by template and non-template variants miRNAs in endocrine cancers have been reviewed: thyroid, adrenal, pancreatic, and pituitary cancers (Lima <i>et al.</i> 2017); endocrine-resistant breast (Muluhngwi & Klinge 2015); prostate, ovarian, and breast (Smith <i>et al.</i> 2017) 	<p>miRBase http://www.mirbase.org miRTarBase: experimentally validated miRNA-target Interactions http://miRTarBase.mbc.nctu.edu.tw miRAD: intragenic miRNA database http://bmi.ana.med.uni-muenchen.de/mirad/ MAGIA http://gencomp.bio.unipd.it/magia/start/ TargetScanHuman search for predicated miRNA targets http://www.targetscan.org/vert_71/ IsomiRage http://cru.genomics.iit.it/Isomirage/</p>
circRNAs	<p>Circular RNAs: 4 types: exonic circRNAs (ecircRNA), circular RNAs from introns, exon-intron circRNAs (EicRNA) and intergenic circRNAs (Meng <i>et al.</i> 2017)</p> <ul style="list-style-type: none"> Cytoplasm Generally formed by alternative splicing of pre-mRNA in which an upstream splice acceptor is joined to a downstream splice donor in a process known as 'backsplicing' (Barrett & Salzman 2016) Expressed in thousands of human genes Stable: half-life >48 h Expressed in a cell-type and tissue-specific manner Act as miRNA sponges, interact with RNA-binding proteins, can be positive regulators of their parental genes Can be translated (Yang <i>et al.</i> 2017c) circRNAs were identified in human papillary thyroid cancer (Peng <i>et al.</i> 2017a) and in human breast tumors (Liang <i>et al.</i> 2017) 	<p>CIRCexplorer (Yang <i>et al.</i> 2017c) https://omictools.com/circexplorer-tool CIRI https://omictools.com/ciri-tool CircRNA_finder https://omictools.com/circrna-finder-tool MapSplice 2 for mapping RNA-seq data to ref genome for splice junction discovery http://www.netlab.uky.edu/p/bioinfo/MapSplice2 CircIncRNAet (RRIC:SCR_015794) https://dknet.org/scicrunch/Resources/record/nlx_144509-1/2d26f6b1-5909-5066-a261-a05ca13e12d9/search?mc_cid=23e8e71213&mc_eid=ed4995675c</p>
eRNAs	<p>Enhancer RNAs are produced from enhancers (Lam <i>et al.</i> 2014)</p> <ul style="list-style-type: none"> Nuclear, 5' methyl-capped, +/- polyadenylation 800–2000 bp Low stability and abundance Usually function in <i>cis</i> to increase transcription by facilitating DNA looping between the enhancer and promoter May facilitate histone modifications and chromatin remodeling (Li <i>et al.</i> 2016b) CGA, the common α-subunit of LH, FSH, and TSH is regulated by an eRNA (Nagarajan <i>et al.</i> 2015) 	
piRNA	<p>PIWI-interacting RNAs</p> <ul style="list-style-type: none"> 24–32 nt ss RNAs Derived from piRNA clusters Nuclear and cytoplasmic Bind PIWI subfamily of Argonaut proteins Involved in gene silencing (Luteijn & Ketting 2013) Protect genome against instability by repressing transposon activity via transcriptional gene silencing Expression mostly restricted to germ cells <p>The transposable element (TE) landscape in postnatal human testis is matched by a relevant piRNA repertoire to silence TEs in spermatogenesis (Gainetdinov <i>et al.</i> 2017)</p>	

(Continued)

Table 1 Continued.

ncRNA	Description, size, location, function and example of findings in endocrine cancers	Websites
Endo-siRNA	<p>Endogenous small-interfering RNAs</p> <ul style="list-style-type: none"> • 20–23 nt • Cytoplasmic • Processed from dsRNAs by DICER • Have a 3' 2-nt overhang • Form an effector complex with AGO1 or AGO2 (Piatek Monica & Werner 2014) • Detected in mammalian embryonic stem cells, oocytes and spermatocytes (Hilz et al. 2016) 	
lncRNAs	<p>Long non-coding RNAs</p> <ul style="list-style-type: none"> • >200 nt • Can be several kb in length • Transcribed by RNA pol II • 5'-Methyl capped • Some are polyadenylated • Expressed in a cell-specific manner • Interact with DNA, RNA, and proteins • Stability generally lower than mRNAs with intronic or promoter-associated lncRNAs less stable than intergenic or antisense lncRNAs (Jarroux et al. 2017) • Located in nucleus, cytoplasm, and mitochondria • Regulate gene expression, chromatin modification and dynamics, protein complex assembly, splicing, and translation • Regulate development and differentiation including embryonic development (Bouckenheimer et al. 2016) • X chromosome silencing by <i>XIST</i>: a 16 kb lncRNA (Cerase et al. 2015) <p>Predicting miRNA/target duplex (Rehmsmeier et al. 2004) has been used to examine miRNA/lncRNA interaction (Zhou et al. 2017)</p>	<p>HUGO Gene Nomenclature Committee lncRNAs https://www.genenames.org/cgi-bin/genefamilies/set/788</p> <p>NONCODE v. 5.0 ncRNA http://www.noncode.org/ lncRNASNP2 is a database of functional SNPs and mutations in human and mouse lncRNAs http://bioinfo.life.hust.edu.cn/lncRNASNP2</p> <p>Websites reviewed in (Fritah et al. 2014)</p> <p>lncpedia 4.1 https://lncpedia.org/ (146,742 human annotated lncRNAs)</p> <p>DIANA-lncBase v.2 includes miRNA:lncRNA interactions (Paraskevopoulou et al. 2016)</p> <p>NONCODE v5.0 http://www.noncode.org/index.php</p> <p>http://bibiserv.techfak.uni-bielefeld.de/rnahybrid/</p>

e.g. very long lincRNA (vlincRNA); (2) being encoded within specific DNA regulator elements and loci, e.g., in telomeres (*TERRA*), centromeres, rDNA loci (*PAPAS*) or 3'UTR-associated RNAs (uaRNA); (3) biogenesis pathway; (4) subcellular localization or origin, e.g., nuclear, cytoplasmic or mitochondrial; (5) function, e.g., scaffold lncRNA (*HOTAIR*) and (6) association with specific biological processes (Jarroux et al. 2017).

lncRNAs show chromatin marks that are typical of transcribed coding genes: H3K4me3 for transcription initiation and H3K36me3 for elongation (Rinn & Chang 2012) and are regulated by transcription factors binding to their promoters. lncRNAs are alternatively spliced, thus generating isoforms. lncRNAs associated with enhancer regions can act in *cis*, leading to increased transcription of neighboring genes, including other lncRNAs (Ørom Ulf & Shiekhattar 2013). Some lncRNAs directly bind target mRNAs which can either target the mRNA for degradation or stabilize the transcript. A class of lncRNAs called 'activating ncRNA-as' specifically activate neighboring coding genes by binding to specific DNA sequences while

associating with the Mediator-cohesin complex to loop enhancer/promoter regions (Dykes & Emanuelli 2017). Despite their length, which can be several kb, lncRNAs do not have a functional protein-encoding capacity, i.e., those that contain small open reading frames (ORFs) do not form a template for translation (Hu et al. 2012).

As indicated in the Introduction, the number of human lncRNAs varies between databases. For example, the NONCODE v5 database includes 96,308 lncRNA genes (Xiyuan et al. 2017) compared with 15,778 in GENECODE and 27,919 in FANTOM (Hon et al. 2017). lncRNAs are expressed in a tissue-specific manner and are usually 5'-end capped and some are 3'-end polyadenylated (Li et al. 2016a). Some lncRNAs, e.g., *MALAT1* and *NEAT1*, have a triple-helical structure at the 3'end that protects them from degradation. The landscape of isoforms of lncRNAs adds to the complexity of understanding lncRNA function (Ziegler & Kretz 2017). In addition, the presence of lncRNA modifications and lncRNA editing has been reported and associated with structural and functional changes, which further increases the variety of




Figure 1
PubMed referenced papers on miRNAs and lncRNAs for endocrine cancers. Values were determined on November 28, 2017.

lncRNAs (reviewed in Ziegler & Kretz 2017). For example, m⁶A modification is required for *XIST1*-mediated gene silencing (Patil *et al.* 2016). Accumulating evidence implicates roles for dysregulated lncRNAs in cancer (Huarte 2015), autoimmune and inflammatory diseases (Chen *et al.* 2017), including type 1 (Mirza *et al.* 2017) and type 2 diabetes (Motterle *et al.* 2016, Leti & DiStefano 2017) and other diseases.

lncRNA function depends on cellular location (Fig. 3). lncRNAs fold in various conformations that allows interaction with RNA, DNA and proteins to

regulate nuclear, cytoplasmic (Delas & Hannon 2017) and mitochondrial function (De Paepe *et al.* 2017). lncRNAs regulate processes including gene transcription, chromatin dynamics, chromatin looping, histone modifications, telomere biology, protein complex assembly, RNA splicing and translation (Mercer & Mattick 2013). Within the nucleus, specific lncRNAs, including *AIRN*, *ANRIL*, *HOTAIR* and *XIST* interact with specific transcription factors, enzymes involved in DNA methylation (DNMT1), chromatin modification (HDAC1, EZH2 (a lysine N-methyltransferase)) and polycomb repressive complex 2 (PRC2) (Nakagawa & Kageyama 2014). lncRNAs are involved in assembly of active e.g., *NEAT1*, or repressed, e.g., *XIST*, nuclear chromatin domains in a cell-dependent manner (Rinn & Guttman 2014). *NEAT1* is an architectural RNA that interacts with >60 RNA-binding proteins and transcription factors in paraspeckles (Yamazaki & Hirose 2015). *NEAT1* is also acts as a ‘sponge’ for miR-214 and miR-101, thus upregulating targets of these miRNAs, i.e., *HMGAI* and *EZH2*, respectively (Qian *et al.* 2017, Wang *et al.* 2017a,c). Thus, many biological processes are regulated by lncRNAs including cell differentiation, proliferation and survival.

Functional roles of lncRNAs in the cytoplasm including acting as a scaffold. For example, the RNA stem-loop structure of *LINC01139* (also called LINK-A) provides a scaffold for HBEGF (heparin-binding epidermal growth factor (EGF)-like growth factor)-triggered, EGF receptor (EGFR):GPNMB (glycoprotein NMB) heterodimer-mediated recruitment of PTK6 (protein tyrosine kinase 6, also called BRK) and LRRK2 (leucine-rich repeat kinase-2) for phosphorylation and stabilization of HIF-1 α in MDA-MB-231 triple-negative breast cancer (TNBC) cells (Lin *et al.* 2016). *LINC01139* also binds phosphatidylinositol-3,4,5-trisphosphate (PIP3) and




Figure 2
Classification of lncRNAs based on their location and transcription in relation to protein-coding genes. Gene structure is diagramed 5’-3’, left to right, for RNA pol II transcription.




Figure 3

Examples of interactions of lncRNAs and miRNAs in endocrine-related cancers. Not all these interactions may be expected in one cell type. lncRNAs can act in cis on an adjacent gene to regulate its transcription and translation. An example is *HOTAIR*'s interaction with AR (Zhang *et al.* 2015a). Nuclear miRNAs can be transported into mitochondria through unknown mechanisms and miRNAs are produced from the mtDNA genome (ds DNA plasmids shown), both are called 'mitomiRs'. lncRNAs are transcribed from mtDNA and are called mitolncRNAs. Pri-miRNAs are processed to pre-miRNAs by the Drosha microprocessor complex and exported by Exportin/RAN GTPase to the cytoplasm where they are cleaved into mature miRNAs by DICER which interacts with the lncRNA *SRA1* (Redfern *et al.* 2013). *LINC01139* acts as a scaffold for PIP3, BRK, LARK and GPNMB, which leads to HIF-1 α phosphorylation (Lin *et al.* 2017). *MALAT1* acts as a ceRNA and 'sponges' miRNAs including miR-101, miR-125b and miR-217 (Table 2 and Supplementary Table 1).

promotes AKT recruitment and phosphorylation (Lin *et al.* 2017). Several lncRNAs modulate mRNA stability (reviewed in Rashid *et al.* 2016). lncRNAs also associate with ribonucleoprotein complexes to regulate translation. For example, *ZFAS1* associates with the 40S ribosome in the cytoplasm of MDA-MB-468 breast cancer (BCa) cells and is downregulated in breast tumors compared with normal breast tissue (Hansji *et al.* 2016).

Examples of lncRNAs dysregulated in endocrine-related cancers

Although many lncRNAs have been identified using bioinformatic approaches, relatively few have been

functionally characterized and little is known about the mechanism(s) of action of even the best characterized lncRNAs. This review is not comprehensive with respect to lncRNAs in endocrine-related cancers, but summarizes representative lncRNAs that are best characterized (Table 2 and described below). Examples of recent reviews of lncRNAs include roles in: female reproductive cancers (Ong *et al.* 2017), BCa (Malih *et al.* 2015, Venkatesh *et al.* 2015, Miano *et al.* 2016, Niknafs *et al.* 2016, Warburton & Boone 2017), prostate cancer (PCa) (Alahari *et al.* 2016, Zhang *et al.* 2016a,c, Yang *et al.* 2017b) and thyroid cancer (Murugan *et al.* 2017).

One of the first characterized lncRNAs, *HOTAIR* (HOX Transcript Antisense RNA) is a 2.2 kb lncRNA

found only in mammals and its structure is conserved, despite changed sequences, especially in its protein-binding segments, implying evolutionary importance (Jarroux *et al.* 2017). *HOTAIR* is transcribed from the HOXC locus from a position intergenic and antisense to the flanking HOXC11 and HOXC12 genes. *HOTAIR* facilitates the HOXD gene cluster for transcriptional repression by recruiting PRC2 (Rinn *et al.* 2007). Analysis of data in the Cancer Genome Atlas (TCGA) revealed that increased *HOTAIR* expression correlates with metastasis and poor prognosis in breast and prostate cancers (Weidle *et al.* 2017). *HOTAIR* is both nuclear and cytoplasmic and considered an onco-lncRNA because of its ability to promote papillary thyroid cancer (PTC) cell proliferation and silencing its expression inhibited cell growth (Zhu *et al.* 2016). *HOTAIR* is a scaffold for the PRC2 and LSD1 complexes affecting H3K27 methylation and H3K4 demethylation for epigenetic gene silencing promoting cancer metastasis (Wu *et al.* 2014). *HOTAIR* decreases oxidative phosphorylation (OXPHOS) complex III subunit VII (UQCRCQ) protein expression, thus reducing OXPHOS efficiency and suggesting that *HOTAIR* is important for mitochondrial function in HeLa cells (Zheng *et al.* 2015).

According to Gene Cards (www.genecards.org), *HOTAIR* has binding sites for 70 proteins involved in transcriptional regulation including AGO1, SIN3A, REST, SUZ12, RELA, ROSL1, MAX, MAZ, NR2C2 (orphan NR TR4), TBP, EGR2, ELK4 and SP2. *HOTAIR* is upregulated by HIF-1 α , histone methylase MLL1 and coactivator p300 (Bhan *et al.* 2017). *HOTAIR* was reported to be induced by E₂ in MCF-7 BCa cells (Bhan *et al.* 2013), but others reported that E₂-ER α directly that repressed *HOTAIR* in MCF-7 cells by binding to a genomic site 14.5 kb upstream of *HOTAIR* (Xue *et al.* 2016b). *HOTAIR* is downregulated by miR-141 (Chiyomaru *et al.* 2014) and is directly targeted by miR-34a in PCa cells (Chiyomaru *et al.* 2013). The 3' domain of *HOTAIR* binds lysine-specific demethylase 1 (LSD1)-CoREST and PRC2 interacts with a 5' domain. *HOTAIR*-PRC2 interaction represses targets, e.g., HOXD10, PTEN, miR-7 and WIF1, which leads to the activation of the Wnt/ β -catenin signaling pathway (Hajjari & Salavaty 2015). *HOTAIR* binds androgen receptor (AR), blocking MDM2 interaction, and preventing AR ubiquitination and protein degradation, thus contributing to castration-resistant prostate cancer (CRPCa) progression (Zhang *et al.* 2015). *HOTAIR* is upregulated in endocrine-resistant BCa cells and its overexpression activates ER α transcriptional

Table 2 Examples of the roles of selected lncRNAs identified in endocrine-related cancers. Name, size, and genome location information is summarized from the sources for lncRNAs in Table 1. Other information is from the references cited.

lncRNA gene	Name, size, location	Interactions	Regulation	Endocrine cancer
<i>PCA3</i>	Prostate cancer-associated 3 23,134 nt 9q21.2 Antisense lincRNA within <i>PRUNE2</i>	<i>PCA3</i> and <i>PRUNE2</i> pre-mRNA are co-expressed and form a double-stranded RNA that recruits adenosine deaminase acting on RNA (ADAR) proteins to form a complex (Zhang <i>et al.</i> 2016a,c)	Increased by androgens in prostate cancer cells (Salameh <i>et al.</i> 2015) Suppressed by ER β (Zhang <i>et al.</i> 2016a,c)	PCa-specific urine biomarker (Laxman <i>et al.</i> 2008); levels inversely correlate with the tumor suppressor <i>PRUNE2</i> in human PCa specimens (Salameh <i>et al.</i> 2015)
<i>GAS5</i>	Growth arrest specific 5 1q25.1 The <i>GAS5</i> gene has 12 exons that are alternatively spliced into 2 mature lncRNAs and <i>GAS5</i> introns encode 10 snoRNAs (reviewed in Pickard & Williams 2014) Tumor suppressor	Forms hairpin structures and interacts with the DNA binding domain of GR to inhibit GR-GRE binding and GR-induced gene transcription (Kino <i>et al.</i> 2010) Represses onco-miR-21 (Zhang <i>et al.</i> 2013) Inhibited miR-103 in EC cells (Guo <i>et al.</i> 2015) Inhibited activation of the AKT/mTOR pathway in PC3 PCa cells (Xue <i>et al.</i> 2016a)		Low expression correlated with poor prognosis in thyroid cancer tissues (Guo <i>et al.</i> 2017); low expression in breast tumors from trastuzumab-treated patient (Li <i>et al.</i> 2016b) Lower in PCa tissues than normal prostate (Xue <i>et al.</i> 2016a)
<i>MEG3</i>	Maternally expressed 3 14q32.2 81,622 nt Tumor suppressor	Binds miR-421 (Zhang <i>et al.</i> 2017c,e) Co-immunoprecipitates with PI3K in HEC-1B EC cells (Sun <i>et al.</i> 2017a)		Downregulated in breast (Zhang <i>et al.</i> 2016a,c, 2017c,e), prostate (Zhang <i>et al.</i> 2016a,c), endometrial (Guo <i>et al.</i> 2016b, Sun <i>et al.</i> 2017a), and PTC (Murugan <i>et al.</i> 2017) tumors

activity independent of ligand (Xue *et al.* 2016a). ER α interacts directly of *HOTAIR* in RNA pulldown assays using nuclear extracts from MCF-7 BCa cells (Xue *et al.* 2017). Overexpression of *HOTAIR* in MCF-7 cells grown under hormone-free (serum-starved) medium conditions increases the number of DNA sites to which ER α binds in chromatin immunoprecipitation assays (ChIP) and increases mRNA expression of some ER α target genes, e.g., *GREB1*, *TFF1*, *PGR* and *CTSD* (Xue *et al.* 2016b). The authors concluded that *HOTAIR* increases ligand-independent ER α transcription. This observation is concordant with a report of increased *HOTAIR* immunostaining in tamoxifen-resistant human breast tumors (Xue *et al.* 2016b).

In contrast, *HOTAIR* was not among the 20 lncRNAs identified as upregulated and 9 downregulated by 'apo-ER α ' (non-ligand-occupied ER α) in a bioinformatic analysis of BCa cell lines and tumor tissues with experimental follow-up in MCF-7 cells (Miano *et al.* 2016). That study identified *DSCAM-AS1* as the most apo-ER α -upregulated lncRNA and its expression was higher in luminal A and B breast tumors compared to normal breast, HER2+ or basal-like tumors. Knockdown of *DSCAM-AS1* stimulated an increase in epithelial-to-mesenchymal (EMT) markers in MCF-7 cells. *DSCAM-AS1* was also upregulated in prostate and lung tumors (Miano *et al.* 2016). A bioinformatics interrogation of 947 breast tumor RNA-seq libraries identified gene sets positively correlated with *DSCAM-AS1* expression as significantly associated with clinical signatures of cancer aggression, tamoxifen resistance, higher grade, stage and metastasis (Niknafs *et al.* 2016).

HOTAIR interacts with ER β in a complex with eNOS on gene promoters to stimulate transcription in LNCaP PCa cells and a primary prostate tumor-derived cell line (called C27IM) and with ER α /eNOS in MCF-7 cells (Aiello *et al.* 2016). A potential concern with regard to the conclusion of the ChIPping of ER β is the notorious lack of specificity of ER β antibodies (Andersson *et al.* 2017, Nelson *et al.* 2017). *HOTAIR* is upregulated in endometrial cancer (EC) (Smolle *et al.* 2015) and contributes to cisplatin resistance by inhibiting autophagy (Sun *et al.* 2017b). *HOTAIR* is highly expressed in ovarian cancer (Luo *et al.* 2017); endometrial, ovarian and cervical cancers (Li *et al.* 2017a) and thyroid cancers (Zhang *et al.* 2017c,e), including PTC (Zhu *et al.* 2016).

MALAT1 (metastasis-associated lung adenocarcinoma transcript 1, 8755 nt) is another well-studied lncRNAs with roles in endocrine-related cancers (reviewed in Zhang *et al.* 2017d). *MALAT1* is expressed in almost all human tissues and is conserved across 33 mammalian species

(Jarroux *et al.* 2017). *MALAT1* lacks a poly A tail, but has a genomically encoded poly A tract and is processed into a 6.7kb nuclear form and a smaller *MALAT1*-associated small cytoplasmic RNA (mascRNA) (Zhang *et al.* 2017d). *MALAT1* is m⁶A modified on 4 of 7 sites located in hairpins of *MALAT1* within HeLa, HEK-293T and MDA-MB-231 cells (Liu *et al.* 2013). Others reported that m⁶A on position 2577 of *MALAT1* alter its secondary structure, enhancing its interaction with the m⁶A reader heterogeneous nuclear ribonucleoprotein C (HNRNPC) (Zhou *et al.* 2016). The DROSHA-DGCR8 microprocessor associates with the 5' end of *MALAT1*. *MALAT1* acts as a sponge for nuclear miR-9 and cytoplasmic miR-133, miR-200s and miR-205, although the exact mechanisms of how *MALAT1* gets to the cytoplasm are unclear (Zhang *et al.* 2017d). In turn, *MALAT1* is post-transcriptionally downregulated by miR-101, miR-125b and miR-217. The crystal structure of *MALAT1* revealed it to be a triple helix RNA that is stable in cells, e.g., $t_{1/2}$ of 7h in HeLa cells (Brown *et al.* 2014). Functionally, nuclear speckle-associated *MALAT1* is involved in alternative splicing by acting as a scaffold (Tripathi *et al.* 2010). *MALAT1* is considered oncogenic and it upregulates the Wnt/ β -catenin pathway (Ong *et al.* 2017). Capture hybridization analysis of RNA targets (CHART) identified *MALAT1* genomic-binding sites to be at actively transcribed loci (reviewed in Zhang *et al.* 2017d). However, *Malat1*-knockout (ko) mice showed no obvious phenotype, were viable and fertile and showed correct localization of nuclear speckles (Nakagawa *et al.* 2012). Expression of the lncRNA *Neat1*, which is located 30 kb upstream (5') to *Malat1*, was lower in *Malat1* ko mice, suggesting a role for *Malat1* as a *cis*-regulator of *Neat1*. RNA-seq studies in adult *Malat1* ko mice revealed upregulation of a small number of adjacent genes, also suggesting *cis* regulation (Zhang *et al.* 2012). One explanation for the lack of concordance between cellular and whole mouse studies is that *Malat1* may only function under stress conditions (reviewed in Zhang *et al.* 2017d). Increased *MALAT1* is found in many cancers including breast tumors (Ellis *et al.* 2012), castration-resistant prostate (Ren *et al.* 2013, Wang *et al.* 2014, Sowalsky *et al.* 2015), EEC (Zhao *et al.* 2014), recurrent ovarian cancer (Yang *et al.* 2017b) and thyroid tumors (Zhang *et al.* 2017a,b). Studies suggest that increased *MALAT1* plays a role in metastasis (reviewed in Zhang *et al.* 2017d). *MALAT1* mutations are frequent in breast tumors (Arun *et al.* 2016, Nik-Zainal *et al.* 2016). *MALAT1* expression was higher in thyroid cancer tissues and in thyroid cancer cells than normal thyroid (Huang *et al.* 2016a). Likewise, *NEAT1* is upregulated in breast (Qian *et al.*

2017), endometrial (Wang *et al.* 2017a,c), ovarian (Chen *et al.* 2016) and thyroid tumors (Li *et al.* 2017b) and is the highest lncRNA in prostate cancer and strongly associated with metastasis (Chakravarty *et al.* 2014).

lincRNA-p21 (gene name *TP53COR1*) is a transcript of ~3kb that is increased by p53 in response to DNA damage and serves as a transcriptional repressor in the p53 pathway by directly interacting with HNRNPK (hnRNP-K), increasing p21 transcription and inhibiting the transcription of pro-survival genes, thus triggering apoptosis (Barsotti & Prives 2010, Dimitrova *et al.* 2014). lincRNA-p21 is increased by hypoxia via HIF-1 α and interacts with and stabilizes HIF-1 α , resulting in increased glycolysis in HeLa cells (Yang *et al.* 2014). lincRNA-p21 is degraded by let-7 (Yoon *et al.* 2014). lincRNA-p21 is transcribed into two isoforms with Alu inverted repeat elements and colocalizes with the lncRNA *NEAT1* in the nucleus of U2OS, HCT-116 and MCF-7 cells (Chillón & Pyle 2016). lincRNA-p21 represses the translation of specific target genes, including *CTNNB1* (β -catenin), through RISC complex activation. Knockdown of heat-shock factor 1 (HSF1) in MDA-MB-231 cells increased lincRNA-p21 expression and the authors identified miR-320a, b and c as dependent on HSF1 that may target lincRNA-p21 through its 5' sequence, although this was not demonstrated (Chou *et al.* 2015). lincRNA-21 is downregulated in PCa and acts as a tumor suppressor in PCa cells (Wang *et al.* 2017a,c). Overexpression of lincRNA-p21 in DU145 and LNCaP PCa cells decreased pyruvate kinase M2 (*PKM2*) expression in a PTEN/AKT/mTOR cascade-dependent manner and transfection with shlincRNAP21 reduced glucose consumption (Wang *et al.* 2017b,d). There are no reports on lincRNA-21/*TP53COR1* in either endometrial or thyroid cancer.

lncRNAs in breast tumors

A recent review of the function of selected lncRNAs upregulated (19) and downregulated (5) in BCa noted that while thousands of dysregulated lncRNAs have been identified, about most relatively little is known (Warburton & Boone 2017). Bioinformatic reanalysis of RNA-seq data from 658 invasive ductal carcinomas from TCGA and sorted by PAM50 assay transcriptomic data identified 1623 lncRNAs expressed at FPKM (fragments per kilobase of non-overlapped exon per million fragments mapped) >1 in 10% of the tumors that the authors postulated are relevant in BCa (Su *et al.* 2014). lncRNA-based unsupervised hierarchical consensus clustering revealed four subgroups that correlated with mRNA transcriptome

PAM50 classification: basal-like, HER2-enriched and luminal A and B. Clustering suggested that *HOTAIR* was higher in HER2-enriched breast tumors. *ACC005152.3*, *RP11-84E24.2* and *HOTAIRM1* were higher in basal-like breast tumors, whereas *RP11-53019.2* and *RP11-473L15.3* were higher in luminal-type tumors. Su and coworkers correlated increases in lncRNAs with increased adjacent gene expression: *HOTAIR* with *HOXC11*; *HOTAIRM1* with *HOXA1*; *RP11-53019.2* and *RP11-473L15.3* with *MRPS30* and *ACC005152.3* and *RP11-84E24.2* with *SOX9* (Su *et al.* 2014). *HOTAIRM1* *SOX9* is a stem cell factor involved in tamoxifen-resistant BCa (Jeselsohn *et al.* 2017). *HOTAIR* and *BCL2* expression were correlated in breast tumors and *HOTAIR* was shown to act as a ceRNA by base-pairing with miR-206, thus relieving miR-206's repression of *BCL2* expression (Ding *et al.* 2017). *HOTAIRM1* expression increased in recurrent ovarian cancer (Yang *et al.* 2017b), but acts as a tumor suppressor in colorectal cancer (Wan *et al.* 2016).

Another group performed a similar analysis on RNA-seq data in the TCGA using ~1000 breast tumors, an independent RNA-seq dataset of 50 pairs of matched tumors and adjacent normal tissues from BCa patients, and 23 normal breast tissues from healthy women (Zhang *et al.* 2016b,d). They identified 2171 lncRNAs expressed in at least one sample, often at low levels, and ultimately identified 83 differentially expressed lncRNAs of which 60 were downregulated in breast tumor samples. Among the upregulated lncRNAs, *GATA-AS1* was associated with higher expression of its adjacent protein-coding gene *GATA3* and both were associated with ER α + breast cancer. *GATA3* regulates the transcription of genes involved in maintaining luminal mammary gland differentiation (Kouros-Mehr *et al.* 2006) and acts as a licensing factor for ER α -chromatin binding (Theodorou *et al.* 2013). *GATA-AS1* may be unique in BCa since no reports were found for the other endocrine cancers discussed in this review. One upregulated lncRNA *RP5-1198O20* was associated with lower survival rates in BCa patients (Zhang *et al.* 2016b,d). How *RP5-1198O20* contributes to BCa mortality is currently unknown.

ANRIL was identified as the lncRNA with the highest expression in SK-BR3 HER2+ BCa cells vs MCF-10A cells in a lncRNA array qPCR study (Lee *et al.* 2017). *ANRIL* expression was also higher in MCF-7 and T47D cells compared with MCF-10A cells. Some lncRNAs are associated with specific metastasis, e.g., lnc-BM (*OR5BMIP*) for BCa brain metastasis by interacting with JAK2 to activate a STAT3-ICAM1 axis that increases cell adhesion to brain capillaries and extravasation (Wang *et al.* 2017b,d).

XIST is best known for X chromosome inactivation (da Rocha & Heard 2017). However, *XIST* interacts on a genome-wide scale with a variety of proteins, including the scaffold attachment factor A (SAF-A), in the non-chromatin nuclear scaffold/matrix (Creamer & Lawrence 2017). *XIST* is downregulated in breast tumors, but its molecular mechanisms as a tumor suppressor are unclear (Huang *et al.* 2016b). One idea is that loss of *XIST* induces reactivation of X-linked genes that contribute to BCa progression (Chaligne *et al.* 2015). However, *XIST*'s role as a tumor suppressor may depend on the type of breast tumor. Patients with high *XIST* mRNA and low protein expression of 53BP1 in BRCA1-like breast tumors showed no benefit from high-dose alkylating chemotherapy and lower disease-free and overall survival (Schouten *et al.* 2016). *XIST* expression was also decreased in uterine papillary serous carcinoma accompanied by widespread X chromosome demethylation (Zhang *et al.* 2014). There are no apparent functional studies of *XIST* in EC cells.

New lncRNAs continue to be discovered. A recent example is *ESRP2-AS* that was first identified in a methylated lncRNA screen in a transgenic mouse model of human BCa (C3(1) SV40Tag) and shown to be higher in human breast tumors compared to normal breast tissue (Heilmann *et al.* 2017). Additional lncRNAs identified in BCa are included in Table 2.

lncRNAs in prostate tumors

Most lncRNAs identified in PCa are upregulated, thus acting as oncogenes (reviewed in Zhang *et al.* 2016a,c). Upregulated lncRNAs include *HOTAIR*, *PCGEM1*, *PRNCR1*, *ANRIL*, *PCAT1*, *MALAT1* and specific for PCa *PCA3*. Downregulated lncRNAs in PCa include *GASS*, *PTENP1* and *MEG3*. Refer Table 2 for details on *PCA3*, *GASS* and *MEG3*. *MEG3* modulates the activity of TGF- β genes by binding to distal regulatory elements forming an RNA–DNA triplex and interacts with the PRC2 complex (Mondal *et al.* 2015). *PCAT1* is the most highly upregulated lncRNA in PCa (Alahari *et al.* 2016). Mechanistically *PCAT1* post-transcriptionally represses tumor suppressor gene *BRCA2*, thus impairing homologous recombination DNA repair. Overexpression of *PCAT1* in DU145 PCa cells increased proliferation, migration, and invasion and post-transcriptionally upregulated *MYC* by interacting with miR-3667-3p (Prensner *et al.* 2014). Another study showed interaction between *PCAT1* and miR-145-5p that resulted in upregulation of *FSCN1*, an actin-binding protein involved in invasion and migration (Xu *et al.* 2017). Overall, *PCAT1* has an oncogenic role

in PCa (Alahari *et al.* 2016). Interestingly, a PCa risk SNP (rs7463708) promotes transformation via upregulation of *PCAT1* lncRNA (Guo *et al.* 2016a). *PCAT1* is also upregulated in hepatocellular carcinoma (HCC) (Zhang *et al.* 2017a,b). Additional lncRNAs in PCa are included in Table 2.

lncRNAs in EC

The death rate for cancer of the uterine corpus, which is associated with obesity, is increasing in the U.S. (Siegel *et al.* 2017). EC is classified as type I or II with most patients having well-differentiated endometrioid endometrial carcinomas (EEC, type I) (Takenaka *et al.* 2016). The identity and characteristics of lncRNAs dysregulated in EEC, including *MALAT1*, *HOTAIR*, *SRA*, *H19*, *OVAAL*, *RP11395G12.3*, *LA16313D11.11*, *CASC2*, *ASLNC04080* and *ENST0000050294* were reviewed (Takenaka *et al.* 2016). The roles and functions of selected lncRNAs, i.e., *MALAT1*, *HOTAIR*, *H19*, *MEG2*, *CCA2*, *ANRIL*, *OCAL*, *BC2200* and *CUDR* in cancers of the female reproductive system including endometrial, ovarian and cervical cancers were recently reviewed (Hosseini *et al.* 2017). *H19* is a well-studied imprinted, maternally expressed lncRNA that encodes miR-675 and is overexpressed in breast, HCC, prostate, colorectal, esophageal, bladder and many other cancers but is downregulated in adrenocortical neoplasms (reviewed in Huarte 2015).

lncRNAs in thyroid cancer

The role of lncRNAs in the pathogenesis of thyroid cancer was recently reviewed (Murugan *et al.* 2017). PTC is the most common thyroid malignancy (Siegel *et al.* 2017). A microarray analysis of lncRNAs in 62 PTC tumors identified 3499 lncRNAs (1192 upregulated, 2307 downregulated) that were differentially expressed compared with paired noncancerous tissue (Lan *et al.* 2015). qPCR confirmed the 'direction' of lncRNA expression in PTC: TCONS_12_00010365, n386477, n340790, lnc-LLPH-2:1 and NR_003225.2 were upregulated and lnc-PSD4-1:14, n335550, lnc-KCMF1-2:1, lnc-PLA2R1-1:1 and ENST00000422494.1 were downregulated (Lan *et al.* 2015). As indicated in Table 2, *MALAT1* is upregulated in thyroid tumors (Zhang *et al.* 2017a,b). *MALAT1* is also upregulated in SW1736, KAT18 and FTC133 thyroid cancer cells and knockdown of *MALAT1* inhibited cell proliferation and invasion by upregulating the expression of the scaffolding protein IQGAP1 (Huang *et al.* 2016a).

Mitochondrial-encoded lncRNAs and lncRNA imported into mitochondria

The human mitochondrial DNA (mtDNA) genome is transcribed as a polycistronic precursor and encodes 13 proteins that function in mt OXPHOS, two rRNAs that are components of the mt ribosome and 22 tRNA (Kim *et al.* 2017b). Seven lncRNAs have been identified to be derived from mtDNA (De Paepe *et al.* 2017). These are referred to as mitolncRNAs (Vendramin *et al.* 2017). Several are chimeric lncRNAs containing nucleotides of mtDNA: *LIPCAR*, *SncmtRNA*, *ASncmtRNA-1* and *ASncmtRNA-2*. *LIPCAR* is cardiac specific and levels are elevated in circulation of patients with chronic heart failure (Kumarswamy *et al.* 2014). *ASncmtRNA-1* and *ASncmtRNA-2* are present in mitochondria and nuclei, suggesting a possible role in retrograde signaling (De Paepe *et al.* 2017). *ASncmtRNA-1* and *ASncmtRNA-2* are downregulated in breast and prostate tumors (Burzio *et al.* 2009), suggesting tumor suppressor activity. However, complete knockdown of *ASncmtRNA-1* and *ASncmtRNA-2* stimulated apoptotic cell death by downregulating the translation of survivin, an anti-apoptotic protein, only in cancer cells through an undefined mechanism, although these lncRNAs were shown to interact with DICER (Vidaurre *et al.* 2014). To date, the mechanism of *ASncmtRNA-1* and *ASncmtRNA-2* remains unknown (De Paepe *et al.* 2017, Dong *et al.* 2017). Three lncRNAs: ncND5, lncND6 and lncCyt b were identified as antisense transcripts of the mtND5, mtND6 and mtCYTB mRNAs but little is known about their function (Rackham *et al.* 2011). There is speculation that these three lncRNAs are transported to the nucleus by unknown RNA-binding proteins (Dong *et al.* 2017). Recently, two novel lncRNAs *MDL1* and *MDLIAS* encoded in the D-loop of human mitochondrial DNA were discovered and identical transcripts were identified in mouse and rat genomes (Gao *et al.* 2017).

RMRP (RNA Component Of Mitochondrial RNA-Processing Endoribonuclease) is a nuclear DNA-encoded lncRNA that binds RNA-binding proteins HUR, PNPASE and GRSF1 for transport into mitochondria where *RMRP* is the RNA component of the mitochondrial RNA-processing endoribonuclease (RNase MRP) for mtDNA replication and RNA processing (Dong *et al.* 2017). *RMRP* acts as a sponge for miR-206 an oncogene in lung cancer (Meng *et al.* 2016). Deep sequencing of 360 primary breast tumors identified mutations in the promoters of lncRNAs *RMRP* and *NEAT1* that increased their expression and noted increased expression of these lncRNAs in breast tumors (Rheinbay *et al.* 2017). However, *RMRP*'s role in

BCa remains to be fully elucidated, and no publications were found with respect to *RMRP* in prostate, endometrial, ovarian or thyroid cancer.

SRA1, steroid receptor RNA activator, was first identified as an RNA transcript acting as an RNA coactivator in a complex with SRC-1 (*NCOA1*) to increase progesterone, glucocorticoid, androgen, estrogen, thyroid hormone, retinoic acid and peroxisome proliferator-activated receptor (PR, GR, AR, ER α , TR β , RAR γ , RXR γ and PPAR γ) transcriptional activity (Lanz *et al.* 1999). Knockout of *Sra1* in mice showed that the *Sra*^{-/-} mice were resistant to high-fat diet-induced obesity and had lower fat mass (Liu *et al.* 2014). In humans, inactivating *SRA1* mutations were identified in three families with a proband with idiopathic hypogonadotropic hypogonadism (Kotan *et al.* 2016). The *SRA1* gene also encodes a protein steroid receptor coactivator protein (SRAP) (reviewed in Leygue 2007) that interacts with AR and is involved in PCa (Kawashima *et al.* 2003). Notably, *SRA1* and SRAP do not interact (McKay *et al.* 2014). *SRA1*'s identity as a lncRNA was revealed in 2012 (Novikova *et al.* 2012). Additional protein-binding partners of *SRA1* were recently reviewed (Liu *et al.* 2016). *SRA1* interacts with DICER (Redfern *et al.* 2013). By its interaction with SLIRP (SRA stem-loop interacting RNA-binding protein), *SRA1* has a repressive function by recruiting corepressors (reviewed in Liu *et al.* 2016). SLIRP represses NR transactivation in an *SRA1*-dependent manner (Hatchell *et al.* 2006). Interestingly, *Slirp*-knockout mice are sub-fertile, and males have defects in sperm motility and mitochondrial morphology (Colley *et al.* 2013). *SRA1* and SLIRP were identified as a BCL2-interacting RNA and proteins, respectively in mitochondria of H1299 lung adenocarcinoma cells (Trisciuglio *et al.* 2016). The SMRT/HDAC1 corepressor complex interacts with *SRA1* (Dong *et al.* 2017). *SRA1* is upregulated in steroid hormone-responsive tumors including breast, endometrial and ovarian (reviewed in Liu *et al.* 2016).

lncRNAs in circulation

Circulating nucleic acids, including miRNAs and lncRNAs, are found in cell-free serum, plasma and other bodily fluids and differ between cancer patients, including those with prostate and breast tumors and non-cancer/normal individuals (reviewed in Qi *et al.* 2016). In addition, virtually all cells release membrane-enclosed extracellular vesicles (EV): exosomes (0.04–0.15 μ m diameter) and microvesicles (0.2–1 μ m diameter); further, apoptotic

cells release apoptotic bodies (0.5–2 µm diameter) that do not contain miRNAs and lncRNAs (Kim *et al.* 2017a). Exosomes, which arise from the endosomal pathway, and microvesicles have been reported to transport mRNAs, ncRNAs, including miRNAs, lncRNAs and circRNAs, as a mechanism of intercellular communication (Kim *et al.* 2017a). Cancer exosomes modulate the immune response and the tumor environment locally and at a distance, i.e., metastasis to secondary sites, by horizontal cargo transfer to recipient cells (Ruivo *et al.* 2017). In a recent study, isolated exosomes from tamoxifen-resistant LCC2 BCa cells were shown to contain ~25-fold higher lncRNA *UCA1* compared to parental MCF-7 cells and incubation of MCF-7 cells with exosomes from LCC2 cells resulted in decreased growth inhibition by tamoxifen (Xu *et al.* 2016), although no uptake of *UCA1* or other lncRNAs or miRNAs was examined. There is great interest in EVs as biomarkers and as tools to deliver therapeutic genetic materials and drugs. Interestingly, a recent study reported that EVs isolated from BCa patients with metastatic disease and whose initial tumor had been ER α + /PR+ /HER2- contained mtRNA from cancer-associated fibroblasts (CAFs) (Sansone *et al.* 2017). Xenograft transplantation studies in immunocompromised mice demonstrated that transfer of mtDNA from CAFs contributed to resistance to fulvestrant *in vivo*. Whether accompanying miRNAs and lncRNAs play a role in these findings was not evaluated.

lncRNA therapeutics

Antisense oligonucleotides (ASO), liposome-delivered RNAi, vector-expressed shRNAs and decoy RNAs or small molecules that compete for domain–domain interactions between lncRNA and proteins or within the lncRNA are of interest in targeting overexpressed lncRNAs in cancer (Weidle *et al.* 2017). Targeting of lncRNAs using lncRNAs and nanoparticle-coated siRNAs has been successful in preclinical mouse models of BCa (Liu *et al.* 2015, Wang *et al.* 2017b,d). Approximately 25 RNAi-based therapeutics are under clinical investigation (Weidle *et al.* 2017).

Overview of miRNA biosynthesis in endocrine-related cancers

miRNAs are small (22 nucleotides), evolutionarily conserved, single-stranded, ncRNAs that regulate mRNA translation or stability by base-pairing with miRNA response elements (MREs) in the 3'UTR of the target transcript within the RNA-induced silencing complex (RISC). The chronology of miRNA discovery was recently

reviewed (Drusco & Croce 2017). Comparative genomics analysis indicated that >60% of human protein-coding genes have been under selective pressure to maintain pairing to miRNAs with >45,000 miRNA-binding sites within 3'UTRs conserved (Friedman *et al.* 2009). The human genome includes 2,588 mature miRNAs (mirBase release 21) <http://www.mirbase.org/> (Kozomara & Griffiths-Jones 2014).

The biogenesis of miRNAs has been reviewed (Ha & Kim 2014, Klinge 2015). In brief, in the canonical pathway of miRNA biogenesis, miRNAs are transcribed as primary-micro-RNAs (pri-miRNAs) by RNA pol II either as independent transcription units or are cotranscribed within introns of pre-mRNAs (~50% of miRNAs) (Saini *et al.* 2007). Most pri-miRNAs are >1 kb, thus meeting the definition of a lncRNA (Dykes & Emanuelli 2017). About half of all miRNAs are produced from non-coding transcripts, i.e., host lncRNAs. One example is *DLEU2* that encodes the miR-15a/16.1 cluster and *MIR155HG* that encodes miR-155 (reviewed in Dykes & Emanuelli 2017). Pri-miRs are capped and polyadenylated (Cai *et al.* 2004). The microprocessor complex of DROSHA (an RNase III endonuclease) and DGCR8, plus additional proteins, cleaves hairpin-loop-containing pri-miR into 60–70 nucleotide (nt) precursor (pre)-miRNAs. The efficiency of pri-miRNA processing depends on stem length (36 ± 3 nt) and the appearance of bulges in the structure (Roden *et al.* 2017). DGCR8 acts as an oncogene in prostate cancer whereas DROSHA has oncogenic or tumor suppressor activity depending on the cancer type (reviewed in Hata & Kashima 2016). Alternative pathways of miRNA biogenesis include miRtrons, which are processed by spliceosomes in a DROSHA/DGCR8-independent/DICER-dependent manner and DROSHA/DGCR8-dependent/DICER-independent yielding miR-451 (reviewed in Dugaard & Hansen 2017). The widespread reduction of miRNAs in human tumors has been attributed to dysregulation of the DROSHA/DGCR8 microprocessor complex as a result of the many pathways that regulate microprocessor components, e.g., p72 is sequestered by YAP in the Hippo pathway in a cell-density-dependent manner (Mori *et al.* 2014). RNA-binding proteins also regulate pri-miRNA accessibility (Fernandez *et al.* 2017).

Pre-miRNAs are exported from the nucleus by Exportin and Ran-GTP or CRM1 to the cytoplasm where they are cleaved by DICER (*DICER1*, an RNase III enzyme) forming ~21–23 nt transiently double-stranded miRNA duplexes (Kurzynska-Kokorniak *et al.* 2015). DICER, in association with cofactors TRBP and PACT, transfer the mature miRNA to RISC, which contains the catalytic argonaute proteins

(AGO1, AGO2, AGO3 and AGO4) (Hock & Meister 2008)) that unwind the miRNA duplexes to form single-stranded miRNAs. The lncRNA *SRA1* binds DICER complex components PACT, TRBP and PKR in various cell lines and also binds NRs, including ER α (Redfern *et al.* 2013). DICER also plays a role in the DNA damage response in which DNA damage results in phosphorylation and binding to double-strand breaks and recruits DNA repair factor MDC1 and 53BP1 (Burger *et al.* 2017). DICER acts as an oncogene in PCa but as a tumor suppressor in breast, endometrial and ovarian cancer (reviewed in Hata & Kashima 2016). DICER is regulated transcriptionally, post-transcriptionally, including by alternative splicing, and by post-translational modifications, e.g. phosphorylation, glycosylation and SUMOylation (Kurzynska-Kokorniak *et al.* 2015). Germline *DICER1* mutations, *DICER1* syndrome, were identified as causative in differentiated thyroid cancer and multinodular goiter (de Kock *et al.* 2014, Rutter *et al.* 2016, Khan *et al.* 2017).

In cancer tissues, overexpressed miRNAs that act as oncogenes are referred to as oncomiRs while downregulated miRNAs are called tumor suppressor miRNAs with their targets, tumor suppressors and oncogenes, respectively dysregulated. However, as we learn more about miRNAs, it is clear that generalization about a particular miRNA being oncogenic or a tumor suppressor depends on cellular context. miRNAs are involved in the post-transcriptional regulation of genes involved in EMT, stemness, cell signaling pathways and regulation of the tumor microenvironment, including the conversion of normal fibroblasts into cancer-associated fibroblasts (CAFs) by transfer of exosomal miRNAs from tumor cells (reviewed in Yang *et al.* 2017a). An issue in the field is the lack of congruence of miRNAs identified for a particular cancer type, e.g., thyroid cancer, even within a subtype, in different studies. This may be attributed to the number of tumor samples examined, tumor heterogeneity, whole tumor RNA isolation vs laser-capture microdissected tumor cells, the technique used to identify the miRNA (microarray, quantitate real-time PCR (qPCR), or NGS/RNA-seq.), the gene normalizer and statistical approaches used. Databases of putative and experimentally verified targets of miRNAs are listed in Table 1.

Despite the large datasets available on miRNAs in endocrine cancer, functional studies are needed for each miRNA and its targets. Functional analysis of overexpression and silencing of individual miRNAs is used to validate the role of deregulated miRNAs in cancer cells *in vitro* and animal tumor models, including human tumor cell xenografts and patient-derived xenograft

(PDX) *in vivo*. An essential step to validate the identify of an mRNA target of a miRNA is to clone the 3'-UTR of the mRNA target downstream of a luciferase reporter and then transfect that reporter into a cell line, with an appropriate loading control either in the same plasmid or another cotransfected reporter, with overexpression and knockdown of the putative miRNA regulator. The investigator should detect reduced luciferase reporter activity and mutation of the MRE in the 3'UTR is required confirm direct miRNA-mRNA target regulation, i.e., validating that the mRNA is a *bona fide* target of that miRNA. Further validation steps include Western blot to demonstrate decrease target protein expression. Target validation in clinical samples is used to demonstrate disease relevance of the miRNA and its target.

Supplementary Table 1 (see section on supplementary data given at the end of this article) summarizes examples of miRNA dysregulated in BCa, PCa, EC and thyroid cancer with examples of their *bona fide*, validated, mRNA targets. Cellular functions associated with the miRNA are listed in Supplementary Table 1. In addition, I have added lncRNAs that either downregulate a miRNA or act as ceRNAs, i.e., endogenous sponge for binding a miRNA. It will be clear to the reader that many of the miRNA:lncRNA interactions listed are from non-endocrine cancers and are single reports requiring further validation.

miRNAs in BCa

In the 12 years since miRNAs were identified to be dysregulated in BCa (Iorio *et al.* 2005), about 3600 papers have been published in miRNAs in BCa (Fig. 1). There are many recent reviews of the identity, regulation and targets of miRNAs in BCa, e.g., (Egeland *et al.* 2015, Klinge 2015, Muluhngwi & Klinge 2015, van Schooneveld *et al.* 2015, Yahya & Elsayed 2015, Li *et al.* 2017c, O'Bryan *et al.* 2017, Smith *et al.* 2017). miRNAs regulate key pathways dysregulated in BCa including apoptosis, cell cycle, cellular energetics, invasion, metabolism and metastasis.

The majority (70%) of primary breast tumors express ER α , and these patients are treated with endocrine therapies including tamoxifen (TAM); the selective estrogen downregulator (SERD), e.g., fulvestrant and aromatase inhibitors (AI), e.g., anastrozole and letrozole (Ring & Dowsett 2004, Clarke *et al.* 2015). Hence, many miRNA studies have examined the correlation of miRNAs with diagnostic markers related to ER α in BCa, estrogen-regulated miRNAs and the potential role of miRNAs in endocrine resistance (Klinge 2015, Muluhngwi & Klinge

2015). ER α is the target of miRNAs including miR-221, miR-222, miR-873, let-7b and let-7i (reviewed in [Muluhngwi & Klinge 2015](#)).

Computational approaches are used to identify putative mRNA and lncRNA targets of miRNAs ([Table 2](#)). In addition, specific algorithms to identify possible targets of dysregulated miRNAs in breast tumors and BCa cell lines have been reported ([O'Day & Lal 2010](#), [Cava et al. 2014](#), [Ru et al. 2015](#), [Liang et al. 2016](#), [Zhang et al. 2016b,d](#)). More recently, various strategies to directly identify miRNA:RNA targets and RNA modifications by protein:RNA immunoprecipitation and high-throughput RNA-seq have been used to examine protein:RNA interactions in human biology, e.g., HITS-CLIP (high-throughput RNA-seq isolated by crosslinking immunoprecipitation) ([Darnell 2010](#)); PAR-CLIP (photoactivatable ribonucleoside-enhanced CLIP) ([Farazi et al. 2014](#)), iCLIP (crosslinking and immunoprecipitation) ([Hong et al. 2015](#)), eCLIP (enhanced CLIP) ([Van Nostrand et al. 2016](#)) and dCLIP (denaturing CLIP) ([Rosenberg et al. 2017](#)). A downside is that CLIP techniques are technically challenging.

Dysregulation of miRNAs in PCa

Dysregulation of miRNA expression is correlated with aggressive PCa phenotypes including tumor stage, Gleason grade and disease recurrence ([Ren et al. 2014](#), [Cha et al. 2016](#)). Altered miRNA expression is associated with physiological changes in tumorigenesis and disease progression in PCa (reviewed in [Chun-Jiao et al. 2017](#), [Vanacore et al. 2017](#)). Changes in miRNA expression contribute to altered expression of genes involved in pathogenesis of metastatic PCa ([Kojima et al. 2017](#)). Changes in miRNA expression in PCa have been reviewed and described below.

miRNAs upregulated by AR in PCa include miR-21, miR-27a, miR-32, miR-125b, miR-135a, miR-141 and AR overexpression in PCa downregulates miR-99a ([Massillo et al. 2017](#)). miR-125b is the most overexpressed miRNA in PCa and downregulates apoptotic genes including *BAK1*. A recent review of 104 studies of miRNAs in PCa tissues identified six miRNA consistently upregulated: miR-34a, miR-106b, miR-183, miR-200a/b and miR-301a; and sixteen miRNAs downregulated: miR-1, miR-23b, miR-27b, miR-34b/c, miR-99b, miR-125b, miR-152, miR-187, miR-199a, miR-204, miR-205, miR-224, miR-452, miR-454 and miR-505 ([Chun-Jiao et al. 2017](#)). Other oncomiRs overexpressed in PCa are miR-21, miR-32,

miR-221, miR-222, miR-181, miR-18a and miR-429 (reviewed in [Kanwal et al. 2017](#)). 'Andro-miRs': let-7c, miR-31, miR-124, miR-185, miR-205 and miR-488 downregulate AR expression ([Massillo et al. 2017](#)). Notably, the top four downregulated miRNAs in PCa tissues are miR-187, miR-205, miR-222 and miR-31 ([Fuse et al. 2012](#)). Additional tumor suppressor miRNAs in PCa and their targets include miR-331-3p that downregulates ERBB2 and AR and is associated with castration-resistant PCa (reviewed in [Kanwal et al. 2017](#)). miRNAs associated with metastasis include downregulation of miR-1, miR-15, miR-16, miR-23a, miR-29b, miR-126, miR-130, miR-132, miR-141, miR-195, miR-200b, miR-200c, miR-203, miR-205, miR-218, miR-375, miR-377, miR-466, miR-573, miR-675, and miR-802 and upregulation of miR-21, miR-22, miR-25, miR-93, miR-106b, miR-154-3p, miR-379, miR-543 and miR-590-3p (reviewed in [Massillo et al. 2017](#)). miRNAs upregulated or downregulated in serum, plasma and urine with potential prognostic value in separating men with potentially lethal PCa vs indolent disease were recently reviewed ([Massillo et al. 2017](#)).

Dysregulation of miRNAs in EC

EC is the most frequently diagnosed gynecological cancer ([Siegel et al. 2017](#)). Among the miRNAs differentially expressed in EC vs normal endometrial tissue are the increased expression of miR-10b, miR-21, miR-31, miR-182, miR-183, miR-205, miR-222, miR-223, miR-410, miR-429, miR-449a, miR-994, and miR-1228; and downregulation of let-7, miR-34b-5p, miR-34c-3p, miR-34c-5p, miR-99b, miR-101, miR-130a, miR-143, miR-145, miR-184, miR-193b, miR-204, miR-340, miR-372, miR-429, miR-449a, miR-490 and miR-495 (reviewed in [Srivastava et al. 2017](#)). In serum samples from women with EC, increased levels of miR-186, miR-222 and miR-223 were identified compared with controls ([Montagnana et al. 2017](#)). Targets of upregulated miRs in EC include the tumor suppressor PTEN that is downregulated by overexpressed miR-21, miR-130a, miR-205, miR-222 and miR-429 ([Srivastava et al. 2017](#)). Although much studied, there are currently no miRNA signatures in use for early detection and screening of gynecological cancers ([Srivastava et al. 2017](#)).

miRNAs in thyroid cancer

The identity and targets of miRNAs dysregulated in PTC and follicular thyroid carcinoma (FTC) have been recently

reviewed (Pallante *et al.* 2013, Saiselet *et al.* 2016, Yoo *et al.* 2016, Celano *et al.* 2017, Lima *et al.* 2017, Pishkari *et al.* 2018). miRNAs upregulated in PTC include miR-21, miR-31, miR-99-3p, miR-128a, miR-128b, miR-139, miR-141, miR-146a, miR-146b-3p, miR-146b-5p, miR-155, miR-181a, miR-181b, miR-187, miR-191, miR-200a, miR-200b, miR-200c, miR-220, miR-221, miR-222, miR-222-5p, miR-224, miR-375, miR-551b (Pallante *et al.* 2013). Downregulated miRNAs in PTC include: let-7f, miR-1, miR-26a-1, miR-30, miR-30c, miR-138, miR-199, miR-219, miR-292, miR-300 and miR-345 (Pallante *et al.* 2013). With RNA-seq, the arm of each miRNA and isoforms can be detected. Thus, RNA-seq has higher resolution and detected increased miR-21-3p, miR-21-5p, miR-31-3p, miR-31-5p, miR-34a-5p, miR-146-3p, miR-146-5p, miR-182-5p, miR-183-5p, miR-221-3p, miR-221-5p, miR-222-3p and miR-551b-3p in PTC (Saiselet *et al.* 2016). RNA-seq identified miRNAs upregulated in FTC include miR-96-5p, miR-182-5p, miR-221-3p, miR-183-5p and miR-222-3p, whereas miR-31-5p, miR-199a-5p and miR-199b-5p showed consistent downregulation in three independent studies (reviewed in Saiselet *et al.* 2016).

miRNAs in mitochondria

Mitochondrial function is critical for metabolic homeostasis. NGS has identified several miRNAs imported into mitochondria including miR-34, miR-181c-5p and miR-146a-5p (Kim *et al.* 2017b). The identity and roles of miRNAs in mitochondrial biology were recently reviewed (Vendramin *et al.* 2017). The term 'mitomiRs' refers to miRNAs functioning in mitochondria, whether nuclear- or mtDNA-encoded (Duarte *et al.* 2015). mitomiRs are 17–25 nt vs the canonical 22 nt for most miRNAs. The exact function and identity of mitomiRs remains unknown and caution is advised since cytoplasmic contamination of mitochondrial extracts is problematic (Vendramin *et al.* 2017). Many ncRNAs appear to be involved in rapid cell stress responses and may be involved in anterograde and retrograde signaling between the nucleus and mitochondria to regulate energy homeostasis and apoptosis. For example, miRNAs appear to provide anterograde regulation of mitochondrial function, apoptosis and cancer cell metabolism (Duarte *et al.* 2015, Cha *et al.* 2017). However, although mitochondrial dysfunction is a hallmark of cancer, the role of ncRNAs in the mitochondrial unfolded protein response (UPR^{mt}) in cancer (reviewed in Kenny & Germain 2017) remains to be examined.

miRNAs in circulation

Screening of free and exosomal-associated miRNAs by high-throughput sequencing platforms has identified changes in breast, ovarian and prostate cancers (Smith *et al.* 2017) and in thyroid cancer (Saiselet *et al.* 2016). The release of exosomes containing miRNAs (as well as other ncRNAs, protein and lipids) and endocytic uptake of the exosomes in recipient cells results in altered regulation of gene translation and has implicated miRNAs as hormones (reviewed in Bayraktar *et al.* 2017). In addition, circulating miRNAs can be transported between cells by microvesicles, apoptotic bodies, lipoproteins (HDL) and ribonucleoproteins, e.g., nucleophosmin 1 and AGO2. Exosomes derived from patients with CRPCa have increased expression of miR-1290 and miR-375 and correlate with decreased overall survival (reviewed in Kanwal *et al.* 2017). CAFs stimulate cancer progression by secreting chemokines, cytokines and growth factors that create the extracellular matrix (ECM) and CAFs secrete EVs containing a different spectrum of miRNAs compared to normal fibroblasts (Bayraktar *et al.* 2017). Among the miRNAs secreted by CAFs that regulate cancer cells are miR-21, miR-143 and miR-378 in BCa (Donnarumma *et al.* 2017); miR-210, miR-409-3p/5p (Yang *et al.* 2017a) and miR-409 (Josson *et al.* 2014) in PCa cells; and miR-21 in ovarian cancer (Au Yeung *et al.* 2016). Tumor-associated macrophages (TAMs) are also donors of miRNAs in cancer (reviewed in Bayraktar *et al.* 2017). Increased TAMs is associated with poor survival in advanced thyroid cancer (Ryder *et al.* 2008).

The mandate from NIH in the Precision Medicine Initiative is to have data-driven care for patients. A current goal of clinical trials is to determine if miRNA signatures obtained in fine needle aspiration biopsies and blood (serum or plasma) or urine will provide early diagnosis and track with therapeutic responses in cancer patients. However, the miRNA expression profile identified in blood samples depends on the processing of the sample, the miRNA extraction method, contamination, including by lysis of blood cells, and which normalizers are used (Saiselet *et al.* 2016). In addition to the ongoing efforts to standardize exosome and miRNA purification from biological fluids, factors that impact variability in miRNAs detected in circulation include diet, physical activity and circadian rhythms (reviewed in Smith *et al.* 2017). Most of the clinical trials involving mRNAs are observational, i.e., examining circulating in PCa, ovarian and breast cancers to monitor therapeutic responses to chemotherapy or other treatment strategies (reviewed in Smith *et al.* 2017).

miRNA therapeutics

Despite the large body of literature supporting miRNAs as biomarkers for diagnosis, prognosis and follow-up in patients with malignancies, the application of miRNA therapeutics into the clinical practice has a number of technical obstacles to overcome include clearance and optimization of bioavailability (reviewed in [Chen *et al.* 2015](#)). Nanoformulations of miRNAs, miRNA mimics, miRNA sponges, anti-miR oligonucleotides (AMOs) and small-molecule inhibitors are being tested in animal models. miRNA mimics to replace the expression of downregulated miRNAs by synthetic oligonucleotides that are combined with hydrogels, liposomes, minicells, nanoparticles, synthetic polymers or viral carriers for better delivery are under investigation ([Drusco & Croce 2017](#)). Recently, a folate-miR-34a-5p conjugate was shown to be taken by MDA-MB-231 cells, which have high folate receptor levels and to inhibit MDA-MB-231 xenograft tumor growth in mice ([Orellana *et al.* 2017](#)). ‘AntagomiRs’ serve to inhibit oncomiR function, e.g., a miR-10b antagomiR to mice did not inhibit primary mammary tumor growth, but inhibited lung metastasis ([Ma *et al.* 2010](#)) and intraperitoneal injection of a miR-92a-LNA inhibitor inhibited human EC cell xenograft growth in mice with no evidence of overt toxicity or metastatic spread ([Torres *et al.* 2016](#)). However, many challenges remain for therapeutics with or against miRNAs, notably the contradictory nature of miRNAs.

Conclusions

lncRNAs and miRNAs dysregulated in endocrine cancers, and they serve as biomarkers and potential therapeutic targets and agents in various cancers and other diseases, e.g., cardiovascular disease and type II diabetes. These and other ncRNAs are interconnected and have complex molecular roles in regulating protein-coding and non-coding nuclear gene transcription: RNA stability, epigenetic processes, chromatin accessibility, translation, gene expression in mitochondria, anterograde and retrograde signaling and intracellular and intercellular signaling. The study of lncRNAs is less mature than that of miRNAs. For example, new lncRNAs continue to be discovered, relatively few studies on any specific lncRNA have been published, and each lncRNA may have multiple isoforms that can be post-transcriptionally modified, adding complexity to evaluating their cellular and physiologic function. lncRNA structures are functionally important, but only a few lncRNAs’ structures are known. Identification of

cell-specific lncRNA-interacting proteins and other RNAs and their subcellular distribution is needed. Both miRNAs and lncRNAs are being actively evaluated as tumor biopsy, plasma, serum, blood and urine biomarkers in the diagnosis and management of endocrine-related cancers as well as therapeutic targets. In addition, lncRNAs and miRNAs may serve as therapeutic agents. Many challenges remain in elucidating the biogenesis and function of ncRNAs in endocrine-related cancers. Our understanding of the functional roles of miRNAs and lncRNAs and their molecular targets is expanding rapidly. However, individual miRNAs and lncRNAs, and their combinations, may have different sets of targets and activities depending on the cell type, microenvironment, microbiome and hormonal milieu; thus, further investigation is required to fully elucidate miRNA and lncRNA function in endocrine-related cancers.

Supplementary data

This is linked to the online version of the paper at <https://doi.org/10.1530/ERC-17-0548>.

Declaration of interest

The author declares that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

References

- Aiello A, Bacci L, Re A, Ripoli C, Pierconti F, Pinto F, Masetti R, Grassi C, Gaetano C, Bassi PE, *et al.* 2016 MALAT1 and HOTAIR long non-coding RNAs play opposite role in estrogen-mediated transcriptional regulation in prostate cancer cells. *Science Reports* **6** 38414. (<https://doi.org/10.1038/srep38414>)
- Alahari SV, Eastlack SC & Alahari SK 2016 Role of long noncoding RNAs in neoplasia: special emphasis on prostate cancer. *International Review of Cell and Molecular Biology* **324** 229–254. (<https://doi.org/10.1016/bs.ircmb.2016.01.004>)
- Andersson R, Gebhard C, Miguel-Escalada I, Hoof I, Bornholdt J, Boyd M, Chen Y, Zhao X, Schmidl C, Suzuki T, *et al.* 2014 An atlas of active enhancers across human cell types and tissues. *Nature* **507** 455–461. (<https://doi.org/10.1038/nature12787>)
- Andersson S, Sundberg M, Pristovsek N, Ibrahim A, Jonsson P, Katona B, Clausson C-M, Zieba A, Ramström M, Söderberg O, *et al.* 2017 Insufficient antibody validation challenges oestrogen receptor beta research. *Nature Communications* **8** 15840. (<https://doi.org/10.1038/ncomms15840>)
- Arun G, Diermeier S, Akerman M, Chang KC, Wilkinson JE, Hearn S, Kim Y, MacLeod AR, Krainer AR, Norton L, *et al.* 2016 Differentiation of mammary tumors and reduction in metastasis upon Malat1

- lncRNA loss. *Genes and Development* **30** 34–51. (<https://doi.org/10.1101/gad.270959.115>)
- Au Yeung CL, Co N-N, Tsuruga T, Yeung T-L, Kwan S-Y, Leung CS, Li Y, Lu ES, Kwan K, Wong K-K, *et al.* 2016 Exosomal transfer of stroma-derived miR21 confers paclitaxel resistance in ovarian cancer cells through targeting APAF1. *Nature Communications* **7** 11150. (<https://doi.org/10.1038/ncomms11150>)
- Barrett SP & Salzman J 2016 Circular RNAs: analysis, expression and potential functions. *Development* **143** 1838. (<https://doi.org/10.1242/dev.128074>)
- Barsotti AM & Prives C 2010 Noncoding RNAs: the missing 'linc' in p53-mediated repression. *Cell* **142** 358–360. (<https://doi.org/10.1016/j.cell.2010.07.029>)
- Bayraktar R, Van Roosbroeck K & Calin GA 2017 Cell-to-cell communication: microRNAs as hormones. *Molecular Oncology* **11** 1673–1686. (<https://doi.org/10.1002/1878-0261.12144>)
- Bhan A, Hussain I, Ansari KI, Kasiri S, Bashyal A & Mandal SS 2013 Antisense transcript long noncoding RNA (lncRNA) HOTAIR is transcriptionally induced by estradiol. *Journal of Molecular Biology* **425** 3707–3722. (<https://doi.org/10.1016/j.jmb.2013.01.022>)
- Bhan A, Deb P, Shihabeddin N, Ansari KI, Brotto M & Mandal SS 2017 Histone methylase MLL1 coordinates with HIF and regulate lncRNA HOTAIR expression under hypoxia. *Gene* **629** 16–28. (<https://doi.org/10.1016/j.gene.2017.07.069>)
- Brown JA, Bulkley D, Wang J, Valenstein ML, Yario TA, Steitz TA & Steitz JA 2014 Structural insights into the stabilization of MALAT1 noncoding RNA by a bipartite triple helix. *Nature Structural and Molecular Biology* **21** 633. (<https://doi.org/10.1038/nsmb.2844>)
- Burger K, Schlackow M, Potts M, Hester S, Mohammed S & Gullerova M 2017 Nuclear phosphorylated Dicer processes double-stranded RNA in response to DNA damage. *Journal of Cell Biology* **216** 2373. (<https://doi.org/10.1083/jcb.201612131>)
- Burzio VA, Villota C, Villegas J, Landerer E, Boccardo E, Villa LL, Martínez R, Lopez C, Gaete F, Toro V, *et al.* 2009 Expression of a family of noncoding mitochondrial RNAs distinguishes normal from cancer cells. *PNAS* **106** 9430–9434. (<https://doi.org/10.1073/pnas.0903086106>)
- Cai X, Hagedorn CH & Cullen BR 2004 Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA* **10** 1957–1966. (<https://doi.org/10.1261/rna.7135204>)
- Cava C, Bertoli G, Ripamonti M, Mauri G, Zoppis I, Della Rosa PA, Gilardi MC & Castiglioni I 2014 Integration of mRNA expression profile, copy number alterations, and microRNA expression levels in breast cancer to improve grade definition. *PLoS ONE* **9** e97681. (<https://doi.org/10.1371/journal.pone.0097681>)
- Celano M, Rosignolo F, Maggiasano V, Pecce V, Iannone M, Russo D & Bulotta S 2017 MicroRNAs as biomarkers in thyroid carcinoma. *International Journal of Genomics* **2017** 6496570. (<https://doi.org/10.1155/2017/6496570>)
- Cerase A, Pintacuda G, Tattermusch A & Avner P 2015 Xist localization and function: new insights from multiple levels. *Genome Biology* **16** 166. (<https://doi.org/10.1186/s13059-015-0733-y>)
- Cha YJ, Lee JH, Han HH, Kim BG, Kang S, Choi YD & Cho NH 2016 MicroRNA alteration and putative target genes in high-grade prostatic intraepithelial neoplasia and prostate cancer: STAT3 and ZEB1 are upregulated during prostate carcinogenesis. *Prostate* **76** 937–947. (<https://doi.org/10.1002/pros.23183>)
- Cha Y, Han M-J, Cha H-J, Zoldan J, Burkart A, Jung JH, Jang Y, Kim C-H, Jeong H-C, Kim B-G, *et al.* 2017 Metabolic control of primed human pluripotent stem cell fate and function by the miR-200c-SIRT2 axis. *Nature Cell Biology* **19** 445. (<https://doi.org/10.1038/ncb3517>)
- Chakravarty D, Sboner A, Nair SS, Giannopoulou E, Li R, Hennig S, Mosquera JM, Pauwels J, Park K, Kossai M, *et al.* 2014 The oestrogen receptor alpha-regulated lncRNA NEAT1 is a critical modulator of prostate cancer. *Nature Communications* **5** 5383. (<https://doi.org/10.1038/ncomms6383>)
- Chaligne R, Popova T, Mendoza-Parra MA, Saleem MA, Gentien D, Ban K, Piolot T, Leroy O, Mariani O, Gronemeyer H, *et al.* 2015 The inactive X chromosome is epigenetically unstable and transcriptionally labile in breast cancer. *Genome Research* **25** 488–503. (<https://doi.org/10.1101/gr.185926.114>)
- Chen Y, Gao D-Y & Huang L 2015 *In vivo* delivery of miRNAs for cancer therapy: challenges and strategies. *Advanced Drug Delivery Reviews* **81** 128–141. (<https://doi.org/10.1016/j.addr.2014.05.009>)
- Chen ZJ, Zhang Z, Xie BB & Zhang HY 2016 Clinical significance of up-regulated lncRNA NEAT1 in prognosis of ovarian cancer. *European Review for Medical and Pharmacological Sciences* **20** 3373–3377.
- Chen YG, Satpathy AT & Chang HY 2017 Gene regulation in the immune system by long noncoding RNAs. *Nature Immunology* **18** 962–972. (<https://doi.org/10.1038/ni.3771>)
- Chillón I & Pyle AM 2016 Inverted repeat Alu elements in the human lincRNA-p21 adopt a conserved secondary structure that regulates RNA function. *Nucleic Acids Research* **44** 9462–9471.
- Chiyomaru T, Yamamura S, Fukuhara S, Yoshino H, Kinoshita T, Majid S, Saini S, Chang I, Tanaka Y, Enokida H, *et al.* 2013 Genistein inhibits prostate cancer cell growth by targeting miR-34a and oncogenic HOTAIR. *PLoS ONE* **8** e70372. (<https://doi.org/10.1371/journal.pone.0070372>)
- Chiyomaru T, Fukuhara S, Saini S, Majid S, Deng G, Shahryari V, Chang I, Tanaka Y, Enokida H, Nakagawa M, *et al.* 2014 Long non-coding RNA HOTAIR is targeted and regulated by miR-141 in human cancer cells. *Journal of Biological Chemistry* **289** 12550–12565. (<https://doi.org/10.1074/jbc.M113.488593>)
- Chou SD, Murshid A, Eguchi T, Gong J & Calderwood SK 2015 HSF1 regulation of [beta]-catenin in mammary cancer cells through control of HuR/elavL1 expression. *Oncogene* **34** 2178–2188. (<https://doi.org/10.1038/onc.2014.177>)
- Chun-Jiao S, Huan C, Li-Zhong C, Guo-Mei R, Jian-Jun G & Qian-Nan D 2017 The potential of microRNAs as human prostate cancer biomarkers: a meta-analysis of related studies. *Journal of Cellular Biochemistry* **119** 2763–2786. (<https://doi.org/10.1002/jcb.26445>)
- Cinghu S, Yang P, Kosak JP, Conway AE, Kumar D, Oldfield AJ, Adelman K & Jothi R 2017 Intragenic enhancers attenuate host gene expression. *Molecular Cell* **68** 104.e106–117.e106. (<https://doi.org/10.1016/j.molcel.2017.09.010>)
- Clarke R, Tyson JJ & Dixon JM 2015 Endocrine resistance in breast cancer – an overview and update. *Molecular and Cellular Endocrinology* **418** 220–234. (<https://doi.org/10.1016/j.mce.2015.09.035>)
- Colley SM, Wintle L, Searles R, Russell V, Firman RC, Smith S, DeBoer K, Merriner DJ, Genevieve B, Bentel JM, *et al.* 2013 Loss of the nuclear receptor corepressor SLIRP compromises male fertility. *PLoS ONE* **8** e70700. (<https://doi.org/10.1371/journal.pone.0070700>)
- Creamer KM & Lawrence JB 2017 XIST RNA: a window into the broader role of RNA in nuclear chromosome architecture. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences* **372** 20160360. (<https://doi.org/10.1098/rstb.2016.0360>)
- da Rocha ST & Heard E 2017 Novel players in X inactivation: insights into Xist-mediated gene silencing and chromosome conformation. *Nature Structural and Molecular Biology* **24** 197–204. (<https://doi.org/10.1038/nsmb.3370>)
- Darnell RB 2010 HITS-CLIP: panoramic views of protein–RNA regulation in living cells. *Wiley Interdisciplinary Reviews: RNA* **1** 266–286. (<https://doi.org/10.1002/wrna.31>)
- Daugaard I & Hansen TB 2017 Biogenesis and function of ago-associated RNAs. *Trends in Genetics* **33** 208–219. (<https://doi.org/10.1016/j.tig.2017.01.003>)
- de Kock L, Sabbaghian N, Soglio DB, Guillerman RP, Park BK, Chami R, Deal CL, Priest JR & Foulkes WD 2014 Exploring the association between DICER1 mutations and differentiated thyroid carcinoma.

- Journal of Clinical Endocrinology and Metabolism* **99** E1072–E1077. (<https://doi.org/10.1210/jc.2013-4206>)
- De Paepe B, Lefever S & Mestdagh P 2017 How long noncoding RNAs enforce their will on mitochondrial activity: regulation of mitochondrial respiration, reactive oxygen species production, apoptosis, and metabolic reprogramming in cancer. *Current Genetics* **64** 163–172. (<https://doi.org/10.1007/s00294-017-0744-1>)
- Delas MJ & Hannon GJ 2017 lncRNAs in development and disease: from functions to mechanisms. *Open Biology* **7** 170121. (<https://doi.org/10.1098/rsob.170121>)
- Dimitrova N, Zamudio Jesse R, Jong Robyn M, Soukup D, Resnick R, Sarma K, Ward Amanda J, Raj A, Lee Jeannie T, Sharp Phillip A, *et al.* 2014 lincRNA-p21 activates p21 in cis to promote polycomb target gene expression and to enforce the G1/S checkpoint. *Molecular Cell* **54** 777–790. (<https://doi.org/10.1016/j.molcel.2014.04.025>)
- Ding W, Ren J, Ren H & Wang D 2017 Long noncoding RNA HOTAIR modulates MiR-206-mediated Bcl-w signaling to facilitate cell proliferation in breast cancer. *Scientific Reports* **7** 17261. (<https://doi.org/10.1038/s41598-017-17492-x>)
- Djebali S, Davis CA, Merkel A, Dobin A, Lassmann T, Mortazavi A, Tanzer A, Lagarde J, Lin W, Schlesinger F, *et al.* 2012 Landscape of transcription in human cells. *Nature* **489** 101–108. (<https://doi.org/10.1038/nature11233>)
- Dong Y, Yoshitomi T, Hu JF & Cui J 2017 Long noncoding RNAs coordinate functions between mitochondria and the nucleus. *Epigenetics and Chromatin* **10** 41. (<https://doi.org/10.1186/s13072-017-0149-x>)
- Donnarumma E, Fiore D, Nappa M, Roscigno G, Adamo A, Iaboni M, Russo V, Affinito A, Puoti I, Quintavalle C, *et al.* 2017 Cancer-associated fibroblasts release exosomal microRNAs that dictate an aggressive phenotype in breast cancer. *Oncotarget* **8** 19592–19608. (<https://doi.org/10.18632/oncotarget.14752>)
- Drusco A & Croce CM 2017 Chapter one – microRNAs and cancer: a long story for short RNAs. In *Advances in Cancer Research*, pp 1–24. Eds CM Croce & PB Fisher. Cambridge, MA, USA: Academic Press.
- Duarte FV, Palmeira CM & Rolo AP 2015 The emerging role of mitomiRNAs in the pathophysiology of human disease. *Advances in Experimental Medicine and Biology* **888** 123–154. (https://doi.org/10.1007/978-3-319-22671-2_8)
- Dykes IM & Emanuelli C 2017 Transcriptional and post-transcriptional gene regulation by long non-coding RNA. *Genomics, Proteomics and Bioinformatics* **15** 177–186. (<https://doi.org/10.1016/j.gpb.2016.12.005>)
- Egeland NG, Lunde S, Jonsdottir K, Lende TH, Cronin-Fenton D, Gilje B, Janssen EA & Soiland H 2015 The role of microRNAs as predictors of response to tamoxifen treatment in breast cancer patients. *International Journal of Molecular Sciences* **16** 24243–24275. (<https://doi.org/10.3390/ijms161024243>)
- Ellis MJ, Ding L, Shen D, Luo J, Suman VJ, Wallis JW, Van Tine BA, Hoog J, Goiffon RJ, Goldstein TC, *et al.* 2012 Whole-genome analysis informs breast cancer response to aromatase inhibition. *Nature* **486** 353–360. (<https://doi.org/10.1038/nature11143>)
- ENCODE Project Consortium 2012 An integrated encyclopedia of DNA elements in the human genome. *Nature* **489** 57–74. (<https://doi.org/10.1038/nature11247>)
- Farazi TA, Ten Hoeve JJ, Brown M, Mihailovic A, Horlings HM, van de Vijver MJ, Tuschl T & Wessels LF 2014 Identification of distinct miRNA target regulation between breast cancer molecular subtypes using AGO2-PAR-CLIP and patient datasets. *Genome Biology* **15** R9. (<https://doi.org/10.1186/gb-2014-15-1-r9>)
- Fernandez N, Cordiner RA, Young RS, Hug N, Macias S & Cáceres JF 2017 Genetic variation and RNA structure regulate microRNA biogenesis. *Nature Communications* **8** 15114. (<https://doi.org/10.1038/ncomms15114>)
- Friedman RC, Farh KK, Burge CB & Bartel DP 2009 Most mammalian mRNAs are conserved targets of microRNAs. *Genome Research* **19** 92–105. (<https://doi.org/10.1101/gr.082701.108>)
- Fritah S, Niclou SP & Azuaje F 2014 Databases for lncRNAs: a comparative evaluation of emerging tools. *RNA* **20** 1655–1665. (<https://doi.org/10.1261/ma.044040.113>)
- Fuse M, Kojima S, Enokida H, Chiyomaru T, Yoshino H, Nohata N, Kinoshita T, Sakamoto S, Naya Y, Nakagawa M, *et al.* 2012 Tumor suppressive microRNAs (miR-222 and miR-31) regulate molecular pathways based on microRNA expression signature in prostate cancer. *Journal of Human Genetics* **57** 691. (<https://doi.org/10.1038/jhg.2012.95>)
- Gainetdinov I, Skvortsova Y, Kondratieva S, Funikov S & Azhikina T 2017 Two modes of targeting transposable elements by piRNA pathway in human testis. *RNA* **23** 1614–1625. (<https://doi.org/10.1261/ma.060939.117>)
- Gao S, Tian X, Chang H, Sun Y, Wu Z, Cheng Z, Dong P, Zhao Q, Ruan J & Bu W 2017 Two novel lncRNAs discovered in human mitochondrial DNA using PacBio full-length transcriptome data. *Mitochondrion* **38** 41–47. (<https://doi.org/10.1016/j.mito.2017.08.002>)
- Guo C, Song WQ, Sun P, Jin L & Dai HY 2015 lncRNA-GASS induces PTEN expression through inhibiting miR-103 in endometrial cancer cells. *Journal of Biomedical Science* **22** 100. (<https://doi.org/10.1186/s12929-015-0213-4>)
- Guo H, Ahmed M, Zhang F, Yao CQ, Li S, Liang Y, Hua J, Soares F, Sun Y, Langstein J, *et al.* 2016a Modulation of long noncoding RNAs by risk SNPs underlying genetic predispositions to prostate cancer. *Nature Genetics* **48** 1142. (<https://doi.org/10.1038/ng.3637>)
- Guo Q, Qian Z, Yan D, Li L & Huang L 2016b lncRNA-MEG3 inhibits cell proliferation of endometrial carcinoma by repressing Notch signaling. *Biomedicine and Pharmacotherapy* **82** 589–594. (<https://doi.org/10.1016/j.biopha.2016.02.049>)
- Guo L-J, Zhang S, Gao B, Jiang Y, Zhang X-H, Tian W-G, Hao S, Zhao J-J, Zhang G, Hu C-Y, *et al.* 2017 Low expression of long non-coding RNA GASS is associated with poor prognosis of patients with thyroid cancer. *Experimental and Molecular Pathology* **102** 500–504. (<https://doi.org/10.1016/j.yexmp.2017.05.008>)
- Ha M & Kim VN 2014 Regulation of microRNA biogenesis. *Nature Reviews Molecular Cell Biology* **15** 509–524. (<https://doi.org/10.1038/nrm3838>)
- Hajjari M & Salavaty A 2015 HOTAIR: an oncogenic long non-coding RNA in different cancers. *Cancer Biology & Medicine* **12** 1–9. (<https://doi.org/10.7497/j.issn.2095-3941.2015.0006>)
- Hansji H, Leung EY, Baguley BC, Finlay GJ, Cameron-Smith D, Figueiredo VC & Askarian-Amiri ME 2016 ZFAS1: a long noncoding RNA associated with ribosomes in breast cancer cells. *Biology Direct* **11** 62. (<https://doi.org/10.1186/s13062-016-0165-y>)
- Hata A & Kashima R 2016 Dysregulation of microRNA biogenesis machinery in cancer. *Critical Reviews in Biochemistry and Molecular Biology* **51** 121–134. (<https://doi.org/10.3109/10409238.2015.1117054>)
- Hatchell EC, Colley SM, Beveridge DJ, Epis MR, Stuart LM, Giles KM, Redfern AD, Miles LEC, Barker A, MacDonald LM, *et al.* 2006 SLIRP, a small SRA binding protein, is a nuclear receptor corepressor. *Molecular Cell* **22** 657–668. (<https://doi.org/10.1016/j.molcel.2006.05.024>)
- Heilmann K, Toth R, Bossmann C, Klimo K, Plass C & Gerhauser C 2017 Genome-wide screen for differentially methylated long noncoding RNAs identifies Esrp2 and lncRNA Esrp2-as regulated by enhancer DNA methylation with prognostic relevance for human breast cancer. *Oncogene* **36** 6446. (<https://doi.org/10.1038/onc.2017.246>)
- Hilz S, Modzelewski AJ, Cohen PE & Grimson A 2016 The roles of microRNAs and siRNAs in mammalian spermatogenesis. *Development* **143** 3061. (<https://doi.org/10.1242/dev.136721>)
- Hock J & Meister G 2008 The Argonaute protein family. *Genome Biology* **9** 210. (<https://doi.org/10.1186/gb-2008-9-2-210>)
- Hon C-C, Ramilowski JA, Harshbarger J, Bertin N, Rackham OJL, Gough J, Denisenko E, Schmeier S, Poulsen TM, Severin J, *et al.* 2017 An atlas of human long non-coding RNAs with accurate 5' ends. *Nature* **543** 199–204. (<https://doi.org/10.1038/nature21374>)

- Hong E, Best A, Gautrey H, Chin J, Razdan A, Curk T, Elliott DJ & Tyson-Capper AJ 2015 Unravelling the RNA-binding properties of SAFB proteins in breast cancer cells. *BioMed Research International* **2015** 395816. (<https://doi.org/10.1155/2015/395816>)
- Hosseini ES, Meryet-Figuere M, Sabzalipoor H, Kashani HH, Nikzad H & Asemi Z 2017 Dysregulated expression of long noncoding RNAs in gynecologic cancers. *Molecular Cancer* **16** 107. (<https://doi.org/10.1186/s12943-017-0671-2>)
- Hu W, Alvarez-Dominguez JR & Lodish HF 2012 Regulation of mammalian cell differentiation by long non-coding RNAs. *EMBO Reports* **13** 971–983. (<https://doi.org/10.1038/embor.2012.145>)
- Huang J-K, Ma L, Song W-H, Lu B-Y, Huang Y-B, Dong H-M, Ma X-K, Zhu Z-Z & Zhou R 2016a MALAT1 promotes the proliferation and invasion of thyroid cancer cells via regulating the expression of IQGAP1. *Biomedicine and Pharmacotherapy* **83** 1–7. (<https://doi.org/10.1016/j.biopha.2016.05.039>)
- Huang YS, Chang CC, Lee SS, Jou YS & Shih HM 2016b Xist reduction in breast cancer upregulates AKT phosphorylation via HDAC3-mediated repression of PHLPP1 expression. *Oncotarget* **7** 43256–43266. (<https://doi.org/10.18632/oncotarget.9673>)
- Huarte M 2015 The emerging role of lncRNAs in cancer. *Nature Medicine* **21** 1253–1261. (<https://doi.org/10.1038/nm.3981>)
- Iorio MV, Ferracin M, Liu CG, Veronese A, Spizzo R & Sabbioni S 2005 MicroRNA gene expression deregulation in human breast cancer. *Cancer Research* **65** 7065–7070. (<https://doi.org/10.1158/0008-5472.CAN-05-1783>)
- Jarroux J, Morillon A & Pinskaya M 2017 History, discovery, and classification of lncRNAs. *Advances in Experimental Medicine and Biology* **1008** 1–46. (https://doi.org/10.1007/978-981-10-5203-3_1)
- Jeselsohn R, Cornwell M, Pun M, Buchwalter G, Nguyen M, Bango C, Huang Y, Kuang Y, Paweletz C, Fu X, et al. 2017 Embryonic transcription factor SOX9 drives breast cancer endocrine resistance. *PNAS* **114** E4482–E4491. (<https://doi.org/10.1073/pnas.1620993114>)
- Josson S, Gururajan M, Sung SY, Hu P, Shao C, Zhou HE, Liu C, Lichterman J, Duan P, Li Q, et al. 2014 Stromal fibroblast-derived miR-409 promotes epithelial-to-mesenchymal transition and prostate tumorigenesis. *Oncogene* **34** 2690. (<https://doi.org/10.1038/onc.2014.212>)
- Kanwal R, Plaga AR, Liu X, Shukla GC & Gupta S 2017 MicroRNAs in prostate cancer: functional role as biomarkers. *Cancer Letters* **407** 9–20. (<https://doi.org/10.1016/j.canlet.2017.08.011>)
- Kawaji H, Kasukawa T, Forrest A, Carninci P & Hayashizaki Y 2017 The FANTOM5 collection, a data series underpinning mammalian transcriptome atlases in diverse cell types. *Scientific Data* **4** 170113. (<https://doi.org/10.1038/sdata.2017.113>)
- Kawashima H, Takano H, Sugita S, Takahara Y, Sugimura K & Nakatani T 2003 A novel steroid receptor co-activator protein (SRAP) as an alternative form of steroid receptor RNA-activator gene: expression in prostate cancer cells and enhancement of androgen receptor activity. *Biochemical Journal* **369** 163. (<https://doi.org/10.1042/bj20020743>)
- Kenny TC & Germain D 2017 mtDNA, metastasis, and the mitochondrial unfolded protein response (UPR^m). *Frontiers in Cell and Developmental Biology* **5** 37. (<https://doi.org/10.3389/fcell.2017.00037>)
- Khan NE, Bauer AJ, Schultz KAP, Doros L, Decastro RM, Ling A, Lodish MB, Harney LA, Kase RG, Carr AG, et al. 2017 Quantification of thyroid cancer and multinodular goiter risk in the DICER1 syndrome: a family-based cohort study. *Journal of Clinical Endocrinology and Metabolism* **102** 1614–1622. (<https://doi.org/10.1210/jc.2016-2954>)
- Kim T-K, Hemberg M & Gray JM 2015 Enhancer RNAs: a class of long noncoding RNAs synthesized at enhancers. *Cold Spring Harbor Perspectives in Biology* **7** a018622. (<https://doi.org/10.1101/cshperspect.a018622>)
- Kim KM, Abdelmohsen K, Mustapic M, Kapogiannis D & Gorospe M 2017a RNA in extracellular vesicles. *Wiley Interdisciplinary Reviews: RNA* **8** e1413. (<https://doi.org/10.1002/wrna.1413>)
- Kim KM, Noh JH, Abdelmohsen K & Gorospe M 2017b Mitochondrial noncoding RNA transport. *BMB Reports* **50** 164–174. (<https://doi.org/10.5483/BMBRep.2017.50.4.013>)
- Kino T, Hurt DE, Ichijo T, Nader N & Chrousos GP 2010 Noncoding RNA Gas5 is a growth arrest- and starvation-associated repressor of the glucocorticoid receptor. *Science Signaling* **3** ra8. (<https://doi.org/10.1126/scisignal.2000568>)
- Klinge CM 2015 miRNAs regulated by estrogens, tamoxifen, and endocrine disruptors and their downstream gene targets. *Molecular and Cellular Endocrinology* **418** 273–297. (<https://doi.org/10.1016/j.mce.2015.01.035>)
- Kojima S, Goto Y & Naya Y 2017 The roles of microRNAs in the progression of castration-resistant prostate cancer. *Journal of Human Genetics* **62** 25–31. (<https://doi.org/10.1038/jhg.2016.69>)
- Kotan LD, Cooper C, Darcan S, Carr IM, Ozen S, Yan Y, Hamedani MK, Gurbuz F, Mengen E, Turan I, et al. 2016 Idiopathic hypogonadotropic hypogonadism caused by inactivating mutations in SRA1. *Journal of Clinical Research in Pediatric Endocrinology* **8** 125–134. (<https://doi.org/10.4274/jcrpe.3248>)
- Kouros-Mehr H, Slorach EM, Sternlicht MD & Werb Z 2006 GATA-3 maintains the differentiation of the luminal cell fate in the mammary gland. *Cell* **127** 1041–1055. (<https://doi.org/10.1016/j.cell.2006.09.048>)
- Kozomara A & Griffiths-Jones S 2014 miRBase: annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res* **42** D68–D73. (<https://doi.org/10.1093/nar/gkt1181>)
- Kumarswamy R, Bauters C, Volkmann I, Maury F, Fetisch J, Holzmann A, Lemesle G, de Groote P, Pinet F & Thum T 2014 Circulating long noncoding RNA, LIPCAR, predicts survival in patients with heart failure. *Circulation Research* **114** 1569–1575. (<https://doi.org/10.1161/CIRCRESAHA.114.303915>)
- Kurzynska-Kokorniak A, Koralewska N, Pokornowska M, Urbanowicz A, Tworak A, Mickiewicz A & Figlerowicz M 2015 The many faces of Dicer: the complexity of the mechanisms regulating Dicer gene expression and enzyme activities. *Nucleic Acids Research* **43** 4365–4380. (<https://doi.org/10.1093/nar/gkv328>)
- Lam MTY, Li W, Rosenfeld MG & Glass CK 2014 Enhancer RNAs and regulated transcriptional programs. *Trends in Biochemical Sciences* **39** 170–182. (<https://doi.org/10.1016/j.tibs.2014.02.007>)
- Lan X, Zhang H, Wang Z, Dong W, Sun W, Shao L, Zhang T & Zhang D 2015 Genome-wide analysis of long noncoding RNA expression profile in papillary thyroid carcinoma. *Gene* **569** 109–117. (<https://doi.org/10.1016/j.gene.2015.05.046>)
- Lanz RB, McKenna NJ, Onate SA, Albrecht U, Wong J, Tsai SY, Tsai MJ & O'Malley BW 1999 A steroid receptor coactivator, SRA, functions as an RNA and is present in an SRC-1 complex. *Cell* **97** 17–27. ([https://doi.org/10.1016/S0092-8674\(00\)80711-4](https://doi.org/10.1016/S0092-8674(00)80711-4))
- Laxman B, Morris DS, Yu J, Siddiqui J, Cao J, Mehra R, Lonigro RJ, Tsodikov A, Wei JT, Tomlins SA, et al. 2008 A first-generation multiplex biomarker analysis of urine for the early detection of prostate cancer. *Cancer Research* **68** 645–649. (<https://doi.org/10.1158/0008-5472.CAN-07-3224>)
- Lee J, Park HY, Kim WW, Lee SJ, Jeong JH, Kang SH, Jung JH & Chae YS 2017 Biological function of long noncoding RNA snaR in HER2-positive breast cancer cells. *Tumour Biology* **39** 1010428317707374. (<https://doi.org/10.1177/1010428317707374>)
- Leti F & DiStefano JK 2017 Long noncoding RNAs as diagnostic and therapeutic targets in type 2 diabetes and related complications. *Genes* **8** E207. (<https://doi.org/10.3390/genes8080207>)
- Leygue E 2007 Steroid receptor RNA activator (SRA1): unusual bifaceted gene products with suspected relevance to breast cancer. *Nuclear Receptor Signaling* **5** e006. (<https://doi.org/10.1621/nrs.050006>)

- Li W, Notani D & Rosenfeld MG 2016a Enhancers as non-coding RNA transcription units: recent insights and future perspectives. *Nature Reviews Genetics* **17** 207–223. (<https://doi.org/10.1038/nrg.2016.4>)
- Li W, Zhai L, Wang H, Liu C, Zhang J, Chen W & Wei Q 2016b Downregulation of lncRNA GAS5 causes trastuzumab resistance in breast cancer. *Oncotarget* **7** 27778–27786. (<https://doi.org/10.18632/oncotarget.8413>)
- Li X, Xiong X & Yi C 2016c Epitranscriptome sequencing technologies: decoding RNA modifications. *Nature Methods* **14** 23. (<https://doi.org/10.1038/nmeth.4110>)
- Li J, Wang J, Zhong Y, Guo R, Chu D, Qiu H & Yuan Z 2017a HOTAIR: a key regulator in gynecologic cancers. *Cancer Cell International* **17** 65. (<https://doi.org/10.1186/s12935-017-0434-6>)
- Li JH, Zhang SQ, Qiu XG, Zhang SJ, Zheng SH & Zhang DH 2017b Long non-coding RNA NEAT1 promotes malignant progression of thyroid carcinoma by regulating miRNA-214. *International Journal of Oncology* **50** 708–716. (<https://doi.org/10.3892/ijco.2016.3803>)
- Li W, Li N, Shi K & Chen Q 2017c Systematic review and meta-analysis of the utility of long non-coding RNA GAS5 as a diagnostic and prognostic cancer biomarker. *Oncotarget* **8** 66414–66425. (<https://doi.org/10.18632/oncotarget.19040>)
- Liang C, Li Y & Luo J 2016 A novel method to detect functional microRNA regulatory modules by bicliques merging. *IEEE/ACM Transactions on Computational Biology and Bioinformatics* **13** 549–556. (<https://doi.org/10.1109/TCBB.2015.2462370>)
- Liang HF, Zhang XZ, Liu BG, Jia GT & Li WL 2017 Circular RNA circ-ABC10 promotes breast cancer proliferation and progression through sponging miR-1271. *American Journal of Cancer Research* **7** 1566–1576.
- Lima CR, Gomes CC & Santos MF 2017 Role of microRNAs in endocrine cancer metastasis. *Molecular and Cellular Endocrinology* **456** 62–75. (<https://doi.org/10.1016/j.mce.2017.03.015>)
- Lin A, Li C, Xing Z, Hu Q, Liang K, Han L, Wang C, Hawke DH, Wang S, Zhang Y, et al. 2016 The LINK-A lncRNA activates normoxic HIF1alpha signalling in triple-negative breast cancer. *Nature Cell Biology* **18** 213–224. (<https://doi.org/10.1038/ncb3295>)
- Lin A, Hu Q, Li C, Xing Z, Ma G, Wang C, Li J, Ye Y, Yao J, Liang K, et al. 2017 The LINK-A lncRNA interacts with PtdIns(3,4,5)P3 to hyperactivate AKT and confer resistance to AKT inhibitors. *Nature Cell Biology* **19** 238. (<https://doi.org/10.1038/ncb3473>)
- Liu N, Parisien M, Dai Q, Zheng G, He C & Pan T 2013 Probing N6-methyladenosine RNA modification status at single nucleotide resolution in mRNA and long noncoding RNA. *RNA* **19** 1848–1856. (<https://doi.org/10.1261/rna.041178.113>)
- Liu S, Sheng L, Miao H, Saunders TL, MacDougald OA, Koenig RJ & Xu B 2014 SRA gene knockout protects against diet-induced obesity and improves glucose tolerance. *Journal of Biological Chemistry* **289** 13000–13009. (<https://doi.org/10.1074/jbc.M114.564658>)
- Liu B, Sun L, Liu Q, Gong C, Yao Y, Lv X, Lin L, Yao H, Su F, Li D, et al. 2015 A cytoplasmic NF-κB interacting long noncoding RNA blocks IκB phosphorylation and suppresses breast cancer metastasis. *Cancer Cell* **27** 370–381. (<https://doi.org/10.1016/j.ccr.2015.02.004>)
- Liu C, Wu H-T, Zhu N, Shi Y-N, Liu Z, Ao B-X, Liao D-F, Zheng X-L & Qin L 2016 Steroid receptor RNA activator: biologic function and role in disease. *Clinica Chimica Acta* **459** 137–146. (<https://doi.org/10.1016/j.cca.2016.06.004>)
- Liz J & Esteller M 2016 lncRNAs and microRNAs with a role in cancer development. *Biochimica et Biophysica Acta (BBA): Gene Regulatory Mechanisms* **1859** 169–176. (<https://doi.org/10.1016/j.bbarm.2015.06.015>)
- Luo P, Liu XF, Wang YC, Li ND, Liao SJ, Yu MX, Liang CZ & Tu JC 2017 Prognostic value of abnormally expressed lncRNAs in ovarian carcinoma: a systematic review and meta-analysis. *Oncotarget* **8** 23927–23936. (<https://doi.org/10.18632/oncotarget.23311>)
- Luteijn MJ & Ketting RF 2013 PIWI-interacting RNAs: from generation to transgenerational epigenetics. *Nature Reviews Genetics* **14** 523–534. (<https://doi.org/10.1038/nrg3495>)
- Ma L, Reinhardt F, Pan E, Soutschek J, Bhat B, Marcusson EG, Teruya-Feldstein J, Bell GW & Weinberg RA 2010 Therapeutic silencing of miR-10b inhibits metastasis in a mouse mammary tumor model. *Nature Biotechnology* **28** 341–347. (<https://doi.org/10.1038/nbt.1618>)
- Macconail LE & Garraway LA 2010 Clinical implications of the cancer genome. *Journal of Clinical Oncology* **28** 5219–5228. (<https://doi.org/10.1200/JCO.2009.27.4944>)
- Malih S, Saidijam M & Malih N 2015 A brief review on long noncoding RNAs: a new paradigm in breast cancer pathogenesis, diagnosis and therapy. *Tumour Biology* **37** 1479–1485. (<https://doi.org/10.1007/s13277-015-4572-y>)
- Marrone AK, Beland FA & Pogribny IP 2014 Noncoding RNA response to xenobiotic exposure: an indicator of toxicity and carcinogenicity. *Expert Opinion on Drug Metabolism and Toxicology* **10** 1409–1422. (<https://doi.org/10.1517/17425255.2014.954312>)
- Massillo C, Dalton GN, Farré PL, De Luca P & De Siervi A 2017 Implications of microRNA dysregulation in the development of prostate cancer. *Reproduction* **154** R81–R97. (<https://doi.org/10.1530/REP-17-0322>)
- McKay DB, Xi L, Barthel KKB & Cech TR 2014 Structure and function of steroid receptor RNA activator protein, the proposed partner of SRA noncoding RNA. *Journal of Molecular Biology* **426** 1766–1785. (<https://doi.org/10.1016/j.jmb.2014.01.006>)
- Meng Q, Ren M, Li Y & Song X 2016 lncRNA-RMRP acts as an oncogene in lung cancer. *PLoS ONE* **11** e0164845. (<https://doi.org/10.1371/journal.pone.0164845>)
- Meng S, Zhou H, Feng Z, Xu Z, Tang Y, Li P & Wu M 2017 CircRNA: functions and properties of a novel potential biomarker for cancer. *Molecular Cancer* **16** 94. (<https://doi.org/10.1186/s12943-017-0663-2>)
- Mercer TR & Mattick JS 2013 Structure and function of long noncoding RNAs in epigenetic regulation. *Nature Structural and Molecular Biology* **20** 300–307. (<https://doi.org/10.1038/nsmb.2480>)
- Miano V, Ferrero G, Reineri S, Caizzi L, Annaratone L, Ricci L, Cutrupi S, Castellano I, Cordero F & De Bortoli M 2016 Luminal long non-coding RNAs regulated by estrogen receptor alpha in a ligand-independent manner show functional roles in breast cancer. *Oncotarget* **7** 3201–3216. (<https://doi.org/10.18632/oncotarget.6420>)
- Mirza AH, Kaur S & Pociot F 2017 Long non-coding RNAs as novel players in β cell function and type 1 diabetes. *Human Genomics* **11** 17. (<https://doi.org/10.1186/s40246-017-0113-7>)
- Mondal T, Subhash S, Vaid R, Enroth S, Uday S, Reinius B, Mitra S, Mohammed A, James AR, Hoberg E, et al. 2015 MEG3 long noncoding RNA regulates the TGF-beta pathway genes through formation of RNA-DNA triplex structures. *Nature Communications* **6** 7743. (<https://doi.org/10.1038/ncomms8743>)
- Montagnana M, Benati M, Danese E, Giudici S, Perfranceschi M, Ruzzenente O, Salvagno GL, Bassi A, Gelati M, Paviati E, et al. 2017 Aberrant microRNA expression in patients with endometrial cancer. *International Journal of Gynecological Cancer* **27** 459–466. (<https://doi.org/10.1097/IGC.0000000000000913>)
- Mori M, Triboulet R, Mohseni M, Schlegelmilch K, Shrestha K, Camargo Fernando D & Gregory Richard I 2014 Hippo signaling regulates microprocessor and links cell-density-dependent miRNA biogenesis to cancer. *Cell* **156** 893–906. (<https://doi.org/10.1016/j.cell.2013.12.043>)
- Morris KV & Mattick JS 2014 The rise of regulatory RNA. *Nature Reviews Genetics* **15** 423–437. (<https://doi.org/10.1038/nrg3722>)
- Motterle A, Sanchez-Parra C & Regazzi R 2016 Role of long non-coding RNAs in the determination of beta-cell identity. *Diabetes, Obesity and Metabolism* **18** (Supplement 1) 41–50. (<https://doi.org/10.1111/dom.12714>)
- Muluhngwi P & Klinge CM 2015 Roles for miRNAs in endocrine resistance in breast cancer. *Endocrine-Related Cancer* **22** R279–R300. (<https://doi.org/10.1530/ERC-15-0355>)

- Murugan AK, Munirajan AK & Alzahrani AS 2017 Long noncoding RNAs: emerging players in thyroid cancer pathogenesis. *Endocrine-Related Cancer* **25** R59–R82. (<https://doi.org/10.1530/ERC-17-0188>)
- Nagarajan S, Benito E, Fischer A & Johnsen SA 2015 H4K12ac is regulated by estrogen receptor- α and is associated with BRD4 function and inducible transcription. *Oncotarget* **6** 7305–7317.
- Nakagawa S & Kageyama Y 2014 Nuclear lncRNAs as epigenetic regulators—beyond skepticism. *Biochimica et Biophysica Acta (BBA): Gene Regulatory Mechanisms* **1839** 215–222. (<https://doi.org/10.1016/j.bbagr.2013.10.009>)
- Nakagawa S, Ip JY, Shioi G, Tripathi V, Zong X, Hirose T & Prasanth KV 2012 Malat1 is not an essential component of nuclear speckles in mice. *RNA* **18** 1487–1499. (<https://doi.org/10.1261/rna.033217.112>)
- Nelson AW, Groen AJ, Miller JL, Warren AY, Holmes KA, Tarulli GA, Tilley WD, Katzenellenbogen BS, Hawse JR, Gnanapragasam VJ, et al. 2017 Comprehensive assessment of estrogen receptor beta antibodies in cancer cell line models and tissue reveals critical limitations in reagent specificity. *Molecular and Cellular Endocrinology* **440** 138–150. (<https://doi.org/10.1016/j.mce.2016.11.016>)
- Nik-Zainal S, Davies H, Staaf J, Ramakrishna M, Glodzik D, Zou X, Martincorena I, Alexandrov LB, Martin S, Wedge DC, et al. 2016 Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature* **534** 47–54. (<https://doi.org/10.1038/nature17676>)
- Niknafs YS, Han S, Ma T, Speers C, Zhang C, Wilder-Romans K, Iyer MK, Pitchiaya S, Malik R, Hosono Y, et al. 2016 The lncRNA landscape of breast cancer reveals a role for DSCAM-AS1 in breast cancer progression. *Nature Communications* **7** 12791. (<https://doi.org/10.1038/ncomms12791>)
- Novikova IV, Hennelly SP & Sanbonmatsu KY 2012 Structural architecture of the human long non-coding RNA, steroid receptor RNA activator. *Nucleic Acids Research* **40** 5034–5051. (<https://doi.org/10.1093/nar/gks071>)
- O'Bryan S, Dong S, Mathis JM & Alahari SK 2017 The roles of oncogenic miRNAs and their therapeutic importance in breast cancer. *European Journal of Cancer* **72** 1–11. (<https://doi.org/10.1016/j.ejca.2016.11.004>)
- O'Day E & Lal A 2010 MicroRNAs and their target gene networks in breast cancer. *Breast Cancer Research* **12** 201. (<https://doi.org/10.1186/bcr2484>)
- Ong MS, Cai W, Yuan Y, Leong HC, Tan TZ, Mohammad A, You ML, Arfuso F, Goh BC, Warriar S, et al. 2017 'Lnc'-ing Wnt in female reproductive cancers: therapeutic potential of long non-coding RNAs in Wnt signalling. *British Journal of Pharmacology* **174** 4684–4700. (<https://doi.org/10.1111/bph.13958>)
- Orellana EA, Tenneti S, Rangasamy L, Lyle LT, Low PS & Kasinski AL 2017 FolamiRs: ligand-targeted, vehicle-free delivery of microRNAs for the treatment of cancer. *Science Translational Medicine* **9** eaam9327. (<https://doi.org/10.1126/scitranslmed.aam9327>)
- Ørom Ulf A & Shiekhattar R 2013 Long noncoding RNAs usher in a new era in the biology of enhancers. *Cell* **154** 1190–1193. (<https://doi.org/10.1016/j.cell.2013.08.028>)
- Pallante P, Battista S, Pierantoni GM & Fusco A 2013 Dereglulation of microRNA expression in thyroid neoplasias. *Nature Reviews Endocrinology* **10** 88. (<https://doi.org/10.1038/nrendo.2013.223>)
- Paraskevopoulou MD, Vlachos IS, Karagkouni D, Georgakilas G, Kanellos I, Vergoulis T, Zagganas K, Tsanakas P, Floros E, Dalamagas T, et al. 2016 DIANA-LncBase v2: indexing microRNA targets on non-coding transcripts. *Nucleic Acids Research* **44** D231–D238. (<https://doi.org/10.1093/nar/gkv1270>)
- Patil DP, Chen C-K, Pickering BE, Chow A, Jackson C, Guttman M & Jaffrey SR 2016 m6A RNA methylation promotes XIST-mediated transcriptional repression. *Nature* **537** 369–373. (<https://doi.org/10.1038/nature19342>)
- Peng N, Shi L, Zhang Q, Hu Y, Wang N & Ye H 2017a Microarray profiling of circular RNAs in human papillary thyroid carcinoma. *PLoS ONE* **12** e0170287. (<https://doi.org/10.1371/journal.pone.0170287>)
- Peng WX, Koirala P & Mo YY 2017b lncRNA-mediated regulation of cell signaling in cancer. *Oncogene* **36** 5661–5667. (<https://doi.org/10.1038/onc.2017.184>)
- Piatek MJ & Werner A 2014 Endogenous siRNAs: regulators of internal affairs. *Biochemical Society Transactions* **42** 1174. (<https://doi.org/10.1042/BST20140068>)
- Pickard MR & Williams GT 2014 Regulation of apoptosis by long non-coding RNA GAS5 in breast cancer cells: implications for chemotherapy. *Breast Cancer Research and Treatment* **145** 359–370. (<https://doi.org/10.1007/s10549-014-2974-y>)
- Pishkari S, Paryan M, Hashemi M, Baldini E & Mohammadi-Yeganeh S 2018 The role of microRNAs in different types of thyroid carcinoma: a comprehensive analysis to find new miRNA supplementary therapies. *Journal of Endocrinological Investigation* **41** 269–283.
- Piwecka M, Glažar P, Hernandez-Miranda LR, Memczak S, Wolf SA, Rybak-Wolf A, Filipchyk A, Klironomos F, Cerda Jara CA, Fenske P, et al. 2017 Loss of a mammalian circular RNA locus causes miRNA deregulation and affects brain function. *Science* **357** eaam8526. (<https://doi.org/10.1126/science.aam8526>)
- Prensner JR, Chen W, Han S, Iyer MK, Cao Q, Kothari V, Evans JR, Knudsen KE, Paulsen MT, Ljungman M, et al. 2014 The long non-coding RNA PCAT-1 promotes prostate cancer cell proliferation through cMyc. *Neoplasia* **16** 900–908. (<https://doi.org/10.1016/j.neo.2014.09.001>)
- Qi P, Zhou X-Y & Du X 2016 Circulating long non-coding RNAs in cancer: current status and future perspectives. *Molecular Cancer* **15** 39. (<https://doi.org/10.1186/s12943-016-0524-4>)
- Qian K, Liu G, Tang Z, Hu Y, Fang Y, Chen Z & Xu X 2017 The long non-coding RNA NEAT1 interacted with miR-101 modulates breast cancer growth by targeting EZH2. *Archives of Biochemistry and Biophysics* **615** 1–9. (<https://doi.org/10.1016/j.abb.2016.12.011>)
- Rackham O, Shearwood AM, Mercer TR, Davies SM, Mattick JS & Filipovska A 2011 Long noncoding RNAs are generated from the mitochondrial genome and regulated by nuclear-encoded proteins. *RNA* **17** 2085–2093. (<https://doi.org/10.1261/rna.029405.111>)
- Rashid F, Shah A & Shan G 2016 Long non-coding RNAs in the cytoplasm. *Genomics, Proteomics and Bioinformatics* **14** 73–80. (<https://doi.org/10.1016/j.gpb.2016.03.005>)
- Redfern AD, Colley SM, Beveridge DJ, Ikeda N, Epis MR, Li X, Foulds CE, Stuart LM, Barker A, Russell VJ, et al. 2013 RNA-induced silencing complex (RISC) proteins PACT, TRBP, and Dicer are SRA binding nuclear receptor coregulators. *PNAS* **110** 6536–6541. (<https://doi.org/10.1073/pnas.1301620110>)
- Rehmsmeier M, Steffen P, Hochsmann M & Giegerich R 2004 Fast and effective prediction of microRNA/target duplexes. *RNA* **10** 1507–1517. (<https://doi.org/10.1261/rna.5248604>)
- Ren S, Liu Y, Xu W, Sun Y, Lu J, Wang F, Wei M, Shen J, Hou J, Gao X, et al. 2013 Long noncoding RNA MALAT-1 is a new potential therapeutic target for castration resistant prostate cancer. *Journal of Urology* **190** 2278–2287. (<https://doi.org/10.1016/j.juro.2013.07.001>)
- Ren Q, Liang J, Wei J, Basturk O, Wang J, Daniels G, Gellert LL, Li Y, Shen Y, Osman I, et al. 2014 Epithelial and stromal expression of miRNAs during prostate cancer progression. *American Journal of Translational Research* **6** 329–339.
- Rheinbay E, Parasuraman P, Grimsby J, Tiao G, Engreitz JM, Kim J, Lawrence MS, Taylor-Weiner A, Rodriguez-Cuevas S, Rosenberg M, et al. 2017 Recurrent and functional regulatory mutations in breast cancer. *Nature* **547** 55–60. (<https://doi.org/10.1038/nature22992>)
- Ring A & Dowsett M 2004 Mechanisms of tamoxifen resistance. *Endocrine-Related Cancer* **11** 643–658. (<https://doi.org/10.1677/erc.1.00776>)
- Rinn JL & Chang HY 2012 Genome regulation by long noncoding RNAs. *Annual Review of Biochemistry* **81** 145–166. (<https://doi.org/10.1146/annurev-biochem-051410-092902>)

- Rinn J & Guttman M 2014 RNA and dynamic nuclear organization. *Science* **345** 1240–1241. (<https://doi.org/10.1126/science.1252966>)
- Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Bruggmann SA, Goodnough LH, Helms JA, Farnham PJ, Segal E, et al. 2007 Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* **129** 1311–1323. (<https://doi.org/10.1016/j.cell.2007.05.022>)
- Roden C, Gaillard J, Kanoria S, Rennie W, Barish S, Cheng J, Pan W, Liu J, Cotsapas C, Ding Y, et al. 2017 Novel determinants of mammalian primary microRNA processing revealed by systematic evaluation of hairpin-containing transcripts and human genetic variation. *Genome Research* **27** 374–384. (<https://doi.org/10.1101/gr.208900.116>)
- Rosenberg M, Blum R, Kesner B, Maier VK, Szanto A & Lee JT 2017 Denaturing CLIP, dCLIP, pipeline identifies discrete RNA footprints on chromatin-associated proteins and reveals that CBX7 targets 3' UTRs to regulate mRNA expression. *Cell Systems* **5** 368.e315–385. e315. (<https://doi.org/10.1016/j.cels.2017.09.014>)
- Ru Y, Kechris KJ, Tabakoff B, Hoffman P, Radcliffe RA, Bowler R, Mahaffey S, Rossi S, Calin GA, Bemis L, et al. 2015 The multiMiR R package and database: integration of microRNA–target interactions along with their disease and drug associations. *Nucleic Acids Research* **42** e133. (<https://doi.org/10.1093/nar/gku631>)
- Ruivo CF, Adem B, Silva M & Melo SA 2017 The biology of cancer exosomes: insights and new perspectives. *Cancer Research* **77** 6480. (<https://doi.org/10.1158/0008-5472.CAN-17-0994>)
- Rutter MM, Jha P, Schultz KA, Sheil A, Harris AK, Bauer AJ, Field AL, Geller J & Hill DA 2016 DICER1 mutations and differentiated thyroid carcinoma: evidence of a direct association. *Journal of Clinical Endocrinology and Metabolism* **101** 1–5. (<https://doi.org/10.1210/jc.2015-2169>)
- Ryder M, Ghossein RA, Ricarte-Filho JCM, Knauf JA & Fagin JA 2008 Increased density of tumor-associated macrophages is associated with decreased survival in advanced thyroid cancer. *Endocrine-Related Cancer* **15** 1069–1074. (<https://doi.org/10.1677/ERC-08-0036>)
- Saini HK, Griffiths-Jones S & Enright AJ 2007 Genomic analysis of human microRNA transcripts. *PNAS* **104** 17719–17724. (<https://doi.org/10.1073/pnas.0703890104>)
- Saiselet M, Pita JM, Augenlicht A, Dom G, Tarabichi M, Fimereli D, Dumont JE, Detours V & Maenhaut C 2016 miRNA expression and function in thyroid carcinomas: a comparative and critical analysis and a model for other cancers. *Oncotarget* **7** 52475–52492. (<https://doi.org/10.18632/oncotarget.9655>)
- Salameh A, Lee AK, Cardó-Vila M, Nunes DN, Efstathiou E, Staquicini FI, Dobroff AS, Marchiò S, Navone NM, Hosoya H, et al. 2015 PRUNE2 is a human prostate cancer suppressor regulated by the intronic long noncoding RNA PCA3. *PNAS* **112** 8403–8408. (<https://doi.org/10.1073/pnas.1507882112>)
- Sansone P, Savini C, Kurelac I, Chang Q, Amato LB, Strillacci A, Stepanova A, Iommarini L, Mastroleo C, Daly L, et al. 2017 Packaging and transfer of mitochondrial DNA via exosomes regulate escape from dormancy in hormonal therapy-resistant breast cancer. *PNAS* **114** E9066–E9075. (<https://doi.org/10.1073/pnas.1704862114>)
- Schouten PC, Vollebergh MA, Opdam M, Jonkers M, Loden M, Wesseling J, Hauptmann M & Linn SC 2016 High *XIST* and low 53BP1 expression predict poor outcome after high-dose alkylating chemotherapy in patients with a *BRCA1*-like breast cancer. *Molecular Cancer Therapeutics* **15** 190. (<https://doi.org/10.1158/1535-7163.MCT-15-0470>)
- Siegel RL, Miller KD & Jemal A 2017 Cancer Statistics, 2017. *CA: A Cancer Journal for Clinicians* **67** 7–30.
- Smith B, Agarwal P & Bhowmick NA 2017 MicroRNA applications for prostate, ovarian and breast cancer in the era of precision medicine. *Endocrine-Related Cancer* **24** R157–R172. (<https://doi.org/10.1530/ERC-16-0525>)
- Smolle MA, Bullock MD, Ling H, Pichler M & Haybaeck J 2015 Long non-coding RNAs in endometrial carcinoma. *International Journal of Molecular Sciences* **16** 26463–26472. (<https://doi.org/10.3390/ijms161125962>)
- Sowalsky AG, Xia Z, Wang L, Zhao H, Chen S, Bubley GJ, Balk SP & Li W 2015 Whole transcriptome sequencing reveals extensive unspliced mRNA in metastatic castration-resistant prostate cancer. *Molecular Cancer Research* **13** 98. (<https://doi.org/10.1158/1541-7786.MCR-14-0273>)
- Srivastava SK, Ahmad A, Zubair H, Miree O, Singh S, Rocconi RP, Scalici J & Singh AP 2017 MicroRNAs in gynecological cancers: Small molecules with big implications. *Cancer Letters* **407** 123–138. (<https://doi.org/10.1016/j.canlet.2017.05.011>)
- Su X, Malouf GG, Chen Y, Zhang J, Yao H, Valero V, Weinstein JN, Spano JP, Meric-Bernstam E, Khayat D, et al. 2014 Comprehensive analysis of long non-coding RNAs in human breast cancer clinical subtypes. *Oncotarget* **5** 9864–9876.
- Sun KX, Wu DD, Chen S, Zhao Y & Zong ZH 2017a lncRNA MEG3 inhibit endometrial carcinoma tumorigenesis and progression through PI3K pathway. *Apoptosis* **22** 1543–1552. (<https://doi.org/10.1007/s10495-017-1426-7>)
- Sun MY, Zhu JY, Zhang CY, Zhang M, Song YN, Rahman K, Zhang LJ & Zhang H 2017b Autophagy regulated by lncRNA HOTAIR contributes to the cisplatin-induced resistance in endometrial cancer cells. *Biotechnology Letters* **39** 1477–1484. (<https://doi.org/10.1007/s10529-017-2392-4>)
- Takenaka K, Chen BJ, Modesitt SC, Byrne FL, Hoehn KL & Janitz M 2016 The emerging role of long non-coding RNAs in endometrial cancer. *Cancer Genetics* **209** 445–455. (<https://doi.org/10.1016/j.cancergen.2016.09.005>)
- Theodorou V, Stark R, Menon S & Carroll JS 2013 GATA3 acts upstream of FOXA1 in mediating ESR1 binding by shaping enhancer accessibility. *Genome Research* **23** 12–22. (<https://doi.org/10.1101/gr.139469.112>)
- Torres A, Kozak J, Korolczuk A, Wdowiak P, Domanska-Glonek E, Maciejewski R & Torres K 2016 In vitro and in vivo activity of miR-92a-Locked Nucleic Acid (LNA)-Inhibitor against endometrial cancer. *BMC Cancer* **16** 822. (<https://doi.org/10.1186/s12885-016-2867-z>)
- Tripathi V, Ellis JD, Shen Z, Song DY, Pan Q, Watt AT, Freier SM, Bennett CF, Sharma A, Bubulya PA, et al. 2010 The nuclear-retained noncoding RNA MALAT1 regulates alternative splicing by modulating SR splicing factor phosphorylation. *Molecular Cell* **39** 925–938. (<https://doi.org/10.1016/j.molcel.2010.08.011>)
- Trisciuglio D, Desideri M, Farini V, De Luca T, Di Martile M, Tupone MG, Urbani A, D'Aguanno S & Del Bufalo D 2016 Affinity purification-mass spectrometry analysis of bcl-2 interactome identified SLIRP as a novel interacting protein. *Cell Death and Disease* **7** e2090. (<https://doi.org/10.1038/cddis.2015.357>)
- Van Nostrand EL, Pratt GA, Shishkin AA, Gelboin-Burkhardt C, Fang MY, Sundararaman B, Blue SM, Nguyen TB, Surka C, Elkins K, et al. 2016 Robust transcriptome-wide discovery of RNA-binding protein binding sites with enhanced CLIP (eCLIP). *Nature Methods* **13** 508–514. (<https://doi.org/10.1038/nmeth.3810>)
- van Schooneveld E, Wildiers H, Vergote I, Vermeulen PB, Dirix LY & Van Laere SJ 2015 Dysregulation of microRNAs in breast cancer and their potential role as prognostic and predictive biomarkers in patient management. *Breast Cancer Research* **17** 21. (<https://doi.org/10.1186/s13058-015-0526-y>)
- Vanacore D, Boccellino M, Rossetti S, Cavaliere C, D'Aniello C, Di Franco R, Romano FJ, Montanari M, La Mantia E, Piscitelli R, et al. 2017 Micromas in prostate cancer: an overview. *Oncotarget* **8** 50240–50251. (<https://doi.org/10.18632/oncotarget.16933>)
- Vendramin R, Marine JC & Leucci E 2017 Non-coding RNAs: the dark side of nuclear-mitochondrial communication. *EMBO Journal* **36** 1123–1133. (<https://doi.org/10.15252/embj.201695546>)

- Venkatesh T, Suresh PS & Tsutsumi R 2015 Non-coding RNAs: functions and applications in endocrine-related cancer. *Molecular and Cellular Endocrinology* **416** 88–96. (<https://doi.org/10.1016/j.mce.2015.08.026>)
- Vidaurre S, Fitzpatrick C, Burzio VA, Briones M, Villota C, Villegas J, Echenique J, Oliveira-Cruz L, Araya M, Borgna V, et al. 2014 Down-regulation of the antisense mitochondrial non-coding RNAs (ncRNAs) is a unique vulnerability of cancer cells and a potential target for cancer therapy. *Journal of Biological Chemistry* **289** 27182–27198. (<https://doi.org/10.1074/jbc.M114.558841>)
- Wan L, Kong J, Tang J, Wu Y, Xu E, Lai M & Zhang H 2016 HOTAIRM1 as a potential biomarker for diagnosis of colorectal cancer functions the role in the tumour suppressor. *Journal of Cellular and Molecular Medicine* **20** 2036–2044. (<https://doi.org/10.1111/jcmm.12892>)
- Wang F, Ren S, Chen R, Lu J, Shi X, Zhu Y, Zhang W, Jing T, Zhang C, Shen J, et al. 2014 Development and prospective multicenter evaluation of the long noncoding RNA MALAT-1 as a diagnostic urinary biomarker for prostate cancer. *Oncotarget* **5** 11091–11102.
- Wang J, Zhao X, Guo Z, Ma X, Song Y & Guo Y 2017a Regulation of NEAT1/miR-214-3p on the growth, migration and invasion of endometrial carcinoma cells. *Archives of Gynecology and Obstetrics* **295** 1469–1475. (<https://doi.org/10.1007/s00404-017-4365-1>)
- Wang S, Liang K, Hu Q, Li P, Song J, Yang Y, Yao J, Mangala LS, Li C, Yang W, et al. 2017b JAK2-binding long noncoding RNA promotes breast cancer brain metastasis. *Journal of Clinical Investigation* **127** 91553. (<https://doi.org/10.1172/JCI91553>)
- Wang X, Ruan Y, Wang X, Zhao W, Jiang Q, Jiang C, Zhao Y, Xu Y, Sun F, Zhu Y, et al. 2017c Long intragenic non-coding RNA lincRNA-p21 suppresses development of human prostate cancer. *Cell Proliferation* **50** e12318.
- Wang X, Xu Y, Wang X, Jiang C, Han S, Dong K, Shen M & Xu D 2017d LincRNA-p21 suppresses development of human prostate cancer through inhibition of PKM2. *Cell Proliferation* **50** e12395.
- Warburton AJ & Boone DN 2017 Insights from global analyses of long noncoding RNAs in breast cancer. *Current Pathobiology Reports* **5** 23–34. (<https://doi.org/10.1007/s40139-017-0122-1>)
- Watanabe T & Lin H 2014 Posttranscriptional regulation of gene expression by piwi proteins and piRNAs. *Molecular Cell* **56** 18–27. (<https://doi.org/10.1016/j.molcel.2014.09.012>)
- Weidle UH, Birzele F, Kollmorgen G & Ruger R 2017 Long non-coding RNAs and their role in metastasis. *Cancer Genomics and Proteomics* **14** 143–160. (<https://doi.org/10.21873/cgp.20027>)
- Wolf JB 2013 Principles of transcriptome analysis and gene expression quantification: an RNA-seq tutorial. *Molecular Ecology Resources* **13** 559–572. (<https://doi.org/10.1111/1755-0998.12109>)
- Wu Y, Zhang L, Wang Y, Li H, Ren X, Wei F, Yu W, Wang X, Zhang L, Yu J, et al. 2014 Long noncoding RNA HOTAIR involvement in cancer. *Tumor Biology* **35** 9531–9538. (<https://doi.org/10.1007/s13277-014-2523-7>)
- Xiyuan L, Dechao B, Liang S, Yang W, Shuangfang F, Hui L, Haitao L, Chunlong L, Wenzheng F, Runsheng C, et al. 2017 Using the NONCODE database resource. *Current Protocols in Bioinformatics* **58** 12.16.11–12.16.19. (<https://doi.org/10.1002/cpbi.25>)
- Xu CG, Yang MF, Ren YQ, Wu CH & Wang LQ 2016 Exosomes mediated transfer of lncRNA UCA1 results in increased tamoxifen resistance in breast cancer cells. *European Review for Medical and Pharmacological Sciences* **20** 4362–4368.
- Xu W, Chang J, Du X & Hou J 2017 Long non-coding RNA PCAT-1 contributes to tumorigenesis by regulating FSCN1 via miR-145-5p in prostate cancer. *Biomedicine and Pharmacotherapy* **95** 1112–1118. (<https://doi.org/10.1016/j.biopha.2017.09.019>)
- Xue D, Zhou C, Lu H, Xu R, Xu X & He X 2016a LncRNA GAS5 inhibits proliferation and progression of prostate cancer by targeting miR-103 through AKT/mTOR signaling pathway. *Tumor Biology* **37** 16187–16197. (<https://doi.org/10.1007/s13277-016-5429-8>)
- Xue X, Yang YA, Zhang A, Fong KW, Kim J, Song B, Li S, Zhao JC & Yu J 2016b LncRNA HOTAIR enhances ER signaling and confers tamoxifen resistance in breast cancer. *Oncogene* **35** 2746–2755. (<https://doi.org/10.1038/onc.2015.340>)
- Xue M, Zhuo Y & Shan B 2017 MicroRNAs, long noncoding RNAs, and their functions in human disease. *Methods in Molecular Biology* **1617** 1–25. (https://doi.org/10.1007/978-1-4939-7046-9_1)
- Yahya SMM & Elsayed GH 2015 A summary for molecular regulations of miRNAs in breast cancer. *Clinical Biochemistry* **48** 388–396. (<https://doi.org/10.1016/j.clinbiochem.2014.12.013>)
- Yamazaki T & Hirose T 2015 The building process of the functional paraspeckle with long non-coding RNAs. *Frontiers in Bioscience* **7** 1–41. (<https://doi.org/10.2741/s420>)
- Yang F, Zhang H, Mei Y & Wu M 2014 Reciprocal regulation of HIF-1 α and lincRNA-p21 modulates the warburg effect. *Molecular Cell* **53** 88–100. (<https://doi.org/10.1016/j.molcel.2013.11.004>)
- Yang F, Ning Z, Ma L, Liu W, Shao C, Shu Y & Shen H 2017a Exosomal miRNAs and miRNA dysregulation in cancer-associated fibroblasts. *Molecular Cancer* **16** 148. (<https://doi.org/10.1186/s12943-017-0718-4>)
- Yang K, Hou Y, Li A, Li Z, Wang W, Xie H, Rong Z, Lou G & Li K 2017b Identification of a six-lncRNA signature associated with recurrence of ovarian cancer. *Scientific Reports* **7** 752. (<https://doi.org/10.1038/s41598-017-00763-y>)
- Yang Y, Fan X, Mao M, Song X, Wu P, Zhang Y, Jin Y, Yang Y, Chen L-L, Wang Y, et al. 2017c Extensive translation of circular RNAs driven by N6-methyladenosine. *Cell Research* **27** 626–641. (<https://doi.org/10.1038/cr.2017.31>)
- Yoo SK, Lee S, Kim SJ, Jee HG, Kim BA, Cho H, Song YS, Cho SW, Won JK, Shin JY, et al. 2016 Comprehensive analysis of the transcriptional and mutational landscape of follicular and papillary thyroid cancers. *PLoS Genetics* **12** e1006239. (<https://doi.org/10.1371/journal.pgen.1006239>)
- Yoon J-H, Abdelmohsen K & Gorospe M 2014 Functional interactions among microRNAs and long noncoding RNAs. *Seminars in Cell and Developmental Biology* **34** 9–14. (<https://doi.org/10.1016/j.semcdb.2014.05.015>)
- Zhang B, Arun G, Mao Yuntao S, Lazar Z, Hung G, Bhattacharjee G, Xiao X, Booth Carmen J, Wu J, Zhang C, et al. 2012 The lncRNA Malat1 is dispensable for mouse development but its transcription plays a cis-regulatory role in the adult. *Cell Reports* **2** 111–123. (<https://doi.org/10.1016/j.celrep.2012.06.003>)
- Zhang Z, Zhu Z, Watabe K, Zhang X, Bai C, Xu M, Wu F & Mo YY 2013 Negative regulation of lncRNA GAS5 by miR-21. *Cell Death and Differentiation* **20** 1558–1568. (<https://doi.org/10.1038/cdd.2013.110>)
- Zhang B, Xing X, Li J, Lowdon RE, Zhou Y, Lin N, Zhang B, Sundaram V, Chiappinelli KB, Hagemann IS, et al. 2014 Comparative DNA methylome analysis of endometrial carcinoma reveals complex and distinct deregulation of cancer promoters and enhancers. *BMC Genomics* **15** 868. (<https://doi.org/10.1186/1471-2164-15-868>)
- Zhang A, Zhao Jonathan C, Kim J, Fong K-W, Yang Yeqing A, Chakravarti D, Mo Y-Y & Yu J 2015 LncRNA HOTAIR enhances the androgen-receptor-mediated transcriptional program and drives castration-resistant prostate cancer. *Cell Reports* **13** 209–221. (<https://doi.org/10.1016/j.celrep.2015.08.069>)
- Zhang A, Zhang J, Kaipainen A, Lucas JM & Yang H 2016a Long non-coding RNA: a newly deciphered 'code' in prostate cancer. *Cancer Letters* **375** 323–330. (<https://doi.org/10.1016/j.canlet.2016.03.003>)
- Zhang W, Le TD, Liu L, Zhou ZH & Li J 2016b Predicting miRNA targets by integrating gene regulatory knowledge with expression profiles. *PLoS ONE* **11** e0152860. (<https://doi.org/10.1371/journal.pone.0152860>)
- Zhang JJ, Guo SH & Jia BQ 2016c Down-regulation of long non-coding RNA MEG3 serves as an unfavorable risk factor for survival of patients with breast cancer. *European Review for Medical and Pharmacological Sciences* **20** 5143–5147.
- Zhang Y, Wagner EK, Guo X, May I, Cai Q, Zheng W, He C & Long J 2016d Long intergenic non-coding RNA expression signature in

- human breast cancer. *Scientific Reports* **6** 37821. (<https://doi.org/10.1038/srep37821>)
- Zhang D, Cao J, Zhong Q, Zeng L, Cai C, Lei L, Zhang W & Liu F 2017a Long noncoding RNA PCAT-1 promotes invasion and metastasis via the miR-129-5p-HMGB1 signaling pathway in hepatocellular carcinoma. *Biomedicine and Pharmacotherapy* **95** 1187–1193. (<https://doi.org/10.1016/j.biopha.2017.09.045>)
- Zhang R, Hardin H, Huang W, Chen J, Asioli S, Righi A, Maletta F, Sapino A & Lloyd RV 2017b MALAT1 long non-coding RNA expression in thyroid tissues: analysis by in situ hybridization and Real-time PCR. *Endocrine Pathology* **28** 7–12. (<https://doi.org/10.1007/s12022-016-9453-4>)
- Zhang W, Shi S, Jiang J, Li X, Lu H & Ren F 2017c LncRNA MEG3 inhibits cell epithelial-mesenchymal transition by sponging miR-421 targeting E-cadherin in breast cancer. *Biomedicine and Pharmacotherapy* **91** 312–319. (<https://doi.org/10.1016/j.biopha.2017.04.085>)
- Zhang Y, Yu S, Jiang L, Wang X & Song X 2017d HOTAIR is a promising novel biomarker in patients with thyroid cancer. *Experimental and Therapeutic Medicine* **13** 2274–2278. (<https://doi.org/10.3892/etm.2017.4231>)
- Zhang X, Hamblin MH & Yin KJ 2017e The long noncoding RNA Malat1: its physiological and pathophysiological functions. *RNA Biology* **14** 1705–1714. (<https://doi.org/10.1080/15476286.2017.1358347>)
- Zhao Y, Yang Y, Trovik J, Sun K, Zhou L, Jiang P, Lau TS, Hoivik EA, Salvesen HB, Sun H, *et al.* 2014 A novel wnt regulatory axis in endometrioid endometrial cancer. *Cancer Research* **74** 5103–5117. (<https://doi.org/10.1158/0008-5472.CAN-14-0427>)
- Zheng P, Xiong Q, Wu Y, Chen Y, Chen Z, Fleming J, Gao D, Bi L & Ge F 2015 Quantitative proteomics analysis reveals novel insights into mechanisms of action of long noncoding RNA hox transcript antisense intergenic RNA (HOTAIR) in HeLa Cells. *Molecular and Cellular Proteomics* **14** 1447–1463. (<https://doi.org/10.1074/mcp.M114.043984>)
- Zhou KI, Parisien M, Dai Q, Liu N, Diatchenko L, Sachleben JR & Pan T 2016 N6-methyladenosine modification in a long noncoding RNA hairpin predisposes its conformation to protein binding. *Journal of Molecular Biology* **428** 822–833. (<https://doi.org/10.1016/j.jmb.2015.08.021>)
- Zhou W, Ye X-L, Xu J, Cao M-G, Fang Z-Y, Li L-Y, Guan G-H, Liu Q, Qian Y-H & Xie D 2017 The lncRNA H19 mediates breast cancer cell plasticity during EMT and MET plasticity by differentially sponging miR-200b/c and let-7b. *Science Signaling* **10** eaak9557. (<https://doi.org/10.1126/scisignal.aak9557>)
- Zhu H, Lv Z, An C, Shi M, Pan W, Zhou L, Yang W & Yang M 2016 Onco-lncRNA HOTAIR and its functional genetic variants in papillary thyroid carcinoma. *Scientific Reports* **6** 31969. (<https://doi.org/10.1038/srep31969>)
- Ziegler C & Kretz M 2017 The more the merrier-complexity in long non-coding RNA loci. *Frontiers in Endocrinology* **8** 90. (<https://doi.org/10.3389/fendo.2017.00090>)

Received in final form 18 December 2017

Accepted 12 February 2018

Accepted Preprint published online 12 February 2018