

Non-contiguous finished genome sequence and description of *Alistipes ihumii* sp. nov.

Anne Pfeleiderer^{1†}, Ajay Kumar Mishra^{1†}, Jean-Christophe Lagier¹, Catherine Robert¹, Aurelia Caputo¹, Didier Raoult^{1,2} and Pierre-Edouard Fournier^{1*}

¹ Aix-Marseille Université, URMITE, UM63, Faculté de médecine, Marseille, France ² King Fahd Medical Research Center, King Abdul Aziz University, Jeddah, Saudi Arabia

* Correspondence: (pierre-edouard.fournier@univ-amud.fr)

[†]These two authors contributed equally to this work

Keywords: *Alistipes ihumii*, genome, culturomics, taxono-genomics

Alistipes ihumii strain AP11^T sp. nov. is the type strain of *A. ihumii* sp. nov., a new species within the genus *Alistipes*. This strain, whose genome is described here, was isolated from the fecal flora of a 21-year-old French Caucasian female, suffering from a severe restrictive form of anorexia nervosa since the age of 12 years. *A. ihumii* is a Gram-negative anaerobic bacillus. Here we describe the features of this organism, together with the complete genome sequence and annotation. The 2,753,264 bp long genome (one chromosome but no plasmid) contains 2,254 protein-coding and 47 RNA genes, including 3 rRNA genes.

Introduction

Alistipes ihumii strain AP11^T (= CSUR P204 = DSM 26107) is the type strain of *A. ihumii* sp. nov. This bacterium is a Gram-negative, non-spore-forming, anaerobic and non-motile bacillus that was isolated from the stool of a 21-year-old French female suffering from anorexia nervosa, and is part of a “culturomics” study aiming at cultivating individually all species within human feces [1-3].

Prokaryotic taxonomy is episodically confronted with the advancement of methodological and conceptual innovations. The current classification methodology for prokaryotes is known as polyphasic taxonomy, and relies on a combination of phenotypic and genotypic characteristics [4]. The number of completely sequenced genomes is geometrically increasing with time, concurrently with the decrease in cost of such techniques. To date, more than 6,000 bacterial genomes have been published and approximately 25,000 genome sequencing projects have been announced [5]. We recently proposed to integrate genomic information in the taxonomic framework for the description of new bacterial species [6-27].

The genus *Alistipes* (Rautio *et al.* 2003) was created in 2003 [28] and is composed of strictly anaerobic Gram-negative rods that resemble the *Bacteroides fragilis* group in that most species are bile-resistant and indole-positive [29]. This genus

is currently comprised of five species with validly published names, including *A. finegoldii*, *A. putredinis* [28], *A. indistinctus* [30], *A. onderdonkii* and *A. shahii* [31], to which we added three proposed new species, *A. senegalensis* [8], *A. timonensis* [9] and *A. obesi* [22].

Here we present a summary classification and a set of features for a new *Alistipes* species, *A. ihumii* sp. nov. strain AP11^T (= CSUR P204 = DSM 26107), together with the description of the complete genomic sequence and its annotation.

Classification and features

A stool sample was collected from a 21-year-old French Caucasian female suffering from severe restrictive form of anorexia nervosa since the age of 12 years. At the time of sample collection, she was hospitalized in our hospital for recent aggravation of her medical condition (BMI: 10.4 kg/m²). The patient gave an informed and signed consent. Both this study and the assent procedure were approved by the Ethics Committee of the Institut Fédératif de Recherche IFR48, Faculty of Medicine, Marseille, France under reference 09-022. Ten other potentially new bacterial species were isolated from this patient's stool, all of which are currently being described. Microbial culturomics also enabled the isolation of several other new

bacterial species from other stool specimens [6-27]. The fecal specimen was stored at -80°C immediately after collection. Strain AP11^T was isolated in November 2011 after 2 days of inoculation in anaerobic blood culture bottle with the addition of 5mL of thioglycolate and further inoculation on Columbia agar (BioMerieux, Marcy l'Etoile, France).

This strain exhibited a 95% 16S rRNA sequence similarity with *A. indistinctus* [30], the

phylogenetically closest *Alistipes* species with a validly published name (Table 1, Figure 1), and 92% with *A. onderdonkii* [28] and *A. putredinis* [31]. This value was in the range of 16S rRNA sequence identities among species within the genus *Alistipes* that range from 90 to 95%, and lower than the 98.7% 16S rRNA gene sequence threshold recommended by Stackebrandt and Ebers to delineate a new species without carrying out DNA-DNA hybridization [41].

Table 1. Classification and general features of *Alistipes ihumii* strain AP11^T according to the MIGS recommendations [32]

MIGS ID	Property	Term	Evidence code ^a
		Domain <i>Bacteria</i>	TAS [33]
		Phylum <i>Bacteroidetes</i>	TAS [34,35]
		Class <i>Bacteroidia</i>	TAS [34,36]
		Order <i>Bacteroidales</i>	TAS [34,37]
		Family <i>Rikenellaceae</i>	TAS [34,38]
		Genus <i>Alistipes</i>	TAS [28,39]
		Species <i>Alistipes ihumii</i>	IDA
	Current classification	Type strain AP11 ^T	IDA
	Gram stain	Negative	IDA
	Cell shape	Rod	IDA
	Motility	nonmotile	IDA
	Sporulation	nonsporulating	IDA
	Temperature range	mesophile	IDA
	Optimum temperature	37°C	IDA
MIGS-6.3	Salinity	unknown	IDA
MIGS-22	Oxygen requirement	anaerobic	IDA
	Carbon source	unknown	
	Energy source	unknown	
MIGS-6	Habitat	human gut	IDA
MIGS-15	Biotic relationship	free living	IDA
	Pathogenicity	unknown	
	Biosafety level	2	
MIGS-14	Isolation	human feces	
MIGS-4	Geographic location	France	IDA
MIGS-5	Sample collection time	November 2011	IDA
MIGS-4.1	Latitude & Longitude	43.296482 & 5.36978	IDA
MIGS-4.3	Depth	surface	IDA
MIGS-4.4	Altitude	0 m above sea level	IDA

^aEvidence codes - IDA: Inferred from Direct Assay; TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [40]. If the evidence is IDA, then the property was directly observed for a live isolate by one of the authors or an expert mentioned in the acknowledgements.




Figure 1. Phylogenetic tree highlighting the position of *Alistipes ihumii* strain AP11^T relative to other type strains within the genus *Alistipes*. GenBank accession numbers are indicated in parentheses. Sequences were aligned using CLUSTALW, and phylogenetic inferences obtained using the maximum-likelihood method within the MEGA software. Numbers at the nodes are percentages of bootstrap values obtained by repeating the analysis 500 times to generate a majority consensus tree. *Bacteroides splanchnicus* was used as the outgroup. The scale bar represents a 2% nucleotide sequence divergence.

Different growth temperatures (25, 30, 37, 45°C) were tested. Growth was observed between 25 and 45°C, with optimal growth at 37°C after 24 hours of inoculation. Colonies were about 0.2 mm in diameter, transparent, and exhibited a β -hemolytic activity on blood-enriched Columbia agar. Growth of the strain was tested on 5% sheep blood agar, under anaerobic and microaerophilic conditions using the GENbag anaer and GENbag microaer systems, respectively (BioMérieux), and under aerobic conditions with or without 5% CO₂. Optimal growth of this strain was obtained anaerobically, weak growth was observed under microaerophilic conditions, and no growth was observed under aerobic atmosphere. The motility test was negative. Cells grown on agar are Gram-negative rods (Figure 2) and have mean diameter and length of 0.72 and 1.69 μ m, respectively, as determined using electron microscopy (Figure 3). Strain AP11^T exhibited oxidase but no catalase activities. Using API 50CH (BioMérieux), we observed that strain AP11^T was asaccharolytic. Using

API 32A (BioMérieux), positive reactions were obtained for α -glucosidase, β -glucosidase, N-acetyl- β -glucosaminidase, mannose and raffinose fermentation, alkaline phosphatase, leucyl glycine arylamidase, alanine arylamidase, and glutamyl glutamic acid arylamidase. Weak reactions were observed for α -galactosidase and glutamic acid decarboxylase. Negative reactions were obtained for urease, arginine dihydrolase, β -galactosidase, 6 phospho- β -galactosidase, α -arabinosidase, β -glucuronidase, α -fucosidase, nitrate reduction, indole production, arginine arylamidase, proline arylamidase, phenylalanine arylamidase, leucine arylamidase, pyroglutamic acid arylamidase, tyrosine arylamidase, glycine arylamidase, histidine arylamidase, and serine arylamidase. *A. ihumii* is susceptible to amoxicillin, imipenem, and clindamycin, but resistant to vancomycin. When compared with representative species from the genus *Alistipes*, strain AP11^T exhibited the phenotypic differences detailed in Table 2.




Figure 2. Gram stain of *A. ihumii* strain AP11^T




Figure 3. Transmission electron microscopy of *A. ihumii* strain AP11^T, using a Morgani 268D (Philips) at an operating voltage of 60kV. The scale bar represents 500 nm.

Table 2. Differential characteristics of *Alistipes* strains[†]

Properties	<i>A. ihumii</i>	<i>A. senegalensis</i>	<i>A. timonensis</i>	<i>A. putredinis</i>	<i>A. indistinctus</i>	<i>A. shahii</i>	<i>A. obesi</i>
Cell diameter (µm)	0.72	0.56	0.62	0.40	0.60	0.15	0.44-0.76
Oxygen requirement	anaerobic	anaerobic	anaerobic	anaerobic	anaerobic	anaerobic	anaerobic
Pigment production	–	+	+	–	+	+	+
Gram stain	–	–	–	–	–	–	–
Salt requirement	na	+	–	–	–	–	–
Motility	–	–	–	–	–	–	+
Endospore formation	–	–	–	–	–	–	na
Production of							
Alkaline phosphatase	na	na	na	+	w	+	+
Catalase	–	+	+	+	+	–	+
Oxidase	+	–	–	–	–	–	–
Nitrate reductase	–	na	na	–	–	–	–
Urease	–	na	na	–	–	+	–
β-galactosidase	–	w	+	–	–	+	+
N-acetylglucosamine	+	na	W	–	+	+	+
Indole	–	w	W	+	–	+	–
Activity for							
Leucyl glycine arylamidase	+	+	+	+	–	+	+
Glutamic acid decarboxylase	w	na	+	+	–	–	–
Glycine arylamidase	–	+	+	na	–	–	–
Chymotrypsin	na	na	na	–	–	–	na
Acid from							
L-Arabinose	na	na	na	–	+	na	na
Raffinose	+	na	–	–	+	+	–
Mannose	+	+	–	–	+	+	–
Mannitol	na	na	na	na	+	na	na
Sucrose	na	na	na	–	+	+	na
D-glucose	na	na	na	–	+	+	na
D-fructose	na	na	na	–	+	+	na
D-maltose	na	na	na	–	+	+	na
D-lactose	na	na	na	–	+	+	na
Hydrolysis of gelatin	na	na	na	+	+	–	na
G+C content (mol%)	57.90	58.40	58.82	55.3	55.2	57.20	58.60
Habitat	human gut	human gut	human gut	appendix of children	human gut	human gut	human gut

na = data not available; w = weak

[†]*Alistipes ihumii* strain AP11^T, *A. senegalensis* strain JC50^T, *A. timonensis* strain JC136^T, *A. putredinis* strain ATCC29800^T, *A. indistinctus* strain YIT12060^T, *A. shahii* strain WAL 8301^T, *A. obesi* strain ph8^T and *A. finegoldii* AHN2437^T

Matrix-assisted laser-desorption/ionization time-of-flight (MALDI-TOF) MS protein analysis was carried out as previously described [42] using a Microflex spectrometer (Bruker Daltonics, Leipzig, Germany). Twelve individual colonies were deposited on a MTP 384 MALDI-TOF target plate (Bruker). The twelve AP11^T spectra were imported into the MALDI BioTyper software (version 2.0, Bruker) and analyzed by standard pattern matching (with default parameter settings) against the main spectra of 4,706 bacteria, including spectra from *A. finegoldii*, *A. onderdonkii*, *A. shahii*, *A.*

senegalensis, *A. obesi* and *A. timonensis*, used as reference data in the BioTyper database. The output score enabled the presumptive identification and discrimination of the tested species from those in the database: a score ≥ 2 with a validated species identifies a strain at the species level; and a score < 1.7 indicates a species-level match was not made. For strain AP11^T, no significant score was obtained, suggesting that our isolate was not a member of any known species (Figures 4 and 5). We added the spectrum from strain AP11^T to our database.




Figure 4. Reference mass spectrum from *A. ihumii* strain AP11^T. Spectra from 12 individual colonies were compared and a reference spectrum was generated.

Genome sequencing information

Genome project history

The organism was selected for sequencing on the basis of its phylogenetic position and 16S rRNA similarity to other members of the *Alistipes* genus, and is part of a “culturomics” study of the human digestive flora aiming at isolating all bacterial species within human feces. It was the eighth se-

quenced genome from an *Alistipes* species and the first from *Alistipes ihumii* sp. nov. A summary of the project information is shown in Table 3. The Genbank accession number is CAPH00000000 and consists of 60 contigs. Table 3 shows the project information and its association with MIGS version 2.0 compliance [43].




Figure 5. Gel view comparing spectra from *Alistipes ihumii* strain AP11^T and other members of the genus *Alistipes* (*A. obesi*, *A. timonensis*, *A. senegalensis*, *A. shahii*, *A. onderdonkii* and *A. finegoldii*). The Gel View displays the raw spectra of all loaded spectrum files arranged in a pseudo-gel like look, with each peak displayed as a band or bar. The peak intensity is reflected by the intensity of the gray color. The right y-axis shows the relationship between the shades of gray and the peak intensity in arbitrary units. The x-axis records the m/z value. The left y-axis displays the running spectrum number originating from subsequent spectra loading.

Table 3. Project information

MIGS ID	Property	Term
MIGS-31	Finishing quality	High-quality draft
MIGS-28	Libraries used	One 454 paired end 3-kb library
MIGS-29	Sequencing platforms	454 GS FLX Titanium
MIGS-31.2	Fold coverage	35×
MIGS-30	Assemblers	Newbler version 2.5.3
MIGS-32	Gene calling method	Prodigal
	Genbank ID	CAPH00000000
	Genbank Date of Release	November 28, 2012
	Gold ID	Gi20720
MIGS-13	Project relevance	Study of the human gut microbiome

Growth conditions and DNA isolation

A. ihumii sp. nov. strain AP11^T, (= CSURP204 = DSM 26107), was grown aerobically on 5% sheep blood agar medium at 37°C. Five Petri dishes were spread and resuspended in 3x100µl of G2 buffer (EZ1 DNA Tissue kit, Qiagen). A first mechanical lysis was performed by glass powder on the Fastprep-24 device (Sample Preparation system, MP Biomedicals, USA) for 2x20 seconds. DNA was treated with 2.5 µg/µL of lysozyme (30 minutes at 37°C) and extracted using the BioRobot EZ 1 Advanced XL (Qiagen). The DNA was concentrated

and purified on a Qiamp kit (Qiagen). The yield and the concentration of DNA was 70.7 ng/µl as measured by using Quant-it Picogreen kit (Invitrogen) on the Genios Tecan fluorometer.

Genome sequencing and assembly

A 3kb paired-end sequencing strategy (Roche, Meylan, France) was used. DNA (5 µg) was mechanically fragmented for the paired-end sequencing, using a Covaris device (Covaris Inc., Woburn, MA, USA) with an enrichment size of 3-4 kb. The DNA fragmentation was visualized through an Ag-

ilent 2100 BioAnalyzer on a DNA Labchip 7500 which yielded an optimal size of 2.3 kb. The library was constructed using the 454 GS FLX Titanium paired-end rapid library protocol. Circularization and nebulization were performed which generated a pattern of optimal size of 457 bp. PCR amplification was performed for 17 cycles followed by double size selection. The single-stranded paired-end library was quantified using a Quant-it Ribogreen Kit (Invitrogen) and the Genios Tecan fluorometer. The library concentration equivalence was calculated as 1.94×10^{10} molecules/ μL . The library was stored at -20°C until further use.

The paired-end library was clonally amplified with 0.5 and 1 cpb in 2 emPCR reactions with the GS Titanium SV emPCR Kit (Lib-L) v2 (Roche). The yield of the shotgun emPCR reactions was 6.24 and 16.24% respectively for the two kinds of paired-end emPCR reactions according to the quality expected (range of 5 to 20%) from the Roche procedure. Two libraries were loaded on the GS Titanium PicoTiterPlates (PTP Kit 70x75, Roche) and pyrosequenced with the GS Titanium Sequencing Kit XLR70 and the GS FLX Titanium sequencer (Roche). The run was performed overnight and then analyzed on the cluster through the gsRunBrowser and Newbler assembler (Roche). A total of 260,838 passed filter wells were obtained and generated 96.3 Mb with an average length of 369 bp. The passed filter sequences were assembled using Newbler with 90% identity and 40 bp as overlap. The final assembly identified 9 scaffolds and 60 contigs ($> 1,500$ bp) and generated a genome size of 2.75 Mb which corresponds to a coverage of $35\times$ genome equivalent.

Genome annotation

Open Reading Frames (ORFs) were predicted using Prodigal [44] with default parameters but the predicted ORFs were excluded if they were spanning a sequencing gap region. The predicted bacterial protein sequences were searched against the GenBank database [45] and the Clusters of Orthologous Groups (COG) databases using BLASTP. The tRNAScan-SE tool [46] was used to find tRNA genes, whereas ribosomal RNAs were found by using RNAmmer [47] and BLASTn

against the GenBank database. Lipoprotein signal peptides and numbers of transmembrane helices were predicted using SignalP [48] and TMHMM [49] respectively. ORFans were identified if their BLASTP *E*-value was lower than $1e^{-03}$ for alignment length greater than 80 amino acids. If alignment lengths were smaller than 80 amino acids, we used an *E*-value of $1e^{-05}$. Such parameter thresholds have already been used in previous works to define ORFans.

Orthologous gene sets composed of one gene from *A. ihumii* compared to each of *A. obesi* strain ph8^T (GenBank accession number CAHA000000000), *A. fingoldii* strain AHN 2437 (CP003274), *A. indistinctus* strain YIT 12060 (ADLD000000000), *A. putredinis* strain DSM 17216 (ABFK000000000), *A. senegalensis* strain JC50^T (CAHI000000000), *A. shahii* strain WAL 8301 (FP929032), and *A. timonensis* strain JC136^T (CAEG000000000) were identified using the Proteinortho software (version 1.4) [50] using a 30% protein identity and an *E*-value of $1e^{-05}$. The average percentage of nucleotide sequence identity of each orthologous set was determined using the Needleman-Wunsch algorithm global alignment technique. Artemis [51] was used for data management and DNA Plotter [52] was used for visualization of genomic features. The Mauve alignment tool was used for multiple genomic sequence alignment and visualization [53].

Genome properties

The genome of *A. ihumii* strain AP11^T is 2,753,264 bp long (1 chromosome, but no plasmid) with a 57.90% G + C content (Figure 6 and Table 4). Of the 2,301 predicted genes, 2,254 were protein-coding genes, and 47 were RNAs. One rRNA operon (one 16S rRNA, one 23S rRNA and one 5S rRNA) and 44 predicted tRNA genes were identified in the genome. A total of 1,465 genes (63.66%) were assigned a putative function. Two hundred thirty-seven genes were identified as ORFans (10.29%). The remaining genes were annotated as hypothetical proteins. The properties and the statistics of the genome are summarized in Tables 4 and 5. The distribution of genes into COGs functional categories is presented in Table 5.




Figure 6. Graphical circular map of the chromosome. From the outside in, the outer two circles show open reading frames oriented in the forward and reverse (colored by COG categories) directions, respectively. The third circle marks the rRNA gene operon (red) and tRNA genes (green). The fourth circle shows the G+C% content plot. The innermost circle shows GC skew, purple and olive indicating negative and positive values, respectively.

Genome comparison with other *Alistipes* species

Here, we compared the genome of *A. ihumii* strain AP11^T to those of *A. obesi* strain ph8^T (GenBank accession number CAHA000000000), *A. finegoldii* strain AHN 2437 (CP003274), *A. indistinctus* strain YIT 12060 (ADLD000000000), *A. putredinis* strain DSM 17216 (ABFK000000000), *A. senegalensis* strain JC50^T (CAHI000000000), *A. shahii* strain WAL 8301 (FP929032), and *A. timonensis* strain JC136^T (CAEG000000000). The draft genome of *A. ihumii* is larger than that of *A. putredinis* (2.75 and 2.55 Mb, respectively) but smaller than those of *A. indistinctus*, *A. obesi*, *A. timonensis*, *A. finegoldii*, *A. shahii* and *A. senegalensis* (2.85, 3.16, 3.49, 3.73,

3.76, and 4.01 Mb, respectively). The G+C content of *A. ihumii* is comparable to that of *A. shahii* (57.90 and 57.60%, respectively), lower than those of *A. timonensis* and *A. senegalensis* (58.8 and 58.4%, respectively) and higher than those of *A. putredinis*, *A. indistinctus* and *A. finegoldii* (53.30, 54.80 and 56.60%, respectively). *A. ihumii* has a smaller gene content than those of *A. putredinis*, *A. indistinctus*, *A. obesi*, *A. timonensis*, *A. shahii*, *A. senegalensis*, and *A. finegoldii* (2,301, 2,335, 2,342, 2,619, 2,709, 3,132, 3,161, and 3,231, respectively). The ratio of genes per MB of *A. ihumii* is higher than those of *A. timonensis*, *A. senegalensis*, *A. indistinctus*, and *A. obesi* (836, 776,

Table 4. Nucleotide content and gene count levels of the genome

Attribute	Value	% of total ^a
Genome size (bp)	2,753,264	
DNA coding region (bp)	2,320,878	84.29
DNA G+C content (bp)	1,594,140	57.90
Number of replicons	1	
Extrachromosomal elements	0	
Total genes	2,301	100
RNA genes	47	2.04
rRNA operons	1	
Protein-coding genes	2,254	97.95
Genes with function prediction	1,540	66.92
Genes assigned to COGs	1,465	63.66
Protein coding genes assigned Pfam domains	1,834	79.70
Genes with peptide signals	296	12.86
Genes with transmembrane helices	457	19.86
CRISPR repeats	1	

^aThe total is based on either the size of the genome in base pairs or the total number of protein coding genes in the annotated genome

Table 5. Number of genes associated with the 25 general COG functional categories

Code	Value	%age ^a	Description
J	143	6.34	Translation
A	0	0	RNA processing and modification
K	88	3.90	Transcription
L	113	5.01	Replication, recombination and repair
B	0	0	Chromatin structure and dynamics
D	19	0.84	Cell cycle control, mitosis and meiosis
Y	0	0	Nuclear structure
V	28	1.24	Defense mechanisms
T	38	1.69	Signal transduction mechanisms
M	161	7.14	Cell wall/membrane biogenesis
N	6	0.27	Cell motility
Z	0	0	Cytoskeleton
W	0	0	Extracellular structures
U	32	1.42	Intracellular trafficking and secretion
O	60	2.66	Posttranslational modification, protein turnover, chaperones
C	111	4.92	Energy production and conversion
G	106	4.70	Carbohydrate transport and metabolism
E	131	5.81	Amino acid transport and metabolism
F	52	2.31	Nucleotide transport and metabolism
H	75	3.33	Coenzyme transport and metabolism
I	49	2.17	Lipid transport and metabolism
P	72	3.19	Inorganic ion transport and metabolism
Q	21	0.93	Secondary metabolites biosynthesis, transport and catabolism
R	235	10.43	General function prediction only
S	91	4.04	Function unknown
-	790	35.05	Not in COGs

^aThe total is based on the total number of protein coding genes in the annotated genome.

788, 821, and 828, respectively), comparable to that of *A. shahii* (833) and smaller than those of *A. finegoldii* and *A. putredinis* (866 and 915, respectively).

The average genomic nucleotide sequence identity between *A. ihumii* and other *Alistipes* species ranged from 70.23 to 74.37%, whereas values ranged from 69.70 to 90.98% among other *Alistipes* species (Table 6).

However, the distribution of genes into COG categories was not entirely similar in all eight compared genomes (Figure 7).

Conclusion

On the basis of phenotypic, phylogenetic and genomic analyses, we formally propose the creation of *Alistipes ihumii* sp. nov. that contains strain AP11^T. This bacterial strain has been isolated from the fecal flora of a patient suffering from anorexia nervosa living in Marseille, France. Several other new bacterial species were also cultivated from this patient as well as fecal samples from other patients using microbial culturomics [6-27], thus suggesting that the human fecal flora from human remains partially unknown.




Figure 7. Distribution of functional classes of predicted genes in *Alistipes ihumii* (colored in green), *A. senegalensis* (pink), *A. timonensis* (yellow), *A. shahii* (brown), *A. finegoldii* (blue), *A. putredinis* (red), *A. obesi* (orange) and *A. indistinctus* (black) chromosomes according to the clusters of orthologous groups of proteins.

Table 6. Numbers of orthologous proteins shared between genomes

	AIH	ASE	AT	AS	AF	AP	AO	AIN
<i>A. ihumii</i>	2,254	1,190	1,164	958	1,150	1,055	1,130	1,147
<i>A. senegalensis</i>	71.16	3,161	1,764	1,739	1,660	1,277	1,405	1,218
<i>A. timonensis</i>	70.90	90.98	2,709	1,650	1,585	1,238	1,377	1,210
<i>A. shahii</i>	71.19	86.33	80.03	3,132	1,674	1,270	1,166	1,155
<i>A. finegoldii</i>	71.62	82.04	81.14	82.90	3,231	1,303	1,385	1,202
<i>A. putredinis</i>	70.23	75.32	75.21	75.50	76.23	2,335	1,182	1,038
<i>A. onderdonkii</i>	71.26	76.42	76.23	77.06	76.31	74.45	2,619	1,137
<i>A. indistinctus</i>	74.37	70.02	70.05	70.00	69.91	69.70	69.91	2,342

Upper right triangle- numbers of orthologous proteins shared between genomes; lower left triangle- average percentage of nucleotide identity between orthologous gene sets shared between genomes; bold- numbers of proteins per genome. Abbreviations: AIH- *A. ihumii*, ASE- *A. senegalensis*, AT- *A. timonensis*, AS- *A. shahii*, AF- *A. finegoldii*, AP- *A. putredinis*, AO- *A. obesi*, AIN- *A. indistinctus*

Description of *Alistipes ihumii* sp. nov.

Alistipes ihumii (i.hum.i'i. N.L. gen. n. ihumii, based on the acronym IHUMI, the Institut Hospitalo-Universitaire Méditerranée-Infection, where the type strain was isolated).

Colonies are 0.2 mm in diameter and are translucent on blood-enriched Columbia agar. Cells are rod-shaped with a mean diameter of 0.72 μm and a mean length of 1.69 μm . Optimal growth is achieved anaerobically. No growth is obtained aerobically but weak growth is observed in microaerophilic conditions. Growth occurs between 25°C and 45°C, with an optimal growth observed at 37°C.

Cells stain Gram-negative, are non motile and are asaccharolytic. Activities present are α -glucosidase, β -glucosidase, N-acetyl- β -glucosaminidase, mannose and raffinose fermentation, alkaline phosphatase, leucyl glycine arylamidase, alanine arylamidase, and glutamyl glutamic acid arylamidase. Cells are negative for urease, arginine dihydrolase, β -galactosidase, 6-phospho- β -galactosidase, α -arabinosidase, β -glucuronidase,

α -fucosidase, nitrate reduction, indole production, arginine arylamidase, proline arylamidase, phenylalanine arylamidase, leucine arylamidase, pyroglutamic acid arylamidase, tyrosine arylamidase, glycine arylamidase, histidine arylamidase, and serine arylamidase. Cells are susceptible to amoxicillin, imipenem, and clindamycin, but resistant to vancomycin. The G+C content of the genome is 57.90%. The 16S rRNA and genome sequences are deposited in Genbank under accession numbers JX101692 and CAPH00000000, respectively.

The type strain AP11^T (= CSUR P204 = DSM 26107) was isolated from the fecal flora of a 21-year-old French Caucasian female suffering from severe anorexia nervosa.

Acknowledgments

The authors thank the Xegen Company for automating the genomic annotation process. This study was funded by the Mediterranee-Infection Foundation.

References

- Lagier JC, Armougom F, Million M, Hugon P, Pagnier I, Robert C, Bittar F, Fournous G, Gimenez G, Maraninchi M, et al. Microbial culturomics: paradigm shift in the human gut microbiome study. *Clin Microbiol Infect* 2012; **18**:1185-1193. [PubMed](#)
- Dubourg G, Lagier JC, Armougom F, Robert C, Hamad I, Brouqui P. The gut microbiota of a patient with resistant tuberculosis is more comprehensively studied by culturomics than by metagenomics. *Eur J Clin Microbiol Infect Dis* 2013; 2013:637-645.
- Pfleiderer A, Lagier JC, Armougom F, Robert C, Vialettes B, Raoult D. Culturomics identified 11 new bacterial species from a single anorexia nervosa stool sample. [Epub ahead of print]. *Eur J Clin Microbiol Infect Dis* 2013.
- Tindall BJ, Rossello-Mora R, Busse HJ, Ludwig W, Kampfer P. Notes on the characterization of pro-

- karyote strains for taxonomic purposes. *Int J Syst Evol Microbiol* 2010; **60**:249-266. [PubMed](#) <http://dx.doi.org/10.1099/ijs.0.016949-0>
5. Genome Online Database. <http://www.genomesonline.org/cgi-bin/GOLD/index.cgi>
 6. Kokcha S, Mishra AK, Lagier JC, Million M, Leroy Q, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Bacillus timonensis* sp. nov. *Stand Genomic Sci* 2012; **6**:346-355. [PubMed](#) <http://dx.doi.org/10.4056/sigs.2776064>
 7. Lagier JC, El Karkouri K, Nguyen TT, Armougom F, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Anaerococcus senegalensis* sp. nov. *Stand Genomic Sci* 2012; **6**:116-125. [PubMed](#) <http://dx.doi.org/10.4056/sigs.2415480>
 8. Mishra AK, Gimenez G, Lagier JC, Robert C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Alistipes senegalensis* sp. nov. *Stand Genomic Sci* 2012; **6**:304-314. [PubMed](#) <http://dx.doi.org/10.4056/sigs.2625821>
 9. Lagier JC, Armougom F, Mishra AK, Ngyuen TT, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Alistipes timonensis* sp. nov. *Stand Genomic Sci* 2012; **6**:315-324. [PubMed](#) <http://dx.doi.org/10.4056/sigs.2685971>
 10. Mishra AK, Lagier JC, Robert C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Clostridium senegalense* sp. nov. *Stand Genomic Sci* 2012; **6**:386-395. [PubMed](#)
 11. Mishra AK, Lagier JC, Robert C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Peptoniphilus timonensis* sp. nov. *Stand Genomic Sci* 2012; **7**:1-11. [PubMed](#) <http://dx.doi.org/10.4056/sigs.2956294>
 12. Mishra AK, Lagier JC, Rivet R, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Paenibacillus senegalensis* sp. nov. *Stand Genomic Sci* 2012; **7**:70-81. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3056450>
 13. Lagier JC, Gimenez G, Robert C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Herbaspirillum massiliense* sp. nov. *Stand Genomic Sci* 2012; **7**:200-209. [PubMed](#)
 14. Roux V, El Karkouri K, Lagier JC, Robert C, Raoult D. Non-contiguous finished genome sequence and description of *Kurthia massiliensis* sp. nov. *Stand Genomic Sci* 2012; **7**:221-232. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3206554>
 15. Kokcha S, Ramasamy D, Lagier JC, Robert C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Brevibacterium senegalense* sp. nov. *Stand Genomic Sci* 2012; **7**:233-245. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3256677>
 16. Ramasamy D, Kokcha S, Lagier JC, N'Guyen TT, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Aeromicrobium massilense* sp. nov. *Stand Genomic Sci* 2012; **7**:246-257. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3306717>
 17. Lagier JC, Ramasamy D, Rivet R, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Cellulomonas massiliensis* sp. nov. *Stand Genomic Sci* 2012; **7**:258-270. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3316719>
 18. Lagier JC, El Karkouri K, Rivet R, Couderc C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Senegalemassilia anaerobia* sp. nov. *Stand Genomic Sci* 2013; **7**:343-356. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3246665>
 19. Mishra AK, Hugon P, Lagier JC, Nguyen TT, Robert C, Couderc C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Peptoniphilus obesi* sp. nov. *Stand Genomic Sci* 2013; **7**:357-369. [PubMed](#) <http://dx.doi.org/10.4056/sigs.32766871>
 20. Mishra AK, Lagier JC, Nguyen TT, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Peptoniphilus senegalensis* sp. nov. *Stand Genomic Sci* 2013; **7**:370-381. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3366764>
 21. Lagier JC, El Karkouri K, Mishra AK, Robert C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Enterobacter massiliensis* sp. nov. *Stand Genomic Sci* 2013; **7**:399-412. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3396830>
 22. Hugon P, Ramasamy D, Lagier JC, Rivet R, Couderc C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Alistipes obesi* sp. nov. *Stand Ge-*

- nomica* *Sci* 2013; **7**:427-439. [PubMed](#)
<http://dx.doi.org/10.4056/sigs.3336746>
23. Mishra AK, Hugon P, Robert C, Couderc C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Peptoniphilus grossensis* sp. nov. *Stand Genomic Sci* 2012; **7**:320-330. [PubMed](#)
24. Mishra AK, Hugon P, Nguyen TT, Raoult D, Fournier PE. Non contiguous-finished genome sequence and description of *Enorma massiliensis* gen. nov., sp. nov., a new member of the Family *Coriobacteriaceae*. *Stand Genomic Sci* 2013; **8**:290-305. [PubMed](#)
<http://dx.doi.org/10.4056/sigs.3426906>
25. Ramasamy D, Lagier JC, Gorlas A, Raoult D, Fournier PE. Non contiguous-finished genome sequence and description of *Bacillus massiliosenegalensis* sp. nov. *Stand Genomic Sci* 2013; **8**:264-278. [PubMed](#)
<http://dx.doi.org/10.4056/sigs.3496989>
26. Ramasamy D, Lagier JC, Nguyen TT, Raoult D, Fournier PE. Non contiguous-finished genome sequence and description of *Dielma fastidiosa* gen. nov., sp. nov., a new member of the Family *Erysipelotrichaceae*. *Stand Genomic Sci* 2013; **8**:336-351. [PubMed](#)
<http://dx.doi.org/10.4056/sigs.3567059>
27. Mishra AK, Lagier JC, Robert C, Raoult D, Fournier PE. Genome sequence and description of *Timonella senegalensis* gen. nov., sp. nov., a new member of the suborder Micrococccinae. *Stand Genomic Sci* 2013; **8**:318-335. [PubMed](#)
<http://dx.doi.org/10.4056/sigs.3476977>
28. Rautio M, Eerola E, Väisänen-Tunkelrott ML, Molitoris D, Lawson P, Collins MD, Jousimies-Somer H. Reclassification of *Bacteroides putredinis* (Weinberg et al., 1937) in a new genus *Alistipes* gen. nov., as *Alistipes putredinis* comb. nov., and description of *Alistipes finegoldii* sp. nov., from human sources. *Syst Appl Microbiol* 2003; **26**:182-188. [PubMed](#)
<http://dx.doi.org/10.1078/072320203322346029>
29. Tyrrell KL, Warren YA, Citron DM, Goldstein EJC. Re-assessment of phenotypic identifications of *Bacteroides putredinis* to *Alistipes* species using molecular methods. *Anaerobe* 2011; **17**:130-134. [PubMed](#)
<http://dx.doi.org/10.1016/j.anaerobe.2011.04.002>
30. Nagai F, Morotomi M, Watanabe Y, Sakon H, Tanaka R. *Alistipes indistinctus* sp. nov. and *Odoribacter laneus* sp. nov., common members of the human intestinal microbiota isolated from faeces. *Int J Syst Evol Microbiol* 2009; **60**:1296-1302. [PubMed](#)
<http://dx.doi.org/10.1099/ijs.0.014571-0>
31. Song Y, Könönen E, Rautio M, Liu C, Bryk A, Eerola E, Finegold SM. *Alistipes onderdonkii* sp. nov. and *Alistipes shahii* sp. nov., of human origin. *Int J Syst Evol Microbiol* 2006; **56**:1985-1990. [PubMed](#)
<http://dx.doi.org/10.1099/ijs.0.64318-0>
32. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547.
33. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains *Archaea*, *Bacteria*, and *Eucarya*. *Proc Natl Acad Sci USA* 1990; **87**:4576-4579. [PubMed](#)
<http://dx.doi.org/10.1073/pnas.87.12.4576>
34. Validation List No. 143. *Int J Syst Evol Microbiol* 2012; **62**:1-4.
<http://dx.doi.org/10.1099/ijs.0.039487-0>
35. Krieg NR, Ludwig W, Euzéby J, Whitman WB. Phylum XIV. *Bacteroidetes* phyl. nov. In: Krieg NR, Staley JT, Brown DR, Hedlund BP, Paster BJ, Ward NL, Ludwig W, Whitman WB (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Volume 4, Springer, New York, 2011, p. 25.
36. Krieg NR. Class I. *Bacteroidia* class. nov. In: Krieg NR, Staley JT, Brown DR, Hedlund BP, Paster BJ, Ward NL, Ludwig W, Whitman WB (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Volume 4, Springer, New York, 2011, p. 25.
37. Krieg NR. Order I. *Bacteroidales* ord. nov. In: Krieg NR, Staley JT, Brown DR, Hedlund BP, Paster BJ, Ward NL, Ludwig W, Whitman WB (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Volume 4, Springer, New York, 2011, p. 25.
38. Krieg NR, Staley JT, Brown DR, Hedlund BP, Paster BJ, Ward NL, Ludwig W, Whitman WB. Family III. *Rikenellaceae* fam. nov. In: Krieg NR, Staley JT, Brown DR, Hedlund BP, Paster BJ, Ward NL, Ludwig W, Whitman WB (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Volume 4, Springer, New York, 2011, p. 54.
39. Validation List no. 94. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. *Int J Syst Evol Microbiol* 2003; **53**:1701-1702. [PubMed](#)
<http://dx.doi.org/10.1099/ijs.0.03001-0>

40. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 2000; **25**:25-29. [PubMed](#) <http://dx.doi.org/10.1038/75556>
41. Stackebrandt E, Ebers J. Taxonomic parameters revisited: tarnished gold standards. *Microbiol Today* 2006; **33**:152-155.
42. Seng P, Drancourt M, Gouriet F, La Scola B, Fournier PE, Rolain JM, Raoult D. Ongoing revolution in bacteriology: routine identification of bacteria by matrix-assisted laser desorption ionization time-of-flight mass spectrometry. *Clin Infect Dis* 2009; **49**:543-551. [PubMed](#) <http://dx.doi.org/10.1086/600885>
43. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547. [PubMed](#) <http://dx.doi.org/10.1038/nbt1360>
44. Prodigal. <http://prodigal.ornl.gov>
45. GenBank database. <http://www.ncbi.nlm.nih.gov/genbank>
46. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 1997; **25**:955-964. [PubMed](#) <http://dx.doi.org/10.1093/nar/25.5.0955>
47. Lagesen K, Hallin P, Rodland EA, Staerfeldt HH, Rognes T, Ussery DW. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 2007; **35**:3100-3108. [PubMed](#) <http://dx.doi.org/10.1093/nar/gkm160>
48. Bendtsen JD, Nielsen H, von Heijne G, Brunak S. Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol* 2004; **340**:783-795. [PubMed](#) <http://dx.doi.org/10.1016/j.jmb.2004.05.028>
49. Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* 2001; **305**:567-580. [PubMed](#) <http://dx.doi.org/10.1006/jmbi.2000.4315>
50. Lechner M, Findeib S, Steiner L, Marz M, Stadler PF, Prohaska SJ. Proteinortho: Detection of (Co-)orthologs in large-scale analysis. *BMC Bioinformatics* 2011; **12**:124. [PubMed](#) <http://dx.doi.org/10.1186/1471-2105-12-124>
51. Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. Artemis: sequence visualization and annotation. *Bioinformatics* 2000; **16**:944-945. [PubMed](#) <http://dx.doi.org/10.1093/bioinformatics/16.10.944>
52. Carver T, Thomson N, Bleasby A, Berriman M, Parkhill J. DNAPlotter: circular and linear interactive genome visualization. *Bioinformatics* 2009; **25**:119-120. [PubMed](#) <http://dx.doi.org/10.1093/bioinformatics/btn578>
53. Darling AC, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 2004; **14**:1394-1403. [PubMed](#) <http://dx.doi.org/10.1101/gr.2289704>