

Annie Raoult

Non-polyconvexity of the stored energy function of a Saint Venant-Kirchhoff material

Aplikace matematiky, Vol. 31 (1986), No. 6, 417–419

Persistent URL: <http://dml.cz/dmlcz/104220>

Terms of use:

© Institute of Mathematics AS CR, 1986

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

NON-POLYCONVEXITY OF THE STORED ENERGY FUNCTION
OF A SAINT VENANT-KIRCHHOFF MATERIAL

ANNIE RAOULT

(Received October 15, 1984)

Summary. A direct proof of the non-polyconvexity of the stored energy function of a Saint Venant-Kirchhoff material is given by means of a simple counter-example.

Keywords: polyconvexity, stored energy function, Saint Venant-Kirchhoff material.

In his famous paper [1] dealing with existence theorems in nonlinear elasticity, John Ball introduced the notion of polyconvexity and proved the existence of an equilibrium state — understood as a minimizer of the total energy function — for hyperelastic materials whose stored energy function is polyconvex, subjected to conservative applied forces. It is well known, for instance, that Ogden's materials are polyconvex materials [2], but as far as we know there has been no direct proof of the non-polyconvexity of the usual Saint Venant-Kirchhoff model. The purpose of this note is to provide such a direct proof by constructing an easy counter-example. However, there exists an “indirect” proof, where the non-polyconvexity is a consequence of the non weak lower semi-continuity of the associated functional (cf. Nečas [4]).

Let M^3 be the set of real matrices of order 3 and let M_+^3 be the subset of matrices with determinant > 0 . Let us recall [1], [2] that a real-valued function W defined on M_+^3 is polyconvex if and only if there exists a convex function g defined on $M^3 \times M^3 \times \mathbb{R}^{+*}$ such that

$$(1) \quad \forall F \in M_+^3, \quad W(F) = g(F, \operatorname{adj} F, \det F)$$

where $\operatorname{adj} F = \det F(F^{-1})$. Notice [1], [3] that $M^3 \times M^3 \times \mathbb{R}^{+*}$ coincides with the convex hull of $\{(F, \operatorname{adj} F, \det F), F \in M_+^3\}$. The stored energy function of a Saint Venant-Kirchhoff material with Lamé's coefficients λ and μ is

$$(2) \quad W(F) = a_1 \operatorname{tr}(F^T F) + a_2 \operatorname{tr}(F^T F)^2 + b_1 \operatorname{tr}(\operatorname{adj}(F^T F))$$

where

$$(3) \quad a_1 = -\frac{3\lambda + 2\mu}{4}, \quad a_2 = \frac{\lambda + 2\mu}{8}, \quad b_1 = \frac{\lambda}{4}.$$

We want to decide whether W is polyconvex or not. It is well-known that for physical reasons λ and μ are positive; therefore the first coefficient in W is negative and this is the first indication that W need not be polyconvex (note that if all coefficients were nonnegative polyconvexity would be immediate [1], [3]).

Theorem. W is not polyconvex.

Proof. Let us construct a counter-example. Let ε be a positive number, and let F and F' be the following elements of M_+^3 :

$$F = \varepsilon I, \quad F' = \varepsilon \operatorname{diag}(1, 1, 3).$$

One immediately obtains

$$\det F = \varepsilon^3, \quad \operatorname{adj} F = \varepsilon^2 I, \quad \det F' = 3\varepsilon^3, \quad \operatorname{adj} F' = \varepsilon^2 \operatorname{diag}(3, 3, 1),$$

$$\frac{F + F'}{2} = \varepsilon \operatorname{diag}(1, 1, 2),$$

$$\det \frac{F + F'}{2} = 2\varepsilon^3, \quad \operatorname{adj} \frac{F + F'}{2} = \varepsilon^2 \operatorname{diag}(2, 2, 1),$$

so that the following relations are satisfied, (of course, they do not hold for arbitrary F and F' in M_+^3):

$$(4) \quad \frac{F + F'}{2} \in M_+^3, \quad \operatorname{adj} \frac{F + F'}{2} = \frac{\operatorname{adj} F + \operatorname{adj} F'}{2}, \quad \det \frac{F + F'}{2} = \frac{\det F + \det F'}{2}.$$

If W were polyconvex, equations (1) and (4) would lead to

$$(5) \quad W\left(\frac{F + F'}{2}\right) \leq \frac{1}{2}(W(F) + W(F')).$$

For the sake of brevity, let us write

$$F^T F = \varepsilon^2 I, \quad F'^T F' = \varepsilon^2 J, \quad \left(\frac{F + F'}{2}\right)^T \left(\frac{F + F'}{2}\right) = \varepsilon^2 K$$

with $J = \operatorname{diag}(1, 1, 9)$, $K = \operatorname{diag}(1, 1, 4)$.

Then using expression (2), where the first term is homogeneous of degree 1 and the remaining terms are homogeneous of degree 2 with respect to $F^T F$, we derive from

inequality (5)

$$a_1 \operatorname{tr} K \varepsilon^2 + (a_2 \operatorname{tr} K^2 + b_1 \operatorname{tr} \operatorname{adj} K) \varepsilon^4 \leq \\ \frac{1}{2}(a_1(\operatorname{tr} I + \operatorname{tr} J) \varepsilon^2 + (a_2(\operatorname{tr} I + \operatorname{tr} J^2) + b_1(\operatorname{tr} I + \operatorname{tr} \operatorname{adj} J)) \varepsilon^4)$$

and this inequality amounts to

$$a_1 + (25a_2 + 2b_1) \varepsilon^2 \geq 0$$

which (recall that a_1 is negative) cannot be true for ε small enough. \square

References

- [1] *J. Ball*: Convexity conditions and existence theorems in nonlinear elasticity. Arch. Rat. Mech. Anal. 63 (1977), p. 337–403.
- [2] *P. G. Ciarlet*: Lectures on three-dimensional elasticity. Tata Institute Lecture Notes, Springer-Verlag, 1983.
- [3] *P. G. Ciarlet*: Topics in mathematical elasticity, vol. I. North-Holland, Amsterdam, 1985.
- [4] *J. Nečas*: Introduction to the theory of nonlinear equations. Teubner Texte fr Mathematik, Band 52, Leipzig.

Souhrn

NEPOLYKONVEXITA FUNKCE VNITŘNÍ ENERGIE SAINT VENANTOVA-KIRCHHOFFOVA MATERIÁLU

ANNIE RAOULT

Je podán protipříklad dokazující, že funkce vnitřní energie Saint Venantova-Kirchhoffova materiálu není polykonvexní.

Резюме

НЕ-ПОЛИВЫПУКЛОСТЬ ВНУТРЕННЕЙ ФУНКЦИИ МАТЕРИАЛА СЕН ВЭНАН-КИРХГОФА

ANNIE RAOULT

Дается пример материала Сен Вэнан-Кирхгофа, функция внутренней энергии которого не является поливыпуклой.

Author's address: Prof. Annie Raoult, Laboratoire d'Analyse Numérique, Université P. et M. Curie, 4 Place Jussieu, 75230 Paris cedex 05, France.