

NCCN Guidelines® Insights

Non–Small Cell Lung Cancer, Version 2.2013

Featured Updates to the NCCN Guidelines

David S. Ettinger, MD¹; Wallace Akerley, MD²; Hossein Borghaei, DO, MS³; Andrew C. Chang, MD⁴; Richard T. Cheney, MD⁵; Lucian R. Chirieac, MD⁶; Thomas A. D'Amico, MD⁷; Todd L. Demmy, MD⁸; Ramaswamy Govindan, MD⁹; Frederic W. Grannis Jr, MD⁹; Stefan C. Grant, MD, JD¹⁰; Leora Horn, MD, MSc¹¹; Thierry M. Jahan, MD¹²; Ritsuko Komaki, MD¹³; Feng-Ming (Spring) Kong, MD, PhD, MPH⁴; Mark G. Kris, MD¹⁴; Lee M. Krug, MD¹⁴; Rudy P. Lackner, MD¹⁵; Inga T. Lennes, MD¹⁶; Billy W. Loo Jr, MD, PhD¹⁷; Renato Martins, MD, MPH¹⁸; Gregory A. Otterson, MD¹⁹; Jyoti D. Patel, MD²⁰; Mary C. Pinder-Schenck, MD²¹; Katherine M. Pisters, MD¹³; Karen Reckamp, MD, MS⁹; Gregory J. Riely, MD, PhD¹⁴; Eric Rohren, MD, PhD¹³; Theresa A. Shapiro, MD, PhD¹; Scott J. Swanson, MD⁶; Kurt Tauer, MD²²; Douglas E. Wood, MD¹⁸; Stephen C. Yang, MD¹; Kristina Gregory, RN, MSN²³; and Miranda Hughes, PhD²³

Abstract

These NCCN Guidelines Insights focus on the diagnostic evaluation of suspected lung cancer. This topic was the subject of a major update in the 2013 NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) for Non–Small Cell Lung Cancer. The NCCN Guidelines Insights focus on the major updates in the NCCN Guidelines and discuss the new updates in greater detail. (*JNCCN* 2013;11:645–653)

From ¹The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins; ²Huntsman Cancer Institute at the University of Utah; ³Fox Chase Cancer Center; ⁴University of Michigan Comprehensive Cancer Center; ⁵Roswell Park Cancer Institute; ⁶Dana-Farber/Brigham and Women's Cancer Center; ⁷Duke Cancer Institute; ⁸Siteman Cancer Center at Barnes-Jewish Hospital and Washington University School of Medicine; ⁹City of Hope Comprehensive Cancer Center; ¹⁰University of Alabama at Birmingham Comprehensive Cancer Center; ¹¹Vanderbilt-Ingram Cancer Center; ¹²UCSF Helen Diller Family Comprehensive Cancer Center; ¹³The University of Texas MD Anderson Cancer Center; ¹⁴Memorial Sloan-Kettering Cancer Center; ¹⁵UNMC Eppley Cancer Center at The Nebraska Medical Center; ¹⁶Massachusetts General Hospital Cancer Center; ¹⁷Stanford Cancer Institute; ¹⁸Fred Hutchinson Cancer Research Center/Seattle Cancer Care Alliance; ¹⁹The Ohio State University Comprehensive Cancer Center - James Cancer Hospital and Solove Research Institute; ²⁰Robert H. Lurie Comprehensive Cancer Center of Northwestern University; ²¹Moffitt Cancer Center; ²²University of Tennessee Cancer Institute; and ²³National Comprehensive Cancer Network.

Disclosures for the NCCN Non–Small Cell Lung Cancer Panel

Individual disclosures of potential conflicts of interest for the NCCN Non–Small Cell Lung Cancer Panel members can be found on page 646.

Please Note

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) are a statement of consensus of the authors regarding their views of currently accepted approaches to treatment. **The NCCN Guidelines® Insights highlight important changes in the NCCN Guidelines® recommendations from previous versions. Colored markings in the algorithm show changes and the discussion aims to further understanding of these changes by summarizing salient portions of the panel's discussion, including the literature reviewed.**

The NCCN Guidelines Insights do not represent the full NCCN Guidelines; further, the National Comprehensive Cancer Network® (NCCN®) makes no representation or warranties of any kind regarding the content, use, or application of the NCCN Guidelines and NCCN Guidelines Insights and disclaims any responsibility for their applications or use in any way.

The full and most current version of these NCCN Guidelines are available at NCCN.org.

© National Comprehensive Cancer Network, Inc. 2013, All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN.

Non–Small Cell Lung Cancer, Version 2.2013

NCCN: Continuing Education

Accreditation Statement

This activity has been designated to meet the educational needs of physicians, nurses, and pharmacists involved in the management of patients with cancer. There is no fee for this article. The National Comprehensive Cancer Network (NCCN) is accredited by the ACCME to provide continuing medical education for physicians. NCCN designates this journal-based CE activity for a maximum of *1.0 AMA PRA Category 1 Credit(s)*[™]. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

NCCN is accredited as a provider of continuing nursing education by the American Nurses Credentialing Center's Commission on Accreditation.

This activity is accredited for 1.0 contact hour. Accreditation as a provider refers to recognition of educational activities only; accredited status does not imply endorsement by NCCN or ANCC of any commercial products discussed/displayed in conjunction with the educational activity. Kristina M. Gregory, RN, MSN, OCN, is our nurse planner for this educational activity.



National Comprehensive Cancer Network is accredited by the Accreditation Council for Pharmacy Education as a provider of continuing pharmacy education. NCCN designates this continuing education activity for 1.0 contact hour(s) (0.1 CEUs) of continuing education credit in states that recognize ACPE accredited providers. This is a knowledge-based activity. UAN: 0836-0000-13-011-H01-P

All clinicians completing this activity will be issued a certificate of participation. To participate in this journal CE activity: 1) review the learning objectives and author disclosures; 2) study the education content; 3) take the posttest with a 70% minimum passing score and complete the evaluation at <http://education.nccn.org/node/22017>; and 4) view/print certificate.

Release date: June 10, 2013; Expiration date: June 10, 2014

Learning Objectives:

Upon completion of this activity, participants will be able to:

- Integrate into professional practice the updates to NCCN Guidelines for NSCLC.
- Describe the rationale behind the decision-making process for developing the NCCN Guidelines for NSCLC.

EDITOR: Kerrin M. Green, MA, Assistant Managing Editor, *JNCCN—Journal of the National Comprehensive Cancer Network*, has disclosed that she has no relevant financial relationships.

CE AUTHORS: Nicole B. Harrold, BS, Manager, Continuing Education and Grants, has disclosed that she has no relevant financial relationships. Kristina M. Gregory, RN, MSN, OCN, Vice President, Clinical Information Operations, has disclosed that she has no relevant financial relationships. James Prazak, RPh, Assistant Director, Continuing Education and Grants, has disclosed the following relationships with commercial interests: Bristol-Myers Squibb Company: Pension; Pfizer, Inc: Stockholder; United Healthcare Group: Stockholder; Johnson & Johnson: Stockholder. Miranda Hughes, PhD, Oncology Scientist/Senior Medical Writer, has disclosed that she has no relevant financial relationships.

Disclosure of Affiliations and Significant Relationships

The following authors have no relevant financial interests to disclose: Dr. Chang, Dr. Demmy, Dr. Grant, Dr. Kong, Dr. Lackner, Dr. Lennes, Dr. Pisters, Dr. Rohren, Dr. Shapiro, Dr. Wood, and Dr. Yang.

The following authors have disclosed the following relevant financial relationships:

Dr. Akerley: PI for Bristol-Myers Squibb Company; Daiichi-Sankyo Co., and Genentech, Inc. Advisory board member for Boehringer Ingelheim GmbH; Genentech, Inc.; Myriad Genetic Laboratories, Inc.; and Biotheranostics. Consultant for BioMarin and Clariant Consulting.

Dr. Borghaei: PI for Genentech, Inc. and Millennium Pharmaceuticals, Inc. Speaker bureau member for Amgen Inc. and Genentech, Inc.

Dr. Cheney: Advisory board member for OptumHealth.

Dr. Chirieac: Advisory board member for Boehringer Ingelheim GmbH. Consultant for Aposense; Infinity; and Shook, Hardy & Bacon.

Dr. D'Amico: Consultant for Scanlan.

Dr. Ettinger: Consultant for Boehringer Ingelheim GmbH; Eli Lilly and Company; Genentech, Inc.; Biodesix; Gilead; and Roche Laboratories, Inc.

Dr. Govindan: Consultant for AstraZeneca Pharmaceuticals LP; Boehringer Ingelheim GmbH; Bristol-Myers Squibb Company; Covidien AG; Genentech, Inc.; GlaxoSmithKline; and Pfizer Inc.

Dr. Grannis: Expert witness for Steven Phillips. Ex-officio member – President of the medical staff for City of Hope National Medical Center Board of Directors.

Dr. Horn: PI for Boehringer Ingelheim GmbH. Advisory board member for Genentech, Inc. and OSI Pharmaceuticals, Inc.

Dr. Jahan: Sub-investigator for Abbott Laboratories. Co-PI for Eli Lilly and Company; Genentech, Inc.; ImClone Systems Incorporated; and Morphotek Inc. Site PI for Aduro Pharmaceuticals and Merrimack Pharmaceuticals. Advisory board member, speaker bureau member, expert witness, or consultant for Novartis Pharmaceuticals Corporation.

Dr. Komaki: PI for Pfizer Inc.

Dr. Kris: Site PI for Boehringer Ingelheim GmbH and Pfizer Inc. Consultant for Boehringer Ingelheim GmbH and Roche Laboratories, Inc. Advisory board member for Genentech, Inc.; Bind Biosciences; Clovis Oncology, Inc.; and Esanex, Inc.

Dr. Krug: PI for Eli Lilly and Company; Genentech, Inc.; Merck & Co., Inc.; Novartis Pharmaceuticals Corporation; and CanBas. Consultant for Genentech, Inc.; MedImmune Inc.; and Morphotek Inc.

Dr. Loo: Workshop participant for Siemens Medical Solutions Diagnostics. Speaker at Annual Users Meeting for Varian Medical Systems, Inc.

Dr. Martins: PI for Bayer HealthCare; Eisai Inc.; Genentech, Inc.; Novartis Pharmaceuticals Corporation; and Pfizer Inc. Receives research funding from Exelixis Inc.

Dr. Otterson: Receives research funding from Abraxis Oncology; Boehringer Ingelheim GmbH; Celgene Corporation; Genentech, Inc.; GlaxoSmithKline; Pfizer Inc; and Pharmacyclics. Consultant for Abraxis Bioscience, Inc. Advisory board member for Genentech, Inc.

Dr. Patel: Receives research funding from Eli Lilly and Company. Advisory board member for Genentech, Inc.

Dr. Pinder-Schenck: Speakers bureau member for Eli Lilly and Company. PI for Genentech, Inc.; Merck & Co., Inc.; and Pfizer Inc.

Dr. Reckamp: PI for Amgen Inc.; Bristol-Myers Squibb Company; GlaxoSmithKline; Astellas; and Pfizer Inc. Consultant for Amgen Inc.

Dr. Riely: Involved in clinical research for GlaxoSmithKline; Millennium Pharmaceuticals, Inc.; Novartis Pharmaceuticals Corporation; and Chugai. PI for Infinity Pharmaceuticals. Consultant for Celgene Corporation and Daiichi-Sankyo Co. Advisory board member for Chugai.

Dr. Swanson: Advisory board member for Covidien AG and Ethicon, Inc.

Dr. Tauer: Speakers bureau member for Eli Lilly and Company.


The NCCN Guidelines Staff have no conflicts to disclose.

Supported by educational grants from Eisai, Inc.; Millennium: The Takeda Oncology Company; Teva Pharmaceuticals; Bayer HealthCare Pharmaceuticals Inc.; Celgene Corporation; Endo Pharmaceuticals and HealthTronics; Genentech; and ARIAD Pharmaceuticals, Inc.

Non–Small Cell Lung Cancer, Version 2.2013

CLINICAL PRESENTATION

RISK ASSESSMENT^b



^aMultidisciplinary evaluation including thoracic surgeons, thoracic radiologists, and pulmonologists to determine the likelihood of a cancer diagnosis and the optimal diagnostic or follow-up strategy.

^bRisk calculators can be used to quantify individual patient and radiologic factors but do not replace evaluation by a multidisciplinary diagnostic team with substantial experience in the diagnosis of lung cancer.

^cSee Principles of Diagnostic Evaluation (DIAG-A 1 of 2).

Version 2.2013 © National Comprehensive Cancer Network, Inc. 2013. All rights reserved. The NCCN Guidelines[®] and this illustration may not be reproduced in any form without the express written permission of NCCN[®].

DIAG-1

NCCN Categories of Evidence and Consensus

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.


All recommendations are category 2A unless otherwise noted.

Clinical trials: NCCN believes that the best management for any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

Overview

Smoking tobacco is the primary risk factor for lung cancer.¹⁻⁴ The risk of lung cancer increases with the number of packs of cigarettes smoked per day and the number of years spent smoking (ie, pack-years of smoking history).⁵⁻⁷ Thus, current smokers should be encouraged to quit (<http://www.smoke-free.gov/>),^{1,8-11} and former smokers should be encouraged to avoid smoking. Agents that can be used to promote smoking cessation include nicotine replacement (eg, gum, inhaler, lozenge, nasal spray, patch), sustained-release bupropion, and varenicline (<http://www.ahrq.gov/clinic/tobacco/medsmoktab.htm>).⁸⁻¹¹ Other risk factors for lung cancer include radon, disease history, family history of lung cancer, and occupational exposure (eg, asbestos, silica, coal smoke, and diesel fumes; see the NCCN Clinical Practice Guidelines in Oncology [NCCN Guidelines] for Lung Cancer Screening, available online at NCCN.org).^{1,4,12,13}

Non–Small Cell Lung Cancer, Version 2.2013



^aMultidisciplinary evaluation including thoracic surgeons, thoracic radiologists, and pulmonologists to determine the likelihood of a cancer diagnosis and the optimal diagnostic or follow-up strategy.

^bRisk calculators can be used to quantify individual patient and radiologic factors but do not replace evaluation by a multidisciplinary diagnostic team with substantial experience in the diagnosis of lung cancer.

^cSee Principles of Diagnostic Evaluation (DIAG-A 1 of 2).

^dA positive PET result is defined as a standard uptake value (SUV) in the lung nodule greater than the baseline mediastinal blood pool. A positive PET scan finding can be caused by infection or inflammation, including absence of lung cancer with localized infection, presence of lung cancer with associated (eg, postobstructive) infection, and presence of lung cancer with related inflammation (nodal, parenchymal, pleural). A false-negative PET scan can be caused by a small nodule, low cellular density (nonsolid nodule or ground glass opacity [GGO]), or low tumor avidity for FDG (eg, adenocarcinoma in situ [previously known as bronchoalveolar carcinoma], carcinoid tumor).

^ePatients with a suspicion of lung cancer after PET-CT require histologic confirmation before any nonsurgical therapy.

^fThe choice of biopsy or surgical excision should be based on the clinical suspicion of lung cancer, location of lesion (feasibility for surgical identification and resection by minimally invasive video-assisted thoracic surgery [VATS]), and patient preferences.

^gPatients with a strong clinical suspicion of stage I or II lung cancer (based on risk factors and radiologic appearance) do not require a biopsy before surgery.

Version 2.2013 © National Comprehensive Cancer Network, Inc. 2013. All rights reserved. The NCCN Guidelines[®] and this illustration may not be reproduced in any form without the express written permission of NCCN[®].

DIAG-2

Currently, most patients with lung cancer are diagnosed at an advanced stage when curative treatment is not possible. Thus, lung cancer has a higher mortality rate than other types of cancer. Only 15.9% of all patients with lung cancer are alive 5 years or more after diagnosis.¹⁴ In 2013, an estimated 228,000 new diagnoses and 159,500 deaths from lung and bronchial cancer will occur in the United States.¹⁵ Screening with low-dose CT has been shown to decrease the mortality rate from lung cancer.¹⁶ Although screening with low-dose CT can be used to identify possible lung cancer, most cases are currently identified through other means (eg, symptoms, including weight loss, long-term cough, dyspnea), because lung cancer screening is not yet widely available and many insurance companies are not currently paying for screening. Early-stage lung cancer is often detected in patients because a chest radiograph or CT scan (obtained for other reasons) incidentally shows a lung nodule or mass. For patients with suspected lung

cancer, diagnostic and treatment strategies should be decided in a multidisciplinary setting and on an individual basis to maximize outcomes.

Data show that low-dose CT can be used to detect lung cancer at an early stage when it is more likely curable.¹⁶ The NCCN Guidelines for Lung Cancer Screening recommend screening with low-dose CT for select high-risk current and former smokers without symptoms of lung cancer¹⁷ (to view the most recent version of these guidelines, visit [NCCN.org](http://www.nccn.org)). Other organizations also recommend screening with low-dose CT for select high-risk patients (eg, American College of Chest Physicians, American Cancer Society, American Lung Association, ASCO, and American Thoracic Society) (<http://www.lung.org/lungdisease/lungcancer/lungcancerscreeningguidelines/>).^{18–20}

The NCCN Guidelines for Non–Small Cell Lung Cancer (NSCLC) discuss management of non-squamous NSCLC (eg, adenocarcinoma), which is the most common type of lung cancer, and also less

Non–Small Cell Lung Cancer, Version 2.2013

PRINCIPLES OF DIAGNOSTIC EVALUATION

- Patients with a strong clinical suspicion of stage I or II lung cancer (based on risk factors and radiologic appearance) do not require a biopsy before surgery.
 - ▶ A biopsy adds time, costs, and procedural risk and may not be needed for treatment decisions.
 - ▶ A preoperative biopsy may be appropriate if a non-lung cancer diagnosis is strongly suspected that can be diagnosed by FNA.
 - ▶ A preoperative biopsy may be appropriate if an intraoperative diagnosis appears difficult or very risky.
 - ▶ If a preoperative tissue diagnosis has not been obtained, then an intraoperative diagnosis (ie, wedge resection or needle biopsy) is necessary before lobectomy, bilobectomy, or pneumonectomy.
- Bronchoscopy should preferably be performed during the planned surgical resection, rather than as a separate procedure.
 - ▶ Bronchoscopy is required before surgical resection (see NSCL-2).
 - ▶ A separate bronchoscopy may not be needed for treatment decisions before the time of surgery and adds time, costs, and procedural risk.
 - ▶ A preoperative bronchoscopy may be appropriate if a central tumor requires pre-resection evaluation for biopsy, surgical planning (eg, potential sleeve resection), or preoperative airway preparation (eg, coring out an obstructive lesion).
- Invasive mediastinal staging is recommended before surgical resection for most patients with clinical stage I or II lung cancer (see NSCL-2).
 - ▶ Patients should preferably undergo invasive mediastinal staging as the initial step before the planned resection (during the same anesthetic procedure), rather than as a separate procedure.
 - ▶ A separate staging procedure adds time, costs, coordination of care, inconvenience, and an additional anesthetic risk.
 - ▶ Preoperative invasive mediastinal staging may be appropriate for a strong clinical suspicion of N2 or N3 nodal disease or when intraoperative cytology or frozen section analysis is not available.
- In patients with suspected NSCLC, many techniques are available for tissue diagnosis.
 - ▶ Diagnostic tools that should be routinely available include:
 - ◊ Sputum cytology
 - ◊ Bronchoscopy with biopsy and transbronchial needle aspiration (TBNA)
 - ◊ Image-guided transthoracic needle aspiration (TTNA)
 - ◊ Thoracentesis
 - ◊ Mediastinoscopy
 - ◊ Video-assisted thoracic surgery (VATS) and open surgical biopsy
 - ▶ Diagnostic tools that provide important additional strategies for biopsy include:
 - ◊ Endobronchial ultrasound (EBUS)–guided biopsy
 - ◊ Navigational bronchoscopy

Version 2.2013 © National Comprehensive Cancer Network, Inc. 2013. All rights reserved. The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

DIAG-A
1 of 2

common types of lung cancer (eg, squamous cell carcinoma, large cell carcinoma).²¹ Other types of thoracic cancer are discussed in the NCCN Guidelines for Malignant Pleural Mesothelioma, Small Cell Lung Cancer, and Thymomas and Thymic Carcinomas (to view the most recent version of these guidelines, visit NCCN.org).

Principles of Diagnostic Evaluation

A new section on diagnostic evaluation of suspected lung cancer was added for the 2013 update (see DIAG-A, pages 649 and 650). This new section includes an algorithm for evaluating suspicious lung nodules observed on chest radiograph or CT scan (see DIAG-1 and DIAG-2, page 647 and 648).

Diagnostic Evaluation of Lung Nodules

This new diagnostic section describes the evaluation of suspicious pulmonary nodules or masses that are seen on chest radiograph and CT. Patient factors

(eg, age, smoking history, occupational exposure) and radiologic characteristics (eg, nodule size, shape, other lung pathology) must be carefully evaluated by a multidisciplinary diagnostic team to help formulate decisions about how to proceed with the workup of a suspicious lung nodule. The NCCN Guidelines recommend biopsy or surgical excision for highly suspicious nodules or further surveillance for nodules with a low suspicion of cancer, depending on the type of nodule and a multidisciplinary evaluation of other patient factors (see DIAG-1, page 647). PET/CT scans also may be used to aid the decision-making process. Several factors, including the size of the pulmonary nodule, whether the nodule is changing in size, and the type of nodule (eg, solid noncalcified nodule vs nonsolid nodule), are used to determine whether the nodule is more or less likely to be lung cancer (see DIAG-2, page 648).^{22–25}

Diagnostic Evaluation of Suspected Lung Cancer

The diagnostic strategy benefits from a multidisci-

Non–Small Cell Lung Cancer, Version 2.2013

PRINCIPLES OF DIAGNOSTIC EVALUATION

- The preferred diagnostic strategy for an individual patient depends on the size and location of the tumor, the presence of mediastinal or distant disease, patient characteristics (such as pulmonary pathology and/or other significant comorbidities), and local experience and expertise.
 - ▶ Factors to be considered in choosing the optimal diagnostic step include:
 - ◊ Anticipated diagnostic yield (sensitivity)
 - ◊ Diagnostic accuracy including specificity and particularly the reliability of a negative diagnostic study (that is, true negative)
 - ◊ Adequate volume of tissue specimen for diagnosis and molecular testing
 - ◊ Invasiveness and risk of procedure
 - ◊ Efficiency of evaluation
 - Access and timeliness of procedure
 - Concomitant staging is beneficial, because it avoids additional biopsies or procedures. It is preferable to biopsy the pathology that would confer the highest stage (that is, to biopsy a suspected metastasis or mediastinal lymph node rather than the pulmonary lesion).
 - ◊ Technologies and expertise available
 - ▶ Decisions about the optimal diagnostic steps for suspected stage I to III lung cancer should be made by thoracic radiologists, interventional radiologists, and board-certified thoracic surgeons who devote a significant portion of their practice to thoracic oncology. Multidisciplinary evaluation may also benefit from involvement of a pulmonologist with experience in advanced bronchoscopic techniques for diagnosis, depending on local expertise.
 - ▶ The least invasive biopsy with the highest yield is preferred as the first diagnostic study.
 - ◊ Patients with central masses and suspected endobronchial involvement should undergo bronchoscopy.
 - ◊ Patients with peripheral (outer one-third) nodules should have navigational bronchoscopy, radial EBUS, or TTNA
 - ◊ Patients with suspected nodal disease should be biopsied by EBUS, navigational bronchoscopy, or mediastinoscopy.
 - Esophageal ultrasound (EUS)-guided biopsy provides additional access to station 5, 7, 8, and 9 lymph nodes if these are clinically suspicious.
 - TTNA and anterior mediastinotomy (that is, Chamberlain procedure) provide additional access to anterior mediastinal (station 5 and 6) lymph nodes if these are clinically suspicious.
 - ◊ Lung cancer patients with an associated pleural effusion should undergo thoracentesis and cytology. A negative cytology result on initial thoracentesis does not exclude pleural involvement. An additional thoracentesis and/or thoracoscopic evaluation of the pleura should be considered before starting curative intent therapy.
 - ◊ Patients suspected of having a solitary site of metastatic disease should preferably have tissue confirmation of that site if feasible.
 - ◊ Patients suspected of having metastatic disease should have confirmation from one of the metastatic sites if feasible.
 - ◊ Patients who may have multiple sites of metastatic disease—based on a strong clinical suspicion—should have biopsy of the primary lung lesion or mediastinal lymph nodes if it is technically difficult or very risky to biopsy a metastatic site.

Version 2.2013 © National Comprehensive Cancer Network, Inc. 2013. All rights reserved. The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.


DIAG-A
2 of 2

plinary evaluation that includes thoracic radiologists, thoracic surgeons, and pulmonary physicians. The NCCN Guidelines recommend that the diagnostic strategy be individualized for each patient depending on the size and location of the tumor or nodule, the presence of mediastinal or distant disease, patient characteristics (eg, comorbidities), and local experience. Decisions regarding whether a biopsy (including what type of biopsy) or surgical excision is appropriate depend on several factors, as outlined in the NCCN algorithm (see DIAG-A, pages 649 and 650). In many situations when the suspicion of lung cancer is high and the patient is a surgical candidate, preoperative biopsy may be unnecessary, because it adds time, cost, and procedural risk but is unlikely to alter the plan for surgical resection (ie, pathologic diagnosis is established intraoperatively). However, if an intraoperative diagnosis is risky or difficult or if there is a significant suspicion of an alternative diag-

nosis, then a preoperative biopsy may be beneficial.

The optimal biopsy approach depends on various factors, including size and location of the nodule, associated lung disease (eg, emphysema), proximity to other structures, and experience and technologies available in the local health care system. Factors to be considered in choosing a biopsy technique include diagnostic yield (sensitivity), diagnostic accuracy (especially true-negative yield), adequacy of tissue volume, risk of the procedure, efficiency (access and timeliness of biopsy), and local experience and expertise. In general, the least-invasive biopsy technique with the highest diagnostic yield should be selected, and the NCCN Guidelines outline general principles to help direct clinicians in this process (see DIAG-A, pages 649 and 650). Radial probe endobronchial ultrasound (EBUS), navigational bronchoscopy, or transthoracic needle aspiration is recommended for patients with suspected peripheral nodules.^{26–32} Routine bronchos-

Non-Small Cell Lung Cancer, Version 2.2013



^aSee Principles of Pathologic Review (NSCL-A).

^bTemel JS, Greer JA, Muzikansky A, et al. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010;363:733-742.

^cBased on the CT of the chest: Peripheral = outer third of lung. Central = inner two thirds of lung.

^dT3, N0 related to size or satellite nodules.

^eFor patients considered to have stage IIB and stage III tumors, where more than one treatment modality (surgery, radiation therapy, or chemotherapy) is usually considered, a multidisciplinary evaluation should be performed.

Version 2.2013 © National Comprehensive Cancer Network, Inc. 2013. All rights reserved. The NCCN Guidelines[®] and this illustration may not be reproduced in any form without the express written permission of NCCN[®].

NSCL-1

copy with (or without) EBUS is preferable for centrally located masses. Mediastinoscopy, EBUS, or navigational bronchoscopy is recommended for patients with suspected nodal disease.^{33,34} Another general principle is to preferentially biopsy the abnormality that would simultaneously confer the highest tumor stage. For example, in a patient with a suspicious liver lesion or pleural effusion, liver biopsy or thoracentesis would most efficiently obtain a tissue diagnosis and establish stage IV disease. Communication among the pathologist, medical oncologist, and practitioner performing the biopsy is of the utmost importance to ensure that sufficient tissue is obtained and processed properly to facilitate molecular testing for genetic alterations (eg, epidermal growth factor receptor mutations, ALK gene rearrangements).

If pathology results from biopsy or surgical excision indicate a diagnosis of NSCLC (see Principles of Pathologic Review in the NCCN Guidelines

for NSCLC; to view the complete version of these guidelines, visit NCCN.org), then further evaluation and clinical staging must be performed so that the patient's health care team can determine the most appropriate and effective treatment plan (see NSCL-1, above). Accurate histology must be determined (eg, adenocarcinoma, squamous cell carcinoma) before initiating systemic therapy; a general diagnosis of NSCLC is not recommended.²¹ Diagnosis, staging, and planned resection (eg, lobectomy) are ideally a single operative procedure for patients with early-stage disease (see DIAG-A, pages 649 and 650). Confirmation and staging of lung cancer are essential before lobectomy or pneumonectomy, although they can be performed with intraoperative diagnosis and concomitant mediastinal staging. The mediastinal lymph nodes must be assessed, because upstaging can occur in patients with presumed early-stage NSCLC based on imaging alone.³⁴⁻³⁶

Non–Small Cell Lung Cancer, Version 2.2013

Conclusions

Most patients with lung cancer are currently identified by symptoms, including weight loss, long-term cough, and dyspnea. For patients with suspected lung cancer, diagnostic and treatment strategies should be decided in a multidisciplinary setting and on an individual basis to maximize outcomes. A new section on diagnostic evaluation of suspected lung cancer was added for the 2013 update of the NCCN Guidelines for NSCLC (to view the most recent version of these guidelines, visit NCCN.org). This new diagnostic section describes the evaluation of suspicious pulmonary nodules or masses that are seen on chest radiograph and CT. Patient factors and radiologic characteristics must be carefully evaluated by a multidisciplinary diagnostic team to help formulate decisions about how to proceed with workup of a suspicious lung nodule. The NCCN Guidelines recommend that the diagnostic strategy be individualized for each patient depending on the size and location of the tumor, the presence of mediastinal or distant disease, patient characteristics, and local experience. The least-invasive biopsy technique with the highest diagnostic yield should be selected; the NCCN Guidelines outline general principles for this process. When obtaining a biopsy specimen, it is essential that the pathologist, medical oncologist, and practitioner confer to ensure that sufficient tissue is obtained and processed properly to facilitate molecular testing. If NSCLC is diagnosed, then further evaluation and clinical staging must be performed so that the patient's health care team can determine the most appropriate and effective treatment plan.

References

- de Groot P, Munden RF. Lung cancer epidemiology, risk factors, and prevention. *Radiol Clin North Am* 2012;50:863–876.
- Alberg AJ, Ford JG, Samet JM. Epidemiology of lung cancer: ACCP evidence-based clinical practice guidelines. 2nd ed. *Chest* 2007;132:29S–55S.
- Dela Cruz CS, Tanoue LT, Matthay RA. Lung cancer: epidemiology, etiology, and prevention. *Clin Chest Med* 2011;32:605–644.
- Secretan B, Straif K, Baan R, et al. A review of human carcinogens—part E: tobacco, areca nut, alcohol, coal smoke, and salted fish. *Lancet Oncol* 2009;10:1033–1034.
- Jha P, Ramasundarahettige C, Landsman V, et al. 21st-century hazards of smoking and benefits of cessation in the United States. *N Engl J Med* 2013;368:341–350.
- Thun MJ, Carter BD, Feskanich D, et al. 50-year trends in smoking-related mortality in the United States. *N Engl J Med* 2013;368:351–364.
- Doll R, Peto R. Mortality in relation to smoking: 20 years' observations on male British doctors. *Br Med J* 1976;2:1525–1536.
- Hays JT, McFadden DD, Ebbert JO. Pharmacologic agents for tobacco dependence treatment: 2011 update. *Curr Atheroscler Rep* 2012;14:85–92.
- Rigotti NA. Strategies to help a smoker who is struggling to quit. *JAMA* 2012;308:1573–1580.
- Fiore MC, Baker TB. Clinical practice. Treating smokers in the health care setting. *N Engl J Med* 2011;365:1222–1231.
- Hurt RD, Ebbert JO, Hays JT, McFadden DD. Preventing lung cancer by treating tobacco dependence. *Clin Chest Med* 2011;32:645–657.
- Straif K, Benbrahim-Tallaa L, Baan R, et al. A review of human carcinogens—part C: metals, arsenic, dusts, and fibres. *Lancet Oncol* 2009;10:453–454.
- Baan R, Grosse Y, Straif K, et al. A review of human carcinogens—Part F: chemical agents and related occupations. *Lancet Oncol* 2009;10:1143–1144.
- Howlander N, Noone A, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2009 (Vintage 2009 Populations) based on November 2011 SEER data submission. Bethesda, MD: National Cancer Institute; 2012. Available at: http://seer.cancer.gov/csr/1975_2009_pops09/.
- Siegel R, Naishadham D, Jemal A. Cancer statistics, 2013. *CA Cancer J Clin* 2013;63:11–30.
- Aberle DR, Adams AM, Berg CD, et al. Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med* 2011;365:395–409.
- Wood DE, Eapen GA, Ettinger DS, et al. Lung cancer screening. *J Natl Compr Canc Netw* 2012;10:240–265.
- Wender R, Fontham ET, Barrera E Jr, et al. American Cancer Society lung cancer screening guidelines. *CA Cancer J Clin* 2013;63:107–117.
- Jaklitsch MT, Jacobson FL, Austin JH, et al. The American Association for Thoracic Surgery guidelines for lung cancer screening using low-dose computed tomography scans for lung cancer survivors and other high-risk groups. *J Thorac Cardiovasc Surg* 2012;144:33–38.
- Bach PB, Mirkin JN, Oliver TK, et al. Benefits and harms of CT screening for lung cancer: a systematic review. *JAMA* 2012;307:2418–2429.
- Travis WD, Brambilla E, Noguchi M, et al. International Association for the Study of Lung Cancer/American Thoracic Society/European Respiratory Society international multidisciplinary classification of lung adenocarcinoma. *J Thorac Oncol* 2011;6:244–285.
- Armato SG 3rd, McLennan G, Bidaut L, et al. The Lung Image Database Consortium (LIDC) and Image Database Resource Initiative (IDRI): a completed reference database of lung nodules on CT scans. *Med Phys* 2011;38:915–931.
- MacMahon H, Austin JH, Gamsu G, et al. Guidelines for management of small pulmonary nodules detected on CT scans: a statement from the Fleischner Society. *Radiology* 2005;237:395–400.
- Naidich DP, Bankier AA, Macmahon H, et al. Recommendations for the management of subsolid pulmonary nodules detected at CT: a statement from the Fleischner Society. *Radiology* 2013;266:304–317.
- Godoy MC, Naidich DP. Overview and strategic management of subsolid pulmonary nodules. *J Thorac Imaging* 2012;27:240–248.
- Yasufuku K, Nakajima T, Fujiwara T, et al. Role of endobronchial ultrasound-guided transbronchial needle aspiration in the management of lung cancer. *Gen Thorac Cardiovasc Surg* 2008;56:268–276.
- Jensen KW, Hsia DW, Seijo LM, et al. Multicenter experience with electromagnetic navigation bronchoscopy for the diagnosis of pulmonary nodules. *J Bronchology Interv Pulmonol* 2012;19:195–199.

Non–Small Cell Lung Cancer, Version 2.2013

28. Chao TY, Chien MT, Lie CH, et al. Endobronchial ultrasonography-guided transbronchial needle aspiration increases the diagnostic yield of peripheral pulmonary lesions: a randomized trial. *Chest* 2009;136:229–236.
29. Paone G, Nicastrì E, Lucantoni G, et al. Endobronchial ultrasound-driven biopsy in the diagnosis of peripheral lung lesions. *Chest* 2005;128:3551–3557.
30. Du Rand IA, Barber PV, Goldring J, et al. Summary of the British Thoracic Society guidelines for advanced diagnostic and therapeutic flexible bronchoscopy in adults. *Thorax* 2011;66:1014–1015.
31. Leong S, Ju H, Marshall H, et al. Electromagnetic navigation bronchoscopy: a descriptive analysis. *J Thorac Dis* 2012;4:173–185.
32. Hsia DW, Jensen KW, Curran-Everett D, Musani AI. Diagnosis of lung nodules with peripheral/radial endobronchial ultrasound guided transbronchial biopsy. *J Bronchology Interv Pulmonol* 2012;19:5–11.
33. Rintoul RC, Tournoy KG, El Daly H, et al. EBUS-TBNA for the clarification of PET positive intra-thoracic lymph nodes—an international multi-centre experience. *J Thorac Oncol* 2009;4:44–48.
34. Defranchi SA, Edell ES, Daniels CE, et al. Mediastinoscopy in patients with lung cancer and negative endobronchial ultrasound guided needle aspiration. *Ann Thorac Surg* 2010;90:1753–1757.
35. Yasufuku K, Pierre A, Darling G, et al. A prospective controlled trial of endobronchial ultrasound-guided transbronchial needle aspiration compared with mediastinoscopy for mediastinal lymph node staging of lung cancer. *J Thorac Cardiovasc Surg* 2011;142:1393–1400.e1.
36. Bille A, Pelosi E, Skanjeti A, et al. Preoperative intrathoracic lymph node staging in patients with non-small-cell lung cancer: accuracy of integrated positron emission tomography and computed tomography. *Eur J Cardiothorac Surg* 2009;36:440–445.

Instructions for Completion

To participate in this journal CE activity: 1) review the learning objectives and author disclosures; 2) study the education content; 3) take the posttest with a 70% minimum passing score and complete the evaluation at <http://education.nccn.org/node/22017>; and 4) view/print certificate. After reading the article, you should be able to answer the following multiple-

choice questions. Credit cannot be obtained for tests completed on paper. You must be a registered user on NCCN.org. If you are not registered on NCCN.org, click on “New Member? Sign up here” link on the left-hand side of the Web site to register. Only one answer is correct for each question. Once you successfully answer all posttest questions you will be able to view and/or print your certificate. Software requirements: Internet.

Posttest Questions

1. For patients who appear to have early-stage lung cancer, which of the following is/are TRUE?
 - a. Diagnosis, staging, and resection can be a single operative procedure.
 - b. Multidisciplinary evaluation is not recommended.
 - c. Preoperative biopsy is essential.
 - d. All of the above
 - e. None of the above
2. Which of the following is/are TRUE about biopsies for lung cancer?
 - a. The least invasive biopsy technique with the highest yield should be selected.
 - b. Radial probe endobronchial ultrasound, navigational bronchoscopy, or transthoracic needle aspiration is recommended for patients with suspected peripheral nodules.
 - c. Diagnostic yield (sensitivity), diagnostic accuracy (especially true negative yield), adequacy of tissue volume, risk of the procedure, efficiency (access and timeliness of biopsy), and local experience and expertise should be considered.
 - d. All of the above
 - e. None of the above
3. True or False: All lung nodules observed on low-dose CT scans should be biopsied as soon as possible.

