



University of
Salford
MANCHESTER

Non-specific amplification compromises environmental DNA metabarcoding with COI

Collins, RA, Bakker, J, Wangenstein Fuentes, OS, Soto, AZ, Corrigan, L, Sims, DW, Genner, MJ and Mariani, S

<http://dx.doi.org/10.1111/2041-210X.13276>

Title	Non-specific amplification compromises environmental DNA metabarcoding with COI
Authors	Collins, RA, Bakker, J, Wangenstein Fuentes, OS, Soto, AZ, Corrigan, L, Sims, DW, Genner, MJ and Mariani, S
Publication title	Methods in Ecology and Evolution
Publisher	British Ecological Society
Type	Article
USIR URL	This version is available at: http://usir.salford.ac.uk/id/eprint/51952/
Published Date	2019

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: library-research@salford.ac.uk.

1 Non-specific amplification compromises 2 environmental DNA metabarcoding with COI

3 Rupert A. Collins*¹, Judith Bakker*^{2,3}, Owen S. Wangensteen^{3,4}, Ana Z. Soto³,
4 Laura Corrigan⁵, David W. Sims^{6,7}, Martin J. Genner¹, and Stefano Mariani^{3,8}

5 ¹School of Biological Sciences, University of Bristol, Life Sciences Building, Tyndall Avenue,
6 Bristol BS8 1TQ, UK

7 ²Department of Biological Sciences, Florida International University, 11200 S.W., 8th Street, Miami,
8 Florida, 33199, USA

9 ³Ecosystems & Environment Research Centre, School of Environment & Life Sciences, University
10 of Salford, Salford M5 4WT, UK

11 ⁴Norwegian College of Fishery Science, UiT The Arctic University of Norway, N-9037, Tromsø,
12 Norway

13 ⁵Environment Agency, Tyneside House, Skinnerburn Road, Newcastle upon Tyne NE4 7AR, UK

14 ⁶Marine Biological Association of the United Kingdom, The Laboratory, Citadel Hill, Plymouth PL1
15 2PB, UK

16 ⁷Ocean and Earth Science, University of Southampton, National Oceanography Centre
17 Southampton, European Way, Southampton SO14 3ZH, UK

18 ⁸School of Natural Sciences & Psychology, Liverpool John Moores University, Liverpool, L3 3AF,
19 UK

20 Corresponding author:

21 Stefano Mariani

22 Email address: s.mariani@salford.ac.uk

*Both authors contributed equally to this work.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi:

10.1111/2041-210X.13276

This article is protected by copyright. All rights reserved.

24 1. Metabarcoding extra-organismal DNA from environmental samples is now a key technique in aquatic
25 biomonitoring and ecosystem health assessment. Of critical consideration when designing experiments,
26 and especially so when developing community standards and legislative frameworks, is the choice of
27 genetic marker and primer set. Mitochondrial cytochrome *c* oxidase subunit I (COI), the standard DNA
28 barcode marker for animals, with its extensive reference library, taxonomic discriminatory power, and
29 predictable sequence variation, is the natural choice for many metabarcoding applications. However, for
30 targeting specific taxonomic groups in environmental samples, the utility of COI has yet to be fully scrutinised.

31
32 2. Here, by using a case study of marine and freshwater fishes from the British Isles, we quan-
33 tify the *in silico* performance of twelve primer pairs from four mitochondrial loci—COI, cytochrome *b*, 12S
34 and 16S—in terms of reference library coverage, taxonomic discriminatory power and primer universality.
35 We subsequently test *in vitro* four primer pairs—three COI and one 12S—for their specificity, reproducibility,
36 and congruence with independent datasets derived from traditional survey methods at five estuarine and
37 coastal sites around the English Channel and North Sea.

38
39 3. Our results show that for aqueous extra-organismal DNA at low template concentrations, both
40 metazoan-targeted and fish-targeted COI primers perform poorly in comparison to 12S, exhibiting low levels
41 of reproducibility due to non-specific amplification of prokaryotic and non-target eukaryotic DNAs.

42
43 4. An ideal metabarcode would have an extensive reference library upon which custom primers
44 could be designed, either for broad assessments of biodiversity, or taxon specific surveys. Such a database
45 is available for COI, but low primer specificity hinders practical application, while conversely, 12S primers
46 offer high specificity, but lack adequate references. The latter, however, can be mitigated by expanding
47 the concept of DNA barcodes to include whole mitochondrial genomes generated by genome-skimming
48 existing tissue collections.

49
50 [Keywords: 12S, COI, eDNA, Environmental DNA, metabarcoding, primer design.]

51 INTRODUCTION

52 DNA barcoding and metabarcoding techniques are now established and indispensable tools for the assessment
53 and monitoring of past and present ecosystems (Leray and Knowlton, 2015; Pedersen et al., 2015; Thomsen and
54 Willerslev, 2015; Valentini et al., 2016), and are being increasingly incorporated into policy and management
55 decisions (Kelly et al., 2014b; Rees et al., 2014; Mariani et al., 2015; Hering et al., 2018). A remarkably wide
56 range of biological substrates can now be sequenced to identify presence of a particular species or reconstruct
57 communities, and can include restaurant sushi meals (Vandamme et al., 2016), deep sea sediments (Guardiola
58 et al., 2015), permafrost ice cores (Willerslev et al., 2003), terrestrial insect collections (Ji et al., 2013), animal
59 faeces (Kartzinel et al., 2015) and seawater samples (Thomsen et al., 2012a).

60 The term “DNA metabarcoding” encompasses two distinct methodologies: (i) bulk sample metabarcoding,
61 which is the direct amplification of a concentrated mixture of organisms, from for example, plankton (Clarke
62 et al., 2017), arthropods (Yu et al., 2012) or gut material (Leray et al., 2013); or (ii) environmental DNA
63 (eDNA) metabarcoding, which is indirect amplification via extra-organismal DNA in water, sediments, or
64 soils (Taberlet et al., 2012). This latter methodology involves first isolating and concentrating DNA using
65 filters, rather than homogenising entire organisms or parts of organisms (Yu et al., 2012; Spens et al., 2017;
66 Macher et al., 2018). The detection of macrobial fauna such as vertebrates and insects using aquatic eDNA
67 has been recognised as a highly sensitive survey technique and a key use-case of metabarcoding (Rees et al.,
68 2014; Valentini et al., 2016). However, DNA from environmental samples such as seawater is likely to be
69 degraded (Collins et al., 2018), and also have a significant quantity of co-extracted microbial DNA that may
70 co-amplify with the targeted metazoan DNA molecules (Stat et al., 2017; Andújar et al., 2018).

71 Early eDNA metabarcoding studies targeting fishes used the cytochrome *b* gene (Minamoto et al., 2012;
72 Thomsen et al., 2012b,a), but more recent studies have used the 12S ribosomal rRNA locus (Kelly et al.,
73 2014a; Hänfling et al., 2016; Port et al., 2016; Stoeckle et al., 2017; Yamamoto et al., 2017; Ushio et al.,
74 2018), and also 16S rRNA (Shaw et al., 2016; Berry et al., 2017; Bylemans et al., 2018; Jeunen et al., 2018;
75 Stat et al., 2018). Various regions of 12S have been proposed as metabarcoding markers, including a ca. 63
76 bp fragment (Valentini et al., 2016), a ca. 106 bp fragment (Riaz et al., 2011; Kelly et al., 2014a), and a ca.
77 171 bp fragment (Miya et al., 2015). Modified versions of some of these primers have been published by
78 Taberlet et al. (2018). Ribosomal genes such as 12S and 16S offer the advantage of conserved priming sites
79 (Deagle et al., 2014; Valentini et al., 2016), and amplification across a broad range of fish taxa (Miya et al.,
80 2015; Bylemans et al., 2018). However, taxonomic resolution can be low (Miya et al., 2015; Hänfling et al.,
81 2016; Andruszkiewicz et al., 2017), with relatively short length ribosomal markers being unable to distinguish
82 commercially important species of the cod family Gadidae, for example (Thomsen et al., 2016). A problem
83 for studies using ribosomal markers are the reference libraries, which are usually poorly populated, and often
84 have to be developed for each project on an *ad hoc* basis (Miya et al., 2015; Thomsen et al., 2016; Stoeckle
85 et al., 2017). Assembling reference libraries for ribosomal genes is further complicated by frequently-used
86 primer sets amplifying different regions, with any two given 12S references taken from GenBank, for example,
87 not necessarily being homologous.

88 For animals, the primary DNA barcode is the 5' “Folmer” region of COI, the cytochrome *c* oxidase subunit
89 I gene (Folmer et al., 1994; Hebert et al., 2003). In comparison to ribosomal markers, the advantages of
90 COI are high interspecific variability (Ward, 2009), an extensive reference database (BOLD; Barcode of Life

91 Database; [Ratnasingham and Hebert, 2007](#)), and due to the protein-coding constraints of the gene, more
92 straightforward bioinformatic procedures such as alignment and denoising ([Andújar et al., 2018](#)). Inside the
93 5' Folmer fragment, multiple primer sets have been developed, targeting shorter regions in the 100–400 bp
94 range. These are more suitable than a full length barcode (ca. 658 bp) for analyses of degraded DNA, or for
95 sequencing on short read platforms such as Illumina ([Leray et al., 2013](#); [Shokralla et al., 2015](#); [Elbrecht and
96 Leese, 2017](#)). However, due to its variability, finding conserved priming regions within the Folmer fragment is
97 difficult, and concerns have been raised about the suitability of some COI primers in terms of species-specific
98 primer-template mismatches, which can result in inefficient, biased amplifications that may hinder quantitative
99 analyses ([Deagle et al., 2014](#)). Addressing this problem with bias requires incorporating a high degree of
100 degeneracy into COI primers ([Leray et al., 2013](#); [Marquina et al., 2019](#)), particularly by the use of multiple
101 inosine sites ([Shokralla et al., 2015](#); [Elbrecht and Leese, 2017](#); [Wangenstein et al., 2018](#)). Despite these issues,
102 [Andújar et al. \(2018\)](#) argue that COI should be the standard marker for metabarcoding, and COI markers are
103 increasingly being used for eDNA metabarcoding ([Bakker et al., 2017](#); [Kelly et al., 2017](#); [Stat et al., 2017](#);
104 [Jeunen et al., 2018](#); [Macher et al., 2018](#); [Singer et al., 2019](#)). However, studies comparing the efficacy of
105 different primer sets have done so in a bulk-sample metabarcoding context ([Clarke et al., 2017](#); [Elbrecht and
106 Leese, 2017](#)), or have compared only ribosomal markers for vertebrate eDNA applications ([Bylemans et al.,
107 2018](#)). Therefore, there lacks a clear assessment of how degenerate COI primers compare to 12S and 16S
108 rRNA when used on low-template-concentration environmental samples where non-target DNA molecules
109 are found in abundance.

110 Given the importance of marker choice in metabarcoding studies ([Alberdi et al., 2018](#)), and the need to
111 thoroughly scrutinise the utility of COI in comparison with the widely used ribosomal markers ([Deagle et al.,
112 2014](#); [Andújar et al., 2018](#)), we use a case study of fishes from the British Isles—a well studied and important
113 group in terms of ecosystem health and human food security—to ask whether COI primer sets can be used for
114 eDNA metabarcoding of aquatic vertebrates, and how they compare to alternative 12S, 16S and cytochrome *b*
115 markers. We survey a range of published primer sets both *in silico* and *in vitro*, including metazoan-targeted
116 COI primers with high levels of degeneracy, and novel fish-targeted COI primers with reduced degeneracy.
117 Using *in silico* methods we assess a number of factors: (i) the reference database coverage for the individual
118 fragments, i.e. how many species and individuals of each species are represented in public databases; (ii) the
119 taxonomic discrimination of each fragment, i.e. is each unique DNA sequence unambiguously associated with
120 a single species name; and (iii) the universality of the primer set, i.e. are all species of the target taxonomic
121 group predicted to amplify equally well. Then, we test using a series of water samples taken from locations
122 with corresponding data from traditional fish survey methods, three COI primer sets against a best performing
123 alternative set, as based upon the results of the *in silico* analyses. By PCR amplifying and sequencing these
124 water samples we compare: (i) the specificity of the primer set, i.e. the proportion of the reads that came from
125 the target taxonomic group; (ii) the power of the primer set, i.e. the total species richness estimated; (iii) the
126 reproducibility of the primer set, i.e. are the same species consistently represented in replicate water samples
127 and PCRs; and (iv) the congruence of the primer set, i.e. are the same species detected in the traditional
128 surveys as the eDNA surveys.

129 METHODS

130 *In silico* analyses

131 **Reference library construction**

132 A list of fish species recorded from the marine and freshwater environments of the British Isles was
133 compiled from three sources: (i) the Global Biodiversity Information Facility (<https://www.gbif.org>; *rg-*
134 *bif v1.1.0*; Chamberlain and Boettiger, 2017); (ii) FishBase (<https://www.fishbase.org>); and (iii) the Eu-
135 ropean Water Framework Directive United Kingdom Technical Advisory Group list of transitional fish
136 species (<https://www.wfduk.org/resources/transitional-waters-fish>; Annex 1). These species were then cross-
137 referenced for all synonyms using *rfishbase v3.0.0* (Boettiger et al., 2012). The subsequent list of valid
138 species names and all their synonyms was then searched using *rentrez v1.2.1* (Winter, 2017) against NCBI
139 GenBank release 230 (nucleotide database; <https://www.ncbi.nlm.nih.gov/nucleotide/>) for any of the following
140 terms: “COI, 12S, 16S, rRNA, ribosomal, cytb, COI, cox1, cytochrome, subunit, COB, CYB, mitochondrial,
141 mitochondrion”. The Barcode of Life Database BOLD (<http://www.boldsystems.org/>) was also searched for
142 the same species using *bold v0.8.6* (Chamberlain, 2018).

143 Hidden Markov models of the alignments of each primer set were then constructed using *HMMER*
144 *v3.1b2* (<http://hmmerr.org/>; Eddy, 1998) and the fish mitochondrial genome database (<http://mitofish.aori.u->
145 [tokyo.ac.jp/](http://mitofish.aori.u-tokyo.ac.jp/); Iwasaki et al., 2013). These profiles were used to extract homologous regions of nucleotides
146 from the total mitochondrial data obtained from the GenBank and BOLD searches. The resulting sequences
147 were then annotated with metadata using *traits v0.4.2* (Chamberlain et al., 2019). A phylogenetic quality
148 control step was then carried out by aligning the sequences in *MAFFT v7.271* (Katoh and Standley, 2013)
149 and constructing a maximum likelihood tree using *RAxML v8.2.12* (Stamatakis et al., 2008). Sequences with
150 putatively spurious annotations—i.e. those indicative of misidentifications—were filtered out if the following
151 criteria were met: (i) individual(s) of species *x* being identical to or nested within a cluster of sequences
152 of species *y*, but with other individuals of species *x* forming an independent cluster; and (ii) the putatively
153 spurious sequences coming from a single study, while the putatively correct sequences of species *x* and *y*
154 coming from multiple studies. Records flagged by NCBI as “unverified” were also omitted.

155 **Primer design**

156 We designed two new COI metabarcoding primers targeting fishes (Table 1): “SeaDNA-short” and “SeaDNA-
157 mid”, which share a forward primer, and are internal to the Folmer fragment. The new primer pairs were
158 designed manually in *Geneious v8.8.1* (Kearse et al., 2012) using the same fish mitochondrial genome dataset
159 as described above, with the assistance of *Primer3* (Untergasser et al., 2012) and the sliding window functions
160 in *spider v1.3.0* (Boyer et al., 2012; Brown et al., 2012). The primers were tested on a range of fish tissue
161 extractions from elasmobranchs and actinopterygians, and produced strong clean PCR amplicons of the
162 expected size.

163 **In silico PCR and taxonomic discrimination**

164 Primers were evaluated using a subset of 955 unique sequences from 184 species obtained in the British Isles
165 fish reference library construction step, for which full mitochondrial genomes were available. Twelve primer
166 pairs were chosen for the *in silico* PCRs, representing COI, cytochrome *b*, ribosomal 12S and ribosomal 16S
167 (Table 1). *MFEprimer v2.0* (Qu et al., 2012) was used to perform the *in silico* PCR on the untagged primers.

168 Amplification universality was estimated using the Primer Pair Coverage (PPC) statistic from *MFEprimer*,
169 where $PPC = \frac{Fm}{Fl} \times \frac{Rm}{Rl} \times (1 - CVfr)$, with *Fl* and *Rl* the length of the forward and reverse primers, and *CVfr*
170 the coefficient of variability of matched lengths *Fm* and *Rm* to the template. Therefore, a PPC value of 100%
171 indicates complete binding of both primers to a template. The highest PPC value was then selected for each
172 species, and averaged over all species to provide the PPC for each primer set. Predicted non-amplifications
173 with a default 5 bp 3' binding stability of $> 0\Delta G$ were set to a PPC of 0%. In order for sufficient RAM to
174 be available to complete the analysis of the highly degenerate Leray-XT primer set, the inosine sites were
175 simplified to double-base ambiguities. This was achieved by choosing the most frequent base combination
176 in the mitogenome alignment. None of the altered inosine sites were within 8 bp of the 3' end of the primer
177 (Table 1).

178 Taxonomic discrimination (= resolution) was assessed first using all available species from the British
179 Isles fish reference library for each primer set individually, and then secondly on a subset of species for which
180 sequences were present for all of the primer sets. Discrimination as a proportion of the total number of species
181 was calculated following [Ficetola et al. \(2010\)](#): “A taxon unambiguously identified by a primer pair owns a
182 barcode sequence associated to this pair that is not shared by any other taxa”.

183 **Primer evaluation *in vitro***

184 ***Field sites and traditional fish survey***

185 Five locations in the United Kingdom were surveyed for fishes using eDNA and traditional methods between
186 October and November of 2016. These included: the River Tees, County Durham (54.631327,-1.164447);
187 two sites within the River Esk estuary, North Yorkshire (54.491633,-0.611833; 54.48975,-0.612617); the
188 River Test, Hampshire (50.901563,-1.440836); and Whitsand Bay, Devon (50.329616,-4.243751). The former
189 four are estuarine sites, while the latter is an inshore coastal area, approximately 1 km from shore. Fish
190 sampling in the River Esk estuary was done by duplicate fyke nets (Esk-fyke) and duplicate beach-seine nets
191 (Esk-seine), in different locations. At the River Tees sampling site, duplicate beach-seine netting and two
192 shallow beam trawls were carried out. The River Test site comprised a 24 h fish impingement survey conducted
193 at Marchwood Power Station. Whitsand Bay was surveyed by four otter trawls, as described in [McHugh et al.](#)
194 (2011). The variety of fishing techniques used in the different sampling locations are part of the currently
195 ongoing fish monitoring programmes implemented by local collaborating organisations (Environment Agency,
196 PISCES Conservation Ltd., Marine Biological Association). Further details are presented in Supporting
197 Information.

198 ***Water processing and DNA extraction***

199 Three 2 L water sample replicates per site were collected immediately prior to the traditional fish survey
200 commencing, using Nalgene HDPE collection bottles pre-sterilised with a 10% bleach solution. Water was
201 pre-strained with a 250 μm nylon mesh filter to remove debris, if required. After collection, the water samples
202 were put into individual sterile plastic bags, and stored in an ice box while being transported back to the
203 laboratory. Within five hours, each 2 L sample was filtered through an 0.22 μm Sterivex-GP PES filter (Merck
204 Millipore) using a 100 mL polypropylene syringe or a peristaltic pump, and cleared of water. When the full 2
205 L could not be passed due to filter clogging, the volume of water was recorded. After filtration, the filters
206 were stored at -20°C . DNA was extracted from the filters using the DNeasy PowerSoil DNA Isolation Kit

207 (MoBio/Qiagen), following the manufacturers' protocol, with the addition of an initial 2 h agitation step to
208 promote the release of DNA from the filter, during which the filter membranes were placed in tubes with lysis
209 buffer C1 and garnet beads from the PowerWater Isolation kit and shaken at 65°C. Filtration blank controls
210 were processed in parallel. All processing was carried out in dedicated eDNA extraction laboratories, and
211 equipment and surfaces were regularly cleaned using a 10% bleach solution. The eDNA extraction, pre-PCR
212 preparations and post-PCR procedures were carried out in separate rooms.

213 **PCR and library preparation**

214 Four primer sets were selected to go forward for *in vitro* testing: three COI primer sets (Leray-XT, SeaDNA-
215 short, SeaDNA-mid), and one best-performing primer set from the *in silico* analysis (12S MiFish-U). All
216 PCR amplifications were done in duplicate reactions each with a unique 7/8-mer oligo-tag barcode, differing
217 by at least three bases (Guardiola et al., 2015). In order to increase variability of the amplicon sequences,
218 a variable number (two, three or four) of fully degenerate positions (Ns) were added at the 5' end of the
219 oligo tags (Wangensteen et al., 2018). For PCR amplification with the newly designed SeaDNA-short and
220 SeaDNA-mid primers, a two-step protocol was used, first using untagged primers, then tagged primers in
221 a second PCR round. The reaction for the first PCR step included 10 μ L AmpliTaq Gold 360 Master Mix
222 (ThermoFisher), with 1 μ L of each 5 μ M forward and reverse primer, 0.16 μ L of bovine serum albumin
223 and 10 ng of purified DNA in a total volume of 20 μ L per sample. Thermocycling profile for the first step
224 included an initial denaturation at 95°C for 10 minutes, then 40 cycles of 94°C for 30 sec, 47°C for 45 sec and
225 72°C for 30 sec, and then a final extension of 72°C for 5 minutes. The profile for the second PCR step was
226 identical, except for the annealing temperature being 50°C instead of 47°C. Amplifications were assessed by
227 electrophoresis on a 1.5% agarose gel, and the field and laboratory controls were checked for the presence of
228 amplicons. Between the first and second PCR step, amplicons were purified using MinElute PCR purification
229 columns (QIAGEN) and diluted by a factor of ten prior to being used as a template for the second PCR. After
230 the second PCR, all tagged amplicons were pooled by marker, purified again using MinElute columns and
231 eluted into a total volume of 45 μ L, in order to concentrate the amplicons approximately 15 times. For 12S
232 MiFish and Leray-XT we used a one-step procedure with tagged PCR primers, with PCR cycling conditions
233 following Miya et al. (2015) and Wangenstein et al. (2018), respectively. Reagents and volumes were the
234 same as for the two-step protocol.

235 Libraries (one for each primer set) were built using the PCR-free NEXTflex library preparation kit (BIOO
236 Scientific). The libraries were quantified using the NEBNext qPCR quantification kit (New England Biolabs)
237 and spiked with with 1% PhiX (Illumina). The libraries were sequenced on an Illumina MiSeq platform,
238 using V3 chemistry (2 \times 75 bp paired-end) for the SeaDNA-short library, which was run along with two other
239 libraries from unrelated projects. For the MiFish-U and SeaDNA-mid libraries, V2 chemistry (2 \times 150 bp
240 paired-end) was used, and these were sequenced in the same run. The Leray-XT library was run using V2
241 chemistry (2 \times 250 bp paired-end) along with another library from an unrelated project.

242 **Bioinformatic processing**

243 Raw sequencing data were converted to fastq format using *bcl2fastq* v2.20 (<https://support.illumina.com/sequencing/sequencing-conversion-software.html>). The remaining bioinformatic steps were carried out using *cutadapt* v2.3 (Martin,
244 2011) and *dada2* v1.10.1 (Callahan et al., 2016). Because a PCR-free library preparation kit was used,
245

246 adapters could have been ligated to either the 5' or the 3' end of the amplicon, and in order to take advantage
247 of the Illumina error profiling in the *dada2* denoising step, the sense- and antisense-orientated sequences were
248 first isolated and processed independently. This was achieved using *cutadapt* by filtering the R1 fastq files for
249 reads with the forward PCR primer, and then for those with the reverse PCR primer. The reads were then
250 demultiplexed by tag, followed by primer and adapter trimming. Quality trimming was carried out in *dada2*
251 using default settings, but with read truncation length “truncLen” determined to give an approximate 30 bp
252 overlap between forward and reverse reads. The reads were then denoised, dereplicated, merged, cleaned
253 of chimaeras and reorientated, using the *dada2* workflow. Our reference library sequences for each primer
254 set were used as priors to avoid low abundance but valid sequences being discarded during denoising. A
255 homology filter was then implemented by aligning the ASVs against a hidden Markov model of the expected
256 fragment using *HMMER hmmsearch*, and the non-homologous reads discarded.

257 Taxonomy assignment of the amplicon sequence variants (ASVs) produced by *dada2* was carried out
258 using a multi-step procedure, incorporating distance-based and phylogenetic methods. First, a preformatted
259 “nt” blast database was downloaded from NCBI (<ftp://ftp.ncbi.nlm.nih.gov/blast/db/v5>; 21 March 2019). Each
260 ASV sequence was then locally blasted against this database using *blastn* v2.9.0 (‘-task blastn -evalue 1000
261 -word_size 11 -max_target_seqs 500’), and the results filtered to obtain a rough taxonomic classification based
262 on the best-scoring blast hit. Next, a more stringent procedure was carried out, with the putative fish sequences
263 extracted from this initial blast result subjected to a second *blastn* search, this time using our curated reference
264 library of British Isles fishes as the blast database (same settings as the “nt” search but with ‘-word_size 7’).
265 The same reads were then run through the Evolutionary Placement Algorithm (*EPA-ng* v0.3.5, *gappa* v0.2.0;
266 [Barbera et al., 2018](#); [Czech and Stamatakis, 2018](#)). Species name(s) were assigned based on either of the
267 following rules: (i) species-level EPA placement same as the best scoring blast hit, with an aligned match
268 length of $\geq 90\%$ of the modal length of the fragment, and an identity of $\geq 97\%$; or (ii) highest likelihood EPA
269 placement same as the best scoring blast hit, with an EPA probability $\geq 90\%$ and blast identity $\geq 90\%$. Rule
270 (i) finds assignments that are congruent between both the EPA and blast methods, but rejects assignments
271 with low similarity and short match lengths. Rule (ii) allows for dissimilar hits, but only ones that have a
272 high phylogenetic probability, and which are usually indicative of low abundance variants with errors. Our
273 prior knowledge of the expected fish fauna of the sites was used to set these cut-off values, with the aim of
274 conservatively minimising false positive assignments. The fish reads were also summarised by OTU clustering
275 using *Swarm* v2.2.2 ([Mahé et al., 2015](#)), with $d = 1$ and the “fastidious” option enabled. This step permitted
276 an evaluation of possible misassigned and unassigned species.

277 RESULTS

278 *In silico* analyses

279 A total of 531 species were identified as part of the United Kingdom marine and freshwater fish fauna. Of
280 these, 176 names were flagged as “common” species, having been identified as relatively widespread marine
281 or freshwater taxa that are likely to be encountered during survey work of coastal and inland habitats ([Kottelat
282 and Freyhof, 2007](#); [Henderson, 2014](#)). The remainder were mostly highly localised species, deep water
283 offshore species, or rare migrants. The combined reference library for all primer sets, after cleaning, duplicate
284 removal and quality control, comprised 43,366 sequences from 491 total species, and 25,799 sequences from

285 172 common species.

286 In terms of reference database coverage for individual primer sets (Table 2), COI primers had the greatest
287 number of reference sequences at 23,911–24,058, covering 91% of species. The “Minamoto-fish” cytochrome
288 *b* set had 15,405 sequences and a species coverage of 65%. Of the ribosomal primer sets, the “Berry-fish”
289 16S set had the greatest number of sequences at 4,089, with species coverage at 77%. Among the 12S
290 sets, the “Riaz-V5” primers had the greatest number of sequences (2,416; species coverage 69%), while
291 the “Valentini-tele01” set had the fewest sequences (1,699; species coverage 51%). The “MiFish” primers
292 and their variants (MiFish-U/E, Taberlet-tele02, Taberlet-elas02) had 1,904 sequences, and a coverage of
293 61%. Per species, the average number of reference sequences was greatest for the COI primer sets (mean
294 49–50; median 24), followed by cytochrome *b* (mean 45; median 7), 16S (mean 9.9; median 4), and then 12S
295 (mean 5.9–6.6; median 2–3). When only the subset of common species was considered, the species coverage
296 increased for all primer sets, as did the average number of sequences per species (Table 2).

297 In terms of taxonomic discrimination of the fragments obtained from each primer set (Table 2), the
298 proportion of British Isles fish species where all individuals could be unambiguously identified was greatest
299 for the Leray-XT COI fragment at 95%, while the shorter SeaDNA-mid and SeaDNA-short COI fragments
300 resolved 91% and 87% respectively. The cytochrome *b* fragment discriminated 91%. The MiFish fragment had
301 the greatest discrimination among the ribosomal primer sets at 93%, with the Berry-fish 16S, Valentini-tele01,
302 and Riaz-V5 pairs having lower rates (89%, 86%, and 79% respectively). When a standardised dataset of
303 species common to all primer sets ($n = 88$) was used, the overall pattern remained similar (Table 2).

304 In terms of primer universality as estimated by *in silico* PCR for British Isles fish species with comparable
305 data available for all markers ($n = 184$; Table 2), the 12S primer sets targeting actinopterygians had a higher
306 mean PPC than all other markers, at between 77.1% (Valentini-tele01) and 92.2% (Riaz-V5), compared to
307 between 19.1% (cytochrome *b*) and 50.9% (16S). The best performing COI marker for actinopterygians
308 (SeaDNA-short) had a PPC value of 34.5%. For elasmobranchs, three 12S primer pairs had the highest
309 mean PPC values, with Taberlet-elas02 at 83.6%, Valentini-tele01 at 68.2%, and MiFish-E at 55%. The
310 12S Riaz-V5 primers, the cytochrome *b* primers, and the 16S primers, had the lowest PPC values (11.2%,
311 20.4% and 0% respectively), while the COI primers had PPC values between 21.5% (SeaDNA-short) and
312 39% (simplified Leray-XT). These patterns remained when only common species were compared (Table 2).

313 ***In vitro* analyses**

314 Total reads from Illumina sequencing (Table 3) varied between 3.4 million (12S MiFish-U) and 14.3 million
315 (COI SeaDNA-mid). After bioinformatic processing, the proportions of reads retained were 46% (COI
316 SeaDNA-short), 54% (COI Leray-XT), 61% (COI SeaDNA-mid) and 63% (12S MiFish-U). Mean cleaned
317 reads recovered per sampling event (triplicate water samples, duplicate PCR tags; $n = 6$) were: 107,458 (SD =
318 46,924) for Leray-XT; 290,104 (SD = 118,592) for SeaDNA-mid; 135,804 (SD = 44,993) for SeaDNA-short;
319 and 71,912 (SD = 13,682) for 12S MiFish-U. Supporting Figure S1 shows distributions of read depths per
320 sample for each site and primer set. The 12S MiFish-U primers provided the greatest proportion of chordate
321 and fish reads (100% and 76% of cleaned reads, respectively), resulting in more than 1.6 million putative
322 fish reads and 156 fish ASVs. From these fish reads, 96% were assigned to 41 species and 67 *Swarm* OTU
323 clusters. A total of 73,377 fish reads comprising 18 *Swarm* OTUs could not be assigned, and in addition to
324 PCR and sequencing artefacts, these likely represent at least eight species not present in the reference library

(Supporting Table S1). For the COI primer sets, chordate reads comprised between 0.2% (Leray-XT) and 6% (SeaDNA-short) of the total cleaned reads, with between 0.1% and 5% putative fish reads comprising between 22 (Leray-XT) and 29 (SeaDNA-short) assigned species. Between 42% (Leray-XT) and 85% (SeaDNA-short) of the putative fish reads were unassigned to species. The non-chordate reads were inferred from the preliminary blast search to consist of DNA from other metazoans (4–10%) and eukaryotes (41–83%), or bacteria (17–59%).

Per sampling location the 12S MiFish-U primer set detected a consistently greater number of total species across sites than the COI markers, at between 2.2 (River Test) and 2.6 (Whitsand Bay) fold higher (Figure 1). The SeaDNA-short primers detected a greater number of species than both the SeaDNA-mid and Leray-XT primers, except at the River Tees site where SeaDNA-mid detected one more.

In terms of reproducibility (Figure 2), the 12S MiFish-U primer set showed a greater proportion of shared species—the top ten species by read abundance at each location—amplified across water sample and PCR replicates, with a 71% mean reproducibility over all sampling locations. The COI primer sets had mean reproducibility values of 36% (SeaDNA-short), 29% (SeaDNA-mid) and 12% (Leray-XT).

When compared to traditional survey methods—with the freshwater species omitted from the eDNA results as they were not expected to be found on the traditional fish surveys of the estuarine and coastal habitats—the 12S MiFish-U primer set showed the greatest congruence (Figure 3), at between 15% (Whitsand Bay) and 54% (River Test). The COI primers were between 9% (Leray-XT) and 13% (SeaDNA-short) congruent overall. The MiFish-U primer set also amplified a greater number of marine/estuarine species to the traditional survey methods at all locations except for Whitsand Bay (26 versus 23 species). The COI primer sets amplified fewer marine/estuarine species than the traditional surveys in all cases, except for the SeaDNA-short primer set at the River Tees and River Esk sites. For each site survey, reads per species (eDNA survey) and individuals per species (traditional survey) are presented in Supporting Tables S2–S6.

DISCUSSION

A single metabarcoding marker for fishes?

Of arguably the greatest importance in the ability of metabarcoding to answer a particular question, is that of the choice of marker and primer (Deagle et al., 2014; Valentini et al., 2016; Clarke et al., 2017; Elbrecht and Leese, 2017; Alberdi et al., 2018). The ideal genetic marker for eDNA metabarcoding marker should be flexible, allowing different primer sets to target different taxonomic groups, but requiring only a single reference library. Each individual primer set must also be designed with the following qualities: (i) it must be universal, i.e. amplifying a large proportion of the target taxonomic group; (ii) it must be specific, i.e. it must not amplify other taxa at the expense of the target group; (iii) it must be unbiased, i.e. not preferentially amplifying a subset of the target group; (iv) it must be discriminatory, i.e. the DNA fragment recovered should differentiate at the appropriate taxonomic level for the question; and (v) it must be replete, i.e. associated with a reference library enabling identifications within the target taxonomic group. Here, we assess these characteristics for COI, cytochrome *b*, 12S, and 16S primer sets using the example of marine and freshwater fishes from the British Isles.

362 Which primers have the best reference library?

363 In terms of reference libraries, the COI primers were substantially better endowed than all other marker genes,
364 with between 1.6 times (cytochrome *b*) and 14 times (Valentini-tele01) more public sequence data available
365 for all species. This was also reflected in the common species coverage, at up to 97% for COI. The 16S (95%),
366 cytochrome *b* (81%), and 12S Riaz-V5 libraries (81%) were also well developed for common species, but
367 coverage for other 12S primer sets was lower, at 56–62%. A reference library with broad taxonomic depth
368 will allow inferences beyond a comparison of anonymous MOTUs, thereby leveraging the wealth of scientific
369 information that a taxonomic name brings with it (Ward et al., 2009). Deep coverage in the COI reference
370 library—i.e. the number and geographic distribution of sequences per species—also has advantages in terms
371 of potential for population level assignments, and for flagging spuriously identified sequences; due to the
372 lesser weight of evidence from the low numbers of sequences, misidentifications were harder to confirm for
373 12S during the quality control step. Furthermore, in terms of voucher specimen and location data etc, much of
374 the ribosomal data on GenBank are not validated to the same standard as COI data on BOLD are (Ward et al.,
375 2009). However, it is important to remember that despite the success of 15 years of the DNA barcode initiative
376 producing COI coverage spanning the majority of northern European fish species, the BOLD database still
377 remains seriously underdeveloped for many other taxonomic groups such as marine invertebrates (Bucklin
378 et al., 2011; Leray and Knowlton, 2016).

379 Which primers best discriminate species?

380 In terms of the discriminatory power for our dataset of British Isles fish species, all primer sets gave a
381 resolution above 90% except for SeaDNA-short (COI), Valentini-tele01 (12S), Riaz-V5 (12S) and Berry-fish
382 (16S). Predictably, the longer COI fragments resolved more species than the shorter ones, at 95% for the 313
383 bp Leray-XT and 87% for the 55 bp SeaDNA-short fragment. The 12S primers did not show this pattern
384 as clearly, with the shorter Valentini-tele01 fragment having a better taxonomic resolution (86%) than the
385 longer Riaz-V5 fragment (79%); the longest, MiFish-U/E and Taberlet-tele02/elas02 primers, had the greatest
386 species resolution at 93%. While discriminatory power may depend on the range of species in that particular
387 library, the observed patterns held up when a dataset of sequences that were shared for all primer sets was
388 used. Discriminatory power also tended to remain the same or increase when only the common species were
389 considered, most likely because rare but genetically similar congeners were excluded.

390 Which primers are most universal?

391 Primer universality as estimated by *in silico* PCR varied greatly. Our results show that the metabarcoding
392 primers targeting protein-coding genes—COI and cytochrome *b*—are likely to exhibit a greater degree of
393 species-level primer bias (i.e. lower universality) than ribosomal 12S and 16S, as indicated by the lower
394 mean PPC values; a mean PPC of 96% was estimated for common actinopterygian species amplified with
395 the Riaz-V5 and Taberlet-tele02 primers. Previous studies have also reported or predicted less primer bias
396 with rRNA targets than protein coding ones (Clarke et al., 2014; Deagle et al., 2014; Elbrecht et al., 2016;
397 Marquina et al., 2019). It is also important to note again that due to the high level of degeneracy the Leray-XT
398 primers were simplified to overcome RAM limitations of the analysis, and therefore the value presented is
399 likely to be an underestimate of their true potential, as highly degenerate COI primers have been shown to
400 reduce bias substantially (Marquina et al., 2019).

401 Regarding higher level taxonomic bias, for the 12S and 16S primers tested here, no set except Valentini-
402 tele01 appeared suited to amplify actinopterygians and elasmobranchs equally. The COI primers were,
403 however, relatively unbiased in regard to higher taxonomic group. The MiFish primers and the [Taberlet et al.](#)
404 (2018) variants of the same sets were both published with actinopterygian (MiFish-U) and elasmobranch
405 (MiFish-E) versions, due to a number of mismatches in the conserved regions ([Miya et al., 2015](#)). Unsur-
406 prisingly, both of these performed substantially better for their respective taxa. The [Taberlet et al. \(2018\)](#)
407 primers were also predicted here to exhibit reduced species-level primer bias compared to the original MiFish
408 versions, for both elasmobranchs and actinopterygians.

409 Many studies computationally predict primer amplification by the number of mismatches between primer
410 and template (e.g. [Riaz et al., 2011](#)), or by the number of mismatches and their type and position (e.g. [Elbrecht](#)
411 [et al., 2017](#)), but often do not fully consider the thermodynamics of a primer-template reaction. We used
412 the thermodynamics-based PCR simulation implemented in MFEprimer ([Qu et al., 2012](#)), but regardless of
413 whether this method is more realistic or accurate than alternative methods, it is important to remember that
414 these are predicted amplifications, and were used here to compare relative performances between primer sets.
415 Therefore, the lower values estimated do not represent amplification failure *per se*, but rather are indicative
416 of increased bias associated with that primer set ([Deagle et al., 2014](#)). For example, the standard COI DNA
417 barcode primers for fishes (Ward-barcode) had a very low PPC, but these are tried-and-tested primers for
418 amplifying a wide range of fish taxa in standard PCR for Sanger sequencing ([Ward et al., 2005](#)). The use
419 of mock communities is an important step in quality controlling an assay if primer bias is suspected ([Piñol](#)
420 [et al., 2015](#); [Elbrecht and Leese, 2017](#); [Bista et al., 2018](#)), but *in silico* PCR has been demonstrated to be an
421 effective proxy in its absence ([Clarke et al., 2014](#)).

422 We used the results of our *in silico* analyses to inform our choices for the *in vitro* experiments. All COI
423 primer sets were selected for testing *in vitro* because of the advantages in terms of reference library and
424 taxonomic discrimination. We chose only one 12S set for comparison, and here we chose the MiFish-U primer
425 pair because this pair had better predicted universality for actinopterygians and more reference sequences
426 available than the Valentini-tele01 primers, and greater taxonomic discrimination than the Riaz-V5 primers.
427 Due to the better predicted universality of the Taberlet-tele02 primer set compared to MiFish-U, these would
428 have been chosen had they been publicly available at the time the experiment was implemented. Despite the
429 well developed reference libraries and good taxonomic discrimination, we did not select cytochrome *b* or 16S
430 because of the lower predicted universality of these primers in comparison to 12S.

431 **Which primers are the most specific?**

432 Despite having the fewest total raw reads, the MiFish-U primer set produced the greatest number and
433 proportion of usable fish reads (76% of processed reads, 48% of raw reads), the greatest overall species
434 richness (41 species), and the greatest proportion of fish reads that were assigned to species (96%). The COI
435 primers amplified a very low proportion of chordate and fish reads compared to the overall sequencing depth
436 (maximum 5% of cleaned reads were fishes). The majority of the SeaDNA-short and SeaDNA-mid reads were
437 estimated by preliminary blast search to have come from bacteria or non-metazoan eukaryotes (86–90%).

438 That the highly degenerate Leray-XT primers produced a low proportion of fish reads is unsurprising
439 given that previous studies on environmental samples using degenerate COI primers have demonstrated that
440 they can amplify widely beyond their target taxa, and can produce large proportions of unassigned reads ([Lim](#)

441 et al., 2016; Stat et al., 2017; Macher et al., 2018; Singer et al., 2019). The proportion of bacterial reads
442 are generally lower when metabarcoding bulk organismal samples, however, with most reads belonging to
443 metazoans (Leray and Knowlton, 2015; Macher et al., 2018; Wangensteen et al., 2018). More surprising was
444 the poor specificity of the SeaDNA-short and SeaDNA-mid primers, which were designed to target fishes, and
445 with minimal degeneracy. These data are, however, consistent with those of an analysis of shark diversity by
446 Bakker et al. (2017), who used COI mini-barcode primers designed on sharks, and reported a similar level of
447 non-specific amplification.

448 The cause of this non-specific amplification is likely to be the extensive homoplasy (nucleotide con-
449 vergence) apparent in the mutationally saturated COI gene and its homologs. Siddall et al. (2009) demon-
450 strated that metazoan-targeted COI primers are likely to co-amplify many marine prokaryote groups—
451 gammaproteobacteria being a particularly diverse and abundant lineage (Sunagawa et al., 2015)—thereby
452 compromising the specificity of these primer sets. Optimisation of PCR protocols or library preparation
453 methods may increase specificity of the assay (Siddall et al., 2009), but it is probably unlikely that it can
454 increase to a level that makes the proportion of usable reads viable for eDNA metabarcoding of targeted
455 taxonomic groups. While this phenomenon was first observed in marine prokaryotes, studies on freshwater
456 and soil faunas have shown a similar pattern, also with large numbers of unassigned reads (Yang et al., 2014;
457 Lim et al., 2016).

458 **Which primers give the most reproducible results?**

459 The low number of usable fish reads for the COI primers is reflected in the reproducibility of the assays across
460 water sample and PCR replicates. For the most frequently amplified species at each site, the COI primers were
461 less consistent than 12S MiFish-U overall. Low quantities of template DNA and stochasticity in early PCR
462 cycles is a known factor in causing poor reproducibility (Leray and Knowlton, 2017; Alberdi et al., 2018;
463 Collins et al., 2018), and can be ameliorated by performing multiple PCR technical replicates (Ficetola et al.,
464 2015). We show that this effect is exacerbated when primer specificity is low and non-target organisms are
465 abundant, as is the case in highly diverse environmental samples such as seawater. For many applications
466 repeatability between assays or sampling sites is a requirement, such as the detection of an endangered or
467 invasive species (Grey et al., 2018). Our results, even considering only the top ten common species, show that
468 detectability can vary between sites with the same genetic marker, and that many more than two PCRs will be
469 required if the rare species are to be detected across multiple PCR and water sample replicates (Dopheide
470 et al., 2018).

471 Species richness estimates at all sampling sites were greatest with 12S MiFish-U, and this was despite
472 the deficiencies in the reference library, at only 61% species coverage. For example, species including the
473 European plaice (*Pleuronectes platessa*) and European flounder (*Platichthys flesus*)—both common fishes
474 present at all sampling locations—were missing from the reference library and therefore not represented when
475 comparing with the traditional fish surveys. Most of the large number of reads that were assigned to American
476 plaice, *Hippoglossoides platessoides* ($n = 198,445$), were likely misassigned and actually belong to European
477 plaice and flounder (Supporting Table S1). The Swarm OTU analysis showed a greater number of clusters
478 (67) than assigned species (41), also suggesting that some species missing from the reference library are
479 likely to have been misassigned. While a small number of the 73,377 unassigned 12S fish reads were low
480 abundance sequences derived from artefacts, almost all could be inferred by phylogenetic analysis

481 or by similarity to geographically disjunct congeners, to belong to at least eight species that were known to
482 be missing from the reference library (Supporting Table S1). Despite this major handicap, the 12S MiFish
483 primers remained superior to COI in terms of congruence with the traditional fish surveys, by recovering a
484 greater overlap of species in all cases. The 12S MiFish primers amplified more species than the traditional
485 surveys at all sites, except Whitsand Bay. This was mainly due to the underrepresentation of the fauna of that
486 site in the 12S reference library, with over half of the surveyed species absent from the library, and a higher
487 proportion of elasmobranchs (five species) than the other sites, which the MiFish-U primers fail to amplify.
488 Overall, no species that were recorded in the traditional surveys were missing from the COI reference libraries,
489 but eighteen species were missing from the 12S MiFish library (37%). The low numbers of species recorded
490 by the traditional surveys at the Esk and Tees sites in comparison to the Whitsand Bay and River Test sites, is
491 partly due to the inherently less diverse fauna of these northerly estuaries, as well as a reflection of the survey
492 techniques, with fyke and seine netting likely to detect fewer species than otter trawling (Whitsand Bay) or
493 a 24 h power station impingement (River Test). It should also be noted that there is no *a priori* assumption
494 that the eDNA and traditional survey data will be completely congruent, as most fish survey methods are
495 imperfect, sampling a moving target of diversity and abundance over difficult-to-define spatio-temporal points.
496 For example, eDNA can be transported in or out by tides, while some species are difficult to sample using
497 particular fishing gears, due to effects of size, behaviour or abundance. Therefore, overlap between eDNA
498 and traditional survey data is best interpreted as a relative measure between the primer sets.

499 CONCLUSIONS

500 While PCR-free methods are being actively investigated, it is clear that despite the limitations in quantification,
501 the majority of environmental metabarcoding will be based around amplicon sequencing, at least for the
502 medium term (Creer et al., 2016; Stat et al., 2017; Bista et al., 2018; Wilcox et al., 2018). Particularly important
503 for regulatory applications, or where researchers wish to compare results over time or between studies, some
504 degree of standardisation is desirable (Hering et al., 2018). Our results—and those of previous studies
505 using similar primer sets (Yang et al., 2014; Lim et al., 2016; Bakker et al., 2017; Stat et al., 2017; Jeunen
506 et al., 2018; Macher et al., 2018; Singer et al., 2019)—show that environmental metabarcoding for restricted
507 taxonomic groups using degenerate COI primers results in excessive volumes of “wasted” sequencing effort.
508 This co-amplification of prokaryotic and non-target eukaryotic DNAs and subsequent lack of specificity is
509 due to the nature of mutation patterns in COI (Siddall et al., 2009). Therefore, while we fully support the
510 arguments presented by Andújar et al. (2018) regarding the overall advantages of COI as a bulk-sample
511 metabarcoding marker, we find it difficult to recommend for metabarcoding environmental samples with low
512 target template concentrations and high microbial and plankton diversity, such as natural water bodies.

513 While the use of multiple primer sets and markers are probably required for a comprehensive view of total
514 biodiversity (Drummond et al., 2015; Stat et al., 2017), for specific taxonomic groups such as fishes a single
515 assay should be a feasible proposition. Unfortunately, no single 12S primer set was shown to be optimal for
516 eDNA fish surveys. The MiFish-U primer set—and *in silico*, the Taberlet et al. (2018) modified versions—
517 performed well in terms of specificity, discriminatory power, and reproducibility. Despite this, MiFish-U is
518 not universal for all fishes, because a separate MiFish-E assay is required to amplify elasmobranchs. The
519 MiFish reference library was also inadequate in this case, missing large numbers of common taxa. The

520 Valentini-tele01 primer set amplifies actinopterygians and elasmobranchs in a single assay, but suffers from
521 an even more poorly populated reference library than MiFish-U, and weaker taxonomic resolution. The
522 Riaz-V5 primers had the most complete reference library of the 12S primer pairs, but also do not amplify
523 elasmobranchs and have the poorest discriminatory power.

524 Because no single alternative primer set to COI will be optimal for all applications, it is clear that the
525 current DNA barcode reference libraries will need to be augmented with data from multiple mitochondrial
526 regions to enable their wider utility for vertebrate metabarcoding. However, rather than sequencing individual
527 12S regions on an ad hoc basis, a better solution is to generate whole mitochondrial genomes which can act as
528 an extended or linking barcode if sequenced from the same collection material (Coissac et al., 2016; Collins
529 and Cruickshank, 2014). Low coverage genome skimming techniques now produce high quality mitogenomes,
530 and are compatible with existing—frequently ethanol-based—tissue collections, and therefore will not require
531 the recollection of specimens (Gillett et al., 2014; Linard et al., 2016). Environmental DNA techniques could
532 potentially be the default survey methodology for aquatic ecosystems, but the existing gap between recovered
533 genotypes and their corresponding phenotypic and historical data can only be filled with substantially more
534 comprehensive reference libraries.

535 **ACKNOWLEDGEMENTS**

536 For assistance with logistics, eDNA sampling and providing the traditional survey data, we thank: Peter
537 Henderson and Robin Somes from Pisces Conservation Ltd.; Tony Gray, Hannah Westoby and the staff at
538 the Environment Agency in Newcastle Upon Tyne; and Aisling Smith and Sophie Rainbird from the Marine
539 Biological Association in Plymouth. This work was co-funded by the Natural Environment Research Council
540 grants NE/N005759/1 and NE/N005937/1 (project SeaDNA), and the University of Salford R&E strategy
541 funding. We thank two reviewers and the MEE editors for their help in improving the manuscript.

542 **AUTHOR CONTRIBUTIONS**

543 SM, MJG, DWS, and LC conceived the study and obtained funding; RAC, JB, OSW, and AZS carried out
544 field work and lab work; RAC, JB, and OSW performed the analyses; RAC and JB drafted the manuscript.
545 All authors contributed critically to the drafts and gave final approval for publication.

546 **DECLARATION OF INTEREST**

547 The authors declare that they have no competing interests.

548 **DATA ACCESSIBILITY**

549 The full reference library and code to reproduce it can be found at <https://doi.org/10.6084/m9.figshare.7464521>.
550 Code to reproduce all other analyses in this study can be found at <https://doi.org/10.6084/m9.figshare.8291660>.
551 Sequence data (fastq format) generated from the *in vitro* analyses can be found at <https://doi.org/10.5061/dryad.b8f6s44>.

552 REFERENCES

- 553 Alberdi, A., Aizpurua, O., Gilbert, M. T. P., and Bohmann, K. (2018). Scrutinizing key steps for reliable metabarcoding of environmental samples. *Methods in Ecology and Evolution*, 9:134–147, DOI: 10.1111/2041-210X.12849.
- 554
555
- 556 Andruszkiewicz, E. A., Starks, H. A., Chavez, F. P., Sassoubre, L. M., Block, B. A., and Boehm, A. B. (2017). Biomonitoring of marine vertebrates in Monterey Bay using eDNA metabarcoding. *PLoS ONE*, 12:e0176343, DOI: 10.1371/journal.pone.0176343.
- 557
558
- 559 Andújar, C., Arribas, P., Yu, D. W., Vogler, A. P., and Emerson, B. C. (2018). Why the COI barcode should be the community DNA metabarcode for the Metazoa. *Molecular Ecology*, 27:3968–3975, DOI: 10.1111/mec.14844.
- 560
561
- 562 Bakker, J., Wangenstein, O. S., Chapman, D. D., Boussarie, G., Buddo, D., Guttridge, T. L., Hertler, H., Mouillot, D., Vigliola, L., and Mariani, S. (2017). Environmental DNA reveals tropical shark diversity in contrasting levels of anthropogenic impact. *Scientific Reports*, 7:16886, DOI: 10.1038/s41598-017-17150-2.
- 563
564
565
- 566 Barbera, P., Kozlov, A. M., Czech, L., Morel, B., Darriba, D., Flouri, T., and Stamatakis, A. (2018). EPA-ng: massively parallel evolutionary placement of genetic sequences. *Systematic Biology*, DOI: 10.1101/291658.
- 567
568
- 569 Berry, T. E., Osterrieder, S. K., Murray, D. C., Coghlan, M. L., Richardson, A. J., Greal, A. K., Stat, M., Bejder, L., and Bunce, M. (2017). Metabarcoding for diet analysis and biodiversity: A case study using the endangered Australian sea lion (*Neophoca cinerea*). *Ecology and Evolution*, pages 1–19, DOI: 10.1002/ece3.3123.
- 570
571
572
- 573 Bista, I., Carvalho, G. R., Tang, M., Walsh, K., Zhou, X., Hajibabaei, M., Shokralla, S., Seymour, M., Bradley, D., Liu, S., Christmas, M., and Creer, S. (2018). Performance of amplicon and shotgun sequencing for accurate biomass estimation in invertebrate community samples. *Molecular Ecology Resources*, 18:1020–1034, DOI: 10.1111/1755-0998.12888.
- 574
575
576
- 577 Boettiger, C., Lang, D. T., and Wainwright, P. C. (2012). rfishbase: exploring , manipulating and visualizing FishBase data from R. *Journal of Fish Biology*, 81:2030–2039, DOI: 10.1111/j.1095-8649.2012.03464.x.
- 578
579
- 580 Boyer, S., Brown, S. D. J., Collins, R. A., Cruickshank, R. H., Lefort, M.-C., Malumbres-Olarte, J., and Wratten, S. D. (2012). Sliding window analyses for optimal selection of mini-barcodes, and application to 454-pyrosequencing for specimen identification from degraded DNA. *PLoS ONE*, 7:e38215, DOI: 10.1371/journal.pone.0038215.
- 581
582
583
- 584 Brown, S. D. J., Collins, R. A., Boyer, S., Lefort, M.-C., Malumbres-Olarte, J., Vink, C. J., and Cruickshank, R. H. (2012). Spider: an R package for the analysis of species identity and evolution, with particular reference to DNA barcoding. *Molecular Ecology Resources*, 12:562–565, DOI: 10.1111/j.1755-0998.2011.03108.x.
- 585
586
587
- 588 Bucklin, A., Steinke, D., and Blanco-Bercial, L. (2011). DNA Barcoding of Marine Metazoa. *Annual Review of Marine Science*, 3:471–508, DOI: 10.1146/annurev-marine-120308-080950.
- 589
- 590 Bylemans, J., Gleeson, D. M., Hardy, C. M., and Furlan, E. (2018). Toward an ecoregion scale evaluation of eDNA metabarcoding primers: A case study for the freshwater fish biodiversity of the Murray-Darling
- 591

592 Basin (Australia). *Ecology and Evolution*, 8:8697–8712, DOI: [10.1002/ece3.4387](https://doi.org/10.1002/ece3.4387).

593 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P. (2016).
594 DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, 13:581–583,
595 DOI: [10.1038/nmeth.3869](https://doi.org/10.1038/nmeth.3869).

596 Chamberlain, S. (2018). bold: Interface to Bold Systems API. [https://cran.r-project.org/
597 package=bold](https://cran.r-project.org/package=bold).

598 Chamberlain, S. and Boettiger, C. (2017). R Python, and Ruby clients for GBIF species occurrence data.
599 *PeerJ PrePrints*, DOI: [10.7287/peerj.preprints.3304v1](https://doi.org/10.7287/peerj.preprints.3304v1).

600 Chamberlain, S., Foster, Z., Bartomeus, I., LeBauer, D., Black, C., and Harris, D. (2019). traits: species trait
601 data from around the web. <https://cran.r-project.org/package=traits>.

602 Clarke, L. J., Beard, J. M., Swadling, K. M., and Deagle, B. E. (2017). Effect of marker choice and thermal
603 cycling protocol on zooplankton DNA metabarcoding studies. *Ecology and Evolution*, 7:873–883, DOI:
604 [10.1002/ece3.2667](https://doi.org/10.1002/ece3.2667).

605 Clarke, L. J., Soubrier, J., Weyrich, L. S., and Cooper, A. (2014). Environmental metabarcodes for insects:
606 In silico PCR reveals potential for taxonomic bias. *Molecular Ecology Resources*, 14:1160–1170, DOI:
607 [10.1111/1755-0998.12265](https://doi.org/10.1111/1755-0998.12265).

608 Coissac, E., Hollingsworth, P. M., Lavergne, S., and Taberlet, P. (2016). From barcodes to genomes: Extending
609 the concept of DNA barcoding. *Molecular Ecology*, 25:1423–1428, DOI: [10.1111/mec.13549](https://doi.org/10.1111/mec.13549).

610 Collins, R. A. and Cruickshank, R. H. (2014). Known knowns, known unknowns, unknown unknowns and
611 unknown knowns in DNA barcoding: A comment on Downton et al. *Systematic Biology*, 63:1005–1009,
612 DOI: [10.1093/sysbio/syu060](https://doi.org/10.1093/sysbio/syu060).

613 Collins, R. A., Wangenstein, O. S., Sims, D. W., Genner, M. J., and Mariani, S. (2018). Per-
614 sistence of environmental DNA in marine systems. *Communications Biology*, 1:185, DOI:
615 [10.1038/s42003-018-0192-6](https://doi.org/10.1038/s42003-018-0192-6).

616 Creer, S., Deiner, K., Frey, S., Porazinska, D., Taberlet, P., Thomas, K., Potter, C., and Bik, H. (2016). The
617 ecologist’s field guide to sequence-based identification of biodiversity. *Methods in Ecology and Evolution*,
618 56:68–74, DOI: [10.1111/2041-210X.12574](https://doi.org/10.1111/2041-210X.12574).

619 Czech, L. and Stamatakis, A. (2018). Scalable methods for post-processing , visualizing , and analyzing
620 phylogenetic placements. *bioRxiv*, pages 1–36, DOI: [10.1101/346353](https://doi.org/10.1101/346353).

621 Deagle, B. E., Jarman, S. N., Coissac, E., Pompanon, F., and Taberlet, P. (2014). DNA metabarcoding and
622 the cytochrome c oxidase subunit I marker: not a perfect match. *Biology Letters*, 10:20140562, DOI:
623 [10.1098/rsbl.2014.0562](https://doi.org/10.1098/rsbl.2014.0562).

624 Dopheide, A., Xie, D., Buckley, T. R., Drummond, A. J., and Newcomb, R. D. (2018). Impacts of DNA
625 extraction and PCR on DNA metabarcoding estimates of soil biodiversity. *Methods in Ecology and
626 Evolution*, DOI: [10.1111/2041-210X.13086](https://doi.org/10.1111/2041-210X.13086).

627 Drummond, A. J., Newcomb, R. D., Buckley, T. R., Xie, D., Dopheide, A., Potter, B. C. M., Heled, J., Ross,
628 H. A., Tooman, L., Grosser, S., Park, D., Demetras, N. J., Stevens, M. I., Russell, J. C., Anderson, S. H.,
629 Carter, A., and Nelson, N. (2015). Evaluating a multigene environmental DNA approach for biodiversity
630 assessment. *GigaScience*, 4:46, DOI: [10.1186/s13742-015-0086-1](https://doi.org/10.1186/s13742-015-0086-1).

631 Eddy, S. R. (1998). Profile hidden Markov models. *Bioinformatics*, 14:755–763, DOI:

632 [10.1093/bioinformatics/14.9.755](https://doi.org/10.1093/bioinformatics/14.9.755).

633 Elbrecht, V. and Leese, F. (2017). Validation and Development of COI Metabarcoding Primers for
634 Freshwater Macroinvertebrate Bioassessment. *Frontiers in Environmental Science*, 5:1–11, DOI:
635 [10.3389/fenvs.2017.00011](https://doi.org/10.3389/fenvs.2017.00011).

636 Elbrecht, V., Taberlet, P., Dejean, T., Valentini, A., Usseglio-Polatera, P., Beisel, J.-N., Coissac, E., Boyer, F.,
637 and Leese, F. (2016). Testing the potential of a ribosomal 16S marker for DNA metabarcoding of insects.
638 *PeerJ*, 4:e1966, DOI: [10.7717/peerj.1966](https://doi.org/10.7717/peerj.1966).

639 Elbrecht, V., Vamos, E. E., Meissner, K., Aroviita, J., and Leese, F. (2017). Assessing strengths and weaknesses
640 of DNA metabarcoding-based macroinvertebrate identification for routine stream monitoring. *Methods in*
641 *Ecology and Evolution*, 8:1265–1275, DOI: [10.1111/2041-210X.12789](https://doi.org/10.1111/2041-210X.12789).

642 Ficetola, G. F., Coissac, E., Zundel, S., Riaz, T., and Shehzad, W. (2010). An in silico approach for the
643 evaluation of DNA barcodes. *BMC Genomics*, 11:434, DOI: [10.1186/1471-2164-11-434](https://doi.org/10.1186/1471-2164-11-434).

644 Ficetola, G. F., Pansu, J., Bonin, A., Coissac, E., Giguët-Covex, C., De Barba, M., Gielly, L., Lopes, C. M.,
645 Boyer, F., Pompanon, F., Rayé, G., and Taberlet, P. (2015). Replication levels, false presences and
646 the estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources*,
647 15:543–556, DOI: [10.1111/1755-0998.12338](https://doi.org/10.1111/1755-0998.12338).

648 Folmer, O., Black, M., Hoeh, W., Lutz, R., and Vrijenhoek, R. (1994). DNA primers for amplification of
649 mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine*
650 *Biology and Biotechnology*, 3:294–299.

651 Gillett, C. P. D. T., Crampton-Platt, A., Timmermans, M. J. T. N., Jordal, B., Emerson, B. C., and
652 Vogler, A. P. (2014). Bulk de novo mitogenome assembly from pooled total DNA elucidates the phy-
653 logeny of weevils (Coleoptera: Curculionoidea). *Molecular Biology and Evolution*, 31:2223–2237, DOI:
654 [10.1093/molbev/msu154](https://doi.org/10.1093/molbev/msu154).

655 Grey, E. K., Bernatchez, L., Cassey, P., Deiner, K., Deveney, M., Howland, K. L., Lacoursière-Roussel, A.,
656 Leong, S. C. Y., Li, Y., Olds, B., Pfrender, M. E., Prowse, T. A., Renshaw, M. A., and Lodge, D. M. (2018).
657 Effects of sampling effort on biodiversity patterns estimated from environmental DNA metabarcoding
658 surveys. *Scientific Reports*, 8:2–11, DOI: [10.1038/s41598-018-27048-2](https://doi.org/10.1038/s41598-018-27048-2).

659 Guardiola, M., Uriz, M. J., Taberlet, P., Coissac, E., Wangenstein, O. S., and Turon, X. (2015). Deep-sea,
660 deep-sequencing: Metabarcoding extracellular DNA from sediments of marine canyons. *PLoS ONE*,
661 10:e0139633, DOI: [10.1371/journal.pone.0139633](https://doi.org/10.1371/journal.pone.0139633).

662 Hänfling, B., Lawson Handley, L., Read, D. S., Hahn, C., Li, J., Nichols, P., Blackman, R. C., Oliver, A., and
663 Winfield, I. J. (2016). Environmental DNA metabarcoding of lake fish communities reflects long-term data
664 from established survey methods. *Molecular Ecology*, 25:3101–3119, DOI: [10.1111/mec.13660](https://doi.org/10.1111/mec.13660).

665 Hebert, P. D. N., Cywinska, A., Ball, S. L., and DeWaard, J. R. (2003). Biological identifications
666 through DNA barcodes. *Proceedings of the Royal Society B: Biological Sciences*, 270:313–321, DOI:
667 [10.1098/rspb.2002.2218](https://doi.org/10.1098/rspb.2002.2218).

668 Henderson, P. A. (2014). *Identification Guide to the Inshore Fish of the British Isles*. Pisces Conservation
669 Ltd., Pennington, UK.

670 Hering, D., Borja, A., Jones, J. I., Pont, D., Boets, P., Bouchez, A., Bruce, K., Drakare, S., Hänfling, B., Kahlert,
671 M., Leese, F., Meissner, K., Mergen, P., Reyjol, Y., Segurado, P., Vogler, A., and Kelly, M. (2018). Imple-

672 mentation options for DNA-based identification into ecological status assessment under the European Water
673 Framework Directive. *Water Research*, 138:192–205, DOI: [10.1016/j.watres.2018.03.003](https://doi.org/10.1016/j.watres.2018.03.003).

674 Iwasaki, W., Fukunaga, T., Isagozawa, R., Yamada, K., Maeda, Y., Satoh, T. P., Sado, T., Mabuchi, K.,
675 Takeshima, H., Miya, M., and Nishida, M. (2013). Mitofish and mitoannotator: A mitochondrial genome
676 database of fish with an accurate and automatic annotation pipeline. *Molecular Biology and Evolution*,
677 30:2531–2540, DOI: [10.1093/molbev/mst141](https://doi.org/10.1093/molbev/mst141).

678 Jeunen, G.-J., Knapp, M., Spencer, H. G., Lamare, M. D., Taylor, H. R., Stat, M., Bunce, M., and
679 Gemmell, N. J. (2018). Environmental DNA (eDNA) metabarcoding reveals strong discrimination
680 among diverse marine habitats connected by water movement. *Molecular Ecology Resources*, DOI:
681 [10.1111/1755-0998.12982](https://doi.org/10.1111/1755-0998.12982).

682 Ji, Y., Ashton, L., Pedley, S. M., Edwards, D. P., Tang, Y., Nakamura, A., Kitching, R., Dolman, P. M., Wood-
683 cock, P., Edwards, F. A., Larsen, T. H., Hsu, W. W., Benedick, S., Hamer, K. C., Wilcove, D. S., Bruce, C.,
684 Wang, X., Levi, T., Lott, M., Emerson, B. C., and Yu, D. W. (2013). Reliable, verifiable and efficient moni-
685 toring of biodiversity via metabarcoding. *Ecology Letters*, 16:1245–1257, DOI: [10.1111/ele.12162](https://doi.org/10.1111/ele.12162).

686 Kartzinel, T. R., Chen, P. A., Coverdale, T. C., Erickson, D. L., Kress, W. J., Kuzmina, M. L., Rubenstein,
687 D. I., Wang, W., and Pringle, R. M. (2015). DNA metabarcoding illuminates dietary niche partitioning
688 by African large herbivores. *Proceedings of the National Academy of Sciences*, 112:8019–8024, DOI:
689 [10.1073/pnas.1503283112](https://doi.org/10.1073/pnas.1503283112).

690 Katoh, K. and Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:
691 improvements in performance and usability. *Molecular Biology and Evolution*, 30:772–780, DOI:
692 [10.1093/molbev/mst010](https://doi.org/10.1093/molbev/mst010).

693 Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A.,
694 Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., and Drummond, A. (2012). Geneious Basic:
695 an integrated and extendable desktop software platform for the organization and analysis of sequence data.
696 *Bioinformatics*, 28:1647–1649, DOI: [10.1093/bioinformatics/bts199](https://doi.org/10.1093/bioinformatics/bts199).

697 Kelly, R. P., Closek, C. J., O’Donnell, J. L., Kralj, J. E., Shelton, A. O., and Samhouri, J. F. (2017). Genetic
698 and manual survey methods yield different and complementary views of an ecosystem. *Frontiers in Marine*
699 *Science*, 3:1–11, DOI: [10.3389/fmars.2016.00283](https://doi.org/10.3389/fmars.2016.00283).

700 Kelly, R. P., Port, J. A., Yamahara, K. M., and Crowder, L. B. (2014a). Using environmental DNA to census
701 marine fishes in a large mesocosm. *PLoS ONE*, 9:e86175, DOI: [10.1371/journal.pone.0086175](https://doi.org/10.1371/journal.pone.0086175).

702 Kelly, R. P., Port, J. A., Yamahara, K. M., Martone, R. G., Lowell, N., Thomsen, P. F., Mach, M. E., Bennett,
703 M., Prahler, E., Caldwell, M. R., and Crowder, L. B. (2014b). Harnessing DNA to improve environmental
704 management. *Science*, 344:1455–1456, DOI: [10.1126/science.1251156](https://doi.org/10.1126/science.1251156).

705 Kottelat, M. and Freyhof, J. (2007). *Handbook of European Freshwater Fishes*. Publications Kottelat, Cornol,
706 Switzerland.

707 Leray, M. and Knowlton, N. (2015). DNA barcoding and metabarcoding of standardized samples reveal
708 patterns of marine benthic diversity. *Proceedings of the National Academy of Sciences*, 112:2076–2081,
709 DOI: [10.1073/pnas.1424997112](https://doi.org/10.1073/pnas.1424997112).

710 Leray, M. and Knowlton, N. (2016). Censusing marine eukaryotic diversity in the twenty-first cen-
711 tury. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371:20150331, DOI:

712 [10.1098/rstb.2015.0331](https://doi.org/10.1098/rstb.2015.0331).

713 Leray, M. and Knowlton, N. (2017). Random sampling causes the low reproducibility of rare eukaryotic
714 OTUs in Illumina COI metabarcoding. *PeerJ*, 5:e3006, DOI: [10.7717/peerj.3006](https://doi.org/10.7717/peerj.3006).

715 Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., Boehm, J. T., and Machida,
716 R. J. (2013). A new versatile primer set targeting a short fragment of the mitochondrial COI region for
717 metabarcoding metazoan diversity: application for characterizing coral reef fish gut contents. *Frontiers in*
718 *Zoology*, 10:34, DOI: [10.1186/1742-9994-10-34](https://doi.org/10.1186/1742-9994-10-34).

719 Lim, N. K. M., Tay, Y. C., Tan, J. W. T., Kwik, J. T. B., Baloglu, B., Meier, R., and Yeo, D. C. J. (2016).
720 Next-generation freshwater bioassessment: eDNA metabarcoding with a conserved metazoan primer reveals
721 species-rich and communities. *Royal Society Open Science*, 3:160635, DOI: [10.1098/rsos.160635](https://doi.org/10.1098/rsos.160635).

722 Linard, B., Arribas, P., Andújar, C., Crampton-Platt, A., and Vogler, A. P. (2016). Lessons from
723 genome skimming of arthropod-preserving ethanol. *Molecular Ecology Resources*, 16:1365–1377, DOI:
724 [10.1111/1755-0998.12539](https://doi.org/10.1111/1755-0998.12539).

725 Macher, J. N., Vivancos, A., Piggott, J. J., Centeno, F. C., Matthaei, C. D., and Leese, F. (2018). Comparison
726 of environmental DNA and bulk-sample metabarcoding using highly degenerate cytochrome c oxidase I
727 primers. *Molecular Ecology Resources*, 18:1456–1468, DOI: [10.1111/1755-0998.12940](https://doi.org/10.1111/1755-0998.12940).

728 Mahé, F., Rognes, T., Quince, C., de Vargas, C., and Dunthorn, M. (2015). Swarm v2: highly-scalable and
729 high-resolution amplicon clustering. *PeerJ*, 3:e1420, DOI: [10.7717/peerj.1420](https://doi.org/10.7717/peerj.1420).

730 Mariani, S., Griffiths, A. M., Velasco, A., Kappel, K., Jerome, M., Perez-Martin, R. I., Schroder, U., Verrez-
731 Bagnis, V., Silva, H., Vandamme, S. G., Boufana, B., Mendes, R., Shorten, M., Smith, C., Hankard, E.,
732 Hook, S. A., Weymer, A. S., Gunning, D., and Sotelo, C. G. (2015). Low mislabeling rates indicate
733 marked improvements in European seafood market operations. *Frontiers in Ecology and the Environment*,
734 13:536–540, DOI: [10.1890/150119](https://doi.org/10.1890/150119).

735 Marquina, D., Andersson, A. F., and Ronquist, F. (2019). New mitochondrial primers for metabarcoding of
736 insects, designed and evaluated using in silico methods. *Molecular Ecology Resources*, 19:90–104, DOI:
737 [10.1111/1755-0998.12942](https://doi.org/10.1111/1755-0998.12942).

738 Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMB-*
739 *net.journal*, 17:10–12, DOI: [10.14806/ej.17.1.200](https://doi.org/10.14806/ej.17.1.200).

740 McHugh, M., Sims, D. W., Partridge, J. C., and Genner, M. J. (2011). A century later: Long-term
741 change of an inshore temperate marine fish assemblage. *Journal of Sea Research*, 65:187–194, DOI:
742 [10.1016/j.seares.2010.09.006](https://doi.org/10.1016/j.seares.2010.09.006).

743 Minamoto, T., Yamanaka, H., Takahara, T., Honjo, M. N., and Kawabata, Z. (2012). Surveil-
744 lance of fish species composition using environmental DNA. *Limnology*, 13:193–197, DOI:
745 [10.1007/s10201-011-0362-4](https://doi.org/10.1007/s10201-011-0362-4).

746 Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J. Y., Sato, K., Minamoto, T., Yamamoto, S., Yamanaka,
747 H., Araki, H., Kondoh, M., and Iwasaki, W. (2015). MiFish, a set of universal PCR primers for metabar-
748 coding environmental DNA from fishes: detection of more than 230 subtropical marine species. *Royal*
749 *Society Open Science*, 2:150088, DOI: [10.1098/rsos.150088](https://doi.org/10.1098/rsos.150088).

750 Pedersen, M. W., Overballe-Petersen, S., Ermini, L., Der Sarkissian, C., Haile, J., Hellstrom, M., Spens,
751 J., Thomsen, P. F., Bohmann, K., Cappellini, E., Schnell, I. B., Wales, N. A., Carøe, C., Campos, F.,

- 752 Schmidt, A. M. Z., Gilbert, M. T. P., Hansen, A. J., Orlando, L., and Willerslev, E. (2015). Ancient and
753 modern environmental DNA. *Philosophical Transactions of the Royal Society B: Biological Sciences*,
754 370:20130383, DOI: [10.1098/rstb.2013.0383](https://doi.org/10.1098/rstb.2013.0383).
- 755 Piñol, J., Mir, G., Gomez-Polo, P., and Agustí, N. (2015). Universal and blocking primer mismatches limit
756 the use of high-throughput DNA sequencing for the quantitative metabarcoding of arthropods. *Molecular*
757 *Ecology Resources*, 15:819–830, DOI: [10.1111/1755-0998.12355](https://doi.org/10.1111/1755-0998.12355).
- 758 Port, J. A., O'Donnell, J. L., Romero-Maraccini, O. C., Leary, P. R., Litvin, S. Y., Nickols, K. J., Yamahara,
759 K. M., and Kelly, R. P. (2016). Assessing vertebrate biodiversity in a kelp forest ecosystem using
760 environmental DNA. *Molecular Ecology*, 25:527–541, DOI: [10.1111/mec.13481](https://doi.org/10.1111/mec.13481).
- 761 Qu, W., Zhou, Y., Zhang, Y., Lu, Y., Wang, X., Zhao, D., Yang, Y., and Zhang, C. (2012). MFEprimer-2.0:
762 A fast thermodynamics-based program for checking PCR primer specificity. *Nucleic Acids Research*,
763 40:205–208, DOI: [10.1093/nar/gks552](https://doi.org/10.1093/nar/gks552).
- 764 Ratnasingham, S. and Hebert, P. D. N. (2007). BOLD: The Barcode of Life
765 Data System (www.barcodinglife.org). *Molecular Ecology Notes*, 7:355–364, DOI:
766 [10.1111/j.1471-8286.2006.01678.x](https://doi.org/10.1111/j.1471-8286.2006.01678.x).
- 767 Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R. M., and Gough, K. C. (2014). The detection
768 of aquatic animal species using environmental DNA - a review of eDNA as a survey tool in ecology.
769 *Journal of Applied Ecology*, 51:1450–1459, DOI: [10.1111/1365-2664.12306](https://doi.org/10.1111/1365-2664.12306).
- 770 Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P., and Coissac, E. (2011). EcoPrimers: Inference of
771 new DNA barcode markers from whole genome sequence analysis. *Nucleic Acids Research*, 39:e145, DOI:
772 [10.1093/nar/gkr732](https://doi.org/10.1093/nar/gkr732).
- 773 Shaw, J. L. A., Clarke, L. J., Wedderburn, S. D., Barnes, T. C., Weyrich, L. S., and Cooper, A. (2016).
774 Comparison of environmental DNA metabarcoding and conventional fish survey methods in a river system.
775 *Biological Conservation*, 197:131–138, DOI: [10.1016/j.biocon.2016.03.010](https://doi.org/10.1016/j.biocon.2016.03.010).
- 776 Shokralla, S., Hellberg, R. S., Handy, S. M., King, I., and Hajibabaei, M. (2015). A DNA mini-
777 barcoding system for authentication of processed fish products. *Scientific Reports*, 5:15894, DOI:
778 [10.1038/srep15894](https://doi.org/10.1038/srep15894).
- 779 Siddall, M. E., Fontanella, F. M., Watson, S. C., Kvist, S., and Erséus, C. (2009). Barcoding bamboozled by
780 bacteria: convergence to metazoan mitochondrial primer targets by marine microbes. *Systematic Biology*,
781 58:445–451, DOI: [10.1093/sysbio/syp033](https://doi.org/10.1093/sysbio/syp033).
- 782 Singer, G. A. C., Fahner, N. A., Barnes, J. G., McCarthy, A., and Hajibabaei, M. (2019). Comprehensive
783 biodiversity analysis via ultra-deep patterned flow cell technology: a case study of eDNA metabarcoding
784 seawater. *Scientific Reports*, 9:5991, DOI: [10.1038/s41598-019-42455-9](https://doi.org/10.1038/s41598-019-42455-9).
- 785 Spens, J., Evans, A. R., Halfmaerten, D., Knudsen, S. W., Sengupta, M. E., Mak, S. S. T., Sigsgaard, E. E.,
786 and Hellström, M. (2017). Comparison of capture and storage methods for aqueous microbial eDNA
787 using an optimized extraction protocol: advantage of enclosed filter. *Methods in Ecology and Evolution*,
788 8:635–645, DOI: [10.1111/2041-210X.12683](https://doi.org/10.1111/2041-210X.12683).
- 789 Stamatakis, A., Hoover, P., and Rougemont, J. (2008). A rapid bootstrap algorithm for the RAxML Web
790 servers. *Systematic Biology*, 57:758–771, DOI: [10.1080/10635150802429642](https://doi.org/10.1080/10635150802429642).
- 791 Stat, M., Huggett, M. J., Bernasconi, R., Dibattista, J. D., Berry, T. E., Newman, S. J., Harvey, E. S., and

- 792 Bunce, M. (2017). Ecosystem biomonitoring with eDNA: metabarcoding across the tree of life in a tropical
793 marine environment. *Scientific Reports*, 7:0–22, DOI: [10.1038/s41598-017-12501-5](https://doi.org/10.1038/s41598-017-12501-5).
- 794 Stat, M., John, J., DiBattista, J. D., Newman, S. J., Bunce, M., and Harvey, E. S. (2018). Combined use of
795 eDNA metabarcoding and video surveillance for the assessment of fish biodiversity. *Conservation Biology*,
796 33:196–205, DOI: [10.1111/cobi.13183](https://doi.org/10.1111/cobi.13183).
- 797 Stoeckle, M. Y., Soboleva, L., and Charlop-Powers, Z. (2017). Aquatic environmental DNA detects
798 seasonal fish abundance and habitat preference in an urban estuary. *PLoS ONE*, 12:e0175186, DOI:
799 [10.1371/journal.pone.0175186](https://doi.org/10.1371/journal.pone.0175186).
- 800 Sunagawa, S., Coelho, L. P., Chaffron, S., Kultima, J. R., Labadie, K., Salazar, G., Djahanschiri, B., Zeller,
801 G., Mende, D. R., Alberti, A., Cornejo-Castillo, F. M., Costea, P. I., Cruaud, C., D’Ovidio, F., Engelen, S.,
802 Ferrera, I., Gasol, J. M., Guidi, L., Hildebrand, F., Kokoszka, F., Lepoivre, C., Lima-Mendez, G., Poulain,
803 J., Poulos, B. T., Royo-Llonch, M., Sarmiento, H., Vieira-Silva, S., Dimier, C., Picheral, M., Searson, S.,
804 Kandels-Lewis, S., Boss, E., Follows, M., Karp-Boss, L., Krzic, U., Reynaud, E. G., Sardet, C., Sieracki,
805 M., Velayoudon, D., Bowler, C., De Vargas, C., Gorsky, G., Grimsley, N., Hingamp, P., Iudicone, D.,
806 Jaillon, O., Not, F., Ogata, H., Pesant, S., Speich, S., Stemmann, L., Sullivan, M. B., Weissenbach, J.,
807 Wincker, P., Karsenti, E., Raes, J., Acinas, S. G., and Bork, P. (2015). Structure and function of the global
808 ocean microbiome. *Science*, 348:1–10, DOI: [10.1126/science.1261359](https://doi.org/10.1126/science.1261359).
- 809 Taberlet, P., Bonin, A., Zinger, L., and Coissac, E. (2018). *Environmental DNA: For Biodiversity Research and*
810 *Monitoring*. Oxford University Press, Oxford, DOI: [10.1093/oso/9780198767220.001.0001](https://doi.org/10.1093/oso/9780198767220.001.0001).
- 811 Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., and Willerslev, E. (2012). Towards next-
812 generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, 21:2045–2050, DOI:
813 [10.1111/j.1365-294X.2012.05470.x](https://doi.org/10.1111/j.1365-294X.2012.05470.x).
- 814 Thomsen, P. F., Kielgast, J., Iversen, L. L., Møller, P. R., Rasmussen, M., and Willerslev, E. (2012a). Detection
815 of a diverse marine fish fauna using environmental DNA from seawater samples. *PLoS ONE*, 7:e41732,
816 DOI: [10.1371/journal.pone.0041732](https://doi.org/10.1371/journal.pone.0041732).
- 817 Thomsen, P. F., Kielgast, J., Iversen, L. L., Wiuf, C., Rasmussen, M., Gilbert, M. T. P., Orlando, L.,
818 and Willerslev, E. (2012b). Monitoring endangered freshwater biodiversity using environmental DNA.
819 *Molecular Ecology*, 21:2565–2573, DOI: [10.1111/j.1365-294X.2011.05418.x](https://doi.org/10.1111/j.1365-294X.2011.05418.x).
- 820 Thomsen, P. F., Møller, P. R., Sigsgaard, E. E., Knudsen, S. W., Jørgensen, O. A., and Willerslev, E. (2016).
821 Environmental DNA from seawater samples correlate with trawl catches of subarctic, deepwater fishes.
822 *PLoS ONE*, 11:e0165252, DOI: [10.1371/journal.pone.0165252](https://doi.org/10.1371/journal.pone.0165252).
- 823 Thomsen, P. F. and Willerslev, E. (2015). Environmental DNA - An emerging tool in conser-
824 vation for monitoring past and present biodiversity. *Biological Conservation*, 183:4–18, DOI:
825 [10.1016/j.biocon.2014.11.019](https://doi.org/10.1016/j.biocon.2014.11.019).
- 826 Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B. C., Remm, M., and Rozen,
827 S. G. (2012). Primer3-new capabilities and interfaces. *Nucleic Acids Research*, 40:e115, DOI:
828 [10.1093/nar/gks596](https://doi.org/10.1093/nar/gks596).
- 829 Ushio, M., Murakami, H., Masuda, R., Sado, T., Miya, M., Sakurai, S., Yamanaka, H., Minamoto, T., and
830 Kondoh, M. (2018). Quantitative monitoring of multispecies fish environmental DNA using high-throughput
831 sequencing. *Metabarcoding and Metagenomics*, 2:1–15, DOI: [10.3897/mbmg.2.23297](https://doi.org/10.3897/mbmg.2.23297).

- 832 Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., Bellemain, E., Besnard, A.,
833 Coissac, E., Boyer, F., Gaboriaud, C., Jean, P., Poulet, N., Roset, N., Copp, G. H., Geniez, P., Pont, D.,
834 Argillier, C., Baudoin, J. M., Peroux, T., Crivelli, A. J., Olivier, A., Acqueberge, M., Le Brun, M., Møller,
835 P. R., Willerslev, E., and Dejean, T. (2016). Next-generation monitoring of aquatic biodiversity using
836 environmental DNA metabarcoding. *Molecular Ecology*, 25:929–942, DOI: [10.1111/mec.13428](https://doi.org/10.1111/mec.13428).
- 837 Vandamme, S. G., Griffiths, A. M., Taylor, S.-A., Di Muri, C., Hankard, E. A., Towne, J. A., Watson,
838 M., and Mariani, S. (2016). Sushi barcoding in the UK: another kettle of fish. *PeerJ*, 4:e1891, DOI:
839 [10.7717/peerj.1891](https://doi.org/10.7717/peerj.1891).
- 840 Wangenstein, O. S., Palacín, C., Guardiola, M., and Turon, X. (2018). DNA metabarcoding of littoral
841 hard-bottom communities: high diversity and database gaps revealed by two molecular markers. *PeerJ*,
842 6:e4705, DOI: [10.7717/peerj.4705](https://doi.org/10.7717/peerj.4705).
- 843 Ward, R. D. (2009). DNA barcode divergence among species and genera of birds and fishes. *Molecular*
844 *Ecology Resources*, 9:1077–1085, DOI: [10.1111/j.1755-0998.2009.02541.x](https://doi.org/10.1111/j.1755-0998.2009.02541.x).
- 845 Ward, R. D., Hanner, R., and Hebert, P. D. N. (2009). The campaign to DNA barcode all fishes, FISH-BOL.
846 *Journal of Fish Biology*, 74:329–56, DOI: [10.1111/j.1095-8649.2008.02080.x](https://doi.org/10.1111/j.1095-8649.2008.02080.x).
- 847 Ward, R. D., Zemlak, T. S., Innes, B. H., Last, P. R., and Hebert, P. D. N. (2005). DNA barcoding Australia's
848 fish species. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360:1847–1857, DOI:
849 [10.1098/rstb.2005.1716](https://doi.org/10.1098/rstb.2005.1716).
- 850 Wilcox, T. M., Zarn, K. E., Piggott, M. P., Young, M. K., McKelvey, K. S., and Schwartz, M. K. (2018).
851 Capture enrichment of aquatic environmental DNA: A first proof of concept. *Molecular Ecology Resources*,
852 18:1392–1401, DOI: [10.1111/1755-0998.12928](https://doi.org/10.1111/1755-0998.12928).
- 853 Willerslev, E., Hansen, A. J., Binladen, J., Brand, T. B., Gilbert, M. T. P., Shapiro, B., Bunce, M., Wiuf, C.,
854 Gilichinsky, D. A., and Cooper, A. (2003). Diverse plant and animal genetic records from Holocene and
855 Pleistocene sediments. *Science*, 300:791–795, DOI: [10.1126/science.1084114](https://doi.org/10.1126/science.1084114).
- 856 Winter, D. J. (2017). rentrez: an R package for the NCBI eUtils API. *The R Journal*, 9:520–526.
- 857 Yamamoto, S., Masuda, R., Sato, Y., Sado, T., Araki, H., Kondoh, M., Minamoto, T., and Miya, M. (2017).
858 Environmental DNA metabarcoding reveals local fish communities in a species-rich coastal sea. *Scientific*
859 *Reports*, 7:40368, DOI: [10.1038/srep40368](https://doi.org/10.1038/srep40368).
- 860 Yang, C., Wang, X., Miller, J. A., De Blécourt, M., Ji, Y., Yang, C., Harrison, R. D., and Yu, D. W. (2014). Us-
861 ing metabarcoding to ask if easily collected soil and leaf-litter samples can be used as a general biodiversity
862 indicator. *Ecological Indicators*, 46:379–389, DOI: [10.1016/j.ecolind.2014.06.028](https://doi.org/10.1016/j.ecolind.2014.06.028).
- 863 Yu, D. W., Ji, Y., Emerson, B. C., Wang, X., Ye, C., Yang, C., and Ding, Z. (2012). Biodiversity soup:
864 metabarcoding of arthropods for rapid biodiversity assessment and biomonitoring. *Methods in Ecology and*
865 *Evolution*, 3:613–623, DOI: [10.1111/j.2041-210X.2012.00198.x](https://doi.org/10.1111/j.2041-210X.2012.00198.x).

Table 1. Primer sets assessed in this study. The approximate fragment length is based upon the length of that region in the *Anguilla anguilla* mitochondrial genome (AP007233.1). The asterisks represent the sequences of the Leray-XT primer set that were simplified by changing inosines to double-base ambiguities to allow an *in silico* assessment with *MFEprimer*. The standard DNA barcode marker for fishes (Ward et al., 2005) is presented for reference.

Primer set	Locus	Primer names	Oligonucleotide 5'–3'	Fragment length (bp)	Reference
Leray-XT	COI	mlCOIintF-XT	GGWACWRGWTGRACWITITAYCCYCC	313	Wangensteen et al. (2018)
		mlCOIintF-XT*	GGWACWRGWTGRACWGTYTAYCCYCC		
		hgHCO2198	TAIACYTCIGGRTGICCRARAAYCA		
		hgHCO2198*	TAKACYTCWGGRTGRCCRAARAAYCA		
SeaDNA-short	COI	coi.175f	GGAGGCTTTGGMAAYTGRYT	55	This study
		coi.226r	GGGGGAAGAARYCARAARCT		
SeaDNA-mid	COI	coi.175f	GGAGGCTTTGGMAAYTGRYT	130	This study
		coi.345r	TAGAGGRGGGTARACWGTYCA		
Ward-barcode	COI	FishF1	TCAACCAACCACAAAGACATTGGCAC	655	Ward et al. (2005)
		FishR1	TAGACTTCTGGGTGGCCAAAGAATCA		
Minamoto-fish	Cytb	L14912-CYB	TTCCCTAGCCATACAYTAYAC	235	Minamoto et al. (2012)
MiFish-U	12S	H15149-CYB	GGTGGCKCCTCAGAAGGACATTTGKCCYCA	171	Miya et al. (2015)
		MiFish-U-F	GTCGGTAAAACCTCGTGCCAGC		
MiFish-E	12S	MiFish-U-R	CATAGTGGGGTATCTAATCCCAGTTG	171	Miya et al. (2015)
		MiFish-E-F	GTTGGTAAATCTCGTGCCAGC		
Taberlet-tele02	12S	MiFish-E-R	CATAGTGGGGTATCTAATCCTAGTTG	167	Taberlet et al. (2018)
		Tele02-f	AAACTCGTGCCAGCCACC		
Taberlet-elas02	12S	Tele02-r	GGGTATCTAATCCCAGTTG	171	Taberlet et al. (2018)
		Elas02-f	GTTGGTHAATCTCGTGCCAGC		
Valentini-tele01	12S	Elas02-r	CATAGTAGGGTATCTAATCCTAGTTG	63	Valentini et al. (2016)
		L1848	ACACCGCCCGTCACTCT		
Riaz-V5	12S	H1913	CTTCCGGTACACTTACCATG	106	Riaz et al. (2011)
		12S-V5f	ACTGGGATTAGATACCCC		
Berry-fish	16S	12S-V5r	TAGAACAGGCTCCTCTAG	219	Berry et al. (2017)
		Fish16sF/D	GACCCATATGGAGCTTTAGAC		
		16s2R	CGCTGTTATCCCTADRGTAACT		

Table 2. Statistics for reference library coverage, taxonomic discriminatory power, and primer universality as estimated by *in silico* PCR, for twelve primer sets from COI, cytochrome *b*, 16S and 12S. Library coverage is calculated as the number of species for which at least one sequence was available out of the total ($n = 531$) or common species subset ($n = 176$) of British Isles marine and freshwater fishes (proportion in parentheses). Library sequences per species is the mean (median in parentheses) number of sequences available for each species. Taxonomic discrimination is the proportion of species for which all individuals can be unambiguously identified by a unique DNA sequence, with values in parentheses showing the proportion for the subset of species that are shared over all primer sets ($n = 221$ for all; $n = 88$ for common). Primer universality represents the mean Primer Pair Coverage (PPC) percent statistic from *MFEprimer*, and was calculated using the 184 British Isles fish species for which data were available for all species. The standard DNA barcode marker for fishes (Ward et al., 2005) is presented for reference. The highly degenerate Leray-XT primers were simplified to overcome analytical RAM limitations (see Table 1).

Locus	Primer pair	Species subset	Total number sequences	Library species coverage	Library sequences per species	Fragment taxonomic discrimination	Primer % universality (Actinopterygii)	Primer % universality (Elasmobranchii)	
COI	Leray-XT	All	24,058	481 (0.91)	50 (24)	0.95 (0.96)	27.8	39	
	SeaDNA-mid		24,045	481 (0.91)	50 (24)	0.91 (0.94)	23.8	23.7	
	SeaDNA-short		23,911	481 (0.91)	49.7 (24)	0.87 (0.9)	34.5	21.5	
	Ward-barcode		23,975	481 (0.91)	49.8 (24)	0.95 (0.97)	6.3	1.2	
CYTB	Minamoto-fish		15,405	344 (0.65)	44.8 (6.5)	0.91 (0.91)	19.1	20.4	
12S	MiFish-U		1,904	322 (0.61)	5.9 (3)	0.93 (0.91)	86.6	3	
	Taberlet-tele02		1,904	322 (0.61)	5.9 (3)	0.93 (0.91)	92.1	8.3	
	MiFish-E		1,904	322 (0.61)	5.9 (3)	0.93 (0.91)	0.6	55	
	Taberlet-elas02		1,904	322 (0.61)	5.9 (3)	0.93 (0.91)	0.5	83.6	
	Valentini-tele01		1,699	273 (0.51)	6.2 (2)	0.86 (0.85)	77.1	68.2	
	Riaz-V5		2,416	364 (0.69)	6.6 (2)	0.79 (0.78)	92.2	11.2	
	16S	Berry-fish		4,089	411 (0.77)	9.9 (4)	0.89 (0.86)	50.9	0
COI	Leray-XT	Common	12,698	170 (0.97)	74.7 (38.5)	0.97 (1)	23.3	49.3	
	SeaDNA-mid		12,639	170 (0.97)	74.3 (37.5)	0.93 (1)	17.6	30	
	SeaDNA-short		12,553	170 (0.97)	73.8 (37.5)	0.93 (1)	32.8	28.9	
	Ward-barcode		12,579	170 (0.97)	74 (37.5)	0.97 (1)	6.3	0	
CYTB	Minamoto-fish		10,936	143 (0.81)	76.5 (16)	0.94 (1)	19.2	12.9	
12S	MiFish-U		941	109 (0.62)	8.6 (3)	0.94 (0.94)	91.8	0	
	Taberlet-tele02		941	109 (0.62)	8.6 (3)	0.94 (0.94)	96.4	0	
	MiFish-E		941	109 (0.62)	8.6 (3)	0.94 (0.94)	0	73.3	
	Taberlet-elas02		941	109 (0.62)	8.6 (3)	0.94 (0.94)	0	100	
	Valentini-tele01		852	99 (0.56)	8.6 (2)	0.93 (0.94)	76.4	68.2	
	Riaz-V5		1,398	143 (0.81)	9.8 (3)	0.85 (0.83)	96.4	0	
	16S	Berry-fish		2,296	167 (0.95)	13.7 (6)	0.87 (0.91)	53.9	0

Table 3. Number of reads remaining after seven bioinformatic steps, as well as the number of estimated reads for taxonomic groups (assignments were carried out on the reads remaining after the homology search step 7). Fish reads (putative) are reads assigned to fishes based on the best scoring *blastn* hit using the NCBI “nt” blast database. Fish reads (assigned) are reads assigned to fish species by the stringent taxonomic identification step using *blastn* and *EPA-ng* on our curated reference library. Fish reads (unassigned) are putative fish reads that could not be assigned to species by the stringent taxonomic identification step.

Filtering step	COI Leray-XT	COI SeaDNA-mid	COI SeaDNA-short	12S MiFish-U
Total passing filter	5,967,313	14,291,168	8,881,088	3,436,278
(1) Detect primers	4,828,799	11,535,904	6,428,030	2,776,073
(2) Demultiplex	4,648,811	10,879,223	5,994,815	2,473,594
(3) Trim primers	4,618,236	10,300,907	5,852,555	2,462,936
(4) Quality filter	4,519,097	10,344,024	5,856,045	2,455,532
(5) Merge	3,395,057	9,658,709	4,804,502	2,383,162
(6) Remove chimaeras	3,225,240	9,404,746	4,416,647	2,271,541
(7) Homology search	3,223,743	8,703,109	4,074,123	2,157,365
Bacteria	1,476,994	1,388,681	2,242,220	4
Eukaryota	1,745,295	7,294,762	1,815,928	2,157,361
Metazoa	321,590	1,161,769	412,871	2,157,361
Chordata	6,351	337,901	250,650	2,157,361
Fish (putative)	2,371	234,219	193,593	1,637,728
Fish (assigned)	1,368	109,486	30,026	1,564,351
Fish (unassigned)	1,003	124,733	163,567	73,377




Figure 1. Fish species richness as estimated by four primer pairs at five sampling locations. Per primer-location combination there are three water sample replicates and two uniquely tagged PCR replicates ($n = 6$). The horizontal represents the median value, the boxes represent the 25–75th percentiles, the whiskers represent the values less than 1.5 times the interquartile range, dots represent the outlying data points, and crosses represent the cumulative number of species.




Figure 2. Reproducibility heatmaps of four primer pairs at five sampling locations for the top ten fish species found at each location by read abundance. Letters A, B, and C represent the three water samples taken, while numbers 1 and 2 represent the independent PCR reactions with uniquely tagged primers. There are ten shades showing 10% increments. The darkest shade shows a reproducibility of 100%, i.e. reads from all of the ten species were common to both PCRs. The lightest shade shows 0% reproducibility, i.e. none of the species were present in both of the PCRs. Diagonals show the proportion of the top ten species amplified in that single PCR.



Figure 3. Overlap between fish species found by eDNA metabarcoding (red) and traditional fish surveying (blue). Sizes of circles are proportional only within each primer-location comparison, and not between. Numbers represent number of species in each set. Only marine and estuarine species are shown; freshwater species recorded by the eDNA surveys were removed to allow an equivalent comparison.