

## REVIEW

## Non-synaptic receptors and transporters involved in brain functions and targets of drug treatment

ES Vizi, A Fekete, R Karoly and A Mike

*Department of Pharmacology, Institute of Experimental Medicine, Hungarian Academy of Sciences, Budapest, Hungary*

Beyond direct synaptic communication, neurons are able to talk to each other without making synapses. They are able to send chemical messages by means of diffusion to target cells via the extracellular space, provided that the target neurons are equipped with high-affinity receptors. While synaptic transmission is responsible for the 'what' of brain function, the 'how' of brain function (mood, attention, level of arousal, general excitability, etc.) is mainly controlled non-synaptically using the extracellular space as communication channel. It is principally the 'how' that can be modulated by medicine. In this paper, we discuss different forms of non-synaptic transmission, localized spillover of synaptic transmitters, local presynaptic modulation and tonic influence of ambient transmitter levels on the activity of vast neuronal populations. We consider different aspects of non-synaptic transmission, such as synaptic–extrasynaptic receptor trafficking, neuron–glia communication and retrograde signalling. We review structural and functional aspects of non-synaptic transmission, including (i) anatomical arrangement of non-synaptic release sites, receptors and transporters, (ii) intravesicular, intra- and extracellular concentrations of neurotransmitters, as well as the spatiotemporal pattern of transmitter diffusion. We propose that an effective general strategy for efficient pharmacological intervention could include the identification of specific non-synaptic targets and the subsequent development of selective pharmacological tools to influence them.

*British Journal of Pharmacology* (2010) **160**, 785–809; doi:10.1111/j.1476-5381.2009.00624.x; published online 8 February 2010

**Keywords:** non-synaptic chemical transmission; extrasynaptic high-affinity receptors; extrasynaptic high-affinity transporters; extracellular space; tonic modulation; presynaptic modulation; receptor and transporter trafficking; targets of medicines

**Abbreviations:** 5-HT, serotonin; ACh, acetylcholine; AMPA,  $\alpha$ -amino-3-hydroxyl-5-methyl-4-isoxazole-propionate; AP, action potential; 2-AG, 2-arachidonoylglycerol; CB1, cannabinoid1; CNS, central nervous system; DA, dopamine; DAT, dopamine transporter; EAAT, excitatory amino acid transporter; GABA,  $\gamma$ -amino-butyric acid; Glu, glutamate; GlyR, glycine receptor; LTP, long-term potentiation; mAChR, muscarinic acetylcholine receptor; mGlu, metabotropic glutamate receptor; nAChR, nicotinic acetylcholine receptor; NA, noradrenaline; NET, noradrenaline transporter; NMDA, N-methyl-D-aspartate; SERT, serotonin transporter

## Introduction

Our understanding of chemical transmission has been primarily influenced by the action of acetylcholine (ACh) at the neuromuscular junction, where ACh is released into the synaptic gap, diffuses 10–20 nm and interacts with intrasynaptic receptors of relatively low affinity. Over the years, functional, neurochemical and morphological evidence has shown that this assumption needs to be redefined, given that many neurotransmitters can diffuse out of the synaptic cleft to distant target receptors, and that the majority of transmitters may also be released from non-synaptic terminals.

The original observations that established the theory of non-synaptic transmission are both of functional (neurochemical) and morphological nature. During the 1950s and 1960s, the picture regarding neuronal communication appeared to be clear. Neurons were believed to communicate using chemicals to bridge the gaps between the pre- and postsynaptic sites of the synapse. Action potentials (APs) were understood to arrive at synaptic terminals in order to release transmitters and to transmit messages from one neuron to another, or between a neuron and its target cell. Based on the all-or-nothing nature of action potentials, the brain was thought to be a predominantly digital device of discrete computational units. This idea was held in general agreement until the end of the 1960s, when some novel discoveries broke the rule of synaptic transmission and challenged the dogma that the synapse is the exclusive site for the cell-to-cell information processing. Neurochemical evidence was obtained that noradrenaline (NA) released from

noradrenergic nerves presynaptically inhibits the release of ACh from the cholinergic interneurons in the Auerbach plexus and cerebral cortex (Vizi, 1968; Paton and Vizi, 1969; Vizi and Knoll, 1971). Light microscopic studies have shown a 'mismatch' between release sites and the locations of corresponding receptors (Herkenham, 1987; 1991). Electron microscopic ultrastructural studies have indicated that the majority of monoaminergic and cholinergic boutons in the CNS lack junctional complexes (Hökfelt, 1968; Descarries *et al.*, 1975; 1977).

These observations led to the suggestion (Vizi, 1974; 1980a,b; 1984) that in addition to synaptic chemical communication, there is another form of interaction among neurons. This interaction occurs when neurons release chemicals that diffuse far away in order to send messages without the need for close synaptic arrangements, thereby challenging the traditional view of synaptic transmission (Vizi, 1974; 1980a,b; 1984; Schmitt, 1984; Agnati *et al.*, 1986; 1995; Fuxe and Agnati, 1991; Bach-y-Rita, 1993; Zoli *et al.*, 1998; Descarries and Mechawar, 2000).

In this review, we aim to emphasize the fact that most CNS drugs primarily affect the non-synaptic system. In the periphery, several drugs are classic agonists and antagonists (such as adrenergic agonists and antagonists, cholinergic agonists, antimuscarinic agents, ganglionic blockers, and neuromuscular blockers); in the CNS, however, the typical drugs are modulators, and their targets are non-synaptic receptors, transporters, enzymes and ion channels. Non-synaptic receptors are more affected not only because they outnumber synaptic ones, but also because they are more sensitive to fine modulation of ambient concentrations of transmitters present in the extracellular space compared with receptors in synapses. In spite of these facts, many textbooks and some reviews (e.g. Krishnan and Nestler, 2008) still fail to discuss this form of interneuronal communication, indicating such ideas as 'increases in the amount of synaptic monoamines induced by antidepressants' have possible therapeutic effects.

This review attempts to summarize the evidence that has started to change and has already begun to shape our understanding of chemical transmission both between neurons and between neurons and non-neuronal target cells.


### Neurochemical and morphological evidence for non-synaptic communication

Neurochemical evidence has demonstrated that there is a functional interaction between the noradrenergic and cholinergic axon terminals, such that NA released from noradrenergic neurons is able to inhibit the release of ACh (Vizi, 1968; 1972; 1974; 1979; 1980a,b; Paton and Vizi, 1969; Knoll and Vizi, 1970; 1971; Vizi and Knoll, 1971) despite the lack of synaptic contact between the two terminals (Silva *et al.*, 1971; Gordon-Weeks, 1982). Noradrenergic inhibition of ACh release can be viewed as presynaptic modulation, and, when compared with regulation at the postsynaptic receptor level, offers the physiologic advantage of economy in chemical transmission and the presynaptic inhibitory interaction. Removal of the noradrenergic control of cholinergic transmis-

sion by chemical lesion of the noradrenergic innervation or the administration of  $\alpha_2$ -adrenoceptor antagonists resulted in a steady state increase of ACh release from the Auerbach plexus (Vizi, 1968; Paton and Vizi, 1969) and from isolated cerebral cortex slices (Vizi, 1974; 1980a). The results indicated that there was a tonic inhibition of ACh release by NA that was being continuously released into the fluid present in the extracellular space surrounding cholinergic release sites. Overall, it was suggested (Vizi, 1974) that '... there is morphological and histological evidence (Silva *et al.*, 1971) that the adrenergic terminals terminate not far from the cholinergic ones. This enables the released NA to reach the cholinergic nerve endings by diffusion and to act there via  $\alpha$ -adrenoceptor'. A similar interaction was shown for dopamine (DA) and ACh release in the striatum (Vizi *et al.*, 1977), where DA inhibited the axonal stimulation-induced release of ACh via tonic activation of the D<sub>2</sub> receptor located on the axons of cholinergic cells. Inhibition of DA release or the removal of the nigrostriatal pathway resulted in a significant increase in ACh release, indicating the tonic influence on the release (Vizi *et al.*, 1977). This was the first evidence of the disinhibitory mechanism with no synaptic contacts among neurons. These observations validate a persistent tonic influence on neuronal activity at the presynaptic level (Vizi and Labos, 1991). Furthermore, they illustrate that even without synaptic connections, it is still possible to create functional interactions between two nerves that are equipped with different transmitter machineries, provided that the target neurons are equipped with receptors sensitive to the transmitter released from another neuron. Further evidence has shown that non-synaptic communication is a fundamental element of neuro-endocrine (Vizi *et al.*, 1992b; 1993) and neuro-immune (Vizi *et al.*, 1995; Elenkov *et al.*, 2000) interactions.

#### *Non-synaptic varicosities*

Over half of the 100 billion neurons in the CNS release glutamate (Glu), and 30–40 billion release GABA ( $\gamma$ -aminobutyric acid) into the synaptic gap (van der Zeyden *et al.*, 2008). Furthermore, almost all neurons express receptors that are sensitive to Glu (van der Zeyden *et al.*, 2008). Glutamatergic and GABAergic neurons were thought to release their transmitters exclusively in synapses (Freund and Buzsaki, 1996). The exception to the rule has recently been discovered for GABAergic varicosities. Neurogliaform cells of the neocortex were found to release GABA from predominantly non-synaptic varicosities (Olah *et al.*, 2009). On the other hand, the overwhelming majority of monoaminergic neurons fail to make synaptic contacts (Descarries and Mechawar, 2000). A common feature of the monoaminergic system is that the cell bodies are located in small subcortical nuclei (for example, the locus coeruleus for noradrenergic, the substantia nigra for dopaminergic and the raphe nuclei for serotonergic neurons), but the varicose axon arborizations diffusely innervate large target areas, thereby creating vast neuronal assemblies (Figure 1). As revealed by concordant light microscopic fluorescence, radioautographic and immunocytochemical studies, preterminal monoaminergic axons branch profusely, and give rise to billions of varicose nerve endings without making synaptic contact (Table 1).



**Figure 1** Scheme of the arborization of different neurons in the hippocampus. Arrows indicate varicosities, while the arrowheads show the bifurcation of branches. The average length of axonal arborization (mm) per cell body, number of boutons per cell body, percentage of non-synaptic boutons and density of boutons are given ( $10^6 \cdot \text{mm}^{-3}$ ). <sup>1</sup>Aznavour *et al.* (2002), <sup>2</sup>Descarries *et al.* (1997), <sup>3</sup>Umbriaco *et al.* (1995), <sup>4</sup>Vizi and Kiss (1998), <sup>5</sup>Hökfelt (1968), <sup>6</sup>Oleskevich *et al.* (1989), <sup>7</sup>Descarries *et al.* (1990), <sup>8</sup>Gulyas *et al.* (1993), <sup>9</sup>Olah *et al.* (2009). The inset shows axonal varicosities of a GABAergic interneuron from the rat hippocampus (A. Barth, N. Hájos and E.S. Vizi, Institute of Experimental Medicine, unpublished data) using a two-photon laser-scanning microscope (for method see Rozsa *et al.* 2004 and Vizi *et al.*, 2004a). Scale bars are 10 and 50  $\mu\text{m}$ , as indicated.

**Table 1** Different neuronal projections in the brain without making synaptic contacts

Projections	% of non-synaptic varicosities	Location	References
Cholinergic	79–90	Somatosensory cortex	(Umbriaco <i>et al.</i> , 1994)
	86	Cerebral cortex	(Descarries <i>et al.</i> , 1997)
	93	Hippocampus (CA1)	
	91	Neostriatum	
	33	Human cerebral cortex	(Smiley <i>et al.</i> , 1997)
Noradrenergic	85	Hippocampus (CA1)	(Hökfelt, 1968; Umbriaco <i>et al.</i> , 1995)
	95	Cerebral cortex	(Descarries <i>et al.</i> , 1977)
	>60	Dorsal horn	(Ridet <i>et al.</i> , 1993)
	90	Frontal cortex	(Aoki <i>et al.</i> , 1998)
Dopaminergic	60–70	Neostriatum	(Descarries <i>et al.</i> , 1996; 2008; Wilson <i>et al.</i> , 1977)
		Median eminence and cerebral cortex	(Calas <i>et al.</i> , 1974; Descarries <i>et al.</i> , 1975; Tao-Cheng and Zhou, 1999)
Serotonergic	82	Raphe nucleus	(Kosofsky and Molliver, 1987)
	97	Somatosensory cortex	(DeFelipe and Jones, 1988)
	>60	Dorsal horn	(Ridet <i>et al.</i> , 1993)
	~50	Suprachiasmatic and supraoptic nuclei of hypothalamus	(Boulaich <i>et al.</i> , 1994)
	79	Hippocampus (CA1)	(Umbriaco <i>et al.</i> , 1995)
Glutamatergic	90	Prefrontal cortex	(Smiley and Goldman-Rakic, 1996)
	>60	Spinal cord	(Alvarez <i>et al.</i> , 1998)
	80	Periaqueductal gray matter	(Lovick <i>et al.</i> , 2000)
	0		(Freund and Gulyas, 1991)
	0		(Freund and Gulyas, 1991; Umbriaco <i>et al.</i> , 1995)
Gabaergic	78	Hippocampus (CA1)	(Olah <i>et al.</i> , 2009)
		Neurogliaform cells in the cortex	

Experiments are carried out in rats if otherwise not stated.

It has been calculated that each noradrenergic cell body that ipsilaterally innervates the CNS (700 per hemisphere in rats; Descarries and Saucier, 1972) has an axonal arborization, with an average of 170 000 varicosities within the hippocam-

pus (Vizi and Kiss, 1998). With regard to cholinergic innervation of the hippocampus, the number is even higher: a single cholinergic cell (located in the median septum, or the diagonal band of Broca) has an axonal arborization that is

~2.1 m in total length within the ~56 mm<sup>3</sup> of the hippocampus, and is equipped with ~840 000 varicosities (Figure 1; Aznavour *et al.*, 2002). The majority of these varicosities exist mainly without synaptic contact (Descarries and Mechawar, 2000). Clearly, when an action potential is generated at the axon hillock, it will not cause release from all the ~840 000 varicosities. Furthermore, the cell body cannot dictate which of the specific varicosities should release transmitter. Rather, release probability is a subject of local modulation, and, as was shown for the innervation of the vas deferens, where ATP and NA are co-transmitters, low release probability is not due to failure of AP propagation but to a variable reliability of depolarization–secretion coupling (Jackson and Cunnane, 2001). This emphasizes the importance of local modulation of transmitter release; even though the AP propagates, whether it evokes release or not depends on the local environment of individual varicosities. It is also possible, however, that release is prevented by shutting down entire axonal branches (Debanne *et al.*, 1997; Kress and Mennerick, 2009). GABA-mediated increases in membrane conductance may be an important mechanism to control AP propagation to axonal branches (Segev, 1990; Verdier *et al.*, 2003).

By means of quantitative autoradiographic studies, the existence of non-synaptic communication in the CNS has been structurally shown. Monoaminergic (Hökfelt, 1968; Descarries *et al.*, 1975; Descarries *et al.*, 1977; Ridet *et al.*, 1993; Descarries *et al.*, 1996; Lovick *et al.*, 2000) and cholinergic axon terminals (Umbriaco *et al.*, 1995; Descarries and Mechawar, 2000) were found without synaptic contact, yet they are still equipped with releasing machinery. Only a small percentage (5–40%) of the varicosities formed synapses (Descarries *et al.*, 1975; Ridet *et al.*, 1993; Umbriaco *et al.*, 1995; Alvarez *et al.*, 1998; Lovick *et al.*, 2000). The overwhelming majority of non-synaptic boutons appear to have all the apparatuses that are associated with AP-dependent vesicular release and are able to release transmitters for long-distance diffusion. These findings indicate that the field of influence for a single neuron, which collects several hundreds or thousands of inputs, could be very large (Figure 1, Table 1).

It is interesting to note that all of the dopaminergic varicosities in the neostriatum or nucleus accumbens, which contain the vesicular transporter for Glu in addition to tyrosine hydroxylase and might therefore release Glu as a co-transmitter, make synaptic contacts (Descarries *et al.*, 2008). Other varicosities that contain only DA in their release machineries, however, fail to make synapses (Descarries *et al.*, 1996). The varicosities of serotonergic innervation that originate from the median raphe always form synapses with hippocampal interneurons (Freund and Buzsaki, 1996; Varga *et al.*, 2009); however, varicosities from the dorsal raphe do not make any synaptic contacts in the hippocampus (Kosofsky and Molliver, 1987). The experiments of Freund *et al.* (1985) may provide some explanation for this discrepancy. After reimplantation of substantia nigra into rats, in which the substantia nigra was previously removed, the nigro-striatal neurons, which otherwise do not synapse with other neurons (Wilson *et al.*, 1977) have made synapses 100% with cholinergic interneurons. It therefore appears very likely

that during development, the time of innervation plays an important role in setting the synapse.

#### *Non-synaptic receptors*

In addition to synaptic localization, there are receptors located non-synaptically. Their presence and capability of receiving chemical messages can be shown by morphological and functional methods (Table 2).

The current terminology for the location-based classification of receptors is inconsistent on several points. In this review, we do not intend to reform the terminology, and we will use the terms that seem the most logical and widely accepted. Still, we cannot avoid pointing out the inconsistencies. There are two major classification schemes based on both subcellular localization and function. One classification of receptors distinguishes synaptic and non-synaptic receptors. Non-synaptic receptors are sometimes referred to as extrasynaptic; however, we prefer to use the term extrasynaptic in a less inclusive manner, distinguishing synaptic, perisynaptic and extrasynaptic localizations (with the latter two groups both being non-synaptic). Perisynaptic receptors are located at the edges of synapses, while extrasynaptic receptors are located farther away. The perisynaptic–extrasynaptic boundary is rather subjective; thus, we propose to define it functionally. Perisynaptic receptors are those that are able to detect individual release events; therefore, the precise timing of digital information coding can be distinguished by these receptors. For extrasynaptic receptors, being far from the release sites, the precise temporal information is lost, and they can only follow the analogue signals, such as tonic changes in the ambient levels of the transmitter. The spatial limits of digital information exchange are termed the 'maximum effective radius' (Rice and Cragg, 2008), which is defined as the distance within which the release of one quantum of a transmitter reaches the EC50 concentration of the receptor. We propose that non-synaptic, somatodendritic receptors within the maximum effective radius should be termed perisynaptic. Unfortunately, we have no appropriate term for non-synaptic receptors located near non-synaptic release sites.

The other classification scheme is to distinguish presynaptic (axonal) and postsynaptic (somatodendritic) receptors. Presynaptic, axonal receptors are responsible for local modulation of depolarization–release coupling, including propagation of APs and modulation of release probability. Postsynaptic, somatodendritic receptors, on the other hand, modulate the integration of inputs, that is they decide whether or not inputs translate into APs. Unfortunately, we have no correct term for axonal and somatodendritic receptors in cases where there are no synapses; therefore, we must call these receptors 'presynaptic' and 'postsynaptic', respectively, despite the lack of morphologically distinguishable synapses. Presynaptic receptors can be further divided into autoreceptors (i.e. receptors that are sensitive to the neuron's own transmitter, which is termed as 'negative feedback' modulation) or heteroreceptors (i.e. receptors conveying the influence of other transmitters and neuromodulators).

Axonal receptors are sometimes divided into two functional categories, presynaptic and preterminal (Wonnacott, 1997). In this terminology, presynaptic receptors are only those



**Table 2** Non-synaptic localization and site of action of different ionotropic and metabotropic receptors (subunits are indicated if needed)

Receptors	Region in the brain	Site of action	References
Ionotropic	GABA <sub>A</sub>	Cerebellum	Synaptic and non-synaptic expression in the somatodendritic compartment of granule cells; no expression in axons, nerve terminals or glial cells (Somogyi <i>et al.</i> , 1989; Nusser <i>et al.</i> , 1995)
		Dorsal lateral geniculate nucleus	Synaptic and non-synaptic expression (Soltesz <i>et al.</i> , 1990)
	GABA <sub>A</sub> (δ)	Cerebellum	Exclusive extrasynaptic location in granule cells (Nusser <i>et al.</i> , 1998)
	GABA <sub>A</sub> (α <sub>4</sub> )	Hippocampus	Mainly perisynaptic; switch to synaptic in a model of ethanol withdrawal and dependence (Liang <i>et al.</i> , 2006)
		Dentate gyrus	
	GABA <sub>A</sub> (β <sub>2</sub> )	Dentate gyrus	Extrasynaptic in granule cells (Herd <i>et al.</i> , 2008)
	GABA <sub>A</sub> (β <sub>3</sub> )	Dentate gyrus	Synaptic in granule cells (Herd <i>et al.</i> , 2008)
	AMPA	Hippocampus	Postsynaptic membrane specialization; in a lower number extrasynaptically on spines, dendrites, somata (Baude <i>et al.</i> , 1995)
		Dentate gyrus	
	AMPA (GluR1)	Cerebellum	Synaptic localization; intracellular and glia cell labeling (Baude <i>et al.</i> , 1994)
	Kainate (GluR6/7)	Striatum	>70% intracellular; presynaptically both in the active zone (~20%) of asymmetric synapses and at non-synaptic sites (~80%); on spines in asymmetric synapses (~40%) and at non-synaptic localization (~60%) (Kieval <i>et al.</i> , 2001)
	Kainate (KA2)	Striatum	>70% intracellular; presynaptically both in the active zone (~30%) of asymmetric synapses and at non-synaptic sites (~70%); on spines in asymmetric synapses (~40%) and at non-synaptic localization (~60%) (Kieval <i>et al.</i> , 2001)
NR2A	Hippocampus Cortex Thalamus	Mainly synaptic localization following postnatal development (Kew <i>et al.</i> , 1998; Tovar and Westbrook, 1999; Liu <i>et al.</i> , 2004b; Thomas <i>et al.</i> , 2006)	
NR2B	Hippocampus Cortex Thalamus	Mainly extrasynaptic localization following postnatal development (synaptic and extrasynaptic until postnatal development) (Kew <i>et al.</i> , 1998; Tovar and Westbrook, 1999; Liu <i>et al.</i> , 2004b; Thomas <i>et al.</i> , 2006)	
NR2C	Cerebellum (granule cell) Cortex (spiny stellate neuron)	Synaptic localization (Rossi <i>et al.</i> , 2002; Binshtok <i>et al.</i> , 2006)	
NR2D	Hippocampus Dorsal horn	Extrasynaptic localization (Momiyama <i>et al.</i> , 1996; Momiyama, 2000; Lozovaya <i>et al.</i> , 2004; Harney <i>et al.</i> , 2008)	
NR3A	Limited expression in rodent, but widespread expression in human	Eriksson <i>et al.</i> , 2007	
nACh (α <sub>7</sub> )	Hippocampus	Presynaptic terminals in cultured neurons (Fabian-Fine <i>et al.</i> , 2001)	
	Ventral tegmental area	Glutamatergic terminals (perisynaptic) (Jones and Wonnacott, 2004)	
	Striatum	Nigro-striatal terminals	
	Cerebellum	Perisynaptic (Caruncho <i>et al.</i> , 1997)	
nACh	Hippocampus	Soma (Zarei <i>et al.</i> , 1999)	
	Ventral tegmental area	Glutamatergic nerve terminals (Jones <i>et al.</i> , 2001)	
nACh	Chicken pretectum	(Ullian and Sargent, 1995)	
Metabotropic (G-protein-coupled)	α <sub>2A</sub>	Prefrontal cortex	Preterminal axon (Aoki <i>et al.</i> , 1998; Abbracchio <i>et al.</i> , 2009)
		Hippocampus	Axon terminal, glia (Smiley <i>et al.</i> , 1994; Milner <i>et al.</i> , 1998)
	D <sub>1</sub>	Cerebral cortex	Spiny dendrite, perikarya (Smiley <i>et al.</i> , 1994; Yung <i>et al.</i> , 1995)
		Basal ganglia	
	D <sub>2</sub>	Midbrain	Axon terminals, spines, dendrites (auto- and heteroreceptors) (Sesack <i>et al.</i> , 1994; Yung <i>et al.</i> , 1995)
		Basal ganglia, subst. Nigra	
	5-HT <sub>1A</sub>	Substantia nigra	Cell body, dendrite (Riad <i>et al.</i> , 2000)
	5-HT <sub>1B</sub>	Substia-nigra	Preterminal axon (Riad <i>et al.</i> , 2000)
		Globus pallidus	
	GABA <sub>B</sub>	Cerebellum	Purkinje cell (Scanziani, 2000; Kulik <i>et al.</i> , 2006; Lujan and Shigemoto, 2006)
Cortex		Dendrites and spines Pyramidal cell dendrite and spines (Lopez-Bendito <i>et al.</i> , 2002)	

Table 2 Continued

Receptors	Region in the brain	Site of action	References
CB1	Hippocampus	Preterminal axon of cholecystokinin-containing gabaergic interneuron	(Katona <i>et al.</i> , 1999)
A <sub>1</sub>	Striatum Hippocampus Widely distributed	Pre- and postsynaptic	(Wu and Saggau, 1997; Ciruela <i>et al.</i> , 2006)
A <sub>2A</sub>	Striatum Hippocampus Cerebral cortex Widely distributed	Pre- and postsynaptic	(Johansson and Fredholm, 1995; Cunha <i>et al.</i> , 1996; Rosin <i>et al.</i> , 1998; Ciruela <i>et al.</i> , 2006)
mGlu1	Hippocampus Cerebellum Neocortex	Concentrated in a perisynaptic annulus of postsynaptic membrane; extrasynaptically at spines, dendrites and somata	(Baude <i>et al.</i> , 1993; Martin <i>et al.</i> , 1992; Lujan <i>et al.</i> , 1996; 1997)
mGlu2	Hippocampus Cerebellum	Extrasynaptically along axons and axon terminals	(Lujan <i>et al.</i> , 1997; Shigemoto <i>et al.</i> , 1997; Ohishi <i>et al.</i> , 1998)
mGlu3	Hippocampus Cerebellum Striatum Thalamus	Extrasynaptically along axons and axon terminals; associated with asymmetrical synapses both in the postsynaptic specialization and in a perisynaptic position, glial processes	(Shigemoto <i>et al.</i> , 1997; Tamaru <i>et al.</i> , 2001)
mGlu4	Striatum Cerebellum Hippocampus Thalamus	Presynaptic active zone, postsynaptic density, spines, dendrites and somata	(Bradley <i>et al.</i> , 1996; Shigemoto <i>et al.</i> , 1997; Corti <i>et al.</i> , 2002)
mGlu5	Hippocampus Cerebellum Striatum	Concentrated in a perisynaptic annulus of postsynaptic membrane; extrasynaptically at spines, dendrite and somata	(Lujan <i>et al.</i> , 1996; 1997; Uchigashima <i>et al.</i> , 2007)
mGlu6	Retina	Somata, Located on dendrites of on-bipolar cells far from the release site (400–800 nm)	(Nakajima <i>et al.</i> , 1993; Nomura <i>et al.</i> , 1994; Vardi <i>et al.</i> , 2000)
mGlu7	Hippocampus Dorsal horn	Presynaptic active zone	(Ohishi <i>et al.</i> , 1995; Shigemoto <i>et al.</i> , 1997)
mGlu8	Hippocampus	Presynaptic active zone	(Shigemoto <i>et al.</i> , 1997)
M1	Striatum Cerebral cortex	Extrasynaptically at somata, spines and dendrites; postsynaptic densities of asymmetric and symmetric synapses	(Mrzljak <i>et al.</i> , 1993; Hersch <i>et al.</i> , 1994; Alcantara <i>et al.</i> , 2001; Narushima <i>et al.</i> , 2007; Uchigashima <i>et al.</i> , 2007)
M2	Striatum Cerebral cortex Nucleus basalis magnocellularis	Extrasynaptically at somata, dendrites; postsynaptic densities of asymmetric and symmetric synapses; presynaptic terminals of asymmetric and symmetric synapses; non-synaptic varicosities and perikarya of cholinergic neurons	(Mrzljak <i>et al.</i> , 1993; Hersch <i>et al.</i> , 1994; Bernard <i>et al.</i> , 1998; Rouse <i>et al.</i> , 2000; Decossas <i>et al.</i> , 2003)
M3	Striatum	Postsynaptic densities of asymmetric synapses; presynaptic terminals of asymmetric synapses	(Hersch <i>et al.</i> , 1994)
M4	Striatum	Extrasynaptically at somata, spines and dendrites; postsynaptic densities of asymmetric synapses; presynaptic terminals of asymmetric synapses; non-synaptic sites of cholinergic perikarya	(Hersch <i>et al.</i> , 1994; Bernard <i>et al.</i> , 1999)
M5			(Yamada <i>et al.</i> , 2003)

located in close proximity to the synapse (possibly even within the synaptic cleft), and their activation results in transmitter release even in the absence of functional axonal sodium channels. Thus, the release is tetrodotoxin (TTX) insensitive. Release evoked by preterminal receptors, on the other hand, requires axonal sodium channels to amplify and conduct the signal; therefore, it is TTX sensitive. This classification, however, can only be used for facilitatory axonal receptors, that is when receptor activation evokes transmitter release. Furthermore, the term presynaptic is more often used in a more general way, including receptors with different axonal locations.

In summary, we can conclude that, thus far, there is no logical terminology for non-synaptic receptors. We have no better option but to call certain groups 'perisynaptic' and

'presynaptic' even when they are connected not to morphologically observable synapses but to non-synaptic release sites. This inconsistency originates from the fact that, until recently, non-synaptic communication was peripheral in scientific thinking, and synaptic communication was used as a synonym of interneuronal communication. Thinking influences terminology, and terminology might have also influenced thinking.

#### *Cell surface exchange of receptors between synaptic and non-synaptic sites (receptor trafficking)*

At the neuromuscular junction the nicotinic acetylcholine receptors (nAChRs) are trapped at the synapse on the postsynaptic site, and are not localized extrasynaptically. Upon

innervation of skeletal muscle, nAChRs congregate in the junctional end-plate area. When the motor nerve has been severed, the nAChRs start to move away from the synapse and the receptors become distributed across the whole surface of the striated muscle (Axelsson and Thesleff, 1959).

A key mechanism regulating synaptic efficiency in the CNS is receptor trafficking composed of the lateral (between synaptic and non-synaptic compartments) and vertical (mobility between intracellular stores and cell surface by endocytosis and exocytosis) mobility of receptors described for many receptor types (Lau and Zukin, 2007; Groc *et al.*, 2009; Jaskolski and Henley, 2009). Formerly, vertical mobility has been considered the mechanism that promotes rapid changes in receptor number at postsynaptic density, however, recently, a complementary hypothesis has emerged. According to this, the mobility of receptors in/out of synapses is composed of two phases: (i) lateral diffusion of receptors in/out of the synapse, (ii) exocytosis/endocytosis of receptors in/out of the non-synaptic compartment (Triller and Choquet, 2005; Yudowski *et al.*, 2007). The finding of endocytic zones on spines lateral from synapse (Blanpied *et al.*, 2002) reveal that lateral diffusion of receptors between synapse and non-synaptic regions may be the part of the process of receptor recycling (Lau and Zukin, 2007).

Examination of Glu, GABA and glycine receptor trafficking is a prominent field of neuroscience. Glu receptor (NMDA, AMPA) trafficking into and out of synapses is influenced by neuronal activity (Lau and Zukin, 2007; Frischknecht *et al.*, 2009), and is an important mechanism to change the number of functional receptors in the synapse during plasticity (Choquet and Triller, 2003). The synaptic scaffolding proteins (e.g. PSD-95, PSD-93, stargazin) are involved in the stabilization of NMDA and AMPA receptors at the synapse (Schnell *et al.*, 2002; Lau and Zukin, 2007; Santos *et al.*, 2009). Synaptic GABA<sub>A</sub> and glycine receptors are also directly recruited from their non-synaptic counterparts and maintained in the synapse by direct binding to the multifunctional protein gephyrin (Meier *et al.*, 2001; Thomas *et al.*, 2005; Jacob *et al.*, 2008). This highly dynamic and regulated exchange between synaptic and non-synaptic membrane (Triller and Choquet, 2003) influences the number of non-synaptic receptors, thereby it may also affect synaptic plasticity and the effectiveness of medicines.

#### *G-protein-coupled receptors (Table 2)*

**Peptidergic receptors.** Neuropeptides are typical non-synaptic transmitters, which are released extrasynaptically (Torrealba and Carrasco, 2004; Wotjak *et al.*, 2008). Because of the large number of neuropeptides, in this paper, we are unable to give a detailed discussion of peptidergic transmission.

**Adrenoceptors.** Noradrenergic neurotransmission is mostly non-synaptic in the CNS (Figure 1, Table 1). Of the nine subtypes of adrenoceptors ( $\alpha_{1A}$ ,  $\alpha_{1B}$  and  $\alpha_{2A}$ ,  $\alpha_{2B}$  and  $\alpha_{2C}$ ,  $\beta_1$ ,  $\beta_2$  and  $\beta_3$ ),  $\alpha_2$  receptors are the most important in presynaptic modulation (Gilsbach and Hein, 2008). An electron microscopic immunocytochemical study has shown that  $\alpha_{2A}$ -adrenoceptors are located mostly on extrasynaptic areas, including axons, dendrites and astrocytic membranes in

the prefrontal cortex (Aoki *et al.*, 1998).  $\beta$ -adrenoceptors expressed on noradrenergic axons (van Zwieten and Timmermans, 1983) and on immune cells (Elenkov *et al.*, 2000) have been shown to be localized extrasynaptically.

**Dopamine receptors.** Morphological evidence for the non-synaptic role of DA in the CNS includes the sparsity of dopaminergic synapses (Table 1) in the striatum and substantia nigra (Wilson *et al.*, 1977), as well as the predominance of extrasynaptic DA receptors (Sesack *et al.*, 1994; Yung *et al.*, 1995; Table 2).


**Serotonin (5-HT, 5-hydroxytryptamine) receptors.** Serotonergic transmission in the CNS is predominantly non-synaptic (Figure 1, Table 1). Using immunogold labelling, extrasynaptic localization was shown for the 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptors. Both receptors are negatively coupled to adenylate cyclase. While 5-HT<sub>1A</sub> immunoreactivity was found on mostly non-synaptic locations of cell bodies and dendrites, 5-HT<sub>1B</sub> receptors were preferentially associated with unmyelinated preterminal axons, but were not found on axon terminals (Riad *et al.*, 2000). These receptors can be both inhibitory auto- and heteroreceptors.

**GABAB receptors.** GABA<sub>B</sub> receptors are largely located non-synaptically (Scanziani, 2000; Kulik *et al.*, 2006), and are also abundant in axons (Kulik *et al.*, 2003; Vigot *et al.*, 2006). On the postsynaptic Purkinje cells of the cerebellum, GABA<sub>B1</sub> and GABA<sub>B2</sub> receptors are localized to the extrasynaptic and perisynaptic plasma membrane (Lujan and Shigemoto, 2006).

**Metabotropic glutamate receptors (mGlu).** The mGlu receptors consist of eight subtypes classified into three subgroups. Group I receptors (mGlu1 and 5) couple to G<sub>q/11</sub> protein and induce LTD (Bellone *et al.*, 2008), group II (mGlu2 and 3) and group III receptors (mGlu4, 6, 7 and 8) inhibit cAMP formation through G<sub>i</sub> protein (Bellone *et al.*, 2008) and suppress transmitter release (Scanziani *et al.*, 1995; Takahashi *et al.*, 1996; Giustizieri *et al.*, 2005; see Figure 2 and Table 2 for localization).

**Muscarinic ACh receptors (mAChRs).** Cholinergic neurotransmission is mainly non-synaptic in the CNS (Figure 1, Table 1). The five mAChR subtypes are referred to as M<sub>1</sub>–M<sub>5</sub>. The M<sub>1</sub>, M<sub>3</sub>, M<sub>5</sub> receptors couple to PLC through G<sub>q/11</sub> activation, while M<sub>2</sub> and M<sub>4</sub> inhibit adenylyl cyclase via G<sub>i</sub> protein (Caulfield and Birdsall, 1998). Species- (Plummer *et al.*, 1999), age- (Decossas *et al.*, 2005), region- and cell-specific differences (Rouse and Levey, 1996; Nathanson, 2008) in the subcellular distribution of various subtypes of mAChRs and caveats in the immunocytochemical techniques (Nathanson, 2008) obscure the ability to distinguish between receptor subtypes (for detailed localization data, see Table 2).

**P2Y receptors.** P2Y receptors have two distinct subgroups. The first contains P2Y<sub>1,2,4,6</sub> and <sub>11</sub>, which couple to G<sub>q/11</sub> protein, and the second constitutes P2Y<sub>12,13,14</sub>, and inhibits adenylyl cyclase through G<sub>i</sub> protein activation (Abbracchio *et al.*, 2009). P2Y receptors have been shown to be present at presynaptic, postsynaptic and extrasynaptic sites in the



**Figure 2** Subcellular localization of mGlu receptors at synaptic and non-synaptic (perisynaptic or extrasynaptic) sites. Group I mGlu receptors are concentrated at the postsynaptic membrane in a perisynaptic annulus, but are also present extrasynaptically. Group II mGlu receptors are located extrasynaptically along axons and axon terminals, while group III mGlu receptors are expressed in the presynaptic active zone. <sup>1</sup>Lujan *et al.* (1996), <sup>2</sup>Ohishi *et al.* (1994), <sup>3</sup>Petralia *et al.* (1996), <sup>4</sup>Lujan *et al.* (1997), <sup>5</sup>Corti *et al.* (2002), <sup>6</sup>Shigemoto *et al.* (1997), <sup>7</sup>Baude *et al.* (1993), <sup>8</sup>Bradley *et al.* (1996), <sup>9</sup>Tamaru *et al.* (2001), <sup>10</sup>Uchigashima *et al.* (2007). \*mGlu3 can be found in a postsynaptic localization in the hippocampus and striatum but not in the ventrobasal nuclear complex of the thalamus.

hippocampus and in glial cells underlying the non-synaptic action of ATP (Fields and Stevens, 2000; Rodrigues *et al.*, 2005; Hussl and Boehm, 2006; Sperlagh *et al.*, 2007a). P2Y receptors also have a prominent role in the sensory pathways (Housley *et al.*, 2009). Many immunocytochemical data, however, should be handled with caution due to the lack of verification of the specificity (Sperlagh *et al.*, 2007a).

**Adenosine receptors.** The extracellular nucleotides are degraded by ectoenzymes to adenosine (Zimmermann, 2008), which in turn acts on postsynaptic metabotropic receptors (A<sub>1</sub>, A<sub>2A</sub>, A<sub>2B</sub>, A<sub>3</sub>) without any postsynaptic morphological arrangements. The high-affinity A<sub>1</sub> and A<sub>2A</sub> receptors (affinities in  $\mu\text{M}$ : A<sub>1</sub>, ~0.07; A<sub>2A</sub>, ~0.15; A<sub>2B</sub>, ~5.1; A<sub>3</sub>, ~6.5) are primarily responsible for the effects of adenosine in the brain, because basal extracellular adenosine concentrations (0.01–0.5  $\mu\text{M}$ ) are sufficient to activate them tonically (Latini *et al.*, 1999; Dunwiddie and Masino, 2001). In addition to their postsynaptic location, A<sub>1</sub> (Johansson and Fredholm, 1995; Wu and Saggau, 1997) and A<sub>2A</sub> (Rosin *et al.*, 1998) receptors have been described presynaptically (Table 2; Ciruela *et al.*, 2006). The

first evidence that adenosine and nucleotides are able to inhibit AP-induced transmitter release from cholinergic terminals was shown in the cerebral cortex and the Auerbach plexus (Vizi and Knoll, 1976). Presynaptic A<sub>1</sub> receptor activation reduces transmitter release mainly through the inhibition of Ca<sup>2+</sup> channels and activation of K<sup>+</sup> conductances (Milusheva *et al.*, 1990; Yawo and Chuhma, 1993; Haas and Selbach, 2000), while A<sub>2A</sub> receptors enhance neurotransmitter release (Popoli *et al.*, 1995; Nishizaki *et al.*, 2002; Cunha, 2008). Adenosine influences immunity via activation of A<sub>2B</sub> receptor (Hasko *et al.*, 2009). A<sub>2A</sub> receptors are also found in glial cells (Hettinger *et al.*, 2001).

**Cannabinoid CB1 receptors.** Major physiological effects of cannabinoids (Freund and Katona, 2007) can be linked to the regulation of neurotransmitter release via presynaptic CB1 receptors (Katona *et al.*, 1999; Freund *et al.*, 2003; Kofalvi *et al.*, 2003; Kofalvi *et al.*, 2005; Schoffelmeer *et al.*, 2006; Katona and Freund, 2008). Immunogold staining methods have shown that CB1 receptors are localized extrasynaptically in the preterminal axons (Katona *et al.*, 1999),



and also in the perisynaptic annulus on the presynaptic terminals, but not in the active zone (Nyiri *et al.*, 2005). Some endocannabinoids (e.g. 2-arachidonoylglycerol, 2-AG) are released from the postsynaptic membrane in response to the activation of primary transmitters (e.g. Glu and 5-HT; Katona *et al.*, 2006; Balazsa *et al.*, 2008). They stimulate extrasynaptic and perisynaptic CB1 receptors and inhibit transmitter release (Wilson and Nicoll, 2002; Lau and Schloss, 2008; Lau *et al.*, 2009) as a function of postsynaptic activity (Katona *et al.*, 2006). Other endocannabinoids such as anandamide, on the other hand, may have a presynaptic origin, since N-acylphosphatidylethanolamine-hydrolysing phospholipase D, which is a recently identified synthesizing enzyme of this bioactive lipid (Okamoto *et al.*, 2007), is localized to the presynaptic site (Nyilas *et al.*, 2008). The presynaptic release of anandamide, however, has not yet been shown.

It is very likely that these non-synaptically located CB1 receptors are the possible targets of endogenous cannabinoids and cannabis, which is used as a recreational drug. The question arises as to whether or not endogenous cannabinoids can tonically activate these CB1 receptors. Since the CB1 antagonist (SR141716A) failed to affect GABA release (Schlicker *et al.*, 1997), it seems likely that there is no tonic influence on GABAergic inhibitory transmission. In contrast, it is very likely that, when taken as a recreational drug of abuse, cannabis can tonically inhibit GABA release, thereby removing the inhibition of DA release by GABA (Ferraro *et al.*, 1996) in the nucleus accumbens. As a consequence of this disinhibitory action of cannabis, the release of DA increases and could then positively influence the reward system (Sperlagh *et al.*, 2009).

#### *Ionotropic receptors (Table 2)*


The majority of synapses within the CNS use mainly ionotropic receptors (receptor ion channels, transmitter-gated channels), because these receptors are equipped for fast (submillisecond) activation, making them able of transmitting information with minimal delay. Nevertheless, it has been discovered, that they are also abundant extrasynaptically (see Table 2).

**Glutamate receptors.** In spite of the predominantly synaptic nature of glutamatergic transmission (Table 1), different types of non-synaptic communication have been proven in various parts of the CNS (Vizi and Mike, 2006). Glutamate that is released into the synaptic gap can 'escape' from the synapse (Asztely *et al.*, 1997; Kullmann, 2000), which enables it to reach extrasynaptic receptors (Pankratov and Krishtal, 2003). This phenomenon is referred to as 'spillover' (Asztely *et al.*, 1997; Rusakov *et al.*, 1999), which could appear at extreme activity of the synapse (Drew *et al.*, 2008).

Ionotropic glutamate receptors are classified into three major groups, AMPA, kainate and NMDA receptors. All three types can be found at extrasynaptic locations (Table 2); however, this does not prove that they are functional extrasynaptically. Extrasynaptic AMPA receptors on neurons (mostly of the same subtype as synaptic receptors – Lu *et al.*, 2009) may only serve as spare receptors: a reservoir for synaptic plasticity. On the other hand, NMDA and kainate recep-

tors show subtype-specific subcellular localization, and different subtypes have been shown to be involved in synaptic and non-synaptic functions.

NMDA receptors have a tetrameric structure that is composed of two NR1 subunits and at least one type of NR2 subunit (A, B, C or D). The NR2A or NR2B subunits (Figure 3)



**Figure 3** Relevance of NMDA receptor localization composed of NR1/NR2A or NR1/NR2B assemblies. It was proposed that NR1/NR2A receptors, which have a lower affinity for glutamate and glycine than NR1/NR2B receptors (Yamakura and Shimoji, 1999), reside in the synapse, while NR1/NR2B receptors are located at extrasynaptic sites. However, it has been shown that this separation evolves only during postnatal development when the NR1/NR2B receptors exchange for NR1/NR2A receptors in the synapse (Tovar and Westbrook, 1999). This dynamic view of NMDA receptor subtypes was completed by the observation that NMDA receptors can move laterally between synaptic and extrasynaptic sites within the plasma membrane (Tovar and Westbrook, 2002; Triller and Choquet, 2005). Groc *et al.* have shown that NR1/NR2A assemblies diffuse laterally more than two orders of magnitude slower than NR1/NR2B receptors (Groc *et al.*, 2009; Groc *et al.*, 2006; see thick purple and thin green arrows representing the mobility of NR1/NR2A and NR1/NR2B assemblies, respectively). The different subcellular locations of NR1/NR2A and NR1/NR2B assemblies determine the source of glutamate being able to activate these receptors. While the NR1/NR2A assemblies are activated by glutamate released in the synapse, the origin of glutamate activating NR1/NR2B assemblies can be: (i) the spillover of synaptic glutamate following high frequency stimulation (Huang, 1998; Vizi and Mike, 2006), (ii) the reversal of glutamate transporters (EAAT1-4) mainly in neurons (Rossi *et al.*, 2007), (iii) the vesicular glutamate release from glial processes in the tripartite synapses (Haydon and Carmignoto, 2006; Perea *et al.*, 2009). The latter is the basis of glia-neuron interaction and can affect both the presynaptic and postsynaptic NR1/NR2B assemblies and either enhances synaptic strength or mediates slow inward currents, respectively (Fellin *et al.*, 2004; Jourdain *et al.*, 2007). Functions of synaptic NR1/NR2A and extrasynaptic NR1/NR2B receptors are opposing to some extent (Haydon and Carmignoto, 2006): (i) NR1/NR2A promote, while NR1/NR2B inhibit AMPA receptor trafficking (Kim *et al.*, 2005), (ii) NR1/NR2A induces the neuroprotective CREB (cAMP response element binding protein) pathway, while NR1/NR2B shuts it down and promotes neuronal death (Hardingham *et al.*, 2002), (iii) NR1/NR2A induces long-term potentiation, while NR1/NR2B seems to be relevant in long-term depression (Liu *et al.*, 2004a; Massey *et al.*, 2004).

are predominant in humans (Wenzel *et al.*, 1997; Cull-Candy *et al.*, 2001). It has been observed that NR2A subunit-containing NMDA receptors are predominantly located synaptically, whereas NR2B subunits are primarily found in extrasynaptic locations (Stocca and Vicini, 1998; Mohrmann *et al.*, 2000; Townsend *et al.*, 2003), although this segregation is not an absolute rule (Thomas *et al.*, 2006; Gardoni *et al.*, 2009). A recent development has shown that NR2B-containing subtype of Glu receptors is also part of the postsynaptic compartment in the synapse, and is involved in the physiological induction of long-term potentiation (LTP; Gardoni *et al.*, 2009). Nevertheless, accumulating evidence indicates that the synaptic NR2A receptors mediate synaptic plasticity and cell survival signals, whereas extrasynaptic NR2B receptors are responsible for excitotoxicity (Hardingham and Bading, 2003; Vanhoutte and Bading, 2003; Zhou and Baudry, 2006; Liu *et al.*, 2007). It has also been shown that NMDA receptors evoke the production of peroxynitrite (product of superoxide and the diffusible nitric oxide interaction), the highly toxic reactive oxygen species, in regions vulnerable to ischemia but not in regions relatively resistant to ischemic injury (Fekete *et al.*, 2008).

As for kainate receptors, although they mediate synaptic transmission by postsynaptic receptors, their predominant role appears to be presynaptic modulation of glutamate and GABA release (Jaskolski *et al.*, 2005; Lerma, 2006; Coussen, 2009). They are located presynaptically near the release site and in the somatodendritic compartment both within the postsynaptic density (e.g. in the hippocampal mossy fiber synapse) and extrasynaptically (Jaskolski *et al.*, 2005; Table 2).

Non-synaptic AMPA and kainate receptors are commonly present in glial cells (astrocytes, oligodendrocytes and microglia; Matute *et al.*, 2001). Different subtypes of NMDA receptors, including NR3A, but not NR3B, have also been shown in the myelinating processes of oligodendrocytes, where the small intracellular space could lead to a large rise in glutamate concentration and mediate excitotoxic injury (Karadottir *et al.*, 2005; Salter and Fern, 2005).

**GABA<sub>A</sub> receptors.** It is generally accepted that the main inhibitory transmitter, GABA is predominantly synaptic (Figure 1, Table 1), and functions as an agonist with high potency but low efficacy. The first evidence that GABA<sub>A</sub> receptors could be localized extrasynaptically was found by Somogyi *et al.* (1989) using an immunogold technique in cerebellar granule cells. This observation was a milestone in neuroscience, because it provided the first convincing morphological evidence that ionotropic receptors are also expressed extrasynaptically. In the same laboratory, it was shown that although the density of GABA<sub>A</sub> receptors is higher within the synapses, with regard to the overall number, extrasynaptically localized receptors outnumber those that are present at the synapse (Nusser *et al.*, 1995). The presence of the  $\delta$  subunit was shown to be a characteristic of GABA<sub>A</sub> receptors that are localized extrasynaptically (Nusser *et al.*, 1998) or perisynaptically (Huang and Pickel, 2002). Herd *et al.* (2008) demonstrated that tonic currents are predominantly mediated by  $\beta_2$  containing GABA<sub>A</sub> receptors, while synaptic currents are mediated by GABA<sub>A</sub> receptors that contain the  $\beta_3$  subunit. These findings indicate that non-synaptic GABA

receptors are different from those that are present at the synapse. While the millimolar concentration of GABA in the synapse (Mody *et al.*, 1994; Jones and Westbrook, 1996) results in a phasic and transient inhibition of the postsynaptic cell, GABA escaping from the synapse can result in a persistent or tonic activation of non-synaptic GABA<sub>A</sub> receptors (Semyanov *et al.*, 2004; Farrant and Nusser, 2005). Subsequent studies have indicated that GABA-mediated tonic conductance exists in many different nerve cells of distinct brain areas (Farrant and Nusser, 2005), such as the cerebellum, hippocampus, dentate gyrus, cortex, thalamus and brainstem (Glykys and Mody, 2007). In fact, GABA-mediated tonic conductance on dendrites and cell bodies has also been found under *in vivo* conditions (Chadderton *et al.*, 2004).

Simulation studies suggest that a decrease in membrane resistance at the opening of GABA<sub>A</sub> receptor channels efficiently prevents propagation of APs into presynaptic terminals (Segev, 1990). Verdier *et al.* (2003) recently reported that GABA<sub>A</sub> receptors could affect AP propagation in branches of afferents in brainstem slices. The  $\delta$  subunit-containing GABA<sub>A</sub> receptors that include either  $\alpha 4$  or  $\alpha 6$  subunits characteristically have a high affinity and a low efficacy for GABA (Saxena and Macdonald, 1996; Brown *et al.*, 2002), and low degree of desensitization to agonists that are continuously present in the extracellular space (Wohlfarth *et al.*, 2002; Bianchi and Macdonald, 2003; Table 2). Their EC<sub>50</sub> is in the tens of nM range, which is well within the concentrations that are present in the extracellular space (Nyitrai *et al.*, 2006).

A novel form of GABAergic inhibition has recently been described (Tamas *et al.*, 2003; Szabadics *et al.*, 2007). A class of cortical GABAergic interneurons, the neurogliaform cells produce slow ionotropic responses (IPSCs lasting for tens of milliseconds), with properties that are intermediate between phasic and tonic inhibitory signalling. Most (78%) axonal varicosities of this cell type lack synapses, but non-synaptic release from its unusually dense axonal arborization can provide an effective widespread inhibition in the local environment (Olah *et al.*, 2009). Slow ionotropic responses have proven to be particularly sensitive to the modulation of reuptake, as well as to certain types of benzodiazepines.

**5-HT<sub>3</sub> receptors.** 5-HT<sub>3</sub> receptors have been shown to be present on axons and presynaptic terminals, and in large dense-core vesicles, however, they do not associate with the synaptic differentiation of terminals (van Hooft and Vijverberg, 2000; Miquel *et al.*, 2002; Huang *et al.*, 2004). Immunoreactive terminals containing abundant dense-core vesicles often lack recognizable synapses, but those containing few dense core vesicles form asymmetric or symmetric synapses (Huang *et al.*, 2004). 5-HT<sub>3</sub> receptors are present on spines and distal portion of dendrites, mainly within the hippocampus. 5-HT<sub>3</sub> receptors are concentrated at the postsynaptic differentiation, but can also be found non-synaptically on spines and dendrites, occasionally in a perisynaptic location (Miquel *et al.*, 2002; Huang *et al.*, 2004), and are also localized on glial processes (Huang *et al.*, 2004).

**Nicotinic receptors.** There are at least 12 different functional pentamer nACh receptors (Lendvai and Vizi, 2008), which can be found in extrasynaptic locations on different neurons

in the hippocampus (Hill *et al.*, 1993; Ullian and Sargent, 1995; Fabian-Fine *et al.*, 2001; Jones and Wonnacott, 2004). It was shown in the ventral tegmental area (Jones and Wonnacott, 2004) with both pre- and post-embedding electron microscopy that 27% of  $\alpha 7$ -nACh receptors are extrasynaptic on the glutamatergic axons, 61% are perisynaptic and only 11% are expressed at the active zone, opposed to the postsynaptic density. Functionally, it was shown that tonic activation of extrasynaptic  $\alpha 7$ -nAChRs by cholinergic agonists – depending on timing – either activates or depresses back-propagating action potentials (Rozsa *et al.*, 2004; 2008; Lendvai and Vizi, 2008) of CA1 stratum radiatum interneurons. A stimulation paradigm that normally failed to produce LTP in the Schaffer collaterals was able to boost LTP production when sufficient depolarization was evoked by  $\alpha 7$ -nAChR stimulation and was timed to coincide with the somatic stimulation. Similarly, the activation of non-synaptic nAChRs resulted in a potentiation of synaptic glutamatergic inputs (Vizi *et al.*, 2004a). Szabo *et al.* (2008) showed that the activation of non-synaptic nAChRs results in  $Ca^{2+}$  transients in pyramidal spines and induces firing of APs. These findings indicate that ACh or its metabolite choline, which itself is an endogenous agonist of the  $\alpha 7$  subtype of nAChRs, are able to produce ‘postsynaptic’ signalling following diffusion far away from their release sites. It seems likely that non-synaptic nAChR activation through ionotropic channels is a potential alternative for metabotropic muscarinic modulation of hippocampal plasticity, which exhibits a rather slow action (Buzsaki, 2002).

**Glycine receptors (GlyRs).** The subunits ( $\alpha(1-4)$  and  $\beta$ ) of pentameric GlyRs form heteromeric or  $\alpha$  homomeric receptors.  $\alpha$  subunits are responsible for agonist binding (Lynch, 2004), while the  $\beta$  subunit binds the anchoring protein gephyrin with high affinity (Kirsch *et al.*, 1993; Meyer *et al.*, 1995), which immobilizes postsynaptic GlyRs at the subsynaptic cytoskeleton. Homomeric  $\alpha$  GlyRs, which do not contain the  $\beta$  subunit, are expected to freely diffuse in the plasma membrane, thus representing extrasynaptic GlyRs.

**P2X receptors.** Adenosine triphosphate (ATP) is a transmitter released from both axon terminals along with another transmitter (Buzsaki, 2002; Zimmermann, 2008; Abbracchio *et al.*, 2009) and from the postsynaptic sites in response to receptor stimulation by the primary transmitter (Vizi *et al.*, 1992a; Sperlagh *et al.*, 2002). Glia is also a major source of extracellular ATP (Cunha, 2008). ATP can act on postsynaptic sites, where it is released from termed ‘cascade’ transmission (Vizi *et al.*, 1992a) or on presynaptic sites. Ionotropic receptors of ATP are called P2X receptors, and seven subunits ( $P2X_{1,2,3,4,5,6,7}$ ) can form homo- or heterotrimeric ion channels (Burnstock, 2008b). P2X receptors are involved in fast synaptic transmission (North and Verkhatsky, 2006), where ATP is co-released with glutamate, and P2X receptors have been shown to be located both within glutamatergic synapses and perisynaptically (Rubio and Soto, 2001). P2X receptors mediating non-synaptic, especially neuro-glia interactions are even more widespread (Abbracchio and Ceruti, 2006). Purinergic receptors are also involved in the modulation of transmitter release (Sperlagh *et al.*, 2007a). P2X receptors typically facilitate release of other transmitters, including GABA

(Sperlagh *et al.*, 2002) and NA (Papp *et al.*, 2004), as well as the carrier-mediated release of Glu from the hippocampus.

*Non-synaptic transporters*

As in the case of the discovery of non-synaptic localization of different receptors, the observation that several transporters are also localized extrasynaptically (Huang *et al.*, 2004; Torres and Amara, 2007; Daws, 2009) has also helped to change our view. There are two ion-coupled co-transporter families that are responsible for the transport of transmitters: the  $Na^+/Cl^-$ -dependent carriers for monoamines (NA, dopamine and serotonin), glycine and GABA, and the  $Na^+$ -dependent transporters for glutamate and aspartate (Torres and Amara, 2007). The role of these transporters is to remove transmitters that have been released into the synaptic gap or are present in the extracellular space in millimolar to nanomolar concentrations and to return them to nerve terminals. This would prevent postsynaptic receptors from becoming desensitized and help the synapses recover so they can relay transmissions again at a high frequency.

It is generally accepted that transporters are equally capable of operating in either the forward or reverse direction, and that they play a critical role in setting the ambient concentrations of different transmitters and the measure of tonic inhibition (Richerson and Wu, 2003). When the transporters operate in the forward direction, they clear neurotransmitters from the synaptic cleft, which recycles the transmitters after release. When they operate in the reverse direction, they release transmitters that are present in the cytoplasm in a  $[Ca^{2+}]_o$ -independent manner (Table 3). This type of release is not the subject of presynaptic inhibition (Vizi *et al.*, 1996), and ensures a permanent and continuous release of transmitters into the extrasynaptic space.

A further characteristic of transporters that are expressed on non-synaptic varicosities is their indiscriminate nature. Monoamine transporters are members of the same structural family with a high degree of homology (Amara and Kuhar, 1993) and are able to take up other transmitters besides their own (Devoto *et al.*, 2001; Moron *et al.*, 2002; Vizi *et al.*, 2004b) from the extracellular space. Monoaminergic varicosities are also able to release different transmitters in response to depolarization. This is also the case with noradrenergic varicosities, which are able to both take up and release DA (Devoto *et al.*, 2001), as well as the case with serotonergic terminals, which are able to take up and release NA (Vizi *et al.*, 2004b). This effect is independent of their machinery, which synthesizes their transmitters. It means that there is a

**Table 3** Origin of transmitters released in response to neuronal firing and reversal of transporter (for details see the text)

	Transmitter release due to	
	Axonal firing	Reversed transporter function
Origin of transmitter	Vesicle	Cytoplasm
$[Ca^{2+}]_o$ -dependence	Yes	No
Presynaptic modulation	Yes	No
Lowering temperature	Not sensitive	Sensitive



**Table 4** Concentrations of transmitters in the vesicle, cytosol, synapse and extracellular space

Concentration of transmitters	Vesicle	Cytosole	Synapse	Extracellular space
Acetylcholine <sup>1</sup>	~100 mM	n.d.	~3 mM	0.01–3 µM
Noradrenaline <sup>2</sup>	~200 mM	n.d.	~1 mM	14–78 nM
Dopamine <sup>3</sup>	300–350 mM	n.d.	n.d.	14 nM–0.7 µM
GABA <sup>4</sup>	~100 mM	1–7 mM	~1 mM	0.2–0.8 µM
Glutamate <sup>5</sup>	~100 mM	2–10 mM	~1 mM	0.03–3 µM
Serotonin <sup>6</sup>	~90 mM	n.d.	~6 mM	1–100 nM

Our calculation based on:

<sup>1</sup>Descarries *et al.* (1997) and Zhang *et al.* (1996).

<sup>2</sup>Abercrombie *et al.* (1988) and Tso *et al.* (2004).

<sup>3</sup>Chen and Budygin (2007), Cragg *et al.* (2001), Jaffe *et al.* (1998), Nirenberg *et al.* (1997) and Pothos *et al.* (1998).

<sup>4</sup>Jones and Westbrook (1995), Lerma *et al.* (1986), Mody *et al.* (1994) and Otsuka *et al.* (1971).

<sup>5</sup>Barbour *et al.* (1994), Cavelier *et al.* (2005), Clements *et al.* (1992), Nedergaard *et al.* (2002) and van der Zeyden *et al.* (2008).

<sup>6</sup>Barnes *et al.* (1992); Bunin and Wightman (1998) and Crespi *et al.* (1988).

n.d., no data available.

functional overlap between different monoaminergic transporters. In the synapse, this phenomenon either does not happen or is negligible, because the concentration of the primary transmitter is about 1000 times higher in the synapse than the concentration of another transmitter diffusing from the extrasynaptic space (see Table 4).

Transporters are able to generate current in response to substrates through a channel-based mechanism (Bringmann *et al.*, 2009). This was strongly supported by electrophysiological evidence for the dopamine transporter (DAT) (Ingram *et al.*, 2002; Carvelli *et al.*, 2004) and for the Glu transporter (Veruki *et al.*, 2006). It was shown that transporter-associated anion current produced by Glu can hyperpolarize the presynaptic terminal and thus reduce transmitter release (Veruki *et al.*, 2006). In contrast, substrates of DAT, such as DA and amphetamine, activate anion conductances and increase the firing rate of dopaminergic neurons (Ingram *et al.*, 2002). It appears likely that the explanation for these observations is that the Cl<sup>-</sup> reversal potential might vary in different neurons.

Transporter trafficking to and from the plasma membrane (Melikian, 2004) could be a response to chronic stress, which would result in a change in neuronal function (Torres *et al.*, 2003; Miner *et al.*, 2006; Robinson, 2006; Torres and Amara, 2007). The finding that chronic stress results in increased plasma membrane expression of the NA transporter (NET) in the prefrontal cortex suggests that there is an activity-dependent change in the subcellular localization of NET (Miner *et al.*, 2006), which is otherwise 85–90% localized in the cytoplasm. Agonists for the serotonin transporter (SERT; Zhu *et al.*, 2004) and NET (Zhou *et al.*, 1998) increase the delivery of transporter molecules to the plasma membrane. These studies illustrate that, even during axonal firing, the transmitter release and uptake machinery of the neuronal network is able to adapt to the new circumstances. Transporter molecules could traffic between intracellular pools and the cell surface (Miner *et al.*, 2003). It remains to be discovered whether or not these translocation changes of transporters occur extrasynaptically or in synapses. More than likely, both scenarios are involved.

**NET.** The NET is present on noradrenergic fibres at non-synaptic sites (Schroeter *et al.*, 2000; Moron *et al.*, 2002; Miner *et al.*, 2003; Liprando *et al.*, 2004; Daws, 2009) and serves as a

target for multiple antidepressants and psychostimulants. Immunogold labelling of the NET was more frequently observed in the cytoplasm than on the plasma membrane (Miner *et al.*, 2003). The NET is also able to take up DA from the extracellular space (Moron *et al.*, 2002). The inhibition of the NET therefore results in an increase in the extracellular levels of both NA and DA.

**Dopamine transporter.** Ultrastructural analysis of DAT immunolabeling revealed that the DAT is preferentially localized non-synaptically (Smiley *et al.*, 1994; Nirenberg *et al.*, 1996; 1997) rather than to the synaptic part of the presynaptic membrane. This was supported by voltammetric investigations that showed the re-uptake of DA at a distance from the original release sites (Garris and Wightman, 1994; Garris *et al.*, 1994). Many current pharmacological or recreational psychotropic drugs, including psychostimulants, antidepressants and neurotoxins, exert their actions, at least in part, via an interaction with the DAT (Sotnikova *et al.*, 2006). Studies in DAT-KO mice have confirmed this hypothesis, showing a total lack of 1-methyl-4-phenyl-1,2,3,6 tetrahydropyridine (MPTP) neurotoxicity in these mice (Gainetdinov *et al.*, 1997; Bezard *et al.*, 1999). The transporter is also of great importance for the toxic action of drugs of abuse (such as methamphetamine and ecstasy), which exert their actions through mobilization of DA from intracellular stores to the extracellular space via DAT-mediated outward transport (Fleckenstein *et al.*, 1997; Fumagalli *et al.*, 1998).

**SERT.** The SERT is mainly localized extrasynaptically on the plasma membrane (Zhou *et al.* 1998; Tao-Cheng and Zhou, 1999) of intervaricosity segments of the axons (Tao-Cheng and Zhou, 1999; Huang and Pickel, 2002; Huang *et al.*, 2004) shown by immunogold immunocytochemistry (Tao-Cheng and Zhou, 1999). This distribution is in contrast to what has been observed in the soma and dendrites, where the labelling is prevalent in the cytoplasm. Neurochemical evidence showed that the SERT is capable of taking up NA from the extracellular space and releasing it in response to axonal stimulation (Vizi *et al.*, 2004b). The non-synaptic SERT, but not those in the synaptic gap, is the main target of the selective transporter inhibitors, as well as many drugs of abuse, such as cocaine, ecstasy and methamphetamine.



**Glutamate transporter (excitatory amino acid transporter, EAAT).** Five EAATs have been identified in the brain. EAAT1 (GLAST) and EAAT2 (GLT-1) are predominantly expressed in astrocytes, while EAAT3 (EAAC1) and EAAT4 are localized in neurons. EAAT5 is expressed primarily in the retina (Amara and Fontana, 2002). The majority of Glu uptake occurs in glia rather than in neurons (Yi and Hazell, 2006). The EAAT3 transporter is expressed both in the synaptic cleft and non-synaptically (Conti *et al.*, 1998). Electrophysiological studies have shown that inhibition of Glu uptake leads to Glu spillover (Diamond, 2002). While the activity of Glu transporters minimally affects synaptic currents (Rusakov and Kullmann, 1998), EAATs represent a major mechanism for removing extrasynaptic Glu in order to ensure a high signal-to-noise ratio and to enable a high frequency of synaptic transmission (Beart and O'Shea, 2007). Although an important role of the EAATs is to maintain the extracellular Glu concentration below neurotoxic levels, the  $[Ca^{2+}]_o$ -independent (Sperlagh *et al.*, 2007b) reversal of Glu transporter activity after ischemic injury, when the  $Na^+/K^+$  electrochemical gradient collapses, seems to be involved in the accumulation of extracellular Glu concentrations.

**GABA transporters.** Inhibition of the GABA transporter results in an excessive accumulation of GABA in the synaptic gap, which is followed by a spillover of GABA (Isaacson *et al.*, 1993). In addition to the vesicular release, GABA can be released also by the reversal of the GABA transporter, which can be induced by pathological or therapeutic conditions, such as high intracellular sodium concentration, which occurs during ischemia, depolarization due to the accumulation of extracellular potassium during seizure activity, or in the presence of the anticonvulsants gabapentin or vigabatrin (Wu *et al.*, 2001; 2007; Richerson and Wu, 2003). The non-synaptic GABA transporter-1 regulates tonic inhibition by controlling the ambient, non-synaptic, extracellular GABA concentrations. The increase in tonic GABAergic inhibition that resulted from the activation of non-synaptic, high-affinity  $GABA_A$  receptors, which do not desensitize, is related to the size of the reversal of the GABA transporter. It is supposed that this mechanism may operate physiologically as well, and can serve as a delicate sensor of intracellular sodium and extracellular GABA concentration (Wu *et al.*, 2007). Release of GABA by reverse operation of GABA transporters have been shown to be evoked by increased glutamate uptake. Increased tonic conductance by GABA receptor activation thus provides a compensatory mechanism for glutamatergic overexcitation (Heja *et al.*, 2009).

**Choline transporter.** Elimination of released acetylcholine is done not by transport but by enzymatic degradation. It is first decomposed by cholinesterase to choline and acetate, followed by the uptake of choline (Ribeiro *et al.*, 2007).

### Mechanisms of transmitter release into the extrasynaptic space

There are several major sources of transmitters (endogenous substances) for non-synaptic release (Figure 4, Table 3).

1 Exocytotic (vesicular) release of transmitters from non-synaptic varicosities of for diffusion (Figure 4). Most typically occur via monoaminergic and cholinergic axons, which can release transmitters via exocytosis into the extracellular space of the brain (Descarries *et al.*, 1997). This release is  $[Ca^{2+}]_o$ -dependent, is related to axonal firing and is subject to presynaptic modulation (Table 3). This type of release has long been known to involve peptides and, in the intestine, ACh.


The terminal axons of the cholinergic Auerbach plexus in the gut have many varicosities and are associated with the accumulations of small vesicles. These represent sites of exocytotic release of ACh; however, there are no post-junctional specializations on the smooth muscle cells (Paton *et al.*, 1971). The distance between varicosities and smooth muscle cells is about 100–200 nm, and ACh that is released from the boutons has to diffuse across these distances (Paton *et al.*, 1971). Muscarinic receptors of the  $M_1$  subtype are evenly distributed on the smooth muscle surface and have a high affinity ( $EC_{50}$  for ACh = ~100 nM).

Similar exocytotic release of NA for diffusion is also known to occur at different neuroeffector transmission sites in the periphery (Burnstock, 2008a). The distance between noradrenergic boutons and effector smooth muscle cells may vary considerably, and clefts range from 15–20 nm (e.g. in the vas deferens and iris) to 1–2  $\mu$ m (e.g. in the large arteries). NA is released *en passage* from varicosities during the conduction of APs along the axon. The APs are propagated along the varicose axon and release transmitters from only some of the varicosities that the action potentials encounter (Jackson and Cunnane, 2001).

Vesicular release of glutamate has been described in glial processes, which participate in the formation of the tripartite synapse (Figure 3).

2 Spillover of transmitters from the synapse (see Figures 3 and 4). Transmitters can 'escape' from the synapse (Asztely *et al.*, 1997), especially during high frequency firing, and spill over, which results in their diffusion far away from the original release sites. These 'escaped' transmitters can then hit extrasynaptic receptors, which could potentially appear as extreme activity of the synapse.

3 Transmitter release by the reverse operation of transporters (carrier-mediated release; Table 3). In certain pathological conditions, such as ischemia or epilepsy, the high  $Na^+$  load (Sheldon *et al.*, 2004; Fekete *et al.*, 2009) can reverse the operation of the transporter for the transmitter, resulting in release instead of reuptake of transmitters as a form of 'carrier-mediated release'. This release is  $[Ca^{2+}]_o$  independent, and is not related to axonal firing or subject to presynaptic modulation (Vizi *et al.*, 1986; Ikeda *et al.*, 1989). It is worth mentioning that transporters are capable of a channel-like operation. In the case of monoamine transporters, not only ions but the transmitter molecule itself is able to cross the membrane through the channel (DeFelice and Goswami, 2007) and to be released into the extracellular space. It has been proposed that reverse operation of transporters may be a normal physiological mechanism for dynamic modulation of ambient levels of certain transmitters (Richerson and Wu, 2003).



**Figure 4** Scheme of different types of chemical communication, including synaptic transmission when the transmitter released into the synaptic gap acts on postsynaptic low affinity receptors located in the synaptic gap, and non-synaptic communication when the transmitter spills over from the synaptic gap or is released from a bouton without making synaptic contact and reaches its target non-synaptic (perisynaptic or extrasynaptic) receptors by diffusion. The distance could be from a few hundred nm to a few hundred μm. Note the significant difference in concentration in the vesicle (~100 mM), in the synaptic gap (~1 mM) and in the extrasynaptic space (~0.01–3 μM). Concentrations in the figure are approximate values from the literature (see Table 4).

4 Other non-exocytotic mechanisms (Vizi and Mike, 2006; Malarkey and Parpura, 2008; van der Zeyden *et al.*, 2008) can also significantly contribute especially to ambient levels of certain transmitters. Included among these mechanisms are *release through gap junction hemichannels* (Ye *et al.*, 2003), *P2X7 purinergic receptors* (Duan *et al.*, 2003), *cystine-Glu exchange* (Cho and Bannai, 1990), *Ca<sup>2+</sup> ion-dependent release* (Bezzi *et al.*, 1998), *volume-sensitive organic anion channels* (Del Arco *et al.*, 2003) and *swelling-induced release* (Kimmelberg *et al.*, 1990). ATP-induced activation of the homomeric P2X<sub>7</sub> receptor releases Glu from astrocytes into the extracellular space via a pore that is able to allow molecules as large as 900 Da to pass through (North, 2002).

This type of release is [Ca<sup>2+</sup>]<sub>o</sub>-independent (Papp *et al.*, 2004; Malarkey and Parpura, 2008).

### Extracellular space as a communication channel

The extracellular space, which accounts for 15–30% of the total brain volume, has been shown to be an important communication channel (Sykova and Nicholson, 2008). It is filled with interstitial fluid that has a composition similar to the cerebrospinal fluid. In addition to its ion content, the extracellular space has an ambient level of various endogenous substances.

In the normal adult brain, the extracellular space may be reduced to 5% during ischemia or epilepsy (Elsen *et al.*, 2006). This type of reduction in the extracellular space can result in a significant increase in cell volume (Franco *et al.*, 2008) and an increase in the concentration of Glu, ions and neuroactive substances that have all been released into the extracellular space (Sykova, 2004).

#### *Levels of transmitters in the extracellular space*

The determinants of extracellular transmitter concentrations include the rate of release, diffusion (regulated by properties of the extracellular space, such as geometry, viscosity, matrix molecules, charged, or transiently binding molecules) and clearing processes (uptake by glia and neurons, enzymatic degradation, and removal through the blood-brain barrier; Sykova and Nicholson, 2008; Sykova and Vargova, 2008). The ambient levels of extrasynaptic transmitters can tonically inhibit or enhance both the synaptic and non-synaptic neuronal functions (release, neuronal excitability; Carmignoto and Fellin, 2006; Semyanov, 2008). A prerequisite of these functions is that neuronal networks are equipped with extrasynaptic receptors.

Although the concentration of transmitters is about 100 mM in the vesicles and 1–3 mM in the synaptic cleft following its release and dilution, the extrasynaptic concentration of transmitters is much lower, about 0.01–3  $\mu$ M (Figure 4, Table 4). When a transmitter is released extrasynaptically by a varicosity, it can freely diffuse into all directions of the space, thus the concentration drops as a cubic function of the distance (Vizi, 2000).

Using Glu microsensors in the hippocampus, the *in vitro* basal extracellular level of Glu was found to be 1.7  $\mu$ M (van der Zeyden *et al.*, 2008). There are, however, large differences in the level of Glu across brain regions and between different studies ranging from 1 to 45  $\mu$ M, which may be partially due to methodological differences (van der Zeyden *et al.*, 2008). In contrast to these studies, Cavelier *et al.* (2005) estimated extrasynaptic, resting Glu concentrations to be much smaller (0.03–0.08  $\mu$ M at 25–35°C) by measurement of the size of tonically activated NMDA receptor (AP-5 suppressible) current. The *in vivo* concentration of Glu was reported to be 18.2  $\mu$ M in the striatum (van der Zeyden *et al.*, 2008), while Rahman *et al.* (2005) found it to be only 1.4–2  $\mu$ M. The basal levels of extracellular GABA and glycine were found to be 0.27, and 4.95  $\mu$ M, respectively (Kennedy *et al.*, 2002). Whereas the resting extracellular adenosine concentration in the brain is 30–300 nM (Rudolphi and Schubert, 1997), it can reach 10–50  $\mu$ M following a 15 min period of ischemia (Hagberg *et al.*, 1987; Pugliese *et al.*, 2009). The extracellular concentration of ATP is as low as 100 nM (Llaudet *et al.*, 2005). Basal extracellular levels of DA, NA and 5-HT are given in Table 4. Uptake inhibitors are able to produce an increase in the extracellular levels of transmitters, showing that transporters have a tonic effect on ambient transmitter level (Segovia and Mora, 2001; Wu *et al.*, 2007).

Neural stimulation results in an increase in the concentration of neurotransmitter level. Local electrical stimulation (20 pulses, 100 Hz) evoked a DA efflux of 315 nM within approximately 1 s in the striatum. In the dorsal raphe, 5-HT peaked at

16 nM under the same stimulation parameters, whereas in the ventral bed nucleus of the stria terminalis, the NA concentration peaked at 78 nM (Tso *et al.*, 2004). Using fast-scan cyclic voltammetry, the concentration of extracellular DA during somatodendritic activation in guinea pig brain slices was site specific, with a significantly higher level in caudal slices (0.48  $\mu$ M) than in rostral slices (0.16  $\mu$ M). In the ventral tegmental area, the stimulated DA levels peaked at 0.74  $\mu$ M (Cragg *et al.*, 1997). Cragg *et al.* (2001) calculated that DA, which is released into the extracellular space in the substantia nigra pars compacta and can diffuse up 200  $\mu$ m from the release site, reached a concentration of 14 nM. This was high enough to activate high-affinity DA receptors ( $EC_{50}$  for  $D_2$  receptor activation = 1–20 nM; Levant, 1997).

#### *Drug concentrations in the extracellular space*

As a general rule, it has been observed that non-synaptic receptors have higher agonist affinity than their synaptic counterparts. This does not mean that they also will have higher affinity to drug molecules as well. However, the general assumption that drugs preferentially affect non-synaptic receptors is nevertheless true, simply because the majority of CNS drugs affect transmitter levels, for example, by acting on the modulation of release (e.g. haloperidol), on the reuptake process (e.g. fluoxetine) or on enzymatic degradation (e.g. selegiline, vigabatrin, moclobemide or galantamine). The effect on transporters and degrading enzymes minimally affect concentrations within the synapse at the time after the release, but has a major role in determining how far the transmitter will diffuse (see next section), that is the relative contribution of non-synaptic receptors.

While neurotransmitters are typically hydrophilic molecules, and therefore are largely confined to the extracellular space, CNS drug molecules are typically amphiphilic or lipophilic (since this is a requirement for blood-brain barrier penetration). Diffusion of these molecules therefore is not restricted to the extracellular space, but involves membrane-water phase partitioning, as well as intramembrane diffusion. For this reason, apparent diffusion coefficients are lower than that of hydrophilic molecules by one to two orders of magnitude (Chesney *et al.*, 2003; Gredell *et al.*, 2004), and free extracellular concentration can be several orders of magnitude lower than total brain concentration (Summerfield *et al.*, 2007; Friden *et al.*, 2009).

#### *Diffusion of transmitters to reach non-synaptic receptors and transporters*

The principal question in the case of any receptor is whether the transmitter can reach the receptors at sufficiently high concentration. The simple question: 'How far can transmitter molecules get within a given time?' addresses three aspects of transmitter diffusion: space, time and concentration.

In order to investigate these questions for different transmitters, we performed simulations of transmitter diffusion. A detailed description of the simulations and a presentation of some typical examples are given in the Supplementary Information (Appendix S1). The files used for simulation are also available.

For a typical transmitter at 1 ms after the release of a single vesicle, 95% of all released molecules will be within a sphere with a radius of  $\sim 2 \mu\text{m}$ . At the highest point of this diffusion wave, the concentration of the transmitter will be  $\sim 5 \mu\text{M}$ . In order to increase the distance 10-fold ( $\sim 20 \mu\text{m}$ ), even if no transporter activity is present, 100-fold this much time (100 ms) will be needed, and the concentration will drop 1000-fold ( $\sim 5 \text{nM}$ ). For the spillover of transmitters to perisynaptic receptors, a two dimensional model may be more appropriate. In this case concentration drops as a quadratic, not cubic function of distance (Slide S1).

We found that differences in the diffusion coefficient, tortuosity and the number of released molecules only had a minor effect on the pattern of diffusion (Slide S2). The most important determinants of the diffusion pattern were the density and binding kinetics of transporters (Slide S3). The third major determinant of the effective distance of non-synaptic transmission was the affinity of the receptor (Slide S4).

The relationship between non-synaptic diffusion and receptor affinity can be characterized by two useful terms: (i) 'maximum effective radius' is defined as the distance within which release of one quantum of transmitter reaches the  $\text{EC}_{50}$  concentration of the receptor, and (ii) 'maximum active lifetime' is the time during which the transmitter concentration remains above  $\text{EC}_{50}$  (Rice and Cragg, 2008). For example, for low affinity ( $\text{EC}_{50}$  between 1 and  $18 \mu\text{M}$ ; Neve and Neve, 1997) synaptic (mainly D1) dopamine receptors, the transmitter concentration that is enough for significant activation is present for  $\sim 1\text{--}2 \text{ms}$  (maximum active lifetime) and within  $\sim 1\text{--}2 \mu\text{m}$  (maximum effective radius) from the release site (Slide S4). In contrast, high affinity ( $\text{EC}_{50}$  between 10 and  $32 \text{nM}$ ; Neve and Neve, 1997) non-synaptic (mainly D2) dopamine receptors can detect release of single quanta from as far as  $\sim 4\text{--}5 \mu\text{m}$ , diffusion takes around  $10\text{--}20 \text{ms}$  to reach this far (see supplementary files). For synaptic AMPA receptors ( $\text{EC}_{50}$  between 300 and  $5000 \mu\text{M}$ ; Erreger *et al.*, 2004), the signal is effectively terminated within the synaptic cleft (maximum effective radius 300 nm, with a maximum active lifetime up to  $\sim 200 \mu\text{s}$  – as judged from the two-dimensional model). In contrast, for perisynaptic and extrasynaptic NMDA receptors ( $\text{EC}_{50}$  between 0.9 and  $13 \mu\text{M}$ ; Erreger *et al.*, 2004), even single quanta can be detected from  $\sim 1.5 \mu\text{m}$  distance, with a maximum active lifetime of  $\sim 1 \text{ms}$  (Slide S4). Similar maximum effective radius and maximum active lifetime values apply to extrasynaptic mGlu receptors as well, because their affinity is within the same range (Cartmell and Schoepp, 2000). The effect of non-instantaneous release was also simulated (Slide S5). Interestingly, the non-instantaneous nature of the release increased, not decreased intrasynaptic concentration within the first 0.2 ms.

## Summary

After slow acceptance, non-synaptic signalling in both the peripheral and central nervous systems is a rapidly expanding field of neuroscience. The results presented in this review may alter the way we think about separate synaptic and non-synaptic functions. Synaptic functions are thought to be responsible for definite, accurate tasks that require high speed

and precision, such as visual pattern recognition or motor pattern execution. Conversely, non-synaptic functions are thought to be responsible for the modulation and tuning of these processes, such as determining the level of arousal, the focus of attention or the emotional background. Note that almost all pharmacological manipulations of the CNS are restricted to this latter realm, predominantly involving non-synaptic functions.

This type of interaction assumes that the released transmitter is able to diffuse within the extracellular space and reach distant targets at relatively low concentrations. The conditions under which this system is efficient imply that the affinity of the targeted receptors for the neurotransmitter is very high, since the concentration of molecules decreases as a cubic function of distance (Vizi, 2000). The existence of non-synaptic receptors and transporters that have affinities in the nanomolar range agrees well with such an arrangement. Moreover, the fact that a transmitter that is released from different terminals diffuses in three spatial dimensions means that the affected receptors and transporters might be located on any part of the surrounding neurons, including the somata, dendrites, axons and varicosities. The non-synaptic localization of receptors and transporters provides an anatomical basis for potential therapeutic implications. These structures, which have no synaptic arrangements, are promiscuous and accessible to chemicals that are released from numerous synapses, non-synaptic boutons or both. In addition, the non-synaptic receptors and transporters are certainly accessible to medications that have an availability in the nanomolar and micromolar concentration range in the brain. Drugs that are given for neuropsychiatric diseases could therefore exert their effects via these structures after diffusion through the extracellular space. Recognition of this general principle indicates that any future drug research should take into account the localization and properties (e.g. affinity, efficacy, etc.) of these non-synaptic receptors and transporters.

## Acknowledgements

This review is dedicated to the memory of Sir William Paton, who was my supervisor (E. S. Vizi) at the Oxford University Department of Pharmacology.

We thank all of our colleagues in the Department of Pharmacology, Institute of Experimental Medicine, Hungarian Academy of Sciences, Budapest, Hungary.

We are grateful to Z. Nusser for the critical reading of this paper. This work was supported by grants by the Hungarian Research Funds OTKA NK 72959 and Jedlik II. OM-00131/2007. Arpad Mike is the recipient of Bolyai-fellowship.

## Conflict of interest

The authors state no conflict of interest.

## References

- Abbracchio MP, Burnstock G, Verkhratsky A, Zimmermann H (2009). Purinergic signalling in the nervous system: an overview. *Trends Neurosci* 32: 19–29.



- Abbracchio MP, Ceruti S (2006). Roles of P2 receptors in glial cells: focus on astrocytes. *Purinergic Signal* 2: 595–604.
- Abercrombie ED, Keller RW, Jr, Zigmond MJ (1988). Characterization of hippocampal norepinephrine release as measured by microdialysis perfusion: pharmacological and behavioral studies. *Neuroscience* 27: 897–904.
- Agnati LF, Fuxe K, Zoli M, Ozini I, Toffano G, Ferraguti F (1986). A correlation analysis of the regional distribution of central enkephalin and beta-endorphin immunoreactive terminals and of opiate receptors in adult and old male rats. Evidence for the existence of two main types of communication in the central nervous system: the volume transmission and the wiring transmission. *Acta Physiol Scand* 128: 201–207.
- Agnati LF, Zoli M, Stromberg I, Fuxe K (1995). Intercellular communication in the brain: wiring versus volume transmission. *Neuroscience* 69: 711–726.
- Alcantara AA, Mrzljak L, Jakab RL, Levey AI, Hersch SM, Goldman-Rakic PS (2001). Muscarinic m1 and m2 receptor proteins in local circuit and projection neurons of the primate striatum: anatomical evidence for cholinergic modulation of glutamatergic prefronto-striatal pathways. *J Comp Neurol* 434: 445–460.
- Alvarez FJ, Pearson JC, Harrington D, Dewey D, Torbeck L, Fyffe RE (1998). Distribution of 5-hydroxytryptamine-immunoreactive boutons on alpha-motoneurons in the lumbar spinal cord of adult cats. *J Comp Neurol* 393: 69–83.
- Amara SG, Fontana AC (2002). Excitatory amino acid transporters: keeping up with glutamate. *Neurochem Int* 41: 313–318.
- Amara SG, Kuhar MJ (1993). Neurotransmitter transporters: recent progress. *Annu Rev Neurosci* 16: 73–93.
- Aoki C, Venkatesan C, Go CG, Forman R, Kurose H (1998). Cellular and subcellular sites for noradrenergic action in the monkey dorsolateral prefrontal cortex as revealed by the immunocytochemical localization of noradrenergic receptors and axons. *Cereb Cortex* 8: 269–277.
- Asztely F, Erdemli G, Kullmann DM (1997). Extrasynaptic glutamate spillover in the hippocampus: dependence on temperature and the role of active glutamate uptake. *Neuron* 18: 281–293.
- Axelsson J, Thesleff S (1959). A study of supersensitivity in denervated mammalian skeletal muscle. *J Physiol* 147: 178–193.
- Aznavour N, Mechawar N, Descarries L (2002). Comparative analysis of cholinergic innervation in the dorsal hippocampus of adult mouse and rat: a quantitative immunocytochemical study. *Hippocampus* 12: 206–217.
- Bach-y-Rita P (1993). Neurotransmission in the brain by diffusion through the extracellular fluid: a review. *Neuroreport* 4: 343–350.
- Balazsa T, Biro J, Gullai N, Ledent C, Sperlagh B (2008). CB1-cannabinoid receptors are involved in the modulation of non-synaptic [3H]serotonin release from the rat hippocampus. *Neurochem Int* 52: 95–102.
- Barbour B, Keller BU, Llano I, Marty A (1994). Prolonged presence of glutamate during excitatory synaptic transmission to cerebellar Purkinje cells. *Neuron* 12: 1331–1343.
- Barnes NM, Cheng CH, Costall B, Ge J, Naylor RJ (1992). Differential modulation of extracellular levels of 5-hydroxytryptamine in the rat frontal cortex by (R)- and (S)-zacopride. *Br J Pharmacol* 107: 233–239.
- Baude A, Molnar E, Latawiec D, McIlhinney RA, Somogyi P (1994). Synaptic and nonsynaptic localization of the GluR1 subunit of the AMPA-type excitatory amino acid receptor in the rat cerebellum. *J Neurosci* 14: 2830–2843.
- Baude A, Nusser Z, Molnar E, McIlhinney RA, Somogyi P (1995). High-resolution immunogold localization of AMPA type glutamate receptor subunits at synaptic and non-synaptic sites in rat hippocampus. *Neuroscience* 69: 1031–1055.
- Baude A, Nusser Z, Roberts JD, Mulvihill E, McIlhinney RA, Somogyi P (1993). The metabotropic glutamate receptor (mGluR1 alpha) is concentrated at perisynaptic membrane of neuronal subpopulations as detected by immunogold reaction. *Neuron* 11: 771–787.
- Beart PM, O'Shea RD (2007). Transporters for l-glutamate: an update on their molecular pharmacology and pathological involvement. *Br J Pharmacol* 150: 5–17.
- Bellone C, Luscher C, Mameli M (2008). Mechanisms of synaptic depression triggered by metabotropic glutamate receptors. *Cell Mol Life Sci* 65: 2913–2923.
- Bernard V, Laribi O, Levey AI, Bloch B (1998). Subcellular redistribution of m2 muscarinic acetylcholine receptors in striatal interneurons in vivo after acute cholinergic stimulation. *J Neurosci* 18: 10207–10218.
- Bernard V, Levey AI, Bloch B (1999). Regulation of the subcellular distribution of m4 muscarinic acetylcholine receptors in striatal neurons in vivo by the cholinergic environment: evidence for regulation of cell surface receptors by endogenous and exogenous stimulation. *J Neurosci* 19: 10237–10249.
- Bezard E, Gross CE, Fournier MC, Dovero S, Bloch B, Jaber M (1999). Absence of MPTP-induced neuronal death in mice lacking the dopamine transporter. *Exp Neurol* 155: 268–273.
- Bezzi P, Carmignoto G, Pasti L, Vesce S, Rossi D, Rizzini BL et al. (1998). Prostaglandins stimulate calcium-dependent glutamate release in astrocytes. *Nature* 391: 281–285.
- Bianchi MT, Macdonald RL (2003). Neurosteroids shift partial agonist activation of GABA(A) receptor channels from low- to high-efficacy gating patterns. *J Neurosci* 23: 10934–10943.
- Binshtok AM, Fleidervish IA, Sprengel R, Gutnick MJ (2006). NMDA receptors in layer 4 spiny stellate cells of the mouse barrel cortex contain the NR2C subunit. *J Neurosci* 26: 708–715.
- Blanpied TA, Scott DB, Ehlers MD (2002). Dynamics and regulation of clathrin coats at specialized endocytic zones of dendrites and spines. *Neuron* 36: 435–449.
- Boulaich S, Daszuta A, Geffard M, Bosler O (1994). Synaptic connectivity of serotonin graft efferents in the suprachiasmatic and supraoptic nuclei of the hypothalamus. *Exp Brain Res* 101: 353–364.
- Bradley SR, Levey AI, Hersch SM, Conn PJ (1996). Immunocytochemical localization of group III metabotropic glutamate receptors in the hippocampus with subtype-specific antibodies. *J Neurosci* 16: 56.
- Bringmann A, Pannicke T, Biedermann B, Francke M, Iandiev I, Grosche J et al. (2009). Role of retinal glial cells in neurotransmitter uptake and metabolism. *Neurochem Int* 54: 143–160.
- Brown N, Kerby J, Bonnert TP, Whiting PJ, Wafford KA (2002). Pharmacological characterization of a novel cell line expressing human alpha(4)beta(3)delta GABA(A) receptors. *Br J Pharmacol* 136: 965–974.
- Bunin MA, Wightman RM (1998). Quantitative evaluation of 5-hydroxytryptamine (serotonin) neuronal release and uptake: an investigation of extrasynaptic transmission. *J Neurosci* 18: 4854–4860.
- Burnstock G (2008a). Non-synaptic transmission at autonomic neuroeffector junctions. *Neurochem Int* 52: 14–25.
- Burnstock G (2008b). Purinergic signalling and disorders of the central nervous system. *Nat Rev Drug Discov* 7: 575–590.
- Buzsaki G (2002). Theta oscillations in the hippocampus. *Neuron* 33: 325–340.
- Calas A, Alonso G, Arnauld E, Vincent JD (1974). Demonstration of indolaminergic fibres in the media eminence of the duck, rat and monkey. *Nature* 250: 241–243.
- Carmignoto G, Fellin T (2006). Glutamate release from astrocytes as a non-synaptic mechanism for neuronal synchronization in the hippocampus. *J Physiol Paris* 99: 98–102.
- Cartmell J, Schoepp DD (2000). Regulation of neurotransmitter release by metabotropic glutamate receptors. *J Neurochem* 75: 889–907.
- Caruncho HJ, Guidotti A, Lindstrom J, Costa E, Pesold C (1997). Subcellular localization of the alpha 7 nicotinic receptor in rat cerebellar granule cell layer. *Neuroreport* 8: 1431–1433.
- Carvelli L, McDonald PW, Blakely RD, Defelice LJ (2004). Dopamine

- transporters depolarize neurons by a channel mechanism. *Proc Natl Acad Sci USA* **101**: 16046–16051.
- Caulfield MP, Birdsall NJ (1998). International Union of Pharmacology. XVII. Classification of muscarinic acetylcholine receptors. *Pharmacol Rev* **50**: 279–290.
- Cavelier P, Hamann M, Rossi D, Mobbs P, Attwell D (2005). Tonic excitation and inhibition of neurons: ambient transmitter sources and computational consequences. *Prog Biophys Mol Biol* **87**: 3–16.
- Chadderton P, Margrie TW, Hausser M (2004). Integration of quanta in cerebellar granule cells during sensory processing. *Nature* **428**: 856–860.
- Chen KC, Budygin EA (2007). Extracting the basal extracellular dopamine concentrations from the evoked responses: re-analysis of the dopamine kinetics. *J Neurosci Methods* **164**: 27–42.
- Chesney MA, Perouansky M, Pearce RA (2003). Differential uptake of volatile agents into brain tissue in vitro. Measurement and application of a diffusion model to determine concentration profiles in brain slices. *Anesthesiology* **99**: 122–130.
- Cho Y, Bannai S (1990). Uptake of glutamate and cysteine in C-6 glioma cells and in cultured astrocytes. *J Neurochem* **55**: 7.
- Choquet D, Triller A (2003). The role of receptor diffusion in the organization of the postsynaptic membrane. *Nat Rev Neurosci* **4**: 251–265.
- Ciruela F, Casado V, Rodrigues RJ, Lujan R, Burgueno J, Canals M et al. (2006). Presynaptic control of striatal glutamatergic neurotransmission by adenosine A1-A2A receptor heteromers. *J Neurosci* **26**: 2080–2087.
- Clements JD, Lester RA, Tong G, Jahr CE, Westbrook GL (1992). The time course of glutamate in the synaptic cleft. *Science* **258**: 1498–1501.
- Conti F, DeBiasi S, Minelli A, Rothstein JD, Melone M (1998). EAAC1, a high-affinity glutamate transporter, is localized to astrocytes and gabaergic neurons besides pyramidal cells in the rat cerebral cortex. *Cereb Cortex* **8**: 108–116.
- Corti C, Aldegheri L, Somogyi P, Ferraguti F (2002). Distribution and synaptic localisation of the metabotropic glutamate receptor 4 (mGluR4) in the rodent CNS. *Neuroscience* **110**: 403–420.
- Coussen F (2009). Molecular determinants of kainate receptor trafficking. *Neuroscience* **158**: 25–35.
- Cragg SJ, Nicholson C, Kume-Kick J, Tao L, Rice ME (2001). Dopamine-mediated volume transmission in midbrain is regulated by distinct extracellular geometry and uptake. *J Neurophysiol* **85**: 1761–1771.
- Cragg S, Rice ME, Greenfield SA (1997). Heterogeneity of electrically evoked dopamine release and reuptake in substantia nigra, ventral tegmental area, and striatum. *J Neurophysiol* **77**: 863–873.
- Crespi F, Martin KF, Marsden CA (1988). Measurement of extracellular basal levels of serotonin in vivo using nafion-coated carbon fibre electrodes combined with differential pulse voltammetry. *Neuroscience* **27**: 885–896.
- Cull-Candy S, Brickley S, Farrant M (2001). NMDA receptor subunits: diversity, development and disease. *Curr Opin Neurobiol* **11**: 327–335.
- Cunha RA (2008). Different cellular sources and different roles of adenosine: A1 receptor-mediated inhibition through astrocytic-driven volume transmission and synapse-restricted A2A receptor-mediated facilitation of plasticity. *Neurochem Int* **52**: 65–72.
- Cunha RA, Johansson B, Constantino MD, Sebastiao AM, Fredholm BB (1996). Evidence for high-affinity binding sites for the adenosine A2A receptor agonist [3H] CGS 21680 in the rat hippocampus and cerebral cortex that are different from striatal A2A receptors. *Naunyn Schmiedeberg Arch Pharmacol* **353**: 261–271.
- Daws LC (2009). Unfaithful neurotransmitter transporters: focus on serotonin uptake and implications for antidepressant efficacy. *Pharmacol Ther* **121**: 89–99.
- Debanne D, Guerineau NC, Gahwiler BH, Thompson SM (1997). Action-potential propagation gated by an axonal I(A)-like K<sup>+</sup> conductance in hippocampus. *Nature* **389**: 286–289.
- Decossas M, Bloch B, Bernard V (2003). Trafficking of the muscarinic m2 autoreceptor in cholinergic basalocortical neurons in vivo: differential regulation of plasma membrane receptor availability and intraneuronal localization in acetylcholinesterase-deficient and -inhibited mice. *J Comp Neurol* **462**: 302–314.
- Decossas M, Doudnikoff E, Bloch B, Bernard V (2005). Aging and subcellular localization of m2 muscarinic autoreceptor in basalocortical neurons in vivo. *Neurobiol Aging* **26**: 1061–1072.
- DeFelice LJ, Goswami T (2007). Transporters as channels. *Annu Rev Physiol* **69**: 87–112.
- DeFelipe J, Jones EG (1988). A light and electron microscopic study of serotonin-immunoreactive fibers and terminals in the monkey sensory-motor cortex. *Exp Brain Res* **71**: 171–182.
- Del Arco A, Segovia G, Fuxe K, Mora F (2003). Changes in dialysate concentrations of glutamate and GABA in the brain: an index of volume transmission mediated actions? *J Neurochem* **85**: 23–33.
- Descarries L, Audet MA, Doucet G, Garcia S, Oleskevich S, Seguela P et al. (1990). Morphology of central serotonin neurons. Brief review of quantified aspects of their distribution and ultrastructural relationships. *Ann N Y Acad Sci* **600**: 81–92.
- Descarries L, Beaudet A, Watkins KC (1975). Serotonin nerve terminals in adult rat neocortex. *Brain Res* **100**: 563–588.
- Descarries L, Berube-Carriere N, Riad M, Bo GD, Mendez JA, Trudeau LE (2008). Glutamate in dopamine neurons: synaptic versus diffuse transmission. *Brain Res Rev* **58**: 290–302.
- Descarries L, Gisiger V, Steriade M (1997). Diffuse transmission by acetylcholine in the CNS. *Prog Neurobiol* **53**: 603–625.
- Descarries L, Mechawar N (2000). Ultrastructural evidence for diffuse transmission by monoamine and acetylcholine neurons of the central nervous system. *Prog Brain Res* **125**: 27–47.
- Descarries L, Saucier G (1972). Disappearance of the locus coeruleus in the rat after intraventricular 6-hydroxydopamine. *Brain Res* **37**: 310–316.
- Descarries L, Watkins KC, Garcia S, Bosler O, Doucet G (1996). Dual character, asynaptic and synaptic, of the dopamine innervation in adult rat neostriatum: a quantitative autoradiographic and immunocytochemical analysis. *J Comp Neurol* **375**: 167–186.
- Descarries L, Watkins KC, Lapierre Y (1977). Noradrenergic axon terminals in the cerebral cortex of rat. III. Topometric ultrastructural analysis. *Brain Res* **133**: 197–222.
- Devoto P, Flore G, Pani L, Gessa GL (2001). Evidence for co-release of noradrenaline and dopamine from noradrenergic neurons in the cerebral cortex. *Mol Psychiatry* **6**: 657–664.
- Diamond JS (2002). A broad view of glutamate spillover. *Nat Neurosci* **5**: 291–292.
- Drew GM, Mitchell VA, Vaughan CW (2008). Glutamate spillover modulates GABAergic synaptic transmission in the rat midbrain periaqueductal grey via metabotropic glutamate receptors and endocannabinoid signaling. *J Neurosci* **28**: 808–815.
- Duan S, Anderson CM, Keung EC, Chen Y, Swanson RA (2003). P2X7 receptor-mediated release of excitatory amino acids from astrocytes. *J Neurosci* **23**: 1320–1328.
- Dunwiddie TV, Masino SA (2001). The role and regulation of adenosine in the central nervous system. *Annu Rev Neurosci* **24**: 31–55.
- Elenkov IJ, Wilder RL, Chrousos GP, Vizi ES (2000). The sympathetic nerve – an integrative interface between two supersystems: the brain and the immune system. *Pharmacol Rev* **52**: 595–638.
- Elsen FP, Liljelund P, Werner DF, Olsen RW, Homanics GE, Harrison NL (2006). GABA(A)-R alpha1 subunit knockin mutation leads to abnormal EEG and anesthetic-induced seizure-like activity in mice. *Brain Res* **1078**: 60–70.
- Eriksson M, Nilsson A, Samuelsson H, Samuelsson EB, Mo L, Akesson E et al. (2007). On the role of NR3A in human NMDA receptors. *Physiol Behav* **92**: 54–59.
- Erreger K, Chen PE, Wyllie DJ, Traynelis SF (2004). Glutamate receptor gating. *Crit Rev Neurobiol* **16**: 187–224.
- Fabian-Fine R, Skehel P, Errington ML, Davies HA, Sher E, Stewart MG

- et al. (2001). Ultrastructural distribution of the alpha7 nicotinic acetylcholine receptor subunit in rat hippocampus. *J Neurosci* 21: 7993–8003.
- Farrant M, Nusser Z (2005). Variations on an inhibitory theme: phasic and tonic activation of GABA(A) receptors. *Nat Rev Neurosci* 6: 215–229.
- Fekete A, Franklin L, Ikemoto T, Rozsa B, Lendvai B, Sylvester Vizi E et al. (2009). Mechanism of the persistent sodium current activator veratridine-evoked Ca elevation: implication for epilepsy. *J Neurochem* 111: 745–756.
- Fekete A, Vizi ES, Kovacs KJ, Lendvai B, Zelles T (2008). Layer-specific differences in reactive oxygen species levels after oxygen-glucose deprivation in acute hippocampal slices. *Free Radical Biology & Medicine* 44: 1010–1022.
- Fellin T, Pascual O, Gobbo S, Pozzan T, Haydon PG, Carmignoto G (2004). Neuronal synchrony mediated by astrocytic glutamate through activation of extrasynaptic NMDA receptors. *Neuron* 43: 729–743.
- Ferraro L, Tanganelli S, O'Connor WT, Antonelli T, Rambert F, Fuxe K (1996). The vigilance promoting drug modafinil decreases GABA release in the medial preoptic area and in the posterior hypothalamus of the awake rat: possible involvement of the serotonergic 5-HT3 receptor. *Neurosci Lett* 220: 5–8.
- Fields RD, Stevens B (2000). ATP: an extracellular signaling molecule between neurons and glia. *Trends Neurosci* 23: 625–633.
- Fleckenstein AE, Metzger RR, Wilkins DG, Gibb JW, Hanson GR (1997). Rapid and reversible effects of methamphetamine on dopamine transporters. *J Pharmacol Exp Ther* 282: 834–838.
- Franco R, Panayiotidis MI, de la Paz LD (2008). Autocrine signaling involved in cell volume regulation: the role of released transmitters and plasma membrane receptors. *J Cell Physiol* 216: 14–28.
- Freund TF, Bolam JP, Bjorklund A, Stenevi U, Dunnett SB, Powell JF et al. (1985). Efferent synaptic connections of grafted dopaminergic neurons reinnervating the host neostriatum: a tyrosine hydroxylase immunocytochemical study. *J Neurosci* 5: 603–616.
- Freund TF, Buzsaki G (1996). Interneurons of the hippocampus. *Hippocampus* 6: 347–470.
- Freund TF, Gulyas AI (1991). GABAergic interneurons containing calbindin D28K or somatostatin are major targets of GABAergic basal forebrain afferents in the rat neocortex. *J Comp Neurol* 314: 187–199.
- Freund TF, Katona I (2007). Perisomatic inhibition. *Neuron* 56: 33–42.
- Freund TF, Katona I, Piomelli D (2003). Role of endogenous cannabinoids in synaptic signaling. *Physiol Rev* 83: 1017–1066.
- Friden M, Winiwarter S, Jerndal G, Bengtsson O, Wan H, Bredberg U et al. (2009). Structure-brain exposure relationships in rat and human using a novel data set of unbound drug concentrations in brain interstitial and cerebrospinal fluids. *J Med Chem* 52: 6233–6243.
- Frischknecht R, Heine M, Perrais D, Seidenbecher CI, Choquet D, Gundelfinger ED (2009). Brain extracellular matrix affects AMPA receptor lateral mobility and short-term synaptic plasticity. *Nat Neurosci* 12: 897–904.
- Fumagalli F, Gainetdinov RR, Valenzano KJ, Caron MG (1998). Role of dopamine transporter in methamphetamine-induced neurotoxicity: evidence from mice lacking the transporter. *J Neurosci* 18: 4861–4869.
- Fuxe K, Agnati LF (eds) (1991). *Volume Transmission in the Brain*. Raven Press: New York.
- Gainetdinov RR, Fumagalli F, Jones SR, Caron MG (1997). Dopamine transporter is required for in vivo MPTP neurotoxicity: evidence from mice lacking the transporter. *J Neurochem* 69: 1322–1325.
- Gardoni F, Mauceri D, Malinverno M, Polli F, Costa C, Tozzi A et al. (2009). Decreased NR2B subunit synaptic levels cause impaired long-term potentiation but not long-term depression. *J Neurosci* 29: 669–677.
- Garris PA, Ciolkowski EL, Wightman RM (1994). Heterogeneity of evoked dopamine overflow within the striatal and striatoamygdaloid regions. *Neuroscience* 59: 417–427.
- Garris PA, Wightman RM (1994). Different kinetics govern dopaminergic transmission in the amygdala, prefrontal cortex, and striatum: an in vivo voltammetric study. *J Neurosci* 14: 442–450.
- Giltsbach R, Hein L (2008). Presynaptic metabotropic receptors for acetylcholine and adrenaline/noradrenaline. *Handb Exp Pharmacol* 184: 261–288.
- Giustizieri M, Bernardi G, Mercuri NB, Berretta N (2005). Distinct mechanisms of presynaptic inhibition at GABAergic synapses of the rat substantia nigra pars compacta. *J Neurophysiol* 94: 1992–2003.
- Glykys J, Mody I (2007). Activation of GABAA receptors: views from outside the synaptic cleft. *Neuron* 56: 763–770.
- Gordon-Weeks PR (1982). Noradrenergic and non-noradrenergic nerves containing small granular vesicles in Auerbach's plexus of the guinea-pig: evidence against the presence of noradrenergic synapses. *Neuroscience* 7: 2925–2936.
- Gredell JA, Turnquist PA, Maciver MB, Pearce RA (2004). Determination of diffusion and partition coefficients of propofol in rat brain tissue: implications for studies of drug action in vitro. *Br J Anaesth* 93: 810–817.
- Groc L, Bard L, Choquet D (2009). Surface trafficking of N-methyl-D-aspartate receptors: physiological and pathological perspectives. *Neuroscience* 158: 4–18.
- Groc L, Heine M, Cousins SL, Stephenson FA, Lounis B, Cognet L et al. (2006). NMDA receptor surface mobility depends on NR2A-2B subunits. *Proc Natl Acad Sci United States America* 103: 18769–18774.
- Gulyas AI, Miles R, Hajos N, Freund TF (1993). Precision and variability in postsynaptic target selection of inhibitory cells in the hippocampal CA3 region. *Eur J Neurosci* 5: 1729–1751.
- Haas HL, Selbach O (2000). Functions of neuronal adenosine receptors. *Naunyn Schmiedeberg's Arch Pharmacol* 362: 375–381.
- Hagberg H, Andersson P, Lacarewicz J, Jacobson I, Butcher S, Sandberg M (1987). Extracellular adenosine, inosine, hypoxanthine, and xanthine in relation to tissue nucleotides and purines in rat striatum during transient ischemia. *J Neurochem* 49: 227–231.
- Hardingham GE, Bading H (2003). The yin and yang of NMDA receptor signalling. *Trends Neurosci* 26: 81–89.
- Hardingham GE, Fukunaga Y, Bading H (2002). Extrasynaptic NMDARs oppose synaptic NMDARs by triggering CREB shut-off and cell death pathways. *Nat Neurosci* 5: 405–414.
- Harney SC, Jane DE, Anwyl R (2008). Extrasynaptic NR2D-containing NMDARs are recruited to the synapse during LTP of NMDAR-EPSCs. *J Neurosci* 28: 11685–11694.
- Hasko G, Csóka B, Nemeth ZH, Vizi ES, Pacher P (2009). A2B adenosine receptors in immunity and inflammation. *Trends Immunol* 30: 263–270.
- Haydon PG, Carmignoto G (2006). Astrocyte control of synaptic transmission and neurovascular coupling. *Physiol Rev* 86: 1009–1031.
- Heja L, Barabas P, Nyitrai G, Kekesi KA, Lasztocki B, Toke O et al. (2009). Glutamate uptake triggers transporter-mediated GABA release from astrocytes. *Plos One* 4: e7153.
- Herd MB, Haythornthwaite AR, Rosahl TW, Wafford KA, Homanics GE, Lambert JJ et al. (2008). The expression of GABAA beta subunit isoforms in synaptic and extrasynaptic receptor populations of mouse dentate gyrus granule cells. *J Physiol* 586: 989–1004.
- Herkenham M (1987). Mismatches between neurotransmitter and receptor localizations in brain: observations and implications. *Neuroscience* 23: 1–38.
- Herkenham M (1991). Mismatches between neurotransmitter and receptor localizations: implications for endocrine functions in brain. In: Fuxe K, Agnati LF (eds). *Volume Transmission in the Brain Novel Mechanisms for Neural Transmission*, Vol. 1. Raven Press: New York, pp. 63–87.
- Hersch SM, Gutekunst CA, Rees HD, Heilman CJ, Levey AI (1994).



- Distribution of m1-m4 muscarinic receptor proteins in the rat striatum: light and electron microscopic immunocytochemistry using subtype-specific antibodies. *J Neurosci* **14**: 3351–3363.
- Hettinger BD, Lee A, Linden J, Rosin DL (2001). Ultrastructural localization of adenosine A2A receptors suggests multiple cellular sites for modulation of GABAergic neurons in rat striatum. *J Comp Neurol* **431**: 331–346.
- Hill JA, Jr, Zoli M, Bourgeois JP, Changeux JP (1993). Immunocytochemical localization of a neuronal nicotinic receptor: the beta 2-subunit. *J Neurosci* **13**: 1551–1568.
- Hökfelt T (1968). In vitro studies on central and peripheral monoamine neurons at the ultrastructural level. *Z Zellforsch Mikrosk Anat* **91**: 1–74.
- van Hooft JA, Vijverberg HP (2000). 5-HT(3) receptors and neurotransmitter release in the CNS: a nerve ending story? *Trends Neurosci* **23**: 605–610.
- Housley GD, Bringmann A, Reichenbach A (2009). Purinergic signaling in special senses. *Trends Neurosci* **32**: 128–141.
- Huang EP (1998). Synaptic transmission: spillover at central synapses. *Current Biology* **8**: R613–R615.
- Huang J, Pickel VM (2002). Serotonin transporters (SERTs) within the rat nucleus of the solitary tract: subcellular distribution and relation to 5HT2A receptors. *J Neurocytol* **31**: 667–679.
- Huang J, Spier AD, Pickel VM (2004). 5-HT3A receptor subunits in the rat medial nucleus of the solitary tract: subcellular distribution and relation to the serotonin transporter. *Brain Res* **1028**: 156–169.
- Hussl S, Boehm S (2006). Functions of neuronal P2Y receptors. *Pflugers Arch* **452**: 538–551.
- Ikeda M, Nakazawa T, Abe K, Kaneko T, Yamatsu K (1989). Extracellular accumulation of glutamate in the hippocampus induced by ischemia is not calcium dependent – in vitro and in vivo evidence. *Neurosci Lett* **96**: 202–206.
- Ingram SL, Prasad BM, Amara SG (2002). Dopamine transporter-mediated conductances increase excitability of midbrain dopamine neurons. *Nat Neurosci* **5**: 971–978.
- Isaacson JS, Solis JM, Nicoll RA (1993). Local and diffuse synaptic actions of GABA in the hippocampus. *Neuron* **10**: 165–175.
- Jackson VM, Cunnane TC (2001). Neurotransmitter release mechanisms in sympathetic neurons: past, present, and future perspectives. *Neurochem Res* **26**: 875–889.
- Jacob TC, Moss SJ, Jurd R (2008). GABA(A) receptor trafficking and its role in the dynamic modulation of neuronal inhibition. *Nature Reviews* **9**: 331–343.
- Jaffe EH, Marty A, Schulte A, Chow RH (1998). Extrasynaptic vesicular transmitter release from the somata of substantia nigra neurons in rat midbrain slices. *J Neurosci* **18**: 3548–3553.
- Jaskolski F, Coussen F, Mulle C (2005). Subcellular localization and trafficking of kainate receptors. *Trends Pharmacol Sci* **26**: 20–26.
- Jaskolski F, Henley JM (2009). Synaptic receptor trafficking: the lateral point of view. *Neuroscience* **158**: 19–24.
- Johansson B, Fredholm BB (1995). Further characterization of the binding of the adenosine receptor agonist [3H]CGS 21680 to rat brain using autoradiography. *Neuropharmacology* **34**: 393–403.
- Jones IW, Bolam JP, Wonnacott S (2001). Presynaptic localisation of the nicotinic acetylcholine receptor beta2 subunit immunoreactivity in rat nigrostriatal dopaminergic neurones. *J Comp Neurol* **439**: 235–247.
- Jones MV, Westbrook GL (1995). Desensitized states prolong GABA channel responses to brief agonist pulses. *Neuron* **15**: 181–191.
- Jones MV, Westbrook GL (1996). The impact of receptor desensitization on fast synaptic transmission. *Trends Neurosci* **19**: 96–101.
- Jones IW, Wonnacott S (2004). Precise localization of alpha7 nicotinic acetylcholine receptors on glutamatergic axon terminals in the rat ventral tegmental area. *J Neurosci* **24**: 11244–11252.
- Jourdain P, Bergersen LH, Bhaukaurally K, Bezzi P, Santello M, Domercq M et al. (2007). Glutamate exocytosis from astrocytes controls synaptic strength. *Nat Neurosci* **10**: 331–339.
- Karadottir R, Cavalier P, Bergersen LH, Attwell D (2005). NMDA receptors are expressed in oligodendrocytes and activated in ischaemia. *Nature* **438**: 1162–1166.
- Katona I, Freund TF (2008). Endocannabinoid signaling as a synaptic circuit breaker in neurological disease. *Nat Med* **14**: 923–930.
- Katona I, Sperlagh B, Sik A, Kafalvi A, Vizi ES, Mackie K et al. (1999). Presynaptically located CB1 cannabinoid receptors regulate GABA release from axon terminals of specific hippocampal interneurons. *J Neurosci* **19**: 4544–4558.
- Katona I, Urban GM, Wallace M, Ledent C, Jung KM, Piomelli D et al. (2006). Molecular composition of the endocannabinoid system at glutamatergic synapses. *J Neurosci* **26**: 5628–5637.
- Kennedy RT, Thompson JE, Vickroy TW (2002). In vivo monitoring of amino acids by direct sampling of brain extracellular fluid at ultralow flow rates and capillary electrophoresis. *J Neurosci Methods* **114**: 39–49.
- Kew JN, Richards JG, Mutel V, Kemp JA (1998). Developmental changes in NMDA receptor glycine affinity and ifenprodil sensitivity reveal three distinct populations of NMDA receptors in individual rat cortical neurons. *J Neurosci* **18**: 1935–1943.
- Kieval JZ, Hubert GW, Charara A, Pare JF, Smith Y (2001). Subcellular and subsynaptic localization of presynaptic and postsynaptic kainate receptor subunits in the monkey striatum. *J Neurosci* **21**: 8746–8757.
- Kim MJ, Dunah AW, Wang YT, Sheng M (2005). Differential roles of NR2A- and NR2B-containing NMDA receptors in Ras-ERK signaling and AMPA receptor trafficking. *Neuron* **46**: 745–760.
- Kimelberg HK, Goderie SK, Higman S, Pang S, Waniewski RA (1990). Swelling-induced release of glutamate, aspartate, and taurine from astrocyte cultures. *J Neurosci* **10**: 1583–1591.
- Kirsch J, Wolters I, Triller A, Betz H (1993). Gephyrin antisense oligonucleotides prevent glycine receptor clustering in spinal neurons. *Nature* **366**: 745–748.
- Knoll J, Vizi ES (1970). Presynaptic inhibition of acetylcholine release by endogenous and exogenous noradrenaline at high rate of stimulation. *Br J Pharmacol* **40**: 554P–555P.
- Knoll J, Vizi ES (1971). Effect of frequency of stimulation on the inhibition by noradrenaline of the acetylcholine output from parasympathetic nerve terminals. *Br J Pharmacol* **42**: 263–272.
- Kofalvi A, Rodrigues RJ, Ledent C, Mackie K, Vizi ES, Cunha RA et al. (2005). Involvement of cannabinoid receptors in the regulation of neurotransmitter release in the rodent striatum: a combined immunohistochemical and pharmacological analysis. *J Neurosci* **25**: 2874–2884.
- Kofalvi A, Vizi ES, Ledent C, Sperlagh B (2003). Cannabinoids inhibit the release of [3H]glutamate from rodent hippocampal synaptosomes via a novel CB1 receptor-independent action. *Eur J Neurosci* **18**: 1973–1978.
- Kosofsky BE, Molliver ME (1987). The serotonergic innervation of cerebral cortex: different classes of axon terminals arise from dorsal and median raphe nuclei. *Synapse* **1**: 153–168.
- Kress GJ, Mennerick S (2009). Action potential initiation and propagation: upstream influences on neurotransmission. *Neuroscience* **158**: 211–222.
- Krishnan V, Nestler EJ (2008). The molecular neurobiology of depression. *Nature* **455**: 894–902.
- Kulik A, Vida I, Fukazawa Y, Guetg N, Kasugai Y, Marker CL et al. (2006). Compartment-dependent colocalization of Kir3.2-containing K<sup>+</sup> channels and GABAB receptors in hippocampal pyramidal cells. *J Neurosci* **26**: 4289–4297.
- Kulik A, Vida I, Lujan R, Haas CA, Lopez-Bendito G, Shigemoto R et al. (2003). Subcellular localization of metabotropic GABA(B) receptor subunits GABA(B1a/b) and GABA(B2) in the rat hippocampus. *J Neurosci* **23**: 11026–11035.
- Kullmann DM (2000). Spillover and synaptic cross talk mediated by glutamate and GABA in the mammalian brain. *Prog Brain Res* **125**: 339–351.



- Latini S, Bordoni F, Corradetti R, Pepeu G, Pedata F (1999). Effect of A2A adenosine receptor stimulation and antagonism on synaptic depression induced by in vitro ischaemia in rat hippocampal slices. *Br J Pharmacol* **128**: 1035–1044.
- Lau T, Horschitz S, Bartsch D, Schloss P (2009). Monitoring mouse serotonin transporter internalization in stem cell-derived serotonergic neurons by confocal laser scanning microscopy. *Neurochem Int* **54**: 271–276.
- Lau T, Schloss P (2008). The cannabinoid CB1 receptor is expressed on serotonergic and dopaminergic neurons. *Eur J Pharmacol* **578**: 137–141.
- Lau CG, Zukin RS (2007). NMDA receptor trafficking in synaptic plasticity and neuropsychiatric disorders. *Nat Rev* **8**: 413–426.
- Lendvai B, Vizi ES (2008). Nonsynaptic chemical transmission through nicotinic acetylcholine receptors. *Physiol Rev* **88**: 333–349.
- Levina J (2006). Kainate receptor physiology. *Curr Opin Pharmacol* **6**: 89–97.
- Leira J, Herranz AS, Herreras O, Abaira V, Martin del Rio R (1986). In vivo determination of extracellular concentration of amino acids in the rat hippocampus. A method based on brain dialysis and computerized analysis. *Brain Res* **384**: 145–155.
- Levant B (1997). The D3 dopamine receptor: neurobiology and potential clinical relevance. *Pharmacol Rev* **49**: 231–252.
- Liang J, Zhang N, Cagetti E, Houser CR, Olsen RW, Spigelman I (2006). Chronic intermittent ethanol-induced switch of ethanol actions from extrasynaptic to synaptic hippocampal GABA receptors. *J Neurosci* **26**: 1749–1758.
- Liprando LA, Miner LH, Blakely RD, Lewis DA, Sesack SR (2004). Ultrastructural interactions between terminals expressing the norepinephrine transporter and dopamine neurons in the rat and monkey ventral tegmental area. *Synapse* **52**: 233–244.
- Liu L, Wong TP, Pozza MF, Lingenhoehl K, Wang Y, Sheng M et al. (2004a). Role of NMDA receptor subtypes in governing the direction of hippocampal synaptic plasticity. *Science (New York, NY)* **304**: 1021–1024.
- Liu XB, Murray KD, Jones EG (2004b). Switching of NMDA receptor 2A and 2B subunits at thalamic and cortical synapses during early postnatal development. *J Neurosci* **24**: 8885–8895.
- Liu Y, Wong TP, Aarts M, Rooyakkers A, Liu L, Lai TW et al. (2007). NMDA receptor subunits have differential roles in mediating excitotoxic neuronal death both in vitro and in vivo. *J Neurosci* **27**: 2846–2857.
- Llaudet E, Hatz S, Droniou M, Dale N (2005). Microelectrode biosensor for real-time measurement of ATP in biological tissue. *Anal Chem* **77**: 3267–3273.
- Lopez-Bendito G, Shigemoto R, Kulik A, Paulsen O, Fairen A, Lujan R (2002). Expression and distribution of metabotropic GABA receptor subtypes GABABR1 and GABABR2 during rat neocortical development. *Eur J Neurosci* **15**: 1766–1778.
- Lovick TA, Parry DM, Stezhka VV, Lumb BM (2000). Serotonergic transmission in the periaqueductal gray matter in relation to aversive behaviour: morphological evidence for direct modulatory effects on identified output neurons. *Neuroscience* **95**: 763–772.
- Lozovaya NA, Grebenyuk SE, Tsintsadze T, Feng B, Monaghan DT, Krishtal OA (2004). Extrasynaptic NR2B and NR2D subunits of NMDA receptors shape 'superslow' afterburst EPSC in rat hippocampus. *J Physiol* **558**: 451–463.
- Lu W, Shi Y, Jackson AC, Bjorgan K, During MJ, Sprengel R et al. (2009). Subunit composition of synaptic AMPA receptors revealed by a single-cell genetic approach. *Neuron* **62**: 254–268.
- Lujan R, Nusser Z, Roberts JD, Shigemoto R, Somogyi P (1996). Perisynaptic location of metabotropic glutamate receptors mGluR1 and mGluR5 on dendrites and dendritic spines in the rat hippocampus. *Eur J Neurosci* **8**: 1488–1500.
- Lujan R, Roberts JD, Shigemoto R, Ohishi H, Somogyi P (1997). Differential plasma membrane distribution of metabotropic glutamate receptors mGluR1 alpha, mGluR2 and mGluR5, relative to neurotransmitter release sites. *J Chem Neuroanat* **13**: 219–241.
- Lujan R, Shigemoto R (2006). Localization of metabotropic GABA receptor subunits GABAB1 and GABAB2 relative to synaptic sites in the rat developing cerebellum. *Eur J Neurosci* **23**: 1479–1490.
- Lynch JW (2004). Molecular structure and function of the glycine receptor chloride channel. *Physiol Rev* **84**: 1051–1095.
- Malarkey EB, Parpura V (2008). Mechanisms of glutamate release from astrocytes. *Neurochem Int* **52**: 142–154.
- Martin LJ, Blackstone CD, Haganir RL, Price DL (1992). Cellular localization of a metabotropic glutamate receptor in rat brain. *Neuron* **9**: 259–270.
- Massey PV, Johnson BE, Moulton PR, Auberson YP, Brown MW, Molnar E et al. (2004). Differential roles of NR2A and NR2B-containing NMDA receptors in cortical long-term potentiation and long-term depression. *J Neurosci* **24**: 7821–7828.
- Matute C, Alberdi E, Domercq M, Perez-Cerda F, Perez-Samartin A, Sanchez-Gomez MV (2001). The link between excitotoxic oligodendroglial death and demyelinating diseases. *Trends Neurosci* **24**: 224–230.
- Meier J, Vannier C, Serge A, Triller A, Choquet D (2001). Fast and reversible trapping of surface glycine receptors by gephyrin. *Nat Neurosci* **4**: 253–260.
- Melikian HE (2004). Neurotransmitter transporter trafficking: endocytosis, recycling, and regulation. *Pharmacol Ther* **104**: 17–27.
- Meyer G, Kirsch J, Betz H, Langosch D (1995). Identification of a gephyrin binding motif on the glycine receptor beta subunit. *Neuron* **15**: 563–572.
- Milner TA, Lee A, Aicher SA, Rosin DL (1998). Hippocampal alpha2A-adrenergic receptors are located predominantly presynaptically but are also found postsynaptically and in selective astrocytes. *J Comp Neurol* **395**: 310–327.
- Milushva E, Sperlagh B, Kiss B, Szporny L, Pasztor E, Pappasova M et al. (1990). Inhibitory effect of hypoxic condition on acetylcholine release is partly due to the effect of adenosine released from the tissue. *Brain Res Bull* **24**: 369–373.
- Miner LH, Jedema HP, Moore FW, Blakely RD, Grace AA, Sesack SR (2006). Chronic stress increases the plasmalemmal distribution of the norepinephrine transporter and the coexpression of tyrosine hydroxylase in norepinephrine axons in the prefrontal cortex. *J Neurosci* **26**: 1571–1578.
- Miner LH, Schroeter S, Blakely RD, Sesack SR (2003). Ultrastructural localization of the norepinephrine transporter in superficial and deep layers of the rat prelimbic prefrontal cortex and its spatial relationship to probable dopamine terminals. *J Comp Neurol* **466**: 478–494.
- Miquel MC, Emerit MB, Nosjean A, Simon A, Rumajogee P, Brisorgueil MJ et al. (2002). Differential subcellular localization of the 5-HT3-A receptor subunit in the rat central nervous system. *Eur J Neurosci* **15**: 449–457.
- Mody I, De Koninck Y, Otis TS, Soltesz I (1994). Bridging the cleft at GABA synapses in the brain. *Trends Neurosci* **17**: 517–525.
- Mohrmann R, Hatt H, Gottmann K (2000). Developmental regulation of subunit composition of extrasynaptic NMDA receptors in neocortical neurones. *Neuroreport* **11**: 1203–1208.
- Momiyama A (2000). Distinct synaptic and extrasynaptic NMDA receptors identified in dorsal horn neurones of the adult rat spinal cord. *J Physiol* **523** (Pt 3): 621–628.
- Momiyama A, Feldmeyer D, Cull-Candy SG (1996). Identification of a native low-conductance NMDA channel with reduced sensitivity to Mg<sup>2+</sup> in rat central neurones. *J Physiol* **494** (Pt 2): 479–492.
- Moron JA, Brockington A, Wise RA, Rocha BA, Hope BT (2002). Dopamine uptake through the norepinephrine transporter in brain regions with low levels of the dopamine transporter: evidence from knock-out mouse lines. *J Neurosci* **22**: 389–395.
- Mrzljak L, Levey AI, Goldman-Rakic PS (1993). Association of m1 and m2 muscarinic receptor proteins with asymmetric synapses in the

- primate cerebral cortex: morphological evidence for cholinergic modulation of excitatory neurotransmission. *Proc Natl Acad Sci USA* **90**: 5194–5198.
- Nakajima Y, Iwakabe H, Akazawa C, Nawa H, Shigemoto R, Mizuno N *et al.* (1993). Molecular characterization of a novel retinal metabotropic glutamate receptor mGluR6 with a high agonist selectivity for L-2-amino-4-phosphonobutyrate. *J Biol Chem* **268**: 11868–11873.
- Narushima M, Uchigashima M, Fukaya M, Matsui M, Manabe T, Hashimoto K *et al.* (2007). Tonic enhancement of endocannabinoid-mediated retrograde suppression of inhibition by cholinergic interneuron activity in the striatum. *J Neurosci* **27**: 496–506.
- Nathanson NM (2008). Synthesis, trafficking, and localization of muscarinic acetylcholine receptors. *Pharmacol Ther* **119**: 33–43.
- Nedergaard M, Takano T, Hansen AJ (2002). Beyond the role of glutamate as a neurotransmitter. *Nat Rev Neurosci* **3**: 748–755.
- Neve KA, Neve RL (eds) (1997). *The Dopamine Receptors in Molecular Biology of Dopamine Receptors*. Humana Press: Totowa, NJ.
- Nirenberg MJ, Chan J, Liu Y, Edwards RH, Pickel VM (1996). Ultrastructural localization of the vesicular monoamine transporter-2 in midbrain dopaminergic neurons: potential sites for somatodendritic storage and release of dopamine. *J Neurosci* **16**: 4135–4145.
- Nirenberg MJ, Chan J, Vaughan RA, Uhl GR, Kuhar MJ, Pickel VM (1997). Immunogold localization of the dopamine transporter: an ultrastructural study of the rat ventral tegmental area. *J Neurosci* **17**: 5255–5262.
- Nishizaki T, Nagai K, Nomura T, Tada H, Kanno T, Tozaki H *et al.* (2002). A new neuromodulatory pathway with a glial contribution mediated via A(2a) adenosine receptors. *Glia* **39**: 133–147.
- Nomura A, Shigemoto R, Nakamura Y, Okamoto N, Mizuno N, Nakanishi S (1994). Developmentally regulated postsynaptic localization of a metabotropic glutamate receptor in rat rod bipolar cells. *Cell* **77**: 361–369.
- North RA (2002). Molecular physiology of P2X receptors. *Physiol Rev* **82**: 1013–1067.
- North RA, Verkhratsky A (2006). Purinergic transmission in the central nervous system. *Pflugers Arch* **452**: 479–485.
- Nusser Z, Roberts JD, Baude A, Richards JG, Somogyi P (1995). Relative densities of synaptic and extrasynaptic GABA<sub>A</sub> receptors on cerebellar granule cells as determined by a quantitative immunogold method. *J Neurosci* **15**: 2948–2960.
- Nusser Z, Sieghart W, Somogyi P (1998). Segregation of different GABA<sub>A</sub> receptors to synaptic and extrasynaptic membranes of cerebellar granule cells. *J Neurosci* **18**: 1693–1703.
- Nyilas R, Dudok B, Urban GM, Mackie K, Watanabe M, Cravatt BF *et al.* (2008). Enzymatic machinery for endocannabinoid biosynthesis associated with calcium stores in glutamatergic axon terminals. *J Neurosci* **28**: 1058–1063.
- Nyiri G, Cserep C, Szabadits E, Mackie K, Freund TF (2005). CB1 cannabinoid receptors are enriched in the perisynaptic annulus and on preterminal segments of hippocampal GABAergic axons. *Neuroscience* **136**: 811–822.
- Nyitrai G, Kekesi KA, Juhasz G (2006). Extracellular level of GABA and Glu: in vivo microdialysis-HPLC measurements. *Curr Top Med Chem* **6**: 935–940.
- Ohishi H, Neki A, Mizuno N (1998). Distribution of a metabotropic glutamate receptor, mGluR2, in the central nervous system of the rat and mouse: an immunohistochemical study with a monoclonal antibody. *Neurosci Res* **30**: 65–82.
- Ohishi H, Nomura S, Ding YQ, Shigemoto R, Wada E, Kinoshita A *et al.* (1995). Presynaptic localization of a metabotropic glutamate receptor, mGluR7, in the primary afferent neurons: an immunohistochemical study in the rat. *Neurosci Lett* **202**: 85–88.
- Ohishi H, Ogawa-Meguro R, Shigemoto R, Kaneko T, Nakanishi S, Mizuno N (1994). Immunohistochemical localization of metabotropic glutamate receptors, mGluR2 and mGluR3, in rat cerebellar cortex. *Neuron* **13**: 55–66.
- Okamoto Y, Wang J, Morishita J, Ueda N (2007). Biosynthetic pathways of the endocannabinoid anandamide. *Chem Biodivers* **4**: 1842–1857.
- Olah S, Fule M, Komlosi G, Varga C, Baldi R, Barzo P *et al.* (2009). Regulation of cortical microcircuits by unitary GABA-mediated volume transmission. *Nature* **461**: 1278–1281.
- Oleskevich S, Descarries L, Lacaille JC (1989). Quantified distribution of the noradrenaline innervation in the hippocampus of adult rat. *J Neurosci* **9**: 3803–3815.
- Otsuka M, Obata K, Miyata Y, Tanaka Y (1971). Measurement of gamma-aminobutyric acid in isolated nerve cells of cat central nervous system. *J Neurochem* **18**: 287–295.
- Pankratov YV, Krishtal OA (2003). Distinct quantal features of AMPA and NMDA synaptic currents in hippocampal neurons: implication of glutamate spillover and receptor saturation. *Biophys J* **85**: 3375–3387.
- Papp L, Balazsa T, Kofalvi A, Erdelyi F, Szabo G, Vizi ES *et al.* (2004). P2X receptor activation elicits transporter-mediated noradrenaline release from rat hippocampal slices. *J Pharmacol Exp Ther* **310**: 973–980.
- Paton WD, Vizi ES (1969). The inhibitory action of noradrenaline and adrenaline on acetylcholine output by guinea-pig ileum longitudinal muscle strip. *Br J Pharmacol* **35**: 10–28.
- Paton WD, Vizi ES, Zar MA (1971). The mechanism of acetylcholine release from parasympathetic nerves. *J Physiol* **215**: 819–848.
- Perea G, Navarrete M, Araque A (2009). Tripartite synapses: astrocytes process and control synaptic information. *Trends Neurosci* **32**: 421–431.
- Petralia RS, Wang YX, Niedzielski AS, Wenthold RJ (1996). The metabotropic glutamate receptors, mGluR2 and mGluR3, show unique postsynaptic, presynaptic and glial localizations. *Neuroscience* **71**: 949–976.
- Plummer KL, Manning KA, Levey AI, Rees HD, Uhlrich DJ (1999). Muscarinic receptor subtypes in the lateral geniculate nucleus: a light and electron microscopic analysis. *J Comp Neurol* **404**: 408–425.
- Popoli P, Betto P, Reggio R, Ricciarello G (1995). Adenosine A<sub>2A</sub> receptor stimulation enhances striatal extracellular glutamate levels in rats. *Eur J Pharmacol* **287**: 215–217.
- Pothos EN, Davila V, Sulzer D (1998). Presynaptic recording of quanta from midbrain dopamine neurons and modulation of the quantal size. *J Neurosci* **18**: 4106–4118.
- Pugliese AM, Traini C, Cipriani S, Gianfriddo M, Mello T, Giovannini MG *et al.* (2009). The adenosine A(2A) receptor antagonist ZM241385 enhances neuronal survival after oxygen-glucose deprivation in rat CA1 hippocampal slices. *Br J Pharmacol* **157**: 818–830.
- Rahman MA, Kwon NH, Won MS, Choe ES, Shim YB (2005). Functionalized conducting polymer as an enzyme-immobilizing substrate: an amperometric glutamate microbiosensor for in vivo measurements. *Anal Chem* **77**: 4854–4860.
- Riad M, Garcia S, Watkins KC, Jodoin N, Doucet E, Langlois X *et al.* (2000). Somatodendritic localization of 5-HT<sub>1A</sub> and preterminal axonal localization of 5-HT<sub>1B</sub> serotonin receptors in adult rat brain. *J Comp Neurol* **417**: 181–194.
- Ribeiro FM, Ferreira LT, Marion S, Fontes S, Gomez M, Ferguson SS *et al.* (2007). SEC14-like protein 1 interacts with cholinergic transporters. *Neurochem Int* **50**: 356–364.
- Rice ME, Cragg SJ (2008). Dopamine spillover after quantal release: rethinking dopamine transmission in the nigrostriatal pathway. *Brain Res Rev* **58**: 303–313.
- Richerson GB, Wu Y (2003). Dynamic equilibrium of neurotransmitter transporters: not just for reuptake anymore. *J Neurophysiol* **90**: 1363–1374.
- Ridet JL, Rajaofetra N, Teilhac JR, Geffard M, Privat A (1993). Evidence for nonsynaptic serotonergic and noradrenergic innervation of the rat dorsal horn and possible involvement of neuron-glia interactions. *Neuroscience* **52**: 143–157.

- Robinson MB (2006). Acute regulation of sodium-dependent glutamate transporters: a focus on constitutive and regulated trafficking. *Handb Exp Pharmacol* 175: 251–275.
- Rodrigues RJ, Almeida T, Richardson PJ, Oliveira CR, Cunha RA (2005). Dual presynaptic control by ATP of glutamate release via facilitatory P2X1, P2X2/3, and P2X3 and inhibitory P2Y1, P2Y2, and/or P2Y4 receptors in the rat hippocampus. *J Neurosci* 25: 6286–6295.
- Rosin DL, Robeva A, Woodard RL, Guyenet PG, Linden J (1998). Immunohistochemical localization of adenosine A2A receptors in the rat central nervous system. *J Comp Neurol* 401: 163–186.
- Rossi DJ, Brady JD, Mohr C (2007). Astrocyte metabolism and signaling during brain ischemia. *Nature Neuroscience* 10: 1377–1386.
- Rossi P, Sola E, Taglietti V, Borchardt T, Steigerwald F, Utvik JK et al. (2002). NMDA receptor 2 (NR2) C-terminal control of NR open probability regulates synaptic transmission and plasticity at a cerebellar synapse. *J Neurosci* 22: 9687–9697.
- Rouse ST, Edmunds SM, Yi H, Gilmor ML, Levey AI (2000). Localization of M(2) muscarinic acetylcholine receptor protein in cholinergic and non-cholinergic terminals in rat hippocampus. *Neurosci Lett* 284: 182–186.
- Rouse ST, Levey AI (1996). Expression of m1-m4 muscarinic acetylcholine receptor immunoreactivity in septohippocampal neurons and other identified hippocampal afferents. *J Comp Neurol* 375: 406–416.
- Rozsa B, Katona G, Kaszas A, Szipocs R, Vizi ES (2008). Dendritic nicotinic receptors modulate backpropagating action potentials and long-term plasticity of interneurons. *Eur J Neurosci* 27: 364–377.
- Rozsa B, Zelles T, Vizi ES, Lendvai B (2004). Distance-dependent scaling of calcium transients evoked by backpropagating spikes and synaptic activity in dendrites of hippocampal interneurons. *J Neurosci* 24: 661–670.
- Rubio ME, Soto F (2001). Distinct Localization of P2X receptors at excitatory postsynaptic specializations. *J Neurosci* 21: 641–653.
- Rudolphi KA, Schubert P (1997). Modulation of neuronal and glial cell function by adenosine and neuroprotection in vascular dementia. *Behav Brain Res* 83: 123–128.
- Rusakov DA, Kullmann DM (1998). Extrasynaptic glutamate diffusion in the hippocampus: ultrastructural constraints, uptake, and receptor activation. *J Neurosci* 18: 3158–3170.
- Rusakov DA, Kullmann DM, Stewart MG (1999). Hippocampal synapses: do they talk to their neighbours? *Trends Neurosci* 22: 382–388.
- Salter MG, Fern R (2005). NMDA receptors are expressed in developing oligodendrocyte processes and mediate injury. *Nature* 438: 1167–1171.
- Santos SD, Carvalho AL, Caldeira MV, Duarte CB (2009). Regulation of AMPA receptors and synaptic plasticity. *Neuroscience* 158: 105–125.
- Saxena NC, Macdonald RL (1996). Properties of putative cerebellar gamma-aminobutyric acid A receptor isoforms. *Mol Pharmacol* 49: 567–579.
- Scanziani M (2000). GABA spillover activates postsynaptic GABA(B) receptors to control rhythmic hippocampal activity. *Neuron* 25: 673–681.
- Scanziani M, Gahwiler BH, Thompson SM (1995). Presynaptic inhibition of excitatory synaptic transmission by muscarinic and metabotropic glutamate receptor activation in the hippocampus: are Ca<sup>2+</sup> channels involved? *Neuropharmacology* 34: 1549–1557.
- Schlicker E, Timm J, Zentner J, Gothert M (1997). Cannabinoid CB1 receptor-mediated inhibition of noradrenaline release in the human and guinea-pig hippocampus. *Naunyn Schmiedeberg's Arch Pharmacol* 356: 583–589.
- Schmitt FO (1984). Molecular regulators of brain function: a new view. *Neuroscience* 13: 991–1001.
- Schnell E, Sizemore M, Karimzadegan S, Chen L, Bredt DS, Nicoll RA (2002). Direct interactions between PSD-95 and stargazin control synaptic AMPA receptor number. *Proc Natl Acad Sci USA* 99: 13902–13907.
- Schoffelmeer AN, Hogenboom F, Wardeh G, De Vries TJ (2006). Interactions between CB1 cannabinoid and mu opioid receptors mediating inhibition of neurotransmitter release in rat nucleus accumbens core. *Neuropharmacology* 51: 773–781.
- Schroeter S, Apparsundaram S, Wiley RG, Miner LH, Sesack SR, Blakely RD (2000). Immunolocalization of the cocaine- and antidepressant-sensitive l-norepinephrine transporter. *J Comp Neurol* 420: 211–232.
- Segev I (1990). Computer study of presynaptic inhibition controlling the spread of action potentials into axonal terminals. *J Neurophysiol* 63: 987–998.
- Segovia G, Mora F (2001). Involvement of NMDA and AMPA/kainate receptors in the effects of endogenous glutamate on extracellular concentrations of dopamine and GABA in the nucleus accumbens of the awake rat. *Brain Res Bull* 54: 153–157.
- Semyanov A (2008). Can diffuse extrasynaptic signaling form a guiding template? *Neurochem Int* 52: 31–33.
- Semyanov A, Walker MC, Kullmann DM, Silver RA (2004). Tonic active GABA A receptors: modulating gain and maintaining the tone. *Trends Neurosci* 27: 262–269.
- Sesack SR, Aoki C, Pickel VM (1994). Ultrastructural localization of D2 receptor-like immunoreactivity in midbrain dopamine neurons and their striatal targets. *J Neurosci* 14: 88–106.
- Sheldon C, Diarra A, Cheng YM, Church J (2004). Sodium influx pathways during and after anoxia in rat hippocampal neurons. *J Neurosci* 24: 11057–11069.
- Shigemoto R, Kinoshita A, Wada E, Nomura S, Ohishi H, Takada M et al. (1997). Differential presynaptic localization of metabotropic glutamate receptor subtypes in the rat hippocampus. *J Neurosci* 17: 7503–7522.
- Silva DG, Ross G, Osborne LW (1971). Adrenergic innervation of the ileum of the cat. *Am J Physiol* 220: 347–352.
- Smiley JF, Goldman-Rakic PS (1996). Serotonergic axons in monkey prefrontal cerebral cortex synapse predominantly on interneurons as demonstrated by serial section electron microscopy. *J Comp Neurol* 367: 431–443.
- Smiley JF, Levey AI, Ciliax BJ, Goldman-Rakic PS (1994). D1 dopamine receptor immunoreactivity in human and monkey cerebral cortex: predominant and extrasynaptic localization in dendritic spines. *Proc Natl Acad Sci USA* 91: 5720–5724.
- Smiley JF, Morrell F, Mesulam MM (1997). Cholinergic synapses in human cerebral cortex: an ultrastructural study in serial sections. *Exp Neurol* 144: 361–368.
- Soltész I, Roberts JD, Takagi H, Richards JG, Mohler H, Somogyi P (1990). Synaptic and nonsynaptic localization of benzodiazepine/GABAA receptor/Cl<sup>-</sup> channel complex using monoclonal antibodies in the dorsal lateral geniculate nucleus of the cat. *Eur J Neurosci* 2: 414–429.
- Somogyi P, Takagi H, Richards JG, Mohler H (1989). Subcellular localization of benzodiazepine/GABAA receptors in the cerebellum of rat, cat, and monkey using monoclonal antibodies. *J Neurosci* 9: 2197–2209.
- Sotnikova TD, Beaulieu JM, Gainetdinov RR, Caron MG (2006). Molecular biology, pharmacology and functional role of the plasma membrane dopamine transporter. *CNS Neurol Disord Drug Targets* 5: 45–56.
- Sperlagh B, Heinrich A, Csolle C (2007a). P2 receptor-mediated modulation of neurotransmitter release—an update. *Purinergic Signal* 3: 269–284.
- Sperlagh B, Kofalvi A, Deuchars J, Atkinson L, Milligan CJ, Buckley NJ et al. (2002). Involvement of P2X7 receptors in the regulation of neurotransmitter release in the rat hippocampus. *J Neurochem* 81: 1196–1211.
- Sperlagh B, Windisch K, Ando RD, Vizi ES (2009). Neurochemical evidence that stimulation of CB1 cannabinoid receptors on GABAergic nerve terminals activates the dopaminergic reward system by increasing dopamine release in the rat nucleus accumbens. *Neurochem Int* 54: 452–457.



- Sperlagh B, Zsilla G, Baranyi M, Illes P, Vizi ES (2007b). Purinergic modulation of glutamate release under ischemic-like conditions in the hippocampus. *Neuroscience* **149**: 99–111.
- Stocca G, Vicini S (1998). Increased contribution of NR2A subunit to synaptic NMDA receptors in developing rat cortical neurons. *J Physiol* **507** (Pt 1): 13–24.
- Summerfield SG, Read K, Begley DJ, Obradovic T, Hidalgo IJ, Coggon S et al. (2007). Central nervous system drug disposition: the relationship between in situ brain permeability and brain free fraction. *J Pharmacol Exp Ther* **322**: 205–213.
- Sykova E (2004). Extrasynaptic volume transmission and diffusion parameters of the extracellular space. *Neuroscience* **129**: 861–876.
- Sykova E, Nicholson C (2008). Diffusion in brain extracellular space. *Physiol Rev* **88**: 1277–1340.
- Sykova E, Vargova L (2008). Extrasynaptic transmission and the diffusion parameters of the extracellular space. *Neurochem Int* **52**: 5–13.
- Szabadics J, Tamas G, Soltesz I (2007). Different transmitter transients underlie presynaptic cell type specificity of GABA<sub>A</sub>, slow and GABA<sub>A</sub>, fast. *Proc Natl Acad Sci USA* **104**: 14831–14836.
- Szabo SI, Zelles T, Lendvai B (2008). Intracellular Ca<sup>2+</sup> dynamics of hippocampal interneurons following nicotinic acetylcholine receptor activation. *Neurochem Int* **52**: 135–141.
- Takahashi T, Forsythe ID, Tsujimoto T, Barnes-Davies M, Onodera K (1996). Presynaptic calcium current modulation by a metabotropic glutamate receptor. *Science* **274**: 594–597.
- Tamaru Y, Nomura S, Mizuno N, Shigemoto R (2001). Distribution of metabotropic glutamate receptor mGluR3 in the mouse CNS: differential location relative to pre- and postsynaptic sites. *Neuroscience* **106**: 481–503.
- Tamas G, Lorincz A, Simon A, Szabadics J (2003). Identified sources and targets of slow inhibition in the neocortex. *Science* **299**: 1902–1905.
- Tao-Cheng JH, Zhou FC (1999). Differential polarization of serotonin transporters in axons versus soma-dendrites: an immunogold electron microscopy study. *Neuroscience* **94**: 821–830.
- Thomas CG, Miller AJ, Westbrook GL (2006). Synaptic and extrasynaptic NMDA receptor NR2 subunits in cultured hippocampal neurons. *J Neurophysiol* **95**: 1727–1734.
- Thomas P, Mortensen M, Hosie AM, Smart TG (2005). Dynamic mobility of functional GABA<sub>A</sub> receptors at inhibitory synapses. *Nat Neurosci* **8**: 889–897.
- Torrealba F, Carrasco MA (2004). A review on electron microscopy and neurotransmitter systems. *Brain Res Brain Res Rev* **47**: 5–17.
- Torres GE, Amara SG (2007). Glutamate and monoamine transporters: new visions of form and function. *Curr Opin Neurobiol* **17**: 304–312.
- Torres GE, Carneiro A, Seamans K, Fiorentini C, Sweeney A, Yao WD et al. (2003). Oligomerization and trafficking of the human dopamine transporter. Mutational analysis identifies critical domains important for the functional expression of the transporter. *J Biol Chem* **278**: 2731–2739.
- Tovar KR, Westbrook GL (1999). The incorporation of NMDA receptors with a distinct subunit composition at nascent hippocampal synapses in vitro. *J Neurosci* **19**: 4180–4188.
- Tovar KR, Westbrook GL (2002). Mobile NMDA receptors at hippocampal synapses. *Neuron* **34**: 255–264.
- Townsend M, Yoshii A, Mishina M, Constantine-Paton M (2003). Developmental loss of miniature N-methyl-D-aspartate receptor currents in NR2A knockout mice. *Proc Natl Acad Sci U S A* **100**: 1340–1345.
- Triller A, Choquet D (2003). Synaptic structure and diffusion dynamics of synaptic receptors. *Biol Cell* **95**: 465–476.
- Triller A, Choquet D (2005). Surface trafficking of receptors between synaptic and extrasynaptic membranes: and yet they do move! *Trends Neurosci* **28**: 133–139.
- Tso MM, Blatchford KL, Callado LF, McLaughlin DP, Stamford JA (2004). Stereoselective effects of ketamine on dopamine, serotonin and noradrenaline release and uptake in rat brain slices. *Neurochem Int* **44**: 1–7.
- Uchigashima M, Narushima M, Fukaya M, Katona I, Kano M, Watanabe M (2007). Subcellular arrangement of molecules for 2-arachidonoyl-glycerol-mediated retrograde signaling and its physiological contribution to synaptic modulation in the striatum. *J Neurosci* **27**: 3663–3676.
- Ullian EM, Sargent PB (1995). Pronounced cellular diversity and extrasynaptic location of nicotinic acetylcholine receptor subunit immunoreactivities in the chicken pretectum. *J Neurosci* **15**: 7012–7023.
- Umbriaco D, Garcia S, Beaulieu C, Descarries L (1995). Relational features of acetylcholine, noradrenaline, serotonin and GABA axon terminals in the stratum radiatum of adult rat hippocampus (CA1). *Hippocampus* **5**: 605–620.
- Umbriaco D, Watkins KC, Descarries L, Cozzari C, Hartman BK (1994). Ultrastructural and morphometric features of the acetylcholine innervation in adult rat parietal cortex: an electron microscopic study in serial sections. *J Comp Neurol* **348**: 351–373.
- Vanhoutte P, Bading H (2003). Opposing roles of synaptic and extrasynaptic NMDA receptors in neuronal calcium signalling and BDNF gene regulation. *Curr Opin Neurobiol* **13**: 366–371.
- Vardi N, Duvoisin R, Wu G, Sterling P (2000). Localization of mGluR6 to dendrites of ON bipolar cells in primate retina. *J Comp Neurol* **423**: 402–412.
- Varga V, Losonczy A, Zemelman BV, Borhegyi Z, Nyiri G, Domonkos A et al. (2009). Fast synaptic subcortical control of hippocampal circuits. *Science* **326**: 449–453.
- Verdier D, Lund JP, Kolta A (2003). GABAergic control of action potential propagation along axonal branches of mammalian sensory neurons. *J Neurosci* **23**: 2002–2007.
- Veruki ML, Morkve SH, Hartveit E (2006). Activation of a presynaptic glutamate transporter regulates synaptic transmission through electrical signaling. *Nat Neurosci* **9**: 1388–1396.
- Vigot R, Barbieri S, Brauner-Osborne H, Turecek R, Shigemoto R, Zhang YP et al. (2006). Differential compartmentalization and distinct functions of GABA<sub>B</sub> receptor variants. *Neuron* **50**: 589–601.
- Vizi ES (1968). The inhibitory action of noradrenaline and adrenaline on release of acetylcholine from guinea-pig ileum longitudinal strips. *Naunyn Schmiedebergs Arch Exp Pathol Pharmacol* **259**: 199–200.
- Vizi ES (1972). Stimulation, by inhibition of (Na<sup>+</sup>-K<sup>+</sup>-Mg<sup>2+</sup>)-activated ATP-ase, of acetylcholine release in cortical slices from rat brain. *J Physiol* **226**: 95–117.
- Vizi ES (1974). Interaction between adrenergic and cholinergic systems: presynaptic inhibitory effect of noradrenaline on acetylcholine release. *J Neural Transm* **11**: 61–78.
- Vizi ES (1979). Presynaptic modulation of neurochemical transmission. *Prog Neurobiol* **12**: 181–290.
- Vizi ES (1980a). Modulation of cortical release of acetylcholine by noradrenaline released from nerves arising from the rat locus coeruleus. *Neuroscience* **5**: 2139–2144.
- Vizi ES (1980b). Non-synaptic modulation of transmitter release: pharmacological implication. *Trends Pharmacol Sci* **1**: 172–175.
- Vizi ES (1984). *Non-Synaptic Interactions between Neurons: Modulation of Neurochemical Transmission. Pharmacological and Clinical Aspects.* John Wiley and Sons, Chichester: New York.
- Vizi ES (2000). Role of high-affinity receptors and membrane transporters in nonsynaptic communication and drug action in the central nervous system. *Pharmacol Rev* **52**: 63–89.
- Vizi ES, Bernath S, Kapocsi J, Serfozo P (1986). Transmitter release from the cytoplasm is of physiological importance but not subject to presynaptic modulation. *J Physiol (Paris)* **81**: 283–288.
- Vizi ES, Kiss JP (1998). Neurochemistry and pharmacology of the major hippocampal transmitter systems: synaptic and nonsynaptic interactions. *Hippocampus* **8**: 566–607.
- Vizi ES, Kiss JP, Lendvai B (2004a). Nonsynaptic communication in the central nervous system. *Neurochem Int* **45**: 443–451.
- Vizi ES, Knoll J (1971). The effects of sympathetic nerve stimulation and guanethidine on parasympathetic neuroeffector transmission;



- the inhibition of acetylcholine release. *J Pharm Pharmacol* 23: 918–925.
- Vizi ES, Knoll J (1976). The inhibitory effect of adenosine and related nucleotides on the release of acetylcholine. *Neuroscience* 1: 391–398.
- Vizi ES, Labos E (1991). Non-synaptic interactions at presynaptic level. *Prog Neurobiol* 37: 145–163.
- Vizi ES, Mike A (2006). Nonsynaptic receptors for GABA and glutamate. *Curr Top Med Chem* 6: 941–948.
- Vizi ES, Mike A, Tarnawa I (1996). 2,3-Benzodiazepines (GYKI 52466 and analogs): negative allosteric modulators of AMPA receptors. *CNS Drug Rev* 2: 91–126.
- Vizi ES, Orso E, Osipenko ON, Hasko G, Elenkov IJ (1995). Neurochemical, electrophysiological and immunocytochemical evidence for a noradrenergic link between the sympathetic nervous system and thymocytes. *Neuroscience* 68: 1263–1276.
- Vizi SE, Ronai A, Harsing L, Jr, Knoll J (1977). Inhibitory effect of dopamine on acetylcholine release from caudate nucleus. *Pol J Pharmacol Pharm* 29: 201–211.
- Vizi ES, Sperlagh B, Baranyi M (1992a). Evidence that ATP released from the postsynaptic site by noradrenaline, is involved in mechanical responses of guinea-pig vas deferens: cascade transmission. *Neuroscience* 50: 455–465.
- Vizi ES, Toth IE, Orso E, Szalay KS, Szabo D, Baranyi M et al. (1993). Dopamine is taken up from the circulation by, and released from, local noradrenergic varicose axon terminals in zona glomerulosa of the rat: a neurochemical and immunocytochemical study. *J Endocrinol* 139: 213–226.
- Vizi ES, Toth IE, Szalay KS, Windisch K, Orso E, Szabo D et al. (1992b). Catecholamines released from local adrenergic axon terminals are possibly involved in fine tuning of steroid secretion from zona glomerulosa cells: functional and morphological evidence. *J Endocrinol* 135: 551–561.
- Vizi ES, Zsilla G, Caron MG, Kiss JP (2004b). Uptake and release of norepinephrine by serotonergic terminals in norepinephrine transporter knock-out mice: implications for the action of selective serotonin reuptake inhibitors. *J Neurosci* 24: 7888–7894.
- Wenzel A, Fritschy JM, Mohler H, Benke D (1997). NMDA receptor heterogeneity during postnatal development of the rat brain: differential expression of the NR2A, NR2B, and NR2C subunit proteins. *J Neurochem* 68: 469–478.
- Wilson CJ, Groves PM, Fifkova E (1977). Monoaminergic synapses, including dendro-dendritic synapses in the rat substantia nigra. *Exp Brain Res* 30: 161–174.
- Wilson RI, Nicoll RA (2002). Endocannabinoid signaling in the brain. *Science* 296: 678–682.
- Wohlfarth KM, Bianchi MT, Macdonald RL (2002). Enhanced neurosteroid potentiation of ternary GABA(A) receptors containing the delta subunit. *J Neurosci* 22: 1541–1549.
- Wonnacott S (1997). Presynaptic nicotinic ACh receptors. *Trends Neurosci* 20: 92–98.
- Wotjak CT, Landgraf R, Engelmann M (2008). Listening to neuropeptides by microdialysis: echoes and new sounds? *Pharmacol Biochem Behav* 90: 125–134.
- Wu H, Jin Y, Buddhala C, Osterhaus G, Cohen E, Jin H et al. (2007). Role of glutamate decarboxylase (GAD) isoform, GAD65, in GABA synthesis and transport into synaptic vesicles—Evidence from GAD65-knockout mice studies. *Brain Res* 1154: 80–83.
- Wu LG, Saggau P (1997). Presynaptic inhibition of elicited neurotransmitter release. *Trends Neurosci* 20: 204–212.
- Wu Y, Wang W, Richerson GB (2001). GABA transaminase inhibition induces spontaneous and enhances depolarization-evoked GABA efflux via reversal of the GABA transporter. *J Neurosci* 21: 2630–2639.
- Yamada M, Basile AS, Fedorova I, Zhang W, Duttaroy A, Cui Y et al. (2003). Novel insights into M5 muscarinic acetylcholine receptor function by the use of gene targeting technology. *Life Sci* 74: 345–353.
- Yamakura T, Shimoji K (1999). Subunit- and site-specific pharmacology of the NMDA receptor channel. *Prog Neurobiol* 59: 279–298.
- Yawo H, Chuhma N (1993). Preferential inhibition of omega-conotoxin-sensitive presynaptic Ca<sup>2+</sup> channels by adenosine autoreceptors. *Nature* 365: 256–258.
- Ye ZC, Wyeth MS, Baltan-Tekkok S, Ransom BR (2003). Functional hemichannels in astrocytes: a novel mechanism of glutamate release. *J Neurosci* 23: 3588–3596.
- Yi JH, Hazell AS (2006). Excitotoxic mechanisms and the role of astrocytic glutamate transporters in traumatic brain injury. *Neurochem Int* 48: 394–403.
- Yudowski GA, Puthenveedu MA, Leonoudakis D, Panicker S, Thorn KS, Beattie EC et al. (2007). Real-time imaging of discrete exocytic events mediating surface delivery of AMPA receptors. *J Neurosci* 27: 11112–11121.
- Yung KK, Bolam JP, Smith AD, Hersch SM, Ciliax BJ, Levey AI (1995). Immunocytochemical localization of D1 and D2 dopamine receptors in the basal ganglia of the rat: light and electron microscopy. *Neuroscience* 65: 709–730.
- Zarei MM, Radcliffe KA, Chen D, Patrick JW, Dani JA (1999). Distributions of nicotinic acetylcholine receptor alpha7 and beta2 subunits on cultured hippocampal neurons. *Neuroscience* 88: 755–764.
- van der Zeyden M, Oldenziel WH, Rea K, Cremers TI, Westerink BH (2008). Microdialysis of GABA and glutamate: analysis, interpretation and comparison with microsensors. *Pharmacol Biochem Behav* 90: 135–147.
- Zhang ZW, Coggan JS, Berg DK (1996). Synaptic currents generated by neuronal acetylcholine receptors sensitive to alpha-bungarotoxin. *Neuron* 17: 1231–1240.
- Zhou M, Baudry M (2006). Developmental changes in NMDA neurotoxicity reflect developmental changes in subunit composition of NMDA receptors. *J Neurosci* 26: 2956–2963.
- Zhou FC, Tao-Cheng JH, Segu L, Patel T, Wang Y (1998). Serotonin transporters are located on the axons beyond the synaptic junctions: anatomical and functional evidence. *Brain Res* 805: 241–254.
- Zhu CB, Hewlett WA, Feoktistov I, Biaggioni I, Blakely RD (2004). Adenosine receptor, protein kinase G, and p38 mitogen-activated protein kinase-dependent up-regulation of serotonin transporters involves both transporter trafficking and activation. *Mol Pharmacol* 65: 1462–1474.
- Zimmermann H (2008). ATP and acetylcholine, equal brethren. *Neurochem Int* 52: 634–648.
- Zoli M, Torri C, Ferrari R, Jansson A, Zini I, Fuxe K et al. (1998). The emergence of the volume transmission concept. *Brain Res Brain Res Rev* 26: 136–147.
- van Zwieten PA, Timmermans PB (1983). Cardiovascular alpha 2-receptors. *J Mol Cell Cardiol* 15: 717–733.

## Supporting information

Additional Supporting Information may be found in the online version of this article:

- Appendix S1** Description of simulations.
- Slide S1** 3D and 2D modeling of glutamate release.
- Slide S2** Dopamine and glutamate.
- Slide S3** The effect of transporters.
- Slide S4** Sensitivity ranges of low- and high-affinity receptors.
- Slide S5** The effect of non-instantaneous release.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.