



ANNALES DE L'INSTITUT FOURIER

Thierry DAUDÉ, NIKY KAMRAN & FRANÇOIS NICOLEAU

**Non-uniqueness results for the anisotropic Calderón problem
with data measured on disjoint sets**

Tome 69, n° 1 (2019), p. 119-170.

http://aif.centre-mersenne.org/item/AIF_2019__69_1_119_0

© Association des Annales de l'institut Fourier, 2019,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.

<http://creativecommons.org/licenses/by-nd/3.0/fr/>



NON-UNIQUENESS RESULTS FOR THE ANISOTROPIC CALDERÓN PROBLEM WITH DATA MEASURED ON DISJOINT SETS

by Thierry DAUDÉ,
Niky KAMRAN & François NICOLEAU (*)

ABSTRACT. — In this paper, we show that there is non-uniqueness in the Calderón problem on Riemannian manifolds when the Dirichlet and Neumann data are measured on disjoint sets of the boundary. We provide counterexamples in the case of two and three dimensional Riemannian manifolds with boundary having the topology of circular cylinders in dimension two and toric cylinders in dimension three. The construction could be easily extended to higher dimensional Riemannian manifolds.

RÉSUMÉ. — Dans cet article, on montre qu'il y a non-unicité pour le problème de Calderón sur des variétés riemanniennes quand les données de Dirichlet et de Neumann sont mesurées sur des sous-ensembles disjoints du bord. On construit des contre-exemples à l'unicité en dimension 2 et 3 pour des variétés riemanniennes à bord topologiquement équivalentes à des cylindres dont les fibres sont des tores. La construction pourrait être aisément étendue à des variétés riemanniennes de dimensions supérieures.

1. Introduction

This paper is devoted to the study of various anisotropic Calderón problems on some simple Riemannian manifolds to be described below. Let us first recall some basic facts about the anisotropic Calderón problem in this

Keywords: Anisotropic Calderón problem, Helmholtz equation on a Riemannian manifold, Sturm–Liouville problems, Weyl–Titchmarsh functions.

2010 *Mathematics Subject Classification:* 81U40, 35P25, 58J50.

(*) The research of the first author is supported by the French National Research Projects AARG, No. ANR-12-BS01-012-01, and Iproblems, No. ANR-13-JS01-0006. The research of the second author is supported by NSERC grant RGPIN 105490-2011. The research of the third author is supported by the French National Research Project NOSEVOL, No. ANR- 2011 BS0101901.

setting. We refer for instance to [10, 11, 15, 16, 25, 31, 32, 33] for important contributions to the subject and to the surveys [17, 26, 42, 45] for the current state of the art.

Let (M, g) be an n dimensional smooth compact Riemannian manifold with smooth boundary ∂M . Let us denote by Δ_{LB} the positive Laplace–Beltrami operator on (M, g) . In a local coordinate system $(x^i)_{i=1, \dots, n}$, the Laplace–Beltrami operator Δ_{LB} has the expression

$$\Delta_{LB} = -\Delta_g = -\frac{1}{\sqrt{|g|}} \partial_i \left(\sqrt{|g|} g^{ij} \partial_j \right),$$

where (g^{ij}) is the matrix inverse of the metric tensor (g_{ij}) , where $|g| = \det(g_{ij})$ is the determinant of g and where we use the Einstein summation convention. We recall that the Laplace–Beltrami operator $-\Delta_g$ with Dirichlet boundary conditions is selfadjoint on $L^2(M, d\text{Vol}_g)$ and has pure point spectrum $\{\lambda_j^2\}_{j \geq 1}$ so that $0 < \lambda_1^2 \leq \dots \leq \lambda_j^2 \rightarrow +\infty$ (see for instance [23]).

We consider the Dirichlet problem at a frequency $\lambda^2 \in \mathbb{R}$ on (M, g) such that $\lambda^2 \notin \{\lambda_j^2\}_{j \geq 1}$. We are interested in the solutions u of

$$(1.1) \quad \begin{cases} -\Delta_g u = \lambda^2 u, & \text{on } M, \\ u = \psi, & \text{on } \partial M. \end{cases}$$

It is well known (see for instance [42]) that for any $\psi \in H^{1/2}(\partial M)$, there exists a unique weak solution $u \in H^1(M)$ of (1.1). This allows us to define the Dirichlet-to-Neumann (DN) map as the operator $\Lambda_g(\lambda^2)$ from $H^{1/2}(\partial M)$ to $H^{-1/2}(\partial M)$ defined for all $\psi \in \mathcal{H}^{1/2}(\partial M)$ by

$$(1.2) \quad \Lambda_g(\lambda^2)(\psi) = (\partial_\nu u)|_{\partial M},$$

where u is the unique solution of (1.1) and $(\partial_\nu u)|_{\partial M}$ is its normal derivative with respect to the unit outer normal vector ν on ∂M . Here $(\partial_\nu u)|_{\partial M}$ is interpreted in the weak sense as an element of $H^{-1/2}(\partial M)$ by

$$\langle \Lambda_g(\lambda^2)\psi | \phi \rangle = \int_M \langle du, dv \rangle_g d\text{Vol}_g,$$

for any $\psi \in H^{1/2}(\partial M)$ and $\phi \in H^{1/2}(\partial M)$ such that u is the unique solution of (1.1) and v is any element of $H^1(M)$ so that $v|_{\partial M} = \phi$. If ψ is sufficiently smooth, we can check that

$$\Lambda_g(\lambda^2)\psi = g(\nu, \nabla u)|_{\partial M} = du(\nu)|_{\partial M} = \nu(u)|_{\partial M},$$

where ν represents the unit outer normal vector to ∂M . Clearly, in that case, an expression in local coordinates for the normal derivative is thus

$$(1.3) \quad \partial_\nu u = \nu^i \partial_i u.$$

We shall be interested in fact in the *partial* DN maps defined as follows. Let Γ_D and Γ_N be two open subsets of ∂M . We then define the partial DN map $\Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2)$ as the restriction of the global DN map $\Lambda_g(\lambda^2)$ to Dirichlet data given on Γ_D and Neumann data measured on Γ_N . Precisely, consider the Dirichlet problem

$$(1.4) \quad \begin{cases} -\Delta_g u = \lambda^2 u, & \text{on } M, \\ u = \psi, & \text{on } \Gamma_D, \\ u = 0, & \text{on } \partial M \setminus \Gamma_D. \end{cases}$$

We thus define $\Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2)$ as the operator acting on the functions $\psi \in H^{1/2}(\partial M)$ with $\text{supp } \psi \subset \Gamma_D$ by

$$(1.5) \quad \Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2)(\psi) = (\partial_\nu u)|_{\Gamma_N},$$

where u is the unique solution of (1.4).

The anisotropic partial Calderón problem can be initially stated as: *does the knowledge of the partial DN map $\Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2)$ at a frequency λ^2 determine uniquely the metric g ?* One can think of three subcases of the above problem:

Full data: $\Gamma_D = \Gamma_N = \partial M$. In that case, we simply denote by $\Lambda_g(\lambda^2)$ the DN map.

Local data: $\Gamma_D = \Gamma_N = \Gamma$, where Γ can be any nonempty open subset of ∂M . In that case, we denote by $\Lambda_{g,\Gamma}(\lambda^2)$ the DN map.

Data on disjoint sets: Γ_D and Γ_N are disjoint open sets of ∂M .

Due to a number of gauge invariances, the answer to the above questions is no. Indeed, it is clear from the definitions (1.4) and (1.5) that in any dimension, the partial DN map $\Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2)$ is invariant under pull-back of the metric by the diffeomorphisms of M that are the identity on $\Gamma_D \cup \Gamma_N$, i.e.

$$(1.6) \quad \forall \phi \in \text{Diff}(M), \phi|_{\Gamma_D \cup \Gamma_N} = \text{Id}, \quad \Lambda_{\phi^*g,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2).$$

In the two dimensional case and for zero frequency $\lambda^2 = 0$, there is another gauge invariance of the DN map due to the conformal invariance of the Laplace–Beltrami operator. More precisely, recall that in dimension 2

$$\Delta_{cg} = \frac{1}{c} \Delta_g,$$

for any smooth function $c > 0$. Therefore, we have in dimension 2

$$(1.7) \quad \forall c \in C^\infty(M), c > 0 \text{ and } c|_{\Gamma_N} = 1, \quad \Lambda_{cg,\Gamma_D,\Gamma_N}(0) = \Lambda_{g,\Gamma_D,\Gamma_N}(0),$$

since the unit outer normal vectors ν_{cg} and ν_g coincide on Γ_N in that case.

Hence the appropriate question (called the *anisotropic Calderón conjecture*) to address is the following.

(Q1). — *Let M be a smooth compact manifold with smooth boundary ∂M and let g_1, g_2 be smooth Riemannian metrics on M . Let Γ_D, Γ_N be any open sets of ∂M and λ^2 be a fixed frequency that does not belong to $\sigma(-\Delta_g)$. If*

$$\Lambda_{g_1, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{g_2, \Gamma_D, \Gamma_N}(\lambda^2),$$

then is it true that

$$g_1 = g_2,$$

up to the invariance (1.6) if $\dim M \geq 3$ and up to the invariances (1.6)–(1.7) if $\dim M = 2$ and $\lambda^2 = 0$?

In $\dim M \geq 3$, we can address another relevant (and simpler!) problem by assuming that the Riemannian manifolds (M, g_1) and (M, g_2) belong to the same conformal class, i.e. there exists a smooth positive function c (called the conformal factor) such that $g_2 = cg_1$. In that case, g_1 is considered as the background known metric and the problem consists in determining the unknown scalar function c from the DN map $\Lambda_{cg_1, \Gamma_D, \Gamma_N}(\lambda^2)$. Precisely, the question becomes:

(Q2). — *Let (M, g) be a smooth compact Riemannian manifold with boundary ∂M and let Γ_D, Γ_N be any open sets of ∂M . Let c be a smooth positive function on M . If*

$$\Lambda_{cg, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2),$$

show that there exists a diffeomorphism $\phi : M \rightarrow M$ with $\phi|_{\Gamma_D \cup \Gamma_N} = \text{Id}$ so that

$$(1.8) \quad \phi^*g = cg.$$

Note that in the case of full data $\Gamma_D = \Gamma_N = \partial M$ or more generally in the case where $\Gamma_D \cup \Gamma_N = \partial M$, it is known that any diffeomorphism $\phi : M \rightarrow M$ which satisfies $\phi|_{\partial M} = \text{Id}$ and $\phi^*g = cg$ must be the identity [35]. Therefore, in either of these particular cases, there is no ambiguity arising from diffeomorphisms and (1.8) should be replaced by the condition

$$(1.9) \quad c = 1, \quad \text{on } M.$$

A last version of the anisotropic Calderón problem which in some sense generalizes (Q2) is the following. Consider the solution of the Schrödinger

equation on (M, g) with potential $V \in L^\infty(M)$

$$(1.10) \quad \begin{cases} (-\Delta_g + V)u = \lambda^2 u, & \text{on } M, \\ u = \psi, & \text{on } \Gamma_D, \\ u = 0, & \text{on } \partial M \setminus \Gamma_D. \end{cases}$$

It is well known (see again for instance [10, 42]) that if λ^2 does not belong to the Dirichlet spectrum of $-\Delta_g + V$, then for any $\psi \in H^{1/2}(\partial M)$, there exists a unique weak solution $u \in H^1(M)$ of (1.10). This allows us to define the partial Dirichlet-to-Neumann map $\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2)$ for all $\psi \in \mathcal{H}^{1/2}(\partial M)$ with $\text{supp } \psi \subset \Gamma_D$ by

$$(1.11) \quad \Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2)(\psi) = (\partial_\nu u)|_{\Gamma_N},$$

where u is the unique solution of (1.10) and $(\partial_\nu u)|_{\Gamma_N}$ is its normal derivative with respect to the unit outer normal vector ν on Γ_N . Once again, we assume that g is a known background metric and the problem consists in determining the unknown potential $V \in L^\infty(M)$ from the DN map $\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2)$. Precisely, the question is:

(Q3). — *Let (M, g) be a smooth compact Riemannian manifold with smooth boundary ∂M and let Γ_D, Γ_N be any open sets of ∂M . Let V_1 and V_2 be potentials in $L^\infty(M)$. If*

$$\Lambda_{g,V_1,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,V_2,\Gamma_D,\Gamma_N}(\lambda^2),$$

is it true that

$$V_1 = V_2?$$

There is a straightforward link between (Q2) and (Q3) in the case of zero frequency $\lambda^2 = 0$ and global data. The main point is the observation that the Laplace–Beltrami operator transforms under conformal scalings of the metric by

$$(1.12) \quad \Delta_{cg}u = c^{-\frac{n+2}{4}} (\Delta_g + q) \left(c^{\frac{n-2}{4}} u \right),$$

where

$$(1.13) \quad q = c^{-\frac{n-2}{4}} \Delta_g c^{\frac{n-2}{4}}.$$

Then we can show that if c is a positive smooth function on M such that

$$c|_{\Gamma_D \cup \Gamma_N} = 1, \quad (\partial_\nu c)|_{\Gamma_N} = 0,$$

then

$$(1.14) \quad \Lambda_{cg,V,\Gamma_D,\Gamma_N}(0) = \Lambda_{g,cV-q,\Gamma_D,\Gamma_N}(0),$$

where q is given by (1.13). The proof of (1.14) is an immediate adaptation of the proof of the Proposition 8.2 in [10]. In particular, we have (using the preceding notations)

$$(1.15) \quad \Lambda_{cg, \Gamma_D, \Gamma_N}(0) = \Lambda_{g, -q, \Gamma_D, \Gamma_N}(0),$$

where q is given by (1.13).

Let us show that (Q3) implies (Q2) in the case of zero frequency and global data (i.e. when $\Gamma_D = \Gamma_N = \partial M$). Assume that (Q3) is true and assume that for two metrics g and cg , we have

$$(1.16) \quad \Lambda_{cg}(0) = \Lambda_g(0).$$

Then by boundary determination ([10, 22, 33], we can show that $c|_{\partial M} = 1$ and $(\partial_\nu c)|_{\partial M} = 0$. Hence, we can use (1.15) to show that (1.16) is equivalent to

$$(1.17) \quad \Lambda_{g, -q}(0) = \Lambda_{g, 0}(0),$$

with q given by (1.13) and $\Lambda_{g, -q}(0)$ stands for the global DN map. Finally, our hypothesis that (Q3) holds true now asserts that $q = 0$, or in other words that $\Delta_g c^{\frac{n-2}{4}} = 0$. Since $c|_{\partial M} = 1$, uniqueness of solutions for the Dirichlet problem shows that $c = 1$ on M and (Q2) is proved.

The most complete results on the anisotropic Calderón problems (Q1), (Q2) and (Q3) have been obtained in the case of full data ($\Gamma_D = \Gamma_N = \partial M$) and local data ($\Gamma_D = \Gamma_N = \Gamma$ with Γ any open subset of M) for *vanishing frequency* $\lambda^2 = 0$. In dimension 2, the anisotropic Calderón problem (Q1) for global and local data with $\lambda^2 = 0$ was shown to be true for connected Riemannian surfaces in [32]. We also refer to [1] for similar results answering (Q1) for global and local data in the case of anisotropic conductivities which are only L^∞ on bounded domains of \mathbb{R}^n . A positive answer to (Q1) for global and local data and zero frequency $\lambda^2 = 0$ in dimension 3 or higher has been given for compact connected real analytic Riemannian manifolds with real analytic boundary first in [33] under some topological assumptions relaxed later in [31, 32] and for compact connected Einstein manifolds with boundary in [15]. Note that Einstein manifolds are real analytic in their interior. Let us point out here that no connectedness assumption on the measurement set Γ was made in the works [15, 32].

The general full or local data anisotropic Calderón problem (Q1) in dimension 3 or higher remains a major open problem. A few deep results concerning the partial questions (Q2) and (Q3) have been obtained recently in [10, 11] for some classes of smooth compact Riemannian manifolds with boundary that are *conformally transversally anisotropic*, i.e. Riemannian

manifolds (M, g) such that

$$M \subset \mathbb{R} \times M_0, \quad g = c(e \oplus g_0),$$

where (M_0, g_0) is a $n - 1$ dimensional smooth compact Riemannian manifold with boundary, e is the Euclidean metric on the real line, and c is a smooth positive function in the cylinder $\mathbb{R} \times M_0$. Under some conditions on the transverse manifold (M_0, g_0) such as for instance simplicity⁽¹⁾, the Riemannian manifold (M, g) is said to be *admissible*. In that framework, the authors of [10, 11] were able to determine uniquely the conformal factor c from the knowledge of the DN map at zero frequency $\lambda^2 = 0$, that is, they answered both (Q2) and (Q3) for the class of admissible Riemannian manifolds. Moreover, these results were extended recently to anisotropic Calderón problems with partial data in [25] (see below). We also refer to [16, 19, 21] for other local data results and to the surveys [17, 26] for more references on the subject.

Concerning the anisotropic Calderón problem with data measured on distinct (not necessarily disjoint) sets Γ_D, Γ_N of ∂M , we refer to [27] for some positive results of (Q3) in the case of bounded domains Ω of $\mathbb{R}^n, n \geq 3$ equipped with the Euclidean metric. Roughly speaking, in [27], the sets Γ_D, Γ_N where the measurements are made must overlap a little bit. Precisely, $\Gamma_D \subset \partial\Omega$ can possibly be very small and Γ_N must then be slightly larger than $\partial\Omega \setminus \Gamma_D$. We also refer to [25] for the generalization of [27] to admissible Riemannian manifolds. To explain the result of [25], we recall first that admissible manifolds admit certain functions φ which are called *limiting Carleman weights* (LCW) and which are useful for constructing complex geometrical optic solutions. We refer to [10] for the definition and properties of limiting Carleman weights on manifolds and their applications. Thanks to the existence of LCW φ , we can decompose the boundary of M as

$$\partial M = \partial M_+ \cup \partial M_{\tan} \cup \partial M_-,$$

where

$$\partial M_{\pm} = \{x \in \partial M : \pm \partial_\nu \varphi(x) > 0\}, \quad \partial M_{\tan} = \{x \in \partial M : \pm \partial_\nu \varphi(x) = 0\}.$$

⁽¹⁾We say that a compact manifold (M_0, g_0) is simple if any two points in M_0 are connected by a unique geodesic depending smoothly on the endpoints and if ∂M_0 is strictly convex (its second fundamental form is positive definite).

Roughly speaking, the authors of [25] show that (Q3) is true⁽²⁾ if the set of Dirichlet data Γ_D contains $\partial M_- \cup \Gamma_a$ and the set of Neumann measurements Γ_N contains $\partial M_+ \cup \Gamma_a$ where Γ_a is some open subset of ∂M_{tan} . Hence in particular, the sets Γ_D and Γ_N must overlap in order to have uniqueness. The only exception occurs in the case where ∂M_{tan} has zero measure. In that case, it is enough to take $\Gamma_D = \partial M_-$ and $\Gamma_N = \partial M_+$ to have uniqueness in (Q3) (see [25, Theorem 2.3]). Note in this case that $\Gamma_D \cap \Gamma_N = \partial M_- \cap \partial M_+ = \emptyset$.

We conclude our survey of anisotropic Calderón problems with the case of data measured on *disjoint sets* for which only a few results are known in the case of zero frequency $\lambda^2 = 0$. We already mentioned above the recent paper [25] which concerns a certain subclass of admissible Riemannian manifolds. The only other result we are aware of is due to Imanuvilov, Uhlmann and Yamamoto [20] who, roughly speaking, showed in the 2-dimensional case, that the potential of a Schrödinger equation on a two-dimensional domain homeomorphic to a disc, where the boundary is partitioned into eight clockwise-ordered parts $\Gamma_1, \Gamma_2, \dots, \Gamma_8$ is determined by boundary measurements with sources supported on $S = \Gamma_2 \cup \Gamma_6$ and fields observed on $R = \Gamma_4 \cup \Gamma_8$, hence answering (Q3) in this particular setting.

Let us also mention some related papers by Rakesh [38] and by Oksanen, Lassas [29, 30] dealing with the *hyperbolic* anisotropic Calderón problem, that is to say in our language, the case where we assume the knowledge of the partial DN map at all frequencies λ^2 . We refer to [23] for a thorough account on hyperbolic anisotropic Calderón problem and to [24] for the link between the hyperbolic DN map and the elliptic DN map at all frequencies. For instance, Oksanen and Lassas showed in [30] that (M, g) is uniquely determined (up to the gauge invariance (1.6)) from the knowledge of $\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2)$ at all frequencies λ^2 under the Hassell–Tao type assumption

$$\exists C_0 > 0, \quad \lambda_j^2 \leq C_0 \|\partial_\nu \phi_j\|_{L^2(\Gamma_D)}^2, \quad \forall j \geq 1, \dots$$

where λ_j^2 are the eigenvalues of $-\Delta_g$ with Dirichlet boundary conditions and ϕ_j are the associated normalized eigenfunctions.

Finally, in [38], Rakesh proved that the coefficients of a wave equation on a one-dimensional interval are determined by boundary measurements with sources supported on one end of the interval and the waves observed

⁽²⁾ In fact, additional geometric assumptions on the transverse manifold (M_0, g_0) are needed to give a full proof of this result. We refer to [25, Theorem 2.1] for the precise statement.

on the other end. Here again, the uniqueness result entails to know the hyperbolic DN map or equivalently the DN map at all frequencies.

In this paper, we are mainly concerned with the anisotropic Calderón problems (Q1) and (Q2) with Dirichlet and Neumann data measured on *disjoint sets*. We provide some simple counterexamples to uniqueness in (Q1) and (Q2) for Riemannian surfaces (in the case of nonzero frequency) and for 3 dimensional Riemannian manifolds (without restriction on the frequency). In fact, similar counterexamples to uniqueness can be found for any n dimensional Riemannian manifold, but we only give the details in the 3 dimensional case to keep things concise.

First, we consider a smooth compact Riemannian surface (\mathcal{S}, g) having the topology of a cylinder $M = [0, 1] \times T^1$ and that is equipped with a Riemannian metric given in global isothermal coordinates (x, y) by

$$(1.18) \quad g = f(x)[dx^2 + dy^2].$$

Here f is a smooth positive function on \mathcal{S} of the variable x only and T^1 stands for the one dimensional torus. The boundary $\partial\mathcal{S}$ of \mathcal{S} is not connected and consists in two copies of T^1 , precisely

$$\partial\mathcal{S} = \Gamma_0 \cup \Gamma_1, \quad \Gamma_0 = \{0\} \times T^1, \quad \Gamma_1 = \{1\} \times T^1.$$

Let Γ_D and Γ_N be nonempty open subsets of ∂M . We denote $\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2)$ the associated partial DN map corresponding to Dirichlet data given on Γ_D and Neumann data measured on Γ_N . We shall prove

THEOREM 1.1. — *Let (\mathcal{S}, g) denote a Riemannian surface of the form (1.18), i.e.*

$$g = f(x)[dx^2 + dy^2],$$

with f a smooth positive function on \mathcal{S} . Let $\lambda^2 \neq 0$ be a fixed frequency. Let Γ_D and Γ_N be nonempty open subsets of ∂M that belong to distinct connected components of ∂M . Then there exists an infinite dimensional family of non-isometric metrics $\tilde{g}_c = cg$ of the form (1.18), parametrized by smooth positive functions $c = c(x)$, which satisfy $c(0) = c(1) = 1$ such that

$$\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{\tilde{g}_c, \Gamma_D, \Gamma_N}(\lambda^2).$$

We emphasize that the family of metrics \tilde{g}_c parametrized by the smooth positive functions $c = c(x)$ satisfying $c(0) = c(1) = 1$ is *explicit*. We refer to the end of the proof of Theorem 3.6 below for its complete description and more particularly to the formulae (3.48) and (3.49) for a countable family of examples. Conversely, at zero frequency $\lambda^2 = 0$ and still for Dirichlet and Neumann data measured on distinct connected components of ∂M ,

we show that a metric $g = f(x)[dx^2 + dy^2]$ can be uniquely determined by $\Lambda_{g, \Gamma_D, \Gamma_N}(0)$ up to the gauge invariance (1.7). We mention finally that in the case where the data Γ_D and Γ_N belong to the same connected component of ∂M , we can show that a metric $g = f(x)[dx^2 + dy^2]$ can be uniquely determined by $\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2)$ for any frequency λ^2 (up to the gauge invariance (1.7) when $\lambda^2 = 0$). This is proved in Theorem 3.2.

In which sense is our family of metrics \tilde{g}_c a counterexample to (Q1) and (Q2) when the data are measured on disjoint sets? Since we work at a fixed nonzero frequency $\lambda^2 \neq 0$, the only gauge invariance for the above partial anisotropic Calderón problem is a priori (1.6), i.e. g and \tilde{g}_c coincide up to the existence of a diffeomorphism $\phi : M \rightarrow M$ such that $\phi|_{\Gamma_D \cup \Gamma_N} = \text{Id}$ and $\phi^*g = \tilde{g}_c = cg$.

Assume the existence of such a diffeomorphism ϕ in the case $\Gamma_D \subset \Gamma_0$ and $\Gamma_N \subset \Gamma_1$. Then ϕ must send the boundary ∂M into itself, i.e. $\phi(\partial M) = \partial M$. More precisely, since by hypothesis $\phi|_{\Gamma_D} = \text{Id}$ and $\phi|_{\Gamma_N} = \text{Id}$, we see that in fact ϕ must send the connected components Γ_0 and Γ_1 of the boundary into themselves, i.e. $\phi(\Gamma_0) = \Gamma_0$ and $\phi(\Gamma_1) = \Gamma_1$. Let us denote now by $g_0 = f(0)dy^2$ and $g_1 = f(1)dy^2$ the metrics on Γ_0 and Γ_1 induced by g . Taking the restrictions to Γ_0 and Γ_1 of $\phi^*g = \tilde{g}_c = cg$, using $c(0) = c(1) = 1$ and our previous result, we get $\phi^*g_0 = g_0$ and $\phi^*g_1 = g_1$. In other words, $\phi|_{\Gamma_0}$ and $\phi|_{\Gamma_1}$ are isometries of (Γ_0, g_0) and (Γ_1, g_1) respectively. But the isometries of (Γ_0, g_0) and (Γ_1, g_1) are simply the transformations $y \mapsto \pm y + a$ for any constant a . Using our hypothesis $\phi|_{\Gamma_D} = \text{Id}$ and $\phi|_{\Gamma_N} = \text{Id}$ again, we see that the unique possibility is $\phi|_{\Gamma_0} = \phi|_{\Gamma_1} = \text{Id}$. We thus have $\phi|_{\partial M} = \text{Id}$. According to [35], the only diffeomorphism ϕ satisfying the previous properties is Id. We thus conclude that our family of metrics $\tilde{g}_c = cg$ cannot come from the pull back of the initial metric g by such a diffeomorphism and thus provide the claimed counterexamples.

We also solve the anisotropic Calderón problem (Q3) in the class of smooth compact Riemannian surfaces (1.18) for potentials $V \in L^\infty$ that only depend on the variable x . We show similarly that if the Dirichlet and Neumann data Γ_D and Γ_N belong to two distinct components of $\partial \mathcal{S}$, then there exists an infinite dimensional family of potentials $\tilde{V} = \tilde{V}(x)$ that satisfies

$$\Lambda_{g, V, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{g, \tilde{V}, \Gamma_D, \Gamma_N}(\lambda^2).$$

This is done in Theorem 3.8.

In 3 dimensions, we consider the family of smooth compact Riemannian manifolds (M, g) which have the topology of a toric cylinder $M = [0, 1] \times T^2$

and that are equipped with a metric

$$(1.19) \quad g = f(x)dx^2 + f(x)dy^2 + h(x)dz^2,$$

where f, h are smooth positive functions on M . Once again, the boundary ∂M of M is disconnected and consists in two copies of T^2 , precisely

$$\partial M = \Gamma_0 \cup \Gamma_1, \quad \Gamma_0 = \{0\} \times T^2, \quad \Gamma_1 = \{1\} \times T^2.$$

Let Γ_D and Γ_N be nonempty open subsets of ∂M . We denote $\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2)$ the associated partial DN map corresponding to Dirichlet data given on Γ_D and Neumann data measured on Γ_N . We shall prove

THEOREM 1.2. — *Let (M, g) and (M, \tilde{g}) denote two generic Riemannian manifolds of the form (1.19), i.e.*

$$g = f(x)dx^2 + f(x)dy^2 + h(x)dz^2, \quad \tilde{g} = \tilde{f}(x)dx^2 + \tilde{f}(x)dy^2 + \tilde{h}(x)dz^2.$$

Let $\lambda^2 \in \mathbb{R}$ be a fixed frequency and let Γ_D and Γ_N be nonempty open subsets of ∂M such that $\Gamma_D \cap \Gamma_N = \emptyset$. Then

- (1) *There exist infinitely many pairs of non isometric Riemannian manifolds (M, g) and (M, \tilde{g}) with $\tilde{g} = cg$ for some smooth positive strictly increasing or decreasing functions $c = c(x)$ satisfying*

- $c(0) = 1$ if $\Gamma_D, \Gamma_N \subset \Gamma_0$,
- $c(1) = 1$ if $\Gamma_D, \Gamma_N \subset \Gamma_1$,
- $c(1)^3 = c(0) \neq 1$ if $\Gamma_D \subset \Gamma_0$ and $\Gamma_N \subset \Gamma_1$,
- $c(0)^3 = c(1) \neq 1$ if $\Gamma_D \subset \Gamma_1$ and $\Gamma_N \subset \Gamma_0$,

such that

$$\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(\lambda^2).$$

- (2) *If moreover $\lambda^2 = 0$, there exists a one parameter family of Riemannian manifolds $(M, \tilde{g}_a)_{a>0}$ non isometric with the given Riemannian manifold (M, g) , satisfying*

$$\Lambda_{g, \Gamma_D, \Gamma_N}(0) = \Lambda_{\tilde{g}_a, \Gamma_D, \Gamma_N}(0).$$

The one parameter family of metrics $(\tilde{g}_a)_{a>0}$ has the form $\tilde{g}_a = c_a g$ where $c_a = c(x, a)$ are smooth positive strictly increasing or decreasing functions satisfying

- $c(0, a) = 1$ if $\Gamma_D, \Gamma_N \subset \Gamma_0$,
- $c(1, a) = 1$ if $\Gamma_D, \Gamma_N \subset \Gamma_1$,
- $c(x, a) > 1$ or $c(x, a) < 1$ if $\Gamma_D \subset \Gamma_0$ (resp. $\Gamma_D \subset \Gamma_1$) and $\Gamma_N \subset \Gamma_1$ (resp. $\Gamma_D \subset \Gamma_0$),

that are explicitly given in terms of g .

As in the 2 dimensional case, the only gauge invariance for the above partial anisotropic Calderón problems (Q1) and (Q2) with data measured on disjoint sets is a priori (1.6). Assume that there exists a diffeomorphism $\phi : M \rightarrow M$ such that $\phi|_{\Gamma_D \cup \Gamma_N} = \text{Id}$ and $\phi^*g = \tilde{g} = cg$. In particular, since ϕ is a diffeomorphism, we see that $\text{Vol}_g(M) = \text{Vol}_{\phi^*g}(M) = \text{Vol}_{cg}(M)$. Hence we must have

$$\int_M \sqrt{|g|} dx dy dz = \int_M c^{3/2} \sqrt{|g|} dx dy dz.$$

But in any case, we know that $c > 1$ or $c < 1$ on $(0, 1)$. Hence the above equality is impossible. We conclude that our family of metrics is not captured by this gauge invariance and thus provides counterexamples to uniqueness.

We emphasize that we could extend the results of Theorem 1.2 to higher dimensional Riemannian manifolds. Consider for instance n dimensional Riemannian manifolds (M, g) which have the topology of a toric cylinder $M = [0, 1] \times T^{n-1}$ and that are equipped with a metric

$$(1.20) \quad g = f(x)dx^2 + f(x)dy_1^2 + h_2(x)dy_2^2 + \cdots + h_{n-1}(x)dy_{n-1}^2,$$

where f, h_2, \dots, h_{n-1} are smooth positive functions on M . The analysis given for the 3 dimensional models extend in a straightforward way to such Riemannian manifolds and the results are basically the same.

We also solve the anisotropic Calderón problem (Q3) in the class of smooth compact Riemannian manifolds (1.19) for potentials $V \in L^\infty$ that only depend on the variable x . Contrary to Theorem 1.2, we show in Theorems 3.8 and 4.10 that if the Dirichlet and Neumann data Γ_D and Γ_N belong to the same connected component of ∂M and if $\Lambda_{g, V, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{g, \tilde{V}, \Gamma_D, \Gamma_N}(\lambda^2)$, (and a technical assumption on Γ_N in Theorem 4.10), then

$$V = \tilde{V}.$$

The anisotropic Calderón problems (Q2) and (Q3) are thus not equivalent in our 3 dimensional models. Moreover, if the Dirichlet and Neumann data Γ_D and Γ_N does not belong to the same connected component of ∂M , we show there exists an explicit infinite dimensional family of potentials \tilde{V} that satisfies $\Lambda_{g, V, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{g, \tilde{V}, \Gamma_D, \Gamma_N}(\lambda^2)$.

The remainder of this paper is devoted to the proofs of Theorems 1.1 and 1.2. In Section 2, we collect some results on the inverse spectral problem of one dimensional Schrödinger operators. In particular, we define the notion of characteristic and Weyl–Titchmarsh functions associated with one dimensional equation Schrödinger equations with certain boundary conditions and state a version of the Borg–Marchenko Theorem that will be

needed for our proof. We refer for instance to [2, 12, 14, 28, 34, 44] for a detailed account of these results.

The interest in one dimensional inverse spectral theory comes from the presence of symmetries in our models. These symmetries allow to decompose the solution of the Dirichlet problem and the DN map onto the Fourier modes $\{e^{imy}\}_{m \in \mathbb{Z}}$ in 2D and the Fourier modes $\{e^{imy}e^{inz}\}_{m,n \in \mathbb{Z}}$ in 3D. Hence we are able to reduce our initial problem to an infinite number of 1D inverse spectral problems for which the results recalled in Section 2 will be useful.

In Section 3, we describe the 2 dimensional models and construct first the global and partial DN maps that we aim to study as well as their restrictions onto the Fourier modes $\{e^{imy}\}_{m \in \mathbb{Z}}$. Using essentially the Complex Angular Momentum Method (see [6, 7, 8, 9, 39, 40]) that consists in allowing the angular momentum m to be complex, we solve in Theorem 3.2 an anisotropic Calderón problem with Dirichlet and Neumann data measured on some open subsets Γ_D and Γ_N of ∂M . We continue with the proof of our main Theorem 1.1 in two dimensions. We finish this Section solving the anisotropic Calderón problem (Q3) in Theorem 3.8 and giving new counterexamples to uniqueness in this setting.

In Section 4, we perform a similar analysis for the 3 dimensional models. We construct first the global and partial DN maps that we aim to study as well as their restrictions onto the Fourier modes $\{e^{imy+inz}\}_{m,n \in \mathbb{Z}}$. In Theorem 4.4, we solve the anisotropic Calderón problem with Dirichlet and Neumann data measured on some overlapping open subsets Γ_D or Γ_N belonging to the same connected component of ∂M . Once again here, the main tool is the Complex Angular Momentum method that makes possible to complexify the angular momenta m and n . We then proceed to prove our second main Theorem 1.2 in three dimensions. We finish this section showing that the anisotropic Calderón problem (Q3) has a unique solution in the case where the Dirichlet and Neumann data belong to the same connected component of the boundary. This is done in Theorem 4.10.

2. Preliminary results on 1D inverse spectral problems

We consider the class of ODE on the interval $[0, 1]$ given by

$$(2.1) \quad -v'' + q(x)v = -\mu^2 v, \quad q \in L^2([0, 1]), \quad q \text{ real},$$

with Dirichlet boundary conditions

$$(2.2) \quad v(0) = 0, \quad v(1) = 0.$$

Since the potential q belongs to $L^2([0, 1]) \subset L^1([0, 1])$, we can define for all $\mu \in \mathbb{C}$ two fundamental systems of solutions (FSS)

$$\{c_0(x, \mu^2), s_0(x, \mu^2)\}, \quad \{c_1(x, \mu^2), s_1(x, \mu^2)\},$$

of (2.1) by imposing the Cauchy conditions

$$(2.3) \quad \begin{cases} c_0(0, \mu^2) = 1, & c_0'(0, \mu^2) = 0, & s_0(0, \mu^2) = 0, & s_0'(0, \mu^2) = 1, \\ c_1(1, \mu^2) = 1, & c_1'(1, \mu^2) = 0, & s_1(1, \mu^2) = 0, & s_1'(1, \mu^2) = 1. \end{cases}$$

It follows from (2.3) that

$$(2.4) \quad W(c_0, s_0) = 1, \quad W(c_1, s_1) = 1, \quad \forall \mu \in \mathbb{C},$$

where $W(u, v) = uv' - u'v$ is the Wronskian of u, v . Moreover, the FSS $\{c_0(x, \mu^2), s_0(x, \mu^2)\}$ and $\{c_1(x, \mu^2), s_1(x, \mu^2)\}$ are entire functions with respect to the variable $\mu^2 \in \mathbb{C}$ by standard properties of ODEs depending analytically on a parameter.

We then define the characteristic function of (2.1) with boundary conditions (2.2) by

$$(2.5) \quad \Delta(\mu^2) = W(s_0, s_1).$$

The characteristic function $\mu^2 \mapsto \Delta(\mu^2)$ is entire with respect to μ^2 and its zeros $(\alpha_k^2)_{k \geq 1}$ correspond to “minus” the eigenvalues of the selfadjoint operator $-\frac{d^2}{dx^2} + q$ with Dirichlet boundary conditions. The eigenvalues α_k^2 are thus real and satisfy $-\infty < \dots < \alpha_2^2 < \alpha_1^2 < \infty$.

We next define two Weyl–Titchmarsh functions by the following classical prescriptions. Let the Weyl solutions Ψ and Φ be the unique solutions of (2.1) having the form

$$(2.6) \quad \begin{aligned} \Psi(x, \mu^2) &= c_0(x, \mu^2) + M(\mu^2)s_0(x, \mu^2), \\ \Phi(x, \mu^2) &= c_1(x, \mu^2) - N(\mu^2)s_1(x, \mu^2), \end{aligned}$$

which satisfy the Dirichlet boundary condition at $x = 1$ and $x = 0$ respectively. Then a short calculation using (2.3) shows that the Weyl–Titchmarsh functions $M(\mu^2)$ and $N(\mu^2)$ are uniquely defined by

$$(2.7) \quad M(\mu^2) = -\frac{W(c_0, s_1)}{\Delta(\mu^2)}, \quad N(\mu^2) = -\frac{W(c_1, s_0)}{\Delta(\mu^2)}.$$

For later use, we introduce the functions $D(\mu^2) = W(c_0, s_1)$ and $E(\mu^2) = W(c_1, s_0)$ which also turn out to be entire functions of μ^2 . We then have the following notation for the Weyl–Titchmarsh functions

$$M(\mu^2) = -\frac{D(\mu^2)}{\Delta(\mu^2)}, \quad N(\mu^2) = -\frac{E(\mu^2)}{\Delta(\mu^2)}.$$

We now collect some analytic results involving the functions $\Delta(\mu^2)$, $D(\mu^2)$, $E(\mu^2)$, $M(\mu^2)$ and $N(\mu^2)$ and give a version of the Borg–Marchenko Theorem we shall need later.

LEMMA 2.1. — *The FSS $\{c_0(x, \mu^2), s_0(x, \mu^2)\}$ and $\{c_1(x, \mu^2), s_1(x, \mu^2)\}$ have the following asymptotics uniformly with respect to $x \in [0, 1]$ as $|\mu| \rightarrow \infty$ in the complex plane \mathbb{C} .*

$$(2.8) \quad \begin{cases} c_0(x, \mu^2) = \cosh(\mu x) + O\left(\frac{e^{|\Re(\mu)|x}}{\mu}\right), \\ c'_0(x, \mu^2) = \mu \sinh(\mu x) + O\left(e^{|\Re(\mu)|x}\right), \\ s_0(x, \mu^2) = \frac{\sinh(\mu x)}{\mu} + O\left(\frac{e^{|\Re(\mu)|x}}{\mu^2}\right), \\ s'_0(x, \mu^2) = \cosh(\mu x) + O\left(\frac{e^{|\Re(\mu)|x}}{\mu}\right), \end{cases}$$

and

$$(2.9) \quad \begin{cases} c_1(x, \mu^2) = \cosh(\mu(1-x)) + O\left(\frac{e^{|\Re(\mu)|(1-x)}}{\mu}\right), \\ c'_1(x, \mu^2) = -\mu \sinh(\mu(1-x)) + O\left(e^{|\Re(\mu)|(1-x)}\right), \\ s_1(x, \mu^2) = -\frac{\sinh(\mu(1-x))}{\mu} + O\left(\frac{e^{|\Re(\mu)|(1-x)}}{\mu^2}\right), \\ s'_1(x, \mu^2) = \cosh(\mu(1-x)) + O\left(\frac{e^{|\Re(\mu)|(1-x)}}{\mu}\right). \end{cases}$$

Proof. — These asymptotics are classical and can be found in [37, Theorem 3, p. 13]. □

COROLLARY 2.2.

- (1) For each fixed $x \in [0, 1]$, the fundamental systems of solutions $\{c_0(x, \mu^2), s_0(x, \mu^2)\}$ and $\{c_1(x, \mu^2), s_1(x, \mu^2)\}$ are entire functions of order $\frac{1}{2}$ with respect to the variable μ^2 .
- (2) The characteristic function $\Delta(\mu^2)$ and the functions $D(\mu^2)$ and $E(\mu^2)$ are entire functions of order $\frac{1}{2}$ with respect to the variable μ^2 .
- (3) We have the following asymptotics in the complex plane \mathbb{C} :

$$\left\{ \begin{array}{l} \Delta(\mu^2) = \frac{\sinh \mu}{\mu} + O\left(\frac{e^{|\Re(\mu)|}}{\mu^2}\right), \\ D(\mu^2) = \cosh(\mu) + O\left(\frac{e^{|\Re(\mu)|}}{\mu}\right), \quad E(\mu^2) = \cosh(\mu) + O\left(\frac{e^{|\Re(\mu)|}}{\mu}\right). \end{array} \right.$$

In particular, $M(\mu^2) = \mp\mu + O(1)$, $N(\mu^2) = \mp\mu + O(1)$ when $\mu \rightarrow \pm\infty$, $\mu \in \mathbb{R}$.

Proof. — The proof of (1), (2) and (3) follows directly from (2.5), (2.7) and Lemma 2.1. \square

COROLLARY 2.3. — *The characteristic function $\Delta(\mu^2)$ and the functions $D(\mu^2)$ and $E(\mu^2)$ can be written as*

$$(2.10) \quad \begin{aligned} \Delta(\mu^2) &= A\mu^{2p} \prod_{k=1}^{\infty} \left(1 - \frac{\mu^2}{\alpha_k^2}\right), \\ D(\mu^2) &= B\mu^{2q} \prod_{k=1}^{\infty} \left(1 - \frac{\mu^2}{\beta_k^2}\right), \\ E(\mu^2) &= C\mu^{2r} \prod_{k=1}^{\infty} \left(1 - \frac{\mu^2}{\gamma_k^2}\right), \end{aligned}$$

where $(\alpha_k^2)_{k \geq 1}$, $(\beta_k^2)_{k \geq 1}$ and $(\gamma_k^2)_{k \geq 1}$ are the zeros of the entire functions $\Delta(\mu^2)$, $D(\mu^2)$ and $E(\mu^2)$ respectively, A, B, C are constants and p, q, r are the multiplicities of the root at the origin.

Proof. — This is a direct consequence of Hadamard's factorization Theorem (see [3, 34, 41]) for the entire functions $\Delta(\mu^2)$, $D(\mu^2)$ and $E(\mu^2)$ of order $\frac{1}{2}$. \square

Let us finally recall here the celebrated Borg–Marchenko Theorem which roughly speaking states that the Weyl–Titchmarsh functions M or N of the differential expression $-\frac{d^2}{dx^2} + q$ with regular boundary condition at $x = 0$ and $x = 1$ and with a real-valued potential q determines uniquely this potential. We refer for instance to [2, 4, 5, 12, 13, 14, 28, 36, 44] for historic and recent developments on the theory of WT functions. We state here a version from [12, Corollary 4.3] adapted to our particular problem.

THEOREM 2.4 (Borg–Marchenko). — *Consider two boundary value problems (2.1) with potentials q and \tilde{q} in $L^1([0, 1])$. Let $M(\mu^2)$ and $\widetilde{M}(\mu^2)$ the Weyl–Titchmarsh functions associated to (2.1) using (2.7). Then, if*

$$M(\mu^2) = \widetilde{M}(\mu^2) + f(\mu^2), \quad \mu^2 \in \mathbb{C} \setminus \mathbb{R},$$

for any f entire function of growth order at most $\frac{1}{2}$ (here the equality is understood on the domains of holomorphy of both sides of the equality, that is in this case, on \mathbb{C} minus the set of discrete eigenvalues of (2.1) that lie on the real axis), then

$$q = \tilde{q}, \quad \text{on } [0, 1].$$

Conversely, if $q = \tilde{q}$ on $[0, 1]$, then obviously $M = \tilde{M}$ on $\mathbb{C} \setminus \mathbb{R}$. Of course, the same results hold with $M(\mu^2)$ and $\tilde{M}(\mu^2)$ replaced by $N(\mu^2)$ and $\tilde{N}(\mu^2)$.

3. The two dimensional case

3.1. The model and the Dirichlet-to-Neumann map

In this Section, we work on a Riemannian surface (\mathcal{S}, g) which has the topology of a cylinder $M = [0, 1] \times T^1$ and that is equipped with a Riemannian metric given in global isothermal coordinates (x, y) by

$$(3.1) \quad g = f(x)[dx^2 + dy^2].$$

Here f is a smooth positive function on \mathcal{S} of the variable x only and T^1 stands for the circle. The metric g obviously possesses one Killing vector field ∂_y that generates the cylindrical symmetry of our surface of revolution. The boundary $\partial\mathcal{S}$ of \mathcal{S} consists in two copies of T^1 , precisely

$$\partial\mathcal{S} = \Gamma_0 \cup \Gamma_1, \quad \Gamma_0 = \{0\} \times T^1, \quad \Gamma_1 = \{1\} \times T^1.$$

In our global coordinates system (x, y) , the positive Laplace–Beltrami operator takes the expression

$$-\Delta_g = \frac{1}{f} (-\partial_x^2 - \partial_y^2).$$

It is well known that the Laplace–Beltrami operator $-\Delta_g$ with Dirichlet boundary conditions is selfadjoint on $L^2(\mathcal{S}, d\text{Vol})$ and has pure point spectrum $\{\lambda_j^2\}_{j \geq 1}$ so that $0 < \lambda_1^2 < \lambda_2^2 \leq \dots \leq \lambda_j^2 \rightarrow +\infty$.

Consider the Dirichlet problem at a frequency λ^2 such that $\lambda \notin \{\lambda_j^2\}_{j \geq 1}$. That is we look at the solutions u of the following PDE

$$-\Delta_g u = \lambda^2 u, \quad \text{on } \mathcal{S},$$

which can be rewritten in our coordinates system as

$$(3.2) \quad -\partial_x^2 u - \partial_y^2 u - \lambda^2 f u = 0, \quad \text{on } \mathcal{S},$$

with the boundary conditions

$$(3.3) \quad u = \psi \quad \text{on } \partial\mathcal{S}.$$

The DN map $\Lambda_g(\lambda^2)$ is then defined by (1.2) as

$$(3.4) \quad \Lambda_g(\lambda^2)(\psi) = (\partial_\nu u)|_{\partial\mathcal{S}},$$

where u is the unique solution in $H^1(\mathcal{S})$ of (3.2) and $(\partial_\nu u)|_{\partial\mathcal{S}} \in H^{-1/2}(\partial\mathcal{S})$ is the weak normal derivative of u on $\partial\mathcal{S}$.

In order to find a nice representation of the DN map, we introduce certain notations. Since the boundary $\partial\mathcal{S}$ of \mathcal{S} has two disjoint components $\partial\mathcal{S} = \Gamma_0 \cup \Gamma_1$, we may decompose the Sobolev spaces $H^s(\partial\mathcal{S})$ as $H^s(\partial\mathcal{S}) = H^s(\Gamma_0) \oplus H^s(\Gamma_1)$ for any $s \in \mathbb{R}$. We also recall that $\Gamma_0 = \Gamma_1 = T^1$. Moreover, we shall use the vector notation

$$\varphi = \begin{pmatrix} \varphi^0 \\ \varphi^1 \end{pmatrix},$$

for all elements φ of $H^s(\partial\mathcal{S}) = H^s(\Gamma_0) \oplus H^s(\Gamma_1)$. Finally, since the DN map is a linear operator from $H^{1/2}(\partial\mathcal{S})$ to $H^{-1/2}(\partial\mathcal{S})$, it has the structure of an operator valued 2×2 matrix

$$\Lambda_g(\lambda^2) = \begin{pmatrix} L(\lambda^2) & T_R(\lambda^2) \\ T_L(\lambda^2) & R(\lambda^2) \end{pmatrix},$$

where $L(\lambda^2), R(\lambda^2), T_R(\lambda^2), T_L(\lambda^2)$ are bounded operators from $H^{1/2}(T^1)$ to $H^{-1/2}(T^1)$. The operators $L(\lambda^2), R(\lambda^2)$ correspond to the partial DN map whose measurements are restricted to Γ_0 and Γ_1 respectively whereas the operators $T_R(\lambda^2), T_L(\lambda^2)$ correspond to the partial DN maps whose measurements are made on the disjoint sets Γ_0 and Γ_1 . For instance, $T_L(\lambda^2)$ is the operator from $H^{1/2}(\Gamma_0)$ to $H^{-1/2}(\Gamma_1)$ given by

$$T_L(\lambda^2)(\psi^0) = (\partial_\nu u)|_{\Gamma_1},$$

where u is the unique solution of (3.2)–(3.3) such that $\text{supp } \psi^0 \subset \Gamma_0$.

To summarize, using the notations in the Introduction, we have the following dictionary

$$\begin{aligned} L(\lambda^2) &= \Lambda_{g,\Gamma_0}(\lambda^2), & R(\lambda^2) &= \Lambda_{g,\Gamma_1}(\lambda^2), \\ T_L(\lambda^2) &= \Lambda_{g,\Gamma_0,\Gamma_1}(\lambda^2), & T_R(\lambda^2) &= \Lambda_{g,\Gamma_1,\Gamma_0}(\lambda^2). \end{aligned}$$

Now we take advantage of the cylindrical symmetry of (\mathcal{S}, g) to find a simple expression of the DN map. We first write $\psi = (\psi^0, \psi^1) \in H^{1/2}(\Gamma_0) \times H^{1/2}(\Gamma_1)$ using their Fourier series representation as

$$\psi^0 = \sum_{m \in \mathbb{Z}} \psi_m^0 Y_m, \quad \psi^1 = \sum_{m \in \mathbb{Z}} \psi_m^1 Y_m,$$

where

$$Y_m(y) = e^{imy}.$$

Note that for any $s \in \mathbb{R}$, the space $H^s(T^1)$ can be described as

$$\left\{ \varphi \in \mathcal{D}'(T^1), \quad \varphi = \sum_{m \in \mathbb{Z}} \varphi_m Y_m, \quad \sum_{m \in \mathbb{Z}} (1 + m^2)^s |\varphi_m|^2 < \infty \right\}.$$

Therefore, the unique solution u of (3.2)–(3.3) can be looked for under the form

$$u = \sum_{m \in \mathbb{Z}} u_m(x) Y_m(y),$$

where for all $m \in \mathbb{Z}$, the functions u_m are the unique solutions of the ODEs (w.r.t. x) with boundary conditions

$$(3.5) \quad \begin{cases} -u_m'' + m^2 u_m - \lambda^2 f(x) u_m = 0, \\ u_m(0) = \psi_m^0, \quad u_m(1) = \psi_m^1. \end{cases}$$

The DN map can now be diagonalized on the Hilbert basis $\{Y_m\}_{m \in \mathbb{Z}}$ and can be shown to have a very simple expression on each Fourier mode. First, a short calculation using the particular form of the metric (3.1) and for smooth enough Dirichlet data ψ shows that

$$\Lambda_g(\lambda^2) \begin{pmatrix} \psi^0 \\ \psi^1 \end{pmatrix} = \begin{pmatrix} (\partial_\nu u)|_{\Gamma_0} \\ (\partial_\nu u)|_{\Gamma_1} \end{pmatrix} = \begin{pmatrix} -\frac{1}{\sqrt{f(0)}} (\partial_x u)|_{x=0} \\ \frac{1}{\sqrt{f(1)}} (\partial_x u)|_{x=1} \end{pmatrix}.$$

Hence, if we let the DN map act on the vector space generated by the Fourier mode Y_m , we get

$$(3.6) \quad \Lambda_g(\lambda^2) \begin{pmatrix} \psi_m^0 Y_m \\ \psi_m^1 Y_m \end{pmatrix} = \begin{pmatrix} -\frac{1}{\sqrt{f(0)}} u'_m(0) Y_m \\ \frac{1}{\sqrt{f(1)}} u'_m(1) Y_m \end{pmatrix}.$$

We denote by

$$\Lambda_g(\lambda^2)|_{\langle Y_m \rangle} = \Lambda_g^m(\lambda^2) = \begin{pmatrix} L^m(\lambda^2) & T_R^m(\lambda^2) \\ T_L^m(\lambda^2) & R^m(\lambda^2) \end{pmatrix},$$

the 2×2 matrix corresponding to the restriction of the DN map to each Fourier mode $\langle Y_m \rangle$. This operator is clearly defined for all $m \in \mathbb{Z}$ and for all $(\psi_m^0, \psi_m^1) \in \mathbb{C} \oplus \mathbb{C}$ by

$$(3.7) \quad \Lambda_g^m(\lambda^2) \begin{pmatrix} \psi_m^0 \\ \psi_m^1 \end{pmatrix} = \begin{pmatrix} -\frac{1}{\sqrt{f(0)}} u'_m(0) \\ \frac{1}{\sqrt{f(1)}} u'_m(1) \end{pmatrix}.$$

We can further simplify the partial DN maps $\Lambda_g^m(\lambda^2)$ by interpreting their coefficients as the characteristic and Weyl–Titchmarsh functions of the 1D equation (3.5) with appropriate boundary conditions. Precisely, consider the ODE

$$(3.8) \quad -v'' + q_\lambda(x)v = -\mu^2 v, \quad q_\lambda = -\lambda^2 f,$$

with boundary conditions

$$(3.9) \quad v(0) = 0, \quad v(1) = 0.$$

Note that the equation (3.8) is nothing but equation (3.5) in which the parameter $-m^2$ is written as $-\mu^2$ and is interpreted as the spectral parameter of the equation. On the other hand, the boundary conditions (3.3) have been replaced by Dirichlet boundary conditions at $x = 0$ and $x = 1$. Since the potential q_λ lies in $L^1([0, 1])$ and is real, the boundary value problem (3.8)–(3.9) lies in the framework recalled in Section 2. Hence, for all $\mu \in \mathbb{C}$, we can define the fundamental systems of solutions

$$\{c_0(x, \mu^2), s_0(x, \mu^2)\}, \quad \{c_1(x, \mu^2), s_1(x, \mu^2)\},$$

of (3.8) by imposing the Cauchy conditions

$$(3.10) \quad \begin{cases} c_0(0, \mu^2) = 1, & c'_0(0, \mu^2) = 0, & s_0(0, \mu^2) = 0, & s'_0(0, \mu^2) = 1, \\ c_1(1, \mu^2) = 1, & c'_1(1, \mu^2) = 0, & s_1(1, \mu^2) = 0, & s'_1(1, \mu^2) = 1. \end{cases}$$

Note that the dependence of the FSS on λ^2 is not written for clarity but implicit.

We then define the characteristic function of (3.8) with boundary conditions (3.9) by

$$(3.11) \quad \Delta(\mu^2) = W(s_0, s_1),$$

whereas the Weyl–Titchmarsh functions are defined by

$$(3.12) \quad \begin{aligned} M(\mu^2) &= -\frac{W(c_0, s_1)}{\Delta(\mu^2)} = -\frac{D(\mu^2)}{\Delta(\mu^2)}, \\ N(\mu^2) &= -\frac{W(c_1, s_0)}{\Delta(\mu^2)} = -\frac{E(\mu^2)}{\Delta(\mu^2)}. \end{aligned}$$

We refer to Section 2 for the notations.

Remark 3.1.

- (1) In the case $\lambda^2 = 0$, we get immediately the following explicit formulas for the characteristic and WT functions.

$$(3.13) \quad \begin{aligned} c_0(x, \mu^2) &= \cosh(\mu x), & s_0(x, \mu^2) &= \frac{\sinh(\mu x)}{\mu}, \\ c_1(x, \mu^2) &= \cosh(\mu(1-x)), & s_1(x, \mu^2) &= -\frac{\sinh(\mu(1-x))}{\mu}, \\ \Delta(\mu^2) &= \frac{\sinh(\mu)}{\mu}, & D(\mu^2) &= E(\mu^2) = \cosh(\mu), \\ M(\mu^2) &= -\mu \coth(\mu), & N(\mu^2) &= -\mu \coth(\mu). \end{aligned}$$

- (2) The WT function $M(\mu^2)$ is meromorphic on \mathbb{C} and has an infinite and discrete set of poles $\{\alpha_k^2\}_{k \geq 1}$ corresponding to “minus” the Dirichlet eigenvalues of $-\frac{d}{dx^2} + q_\lambda$ or equivalently, corresponding

to the zeros of the characteristic function $\Delta(\mu^2)$. Let us show that the integers $m^2, m \in \mathbb{Z}$ cannot be poles of $M(\mu^2)$ under our general assumption. Assume the converse, i.e. there exists $m \in \mathbb{Z}$ such that m^2 is a pole of $M(\mu^2)$. Thus $-m^2$ is a Dirichlet eigenvalue of $-\frac{d}{dx^2} + q_\lambda$ and we denote by $u_m(x)$ the associated eigenfunction solution of (3.5). Then the function $u(x, y) = u_m(x)Y_m(y)$ is a nontrivial solution of (3.2) with Dirichlet boundary conditions. We conclude that u is an eigenfunction of $-\Delta_g$ associated to the Dirichlet eigenvalue λ^2 . But this case has been ruled out from the beginning since we assume that λ^2 is not a Dirichlet eigenvalue of Δ_g . Whence the contradiction.

In particular, we see that 0 cannot be a pole of $M(\mu^2)$ or in other words, that the characteristic function $\Delta(\mu^2)$ doesn't vanish at 0. Using Corollary 2.3, we have thus the following factorization for $\Delta(\mu^2)$ that will be used later.

$$(3.14) \quad \Delta(\mu^2) = \Delta(0) \prod_{k=1}^{\infty} \left(1 - \frac{\mu^2}{\alpha_k^2}\right).$$

We now come back to the expression (3.7) of the partial DN map $\Lambda_g^m(\lambda^2)$. For all $m \in \mathbb{Z}$, we need to express $u'_m(0)$ and $u'_m(1)$ in terms of ψ_m^0 and ψ_m^1 in order to find the expressions of the coefficients $L^m(\lambda^2), T_R^m(\lambda^2), T_L^m(\lambda^2)$ and $R^m(\lambda^2)$. But the solution u_m of (3.5) can be written as linear combinations of the FSS $\{c_0(x, m^2), s_0(x, m^2)\}$ and $\{c_1(x, m^2), s_1(x, m^2)\}$. Precisely, we have

$$u_m(x) = \alpha c_0(x, m^2) + \beta s_0(x, m^2) = \gamma c_1(x, m^2) + \delta s_1(x, m^2),$$

for some constants $\alpha, \beta, \gamma, \delta$. Using (3.5) and (3.10), we first get

$$(3.15) \quad \begin{pmatrix} u_m(0) \\ u_m(1) \end{pmatrix} = \begin{pmatrix} \psi_m^0 \\ \psi_m^1 \end{pmatrix} = \begin{pmatrix} \alpha \\ \gamma \end{pmatrix} = \begin{pmatrix} \gamma c_1(0, m^2) + \delta s_1(0, m^2) \\ \alpha c_0(1, m^2) + \beta s_0(1, m^2) \end{pmatrix}.$$

From (3.15), we obtain in particular

$$(3.16) \quad \begin{pmatrix} \beta \\ \delta \end{pmatrix} = \begin{pmatrix} -\frac{c_0(1, m^2)}{s_0(1, m^2)} & \frac{1}{s_0(1, m^2)} \\ \frac{1}{s_1(0, m^2)} & -\frac{c_1(0, m^2)}{s_1(0, m^2)} \end{pmatrix} \begin{pmatrix} \psi_m^0 \\ \psi_m^1 \end{pmatrix}.$$

Also using (3.10) and (3.16), we have

$$(3.17) \quad \begin{pmatrix} u'_m(0) \\ u'_m(1) \end{pmatrix} = \begin{pmatrix} \beta \\ \delta \end{pmatrix} = \begin{pmatrix} -\frac{c_0(1, m^2)}{s_0(1, m^2)} & \frac{1}{s_0(1, m^2)} \\ \frac{1}{s_1(0, m^2)} & -\frac{c_1(0, m^2)}{s_1(0, m^2)} \end{pmatrix} \begin{pmatrix} \psi_m^0 \\ \psi_m^1 \end{pmatrix}.$$

Therefore by (3.7) and (3.15)–(3.17), we obtain.

$$(3.18) \quad \Lambda_g^m(\lambda^2) \begin{pmatrix} \psi_m^0 \\ \psi_m^1 \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{f(0)}} \frac{c_0(1, m^2)}{s_0(1, m^2)} & -\frac{1}{\sqrt{f(0)}} \frac{1}{s_0(1, m^2)} \\ \frac{1}{\sqrt{f(1)}} \frac{1}{s_1(0, m^2)} & -\frac{1}{\sqrt{f(1)}} \frac{c_1(0, m^2)}{s_1(0, m^2)} \end{pmatrix} \begin{pmatrix} \psi_m^0 \\ \psi_m^1 \end{pmatrix}.$$

Hence, we obtain the following expression for the partial DN map $\Lambda_g^m(\lambda^2)$.

$$(3.19) \quad \Lambda_g^m(\lambda^2) = \begin{pmatrix} \frac{1}{\sqrt{f(0)}} \frac{c_0(1, m^2)}{s_0(1, m^2)} & -\frac{1}{\sqrt{f(0)}} \frac{1}{s_0(1, m^2)} \\ \frac{1}{\sqrt{f(1)}} \frac{1}{s_1(0, m^2)} & -\frac{1}{\sqrt{f(1)}} \frac{c_1(0, m^2)}{s_1(0, m^2)} \end{pmatrix}.$$

Finally, using (3.10) again, we easily show that

$$(3.20) \quad \begin{aligned} \Delta(m^2) &= s_0(1, m^2) = -s_1(0, m^2), \\ M(m^2) &= -\frac{c_0(1, m^2)}{s_0(1, m^2)}, \\ N(m^2) &= \frac{c_1(0, m^2)}{s_1(0, m^2)}. \end{aligned}$$

Therefore

$$(3.21) \quad \Lambda_g^m(\lambda^2) = \begin{pmatrix} -\frac{1}{\sqrt{f(0)}} M(m^2) & -\frac{1}{\sqrt{f(0)}} \frac{1}{\Delta(m^2)} \\ -\frac{1}{\sqrt{f(1)}} \frac{1}{\Delta(m^2)} & -\frac{1}{\sqrt{f(1)}} N(m^2) \end{pmatrix}.$$

We emphasize that the quantities $\Delta(m^2)$ and $M(m^2)$ are well defined for all $m \in \mathbb{Z}$ thanks to our assumption on λ^2 , i.e. λ^2 is not a Dirichlet eigenvalue of Δ_g (see Remark 3.1 for this point).

The result above shows that the coefficients of $\Lambda_g^m(\lambda^2)$ which correspond to the global DN map $\Lambda_g(\lambda^2)$ restricted to the Fourier mode $\langle Y_m \rangle$, can be simply interpreted in terms of the characteristic and Weyl–Titchmarsh functions $\Delta(m^2)$, $M(m^2)$ and $N(m^2)$ associated to the simple ODE (3.8) with boundary data (3.9). In particular, the partial DN maps $L(\lambda^2)$ and $R(\lambda^2)$ which correspond to the global DN map with data restricted to Γ_0 and Γ_1 respectively, only depend on the Weyl–Titchmarsh functions $M(m^2)$ and $N(m^2)$ respectively (modulo some explicit constants). In fact, it is clear in our model that the knowledge of $L(\lambda^2)$ and $R(\lambda^2)$ is equivalent to the knowledge of the sequence of Weyl–Titchmarsh functions $\{M(m^2)\}_{m \in \mathbb{Z}}$ and $\{N(m^2)\}_{m \in \mathbb{Z}}$ respectively (modulo some constants).

Similarly, the knowledge of the partial DN maps $T_L(\lambda^2)$ and $T_R(\lambda^2)$ which correspond to the global DN map whose data are measured on the disjoint sets Γ_0 and Γ_1 , is equivalent to the knowledge of the characteristic functions $\{\Delta(m^2)\}_{m \in \mathbb{Z}}$ (modulo some explicit constants). We emphasize

that this is the key point that explains our non-uniqueness result for the Calderón problem with data measured on disjoint sets belonging to two distinct connected components of the boundary. Indeed, it is well known that the characteristic function $\Delta(\mu^2)$ does not contain enough information to determine uniquely the potential q_λ in (3.8) and thus the metric g .

3.2. The 2D anisotropic Calderón problem with data measured on the same connected component.

In this Section, we solve the Calderón problem from the knowledge of the partial DN map $\Lambda_{g,\Gamma_D,\Gamma_N}$, where Γ_D and Γ_N are any open subsets living in the same connected component of the boundary, (i.e, we assume that Γ_D, Γ_N are subsets of, either $\{0\} \times T^1$, or $\{1\} \times T^1$). Precisely, we prove the following result.

THEOREM 3.2. — *Let (\mathcal{S}, g) and (\mathcal{S}, \tilde{g}) two Riemannian surfaces of the form (3.1), i.e.*

$$g = f(x)[dx^2 + dy^2], \quad \tilde{g} = \tilde{f}(x)[dx^2 + dy^2].$$

We shall add the subscript \sim to all the quantities referring to (\mathcal{S}, \tilde{g}) . Let the frequency λ^2 be fixed and let Γ_D, Γ_N be subsets belonging to the same connected component of $\partial\mathcal{S}$. We assume that

$$\Lambda_{g,\Gamma_D,\Gamma_N} = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}.$$

Then

- (1) *If $\lambda^2 = 0$, there exists a function $c > 0$ with $c(0) = 1$ if $\Gamma_D, \Gamma_N \subset \{0\} \times T^1$, and $c(1) = 1$ if $\Gamma_D, \Gamma_N \subset \{1\} \times T^1$ such that*

$$\tilde{g} = cg.$$

- (2) *If $\lambda^2 \neq 0$, we have $\tilde{g} = g$.*

Remark 3.3. — In the case $\lambda^2 = 0$ and when Γ_D and Γ_N are not disjoint, this result is well known and has been proved first for any smooth Riemannian surface by Lassas and Uhlmann in [32]. We also refer to the work of Guillarmou and Tzou [16] for similar results and to [17] for a recent survey on 2D inverse Calderón problem.

Proof. — Assume for instance that $\Gamma_D, \Gamma_N \subset \{0\} \times T^1$. As usually, we identify $\{0\} \times T^1$ with $[-\pi, \pi]$. Without loss of generality, we can always

assume that $0 \in \Gamma_D$ and $\pm\pi \in \Gamma_N$. The general case requires minor modifications. It follows from the hypothesis that, for all $\Psi \in C_0^\infty(-R, R)$, R small enough, ($R < \pi$), and for all $y \in \Gamma_N$,

$$(3.22) \quad \sum_{m \in \mathbb{Z}} \frac{1}{\sqrt{f(0)}} M(m^2) \hat{\Psi}(m) e^{imy} = \sum_{m \in \mathbb{Z}} \frac{1}{\sqrt{\tilde{f}(0)}} \tilde{M}(m^2) \hat{\Psi}(m) e^{imy},$$

where

$$(3.23) \quad \hat{\Psi}(m) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \Psi(y) e^{-imy} dy$$

is the m^{th} Fourier coefficient of Ψ . We set

$$(3.24) \quad \Phi(y) = \sum_{m \in \mathbb{Z}} \left(\frac{1}{\sqrt{f(0)}} M(m^2) - \frac{1}{\sqrt{\tilde{f}(0)}} \tilde{M}(m^2) \right) \hat{\Psi}(m) e^{imy}.$$

From (3.22), we deduce that $\text{supp}\Phi \subset (-R', R')$ for some $R' < \pi$, and clearly, the m^{th} Fourier coefficient of Φ is

$$(3.25) \quad \hat{\Phi}(m) = \left(\frac{1}{\sqrt{f(0)}} M(m^2) - \frac{1}{\sqrt{\tilde{f}(0)}} \tilde{M}(m^2) \right) \hat{\Psi}(m).$$

By the Paley–Wiener Theorem ([3, Theorem 6.8.1]), $\hat{\Psi}(m)$, (resp. $\hat{\Phi}(m)$) can be extended as an entire function, $\mu \rightarrow \hat{\Psi}(\mu)$, (resp. $\mu \rightarrow \hat{\Phi}(\mu)$) of order 1, of type strictly less than π . More precisely, there exists $A > 0$ and $B < \pi$ such that

$$(3.26) \quad |\hat{\Psi}(\mu)| \leq Ae^{B|\mu|}, |\hat{\Phi}(\mu)| \leq Ae^{B|\mu|}, \quad \forall \mu \in \mathbb{C}.$$

Now, we can establish the following elementary lemma:

LEMMA 3.4. — For all $\mu \in \mathbb{C}$,

$$F(\mu) := \Delta(\mu^2) \tilde{\Delta}(\mu^2) \hat{\Phi}(\mu) + \left(\frac{1}{\sqrt{f(0)}} D(\mu^2) \tilde{\Delta}(\mu^2) - \frac{1}{\sqrt{\tilde{f}(0)}} \tilde{D}(\mu^2) \Delta(\mu^2) \right) \hat{\Psi}(\mu) = 0.$$

Proof. — Using (3.26) and Corollary 2.2, we see that $F(\mu)$ is an entire function of order 1 satisfying the following estimate on the imaginary axis:

$$(3.27) \quad |F(iy)| \leq Ce^{B|y|}, \quad \forall y \in \mathbb{R},$$

for some $C > 0$. Moreover, since $M(m^2) = -\frac{D(m^2)}{\Delta(m^2)}$, we deduce from (3.25) that $F(m) = 0, \forall m \in \mathbb{Z}$. Since $B < \pi$, using Carlson’s theorem, (see [3, Theorem 9.2.1]), we have $F(\mu) = 0$ for all $\mu \in \mathbb{C}$. \square

We can deduce from this lemma:

PROPOSITION 3.5. — Assume that $\Lambda_{g,\Gamma_D,\Gamma_N} = \widetilde{\Lambda}_{g,\Gamma_D,\Gamma_N}$, with

$$\Gamma_D, \Gamma_N \subset \{0\} \times T^1.$$

Then:

(1) The zeros of $\Delta(\mu^2)$ and $\widetilde{\Delta}(\mu^2)$ coincide, and in particular, for all $\mu \in \mathbb{C}$,

$$(3.28) \quad \Delta(\mu^2) = \widetilde{\Delta}(\mu^2).$$

(2) For all $\mu \in \mathbb{C} \setminus \mathbb{R}$,

$$(3.29) \quad M(\mu^2) - M(0) = \widetilde{M}(\mu^2) - \widetilde{M}(0)$$

Proof. — First, we recall that the zeros of $\Delta(\mu^2)$, (resp. $\widetilde{\Delta}(\mu^2)$) are simple (see for instance [37, Theorem 2, p. 30]). Second, let us show that the zeros of $\Delta(\mu^2)$ and $\widetilde{\Delta}(\mu^2)$ coincide. For instance, assume that $\Delta(\alpha_k^2) = 0, k \geq 1$. Taking in Lemma 3.4, $\Psi \in C_0^\infty(-R, R)$ such that $\widehat{\Psi}(\alpha_k) = 1$, we obtain

$$D(\alpha_k^2)\widetilde{\Delta}(\alpha_k^2) = 0.$$

But, by definition, $D(\mu^2)$ and $\Delta(\mu^2)$ cannot vanish at the same time. We deduce then that $\widetilde{\Delta}(\alpha_k^2) = 0$.

Hence we infer from Corollary 2.3 and (3.14) that

$$(3.30) \quad \Delta(\mu^2) = \frac{\Delta(0)}{\widetilde{\Delta}(0)}\widetilde{\Delta}(\mu^2), \quad \forall \mu \in \mathbb{C}.$$

But we know from Corollary 2.2 that $\Delta(\mu^2) \sim \widetilde{\Delta}(\mu^2)$ when $\mu \rightarrow +\infty$. We deduce that $\Delta(0) = \widetilde{\Delta}(0)$ and then $\Delta(\mu^2) = \widetilde{\Delta}(\mu^2)$ for all $\mu \in \mathbb{C}$. This equality will be useful in the next section.

Now, let us prove the assertion (2) of Proposition 3.5. First, using Lemma 3.4, we obtain: for all $\mu \in \mathbb{C} \setminus \mathbb{R}$,

$$(3.31) \quad \widehat{\Phi}(\mu) + \left(\frac{1}{\sqrt{f(0)}}M(\mu^2) - \frac{1}{\sqrt{\widetilde{f}(0)}}\widetilde{M}(\mu^2) \right) \widehat{\Psi}(\mu) = 0$$

We set $m(\mu) = M(\mu^2)$, (resp. $\widetilde{m}(\mu) = \widetilde{M}(\mu^2)$). Clearly, m , (resp. \widetilde{m}) is a meromorphic function with simple poles $\pm\alpha_k$, (we recall that $\alpha_k \neq 0$). Moreover, $\text{Res}(m; -\alpha_k) = -\text{Res}(m; \alpha_k)$, (resp. $\text{Res}(\widetilde{m}; -\alpha_k) = -\text{Res}(\widetilde{m}; \alpha_k)$). Now, in (3.31), taking again a function Ψ such that $\widehat{\Psi}(\alpha_k) =$

1 or $\hat{\Psi}(-\alpha_k) = 1$, and since $\hat{\Phi}(\mu)$ is entire, we see that the residues at $\pm\alpha_k$ of $\frac{1}{\sqrt{f(0)}}m(\mu)$ and $\frac{1}{\sqrt{\tilde{f}(0)}}\tilde{m}(\mu)$ must be the same. So, we have proved that:

$$(3.32) \quad \frac{1}{\sqrt{f(0)}} \operatorname{Res}(m; \pm\alpha_k) = \frac{1}{\sqrt{\tilde{f}(0)}} \operatorname{Res}(\tilde{m}; \pm\alpha_k).$$

Secondly, we set

$$(3.33) \quad F(\mu) = \frac{m(\mu) - m(0)}{\mu}.$$

Clearly, $F(\mu)$ is a meromorphic function with simple poles at the zeros $\pm\alpha_k$, $k \geq 1$. Moreover, using the asymptotics of $D(\mu^2)$ and $\Delta(\mu^2)$ given in Corollary 2.2, a standard calculation (see for instance, [43, Chapter 7, p. 227]) shows there exists a constant $C > 0$ such that for all $N \in \mathbb{N}$:

$$(3.34) \quad |F(z)| \leq C, \quad \forall z \in C_N,$$

where C_N is the square with vertices $z = (\pm 1 \pm i)(N + \frac{1}{2})\pi$. We deduce from (3.34) that

$$(3.35) \quad \lim_{N \rightarrow +\infty} \int_{C_N} \frac{F(z)}{z(z - \mu)} dz = 0,$$

where $\mu \neq 0, \pm\alpha_k$ is fixed. Then, using the Cauchy's residue theorem, we obtain the well-known Mittag-Leffler's expansion (see [43, Chapter 7, p. 210] for details):

$$(3.36) \quad F(\mu) = F(0) + \sum_{k=1}^{+\infty} \operatorname{Res}(F; \alpha_k) \left(\frac{1}{\mu - \alpha_k} + \frac{1}{\alpha_k} \right) \\ + \sum_{k=1}^{+\infty} \operatorname{Res}(F; -\alpha_k) \left(\frac{1}{\mu + \alpha_k} - \frac{1}{\alpha_k} \right).$$

Clearly, $F(0) = 0$ since $m(\mu)$ is an even function of μ and we also have:

$$(3.37) \quad \operatorname{Res}(F; \pm\alpha_k) = \pm \frac{\operatorname{Res}(m; \pm\alpha_k)}{\alpha_k}.$$

So, we can rewrite (3.36) as

$$(3.38) \quad M(\mu^2) - M(0) = 2 \sum_{k=1}^{+\infty} \frac{\operatorname{Res}(m; \alpha_k)}{\alpha_k} \frac{\mu^2}{\mu^2 - \alpha_k^2}.$$

Thus, with the help of (3.32), we obtain:

$$(3.39) \quad \frac{1}{\sqrt{f(0)}} (M(\mu^2) - M(0)) = \frac{1}{\sqrt{\tilde{f}(0)}} \left(\tilde{M}(\mu^2) - \tilde{M}(0) \right).$$

Using the asymptotics of the Weyl–Titchmarsh function $M(\mu^2)$ as $\mu \rightarrow \infty$ given in Corollary 2.2, we get $f(0) = \tilde{f}(0)$ and the Proposition is proved. \square

Now, we can finish the proof of Theorem 3.2. First, let us examine the obvious case $\lambda^2 = 0$, (which can probably be done differently). When $\lambda^2 = 0$, the WT functions $M(\mu^2)$, (resp. $\tilde{M}(\mu^2)$) do not depend on the metric, (see (3.13)). So the unique condition we can get from the hypothesis $\Lambda_{g, \Gamma_D, \Gamma_N} = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}$ is $f(0) = \tilde{f}(0)$, which was obtained above. Thus, we have:

$$\tilde{g} = cg, \quad c(x) = \frac{\tilde{f}(x)}{f(x)},$$

with

$$c(0) = 1.$$

Now, let us study the more interesting case $\lambda^2 \neq 0$. Recalling that the WT functions $M(\mu^2)$ and $\tilde{M}(\mu^2)$ are associated to equation (3.8) with potentials q_λ and \tilde{q}_λ , we thus conclude from (3.29) and the Borg–Marchenko Theorem 2.4 that

$$(3.40) \quad q_\lambda(x) = \tilde{q}_\lambda(x), \quad \forall x \in [0, 1].$$

In other words, $f(x) = \tilde{f}(x), \forall x \in [0, 1]$ and thus $\tilde{g} = g$. This proves the result. \square

3.3. The 2D anisotropic Calderón problem with data measured on distinct connected components.

In this Section, we prove our first main theorem, namely we give a counterexample to uniqueness for the anisotropic Calderón problem, at a nonzero frequency, for Riemannian surfaces with partial data measured on sets belonging to two distinct connected components. Precisely, we prove:

THEOREM 3.6. — *Let (\mathcal{S}, g) and (\mathcal{S}, \tilde{g}) two Riemannian surfaces of the form (3.1), i.e.*

$$g = f(x)[dx^2 + dy^2], \quad \tilde{g} = \tilde{f}(x)[dx^2 + dy^2].$$

We shall add the subscript $\tilde{\cdot}$ to all the quantities referring to (\mathcal{S}, \tilde{g}) . Let the frequency λ^2 be fixed and let Γ_D, Γ_N be subsets belonging to distinct connected components of $\partial\mathcal{S}$. Then

- (1) If $\lambda^2 = 0$ and $\Lambda_{g,\Gamma_D,\Gamma_N}(0) = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}(0)$, there exists a function $c > 0$ with $c(0) = 1$ if $\Gamma_D \subset \{1\} \times T^1$, and $\Gamma_N \subset \{0\} \times T^1$, (resp. $c(1) = 1$ if $\Gamma_D \subset \{0\} \times T^1$ and $\Gamma_N \subset \{1\} \times T^1$) such that

$$\tilde{g} = cg.$$

- (2) If $\lambda^2 \neq 0$, there exists an explicit infinite dimensional family of metrics $\tilde{g} = cg$ with $c > 0$ and $c(0) = c(1) = 1$ that satisfies

$$\Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}(\lambda^2).$$

Proof. — Assume for instance that $\Gamma_D \subset \{1\} \times T^1$ and $\Gamma_N \subset \{0\} \times T^1$. We follow the same strategy as in the previous section and we use the same notation. We assume that $0 \in \Gamma_D$ and $\pm\pi \in \Gamma_N$ where T^1 is identified with $[-\pi, \pi]$. Thus, for all $\Psi \in C_0^\infty(-R, R)$, R small enough, ($R < \pi$), and for all $y \in \Gamma_N$,

$$(3.41) \quad \sum_{m \in \mathbb{Z}} \frac{1}{\sqrt{f(0)}} \frac{1}{\Delta(m^2)} \hat{\Psi}(m) e^{imy} = \sum_{m \in \mathbb{Z}} \frac{1}{\sqrt{\tilde{f}(0)}} \frac{1}{\tilde{\Delta}(m^2)} \hat{\Psi}(m) e^{imy}.$$

Then, repeating exactly the argument of the previous section, we see that

$$(3.42) \quad f(0) = \tilde{f}(0),$$

and

$$(3.43) \quad \alpha_k^2 = \tilde{\alpha}_k^2, \quad \forall k \geq 1,$$

where $(\alpha_k^2)_{k \geq 1}$ and $(\tilde{\alpha}_k^2)_{k \geq 1}$ are the zeros of $\Delta(\mu^2)$ and $\tilde{\Delta}(\mu^2)$.

Obviously, in the case $\lambda^2 = 0$, since the characteristic functions $\Delta(\mu^2)$ and $\tilde{\Delta}(\mu^2)$ do not depend on the metrics (see (3.13)), the unique condition we can get from the hypothesis $\Lambda_{g,\Gamma_D,\Gamma_N} = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}$ is $f(0) = \tilde{f}(0)$. Hence, we have

$$\tilde{g} = cg, \quad c(x) = \frac{\tilde{f}(x)}{f(x)},$$

with

$$c(0) = 1.$$

This proves the assertion (1).

Now, let us examine the case $\lambda^2 \neq 0$. First, we make the following remark: if the conditions (3.42) and (3.43) are satisfied, we can prove easily that $\Lambda_{g,\Gamma_D,\Gamma_N} = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}$. Indeed, from the proof of Proposition 3.5 and (3.43), we deduce that

$$(3.44) \quad \Delta(\mu^2) = \tilde{\Delta}(\mu^2), \forall \mu \in \mathbb{C}.$$

It follows that, under the conditions (3.42) and (3.43), we have $T_R^m(\lambda^2) = \tilde{T}_R^m(\lambda^2)$, for all $m \in \mathbb{Z}$, or equivalently $T_R(\lambda^2) = \tilde{T}_R(\lambda^2)$ which is a stronger

condition than $\Lambda_{g,\Gamma_D,\Gamma_N} = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}$. To summarize, this shows that the hypothesis $\Lambda_{g,\Gamma_D,\Gamma_N} = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}$ is equivalent to both conditions (3.42) and (3.43).

We would like to emphasize now that the condition (3.43) is nothing but a condition of isospectrality of the potentials q_λ and \tilde{q}_λ appearing in equation (3.8). Indeed, the sequences $(\alpha_k^2)_{k \geq 1}$ and $(\tilde{\alpha}_k^2)_{k \geq 1}$ correspond to “minus” the Dirichlet spectra of the Schrödinger operators $-\frac{d^2}{dx^2} + q_\lambda$ and $-\frac{d^2}{dx^2} + \tilde{q}_\lambda$. Hence, condition (3.43) means exactly that the Dirichlet spectra of $-\frac{d^2}{dx^2} + q_\lambda$ and $-\frac{d^2}{dx^2} + \tilde{q}_\lambda$ coincide, (we recall that the eigenvalues of these Schrödinger operators are always simple, (see for instance [37, Theorem 2, p. 30] or [46, Theorem 4.3.1])). In other words, the potentials q_λ and \tilde{q}_λ are isospectral.

Hence, the hypothesis $\Lambda_{g,\Gamma_D,\Gamma_N} = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}$ is equivalent to both conditions (3.42) and $q_\lambda, \tilde{q}_\lambda$ are isospectral.

It turns out that the isospectral class in $L^2([0, 1])$ associated to a given potential q and equation (3.8) has been the subject of intensive studies. We refer to the beautiful book [37] for a clear presentation of the results concerning this problem. For instance, the isospectral class in $L^2([0, 1])$ of a given potential q has been shown to be a real analytic submanifold of $L^2([0, 1])$ lying in the hyperplane of all functions with mean $[q] = \int_0^1 q(s)ds$ (see [37, Theorem 4.1, p. 69]).

Even better, we have the following explicit characterization of the isospectral class of a given potential $q \in L^2([0, 1])$.

THEOREM 3.7 ([37, Theorem 5.2, p. 102]). — *Let $q \in L^2([0, 1])$. Denote by $(v_k)_{k \geq 1} = (v_k(x, q))_{k \geq 1}$ the normalized eigenfunctions of the boundary value problem (3.8)–(3.9) with the potential q . The eigenfunctions $(v_k)_{k \geq 1}$ are associated to the eigenvalues $(\alpha_k^2)_{k \geq 1}$.*

Let $\xi \in l_1^2 = \{\xi = (\xi_1, \xi_2, \dots), \sum_{k \geq 1} k^2 |\xi_k|^2 < \infty\}$. Define the infinite matrix $\Theta(x, \xi, q) = (\theta_{ij}(x, \xi, q))_{i,j \geq 1}$ by

$$\theta_{ij}(x, \xi, q) = \delta_{ij} + (e^{\xi_i} - 1) \int_x^1 v_i(s)v_j(s)ds.$$

Define also the determinant of Θ as the limit of the determinants of its principal minors, i.e.

$$\det \Theta = \lim_{n \rightarrow \infty} \det \Theta^n, \quad \Theta^n = (\theta_{ij})_{1 \leq i,j \leq n}.$$

Then the isospectral class of q is completely and explicitly described by the family of potentials

$$q_\xi(x) = q(x) - 2 \frac{d^2}{dx^2} \log \det \Theta(x, \xi, q),$$

where $\xi \in l_1^2$. It is implicit in the statement of this result that the determinant of Θ always exists and never vanishes.

In particular, to each sequence $\xi^k \in l_1^2$ defined by $\xi^k = (\xi_j^k)_{j \geq 1}$ with $\xi_j^k = t \delta_{kj}$, $t \in \mathbb{R}$, we can find a one parameter family of explicit isospectral potentials to $q \in L^2([0, 1])$ by

$$(3.45) \quad q_{k,t}(x) = q(x) - 2 \frac{d^2}{dx^2} \log \theta_{k,t}(x), \quad \forall t \in \mathbb{R},$$

where

$$(3.46) \quad \theta_{k,t}(x) = 1 + (e^t - 1) \int_x^1 v_k^2(s) ds.$$

Let us make a few remarks on the family of potentials $(q_{k,t})_{k \geq 1, t \in \mathbb{R}}$ under the extra assumption that the potential q is smooth on $[0, 1]$.

- (1) The potentials $q_{k,t}$ are smooth on $[0, 1]$ for all $k \geq 1$ and for all $t \in \mathbb{R}$. Indeed, the eigenfunctions $v_k(x, q)$ are smooth on $[0, 1]$ by elliptic regularity. Hence, the functions $\theta_{k,t}$ are also smooth and never vanish on $[0, 1]$ for all $k \geq 1$ and for all $t \in \mathbb{R}$ by (3.46). This proves the smoothness of $q_{k,t}$ thanks to (3.45).
- (2) For all $k \geq 1$ and for all $t \in \mathbb{R}$, $q_{k,t}(0) = q(0)$ and $q_{k,t}(1) = q(1)$. Indeed, a short calculation shows that $2 \frac{d^2}{dx^2} \log \theta_{k,t}(x)$ is equal to

$$(3.47) \quad -2 \frac{(e^t - 1) \left[1 + (e^t - 1) \int_x^1 v_k^2(s) ds \right] v_k'(x) v_k(x) + (e^t - 1)^2 v_k^4(x)}{\left[1 + (e^t - 1) \int_x^1 v_k^2(s) ds \right]^2}.$$

Hence, since $v_k(0) = 0$ and $v_k(1) = 0$ by definition, we get the result using (3.45).

- (3) If moreover $q > 0$ (resp. $q < 0$), then for all $k \geq 1$, there exists $T_k > 0$ such that $q_{k,t} > 0$ (resp. $q_{k,t} < 0$) for all $-T_k < t < T_k$. Indeed, from (3.47), it is clear that for a fixed $k \geq 1$, the function $2 \frac{d^2}{dx^2} \log \theta_{k,t}(x)$ can be made arbitrarily small as $t \rightarrow 0$ uniformly w.r.t. $x \in [0, 1]$. Whence the result thanks to (3.45).

We now come back to our initial problem, that is given a frequency $\lambda^2 \neq 0$ and a smooth positive function $f(x)$, find all the smooth positive functions \tilde{f} such that (3.42) and (3.43) are satisfied. Define the smooth potential $q_\lambda = -\lambda^2 f$ as in (3.8). The potential q_λ is either positive or

negative according to the sign of λ^2 . Then, as discussed above, (3.43) is equivalent to finding the smooth positive or negative potentials $\tilde{q}_\lambda = -\lambda^2 \tilde{f}$ that are isospectral to q_λ . Thus in order to prove our result and according to Theorem 3.7, it suffices to show that the intersection of the isospectral class of q_λ in $L^2([0, 1])$ with the set of smooth positive or negative potentials satisfying (3.42) is infinite dimensional.

But using the three remarks above, we see that for all $k \geq 1$, there exists $T_k > 0$ such that the potentials $q_{\lambda,k,t}$ given by (3.45)–(3.46) with q replaced by q_λ are smooth positive or negative (according to the sign of λ^2) on $[0, 1]$ for all $-T_k < t < T_k$ and satisfy (3.42). We thus conclude that the infinite dimensional family of metrics (3.1) parametrized by the positive functions

$$(3.48) \quad f_{\lambda,k,t}(x) = f(x) + \frac{2}{\lambda^2} \frac{d^2}{dx^2} \log \theta_{k,t}(x), \quad \forall k \geq 1, \quad -T_k < t < T_k,$$

with

$$(3.49) \quad \theta_{k,t}(x) = 1 + (e^t - 1) \int_x^1 v_k^2(s) ds,$$

where v_k is the normalized eigenfunction of (3.8)–(3.9) associated to the eigenvalues α_k^2 , has the same partial DN map $\Lambda_{g,\Gamma_D,\Gamma_N}$ as the metric (3.1) associated to f . This finishes the proof of the theorem. \square

3.4. The 2D anisotropic Calderón with a potential

In this Section, we treat the anisotropic Calderón problem (Q3) with a potential $V \in L^\infty(\mathcal{S})$ such that $V = V(x)$ for our family of metrics (3.1). The global DN map $\Lambda_{g,V}(\lambda^2)$ associated to the Dirichlet problem

$$(3.50) \quad \begin{cases} (-\Delta_g + V)u = \lambda^2 u, & \text{on } \mathcal{S}, \\ u = \psi, & \text{on } \partial\mathcal{S}, \end{cases}$$

with λ^2 not belonging to the Dirichlet spectrum of $-\Delta_g + V$ can be constructed in the same way as in Section 3 since $V = V(x)$ respects the symmetry of (\mathcal{S}, g) . On each Fourier modes $Y_m = e^{imy}$, we get the following expression for the induced DN map:

$$(3.51) \quad \begin{aligned} \Lambda_{g,V}^m(\lambda^2) &= \begin{pmatrix} L_V^m(\lambda^2) & T_{R,V}^m(\lambda^2) \\ T_{L,V}^m(\lambda^2) & R_V^m(\lambda^2) \end{pmatrix} \\ &= \begin{pmatrix} -\frac{1}{\sqrt{f(0)}} M_V(m^2) & -\frac{1}{\sqrt{f(0)}} \frac{1}{\Delta_V(m^2)} \\ -\frac{1}{\sqrt{f(1)}} \frac{1}{\Delta_V(m^2)} & -\frac{1}{\sqrt{f(1)}} N_V(m^2) \end{pmatrix}, \end{aligned}$$

where the characteristic and Weyl–Titchmarsh functions $\Delta_V(m^2)$, $M_V(m^2)$ and $N_V(m^2)$ defined by (3.11)–(3.12) are associated to the radial ODE

$$(3.52) \quad -v'' + q_{\lambda,V}(x)v = -\mu^2 v, \quad q_{\lambda,V} = (V - \lambda^2)f,$$

with boundary conditions

$$(3.53) \quad v(0) = 0, \quad v(1) = 0.$$

We also recall the dictionary between the above coefficients of the DN map and the notations used in the Introduction

$$\begin{aligned} L_V(\lambda^2) &= \Lambda_{g,V,\Gamma_0}(\lambda^2), & R_V(\lambda^2) &= \Lambda_{g,V,\Gamma_1}(\lambda^2), \\ T_{L,V}(\lambda^2) &= \Lambda_{g,V,\Gamma_0,\Gamma_1}(\lambda^2), & T_{R,V}(\lambda^2) &= \Lambda_{g,V,\Gamma_1,\Gamma_0}(\lambda^2), \end{aligned}$$

where $\Gamma_0 = \{0\} \times T^1$ and $\Gamma_1 = \{1\} \times T^1$.

We prove

THEOREM 3.8. — *Let (\mathcal{S}, g) a Riemannian surface of the form (3.1), i.e.*

$$g = f(x)[dx^2 + dy^2].$$

Let $V, \tilde{V} \in L^\infty(\mathcal{S})$ be two potentials that only depend on the variable x . Let the frequency λ^2 be fixed and not belonging to the Dirichlet spectra of $-\Delta_g + V$ and $-\Delta_g + \tilde{V}$. Let Γ_D, Γ_N be nonempty open subsets belonging to $\partial\mathcal{S}$. Then

(1) *If Γ_D, Γ_N belong to the same connected component of $\partial\mathcal{S}$ and*

$$\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,\tilde{V},\Gamma_D,\Gamma_N}(\lambda^2),$$

then

$$\tilde{V} = V.$$

(2) *If Γ_D, Γ_N belong to distinct connected components of $\partial\mathcal{S}$, then there exists an explicit infinite dimensional family of potentials \tilde{V} that satisfies*

$$\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,\tilde{V},\Gamma_D,\Gamma_N}(\lambda^2).$$

Proof.

(1). — Assume for instance that $\Gamma_D, \Gamma_N \subset \Gamma_0$ and

$$\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,\tilde{V},\Gamma_D,\Gamma_N}(\lambda^2).$$

Then the same proof as in Theorem 3.2 shows first that

$$(3.54) \quad M_V(\mu^2) - M_V(0) = M_{\tilde{V}}(\mu^2) - M_{\tilde{V}}(0), \quad \forall \mu \in \mathbb{C} \setminus \mathbb{R}.$$

Hence the Borg–Marchenko Theorem 2.4 entails

$$(3.55) \quad q_{\lambda,V} = q_{\lambda,\tilde{V}}, \quad \text{on } [0, 1].$$

We conclude using (3.55) and (3.52) that $V = \tilde{V}$ on $[0, 1]$.

(2). — Assume that Γ_D and Γ_N belong to distinct connected components of $\partial\mathcal{S}$ and that

$$\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,\tilde{V},\Gamma_D,\Gamma_N}(\lambda^2).$$

Then the same proof as in Theorem 3.6 shows first that our hypothesis is equivalent to the unique condition

$$(3.56) \quad \Delta_V(\mu^2) = \Delta_{\tilde{V}}(\mu^2), \quad \forall \mu \in \mathbb{C}.$$

But, as explained in the proof of Theorem 3.6, this condition is in turn equivalent to the fact that the potentials $q_{\lambda,V}$ and $q_{\lambda,\tilde{V}}$ are isospectral. We deduce thus from [37] that given a potential $V \in L^\infty(\mathcal{S})$ as above, there exists an infinite dimensional family of explicit potentials \tilde{V} satisfying (3.56). More precisely, the family

$$(3.57) \quad \tilde{V}_{\lambda,k,t}(x) = V(x) - \frac{2}{f(x)} \frac{d^2}{dx^2} \log \theta_{k,t}(x), \quad \forall k \geq 1, \quad t \in \mathbb{R},$$

with

$$(3.58) \quad \theta_{k,t}(x) = 1 + (e^t - 1) \int_x^1 v_k^2(s) ds,$$

where v_k is the normalized eigenfunction of (3.52)–(3.53) associated to the eigenvalues α_k^2 satisfies

$$\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,\tilde{V}_{\lambda,n,t},\Gamma_D,\Gamma_N}(\lambda^2).$$

Note here that there is no restriction in the range of the parameter $t \in \mathbb{R}$ since we work in the class of potentials $V \in L^\infty([0, 1])$. In particular, we don't need \tilde{V} to be smooth anylonger. □

4. The three dimensional case

4.1. The model and the Dirichlet-to-Neumann map

In this Section, we work on a Riemannian manifold (M, g) which has the topology of a cylinder $M = [0, 1] \times T^2$ and that is equipped with a Riemannian metric

$$(4.1) \quad g = f(x)dx^2 + f(x)dy^2 + h(x)dz^2,$$

where f, h are smooth positive functions on M . Here (x, y, z) is a global coordinates system on M and T^2 stands for the two dimensional torus. The metric g obviously possesses two commuting Killing vector fields ∂_y

and ∂_z that generate a two-parameter Abelian group of isometries on M . These isometries will be referred to as the toroidal symmetries of M . The boundary ∂M of M is disconnected and consists in the disjoint union of two copies of T^2 , precisely

$$\partial M = \Gamma_0 \cup \Gamma_1, \quad \Gamma_0 = \{0\} \times T^2, \quad \Gamma_1 = \{1\} \times T^2.$$

In our global coordinate system (x, y, z) , the positive Laplace–Beltrami operator has the expression

$$-\Delta_g = \frac{1}{f} \left(-\partial_x^2 - \frac{1}{2}(\log h)' \partial_x - \partial_y^2 - \frac{f}{h} \partial_z^2 \right).$$

We look at the Dirichlet problem at a frequency λ^2 on M such that $\lambda^2 \notin \{\lambda_j^2\}_{j \geq 1}$ where $\{\lambda_j^2\}_{j \geq 1}$ is the Dirichlet spectrum of $-\Delta_g$. We consider thus the solutions u of

$$(4.2) \quad -\partial_x^2 u - \frac{1}{2}(\log h)' \partial_x u - \partial_y^2 u - \frac{f}{h} \partial_z^2 u - \lambda^2 f u = 0, \quad \text{on } M,$$

together with the boundary condition

$$(4.3) \quad u = \psi \quad \text{on } \partial M.$$

In order to construct the DN map, we shall use the notations introduced in Section 3. Recall that the boundary ∂M of M has two disjoint components $\partial M = \Gamma_0 \cup \Gamma_1$ where $\Gamma_0 \simeq \Gamma_1 \simeq T^2$. Hence, we can decompose the Sobolev spaces $H^s(\partial M)$ as $H^s(\partial M) = H^s(\Gamma_0) \oplus H^s(\Gamma_1)$ for any $s \in \mathbb{R}$ and we shall use the vector notation

$$\varphi = (\varphi^0 \varphi^1),$$

for all elements φ of $H^s(\partial M) = H^s(\Gamma_0) \oplus H^s(\Gamma_1)$. The DN map is a linear operator from $H^{1/2}(\partial M)$ to $H^{-1/2}(\partial M)$ and thus has the structure of an operator valued 2×2 matrix

$$\Lambda_g(\lambda^2) = \begin{pmatrix} L(\lambda^2) & T_R(\lambda^2) T_L(\lambda^2) \\ R(\lambda^2) & \end{pmatrix},$$

where $L(\lambda^2), R(\lambda^2), T_R(\lambda^2), T_L(\lambda^2)$ are bounded operators from $H^{1/2}(T^2)$ to $H^{-1/2}(T^2)$. Finally, we recall the dictionary established in Section 3

$$\begin{aligned} L(\lambda^2) &= \Lambda_{g, \Gamma_0}(\lambda^2), & R(\lambda^2) &= \Lambda_{g, \Gamma_1}(\lambda^2), \\ T_L(\lambda^2) &= \Lambda_{g, \Gamma_0, \Gamma_1}(\lambda^2), & T_R(\lambda^2) &= \Lambda_{g, \Gamma_1, \Gamma_0}(\lambda^2), \end{aligned}$$

which makes the link with the notations used in the Introduction.

Now we use the toroidal symmetry of (M, g) to find a simple expression of the DN map. We first write $\psi = (\psi^0, \psi^1) \in H^{1/2}(\Gamma_0) \times H^{1/2}(\Gamma_1)$ using

their Fourier series as

$$\psi^0 = \sum_{m,n \in \mathbb{Z}} \psi_{mn}^0 Y_{mn}, \quad \psi^1 = \sum_{m,n \in \mathbb{Z}} \psi_{mn}^1 Y_{mn},$$

where

$$Y_{mn}(y, z) = e^{imy} e^{inz}.$$

Note that for any $s \in \mathbb{R}$, the space $H^s(T^2)$ can be described as

$$\left\{ \varphi \in \mathcal{D}'(T^2), \varphi = \sum_{m,n \in \mathbb{Z}} \varphi_{mn} Y_{mn}, \quad \sum_{m,n \in \mathbb{Z}} (1 + m^2 + n^2)^s |\varphi_{mn}|^2 < \infty \right\}.$$

Next, the unique solution u of (4.2)–(4.3) can be looked for in the form

$$u = \sum_{m,n \in \mathbb{Z}} u_{mn}(x) Y_{mn}(y, z),$$

and for all $m, n \in \mathbb{Z}$, the function u_{mn} is the unique solution of the ODE (w.r.t. x) with boundary conditions

$$(4.4) \quad \begin{cases} -u''_{mn} - \frac{1}{2}(\log h)'u'_{mn} + m^2 u_{mn} + n^2 \frac{f}{h} u_{mn} - \lambda^2 f u_{mn} = 0, \\ u_{mn}(0) = \psi_{mn}^0, \quad u_{mn}(1) = \psi_{mn}^1. \end{cases}$$

For later use, we get rid of the term of order 1 in (4.4) by introducing the new functions

$$(4.5) \quad v_{mn} = (h)^{1/4} u_{mn},$$

which then satisfy

$$(4.6) \quad \begin{cases} -v''_{mn} + \frac{[(\log h)']^2}{16} v_{mn} + \frac{(\log h)''}{4} v_{mn} + m^2 v_{mn} + n^2 \frac{f}{h} v_{mn} - \lambda^2 f v_{mn} = 0, \\ v_{mn}(0) = (h(0))^{1/4} \psi_{mn}^0, \quad v_{mn}(1) = (h(1))^{1/4} \psi_{mn}^1. \end{cases}$$

The DN map is now diagonalized on the Hilbert basis $\{Y_{mn}\}_{m,n \in \mathbb{Z}}$ and is shown to have a very simple expression on each harmonic. First, a short calculation using the particular form of the metric (4.1) shows that for smooth enough boundary data ψ

$$\Lambda_g(\lambda^2) \begin{pmatrix} \psi^0 \\ \psi^1 \end{pmatrix} = \begin{pmatrix} (\partial_\nu u)|_{\Gamma_0} \\ (\partial_\nu u)|_{\Gamma_1} \end{pmatrix} = \begin{pmatrix} -\frac{1}{\sqrt{f(0)}}(\partial_x u)|_{x=0} \\ \frac{1}{\sqrt{f(1)}}(\partial_x u)|_{x=1} \end{pmatrix}.$$

Hence, if we make the DN map act on the vector space generated by the harmonic Y_{mn} , we get

$$(4.7) \quad \Lambda_g(\lambda^2) \begin{pmatrix} \psi_{mn}^0 Y_{mn} \\ \psi_{mn}^1 Y_{mn} \end{pmatrix} = \begin{pmatrix} -\frac{1}{\sqrt{f(0)}} u'_{mn}(0) Y_{mn} \\ \frac{1}{\sqrt{f(1)}} u'_{mn}(1) Y_{mn} \end{pmatrix}.$$

We denote

$$\Lambda_g(\lambda^2)|_{\langle Y_{mn} \rangle} = \Lambda_g^{mn}(\lambda^2) = \begin{pmatrix} L^{mn}(\lambda^2) & T_R^{mn}(\lambda^2) \\ T_L^{mn}(\lambda^2) & R^{mn}(\lambda^2) \end{pmatrix},$$

the restriction of the global DN map to each harmonic $\langle Y_{mn} \rangle$. This operator has the structure of a 2×2 matrix and satisfies for all $m, n \in \mathbb{Z}$

$$(4.8) \quad \begin{pmatrix} L^{mn}(\lambda^2) & T_R^{mn}(\lambda^2) \\ T_L^{mn}(\lambda^2) & R^{mn}(\lambda^2) \end{pmatrix} \begin{pmatrix} \psi_{mn}^0 \\ \psi_{mn}^1 \end{pmatrix} = \begin{pmatrix} -\frac{1}{\sqrt{f(0)}}u'_{mn}(0) \\ \frac{1}{\sqrt{f(1)}}u'_{mn}(1) \end{pmatrix}.$$

Using the change of functions (4.5), we get the equivalent expression for the action of the partial DN maps $\Lambda_g^{mn}(\lambda^2)$ on vectors $(\psi_{mn}^0, \psi_{mn}^1) \in \mathbb{C}^2$

$$(4.9) \quad \Lambda_g^{mn}(\lambda^2) \begin{pmatrix} \psi_{mn}^0 \\ \psi_{mn}^1 \end{pmatrix} = \begin{pmatrix} -\frac{1}{\sqrt{f(0)}} \left(-\frac{h'(0)}{4h^{5/4}(0)}v_{mn}(0) + \frac{1}{h^{1/4}(0)}v'_{mn}(0) \right) \\ \frac{1}{\sqrt{f(1)}} \left(-\frac{h'(1)}{4h^{5/4}(1)}v_{mn}(1) + \frac{1}{h^{1/4}(1)}v'_{mn}(1) \right) \end{pmatrix}.$$

As in Section 3, we can further simplify the partial DN maps $\Lambda_g^{mn}(\lambda^2)$ by interpreting their coefficients as the characteristic and Weyl–Titchmarsh functions of the ODE (4.6) with appropriate boundary conditions. Here is the procedure. First fix $n \in \mathbb{Z}$ and consider the ODE

$$(4.10) \quad -v'' + q_{\lambda n}(x)v = -\mu^2v, \quad q_{\lambda n} = \frac{[(\log h)']^2}{16} + \frac{(\log h)''}{4} + n^2 \frac{f}{h} - \lambda^2 f,$$

with boundary conditions

$$(4.11) \quad v(0) = 0, \quad v(1) = 0.$$

Note that the equation (4.10) is nothing but equation (4.6) in which the parameter $-m^2$ is written as $-\mu^2$ and is interpreted as the spectral parameter of the equation. On the other hand, the boundary conditions (4.3) have been replaced by Dirichlet boundary conditions at $x = 0$ and $x = 1$. Since the potential $q_{\lambda n} \in L^1([0, 1])$ and is real, we can define for all $n \in \mathbb{Z}$ and all $\mu \in \mathbb{C}$ the fundamental systems of solutions

$$\{c_0(x, \mu^2, n^2), s_0(x, \mu^2, n^2)\}, \quad \{c_1(x, \mu^2, n^2), s_1(x, \mu^2, n^2)\},$$

of (4.10) by imposing the Cauchy conditions

$$(4.12) \quad \begin{aligned} c_0(0, \mu^2, n^2) &= 1, & c'_0(0, \mu^2, n^2) &= 0, & s_0(0, \mu^2, n^2) &= 0, & s'_0(0, \mu^2, n^2) &= 1, \\ c_1(1, \mu^2, n^2) &= 1, & c'_1(1, \mu^2, n^2) &= 0, & s_1(1, \mu^2, n^2) &= 0, & s'_1(1, \mu^2, n^2) &= 1. \end{aligned}$$

Note that the dependence of the FSS on λ^2 is not written for clarity but implicit. Also, it is clear that (2.4) is satisfied and that the solutions $\{c_0(x, \mu^2, n^2), s_0(x, \mu^2, n^2)\}$ and $\{c_1(x, \mu^2, n^2), s_1(x, \mu^2, n^2)\}$ are entire functions with respect to the variable $\mu^2 \in \mathbb{C}$.

Following Section 2 we thus define the characteristic function of (4.10) with boundary conditions (4.11) by

$$(4.13) \quad \Delta(\mu^2, n^2) = W(s_0, s_1),$$

while we define the Weyl–Titchmarsh functions by

$$(4.14) \quad \begin{aligned} M(\mu^2, n^2) &= -\frac{W(c_0, s_1)}{\Delta(\mu^2, n^2)} = -\frac{D(\mu^2, n^2)}{\Delta(\mu^2, n^2)}, \\ N(\mu^2, n^2) &= -\frac{W(c_1, s_0)}{\Delta(\mu^2, n^2)} = -\frac{E(\mu^2, n^2)}{\Delta(\mu^2, n^2)}. \end{aligned}$$

Remark 4.1. — For all $n \in \mathbb{Z}$, the WT function $\mu^2 \mapsto M(\mu^2, n^2)$ is meromorphic on \mathbb{C} and has an infinite and discrete set of poles $\{\alpha_{n,j}^2\}_{j \geq 1}$ corresponding to “minus” the Dirichlet eigenvalues of $-\frac{d}{dx^2} + q_{\lambda n}$ or equivalently, corresponding to the zeros of the characteristic function $\mu^2 \mapsto \Delta(\mu^2, n^2)$. Let us show that the integers $m^2, m \in \mathbb{Z}$ cannot be poles of $\mu^2 \mapsto M(\mu^2, n^2)$ under our general assumption. Assume the converse, i.e. there exists $m \in \mathbb{Z}$ such that m^2 is a pole of $M(\mu^2, n^2)$. Thus $-m^2$ is a Dirichlet eigenvalue of $-\frac{d}{dx^2} + q_{\lambda n}$ and we denote by $u_{mn}(x)$ the associated eigenfunction solution of (4.6). Then the function $u(x, y) = u_{mn}(x)Y_{mn}(y, z)$ is a nontrivial solution of (4.2) with Dirichlet boundary conditions. We conclude that such a u is an eigenfunction of $-\Delta_g$ associated to the Dirichlet eigenvalue λ^2 . But this case has been ruled out from the beginning since we assume that λ^2 is not a Dirichlet eigenvalue of Δ_g . Whence the contradiction.

We now come back to the expression (4.9) of the partial DN map $\Lambda_g^{mn}(\lambda^2)$. For all $m, n \in \mathbb{Z}$, we need to express $v'_{mn}(0)$ and $v'_{mn}(1)$ in terms of ψ_{mn}^0 and ψ_{mn}^1 in order to find the expressions of the coefficients $L^{mn}(\lambda^2)$, $T_R^{mn}(\lambda^2)$, $T_L^{mn}(\lambda^2)$ and $R^{mn}(\lambda^2)$. But the solution v_{mn} of (4.6) can be written as linear combinations of the FSS

$$\{c_0(x, m^2, n^2), s_0(x, m^2, n^2)\}, \quad \{c_1(x, m^2, n^2), s_1(x, m^2, n^2)\}.$$

Precisely, we write

$$\begin{aligned} v_{mn}(x) &= \alpha c_0(x, m^2, n^2) + \beta s_0(x, m^2, n^2) \\ &= \gamma c_1(x, m^2, n^2) + \delta s_1(x, m^2, n^2), \end{aligned}$$

for some constants $\alpha, \beta, \gamma, \delta$. Using similar calculations as in Section 3, we get for $\Lambda_g^{mn}(\lambda^2)$ the expression

$$(4.15) \quad \Lambda_g^{mn}(\lambda^2) = \begin{pmatrix} \frac{(\log h)'(0)}{4\sqrt{f(0)}} - \frac{M(m^2, n^2)}{\sqrt{f(0)}} & -\frac{h^{1/4}(1)}{h^{1/4}(0)\sqrt{f(0)\Delta(m^2, n^2)}} \\ -\frac{h^{1/4}(0)}{h^{1/4}(1)\sqrt{f(1)\Delta(m^2, n^2)}} & -\frac{(\log h)'(1)}{4\sqrt{f(1)}} - \frac{\Delta(m^2, n^2)}{\sqrt{f(1)}} \end{pmatrix}.$$

Remark 4.2.

- (1) For all $m, n \in \mathbb{Z}$, the characteristic and WT functions $\Delta(m^2, n^2)$ and $M(m^2, n^2)$ are well defined thanks to our assumption stating that λ^2 is not a Dirichlet eigenvalue of Δ_g (see Remark 4.1 for this point).
- (2) There is of course no importance in the choice of the parameters m or n to be fixed. If $m \in \mathbb{Z}$ was to be fixed, we could define the same objects with respect to the parameter $\nu^2 \in \mathbb{C}$ which would replace the parameter n^2 in (4.10).

The result above shows that the coefficients of the partial DN map $\Lambda_g^{mn}(\lambda^2)$ can be simply interpreted in terms of the characteristic and Weyl–Titchmarsh functions associated to the ODE (4.10)–(4.11). In particular, the partial DN maps $L(\lambda^2)$ and $R(\lambda^2)$ which correspond to the global DN map restricted to Γ_0 and Γ_1 respectively, only depend on the Weyl–Titchmarsh functions $M(m^2, n^2)$ (modulo some explicit constants). In fact, in our model, the knowledge of $L(\lambda^2)$ and $R(\lambda^2)$ is clearly equivalent to the knowledge of the sequence of Weyl–Titchmarsh functions $\{M(m^2, n^2)\}_{m, n \in \mathbb{Z}}$ (modulo some constants).

Similarly, the knowledge of the partial DN maps $T_L(\lambda^2)$ and $T_R(\lambda^2)$ which correspond to the global DN map whose data are measured on the disjoint sets Γ_0 and Γ_1 , is equivalent to the knowledge of the characteristic functions $\Delta(m^2, n^2)$ (modulo some explicit constants).

We finish this Section with a simple result that is the key point of our non-uniqueness results for the anisotropic Calderón problem with data measured on disjoint sets.

LEMMA 4.3. — *Consider two potentials q_{λ_n} and \tilde{q}_{λ_n} of the form (4.10) associated to two metrics g and \tilde{g} of the form (4.1). Assume that*

$$q_{\lambda_n} = \tilde{q}_{\lambda_n},$$

for at least two different $n \in \mathbb{Z}$. Then

$$(4.16) \quad \tilde{f}(x) = c^4(x)f(x), \quad \tilde{h}(x) = c^4(x)h(x),$$

where the positive function c satisfies the non-linear ODE

$$(4.17) \quad c'' + \frac{1}{2}(\log h)'c' + \lambda^2 f(c - c^5) = 0.$$

If moreover, $\lambda^2 = 0$, then we can solve (4.17) explicitly and get

$$(4.18) \quad c(x) = \left(A + B \int_0^x \frac{ds}{\sqrt{h(s)}} \right),$$

where (A, B) are any constants such that $c(x) > 0$ for all $x \in [0, 1]$.

Proof. — Assume that $q_{\lambda n} = \tilde{q}_{\lambda n}$ for two different $n \in \mathbb{Z}$. Then using the definition (4.10) for these two different values of n , we immediately get

$$(4.19) \quad \begin{cases} \frac{f}{h} = \frac{\tilde{f}}{\tilde{h}}, \\ \frac{(\log h)'^2}{16} + \frac{(\log h)''}{4} - \lambda^2 f = \frac{(\log \tilde{h})'^2}{16} + \frac{(\log \tilde{h})''}{4} - \lambda^2 \tilde{f}. \end{cases}$$

We set $c = \left(\frac{\tilde{h}}{h}\right)^{1/4}$. Hence a short calculation shows that c must satisfy (4.17). Moreover we get easily (4.16) from (4.19). Finally, when $\lambda^2 = 0$, the ODE (4.17) becomes a first order linear ODE in the unknown c' and its solution leads to (4.18). □

4.2. The 3D anisotropic Calderón problem with data measured on the same connected component

In this Section, we solve the anisotropic Calderón problem in the case where the Dirichlet and Neumann data are measured on non-empty open sets Γ_D and Γ_N belonging to the same connected component of ∂M . Precisely, we prove

THEOREM 4.4. — *Let (M, g) and (M, \tilde{g}) denote two Riemannian manifolds of the form (4.1), i.e.*

$$g = f(x)dx^2 + f(x)dy^2 + h(x)dz^2, \quad \tilde{g} = \tilde{f}(x)dx^2 + \tilde{f}(x)dy^2 + \tilde{h}(x)dz^2.$$

We shall add the subscript $\tilde{\cdot}$ to all the quantities referring to (M, \tilde{g}) . Let $\lambda^2 \in \mathbb{R}$. Let Γ_D and Γ_N be non-empty open sets belonging to the same connected component of ∂M . Assume that $\Gamma_D \cap \Gamma_N \neq \emptyset$, with Γ_N containing an annular region of the type

$$(y_0 - \delta, y_0 + \delta) \times T^1, \quad \text{or} \quad T^1 \times (z_0 - \delta, z_0 + \delta),$$

where $\delta > 0$. Assume moreover that $\Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}(\lambda^2)$. Then

$$\tilde{g} = g.$$

To prove this theorem, we shall use the Complex Angular Momentum (CAM) method with respect to the parameter m^2 , that is to say that we shall allow m^2 to be complex and use the beautiful analytic properties of the different objects related to the DN map (such as the characteristic function $\Delta(m^2, n^2)$ and the Weyl–Titchmarsh functions $M(m^2, n^2)$ and $N(m^2, n^2)$). In what follows, we shall use freely the results recalled in Section 2 on the functions $\Delta(\mu^2, n^2), D(\mu^2, n^2), E(\mu^2, n^2), M(\mu^2, n^2), \mathbb{N}(\mu^2, n^2)$ with $\mu \in \mathbb{C}$ and for fixed $n \in \mathbb{Z}$.

Proof. — We assume that $\Gamma_D, \Gamma_N \subset \Gamma_0 = \{0\} \times T^2$. The proof for $\Gamma_D, \Gamma_N \subset \Gamma_1 = \{1\} \times T^2$ is the same and we omit it. Assume also (without loss of generality) that Γ_N contains the annular open set

$$([-\pi, -\pi + \delta] \times T^1) \cup (\pi - \delta, \pi] \times T^1),$$

for a small $\delta > 0$ and that $\Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}(\lambda^2)$. According to (4.15) and the discussion after it, this assumption is equivalent to

(4.20)

$$\Phi(y, z) := \sum_{m,n} \left[A - \frac{M(m^2, n^2)}{\sqrt{f(0)}} - \tilde{A} + \frac{\tilde{M}(m^2, n^2)}{\sqrt{\tilde{f}(0)}} \right] \hat{\psi}(m, n) e^{imy+inz} = 0,$$

for all $(y, z) \in \Gamma_N$ and for all $\psi \in H^{1/2}(T^2)$ with $\text{supp } \psi \subset \Gamma_D$. Here $\hat{\psi}(m, n)$ denotes the Fourier coefficients of ψ and

$$A = \frac{(\log h)'(0)}{4\sqrt{f(0)}}, \quad \tilde{A} = \frac{(\log \tilde{h})'(0)}{4\sqrt{\tilde{f}(0)}}.$$

Let us assume from now on that ψ is smooth enough (say $\psi \in H^1(T^1)$) and that $\text{supp } \psi \subset \Gamma_D \cap [-r, r] \times T^1$ where $r < \pi$. This is always possible since Γ_D is open in T^2 and up to consider ψ with smaller support than Γ_D . Then we can extract several informations from (4.20).

First the functions Φ and ψ are in $L^2(T^2)$. Second, $\text{supp } \Phi \subset [-R, R] \times T^1$ with $R < \pi$ and $\text{supp } \psi \subset [-r, r] \times T^1$ where $r < \pi$. We thus conclude from the multivariable Paley–Wiener Theorem (see [18, Theorem 7.3.1, p. 181]) that the Fourier transforms of Φ and ψ are entire functions on \mathbb{C}^2 that satisfy the estimates

$$(4.21) \quad \begin{cases} c|\hat{\psi}(\mu, \nu)| \leq C e^{r|\Im(\mu)| + \pi|\Im(\nu)|}, \\ |\hat{\Phi}(\mu, \nu)| \leq C e^{R|\Im(\mu)| + \pi|\Im(\nu)|}, \end{cases} \quad \forall (\mu, \nu) \in \mathbb{C}^2.$$

Third, we can check directly that for all $m, n \in \mathbb{Z}$

$$(4.22) \quad \hat{\Phi}(m, n) = \left[A - \frac{M(m^2, n^2)}{\sqrt{f(0)}} - \tilde{A} + \frac{\tilde{M}(m^2, n^2)}{\sqrt{\tilde{f}(0)}} \right] \hat{\psi}(m, n).$$

Recalling the definition (4.14) of the Weyl–Titchmarsh functions M and \tilde{M} , this latter equality can be rewritten as

$$(4.23) \quad \Delta(m^2, n^2) \tilde{\Delta}(m^2, n^2) \hat{\Phi}(m, n) - \left[\Delta(m^2, n^2) \tilde{\Delta}(m^2, n^2) (A - \tilde{A}) - \frac{D(m^2, n^2) \tilde{\Delta}(m^2, n^2)}{\sqrt{f(0)}} + \frac{\tilde{D}(m^2, n^2) \Delta(m^2, n^2)}{\sqrt{\tilde{f}(0)}} \right] \hat{\psi}(m, n) = 0,$$

for all $m, n \in \mathbb{Z}$.

Let us fix $n \in \mathbb{Z}$. We denote by $F(\mu, n)$ the function in the left-hand-side of (4.23) where m^2 is replaced by μ . Clearly, F is entire of order 1 with respect to μ and satisfies the estimate on the imaginary axis:

$$|F(iy, n)| \leq C e^{\max(r, R)|y|}, \quad \forall y \in \mathbb{R},$$

thanks to Corollary 2.2 and (4.21). Moreover, F vanishes on the integers by (4.22). Since $\max(r, R) < \pi$, we deduce from Carlson’s Theorem (see [3, Theorem 9.2.1]) that $F(\mu, n) = 0$ for all $\mu \in \mathbb{C}, n \in \mathbb{Z}$. This can be written equivalently as

$$(4.24) \quad \hat{\Phi}(\mu, n) = \left[A - \frac{M(\mu^2, n^2)}{\sqrt{f(0)}} - \tilde{A} + \frac{\tilde{M}(\mu^2, n^2)}{\sqrt{\tilde{f}(0)}} \right] \hat{\psi}(\mu, n), \quad \forall \mu \in \mathbb{C}, n \in \mathbb{Z}.$$

Since the function $\hat{\Phi}(\mu, n)$ is entire w.r.t. μ , the function

$$\left[A - \frac{M(\mu^2, n^2)}{\sqrt{f(0)}} - \tilde{A} + \frac{\tilde{M}(\mu^2, n^2)}{\sqrt{\tilde{f}(0)}} \right] \hat{\psi}(\mu, n),$$

must also be entire in μ for all ψ smooth enough such that $\text{supp } \psi \subset \Gamma_D \cap [-r, r] \times T^1$ where $r < \pi$. In other words, the poles of $\frac{M(\mu^2, n^2)}{\sqrt{f(0)}}$ and $\frac{\tilde{M}(\mu^2, n^2)}{\sqrt{\tilde{f}(0)}}$ as well their residues must coincide. In fact, an easy adaption then of the argument given in the proof of Proposition 3.5 shows that (4.24)

implies

$$(4.25) \quad \Delta(\mu^2, n^2) = \tilde{\Delta}(\mu^2, n^2), \quad \forall \mu \in \mathbb{C}, n \in \mathbb{Z}.$$

$$(4.26) \quad M(\mu^2, n^2) - M(0, n^2) = \tilde{M}(\mu^2, n^2) - \tilde{M}(0, n^2), \quad \forall \mu \in \mathbb{C} \setminus \mathbb{R}, n \in \mathbb{Z},$$

and

$$(4.27) \quad f(0) = \tilde{f}(0).$$

We can now finish the proof as follows. From (4.26) and the Borg–Marchenko Theorem 2.4, we deduce first that

$$(4.28) \quad q_{\lambda n}(x) = \tilde{q}_{\lambda n}(x), \quad \forall x \in [0, 1], n \in \mathbb{Z},$$

which the main assumption of Lemma 4.3. In particular, we know that $\tilde{f} = c^4 f$ and $\tilde{h} = c^4 h$ where c is a function satisfying the ODE (4.17).

But the equality (4.28) in turn implies that $M(\mu^2, n^2) = \tilde{M}(\mu^2, n^2)$ for all $\mu \in \mathbb{C}, n \in \mathbb{Z}$. Putting this in (4.20), we get

$$(4.29) \quad (A - \tilde{A})\psi(y, z) = 0, \quad \forall (y, z) \in \Gamma_N,$$

and for all ψ such that $\text{supp } \psi \subset \Gamma_D$. Since $\Gamma_D \cap \Gamma_N \neq \emptyset$, we conclude from (4.29) that

$$(4.30) \quad A = \tilde{A}.$$

Together with (4.27), (4.30) implies that the function $c = \left(\frac{\tilde{h}}{h}\right)^{1/4}$ satisfies the Cauchy conditions

$$(4.31) \quad c(0) = 1, \quad c'(0) = 0.$$

Hence, we deduce from (4.31) that $c = 1$ is the unique solution of the ODE (4.17). In other words, we have proved that $\tilde{f} = f$ and $\tilde{h} = h$ or equivalently, $\tilde{g} = g$ which concludes the proof of the theorem. \square

We now state our first non-uniqueness result for three dimensional Riemannian manifolds.

THEOREM 4.5. — *Let (M, g) and (M, \tilde{g}) denote Riemannian manifolds of the form (4.1), i.e.*

$$g = f(x)dx^2 + f(x)dy^2 + h(x)dz^2, \quad \tilde{g} = \tilde{f}(x)dx^2 + \tilde{f}(x)dy^2 + \tilde{h}(x)dz^2.$$

Let $\lambda^2 \in \mathbb{R}$. Let Γ_D and Γ_N be non-empty open sets belonging to the same connected component of ∂M . Assume that $\Gamma_D \cap \Gamma_N = \emptyset$. Then there exists infinitely many pairs of non-isometric metrics (g, \tilde{g}) given by $\tilde{g} = c^4 g$ where c are smooth positive strictly increasing or decreasing functions such that $c(0) = 1$ if $\Gamma_D, \Gamma_N \subset \Gamma_0$ and $c(1) = 1$ if $\Gamma_D, \Gamma_N \subset \Gamma_1$, satisfying

$$\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(\lambda^2).$$

Proof. — Let us assume that $\Gamma_D, \Gamma_N \subset \Gamma_0 = \{0\} \times T^1$. We construct pairs of metrics (g, \tilde{g}) satisfying $\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(\lambda^2)$ as follows. Let f and c be any smooth positive function on $[0, 1]$ such that $c(0) = 1$ and $c'(x) \neq 0$ for all $x \in [0, 1]$. In other words, $c(x)$ is a strictly monotonic function. Define

$$(4.32) \quad h = Ce^{-2 \int_0^x \frac{c''(s) + \lambda^2 f(s)(c(s) - c^5(s))}{c'(s)} ds}, \quad \tilde{f} = c^4 f, \quad \tilde{h} = c^4 h.$$

Clearly, we have then $\tilde{g} = c^4 g$. Using Lemma 4.3, it is immediate to check that $q_{\lambda n} = \tilde{q}_{\lambda n}$ for all $n \in \mathbb{Z}$ where $q_{\lambda n}$ and $\tilde{q}_{\lambda n}$ are given by (4.10). In particular, for such choices of metrics (g, \tilde{g}) , we always have $M(m^2, n^2) = \tilde{M}(m^2, n^2)$ for all $m, n \in \mathbb{Z}$. Moreover, our assumption $c(0) = 1$ implies that $f(0) = \tilde{f}(0)$.

Now, using (4.15) and the discussion after it, we see that for all $\psi \in H^{1/2}(T^2)$ with $\text{supp } \psi \subset \Gamma_D$, we have

$$(4.33) \quad \begin{aligned} &\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2)(\psi) \\ &:= \left[\sum_{m, n} \left(\frac{(\log h)'(0)}{4\sqrt{f(0)}} - \frac{M(m^2, n^2)}{\sqrt{f(0)}} \right) \hat{\psi}(m, n) e^{imy+inz} \right]_{|(y, z) \in \Gamma_N} \\ &= \left[\frac{(\log h)'(0)}{4\sqrt{f(0)}} \psi(y, z) - \sum_{m, n} \frac{M(m^2, n^2)}{\sqrt{f(0)}} \hat{\psi}(m, n) e^{imy+inz} \right]_{|(y, z) \in \Gamma_N}. \end{aligned}$$

But since $\Gamma_D \cap \Gamma_N = \emptyset$, we thus deduce that

$$\frac{(\log h)'(0)}{4\sqrt{f(0)}} \psi(y, z) = 0,$$

for all $(y, z) \in \Gamma_N$. Hence we obtain

$$(4.34) \quad \begin{aligned} \Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2)(\psi) &= - \left[\sum_{m, n} \frac{M(m^2, n^2)}{\sqrt{f(0)}} \hat{\psi}(m, n) e^{imy+inz} \right]_{|(y, z) \in \Gamma_N} \\ &= - \left[\sum_{m, n} \frac{\tilde{M}(m^2, n^2)}{\sqrt{\tilde{f}(0)}} \hat{\psi}(m, n) e^{imy+inz} \right]_{|(y, z) \in \Gamma_N}, \\ &= \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(\lambda^2)(\psi), \end{aligned}$$

for all $\psi \in H^{1/2}(T^2)$ with $\text{supp } \psi \subset \Gamma_D$. This proves the result. □

Remark 4.6. — If we fix a Riemannian manifold (M, g) of the form (4.1), we don't know a priori whether there exists a metric \tilde{g} such that

$\Lambda_{g,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}(\lambda^2)$ with $\Gamma_D \cap \Gamma_N = \emptyset$ except if the original metric g has the form

$$g = f(x)dx^2 + f(y)dy^2 + h(x)dz^2,$$

where h given by (4.32).

In the case of zero frequency $\lambda^2 = 0$, we can do a little better since we can solve explicitly the ODE (4.17). Since this result is interesting in its own sake, we state it as a theorem

THEOREM 4.7. — *Let (M, g) denotes a Riemannian manifold of the form (4.1), i.e.*

$$g = f(x)dx^2 + f(y)dy^2 + h(x)dz^2.$$

Let Γ_D and Γ_N be non-empty open sets belonging to the same connected component of ∂M . Assume that $\Gamma_D \cap \Gamma_N = \emptyset$. Then

- (1) *if $\Gamma_D, \Gamma_N \subset \Gamma_0$, there exists a one parameter family of metrics \tilde{g} given by*

$$\tilde{g} = \left[1 + B \int_0^x \frac{ds}{\sqrt{h(s)}} \right]^4 g, \quad B > 0,$$

that satisfies $\Lambda_{g,\Gamma_D,\Gamma_N}(0) = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}(0)$.

- (2) *if $\Gamma_D, \Gamma_N \subset \Gamma_1$, there exists a one parameter family of metrics \tilde{g} given by*

$$\tilde{g} = \left[1 + B \int_x^1 \frac{ds}{\sqrt{h(s)}} \right]^4 g, \quad B > 0,$$

that satisfies $\Lambda_{g,\Gamma_D,\Gamma_N}(0) = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}(0)$.

Proof. — We only prove (1). Assume that $\lambda^2 = 0$. For arbitrary smooth positive functions f, h on $[0, 1]$, the function c solution of (4.17) is explicitly given by

$$c(x) = A + B \int_0^x \frac{ds}{\sqrt{h(s)}},$$

for some constants A, B . Since we also demand $c(0) = 1$ and $c^4 > 0$, we only consider the one parameter solutions

$$c(x) = 1 + B \int_0^x \frac{ds}{\sqrt{h(s)}},$$

for some constant $B > 0$. Then we set $\tilde{g} = c^4 g$ and we use the same proof as in Theorem 4.5 to conclude that $\Lambda_{g,\Gamma_D,\Gamma_N}(0) = \Lambda_{\tilde{g},\Gamma_D,\Gamma_N}(0)$. □

4.3. The 3D anisotropic Calderón problem with data measured on distinct connected components

In this Section, we prove our third main theorem, namely we give a counterexample to uniqueness for the anisotropic Calderón problem for three dimensional Riemannian manifolds with data measured on two different connected components of the boundary.

THEOREM 4.8. — *Let (M, g) and (M, \tilde{g}) denote Riemannian manifolds of the form (4.1), i.e.*

$$g = f(x)dx^2 + f(x)dy^2 + h(x)dz^2, \quad \tilde{g} = \tilde{f}(x)dx^2 + \tilde{f}(x)dy^2 + \tilde{h}(x)dz^2.$$

Let $\lambda^2 \in \mathbb{R}$. Let Γ_D and Γ_N be non-empty open sets belonging to two different connected components of ∂M . Then there exist infinitely many pairs of non-isometric metrics (g, \tilde{g}) given by $\tilde{g} = c^4g$ where c are smooth positive strictly increasing or decreasing functions such that $c(1)^3 = c(0) \neq 1$ if $\Gamma_D \subset \Gamma_0$ and $\Gamma_N \subset \Gamma_1$ or $c(0)^3 = c(1) \neq 1$ if $\Gamma_D \subset \Gamma_1$ and $\Gamma_N \subset \Gamma_0$, satisfying

$$\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(\lambda^2).$$

Proof. — The proof is almost the same as that of Theorem 4.5.

Let us assume that $\Gamma_D \subset \Gamma_0 = \{0\} \times T^1$ and $\Gamma_N \subset \Gamma_1 = \{1\} \times T^1$. We construct pairs of metrics (g, \tilde{g}) satisfying $\Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(\lambda^2)$ as follows. Let f and c be any smooth positive function on $[0, 1]$ such that $c(1)^3 = c(0)$ and $c'(x) \neq 0$ for all $x \in [0, 1]$.

Define

$$(4.35) \quad h = Ce^{-2} \int_0^x \frac{c''(s) + \lambda^2 f(s)(c(s) - c^5(s))}{c'(s)} ds, \quad \tilde{f} = c^4 f, \quad \tilde{h} = c^4 h.$$

Clearly, we have then $\tilde{g} = c^4g$. Using Lemma 4.3, it is immediate to check that $q_{\lambda n} = \tilde{q}_{\lambda n}$ for all $n \in \mathbb{Z}$ where $q_{\lambda n}$ and $\tilde{q}_{\lambda n}$ are given by (4.10). In particular, for such choices of metrics (g, \tilde{g}) , we always have $\Delta(m^2, n^2) = \tilde{\Delta}(m^2, n^2)$ for all $m, n \in \mathbb{Z}$. Moreover, our assumption $c(1)^3 = c(0)$ is then equivalent to

$$\frac{1}{\sqrt{f(1)}} \left(\frac{h(0)}{h(1)} \right)^{1/4} = \frac{1}{\sqrt{\tilde{f}(1)}} \left(\frac{\tilde{h}(0)}{\tilde{h}(1)} \right)^{1/4}.$$

Now, using (4.15) and the ensuing discussion, we see that for all $\psi \in H^{1/2}(T^2)$ with $\text{supp } \psi \subset \Gamma_D$, we have

$$\begin{aligned}
 (4.36) \quad & \Lambda_{g, \Gamma_D, \Gamma_N}(\lambda^2)(\psi) \\
 & := - \left[\sum_{m,n} \left(\frac{1}{\sqrt{f(1)}} \left(\frac{h(0)}{h(1)} \right)^{1/4} \frac{1}{\Delta(m^2, n^2)} \right) \hat{\psi}(m, n) e^{imy+inz} \right]_{|(y,z) \in \Gamma_N} \\
 & = - \left[\sum_{m,n} \left(\frac{1}{\sqrt{\tilde{f}(1)}} \left(\frac{\tilde{h}(0)}{\tilde{h}(1)} \right)^{1/4} \frac{1}{\tilde{\Delta}(m^2, n^2)} \right) \hat{\psi}(m, n) e^{imy+inz} \right]_{|(y,z) \in \Gamma_N} \\
 & = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(\lambda^2)(\psi).
 \end{aligned}$$

Since the above equality holds for all $\psi \in H^{1/2}(T^2)$ with $\text{supp } \psi \subset \Gamma_D$, the result is proved. \square

Finally, we can slightly improve our non-uniqueness result in the case of zero frequency. We have

THEOREM 4.9. — *Let (M, g) denotes a Riemannian manifold of the form (4.1), i.e.*

$$g = f(x)dx^2 + f(x)dy^2 + h(x)dz^2.$$

Let Γ_D and Γ_N be non-empty open sets belonging to two different connected component of ∂M . Then

- (1) *if $\Gamma_D \subset \Gamma_0$ and $\Gamma_N \subset \Gamma_1$, there exists a one parameter family of metrics \tilde{g} given by*

$$\tilde{g} = \left[A + \frac{A^3 - A}{\int_0^1 \frac{ds}{\sqrt{h(s)}}} \int_x^1 \frac{ds}{\sqrt{h(s)}} \right]^4 g, \quad A > 0,$$

that satisfies $\Lambda_{g, \Gamma_D, \Gamma_N}(0) = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(0)$.

- (2) *if $\Gamma_D \subset \Gamma_1$ and $\Gamma_N \subset \Gamma_0$, there exists a one parameter family of metrics \tilde{g} given by*

$$\tilde{g} = \left[A + \frac{A^3 - A}{\int_0^1 \frac{ds}{\sqrt{h(s)}}} \int_0^x \frac{ds}{\sqrt{h(s)}} \right]^4 g, \quad A > 0,$$

that satisfies $\Lambda_{g, \Gamma_D, \Gamma_N}(0) = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(0)$.

Proof. — We only prove (1). Assume that $\lambda^2 = 0$. For arbitrary smooth positive functions f, h on $[0, 1]$, the function c solution of (4.17) is explicitly

given by

$$c(x) = A + B \int_x^1 \frac{ds}{\sqrt{h(s)}},$$

for some constants A, B . Since we also requires that $c(1)^3 = c(0)$, we only consider the one parameter family of functions

$$c(x) = A + \frac{A^3 - A}{\int_0^1 \frac{ds}{\sqrt{h(s)}}} \int_x^1 \frac{ds}{\sqrt{h(s)}}$$

for some constant $A > 0$. Then we set $\tilde{g} = c^4 g$ and we use the same proof as in Theorem 4.5 to conclude that $\Lambda_{g, \Gamma_D, \Gamma_N}(0) = \Lambda_{\tilde{g}, \Gamma_D, \Gamma_N}(0)$. \square

4.4. The 3D anisotropic Calderón problem with a potential

In this Section, we treat the anisotropic Calderón problem (Q3) with a potential $V = V(x) \in L^\infty(M)$, that is a potential depending only on the variable x , for our family of metrics (4.1). The global DN map $\lambda_{g,V}(\lambda^2)$ associated to the Dirichlet problem

$$(4.37) \quad \begin{cases} (-\Delta_g + V)u = \lambda^2 u, & \text{on } M, \\ u = \psi, & \text{on } \partial M, \end{cases}$$

with λ^2 not belonging to the Dirichlet spectrum of $-\Delta_g + V$ can be constructed in the same way as in Section 4 since $V = V(x)$ respects the symmetry of (M, g) . On each Fourier mode $Y_{mn} = e^{imy+inz}$, we get the following expression for the induced DN map:

$$(4.38) \quad \Lambda_{g,V}^{mn}(\lambda^2) = \begin{pmatrix} \frac{(\log h)'(0)}{4\sqrt{f(0)}} - \frac{M_V(m^2, n^2)}{\sqrt{f(0)}} & -\frac{h^{1/4}(1)}{h^{1/4}(0)\sqrt{f(0)}\Delta_V(m^2, n^2)} \\ -\frac{h^{1/4}(0)}{h^{1/4}(1)\sqrt{f(1)}\Delta_V(m^2, n^2)} & -\frac{(\log h)'(1)}{4\sqrt{f(1)}} - \frac{N_V(m^2, n^2)}{\sqrt{f(1)}} \end{pmatrix},$$

where the characteristic function $\Delta_V(m^2, n^2)$ and the Weyl–Titchmarsh functions $M_V(m^2, n^2)$ and $N_V(m^2, n^2)$ defined by (4.13)–(4.14) are associated to the radial ODE

$$(4.39) \quad -v'' + q_{\lambda,V,n}(x)v = -\mu^2 v,$$

with potential $q_{\lambda,V,n} = \frac{[(\log h)']^2}{16} + \frac{(\log h)''}{4} + n^2 \frac{f}{h} + (V - \lambda^2)f$ and boundary conditions

$$(4.40) \quad v(0) = 0, \quad v(1) = 0.$$

We also recall the dictionary between the above coefficients of the DN map and the notations used in the Introduction

$$\begin{aligned}
 L_V^{mn}(\lambda^2) &= \frac{(\log h)'(0)}{4\sqrt{f(0)}} - \frac{1}{\sqrt{f(0)}} M_V(m^2, n^2) = \Lambda_{g,V,\Gamma_0}^{mn}(\lambda^2), \\
 R_V^{mn}(\lambda^2) &= -\frac{(\log h)'(1)}{4\sqrt{f(1)}} - \frac{1}{\sqrt{f(1)}} N_V(m^2, n^2) = \Lambda_{g,V,\Gamma_1}^{mn}(\lambda^2), \\
 T_{L,V}^{mn}(\lambda^2) &= -\frac{1}{\sqrt{f(1)}} \frac{h^{1/4}(0)}{h^{1/4}(1)} \frac{1}{\Delta_V(m^2, n^2)} = \Lambda_{g,V,\Gamma_0,\Gamma_1}^{mn}(\lambda^2), \\
 T_{R,V}^{mn}(\lambda^2) &= -\frac{1}{\sqrt{f(0)}} \frac{h^{1/4}(1)}{h^{1/4}(0)} \frac{1}{\Delta_V(m^2, n^2)} = \Lambda_{g,V,\Gamma_1,\Gamma_0}^{mn}(\lambda^2),
 \end{aligned}$$

where $\Gamma_0 = \{0\} \times T^1$ and $\Gamma_1 = \{1\} \times T^1$.

We prove:

THEOREM 4.10. — *Let (M, g) a smooth compact Riemannian manifold of the form (4.1), i.e.*

$$g = f(x)dx^2 + f(x)dy^2 + h(x)dz^2.$$

Let $V, \tilde{V} \in L^\infty(M)$ be two potentials that only depend on the variable x . Let the frequency λ^2 be fixed and not belonging to the Dirichlet spectra of $-\Delta_g + V$ and $-\Delta_g + \tilde{V}$. Let Γ_D, Γ_N be nonempty open subsets belonging to the same connected component of ∂M , with Γ_N containing an annular region of the type

$$(y_0 - \delta, y_0 + \delta) \times T^1, \quad \text{or} \quad T^1 \times (z_0 - \delta, z_0 + \delta),$$

where $\delta > 0$. Assume moreover

$$\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,\tilde{V},\Gamma_D,\Gamma_N}(\lambda^2).$$

Then

$$\tilde{V} = V.$$

Proof. — Assume for instance that $\Gamma_D, \Gamma_N \subset \Gamma_0$ and

$$\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,\tilde{V},\Gamma_D,\Gamma_N}(\lambda^2).$$

Then the same proof as in Theorem 4.4 shows first that

$$\begin{aligned}
 (4.41) \quad M_V(\mu^2, n^2) - M_V(0, n^2) &= M_{\tilde{V}}(\mu^2, n^2) - M_{\tilde{V}}(0, n^2), \\
 &\forall \mu \in \mathbb{C} \setminus \mathbb{R}, \forall n \in \mathbb{Z}.
 \end{aligned}$$

Hence the Borg–Marchenko Theorem 2.4 gives

$$(4.42) \quad q_{\lambda,V,n} = q_{\lambda,\tilde{V},n}, \quad \text{on } [0, 1], \quad \forall n \in \mathbb{Z}.$$

We conclude using (4.42) and (4.39) that $V = \tilde{V}$ on $[0, 1]$. □

In the case where the Dirichlet and Neumann data Γ_D and Γ_N do not belong to the same connected component of ∂M , we are able to give very simple counterexamples to uniqueness, in the case where $f = h$. Precisely, we prove:

THEOREM 4.11. — *Let (M, g) a smooth compact Riemannian manifold of the form*

$$g = f(x)[dx^2 + dy^2 + dz^2].$$

Let $V, \tilde{V} \in L^\infty(M)$ be two potentials that only depend on the variable x . Let the frequency λ^2 be fixed and not belonging to the Dirichlet spectra of $-\Delta_g + V$ and $-\Delta_g + \tilde{V}$. Let Γ_D, Γ_N be nonempty open subsets belonging to distinct connected components of ∂M . Then, there exists an infinite dimensional family of explicit potentials $\tilde{V} \in L^\infty([0, 1])$ that satisfy

$$\Lambda_{g, V, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{g, \tilde{V}, \Gamma_D, \Gamma_N}(\lambda^2).$$

Proof. — In the case where $f = h$, the potentials $q_{\lambda, V, n}$ and $q_{\lambda, \tilde{V}, n}$ have the following simple form

$$\begin{aligned} q_{\lambda, V, n} &= \frac{[(\log f)']^2}{16} + \frac{(\log f)''}{4} + n^2 + (V(x) - \lambda^2)f(x), \\ q_{\lambda, \tilde{V}, n} &= \frac{[(\log f)']^2}{16} + \frac{(\log f)''}{4} + n^2 + (\tilde{V}(x) - \lambda^2)f(x). \end{aligned}$$

Clearly, since n^2 does not depend on x , the potentials $q_{\lambda, V, n}, q_{\lambda, \tilde{V}, n}$ are isospectral if and only if the potentials $\frac{[(\log f)']^2}{16} + \frac{(\log f)''}{4} + (V(x) - \lambda^2)f(x)$ and $\frac{[(\log f)']^2}{16} + \frac{(\log f)''}{4} + (\tilde{V}(x) - \lambda^2)f(x)$ are also isospectral. We recall that this is also equivalent (see the proofs of Theorems 3.8 and 4.4) to

$$(4.43) \quad \Delta_V(\mu^2, n^2) = \Delta_{\tilde{V}}(\mu^2, n^2), \quad \forall \mu \in \mathbb{C}, \forall n \in \mathbb{Z},$$

which would imply that

$$(4.44) \quad \Lambda_{g, V, \Gamma_D, \Gamma_N}(\lambda^2) = \Lambda_{g, \tilde{V}, \Gamma_D, \Gamma_N}(\lambda^2),$$

according to (4.38).

We deduce thus from [37] that given a potential $V \in L^\infty(M)$ as above, there exists an infinite dimensional family of explicit potentials \tilde{V} satisfying (4.43) and thus (4.44). More precisely, the family

$$(4.45) \quad \tilde{V}_{\lambda, k, t}(x) = V(x) - \frac{2}{f(x)} \frac{d^2}{dx^2} \log \theta_{k, t}(x), \quad \forall k \geq 1, \quad t \in \mathbb{R},$$

with

$$(4.46) \quad \theta_{k,t}(x) = 1 + (e^t - 1) \int_x^1 v_k^2(s) ds,$$

where v_k is the normalized eigenfunction of (4.39)–(4.40) associated to the k th-eigenvalue α_k^2 satisfies

$$\Lambda_{g,V,\Gamma_D,\Gamma_N}(\lambda^2) = \Lambda_{g,\tilde{V}_{\lambda,k,t},\Gamma_D,\Gamma_N}(\lambda^2), \quad \forall k \geq 1, \quad t \in \mathbb{R}. \quad \square$$

BIBLIOGRAPHY

- [1] K. ASTALA, L. PÄIVÄRINTA & M. LASSAS, “Calderón’s inverse problem for anisotropic conductivity in the plane”, *Commun. Partial Differ. Equations* **30** (2005), no. 1-3, p. 207-224.
- [2] C. BENNEWITZ, “A proof of the local Borg-Marchenko theorem”, *Commun. Math. Phys.* **218** (2001), no. 1, p. 131-132.
- [3] R. P. BOAS, JR., *Entire functions*, Pure and Applied Mathematics, vol. 5, Academic Press Inc., 1954, x+276 pages.
- [4] G. BORG, “Eine Umkehrung der Sturm-Liouvilleschen Eigenwertaufgabe. Bestimmung der Differentialgleichung durch die Eigenwerte”, *Acta Math.* **78** (1946), p. 1-96.
- [5] ———, “Uniqueness theorems in the spectral theory of $y'' + (\lambda - q(x))y = 0$ ”, in *Den 11te Skandinaviske Matematikerkongress, Trondheim, 1949*, Johan Grundt Tanums Forlag, 1952, p. 276-287.
- [6] T. DAUDÉ, D. GOBIN & F. NICOLEAU, “Local inverse scattering at fixed energy in spherically symmetric asymptotically hyperbolic manifolds”, *Inverse Probl. Imaging* **10** (2016), no. 3, p. 659-688.
- [7] T. DAUDÉ, N. KAMRAN & F. NICOLEAU, “Inverse scattering at fixed energy on asymptotically hyperbolic Liouville surfaces”, *Inverse Probl.* **31** (2015), no. 12, Art. ID 125009, 37 p.
- [8] T. DAUDÉ & F. NICOLEAU, “Inverse scattering at fixed energy in de Sitter-Reissner-Nordström black holes”, *Ann. Henri Poincaré* **12** (2011), no. 1, p. 1-47.
- [9] ———, “Direct and inverse scattering at fixed energy for massless charged Dirac fields by Kerr-Newman-de Sitter black holes”, *Mem. Am. Math. Soc.* **247** (2017), no. 1170, p. iv+113.
- [10] D. DOS SANTOS FERREIRA, C. KENIG, M. SALO & G. UHLMANN, “Limiting Carleman weights and anisotropic inverse problems”, *Invent. Math.* **178** (2009), no. 1, p. 119-171.
- [11] D. DOS SANTOS FERREIRA, Y. KURYLEV, M. LASSAS & M. SALO, “The Calderón problem in transversally anisotropic geometries”, *J. Eur. Math. Soc.* **18** (2016), no. 11, p. 2579-2626.
- [12] J. ECKHARDT & G. TESCHL, “Uniqueness results for one-dimensional Schrödinger operators with purely discrete spectra”, *Trans. Am. Math. Soc.* **365** (2013), no. 7, p. 3923-3942.
- [13] G. FREILING & V. YURKO, “Inverse problems for differential operators with singular boundary conditions”, *Math. Nachr.* **278** (2005), no. 12-13, p. 1561-1578.
- [14] F. GESZTESY & B. SIMON, “On local Borg-Marchenko uniqueness results”, *Commun. Math. Phys.* **211** (2000), no. 2, p. 273-287.
- [15] C. GUILLARMOU & A. SÁ BARRETO, “Inverse problems for Einstein manifolds”, *Inverse Probl. Imaging* **3** (2009), no. 1, p. 1-15.

- [16] C. GUILLARMOU & L. TZOU, “Calderón inverse problem with partial data on Riemann surfaces”, *Duke Math. J.* **158** (2011), no. 1, p. 83-120.
- [17] ———, “The Calderón inverse problem in two dimensions”, in *Inverse problems and applications: inside out. II*, Mathematical Sciences Research Institute Publications, vol. 60, Cambridge University Press, 2013, p. 119-166.
- [18] L. HÖRMANDER, *The analysis of linear partial differential operators. I: Distribution theory and Fourier analysis*, Classics in Mathematics, Springer, 2003, x+440 pages.
- [19] O. Y. IMANUVILOV, G. UHLMANN & M. YAMAMOTO, “The Calderón problem with partial data in two dimensions”, *J. Am. Math. Soc.* **23** (2010), no. 3, p. 655-691.
- [20] ———, “Inverse boundary value problem by measuring Dirichlet data and Neumann data on disjoint sets”, *Inverse Probl.* **27** (2011), no. 8, Art. ID 085007, 26 p.
- [21] V. ISAKOV, “On uniqueness in the inverse conductivity problem with local data”, *Inverse Probl. Imaging* **1** (2007), no. 1, p. 95-105.
- [22] H. KANG & K. YUN, “Boundary determination of conductivities and Riemannian metrics via local Dirichlet-to-Neumann operator”, *SIAM J. Math. Anal.* **34** (2002), no. 3, p. 719-735.
- [23] A. KATCHALOV, Y. KURYLEV & M. LASSAS, *Inverse boundary spectral problems*, Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics, vol. 123, Chapman & Hall/CRC Press, 2001, xx+290 pages.
- [24] A. KATCHALOV, Y. KURYLEV, M. LASSAS & N. MANDACHE, “Equivalence of time-domain inverse problems and boundary spectral problems”, *Inverse Probl.* **20** (2004), no. 2, p. 419-436.
- [25] C. KENIG & M. SALO, “The Calderón problem with partial data on manifolds and applications”, *Anal. PDE* **6** (2013), no. 8, p. 2003-2048.
- [26] ———, “Recent progress in the Calderón problem with partial data”, in *Inverse problems and applications*, Contemporary Mathematics, vol. 615, American Mathematical Society, 2014, p. 193-222.
- [27] C. KENIG, J. SJÖSTRAND & G. UHLMANN, “The Calderón problem with partial data”, *Ann. Math.* **165** (2007), no. 2, p. 567-591.
- [28] A. KOSTENKO, A. SAKHNOVICH & G. TESCHL, “Weyl-Titchmarsh theory for Schrödinger operators with strongly singular potentials”, *Int. Math. Res. Not.* (2012), no. 8, p. 1699-1747.
- [29] M. LASSAS & L. OKSANEN, “An inverse problem for a wave equation with sources and observations on disjoint sets”, *Inverse Probl.* **26** (2010), no. 8, Art. ID 085012, 19 p.
- [30] ———, “Inverse problem for the Riemannian wave equation with Dirichlet data and Neumann data on disjoint sets”, *Duke Math. J.* **163** (2014), no. 6, p. 1071-1103.
- [31] M. LASSAS, M. TAYLOR & G. UHLMANN, “The Dirichlet-to-Neumann map for complete Riemannian manifolds with boundary”, *Commun. Anal. Geom.* **11** (2003), no. 2, p. 207-221.
- [32] M. LASSAS & G. UHLMANN, “On determining a Riemannian manifold from the Dirichlet-to-Neumann map”, *Ann. Sci. Éc. Norm. Supér.* **34** (2001), no. 5, p. 771-787.
- [33] J. M. LEE & G. UHLMANN, “Determining anisotropic real-analytic conductivities by boundary measurements”, *Commun. Pure Appl. Math.* **42** (1989), no. 8, p. 1097-1112.
- [34] B. Y. LEVIN, *Lectures on entire functions*, Translations of Mathematical Monographs, vol. 150, American Mathematical Society, 1996, In collaboration with and with a preface by Yu. Lyubarskii, M. Sodin and V. Tkachenko, Translated from the Russian manuscript by Tkachenko, xvi+248 pages.

- [35] W. R. B. LIONHEART, “Conformal uniqueness results in anisotropic electrical impedance imaging”, *Inverse Probl.* **13** (1997), no. 1, p. 125-134.
- [36] V. A. MARČENKO, “Some questions of the theory of one-dimensional linear differential operators of the second order. I”, *Trudy Moskov. Mat. Obšč.* **1** (1952), p. 327-420.
- [37] J. PÖSCHEL & E. TRUBOWITZ, *Inverse spectral theory*, Pure and Applied Mathematics, vol. 130, Academic Press Inc., 1987, x+192 pages.
- [38] RAKESH, “Characterization of transmission data for Webster’s horn equation”, *Inverse Probl.* **16** (2000), no. 2, p. L9-L24.
- [39] A. G. RAMM, “An inverse scattering problem with part of the fixed-energy phase shifts”, *Commun. Math. Phys.* **207** (1999), no. 1, p. 231-247.
- [40] T. REGGE, “Introduction to complex orbital momenta”, *Nuovo Cimento* **14** (1959), p. 951-976.
- [41] W. RUDIN, *Real and complex analysis*, 3rd ed., McGraw-Hill Book Co., 1987, xiv+416 pages.
- [42] M. SALO, “The Calderón problem on Riemannian manifolds”, in *Inverse problems and applications: inside out. II*, Mathematical Sciences Research Institute Publications, vol. 60, Cambridge University Press, 2013, p. 167-247.
- [43] M. SPIEGEL, *Theory and problems of complex variables*, 2nd ed., McGraw-Hill Book Co., 2009.
- [44] G. TESCHL, *Mathematical methods in quantum mechanics. With applications to Schrödinger operators*, 2nd ed., Graduate Studies in Mathematics, vol. 157, American Mathematical Society, 2014, xiv+358 pages.
- [45] G. UHLMANN, “Electrical impedance tomography and Calderón’s problem”, *Inverse Probl.* **25** (2009), no. 12, Art. ID 123011, 39 p.
- [46] A. ZETTL, *Sturm-Liouville theory*, Mathematical Surveys and Monographs, vol. 121, American Mathematical Society, 2005, xii+328 pages.

Manuscrit reçu le 14 octobre 2016,
 accepté le 14 septembre 2017.

Thierry DAUDÉ
 Département de Mathématiques, UMR CNRS 8088
 Université de Cergy-Pontoise
 95302 Cergy-Pontoise (France)
 thierry.daude@u-cergy.fr

Niky KAMRAN
 Department of Mathematics and Statistics
 McGill University
 Montreal, QC, H3A 2K6 (Canada)
 nkamran@math.mcgill.ca

François NICOLEAU
 Laboratoire de Mathématiques Jean Leray, UMR
 CNRS 6629
 2 Rue de la Houssinière BP 92208
 44322 Nantes Cedex 03 (France)
 francois.nicoleau@math.univ-nantes.fr