

 Open access • Journal Article • DOI:10.30757/ALEA.V18-07

Non universality of fluctuations of outlier eigenvectors for block diagonal deformations of Wigner matrices — [Source link](#)

Mireille Capitaine, Catherine Donati-Martin

Institutions: Institut de Mathématiques de Toulouse

Published on: 01 Jan 2021 - ALEA-Latin American Journal of Probability and Mathematical Statistics (Institute for Applied and Pure Mathematics (IMPA))

Topics: Block matrix, Random matrix, Matrix (mathematics), Eigenvalues and eigenvectors and Hermitian matrix

Related papers:

- [Non universality of fluctuations of outlier eigenvectors for block diagonal deformations of Wigner matrices](#)
- [Central limit theorem for mesoscopic eigenvalue statistics of deformed Wigner matrices and sample covariance matrices](#)
- [The Isotropic Semicircle Law and Deformation of Wigner Matrices](#)
- [Local semicircle law and complete delocalization for Wigner random matrices](#)
- [Edge universality for deformed Wigner matrices](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/non-universality-of-fluctuations-of-outlier-eigenvectors-for-42hl1943yz>



HAL
open science

Non universality of fluctuations of outlier eigenvectors for block diagonal deformations of Wigner matrices

Mireille Capitaine, Catherine Donati-Martin

► **To cite this version:**

Mireille Capitaine, Catherine Donati-Martin. Non universality of fluctuations of outlier eigenvectors for block diagonal deformations of Wigner matrices. *ALEA : Latin American Journal of Probability and Mathematical Statistics*, Instituto Nacional de Matemática Pura e Aplicada, 2021, 18 (1), pp.129-165. 10.30757/ALEA.v18-07. hal-03357320

HAL Id: hal-03357320

<https://hal.archives-ouvertes.fr/hal-03357320>

Submitted on 18 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non universality of fluctuations of outlier eigenvectors for block diagonal deformations of Wigner matrices

M. Capitaine* and C. Donati-Martin†

Abstract

In this paper, we investigate the fluctuations of a unit eigenvector associated to an outlier in the spectrum of a spiked $N \times N$ complex Deformed Wigner matrix M_N . M_N is defined as follows: $M_N = W_N/\sqrt{N} + A_N$ where W_N is an $N \times N$ Hermitian Wigner matrix whose entries have a law μ satisfying a Poincaré inequality and the matrix A_N is a block diagonal matrix, with an eigenvalue θ of multiplicity one, generating an outlier in the spectrum of M_N . We prove that the fluctuations of the norm of the projection of a unit eigenvector corresponding to the outlier of M_N onto a unit eigenvector corresponding to θ are not universal. Indeed, we take away a fit approximation of its limit from this norm and prove the convergence to zero as N goes to ∞ of the Lévy–Prohorov distance between this rescaled quantity and the convolution of μ and a centered Gaussian distribution (whose variance may depend upon N and may not converge).

Key words: Random matrices; Outliers; eigenvectors; Fluctuations; Nonuniversality, Free probability.

Mathematics Subject Classification 2000: 15A18, 15A52, 60F05, 60B12, 46L54.

1 Introduction

To begin with, we introduce some notations.

* *Institut de Mathématiques de Toulouse; UMR5219; Université de Toulouse; CNRS; UPS, 118 rte de Narbonne F-31062 Toulouse, FRANCE. E-mail: mireille.capitaine@math.univ-toulouse.fr*

† *Laboratoire de Mathématiques de Versailles, UVSQ, CNRS, Université Paris-Saclay, 78035-Versailles Cedex, France, E-mail: catherine.donati-martin@uvsq.fr*

- $M_N(\mathbb{C})$ is the set of $N \times N$ matrices with complex entries, $M_N^{sa}(\mathbb{C})$ the subset of self-adjoint elements of $M_N(\mathbb{C})$ and I_N the identity matrix.
- Tr_N denotes the trace and $\text{tr}_N = \frac{1}{N} \text{Tr}_N$ the normalized trace on $M_N(\mathbb{C})$.
- $\|\cdot\|$ denotes the operator norm on $M_N(\mathbb{C})$.
- For any $X \in M_N^{sa}(\mathbb{C})$, $(\lambda_1(X), \dots, \lambda_N(X))$ denote the eigenvalues of X ranked in decreasing order and the empirical spectral measure of X is defined by

$$\mu_X := \frac{1}{N} \sum_{i=1}^N \delta_{\lambda_i(X)}.$$

- For a probability measure τ on \mathbb{R} , $\text{supp}(\tau)$ denotes the support of τ and $g_\tau : z \in \mathbb{C} \setminus \text{supp}(\tau) \mapsto \int \frac{1}{z-x} d\tau(x)$ is the Stieltjes transform of τ .
- d_{LP} denotes the Lévy-Prohorov distance, which is a metric for the topology of the convergence in distribution.

1.1 Wigner matrices

Wigner matrices are complex Hermitian random matrices whose entries are independent (up to the symmetry condition). They were introduced by Wigner in the fifties, in connection with nuclear physics. Here, we will consider Hermitian Wigner matrices of the following form :

$$X_N = \frac{1}{\sqrt{N}} W_N$$

where W_N is an Hermitian matrix, $\{W_{ii}, \sqrt{2}\mathcal{R}W_{ij}, \sqrt{2}\mathcal{I}W_{ij}\}_{1 \leq i < j}$ are independent identically distributed random variables with law μ , with mean zero and variance σ^2 . If the entries are independent Gaussian variables, $X_N =: X_N^G$ is a matrix from the Gaussian Unitary Ensemble (G.U.E.).

There is currently a quite precise knowledge of the asymptotic spectral properties (i.e. when the dimension of the matrix tends to infinity) of Wigner matrices. This understanding covers both the so-called global regime (asymptotic behavior of the spectral measure) and the local regime (asymptotic behavior of the extreme eigenvalues and eigenvectors, spacings...). Wigner proved that a precise description of the limiting spectrum of these matrices can be achieved.

Theorem 1. [39, 40]

$$\mu_{X_N} \xrightarrow{w} \mu_{sc} \text{ a.s. when } N \rightarrow +\infty$$

where

$$\frac{d\mu_{sc}}{dx}(x) = \frac{1}{2\pi\sigma^2} \sqrt{4\sigma^2 - x^2} 1_{[-2\sigma, 2\sigma]}(x) \quad (1)$$

is the so-called semi-circular distribution.

A priori, the convergence of the spectral measure does not prevent an asymptotically negligible fraction of eigenvalues from going away from the limiting support (called *outliers* in the following). Actually, it turns out that Wigner matrices do not exhibit outliers.

Theorem 2. [5] Assume that the entries of W_N has finite fourth moment, then almost surely,

$$\lambda_1(X_N) \rightarrow 2\sigma \text{ and } \lambda_N(X_N) \rightarrow -2\sigma \text{ when } N \rightarrow +\infty.$$

In [37], Tracy and Widom derived the limiting distribution (called the Tracy-Widom law) of the largest eigenvalue of a G.U.E. matrix.

Theorem 3. Let $q : \mathbb{R} \rightarrow \mathbb{R}$ be the unique solution of the differential equation

$$q''(x) = xq(x) + 2q(x)^3$$

such that $q(x) \sim_{x \rightarrow +\infty} Ai(x)$ where Ai is the Airy function, unique solution on \mathbb{R} of the differential equation $f''(x) = xf(x)$ satisfying $f(x) \sim_{x \rightarrow +\infty} (4\pi\sqrt{x})^{1/2} \exp(-2/3x^{3/2})$. Then

$$\lim_{N \rightarrow +\infty} \mathbb{P} \left(\frac{N^{2/3}}{\sigma} \left(\frac{\lambda_1(X_N^G)}{\sqrt{N}} - 2\sigma \right) \leq s \right) = F_2(s),$$

where $F_2(s) = \exp \left(- \int_s^{+\infty} (x-s)q^2(x)dx \right)$.

The first main step to prove the universality conjecture for fluctuations of the largest eigenvalue of Wigner matrices has been achieved by Soshnikov [35]; in [28], a necessary and sufficient condition on off-diagonal entries of the Wigner matrix is established for the distribution of the largest eigenvalue to weakly converge to the Tracy-Widom distribution. We also refer to these papers for references on investigations on edge universality.

In regards to eigenvectors, it is well known that the matrix whose columns are the eigenvectors of a G.U.E. matrix can be chosen to be distributed according to the Haar measure on the unitary group. In the non-Gaussian case,

the exact distribution of the eigenvectors cannot be computed. However, the eigenvectors of general Wigner matrices have been the object of a growing interest and in several papers, a delocalization and universality property were shown for the eigenvectors of these standard models (see among others [13, 22, 23, 26, 36] and references therein). Heuristically, delocalization for a random matrix means that its normalized eigenvectors look like the vectors uniformly distributed over the unit sphere. Let us state for instance the following sample result.

Theorem 4. (*Isotropic delocalization, Theorem 2.16 from [13]*). *Let X_N be a $N \times N$ Wigner matrix satisfying some technical assumptions. Let $v(1), \dots, v(N)$ denote the normalized eigenvectors of X_N . Then, for any $C_1 > 0$ and $0 < \epsilon < 1/2$, there exists $C_2 > 0$ such that*

$$\sup_{1 \leq i \leq N} |\langle v(i), u \rangle| \leq \frac{N^\epsilon}{\sqrt{N}},$$

for any fixed unit vector $u \in \mathbb{C}^N$, with probability at least $1 - C_2 N^{-C_1}$.

1.2 Deformed Wigner matrices

Practical problems (in the theory of statistical learning, signal detection etc.) naturally lead to wonder about the spectrum reaction of a given random matrix after a deterministic perturbation. In those applications, the random matrix is the noise and the perturbed matrix is a noisy version of the information; the question is to know whether the observation of the spectral properties of the perturbed matrix can give access to significant parameters on the information. Theoretical results on these deformed random models may allow to establish statistical tests on these parameters. A typical illustration is the so-called BBP phenomenon (after Baik, Ben Arous, P  ch  ) which put forward outliers (eigenvalues that move away from the rest of the spectrum) and their Gaussian fluctuations for spiked covariance matrices in [6] and for low rank deformations of G.U.E. in [33].

In this paper, we consider additive perturbations of Wigner matrices. The pionner works on additive deformations go back to Pastur [32] for the behavior of the limiting spectral distribution and to F  redi and Koml  s [24] for the behavior of the largest eigenvalue.

We refer to [17] and the references therein for a survey on spectral properties of deformed random matrices.

The model studied is as follows :

$$M_N := \frac{W_N}{\sqrt{N}} + A_N, \tag{2}$$

where

(W) W_N is a complex Wigner matrix, that is a $N \times N$ random Hermitian matrix such that $\{W_{ii}, \sqrt{2}\mathcal{R}W_{ij}, \sqrt{2}\mathcal{I}W_{ij}\}_{1 \leq i < j}$ are independent identically distributed random variables with law μ . We assume that μ is a distribution with mean zero, variance σ^2 , and satisfies a Poincaré inequality (see Appendix). Note that this condition implies that μ has moments of any order (see Corollary 3.2 and Proposition 1.10 in [27]).

(A) A_N is a $N \times N$ deterministic Hermitian matrix, whose spectral measure μ_{A_N} converges to a compactly supported probability measure ν . We assume that A_N has a fixed number q of eigenvalues, not depending on N , outside the support of ν called spikes, whereas the distance of the other eigenvalues to the support of ν goes to 0.

The empirical spectral distribution μ_{M_N} converges a.s. towards the probability measure $\lambda := \mu_{sc} \boxplus \nu$ where μ_{sc} is the semicircular distribution with variance σ^2 and \boxplus denotes the free convolution, see [32] (in this paper, the limiting distribution is given via a functional equation for its Stieltjes transform), [1, Theorem 5.4.5]. We refer to [25, 30] for an introduction to free probability theory.

Concerning extremal eigenvalues, [20] proved that the spikes of A_N can generate outliers for the limiting spectrum of M_N , i.e. eigenvalues outside the support of the limiting distribution λ . More precisely, [20] proved the following (see [20, Theorem 8.1] for a more general statement).

Proposition 1. [20] *Assume that a spike θ with a fixed multiplicity k_0 in the spectrum of A_N satisfies :*

$$\theta \in \Theta_{\sigma, \nu} := \left\{ u \in \mathbb{R} \setminus \text{supp}(\nu), \int_{\mathbb{R}} \frac{d\nu(x)}{(u-x)^2} < \frac{1}{\sigma^2} \right\}. \quad (3)$$

Denote by n_0+1, \dots, n_0+k_0 the descending ranks of θ among the eigenvalues of A_N . Then the k_0 eigenvalues $(\lambda_{n_0+i}(M_N), 1 \leq i \leq k_0)$ converge almost surely outside the support of λ towards $\rho_\theta := \theta + \sigma^2 g_\nu(\theta)$. Moreover, these eigenvalues asymptotically separate from the rest of the spectrum since (with the conventions that $\lambda_0(M_N) = +\infty$ and $\lambda_{N+1}(M_N) = -\infty$) there exists $0 < \delta_0$ such that almost surely for all large N ,

$$\lambda_{n_0}(M_N) > \rho_\theta + \delta_0 \quad \text{and} \quad \lambda_{n_0+k_0+1}(M_N) < \rho_\theta - \delta_0. \quad (4)$$

Note that [20] assumes that the distribution μ is symmetric but this assumption can be removed. Indeed, this assumption is used for establishing

Theorem 5.1 in [20] that is now generalized by Theorem 1.1 in [8]. Note also that [20] assumes moreover that the support of ν has a finite number of connected components in order to prove Theorem 6.1 in [20] but this assumption is removed in Theorem 2.3 in [21].

Remark 1. *Note that*

$$\{u \in \mathbb{R} \setminus \text{supp}(\nu), \int_{\mathbb{R}} \frac{d\nu(x)}{(u-x)^2} < \frac{1}{\sigma^2}\} = \{u \in \mathbb{R} \setminus \text{supp}(\nu), H'(u) > 0\}$$

and for any θ in this set, $\rho_\theta = \theta + \sigma^2 g_\nu(\theta) = H(\theta)$, where H is defined by (67) (see Section 2.4 below).

It turns out that we can also describe the angle between the eigenvector associated to the outlier of M_N and the corresponding eigenvector associated to the spike θ . Capitaine [14] (see also [17]) proved

Proposition 2. [14] *We keep the notation and hypothesis of Proposition 1. Let ξ be a unit eigenvector associated to one of the eigenvalues $(\lambda_{n_0+i}(M_N), 1 \leq i \leq k_0)$. Then, a.s.*

$$\|P_{\text{Ker}(A_N - \theta I)}(\xi)\|^2 \xrightarrow{N \rightarrow +\infty} \tau(\theta) := 1 - \sigma^2 \int \frac{1}{(\theta - x)^2} d\nu(x), \quad (5)$$

where P_E denotes the orthogonal projection onto any subspace E .

Note that fluctuations of outliers for deformed non-Gaussian Wigner matrices have been more extensively studied in the case of perturbations A_N of fixed rank r . We emphasize that the limiting distribution in the CLT for outliers depends on the localisation/delocalisation of the eigenvector of the spike. Roughly speaking, in the delocalized case, the limiting distribution of the fluctuations of the corresponding outliers is Gaussian. In the localized case, the limiting distribution depends on the distribution μ of the entries and thus, this uncovers a non universality phenomenon. We refer to [19] for these results.

We first recall the fluctuations of the largest eigenvalue $\lambda_1(M_N)$ when the matrix A_N is a diagonal matrix of rank 1 in the localized case.

Proposition 3. [18] *Assume that $A_N = \text{diag}(\theta, 0, \dots, 0)$ with $\theta > \sigma$. The fluctuations of $\lambda_1(M_N)$ around $\rho_\theta = \theta + \frac{\sigma^2}{\theta}$ are given by*

$$c_\theta \sqrt{N} (\lambda_1(M_N) - \rho_\theta) \xrightarrow[N \rightarrow \infty]{(law)} \mu \star N(0, v_\theta^2)$$

where $c_\theta = (1 - \frac{\sigma^2}{\theta^2})^{-1}$, $v_\theta^2 = \frac{1}{2} \frac{m_4 - 3\sigma^4}{\theta^2} + \frac{\sigma^4}{\theta^2 - \sigma^2}$ and m_4 denotes the fourth moment of μ .

Capitaine and P ech e [21] proved a fluctuation result for any outlier of a full rank deformation of a G.U.E. matrix. Their result yields the following

Proposition 4. [21] *Assume that W_N is a G.U.E. matrix (that is $\mu = \mathcal{N}(0, \sigma^2)$) and that A_N is a diagonal matrix with a spike $\lambda_{i_0}(A_N) = \theta \in \Theta_{\sigma, \nu}$ of multiplicity one and limiting spectral distribution ν . The fluctuations of $\lambda_{i_0}(M_N)$ around*

$$\rho_\theta^{(N)} = \theta + \sigma^2 \frac{1}{N-1} \sum_{\lambda_j(A_N) \neq \theta} \frac{1}{\theta - \lambda_j(A_N)}$$

are given by¹:

$$c_{\theta, \nu} \sqrt{N} (\lambda_1(M_N) - \rho_\theta^{(N)}) \xrightarrow[N \rightarrow \infty]{(law)} \mathcal{N}(0, \sigma_{\theta, \nu}^2)$$

where $c_{\theta, \nu} = \left(1 - \sigma^2 \int \frac{1}{(\theta - x)^2} d\nu(x)\right)^{-1}$ and

$$\sigma_{\theta, \nu}^2 = \sigma^2 \left(1 - \sigma^2 \int \frac{1}{(\theta - x)^2} d\nu(x)\right)^{-1}.$$

1.3 Main results

In the following, we consider block diagonal perturbations. We focus on spikes of the perturbation with multiplicity one generating an outlier in the spectrum of the deformed Wigner model. Therefore, we shall consider the following assumption on A_N throughout the paper:

(A') A_N satisfies **(A)** and

$$A_N = \text{diag}(A_p, A_{N-p}),$$

with A_{N-p} a $(N-p) \times (N-p)$ Hermitian matrix for some fixed integer p , $A_p = PDP^*$ is a fixed matrix (independent of N) where P a $p \times p$ unitary matrix and D is a diagonal matrix.

Assume that A_N has a spike of multiplicity one, which is an eigenvalue of A_p , $\theta = \lambda_{i_0}(A_N)$ for some i_0 , satisfying (3). Without loss of generality, we can assume that $\theta = D_{11}$.

We set

$$W_N = \begin{pmatrix} W_p & Y^* \\ Y & W_{N-p} \end{pmatrix},$$

¹They consider fluctuations around this point depending on N in order to not prescribe speed of convergence of μ_{A_N} to ν .

where $W_p \in M_p(\mathbb{C})$, $Y \in M_{(N-p) \times p}(\mathbb{C})$ and $W_{N-p} \in M_{N-p}(\mathbb{C})$.

The main results of this paper are the following Theorems 6 and 7 on non universality of fluctuations of an eigenvector associated with such an outlier. But, sticking to the approach of [16], we first establish Theorem 5 below, which is an extension in the non-Gaussian case of Proposition 3 and Proposition 4, in the block diagonal case.

Let M_N be defined by (2) with assumptions **(W)** and **(A')**. By Proposition 1,

$$\lambda_{i_0}(M_N) \xrightarrow{N \rightarrow +\infty} \rho_\theta \text{ a.s.} \quad (6)$$

Define

$$\rho_N = \theta + \sigma^2 g_{\mu_{A_{N-p}}}(\theta). \quad (7)$$

Note that

$$\rho_N \xrightarrow{N \rightarrow +\infty} \rho_\theta. \quad (8)$$

Theorem 5. *Let M_N be defined by (2) with assumptions **(W)** and **(A')**.*

Define

$$\mathbf{C}_p = \begin{cases} {}^t \text{com}(\theta I_p - A_p), & \text{if } \theta I_p - A_p \neq 0 \\ I_p & \text{else} \end{cases}, \quad (9)$$

where $\text{com}(B)$ denotes the comatrix of a matrix B , and

$$c_\rho^{(1)} = \left(1 + \sigma^2 \int \frac{d\lambda(x)}{(\rho_\theta - x)^2} \right) \text{Tr}_p(\mathbf{C}_p).$$

Then

$$d_{LP}(c_\rho^{(1)} \sqrt{N}(\lambda_{i_0}(M_N) - \rho_N), \Phi_N) \xrightarrow{N \rightarrow \infty} 0 \quad (10)$$

where $\Phi_N = \text{Tr}_p(\mathbf{C}_p W_p) + \mathcal{Z}_N$, W_p is the $p \times p$ upper left corner of the Wigner matrix W_N , \mathcal{Z}_N is a Gaussian random variable, independent from W_p , with mean 0 and variance $v_\rho(N)$, with

$$v_\rho(N) = \text{Tr}_p(\mathbf{C}_p^2) \sigma^4 \int \frac{d\lambda(x)}{(\rho_\theta - x)^2} + \frac{1}{2} (m_4 - 3\sigma^4) \kappa_N \sum_{i=1}^p ((\mathbf{C}_p)_{ii})^2,$$

$$\lambda = \mu_{sc} \boxplus \nu, \quad \kappa_N = \frac{1}{N-p} \sum_{i=1}^{N-p} (((\theta I_{N-p} - A_{N-p})^{-1})_{ii})^2.$$

In particular, if A_{N-p} is diagonal, $c_\rho^{(1)} \sqrt{N}(\lambda_{i_0}(M_N) - \rho_N)$ converges in distribution to $\text{Tr}_p(\mathbf{C}_p W_p) + \mathcal{Z}$ where \mathcal{Z} is a Gaussian random variable, independent from W_p , with mean 0 and variance v_ρ , with

$$v_\rho = \text{Tr}_p(\mathbf{C}_p^2) \sigma^4 \int \frac{d\lambda(x)}{(\rho_\theta - x)^2} + \frac{1}{2} (m_4 - 3\sigma^4) \int \frac{d\nu(x)}{(\theta - x)^2} \sum_{i=1}^p ((\mathbf{C}_p)_{ii})^2.$$

The aim of Theorems 6 and 7 below is to study the fluctuations associated to the a.s. convergence given in Proposition 2 above, for block diagonal perturbations. We first state an approximation result in distribution, in the spirit of [31] for perturbations A_N satisfying **(A')**.

Theorem 6. *Let M_N be defined by (2) with assumptions **(W)** and **(A')**. Let u_{i_0} , resp. v_{i_0} be a unit eigenvector associated to the spike θ of A_N , resp. the outlier $\lambda_{i_0}(M_N)$. Define $\tau_N(\theta)$ an approximation of $\tau(\theta)$ by*

$$\tau_N(\theta) = 1 - \sigma^2 \int \frac{1}{(\theta - x)^2} d\mu_{A_{N-p}}(x). \quad (11)$$

Then

$$d_{LP}(\sqrt{N}(|\langle u_{i_0}, v_{i_0} \rangle|^2 - \tau_N(\theta)), \Psi_N) \xrightarrow{N \rightarrow \infty} 0$$

where the r.v. Ψ_N is given by

$$\Psi_N = (P^*(c_{\theta, \sigma} W_p + Z_{p, N})P)_{11},$$

where

$$c_{\theta, \sigma} = \sigma^2 g_\nu''(\theta), \quad (12)$$

W_p is the $p \times p$ upper left corner of the Wigner matrix W_N , $Z_{p, N}$ is a centered Gaussian Hermitian matrix of size p , independent from W_p , with independent entries (modulo the symmetry conditions). The diagonal coefficients are i.i.d. with variance

$$\sigma^4 B_{\theta, \nu} + \frac{1}{2}(m_4 - 3\sigma^4)A_{\theta, \nu, N} \quad (13)$$

where

$$B_{\theta, \nu} = -\frac{1}{6}g_\nu'''(\theta) - \frac{\sigma^2}{2}(g_\nu''(\theta))^2 \frac{1 + 2\sigma^2 g_\nu'(\theta)}{1 + \sigma^2 g_\nu'(\theta)}, \quad (14)$$

$$A_{\theta, \nu, N} = \quad (15)$$

$$\frac{1}{N-p} \sum_{i=1}^{N-p} (\sigma^2 g_\nu''(\theta) [(\theta I_{N-p} - A_{N-p})^{-1}]_{ii} - (1 + \sigma^2 g_\nu'(\theta)) [(\theta I_{N-p} - A_{N-p})^{-2}]_{ii})^2$$

m_4 is the fourth moment of μ , g_ν is the Stieltjes transform of ν . The off diagonal elements $Z_{p, N}(i, j)$, $i < j$ are iid complex Gaussian with distribution Z such that $\mathbb{E}(Z^2) = 0$ and $\mathbb{E}(|Z|^2) = \sigma^4 B_{\theta, \nu}$.

In the case where the matrix A_{N-p} is a diagonal matrix, the sequence $(A_{\theta, \nu, N})_N$ defined in (15) converges as N tends to infinity. This leads to the following fluctuations result :

Theorem 7. Let M_N be defined by (2) with assumptions **(W)**, **(A')** and A_{N-p} diagonal.

Let u_{i_0} , resp. v_{i_0} be a unit eigenvector associated to the spike θ of A_N , resp. to the outlier $\lambda_{i_0}(M_N)$. Then,

$$\sqrt{N}(|\langle u_{i_0}, v_{i_0} \rangle|^2 - \tau_N(\theta)) \xrightarrow[N \rightarrow \infty]{(law)} (P^*(c_{\theta, \sigma} W_p + Z_p)P)_{11} \quad (16)$$

where $\tau_N(\theta)$ is defined by (11). W_p is the $p \times p$ upper left corner of the Wigner matrix W_N , Z_p is a centered Gaussian Hermitian matrix of size p , independent from W_p , with independent entries (modulo the symmetry condition). The diagonal coefficients are i.i.d. with variance

$$\frac{1}{2}(m_4 - 3\sigma^4)A_{\theta, \nu} + \sigma^4 B_{\theta, \nu} \quad (17)$$

and the off diagonal elements are iid complex Gaussian with distribution Z such that $\mathbb{E}(Z^2) = 0$ and $\mathbb{E}(|Z|^2) = \sigma^4 B_{\theta, \nu}$ where

$$\begin{cases} c_{\theta, \nu} = \sigma^2 g_\nu''(\theta), \\ A_{\theta, \nu} = -\frac{1}{6} g_\nu'''(\theta)(1 + \sigma^2 g_\nu'(\theta))^2 - 2\sigma^4 (g_\nu''(\theta))^2 g_\nu'(\theta) - \sigma^2 (g_\nu''(\theta))^2, \\ B_{\theta, \nu} = -\frac{1}{6} g_\nu'''(\theta) - \frac{\sigma^2}{2} (g_\nu''(\theta))^2 \frac{1 + 2\sigma^2 g_\nu'(\theta)}{1 + \sigma^2 g_\nu'(\theta)}, \end{cases} \quad (18)$$

m_4 is the fourth moment of μ and g_ν is the Stieltjes transform of ν .

In particular, we readily deduce the following corollary.

Corollary 1. Let M_N be defined by (2) with assumptions **(W)** and **(A')**. Assume moreover that A_N is a diagonal matrix ($p = 1$). Let u_{i_0} , resp. v_{i_0} be a unit eigenvector associated to the spike θ of A_N , resp. the outlier $\lambda_{i_0}(M_N)$. Then,

$$\sqrt{N}(|\langle u_{i_0}, v_{i_0} \rangle|^2 - \tau_N(\theta)) \xrightarrow[N \rightarrow \infty]{(law)} c_{\theta, \nu} W_{11} + Z \quad (19)$$

where Z is a centered Gaussian variable, independent from W_{11} , with variance :

$$\frac{1}{2}(m_4 - 3\sigma^4)A_{\theta, \nu} + \sigma^4 B_{\theta, \nu}. \quad (20)$$

See Eq.(18) for the definitions of $c_{\theta, \sigma}$, $A_{\theta, \nu}$, $B_{\theta, \nu}$.

Note that when the Wigner matrix is Gaussian, the choice of a diagonal matrix for A_N is not a restriction, due to the unitary invariance of the

G.U.E..

The proof of Theorem 6 relies upon a representation, through Helffer-Sjöstrand formula, of the variable $|\langle u_{i_0}, v_{i_0} \rangle|^2$ in terms of the $p \times p$ -matrix valued process $\{G_p(z), z \in \mathbb{C} \setminus \mathbb{R}\}$ where $G_p(z)$ denotes the principal submatrix of size p of the resolvent matrix $G(z) = (zI_N - M_N)^{-1}$. Then, the fluctuations of the process $\{G_p(z), z \in \mathbb{C} \setminus \mathbb{R}\}$ are analysed using Schur's formula which enables to express $\{G_p(z), z \in \mathbb{C} \setminus \mathbb{R}\}$ in terms of random sesquilinear forms. This approach is described in Section 4 where we also prove Theorem 7. Section 2 gathers preliminary results used in the proof of the main results. Therein, first we recall classical algebraic identities, Helffer-Sjöstrand's calculus, tightness criterion for random analytic process and some basic facts on free convolution with a semicircular distribution; later we establish some extension of central limit theorem for random quadratic forms and recall or deduce some results on deformed Wigner matrices from [8, 16]. In Section 3, we first establish Theorem 5, sticking to the approach of [16]. The last Section is an appendix reminding the reader about Poicaré inequality and concentration phenomenon.

For any integer number k , we will say that a matrix-valued function f_N on $\mathbb{C} \setminus \mathbb{R}$ is $O\left(\frac{1}{N^k}\right)$ if there exists a polynomial Q with nonnegative coefficients and an integer number d such that for all large N , for any z in $\mathbb{C} \setminus \mathbb{R}$,

$$\|f_N(z)\| \leq \frac{Q(|\Im z|^{-1})(|z| + 1)^d}{N^k}.$$

For a family of functions $f_N^{(i)}$, $i \in \{1, \dots, N\}^2$, we will set $f_N^{(i)} = O^{(u)}\left(\frac{1}{N^k}\right)$ if for each i , $f_N^{(i)} = O\left(\frac{1}{N^k}\right)$ and moreover one can find a bound of the norm of each $f_N^{(i)}$ as above involving a common polynomial Q and a common d , that is not depending on i .

For two sequences $(X_N)_N$ and $(Y_N)_N$ of random variables, $X_N = Y_N + o_{\mathbb{P}}(1)$ means that $X_N - Y_N \xrightarrow{N \rightarrow \infty} 0$ in probability.

2 Preliminaries

2.1 Basic identities and inequalities

For a matrix $A \in M_N(\mathbb{C})$ and I and J non empty subsets of $\{1, \dots, N\}$, we denote by $A_{I \times J}$ the submatrix of A obtained by keeping the rows with indices $i \in I$ and columns with indices $j \in J$. We set $A_I := A_{I \times I}$.

Proposition 5 (Schur inversion formula). *Let I be a non-empty subset of $\{1, \dots, N\}$ and $A \in M_N(\mathbb{C})$ such that A_I is invertible, then A is invertible if and only if $A_{I^c} - A_{I^c \times I} A_I^{-1} A_{I \times I^c}$ is invertible, in which case the following formulas hold:*

$$\begin{aligned} (A^{-1})_I &= (A_I)^{-1} + (A_I)^{-1} A_{I \times I^c} (A_{I^c} - A_{I^c \times I} A_I^{-1} A_{I \times I^c})^{-1} A_{I^c \times I} (A_I)^{-1}, \\ (A^{-1})_{I \times I^c} &= -(A_I)^{-1} A_{I \times I^c} (A_{I^c} - A_{I^c \times I} A_I^{-1} A_{I \times I^c})^{-1}, \\ (A^{-1})_{I^c \times I} &= -(A_{I^c} - A_{I^c \times I} A_I^{-1} A_{I \times I^c})^{-1} A_{I^c \times I} (A_I)^{-1}, \\ (A^{-1})_{I^c} &= (A_{I^c} - A_{I^c \times I} A_I^{-1} A_{I \times I^c})^{-1}. \end{aligned}$$

Lemma 1. *For any matrix $B \in M_N(\mathbb{C})$ and for any fixed k , we have*

$$\sum_{l=1}^N |B_{lk}|^2 \leq \|B\|^2 \quad (21)$$

(or equivalently

$$\sum_{l=1}^N |B_{kl}|^2 \leq \|B\|^2.) \quad (22)$$

Therefore, we have

$$\frac{1}{N} \sum_{k,l=1}^N |B_{kl}|^2 \leq \|B\|^2. \quad (23)$$

Proof. Note that

$$\begin{aligned} \sum_{l=1}^N |B_{lk}|^2 &= \text{Tr}_N(BE_{kk}B^*) \\ &= \text{Tr}_N(B^*BE_{kk}) \\ &\leq \|B\|^2 \text{Tr}_N(E_{kk}) = \|B\|^2. \end{aligned}$$

Now, since

$$\sum_{l=1}^N |B_{kl}|^2 = \sum_{l=1}^N |B_{kl}^*|^2 = \sum_{l=1}^N |(B^*)_{lk}|^2 \text{ and } \|B^*\| = \|B\|,$$

(21) and (22) can be deduced from each other thanks to conjugate transposition. Finally (22) readily yields (23). \square

Lemma 2. [Lemma A2 [16]] Let A and H be $m \times m$ matrices such that, for some $K > 0$,

$$\|A\| \leq K, \|H\| \leq K. \quad (24)$$

Then

$$\det(A + H) = \det(A) + \text{Tr}_m({}^t \text{com}(A)H) + \epsilon,$$

where $\text{com}(A)$ denotes the comatrix of A , and there exists a constant $C_{m,K} > 0$, only depending on m and K , such that $|\epsilon| \leq C_{m,K} \|H\|^2$.

We will often use the following obvious facts that for any $z \in \mathbb{C} \setminus \mathbb{R}$, for any $N \times N$ Hermitian matrix H ,

$$\|(zI_N - H)^{-1}\| \leq |\Im z|^{-1} \quad (25)$$

and for any probability measure ν on \mathbb{R} , the Stieltjes transform g_ν satisfies for any $z \in \mathbb{C} \setminus \mathbb{R}$, $\Im z \Im g_\nu(z) < 0$, $|g_\nu(z)| \leq |\Im z|^{-1}$, and for any $\alpha \geq 0$, any $x \in \mathbb{R}$,

$$\frac{1}{|z - \alpha g_\nu(z) - x|} \leq |\Im z|^{-1}. \quad (26)$$

2.2 Helffer-Sjöstrand's calculus

2.2.1 Helffer-Sjöstrand's representation formula

We recall Helffer-Sjöstrand's representation formula (see [9, Proposition C.1]): let $f \in C^{k+1}(\mathbb{R})$ with compact support and M be a Hermitian matrix; we have

$$f(M) = \frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(f)(z) (M - z)^{-1} d^2 z \quad (27)$$

where $d^2 z$ denotes the Lebesgue measure on \mathbb{C} ,

$$F_k(f)(x + iy) = \sum_{l=0}^k \frac{(iy)^l}{l!} f^{(l)}(x) \chi(y) \quad (28)$$

where $\chi : \mathbb{R} \rightarrow \mathbb{R}^+$ is a smooth compactly supported function such that $\chi \equiv 1$ in a neighborhood of 0, and $\bar{\partial} = \frac{1}{2}(\partial_x + i\partial_y)$.

The function $F_k(f)$ coincides with f on the real axis and is an extension to the complex plane.

Moreover

$$\bar{\partial} F_k(f)(x + iy) = \frac{1}{2} \frac{(iy)^k}{k!} f^{(k+1)}(x) \chi(y) + \frac{i}{2} \sum_{l=0}^k \frac{(iy)^l}{l!} f^{(l)}(x) \chi'(y) \quad (29)$$

Thus, since $\chi \equiv 1$ in a neighborhood of 0, we have that, in a neighborhood of the real axis,

$$\bar{\partial}F_k(f)(x + iy) = \frac{1}{2} \frac{(iy)^k}{k!} f^{(k+1)}(x) = O(|y|^k) \text{ as } y \rightarrow 0. \quad (30)$$

2.2.2 Computation of Helffer-Sjöstrand's integral

Proposition 6. *Let h be a smooth function with compact support in $(\rho_\theta - 2\delta, \rho_\theta + 2\delta)$ and satisfying $h \equiv 1$ on $[\rho_\theta - \delta, \rho_\theta + \delta]$. Let χ be a compactly supported function on $(-L, L)$, and $\chi = 1$ around 0. We denote by $D = (\rho_\theta - 2\delta, \rho_\theta + 2\delta) \times (-L, L)$.*

Let ϕ be a meromorphic function in D , with a pole in ρ_θ . Then,

$$I(\phi) := \frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial}F_k(h)(z) \phi(z) d^2z = -\text{Res}(\phi, \rho_\theta) \quad (31)$$

where $F_k(h)$ is defined in (28). $\text{Res}(\phi, \rho_\theta)$ denotes the residue of the function ϕ at the point ρ_θ .

Proof. Let ϵ small enough such that $F_k(h)(z) = 1$ for $z \in B(\rho_\theta, \epsilon)$. Set $D_\epsilon = D \setminus B(\rho_\theta, \epsilon)$. ϕ is holomorphic on D_ϵ . Since $F_k(h)$ has compact support in D , we have,

$$\begin{aligned} 0 &= \int_{\partial D} F_k(h)(z) \phi(z) dz = \int_{\partial D_\epsilon} F_k(h)(z) \phi(z) dz + \int_{\partial B(\rho_\theta, \epsilon)} F_k(h)(z) \phi(z) dz \\ &= 2i \int_{D_\epsilon} \bar{\partial}F_k(h)(z) \phi(z) d^2z + \int_{\partial B(\rho_\theta, \epsilon)} \phi(z) dz \end{aligned}$$

where the first term is obtained by Green's formula using that $\bar{\partial}\phi(z) = 0$ on D_ϵ .

$$\int_{D_\epsilon} \bar{\partial}F_k(h)(z) \phi(z) d^2z \xrightarrow{\epsilon \rightarrow 0} \pi I(\phi)$$

and

$$\int_{\partial B(\rho_\theta, \epsilon)} \phi(z) dz = 2i\pi \text{Res}(\phi, \rho_\theta).$$

□

2.3 Tightness criterion for a sequence of random analytic processes

We recall here some results from [34]. Let $D \subset \mathbb{C}$ be an open set in the complex plane. Denote by $\mathcal{H}(D)$ the space of complex analytic functions

in D , endowed with the uniform topology on compact set. For $f \in \mathcal{H}(D)$ and K a compact set of D , we denote $\|f\|_K = \sup_{z \in K} |f(z)|$. The space $\mathcal{H}(D)$ is equipped with the (topological) Borel σ -field $\mathcal{B}(\mathcal{H}(D))$ and the set of probability measures on $(\mathcal{H}(D); \mathcal{B}(\mathcal{H}(D)))$ is denoted by $\mathcal{P}(\mathcal{H}(D))$. By a random analytic function on D we mean an $\mathcal{H}(D)$ -valued random variable on a probability space. The probability law of a random analytic function is uniquely determined by its finite dimensional distributions.

Proposition 7. (*Proposition 2.5. in [34]*) *Let f_n be a sequence of random analytic functions in D . If $\|f_n\|_K$ is tight for any compact set K , then $\mathcal{L}(f_n)$ is tight in $\mathcal{P}(\mathcal{H}(D))$.*

Using that, by Markov's inequality, for any $C > 0$ and any $r > 0$,

$$\mathbb{P}(\|f_n\|_K > C) \leq \frac{1}{C^r} \mathbb{E}(\|f_n\|_K^r), \quad (32)$$

the following lemma turns out to be useful to prove tightness results.

Lemma 3. (*lemma 2.6 [34]*) *For any compact set K in D , there exists $\delta > 0$ such that*

$$\|f\|_K^r \leq (\pi\delta^2)^{-1} \int_{\overline{K_\delta}} |f(z)|^r m(dz), \quad f \in \mathcal{H}(D),$$

for any $r > 0$, where $\overline{K_\delta} \subset D$ is the closure of the δ -neighborhood of K and m denotes the Lebesgue measure.

2.4 Free convolution with a semicircular distribution

Let τ be a probability measure on \mathbb{R} . Its Stieltjes transform $g_\tau : z \mapsto \int_{\mathbb{R}} \frac{1}{z-x} d\tau(x)$ is analytic on the complex upper half-plane \mathbb{C}^+ . There exists a domain

$$D_{\alpha,\beta} = \{u + iv \in \mathbb{C}, |u| < \alpha v, v > \beta\}$$

on which g_τ is univalent. Let K_τ be its inverse function, defined on $g_\tau(D_{\alpha,\beta})$, and

$$R_\tau(z) = K_\tau(z) - \frac{1}{z}.$$

Given two probability measures τ and ν , there exists a unique probability measure λ such that

$$R_\lambda = R_\tau + R_\nu$$

on a domain where these functions are defined. The probability measure λ is called the free convolution of τ and ν and denoted by $\tau \boxplus \nu$.

The free convolution of probability measures has an important property, called subordination, which can be stated as follows: let τ and ν be two probability measures on \mathbb{R} ; there exists an analytic map $\omega_{\tau,\nu} : \mathbb{C}^+ \rightarrow \mathbb{C}^+$ such that

$$\forall z \in \mathbb{C}^+, \quad g_{\tau \boxplus \nu}(z) = g_\nu(\omega_{\tau,\nu}(z)).$$

This phenomenon was first observed by D. Voiculescu under a genericity assumption in [38], and then proved in generality in [11] Theorem 3.1. Later, a new proof of this result was given in [7], using a fixed point theorem for analytic self-maps of the upper half-plane.

In [10], P. Biane provides a deep study of the free convolution by a semicircular distribution. We first recall here some of his results that will be useful in our approach. Let ν be a probability measure on \mathbb{R} and μ_{sc} the semicircular distribution defined by (1). For any $z \in \mathbb{C} \setminus \text{supp}(\mu_{sc} \boxplus \nu)$, the subordination function $\omega_{\mu_{sc},\nu}$ is given by

$$\omega_{\mu_{sc},\nu}(z) = z - \sigma^2 g_{\mu_{sc} \boxplus \nu}(z). \quad (33)$$

In the following, we will denote $\omega_{\mu_{sc},\nu}$ by ω . For any $x \in \mathbb{C} \setminus \text{supp}(\nu)$, define

$$H(x) = x + \sigma^2 g_\nu(x). \quad (34)$$

We have for any $x \in \mathbb{C} \setminus \text{supp}(\mu_{sc} \boxplus \nu)$,

$$H(\omega(x)) = x. \quad (35)$$

More precisely, the following one to one correspondance holds:

$$\mathbb{R} \setminus \text{supp}(\mu_{sc} \boxplus \nu) \xrightleftharpoons[H]{\omega} \left\{ u \in \mathbb{R} \setminus \text{supp} \nu, \int \frac{1}{(u-x)^2} d\nu(x) < \frac{1}{\sigma^2} \right\}. \quad (36)$$

Assume that A_N satisfies **(A)**. Let $\{\theta_j, 1 \leq j \leq q\}$ be the spiked eigenvalue of A_N outside $\text{supp}(\nu)$. Furthermore, for all $\theta_j \in \Theta_{\sigma,\nu}$, where $\Theta_{\sigma,\nu}$ is defined by (3), we set

$$\rho_{\theta_j} := H(\theta_j) = \theta_j + \sigma^2 g_\nu(\theta_j) \quad (37)$$

which is outside the support of $\mu_{sc} \boxplus \nu$ according to (36), and we define

$$K_{\sigma,\nu}(\theta_1, \dots, \theta_q) := \text{supp}(\mu_{sc} \boxplus \nu) \cup \{\rho_{\theta_j}, \theta_j \in \Theta_{\sigma,\nu}\}.$$

An important consequence of (36) is the following

Proposition 8. [Theorem 2.3 [21]] For any $\delta > 0$,

$$\text{supp}(\mu_{sc} \boxplus \mu_{A_N}) \subset K_{\sigma,\nu}(\theta_1, \dots, \theta_q) + (-\delta, \delta),$$

when N is large enough.

2.5 Central limit theorem for processes of matrix valued random quadratic forms

Lemma 4 (cf Lemma 2.7 [2]). *There exists $C > 0$ such that for any $N \times N$*

deterministic matrix $B = (b_{ij})_{1 \leq i, j \leq N}$, any random vectors $Y = \begin{pmatrix} y_1 \\ \vdots \\ y_N \end{pmatrix}$ in

\mathbb{C}^N with i.i.d. standardized entries ($\mathbb{E}(y_i) = 0$, $\mathbb{E}(|y_i|^2) = 1$, $\mathbb{E}(y_i^2) = 0$) such that $\mathbb{E}(|y_i|^4) < \kappa$ and any independent copy X of Y , one has

$$\mathbb{E}(|Y^* B Y - \text{Tr}_N(B)|^2) \leq C \kappa \text{Tr}_N(B^* B),$$

$$\mathbb{E}(|Y^* B X|^2) \leq C \kappa \text{Tr}_N(B^* B).$$

In [31], the authors establish the following variation around the central limit theorem for martingales.

Lemma 5 (Lemma 5.6 [31]). *Suppose that for each n ($Y_{nj}; 1 \leq j \leq r_n$) is a \mathbb{C}^d -valued martingale difference sequence with respect to the increasing σ -field $\{\mathcal{G}_{n,j}; 1 \leq j \leq r_n\}$ having second moments. Write*

$$Y_{nj}^T = (Y_{nj}^1, \dots, Y_{nj}^d).$$

Assume moreover that $(\Theta_n(k, l))_n$ and $(\tilde{\Theta}_n(k, l))_n$ are uniformly bounded sequences of complex numbers, for $1 \leq k, l \leq d$. If

$$\sum_{j=1}^{r_n} \mathbb{E} \left(Y_{nj}^k \bar{Y}_{nj}^l \mid \mathcal{G}_{n,j-1} \right) - \Theta_n(k, l) \xrightarrow[n \rightarrow +\infty]{P} 0, \quad (38)$$

$$\sum_{j=1}^{r_n} \mathbb{E} \left(Y_{nj}^k Y_{nj}^l \mid \mathcal{G}_{n,j-1} \right) - \tilde{\Theta}_n(k, l) \xrightarrow[n \rightarrow +\infty]{P} 0, \quad (39)$$

and for each $\epsilon > 0$, $\sum_{j=1}^{r_n} \mathbb{E} \left(\|Y_{nj}\|^2 \mathbf{1}_{\|Y_{nj}\| > \epsilon} \right) \rightarrow_{n \rightarrow +\infty} 0$, then, for every bounded continuous function $f : \mathbb{C}^d \rightarrow \mathbb{R}$,

$$\mathbb{E} f \left(\sum_{j=1}^{r_n} Y_{nj} \right) - \mathbb{E} f(Z_n) \xrightarrow[n \rightarrow +\infty]{} 0, \quad (40)$$

where Z_n is a \mathbb{C}^d -valued centered Gaussian random vector with parameters

$$\mathbb{E}(Z_n Z_n^*) = (\Theta_n(k, l))_{k,l} \text{ and } \mathbb{E}(Z_n Z_n^T) = (\tilde{\Theta}_n(k, l))_{k,l}.$$

Following the lines of the proof of the central limit theorem for quadratic forms by Baik and Silverstein in the appendix of [18] and using Lemma 5, we will establish the following extension.

Proposition 9. *Let (z_1, \dots, z_q) be in I^q , where I is a subset of \mathbb{C} such that $\forall z \in I, \bar{z} \in I$. For any z in $\{z_1, \dots, z_q\}$, let $B(z) = (b_{ij}(z))$ be a $N \times N$ matrix such that $(B(z))^* = B(\bar{z})$ and there exists a constant $a > 0$ (not depending on N) such that for any z in $\{z_1, \dots, z_q\}$, $\|B(z)\| \leq a$. Let p be a fixed integer number and $Y_N = (y_{ij})_{1 \leq i \leq N, 1 \leq j \leq p}$ be a $N \times p$ matrix which contains i.i.d. complex standardized entries with bounded fourth moment and such that $\mathbb{E}(y_{11}^2) = 0$. Set*

$$V_N = \left((1/\sqrt{N})(Y_N^* B(z_1) Y_N - \text{Tr}_N(B(z_1)) I_p), \dots, (1/\sqrt{N})(Y_N^* B(z_q) Y_N - \text{Tr}_N(B(z_q)) I_p) \right).$$

If there exists uniformly bounded sequences $(f_N)_N$ and $(g_N)_N$ of functions on I^2 such that for any z_l, z_k ,

$$\frac{1}{N} \sum_{i=1}^N (B(z_l))_{ii} (B(z_k))_{ii} = f_N(z_l, z_k) + o(1),$$

and

$$\text{tr}_N(B(z_l) B(z_k)) = g_N(z_l, z_k) + o(1),$$

then $d_{LP}(V_N, \mathcal{V}_N) \rightarrow 0$ where $\mathcal{V}_N = (\mathcal{G}_N(z_1), \dots, \mathcal{G}_N(z_q))$, \mathcal{G}_N is a centered Gaussian matrix valued process whose distribution is given as follows :

- 1) the processes $((\mathcal{G}_N(z))_{ij})_z$ for $1 \leq i \leq j \leq p$ are independent, and $(\mathcal{G}_N(z))_{ji} = \overline{(\mathcal{G}_N(\bar{z}))_{ij}}$.
- 2) For $i \leq p$,

$$\begin{aligned} \mathbb{E}((\mathcal{G}_N(z_l))_{ii} (\mathcal{G}_N(z_k))_{ii}) &= (\mathbb{E}(|y_{11}|^4) - 2) (f_N(z_l, z_k)) \\ &\quad + g_N(z_l, z_k) \end{aligned}$$

and

$$\mathbb{E}((\mathcal{G}_N(z_l))_{ii} \overline{(\mathcal{G}_N(z_k))_{ii}}) = \mathbb{E}((\mathcal{G}_N(z_1))_{ii} (\mathcal{G}_N(\bar{z}_2))_{ii}) \quad (41)$$

- 2) For $1 \leq i \neq j \leq p$,

$$\mathbb{E}((\mathcal{G}_N(z_l))_{ij} (\mathcal{G}_N(z_k))_{ij}) = 0$$

and

$$\mathbb{E}((\mathcal{G}_N(z_l))_{ij} \overline{(\mathcal{G}_N(z_k))_{ij}}) = g_N(z_l, \bar{z}_k). \quad (42)$$

Proof. First, for any B in $\{B(z), z = z_1, \dots, z_q\}$ and any $1 \leq s, t \leq p$, one can write $(1/\sqrt{N})(Y_N^* B Y_N - \text{Tr}(B)I_p)_{st}$ as a sum of martingale differences:

$$\begin{aligned}
& (1/\sqrt{N})(Y_N^* B Y_N - \text{Tr}(B)I_p)_{st} \\
&= (1/\sqrt{N}) \sum_{i=1}^N \left((\bar{y}_{is} y_{it} - \delta_{st}) b_{ii} + \bar{y}_{is} \sum_{j<i} y_{jt} b_{ij} + \bar{y}_{is} \sum_{j>i} y_{jt} b_{ij} \right) \\
&= (1/\sqrt{N}) \sum_{i=1}^N \left((\bar{y}_{is} y_{it} - \delta_{st}) b_{ii} + \bar{y}_{is} \sum_{j<i} y_{jt} b_{ij} + y_{it} \sum_{j<i} \bar{y}_{js} b_{ji} \right) \\
&= \sum_{i=1}^N (Z_i(B))_{st}
\end{aligned}$$

where

$$(Z_i(B))_{st} = (1/\sqrt{N}) \left((\bar{y}_{is} y_{it} - \delta_{st}) b_{ii} + \bar{y}_{is} \sum_{j<i} y_{jt} b_{ij} + y_{it} \sum_{j<i} \bar{y}_{js} b_{ji} \right).$$

Let $\mathcal{F}_{N,i}$ be the σ -field generated by $\{y_{1s}, \dots, y_{is}, 1 \leq s \leq p\}$. Let also $\mathbb{E}_i(\cdot)$ denote conditional expectation with respect to $\mathcal{F}_{N,i}$. It is clear that for any B in $\{B(z), z \in I\}$, $Z_i(B) \in M_p(\mathbb{C}) \simeq \mathbb{C}^{p^2}$ is measurable with respect to $\mathcal{F}_{N,i}$ and satisfies $\mathbb{E}_{i-1}(Z_i(B)) = 0$.

We will show that the conditions of Lemma 5 are met for the \mathbb{C}^{qp^2} -valued martingale difference sequence $Y_{N,i}^T = (Z_i(B(z_1)), \dots, Z_i(B(z_q)))$. Write $(Z_i(B))_{st} = X_1^i + X_2^i$, with $X_1^i = (1/\sqrt{N})(\bar{y}_{is} y_{it} - \delta_{st}) b_{ii}$. Then for $\epsilon > 0$,

$$\sum_{i=1}^N \mathbb{E}(|X_1^i|^2 \mathbf{1}_{(|X_1^i| \geq \epsilon)}) \leq a^2 \mathbb{E}(|\bar{y}_{1s} y_{1t} - \delta_{st}|^2 \mathbf{1}_{\{|\bar{y}_{1s} y_{1t} - \delta_{st}|^2 \geq \sqrt{N}\epsilon/a\}}) \rightarrow 0 \quad (43)$$

as $N \rightarrow \infty$, by dominated convergence theorem.

We have

$$\begin{aligned}
& \mathbb{E} \left| \sum_{j<i} y_{jt} b_{ij} \right|^4 \\
&= \mathbb{E}(|y_{1t}|^4 \sum_{j<i} |b_{ij}|^4) + 2\mathbb{E} \left(\sum_{j_1 < j_2} |b_{ij_1}|^2 |b_{ij_2}|^2 \right) + \mathbb{E}(|y_{1t}|^2 \sum_{j_1 < j_2} |b_{ij_1}|^2 |b_{ij_2}|^2) \\
&\leq \mathbb{E}|y_{1t}|^4 \mathbb{E}[(\max_{i,j} |b_{ij}|)^2 \max_i (BB^*)_{ii}] + (2 + \mathbb{E}|y_{1t}|^2) \mathbb{E}[(BB^*)_{ii}^2]
\end{aligned}$$

$$\leq a^4 [\mathbb{E}|y_{1t}|^4 + 2 + \mathbb{E}|y_{1t}^2|^2]$$

where the sum \sum_* is over $\{j_1 < i, j_2 < i, j_1 \neq j_2\}$. Therefore $\mathbb{E}|X_2^i|^4 = o(N^{-1})$ so that for any $\epsilon > 0$,

$$\sum_{i=1}^N \mathbb{E}(|X_2^i|^2 1_{(|X_2^i| \geq \epsilon)}) \leq (1/\epsilon^2) \sum_{i=1}^N \mathbb{E}|X_2^i|^4 \rightarrow 0 \quad \text{as } N \rightarrow \infty. \quad (44)$$

Thus, by (43), (44) and (A.4) in [18], $\{Z_i(B)\}$ satisfies the Lindeberg condition of Lemma 5.

Now, we shall verify condition (39) of Lemma 5. We have for any B and C in $\{B(z), z \in I\}$, for any $1 \leq s, t, s', t' \leq p$,

$$\sum_{i=1}^N \mathbb{E}_{i-1}(Z_i(B))_{st} Z_i(C)_{s't'} \quad (45)$$

$$= (1/N) \sum_{i=1}^N \left\{ (\mathbb{E}|y_{11}|^4 - 1) \delta_{st} \delta_{ss'} \delta_{s't'} + \delta_{s't'} \delta_{s't} (1 - \delta_{ss'}) \right\} b_{ii} c_{ii} \quad (46)$$

$$\begin{aligned} & + \mathbb{E}(|y_{11}|^2 \bar{y}_{11}) \left[\delta_{st} \delta_{ss'} \sum_{j < i} y_{jt'} c_{ij} b_{ii} + \delta_{s't'} \delta_{ss'} \sum_{j < i} y_{jt} b_{ij} c_{ii} \right] \\ & + \mathbb{E}(|y_{11}|^2 y_{11}) \left[\delta_{st} \delta_{tt'} \sum_{j < i} \bar{y}_{js'} c_{ji} b_{ii} + \delta_{s't'} \delta_{tt'} \sum_{j < i} \bar{y}_{js} b_{ji} c_{ii} \right] \quad (47) \\ & + \delta_{s't} \sum_{j < i} \bar{y}_{js} b_{ji} \sum_{j < i} y_{jt'} c_{ij} + \delta_{s't'} \sum_{j < i} \bar{y}_{js'} c_{ji} \sum_{j < i} y_{jt} b_{ij} \}. \end{aligned}$$

Let B_L (resp. B_U) denote the strictly lower (resp. upper) triangular part of

B. We have using Cauchy-Schwarz's inequality that

$$\begin{aligned}
\mathbb{E} \left| (1/N) \sum_{i=1}^N c_{ii} \sum_{j<i} y_{jt} b_{ij} \right|^2 &= \mathbb{E} \left| (1/N) \sum_{j=1}^{N-1} y_{jt} \sum_{i>j} c_{ii} b_{ij} \right|^2 \\
&= (1/N^2) \mathbb{E} \left(\sum_{j=1}^{N-1} \sum_{i>j} c_{ii} b_{ij} \sum_{i>j} \bar{c}_{ii} \bar{b}_{ij} \right) \\
&= (1/N^2) \mathbb{E} \left(\sum_{\underline{ii}} c_{ii} \bar{c}_{ii} (B_L B_L^*)_{\underline{ii}} \right) \\
&\leq \mathbb{E} \left[(\max_i |c_{ii}|)^2 (1/N) \left(\sum_{\underline{ii}} |(B_L B_L^*)_{\underline{ii}}|^2 \right)^{1/2} \right] \\
&= \mathbb{E} \left[(\max_i |c_{ii}|)^2 (1/N) \text{Tr}((B_L B_L^*)^2)^{1/2} \right] \\
&\leq \mathbb{E} \left[(\max_i |c_{ii}|)^2 (1/\sqrt{N}) \|B_L\|^2 \right].
\end{aligned}$$

We apply the following bound (due to R. Mathias, see [29]): $\|B_L\| \leq \gamma_N \|B\|$ where $\gamma_N = O(\ln N)$, and the bounds $\|B\| \leq a$, $\|C\| \leq a$, to conclude that

$$(1/N) \sum_{i=1}^N c_{ii} \sum_{j<i} y_{jt} b_{ij} \xrightarrow[N \rightarrow +\infty]{P} 0.$$

Thus,

$$\begin{aligned}
&\sum_{i=1}^N \mathbb{E}_{i-1} (Z_i(B))_{st} Z_i(C)_{s't'} \\
&= (1/N) \sum_{i=1}^N \left\{ [(\mathbb{E}|y_{11}|^4 - 1) \delta_{st} \delta_{ss'} \delta_{s't'} + \delta_{st'} \delta_{s't} (1 - \delta_{ss'})] b_{ii} c_{ii} \right. \\
&\quad \left. + \delta_{s't} \sum_{j<i} \bar{y}_{js} b_{ji} \sum_{j<i} y_{jt'} c_{ij} + \delta_{st'} \sum_{j<i} \bar{y}_{j's'} c_{ji} \sum_{j<i} y_{jt} b_{ij} \right\} + o_{\mathbb{P}}(1) \\
&= [(\mathbb{E}|y_{11}|^4 - 1) \delta_{st} \delta_{ss'} \delta_{s't'} + \delta_{st'} \delta_{s't} (1 - \delta_{ss'})] \frac{1}{N} \sum_{i=1}^N b_{ii} c_{ii} \\
&\quad + \delta_{s't} (1/N) (Y_N)_s^* B_U C_L (Y_N)_{t'} + \delta_{st'} (1/N) (Y_N)_{s'}^* C_U B_L (Y_N)_t + o_{\mathbb{P}}(1)
\end{aligned}$$

Besides, from Lemma 4 we have

$$\begin{aligned}
\mathbb{E} \left| (1/N) ((Y_N)_s^* B_U C_L (Y_N)_{t'} - \delta_{st'} \text{Tr}(B_U C_L)) \right|^2 &\leq (1/N^2) \mathbb{E} (\text{Tr}(C_L^* B_U^* B_U C_L)) \\
&\leq K \mathbb{E} \|B\|^2 \|C\|^2 \frac{\ln^4 N}{N} \rightarrow_{N \rightarrow \infty} 0.
\end{aligned}$$

Hence,

$$\begin{aligned}
& \sum_{i=1}^N \mathbb{E}_{i-1}(Z_i(B))_{st}(Z_i(C))_{s't'} \\
&= [(\mathbb{E}|y_{11}|^4 - 1)\delta_{st}\delta_{ss'}\delta_{s't'} + \delta_{st'}\delta_{s't}(1 - \delta_{ss'})] \frac{1}{N} \sum_{i=1}^N b_{ii}c_{ii} \\
&\quad + \delta_{st'}\delta_{s't} \frac{1}{N} \sum_{j<i} b_{ij}c_{ji} + \delta_{s't}\delta_{st'} \frac{1}{N} \sum_{j<i} b_{ji}c_{ij} + o_{\mathbb{P}}(1) \\
&= [(\mathbb{E}|y_{11}|^4 - 1)\delta_{st}\delta_{ss'}\delta_{s't'} + \delta_{st'}\delta_{s't}(1 - \delta_{ss'})] \frac{1}{N} \sum_{i=1}^N b_{ii}c_{ii} \\
&\quad + \delta_{st'}\delta_{s't} \frac{1}{N} \text{Tr}(BC) - \delta_{s't}\delta_{st'} \frac{1}{N} \sum_i b_{ii}c_{ii} + o_{\mathbb{P}}(1) \\
&= [(\mathbb{E}|y_{11}|^4 - 1)\delta_{st}\delta_{ss'}\delta_{s't'} - \delta_{st'}\delta_{s't}\delta_{ss'}] \frac{1}{N} \sum_{i=1}^N b_{ii}c_{ii} \\
&\quad + \delta_{st'}\delta_{s't} \text{tr}_N(BC) + o_{\mathbb{P}}(1).
\end{aligned}$$

Proposition 9 readily follows. \square

2.6 Preliminary results on deformed Wigner matrices

2.6.1 Preliminary results from [8]

Note that, in Section 5 in [8], the authors consider for any fixed integer numbers m, r, t and any fixed $m \times m$ Hermitian matrices $\gamma, \alpha_1, \dots, \alpha_r, \beta_1, \dots, \beta_t$, the matrix model in $M_m(\mathbb{C}) \otimes M_N(\mathbb{C})$, $\gamma \otimes I_N + \sum_{v=1}^r \alpha_v \otimes \frac{W_N^{(v)}}{\sqrt{N}} + \sum_{u=1}^t \beta_u \otimes A_N^{(u)}$ where the $W_N^{(v)}$'s are independent more general Wigner matrices than ours and the $A_N^{(u)}$'s are deterministic matrices such that $\sup_N \|A_N^{(u)}\| < \infty$. Therefore, the results therein apply to our model by choosing $m = 1, r = t = 1$ and $\alpha_1 = 1 = \beta_1$.

For any $z \in \mathbb{C} \setminus \mathbb{R}$, $G(z) = [G_{ij}(z)]_{1 \leq i, j \leq N} = (zI_N - M_N)^{-1}$ denotes the resolvent of M_N . Note that by (25),

$$\|G(z)\| \leq |\Im z|^{-1}. \quad (48)$$

For any $z \in \mathbb{C} \setminus \mathbb{R}$, define $g_N(z) = \mathbb{E}(\text{tr}_N G(z))$ and denote by $\tilde{g}_N(z)$ the Stieltjes transform of $\mu_{sc} \boxplus \mu_{A_N}$.

Lemma 6. For any $(i, j) \in \{1, \dots, N\}^2$,

$$\mathbb{E}(G_{ij}(z)) = \left[((z - \sigma^2 \tilde{g}_N(z))I_N - A_N)^{-1} \right]_{ij} + O^{(u)}\left(\frac{1}{\sqrt{N}}\right).$$

Proof. According to Corollary 5.5 in [8], for any $(i, j) \in \{1, \dots, N\}^2$,

$$\begin{aligned} \mathbb{E}(G_{ij}(z)) &= (Y_N(z))_{ij} \\ &+ \frac{(1 - \sqrt{-1})\kappa_3}{2\sqrt{2N}\sqrt{N}} \sum_{s,l=1}^N (Y_N(z))_{il} (Y_N(z))_{ss} (Y_N(z))_{ll} \mathbb{E}(G_{sj}(z)) \\ &+ O_{ij}^{(u)}\left(\frac{1}{N}\right) \end{aligned}$$

where

$$Y_N(z) = ((z - g_N(z))I_N - A_N)^{-1}$$

and κ_3 is the third classical cumulant of μ . Note that $|\Im(z - g_N(z))| \geq |\Im z|$ so that, by (25)

$$\|Y_N(z)\| \leq \frac{1}{|\Im z|}. \quad (49)$$

Now,

$$\begin{aligned} &\left| \sum_{s,l=1}^N (Y_N(z))_{il} (Y_N(z))_{ss} (Y_N(z))_{ll} \mathbb{E}(G_{sj}(z)) \right| \\ &\leq \sum_{s,l=1}^N |(Y_N(z))_{il}| |(Y_N(z))_{ss}| |(Y_N(z))_{ll}| |\mathbb{E}(G_{sj}(z))| \\ &\leq |\Im z|^{-2} N \left(\sum_{s=1}^N |\mathbb{E}(G_{sj}(z))|^2 \right)^{1/2} \left(\sum_{l=1}^N |(Y_N(z))_{il}|^2 \right)^{1/2} \\ &\leq N |\Im z|^{-4} \end{aligned}$$

where we used Lemma 1, (49) and (48). Therefore

$$\mathbb{E}(G_{ij}(z)) = (Y_N(z))_{ij} + O_{ij}^{(u)}\left(\frac{1}{\sqrt{N}}\right).$$

Now, according to (5.56) in [8],

$$\|Y_N(z) - \tilde{Y}_N(z)\| = O\left(\frac{1}{\sqrt{N}}\right)$$

where

$$\tilde{Y}_N(z) = ((z - \tilde{g}_N(z))I_N - A_N)^{-1}. \quad (50)$$

Lemma 6 readily follows. \square

Lemma 7. *We have*

$$\forall z \in \mathbb{C} \setminus \mathbb{R}, g_N(z) = \tilde{g}_N(z) + O\left(\frac{1}{N}\right).$$

Proof. According to Proposition 5.8 in [8], we have

$$g_N(z) - \tilde{g}_N(z) = \left(1 - \tilde{g}'_N(z)\right) \tilde{L}_N(z) + O\left(\frac{1}{N\sqrt{N}}\right)$$

where

$$\begin{aligned} \tilde{L}_N(z) &= \frac{\kappa_4}{2N^3} \sum_{i,l=1}^N \left(\tilde{Y}_N(z)^2\right)_{ll} \left[\left(\tilde{Y}_N(z)\right)_{ii}\right]^2 \left(\tilde{Y}_N(z)\right)_{ll} \\ &+ \frac{\kappa_3(1 + \sqrt{-1})}{2\sqrt{2}N^2\sqrt{N}} \sum_{i,l=1}^N \left(\tilde{Y}_N(z)^2\right)_{ll} \left(\tilde{Y}_N(z)\right)_{ii} \left(\tilde{Y}_N(z)\right)_{li} \\ &+ \frac{\kappa_3(1 - \sqrt{-1})}{2\sqrt{2}N^2\sqrt{N}} \sum_{i,l=1}^N \left(\tilde{Y}_N(z)^2\right)_{il} \left(\tilde{Y}_N(z)\right)_{ii} \left(\tilde{Y}_N(z)\right)_{ll} \\ &+ \frac{\kappa_3(1 - \sqrt{-1})}{2\sqrt{2}N^2\sqrt{N}} \sum_{i,l=1}^N \left(\tilde{Y}_N(z)^2\right)_{ll} \left(\tilde{Y}_N(z)\right)_{il} \left(\tilde{Y}_N(z)\right)_{ii}, \end{aligned}$$

and \tilde{Y}_N is defined by (50). (Note that in Proposition 5.8 in [8], in full generality the $\left(\tilde{Y}_N(z)\right)_{il}$'s are $m \times m$ matrices which a priori do not commute and $\tilde{G}_N(zI_m)$ is a $m \times m$ matrix too. But in the present paper, since $m = 1$, $\left(\tilde{Y}_N(z)\right)_{il}$'s are scalar and obviously commute and $\tilde{G}_N(zI_m) = \tilde{g}_N(z)$.) Note that, $|\Im(z - \tilde{g}_N(z))| \geq |\Im z|$ so that, by (25),

$$\|\tilde{Y}_N(z)\| \leq |\Im z|^{-1}.$$

Lemma 7 readily follows by using Cauchy-Schwarz's inequality and (23). \square

Lemma 8.7 in [8] implies in particular the following variance estimates.

Lemma 8.

$$\text{Var}(G_{ij}(z)) = O^{(u)}(1/N).$$

Lemma 9.

$$\text{Var}(\text{tr}_N G(z)) = O(1/N^2)$$

The following result is a corollary of Theorem 1.1 in [8].

Proposition 10. [Theorem 1.1 [8]] *Let $[b; c]$ be a real interval such that there exists $\delta > 0$ such that, for any large N , $[b + \delta; c + \delta]$ lies outside the support of $\mu_{sc} \boxplus \mu_{A_N}$. Then, almost surely, for all large N , there is no eigenvalue of M_N in $[b; c]$.*

2.6.2 Quantitative asymptotic freeness

Let $\epsilon_0 > 0$ be fixed such that $d(\rho_\theta, \text{supp}(\mu_{sc} \boxplus \nu) \cup \{\rho_{\theta_j}, \theta_j \neq \theta\}) > \epsilon_0$ and $d(\theta, \text{supp}(\nu) \cup \{\theta_j, \theta_j \neq \theta\}) > \epsilon_0$. Let Ω_N be the event on which there is no eigenvalue of $\frac{W_{N-p}}{\sqrt{N}} + A_{N-p}$ in $]\rho_\theta - \epsilon_0; \rho_\theta + \epsilon_0[$, $\lambda_{i_0}(M_N)$ is the unique eigenvalue of M_N in $]\rho_\theta - \epsilon_0/2; \rho_\theta + \epsilon_0/2[$ and $\left\| \frac{W_N}{\sqrt{N}} \right\| \leq 3$. Propositions 8, 10 applied to M_{N-p} , Proposition 1 and Bai-Yin's theorem lead that

$$\lim_{N \rightarrow +\infty} \mathbf{1}_{\Omega_N} = 1, \text{ a.s.} \quad (51)$$

Denoting by \hat{G}_{N-p} the resolvent of the lower right submatrix of size $N - p$ of M_N and ρ_N being defined by (7), we have the following

Proposition 11.

$$\sqrt{N} \left\{ \text{tr}_{N-p} \hat{G}_{N-p}(\rho_N) \mathbf{1}_{\Omega_N} - \int \frac{d\mu_{sc} \boxplus \mu_{A_{N-p}}(x)}{(\rho_N - x)} \right\}$$

goes to zero in probability.

Proof. We stick to the proof of Proposition 5.5 in [16]. Using Proposition 8, for N large enough,

$$d(\rho_N, \text{supp}(\mu_{sc} \boxplus \mu_{A_{N-p}})) > \epsilon_0/2 \quad (52)$$

and on Ω_N , $d(\{\rho_N, \lambda_{i_0}(M_N)\}, \text{spect}\left(\frac{W_{N-p}}{\sqrt{N}} + A_{N-p}\right)) > \epsilon_0/2$, so that

$$\left\| \hat{G}_{N-p}(\rho_N) \right\| \leq \frac{2}{\epsilon_0}, \quad \left\| \hat{G}_{N-p}(\lambda_{i_0}(M_N)) \right\| \leq \frac{2}{\epsilon_0}. \quad (53)$$

Moreover, there exists $K > 0$ such that for any $x \in \text{supp}(\mu_{sc} \boxplus \mu_{A_{N-p}})$,

$$|\rho_N - x| \leq K \text{ and on } \Omega_N, \left\| \left(\rho_N I_{N-p} - \frac{W_{N-p}}{\sqrt{N}} - A_{N-p} \right) \right\| \leq K.$$

Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a \mathcal{C}^∞ function with support in $\{\epsilon_0/4 \leq |x| \leq 2K\}$ and such that $g \equiv 1$ on $\{\epsilon_0/2 \leq |x| \leq K\}$. $f : x \mapsto \frac{g(x)}{x}$ is a \mathcal{C}^∞ function with compact support. Note that

$$\int \frac{d\mu_{sc} \boxplus \mu_{A_{N-p}}(x)}{(\rho_N - x)} = \int f(\rho_N - x) d\mu_{sc} \boxplus \mu_{A_{N-p}}(x) \quad (54)$$

$$\text{and on } \Omega_N, \hat{G}_{N-p}(\rho_N) = f\left(\rho_N I_{N-p} - \frac{W_{N-p}}{\sqrt{N}} - A_{N-p}\right). \quad (55)$$

According to Lemma 7, for any $z \in \mathbb{C} \setminus \mathbb{R}$,

$$\sqrt{N} \operatorname{tr}_{N-p} \mathbb{E} \left[\hat{G}_{N-p}(\rho_N - z) \right] = \sqrt{N} \frac{d\mu_{sc} \boxplus \mu_{A_{N-p}}(x)}{(\rho_N - z - x)} + o^{(z)}(1), \quad (56)$$

where there exist polynomials Q_1 and Q_2 with non negative coefficients and $(d, k) \in \mathbb{N}^2$ such that

$$\|o^{(z)}(1)\| \leq \frac{Q_1(|\Im z|^{-1})(|z| + 1)^d}{\sqrt{N}} \leq \frac{1}{\sqrt{N}} \frac{Q_2(|\Im z|)(|z| + 1)^d}{|\Im z|^k}. \quad (57)$$

Therefore, by Helffer-Sjöstrand functional calculus (see Section 2.2),

$$\begin{aligned} & \sqrt{N} \operatorname{tr}_{N-p} \mathbb{E} \left(f \left(\rho_N I_{N-p} - \frac{W_{N-p}}{\sqrt{N}} - A_{N-p} \right) \right) \\ &= \frac{1}{\pi} \int_{\mathbb{C} \setminus \mathbb{R}} \bar{\partial} F_k(f)(z) \sqrt{N} \operatorname{tr}_{N-p} \mathbb{E} \left[\hat{G}_{N-p}(\rho_N - z) \right] d^2 z \end{aligned}$$

and

$$\sqrt{N} \int f((\rho_N - x) d\mu_{sc} \boxplus \mu_{A_{N-p}}(x) = \frac{1}{\pi} \int_{\mathbb{C} \setminus \mathbb{R}} \bar{\partial} F_k(f)(z) \sqrt{N} \frac{d\mu_{sc} \boxplus \mu_{A_{N-p}}(x)}{(\rho_N - z - x)} d^2 z.$$

Hence, using (56) and (54), we can deduce that

$$\begin{aligned} & \sqrt{N} \operatorname{tr}_{N-p} \mathbb{E} \left(f \left(\rho_N I_{N-p} - \frac{W_{N-p}}{\sqrt{N}} - A_{N-p} \right) \right) \\ &= \sqrt{N} \int \frac{d\mu_{sc} \boxplus \mu_{A_{N-p}}(x)}{(\rho_N - x)} + \frac{1}{\pi} \int_{z \in \mathbb{C} \setminus \mathbb{R}} \bar{\partial} F_k(f)(z) o^{(z)}(1) d^2 z. \end{aligned}$$

Note that since f and χ are compactly supported, the last integral is an integral on a bounded set of \mathbb{C} and according to (57) and (30),

$$\left\| \frac{1}{\pi} \int_{\mathbb{C} \setminus \mathbb{R}} \bar{\partial} F_k(f)(z) o^{(z)}(1) d^2 z \right\| \leq \frac{C}{\sqrt{N}}.$$

Thus,

$$\sqrt{N} \left\{ \mathbb{E} \operatorname{tr}_{N-p} \left(f \left(\rho_N I_{N-p} - \frac{W_{N-p}}{\sqrt{N}} - A_{N-p} \right) \right) - \int \frac{d\mu_{sc} \boxplus \mu_{A_{N-p}}(x)}{(\rho_N - x)} \right\} \xrightarrow{N \rightarrow +\infty} 0. \quad (58)$$

Define $k : M_{N-p}^{sa}(\mathbb{C}) \rightarrow \mathbb{C}$ by

$$k(X) = \operatorname{tr}_{N-p} [f(\rho_N I_{N-p} - X - A_{N-p})].$$

Applying Poincaré inequality, we get that

$$\mathbb{E} \left(\left| k \left(\frac{W_{N-p}}{\sqrt{N}} \right) - \mathbb{E} \left(k \left(\frac{W_{N-p}}{\sqrt{N}} \right) \right) \right|^2 \right) \leq \frac{C}{N} \mathbb{E} \left(\left\| \operatorname{grad} k \left(\frac{W_{N-p}}{\sqrt{N}} \right) \right\|_e^2 \right),$$

with

$$\|\operatorname{grad} k(X)\|_e^2 = \sup_{w \in S_1(M_{N-p}^{sa}(\mathbb{C}))} \left| \frac{d}{dt} k(X + tw) \Big|_{t=0} \right|^2.$$

Since f is a Lipschitz function on \mathbb{R} with Lipschitz constant C_L , its extension on Hermitian matrices is C_L -Lipschitz with respect to the norm $\|M\|_e = (\operatorname{Tr}_{N-p} M^2)^{1/2}$. Therefore,

$$\sup_{w \in S_1(M_{N-p}^{sa}(\mathbb{C}))} \left| \frac{d}{dt} k(X + tw) \Big|_{t=0} \right|^2 \leq \frac{C}{N},$$

and then

$$\mathbb{E} \left(\left| \sqrt{N} \left\{ k \left(\frac{W_{N-p}}{\sqrt{N}} \right) - \mathbb{E} \left(k \left(\frac{W_{N-p}}{\sqrt{N}} \right) \right) \right\} \right|^2 \right) \leq \frac{C}{N}.$$

It readily follows that

$$\sqrt{N} \operatorname{tr}_{N-p} \left\{ f \left(\rho_N I_{N-p} - \frac{W_{N-p}}{\sqrt{N}} - A_{N-p} \right) - \mathbb{E} \left(f \left(\rho_N I_{N-p} - \frac{W_{N-p}}{\sqrt{N}} - A_{N-p} \right) \right) \right\} = o_{\mathbb{P}}(1). \quad (59)$$

Proposition 11 follows from (55), (58), (59) and (51). \square

3 Proof of Theorem 5

The approach to prove (10) is the one of [16]. On Ω_N , defined at the beginning of Section 2.6.2, we have by Proposition 5,

$$\det(X_p(N)) = 0, \quad (60)$$

where

$$X_p(N) = \lambda_{i_0}(M_N)I_p - \frac{W_p}{\sqrt{N}} - A_p - \frac{1}{N}Y^*\hat{G}_{N-p}(\lambda_{i_0}(M_N))Y,$$

and \hat{G}_{N-p} is the resolvent of $\frac{W_{N-p}}{\sqrt{N}} + A_{N-p}$. Let ρ_N be as defined by (7). Using the identity

$$\hat{G}_{N-p}(\rho_N) - \hat{G}_{N-p}(\lambda_{i_0}(M_N)) = (\lambda_{i_0}(M_N) - \rho_N)\hat{G}_{N-p}(\rho_N)\hat{G}_{N-p}(\lambda_{i_0}(M_N)),$$

we have

$$X_p(N) = H_p(N) + X_p^{(0)},$$

where

$$X_p^{(0)} = \theta I_p - A_p,$$

$$\begin{aligned} H_p(N) &= (\lambda_{i_0}(M_N) - \rho_N)I_p - \Delta_1(N) - \Delta_2(N) \\ &\quad + (\lambda_{i_0}(M_N) - \rho_N)r_1(N) - \frac{W_p}{\sqrt{N}} - (\lambda_{i_0}(M_N) - \rho_N)^2 r_2(N) \end{aligned}$$

with

$$r_1(N) = \frac{1}{N}Y^*\hat{G}_{N-p}(\rho_N)^2\mathbf{1}_{\Omega_N}Y,$$

$$r_2(N) = \frac{1}{N}Y^*\hat{G}_{N-p}(\rho_N)^2\hat{G}_{N-p}(\lambda_{i_0}(M_N))\mathbf{1}_{\Omega_N}Y,$$

$$\Delta_1(N) = \frac{1}{N}Y^*\hat{G}_{N-p}(\rho_N)\mathbf{1}_{\Omega_N}Y - \frac{\sigma^2}{N}\mathrm{Tr}_{N-p}\left(\hat{G}_{N-p}(\rho_N)\mathbf{1}_{\Omega_N}\right),$$

$$\Delta_2(N) = \frac{1}{N}\mathrm{Tr}_{N-p}\left(\hat{G}_{N-p}(\rho_N)\mathbf{1}_{\Omega_N}\right) - \int \frac{d\mu_{sc} \boxplus \mu_{A_{N-p}}(x)}{(\rho_N - x)}.$$

First, by Lemma 4 we have that,

$$r_1(N) - \sigma^2 \frac{1}{N}\mathrm{Tr}_{N-p}(\hat{G}_{N-p}(\rho_N)^2)\mathbf{1}_{\Omega_N}I_p = o_{\mathbb{P}}(1).$$

By (51), (55), (8), (52) and asymptotic freeness of $\frac{W_{N-p}}{\sqrt{N}}$ and A_{N-p} (see Theorem 5.4.5 [1]),

$$\frac{1}{N}\mathrm{Tr}_{N-p}(\hat{G}_{N-p}(\rho_N)^2)\mathbf{1}_{\Omega_N} \xrightarrow{N \rightarrow \infty} \int \frac{d\lambda(x)}{(\rho_\theta - x)^2} \text{ almost surely.} \quad (61)$$

Therefore,

$$r_1(N) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} \sigma^2 \int \frac{d\lambda(x)}{(\rho_\theta - x)^2} I_p. \quad (62)$$

Now on Ω_N , using (53),

$$\|r_2(N)\| \leq \left\| \hat{G}_{N-p}(\rho_N) \mathbf{1}_{\Omega_N} \right\|^2 \left\| \hat{G}_{N-p}(\lambda_{i_0}(M_N)) \mathbf{1}_{\Omega_N} \right\| \frac{\|Y\|^2}{N} \leq 9 \left(\frac{2}{\epsilon_0} \right)^2. \quad (63)$$

By Lemma 4,

$$\Delta_1(N) = o_{\mathbb{P}}(1). \quad (64)$$

Lemma 10.

$$\frac{1}{N-p} \sum_{j=1}^{N-p} \left[\hat{G}_{N-p}(\rho_N)_{jj} \right]^2 \mathbf{1}_{\Omega_N} = \frac{1}{N-p} \sum_{j=1}^{N-p} \left[[(\theta - A_{N-p})^{-1}]_{jj} \right]^2 + o_{\mathbb{P}}(1).$$

Proof. By (96) which will be proved below, for any $r > 0$,

$$\frac{1}{N-p} \sum_{j=1}^{N-p} \left[\hat{G}_{N-p}(\rho_\theta + \frac{i}{r})_{jj} \right]^2 \mathbf{1}_{\Omega_N} = \frac{1}{N-p} \sum_{j=1}^{N-p} \left[[(\omega(\rho_\theta + \frac{i}{r}) - A_{N-p})^{-1}]_{jj} \right]^2 + o_{\mathbb{P}}(1),$$

with ω defined by (33). Now, using resolvent identity, one can easily obtain that there exists some constant $C(\epsilon_0)$ such that for any $r > 0$ and any N ,

$$\left| \frac{1}{N-p} \sum_{j=1}^{N-p} \left[\hat{G}_{N-p}(\rho_N)_{jj} \right]^2 \mathbf{1}_{\Omega_N} - \frac{1}{N-p} \sum_{j=1}^{N-p} \left[\hat{G}_{N-p}(\rho_\theta + \frac{i}{r})_{jj} \right]^2 \mathbf{1}_{\Omega_N} \right| \leq C(\epsilon_0) \left(\frac{1}{r} + \rho_N - \rho \right).$$

Moreover, $\omega(\rho_\theta) = \theta$ so that for all large N , $d(\omega(\rho_\theta), \text{supp}(\mu_{A_{N-p}})) > \epsilon_0/2$. Now, choose r_0 large enough such that for all large N , $\forall r \geq r_0$, $d(\omega(\rho_\theta + \frac{i}{r}), \text{supp}(\mu_{A_{N-p}})) > \epsilon_0/4$. Using resolvent identity, one can easily obtain that there exists some constant $C(\epsilon_0)$ such that for all large N , $\forall r \geq r_0$,

$$\left| \frac{1}{N-p} \sum_{j=1}^{N-p} \left[[(\omega(\rho_\theta + \frac{i}{r}) - A_{N-p})^{-1}]_{jj} \right]^2 - \frac{1}{N-p} \sum_{j=1}^{N-p} \left[[(\omega(\rho_\theta) - A_{N-p})^{-1}]_{jj} \right]^2 \right| \leq C(\epsilon_0)/r.$$

Lemma 10 follows by letting N go to infinity and then r go to infinity. \square

(61), Lemma 10 and Proposition 9 yield that

$$\sqrt{N}\Delta_1(N)^2 = o_{\mathbb{P}}(1), \quad (65)$$

Now, one can prove that by Proposition 11, we have

$$\sqrt{N}\Delta_2(N) = o_{\mathbb{P}}(1). \quad (66)$$

Thus (6), (8), (62), (63), (64) and (66) yield that

$$H_p(N) = o_{\mathbb{P}}(1). \quad (67)$$

Therefore, according to Lemma 2 (using (67)) and (60), with a probability going to one as N goes to infinity,

$$\begin{aligned} 0 &= \det X_p(N) \\ &= \det(X_p^{(0)} + H_p(N)) \\ &= \det(X_p^{(0)}) + \text{Tr}_p \left[B_{X_p^{(0)}} H_p(N) \right] + \epsilon_N \\ &= \text{Tr}_p \left[B_{X_p^{(0)}} H_p(N) \right] + \epsilon_N, \end{aligned}$$

where

$$\begin{aligned} B_{X_p^{(0)}} &= {}^t \text{com}(X_p^{(0)}), \\ \epsilon_N &= O(\|H_p(N)\|^2). \end{aligned}$$

Thus, using (6), (8), (62), (63), (66) and (65),

$$\sqrt{N}\epsilon_N = o_{\mathbb{P}}(\sqrt{N}(\lambda_{i_0}(M_N) - \rho_N)) + o_{\mathbb{P}}(1).$$

Hence, with a probability going to one as N goes to infinity,

$$\begin{aligned} &\sqrt{N}(\lambda_{i_0}(M_N) - \rho_N) \left[\text{Tr}_p B_{X_p^{(0)}} (I_p + r_1(N)) + o_{\mathbb{P}}(1) \right] \\ &= \text{Tr}_p \left[B_{X_p^{(0)}} \left(\sqrt{N}\Delta_1(N) + W_p \right) \right] + o_{\mathbb{P}}(1). \end{aligned}$$

(10) readily follows from (61), Lemma 10, Proposition 9, the independence of $\Delta_1(N)$ and W_p . When A_{N-p} is diagonal, the result follows using (96).

4 Proofs of Theorems 6 and 7

Let M_N be defined by (2) with assumptions **(W)** and **(A')**. We denote by $\lambda_i(A_N)$, resp. $\lambda_i(M_N)$, the eigenvalues of A_N , resp. M_N and u_i , resp. v_i the normalized associated eigenvectors. According to assumption **(A)**, there exists $\delta > 0$ such that for all large N , the distance from θ to the rest of the spectrum (that is the other eigenvalues of A_N except θ) is greater than δ . Moreover, from Proposition 1, we know that a.s.

$$\lambda_{i_0}(M_N) \xrightarrow{N \rightarrow +\infty} \rho_\theta = \theta + \sigma^2 g_\nu(\theta). \quad (68)$$

and there exists $\delta_0 > 0$ such that almost surely for all large N , the distance from ρ_θ to the rest of the spectrum (that is the other eigenvalues of M_N except $\lambda_{i_0}(M_N)$) is greater than δ_0 .

Throughout this section, h is a smooth function with support in $]\rho_\theta - \delta_0/2; \rho_\theta + \delta_0/2[$ which is equal to 1 near ρ_θ .

4.1 Representation in terms of resolvent

The aim of this section is to be brought back to the study of the fluctuations of the $p \times p$ -matrix valued process $\{G_p(z), z \in \mathbb{C} \setminus \mathbb{R}\}$ where $G_p(z)$ denotes the principal submatrix of size p of the resolvent matrix $G(z) = (zI_N - M_N)^{-1}$.

Proposition 12. *Almost surely, for all large N ,*

$$\begin{aligned} & \sqrt{N}(|\langle u_{i_0}, v_{i_0} \rangle|^2 - \tau_N(\theta)) \\ &= -\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) \left(P^* \sqrt{N} (G_p(z) - \Lambda_p(z)) P \right)_{11} d^2 z, \end{aligned}$$

where $\tau_N(\theta)$ is defined by (11),

$$\Lambda_p(z) = (zI_p - A_p - \sigma^2 \tilde{g}_{N-p}(z) I_p)^{-1} \quad (69)$$

and $\tilde{g}_{N-p}(z)$ is the Stieltjes transform of $\mu_{sc} \boxplus \mu_{A_{N-p}}$.

Proposition 12 readily follows from the two preliminaries Lemmas 11 and 12.

Lemma 11. *Almost surely, for all large N , for any integer number k ,*

$$|\langle u_{i_0}, v_{i_0} \rangle|^2 = -\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) (P^* G_p(z) P)_{11} d^2 z, \quad (70)$$

where $F_k(h)$ is defined by (28).

Proof. Let f be any smooth function with support in $]\theta - \delta/2; \theta + \delta/2[$ which is equal to 1 near θ . From the formula

$$\mathrm{Tr}_N(h(M_N)f(A_N)) = \sum_{i,j=1}^N h(\lambda_i(M_N))f(\lambda_j(A_N))|\langle u_j, v_i \rangle|^2, \quad (71)$$

we easily deduce that, almost surely, for all large N ,

$$|\langle u_{i_0}, v_{i_0} \rangle|^2 = \sum_{i,j=1}^p (P^*)_{1i} h(M_N)_{ij} P_{j1}. \quad (72)$$

By Helffer-Sjöstrand's representation formula (27), we can write $h(M_N)_{ij}$ as

$$h(M_N)_{ij} = -\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) G_{ij}(z) d^2 z,$$

so that

$$\sum_{i,j=p}^p (P^*)_{1i} h(M_N)_{ij} P_{j1} = -\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) (P^* G_p(z) P)_{11} d^2 z. \quad (73)$$

Lemma 11 follows from (72) and (73). \square

Lemma 12. *For N large enough,*

$$\begin{aligned} \tau_N(\theta) &= 1 - \sigma^2 \int \frac{1}{(\theta - x)^2} d\mu_{A_{N-p}}(x) \\ &= -\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) (P^*(zI_p - A_p - \sigma^2 \tilde{g}_{N-p}(z) I_p)^{-1} P)_{11} d^2 z \end{aligned}$$

where $\tilde{g}_{N-p}(z)$ is the Stieltjes transform of $\mu_{sc} \boxplus \mu_{A_{N-p}}$.

Proof. Note that

$$(P^*(zI_p - A_p - \sigma^2 \tilde{g}_{N-p}(z) I_p)^{-1} P)_{11} = \frac{1}{z - \sigma^2 \tilde{g}_{N-p}(z) - \theta}.$$

Let us define for any $z \in \mathbb{C} \setminus \mathrm{supp}(\mu_{A_{N-p}})$,

$$H_{N-p}(z) = z + \sigma^2 g_{\mu_{A_{N-p}}}(z) \quad (74)$$

and for any $z \in \mathbb{C} \setminus \mathrm{supp}(\mu_{sc} \boxplus \mu_{A_{N-p}})$,

$$\omega_{N-p}(z) = z - \sigma^2 g_{\mu_{sc} \boxplus \mu_{A_{N-p}}}(z) = z - \sigma^2 \tilde{g}_{N-p}(z). \quad (75)$$

Note that, for any $z \in \mathbb{C} \setminus \mathbb{R}$, $|\Im \omega_{N-p}(z)| \geq |\Im z| > 0$. Moreover, according to (36), the following one to one correspondance holds:

$$\mathbb{R} \setminus \text{supp}(\mu_{sc} \boxplus \mu_{A_{N-p}}) \xrightleftharpoons[H_{N-p}]{\omega_{N-p}} \left\{ u \in \mathbb{R} \setminus \text{supp}(\mu_{A_{N-p}}), \int \frac{1}{(u-x)^2} d\mu_{A_{N-p}}(x) < \frac{1}{\sigma^2} \right\}$$

and for any $x \in \mathbb{C} \setminus \text{supp}(\mu_{sc} \boxplus \mu_{A_{N-p}})$, $H_{N-p}(\omega_{N-p}(x)) = x$. Hence $\rho_N = H_{N-p}(\theta) = \theta + \sigma^2 g_{\mu_{A_{N-p}}}(\theta)$ is the single pole of $\frac{1}{z - \sigma^2 \tilde{g}_{N-p}(z) - \theta}$ in \mathbb{C} . Therefore, (8) and Proposition 6 (used with $\phi(z) = \frac{1}{\omega_{N-p}(z) - \theta}$) imply that

$$-\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) \frac{1}{z - \sigma^2 \tilde{g}_{N-p}(z) - \theta} d^2 z = \frac{1}{\omega'_{N-p}(\rho_N)} = H'_{N-p}(\theta) = \tau_N(\theta). \quad (76)$$

□

We now consider the process

$$\xi_N(z) = \left(P^* \sqrt{N} (G_p(z) - \Lambda_p(z)) P \right)_{11}. \quad (77)$$

where Λ_p is defined by (69).

4.2 Tightness of the sequence of processes $\{\xi_N\}_N$

Proposition 13. ξ_N is tight on $\mathcal{H}(\mathbb{C} \setminus \mathbb{R})$.

Proof. $\xi_N : z \mapsto \left(P^* \sqrt{N} (G_p(z) - \Lambda_p(z)) P \right)_{11}$ is analytic on $\mathbb{C} \setminus \mathbb{R}$. Let K be a compact set in $\mathbb{C} \setminus \mathbb{R}$. According to Lemma 3, there exists $\delta > 0$ such that $\overline{K_\delta} \subset \mathbb{C} \setminus \mathbb{R}$ and for any $r > 0$,

$$\|\xi_N\|_K^r \leq (\pi \delta^2)^{-1} \int_{\overline{K_\delta}} |\xi_N(z)|^r m(dz).$$

Therefore

$$\mathbb{E}(\|\xi_N\|_K^r) \leq (\pi \delta^2)^{-1} \int_{\overline{K_\delta}} \mathbb{E}(|\xi_N(z)|^r) m(dz) \quad (78)$$

$$\leq (\pi \delta^2)^{-1} \sup_{z \in \overline{K_\delta}} \mathbb{E}(|\xi_N(z)|^r) m(\overline{K_\delta}). \quad (79)$$

In order to prove the tightness of ξ_N , using (32) and (79), we are going to show that, for any compact set $K \subset \mathbb{C} \setminus \mathbb{R}$, there exists a constant $C' > 0$ such that for all large N ,

$$\sup_{z \in K} \mathbb{E}(|\xi_N(z)|^2) < C'. \quad (80)$$

We have for any z_1 and z_2 in $\mathbb{C} \setminus \mathbb{R}$,

$$\begin{aligned} & \mathbb{E}(\xi_N(z_1)\xi_N(z_2)) \\ &= N \sum_{i,j,u,v=1}^p P_{1i}^* P_{j1} P_{1u}^* P_{v1} \mathbb{E}((G_{ij}(z_1) - (\Lambda_p(z_1))_{ij})(G_{uv}(z_2) - (\Lambda_p(z_2))_{uv})), \end{aligned}$$

and

$$\begin{aligned} & \mathbb{E}((G_{ij}(z_1) - (\Lambda_p(z_1))_{ij})(G_{uv}(z_2) - (\Lambda_p(z_2))_{uv})) \\ &= \mathbb{E}((G_{ij}(z_1) - \mathbb{E}(G_{ij}(z_1)))(G_{uv}(z_2) - \mathbb{E}(G_{uv}(z_2)))) \\ & \quad + [\mathbb{E}(G_{ij}(z_1)) - (\Lambda_p(z_1))_{ij}][\mathbb{E}(G_{uv}(z_2)) - (\Lambda_p(z_2))_{uv}]. \end{aligned}$$

According to Lemma 6 and the block diagonal structure of A_N , we have for any $l = 1, 2$,

$$\mathbb{E}(G_p(z_l)) = ((z_l - \sigma^2 \tilde{g}_N(z_l))I_p - A_p)^{-1} + O(1/\sqrt{N}), \quad (81)$$

where \tilde{g}_N is the Stieltjes transform of $\mu_{A_N} \boxplus \mu_{sc}$. First set

$$\check{G}_{N-p}(w) = \left(zI_{N-p} - \frac{W_{N-p}}{\sqrt{N-p}} - A_{N-p} \right)^{-1}.$$

By (25), we have

$$\|\check{G}_{N-p}(z)\| \leq |\Im z|^{-1}. \quad (82)$$

Note that,

$$\begin{aligned} \hat{G}_{N-p}(z) &= \check{G}_{N-p}(z) \\ & \quad + \frac{p}{\sqrt{N-p}(\sqrt{N} + \sqrt{N-p})} \\ & \quad \times \left(I_{N-p} - \hat{G}_{N-p}(z)(zI_{N-p} - A_{N-p}) \right) \check{G}_{N-p}(z) \end{aligned} \quad (83)$$

Now, Lemma 7 yields that

$$\tilde{g}_N(z) = \mathbb{E}(\text{tr}_N G(z)) + O(1/N)$$

and

$$\tilde{g}_{N-p}(z) = \mathbb{E}(\text{tr}_{N-p} \check{G}_{N-p}(z)) + O(1/N),$$

and then, using (83), that

$$\tilde{g}_{N-p}(z) = \mathbb{E}(\text{tr}_{N-p} \hat{G}_{N-p}(z)) + O(1/N). \quad (84)$$

Since by (A.1.12) in [3], we have $\mathbb{E}(\text{tr}_N G(z)) = \mathbb{E}(\text{tr}_{N-p} \hat{G}_{N-p}(z)) + O(1/N)$, we can deduce that $\tilde{g}_N(z) = \tilde{g}_{N-p}(z) + O(1/N)$, and thus that, for any $1 \leq i, j \leq p$,

$$\mathbb{E}(G_{ij}(z_l)) = (\Lambda_p(z_l))_{ij} + O(1/\sqrt{N}).$$

Moreover, by Lemma 8,

$$\text{Var}(G_{ij}(z_l)) = O(1/N).$$

It readily follows that there exist polynomials P_1 and P_2 with nonnegative coefficients such that

$$\mathbb{E}(\xi_N(z_1)\xi_N(z_2)) \leq P_1(|\Im z_1|^{-1}) P_2(|\Im z_2|^{-1})$$

and then there exists some polynomial P_3 with nonnegative coefficients such that for all large N and all $z \in \mathbb{C} \setminus \mathbb{R}$,

$$\mathbb{E}(|\xi_N(z)|^2) \leq P_3(|\Im z|^{-1}). \quad (85)$$

This implies (80). Therefore, for any compact subset K in $\mathbb{C} \setminus \mathbb{R}$, there exists a constant $C > 0$ such that for all large N , $\mathbb{E}(\|\xi_N(z)\|_K^2) < C$ and the tightness of ξ_N in $\mathcal{H}(\mathbb{C} \setminus \mathbb{R})$ follows from (32) and Proposition 7. \square

4.3 Finite dimensional distributions of ξ_N

Set

$$\nabla_N(z) = \sqrt{N}(G_p(z) - \Lambda_p(z)).$$

We use Proposition 5 for the inversion of

$$zI_N - M_N = \begin{pmatrix} zI_p - \frac{1}{\sqrt{N}}W_p - A_p & -\frac{1}{\sqrt{N}}Y^* \\ -\frac{1}{\sqrt{N}}Y & zI_{N-p} - M_{N-p} \end{pmatrix}$$

where M_{N-p} is the lower right submatrix of size $N-p$ of M_N , leading to:

$$G_p(z) = (zI_p - \frac{1}{\sqrt{N}}W_p - A_p - \frac{1}{N}Y^*\hat{G}_{N-p}(z)Y)^{-1} \quad (86)$$

where \hat{G}_{N-p} is the resolvent of M_{N-p} . Thus,

$$\nabla_N(z) = G_p(z)(W_p + \sqrt{N}(\frac{1}{N}Y^*\hat{G}_{N-p}(z)Y - \sigma^2\tilde{g}_{N-p}(z)I_p))\Lambda_p(z). \quad (87)$$

Lemma 13. Define for $z \in \mathbb{C} \setminus \mathbb{R}$,

$$\tilde{\nabla}_N(z) = (zI_p - A_p - \sigma^2 g(z)I_p)^{-1} (W_p + Q_N(z)) (zI_p - A_p - \sigma^2 g(z)I_p)^{-1}, \quad (88)$$

where $Q_N(z)$ is the following matrix of size p :

$$Q_N(z) = \frac{1}{\sqrt{N}} (Y^* \hat{G}_{N-p}(z) Y - \sigma^2 \text{Tr}_{N-p}(\hat{G}_{N-p}(z)) I_p), \quad z \in \mathbb{C} \setminus \mathbb{R} \quad (89)$$

For any $z \in \mathbb{C} \setminus \mathbb{R}$,

$$\nabla_N(z) - \tilde{\nabla}_N(z) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} 0.$$

Proof. Obviously, we have

$$\Lambda_p(z) \xrightarrow[N \rightarrow \infty]{} (zI_p - A_p - \sigma^2 g(z)I_p)^{-1}, \quad (90)$$

and (81) and Lemma 8 yield that

$$G_p(z) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} (zI_p - A_p - \sigma^2 g(z)I_p)^{-1}. \quad (91)$$

We write

$$\begin{aligned} & \sqrt{N} \left(\frac{1}{N} Y^* \hat{G}_{N-p}(z) Y - \sigma^2 \tilde{g}_{N-p}(z) I_p \right) \\ &= \frac{1}{\sqrt{N}} (Y^* \hat{G}_{N-p}(z) Y - \sigma^2 \text{Tr}_{N-p}(\hat{G}_{N-p}(z)) I_p) \\ & \quad + \sqrt{N} \sigma^2 (\text{tr}_{N-p}(\hat{G}_{N-p}(z)) I_p - \tilde{g}_{N-p}(z) I_p). \end{aligned} \quad (92)$$

(84) and Lemma 9 yield that

$$\sqrt{N} \sigma^2 (\text{tr}_{N-p}(\hat{G}_{N-p}(z)) I_p - \tilde{g}_{N-p}(z) I_p) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} 0. \quad (93)$$

Lemma 13 readily follows from (87), (92), (90), (91), (93) and the tightness of $Q_N(z)$ (which readily follows from Lemma 4), by using Slutsky's theorem and classical operations on convergence in probability. \square

We now state an approximation result in distribution for the finite dimensional distributions of the process $\{Q_N(z), z \in \mathbb{C} \setminus \mathbb{R}\}$.

Lemma 14. Let (z_1, \dots, z_q) be in $(\mathbb{C} \setminus \mathbb{R})^q$. Set $V_N = (Q_N(z_1), \dots, Q_N(z_q))$. Then, under the assumptions of Theorem 6

$$d_{LP}(V_N, (\mathcal{G}_N(z_1), \dots, \mathcal{G}_N(z_q))) \rightarrow 0$$

where \mathcal{G}_N is a centered matrix valued Gaussian process whose distribution is given as follows :

- 1) the processes $((\mathcal{G}_N)_{ij}(z))_z$ for $1 \leq i \leq j \leq p$ are independent, and $(\mathcal{G}_N)_{ji}(z) = \overline{(\mathcal{G}_N)_{ij}(\bar{z})}$.
- 2) For $i \leq p$,

$$\mathbb{E}((\mathcal{G}_N)_{ii}(z_k)(\mathcal{G}_N)_{ii}(z_l)) \quad (94)$$

$$\begin{aligned} &= \frac{(m_4 - 3\sigma^4)}{2(N-p)} \sum_{i=1}^{N-p} ((z_k - \sigma^2 g(z_k) - A_{N-p})^{-1})_{ii} ((z_l - \sigma^2 g(z_l) - A_{N-p})^{-1})_{ii} \\ &\quad + \sigma^4 \int \frac{1}{(z_k - x)(z_l - x)} d\lambda(x) \end{aligned} \quad (95)$$

- 2) For $1 \leq i \neq j \leq p$,

$$\mathbb{E}((\mathcal{G}_N)_{ij}(z_k)(\mathcal{G}_N)_{ij}(z_l)) = 0$$

and

$$\mathbb{E}((\mathcal{G}_N)_{ij}(z_k) \overline{(\mathcal{G}_N)_{ij}(z_l)}) = \sigma^4 \int \frac{1}{(z_k - x)(\bar{z}_l - x)} d\lambda(x).$$

Proof. We apply Proposition 9 to the matrices $B(z) = \hat{G}_{N-p}(z)$, $z \in \mathbb{C} \setminus \mathbb{R}$. Note that these matrices are random but they are independent of the $N \times p$ matrix Y .

In order to conclude, we need to show that

$$I_N := \frac{1}{N-p} \sum_{i=1}^{N-p} ((\hat{G}_{N-p}(z_1))_{ii} (\hat{G}_{N-p}(z_2))_{ii}) = f_N(z_1, z_2) + o(1) \quad (96)$$

with

$$f_N(z_1, z_2) = \frac{1}{N-p} \sum_{i=1}^{N-p} ((z_k - \sigma^2 g(z_k) - A_{N-p})^{-1})_{ii} ((z_l - \sigma^2 g(z_l) - A_{N-p})^{-1})_{ii}$$

and

$$J_N := \text{tr}_{N-p}(\hat{G}_{N-p}(z_1)\hat{G}_{N-p}(z_2)) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} \int \frac{1}{(z_1 - x)(z_2 - x)} d\lambda(x).$$

The second convergence follows from the convergence of $\mu_{M_{N-p}}$ towards λ . For the first one, Lemma 6 and (83) yield that, for $k \leq N-p$,

$$\begin{aligned} \mathbb{E} \left((\hat{G}_{N-p}(z))_{kk} \right) &= \left[((z - \sigma^2 \tilde{g}_{N-p}(z)) I_N - A_{N-p})^{-1} \right]_{kk} + O \left(\frac{1}{\sqrt{N}} \right) \\ &= \left[((z - \sigma^2 g(z)) I_N - A_{N-p})^{-1} \right]_{kk} + o^{(u)}(1) \end{aligned}$$

Thus, using Lemma 8, we can deduce that

$$\begin{aligned} & \mathbb{E}((\hat{G}_{N-p}(z_1))_{kk}(\hat{G}_{N-p}(z_2))_{kk}) \\ &= [(z_1 - \sigma^2 g(z_1) - A_{N-p})^{-1}]_{kk} [(z_2 - \sigma^2 g(z_2) - A_{N-p})^{-1}]_{kk} \\ & \quad + o^{(u)}(1) \end{aligned} \tag{97}$$

From Lemma 17 in the Appendix, using that $f_N(W) = \frac{1}{N-p} \sum_{i=1}^{N-p} [(z_1 - W - A_{N-p})^{-1}]_{ii} [(z_2 - W - A_{N-p})^{-1}]_{ii}$ is Lipschitz with constant $|\Im(z_1)|^{-2} |\Im(z_2)|^{-1} + |\Im(z_1)|^{-1} |\Im(z_2)|^{-2}$, we can deduce that

$$\frac{1}{N-p} \sum_{i=1}^{N-p} (\hat{G}_{N-p}(z_1))_{ii} (\hat{G}_{N-p}(z_2))_{ii} = \frac{1}{N-p} \sum_{i=1}^{N-p} \mathbb{E} \left((\hat{G}_{N-p}(z_1))_{ii} (\hat{G}_{N-p}(z_2))_{ii} \right) + o_{\mathbb{P}}(1).$$

We also use to obtain (95) from Proposition 9 that for $\sigma^2 = 1$,

$$\mathbb{E}(|y_{11}|^4) - 2 = \frac{1}{2}(m_4 - 3),$$

where we recall that μ is the distribution of $\sqrt{2}\Re y_{11}$ and $\sqrt{2}\Im y_{11}$. \square

Corollary 2. *Under the assumptions of Theorem 6, for any (z_1, \dots, z_q) be in $(\mathbb{C} \setminus \mathbb{R})^q$,*

$$d_{LP}((\xi_N(z_1), \dots, \xi_N(z_q)), (\mathcal{T}_N(z_1), \dots, \mathcal{T}_N(z_q))) \rightarrow 0$$

where ξ_N is the process defined by (77) and

$$\mathcal{T}_N(z) = (z - \theta - \sigma^2 g(z))^{-2} (P^*(W_p + \mathcal{G}_N(z))P)_{11} \tag{98}$$

\mathcal{G}_N being a centered Gaussian matrix valued process independent from W_p whose distribution is described in Lemma 14.

4.4 Fluctuations of the eigenvector

Recall from Proposition 12 that, for any $k \in \mathbb{N}^*$, Φ_N defined as

$$\sqrt{N}(|\langle u_{i_0}, v_{i_0} \rangle|^2 - \tau_N(\theta))$$

has the representation

$$\Phi_N = -\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) \xi_N(z) d^2 z. \tag{99}$$

We follow the proof of Lemma 6.3 in [31] based upon the following estimates, from (85) :

$$\sup_N \mathbb{E}(|\xi_N(z)|) \leq P_3(|\Im(z)|^{-1}), \quad (100)$$

and from Lemma 14 and (26), (\mathcal{T}_N being defined by (98))

$$\sup_N \mathbb{E}(|\mathcal{T}_N(z)|) \leq P_5(|\Im(z)|^{-1}), \quad (101)$$

where P_3 , P_4 and P_5 are some polynomial with nonnegative coefficients. Hence in the following, in (99), we choose k greater than the degrees of P_3 , P_4 and P_5 .

Proposition 14. *Under the assumptions of Theorem 6, $d_{LP}(\Phi_N, \tilde{\Phi}_N) \rightarrow 0$ where $\tilde{\Phi}_N$ is given by*

$$\tilde{\Phi}_N = (P^*(c_{\theta,\nu}W_p + Z_{p,N})P)_{11},$$

where W_p is a Wigner matrix of size p , $Z_{p,N}$ is a centered Gaussian Hermitian matrix of size p with independent entries (modulo the symmetry condition); the diagonal coefficients are iid with variance

$$\frac{1}{2}(m_4 - 3\sigma^4)A_{\theta,\nu,N} + \sigma^4 B_{\theta,\nu} \quad (102)$$

and the off diagonal elements are i.i.d. complex Gaussian with distribution Z such that $\mathbb{E}(Z^2) = 0$ and $\mathbb{E}(|Z|^2) = \sigma^4 B_{\theta,\nu}$.

See Eq.(18) for the definitions of $c_{\theta,\sigma}$, $B_{\theta,\nu}$ and (15) for the definition of $A_{\theta,\nu,N}$.

Proof. For the reader's convenience, we repeat here the strategy of Lemma 6.3 in [31]. For any $0 < \epsilon < 1$ small enough such that $\chi \equiv 1$ on $] - \epsilon; \epsilon[$, define

$$D_\epsilon = \{z \in \mathbb{C}, |\Im(z)| \geq \epsilon\}.$$

Set

$$U_N = \int_{\mathbb{C}} \bar{\partial}F_k(h)(z)\xi_N(z)d^2z, \quad U_N^\epsilon = \int_{D_\epsilon} \bar{\partial}F_k(h)(z)\xi_N(z)d^2z$$

and

$$V_N = \int_{\mathbb{C}} \bar{\partial}F_k(h)(z)\mathcal{T}_N(z)d^2z, \quad V_N^\epsilon = \int_{D_\epsilon} \bar{\partial}F_k(h)(z)\mathcal{T}_N(z)d^2z,$$

where \mathcal{T}_N is defined by (98). Let f be a bounded continuous complex function on \mathbb{C} .

We have

$$\begin{aligned} |\mathbb{E}(f(U_N)) - \mathbb{E}(f(V_N))| &\leq |\mathbb{E}(f(U_N)) - \mathbb{E}(f(U_N^\epsilon))| \\ &\quad + |\mathbb{E}(f(U_N^\epsilon)) - \mathbb{E}(f(V_N^\epsilon))| \\ &\quad + |\mathbb{E}(f(V_N^\epsilon)) - \mathbb{E}(f(V_N))| \end{aligned}$$

Let $\delta > 0$.

i) For any $\eta > 0$ and $K > 0$, we have

$$\begin{aligned} &|\mathbb{E}(f(U_N)) - \mathbb{E}(f(U_N^\epsilon))| \\ &\leq \left| \mathbb{E}(f(U_N) - f(U_N^\epsilon)) \mathbf{1}_{|U_N - U_N^\epsilon| > \eta} \right| \end{aligned} \quad (103)$$

$$+ \left| \mathbb{E}(f(U_N) - f(U_N^\epsilon)) \mathbf{1}_{|U_N - U_N^\epsilon| \leq \eta, |U_N| \vee |U_N^\epsilon| > K} \right| \quad (104)$$

$$+ \left| \mathbb{E}(f(U_N) - f(U_N^\epsilon)) \mathbf{1}_{|U_N - U_N^\epsilon| \leq \eta, |U_N| \vee |U_N^\epsilon| \leq K} \right| \quad (105)$$

In the following, the constant $C > 0$ may vary from line to line. By (30), for any $z = x + iy$ in a neighborhood of the real axis,

$$|\bar{\partial}F_k(h)(z)| \leq C|y|^k. \quad (106)$$

(29), (100) and (106) readily yield that for any $\epsilon > 0$, for any N ,

$$\mathbb{E}(|U_N|) \vee \mathbb{E}(|U_N^\epsilon|) \leq \int_{\mathbb{C}} |\bar{\partial}F_k(h)(z)| \mathbb{E}|\xi_N(z)| d^2z \leq C,$$

and therefore

$$\begin{aligned} \mathbb{P}(|U_N| \vee |U_N^\epsilon| > K) &\leq \mathbb{P}(|U_N| > K) + \mathbb{P}(|U_N^\epsilon| > K) \\ &\leq \frac{2}{K} \mathbb{E}(|U_N|) \vee \mathbb{E}(|U_N^\epsilon|) \leq \frac{C}{K}. \end{aligned}$$

Thus, we can choose K such that, for any $\epsilon > 0$, any $\eta > 0$ and any N , the RHS in (104) is smaller than δ . Now, since f is uniformly continuous on $\{z \in \mathbb{C} \setminus \mathbb{R}, |z| \leq K\}$, one can choose η small enough such that, for any $\epsilon > 0$ and any N , (105) are smaller than δ . Finally, by using (100) and (106), for any ϵ small enough, for any N ,

$$\mathbb{E}|U_N - U_N^\epsilon| = \mathbb{E} \left| \int_{\{z, |\Im(z)| < \epsilon\}} \bar{\partial}F_k(h)(z) \xi_N(z) d^2z \right| \leq C\epsilon, \quad (107)$$

and then

$$\mathbb{P}(|U_N - U_N^\epsilon| > \eta) \leq \frac{C}{\eta} \epsilon.$$

The term $|\mathbb{E}(f(V_N^\epsilon)) - \mathbb{E}(f(V_N))|$ is treated in the same way, using (101).
ii) Note that since h and χ are compactly supported, $\int_{D_\epsilon} \bar{\partial} F_k(h)(z) \xi_N(z) d^2 z$ may be seen as an integral on a fixed compact set $K_\epsilon \subset D_\epsilon$. Proposition 13 readily yields that ξ_N is tight on the set $\mathcal{C}(K_\epsilon)$ of complex continuous functions on K_ϵ . Thus, using Corollary 2, since $f \mapsto \int_{D_\epsilon} \bar{\partial} F_k(h)(z) f(z) d^2 z$ is continuous on $\mathcal{C}(K_\epsilon)$, it remains to prove the tightness of the process $\{\mathcal{T}_N(\cdot)\}$ on any compact set K in $\{z, |\Im z| \geq \epsilon\}$ to deduce from Lemma 5.7 in [31] that :

$$|\mathbb{E}(f(U_N^\epsilon)) - \mathbb{E}(f(V_N^\epsilon))| \rightarrow_{N \rightarrow +\infty} 0.$$

We postpone the proof of the tightness of the process $\{\mathcal{T}_N(\cdot)\}$ (see Lemma 15 below) Thus, up to the proof of Lemma 15, the convergence to 0 of $d_{LP}(\Phi_n, -\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) \mathcal{T}_N(z) d^2 z)$ follows.

It remains to compute $c_{\theta, \nu}, B_{\theta, \nu}, A_{\theta, \nu, N}$.

The computation of $c_{\theta, \nu}$ follows from Proposition 6 :

$$c_{\theta, \nu} = -\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) \frac{1}{(z - \sigma^2 g(z) - \theta)^2} dz = \text{Res}\left(\frac{1}{(z - \sigma^2 g(z) - \theta)^2}, \rho_\theta\right).$$

A straightforward computation gives

$$\text{Res}\left(\frac{1}{(z - \sigma^2 g(z) - \theta)^2}, \rho_\theta\right) = -\frac{\omega''(\rho_\theta)}{(\omega'(\rho_\theta))^3} = H''(\theta)$$

where $\omega(z) = z - \sigma^2 g(z)$ ($g := g_\lambda$) and $H(z) = z + \sigma^2 g_\nu(z)$.

From Lemma 14 and using Fubini theorem, the diagonal entries of Z_N have variance equal to

$$\frac{1}{2}(m_4 - 3\sigma^4)A_{\theta, \nu, N} + \sigma^4 B_{\theta, \nu}$$

and the off diagonal entries of Z_N have variance equal to $\sigma^4 B_{\theta, \nu}$ where

$$A_{\theta, \nu, N} = \frac{1}{N-p} \sum_{i=1}^{N-p} \left(\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) \frac{[(z - \sigma^2 g(z) - A_{N-p})^{-1}]_{ii}}{(z - \sigma^2 g(z) - \theta)^2} d^2 z \right)^2 d\nu(x)$$

$$B_{\theta, \nu} = \int_{\mathbb{R}} \left(\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} F_k(h)(z) \frac{1}{(z - \sigma^2 g(z) - \theta)^2 (z - x)} d^2 z \right)^2 d\lambda(x).$$

The functions $\phi_i(z) = \frac{[(z - \sigma^2 g(z) - A_{N-p})^{-1}]_{ii}}{(z - \sigma^2 g(z) - \theta)^2}$ for $i \leq N - p$ and $\phi_x(z) = \frac{1}{(z - \sigma^2 g(z) - \theta)^2 (z - x)}$ for $x \in \text{supp}(\lambda)$ satisfy the hypothesis of Proposition 6. Straightforward computations lead to :

$$\begin{aligned}
A_{\theta, \nu, N} &= \frac{1}{N-p} \sum_{i=1}^{N-p} (\text{Res}(\frac{[(z - \sigma^2 g(z) - A_{N-p})^{-1}]_{ii}}{(z - \sigma^2 g(z) - \theta)^2}, \rho_\theta))^2 \\
&= \frac{1}{N-p} \sum_{i=1}^{N-p} (H''(\theta)[(\theta I_{N-p} - A_{N-p})^{-1}]_{ii} - H'(\theta)[(\theta I_{N-p} - A_{N-p})^{-2}]_{ii})^2 \\
&= \frac{1}{N-p} \sum_{i=1}^{N-p} (\sigma^2 g''_\nu(\theta)[(\theta I_{N-p} - A_{N-p})^{-1}]_{ii} - (1 + \sigma^2 g'_\nu(\theta))[(\theta I_{N-p} - A_{N-p})^{-2}]_{ii})^2
\end{aligned}$$

and

$$\begin{aligned}
B_{\theta, \nu} &= \int_{\mathbb{R}} (\text{Res}(\frac{1}{(z - \sigma^2 g(z) - \theta)^2 (z - x)}, \rho_\theta))^2 d\lambda(x) \\
&= -\frac{1}{6} g'''_\nu(\theta) - \frac{\sigma^2}{2} (g''_\nu(\theta))^2 \frac{1 + 2\sigma^2 g'_\nu(\theta)}{1 + \sigma^2 g'_\nu(\theta)}.
\end{aligned}$$

□

We now prove the following Lemma, used in the proof of the above Proposition.

Lemma 15. *Let K be a compact subset in $\{z, |\Im z| \geq \epsilon\}$, for some $\epsilon > 0$. The process $\{\mathcal{T}_N(\cdot)\}$ defined in (98) is a tight sequence on K , more precisely,*

$$\sup_{z_1, z_2 \in K, n \in \mathbb{N}} \frac{\mathbb{E}(|\mathcal{T}_N(z_1) - \mathcal{T}_N(z_2)|^2)}{|z_1 - z_2|^2} < \infty. \quad (108)$$

Proof. From Lemma 14,

$$\begin{aligned}
&\mathbb{E}(|(\mathcal{G}_N)_{ij}(z_1) - (\mathcal{G}_N)_{ij}(z_2)|^2) \\
&= \delta_{ij} \frac{1}{2} (m_4 - 3\sigma^4) \frac{1}{N-p} \sum_{i=1}^{N-p} (|((z_1 - \sigma^2 g(z_1) - A_{N-p})^{-1})_{ii} - ((z_2 - \sigma^2 g(z_2) - A_{N-p})^{-1})_{ii}|^2) \\
&\quad + \sigma^4 \int \left| \frac{1}{z_1 - x} - \frac{1}{z_2 - x} \right|^2 d\lambda(x).
\end{aligned}$$

From the resolvent identity,

$$\begin{aligned}
&(z_1 - \sigma^2 g(z_1) - A_{N-p})^{-1} - (z_2 - \sigma^2 g(z_2) - A_{N-p})^{-1} \\
&= (z_2 - z_1 - \sigma^2 (g(z_2) - g(z_1))) (z_1 - \sigma^2 g(z_1) - A_{N-p})^{-1} (z_2 - \sigma^2 g(z_2) - A_{N-p})^{-1},
\end{aligned}$$

thus,

$$|((z_1 - \sigma^2 g(z_1) - A_{N-p})^{-1})_{ii} - ((z_2 - \sigma^2 g(z_2) - A_{N-p})^{-1})_{ii}| \leq \frac{1}{\varepsilon^2} (1 + \frac{\sigma^2}{\varepsilon^2}) |z_1 - z_2|$$

and thus,

$$\frac{1}{N-p} \sum_{i=1}^{N-p} (|((z_1 - \sigma^2 g(z_1) - A_{N-p})^{-1})_{ii} - ((z_2 - \sigma^2 g(z_2) - A_{N-p})^{-1})_{ii}|^2) \leq \frac{1}{\varepsilon^4} (1 + \frac{\sigma^2}{\varepsilon^2})^2 |z_1 - z_2|^2.$$

Since moreover

$$\int \left| \frac{1}{z_1 - x} - \frac{1}{z_2 - x} \right|^2 d\lambda(x) \leq |\Im z_1|^2 |z_2|^2 |z_1 - z_2|^2 \leq \frac{1}{\varepsilon^4} |z_1 - z_2|^2$$

(108) readily follows. The tightness follows from Kolmogorov's criterion (see [12]). \square

Proposition 14 and Proposition 12 readily yield Theorem 6.

4.5 Proof of Theorem 7

Theorem 7 follows from Theorem 6 once we proved that $A_{\theta, \nu, N}$ converge to $A_{\theta, \nu}$.

Lemma 16. *Assume that the matrix A_N satisfies (A') with A_{N-p} diagonal. Then, the sequence $(A_{\theta, \nu, N})_N$ defined by (15) converges to $A_{\theta, \nu}$ defined by (18).*

Proof. Denote by d_i the eigenvalues of A_{N-p} .

$$\begin{aligned} A_{\theta, \nu, N} &= \frac{1}{N-p} \sum_{i=1}^{N-p} (H''(\theta)(\theta - d_i)^{-1} - H'(\theta)(\theta - d_i)^{-2})^2 \\ &\xrightarrow{N \rightarrow \infty} (H''(\theta))^2 \int \frac{1}{(\theta - x)^2} d\nu(x) - 2H'(\theta)H''(\theta) \int \frac{1}{(\theta - x)^3} d\nu(x) \\ &\quad + (H'(\theta))^2 \int \frac{1}{(\theta - x)^4} d\nu(x) \\ &= -(H''(\theta))^2 g'_\nu(\theta) - H'(\theta)H''(\theta)g''_\nu(\theta) - \frac{1}{6}(H'(\theta))^2 g'''_\nu(\theta) \end{aligned}$$

Using, $H'(\theta) = 1 + \sigma^2 g'_\nu(\theta)$ and $H''(\theta) = \sigma^2 g''_\nu(\theta)$, we obtain the formula for $A_{\theta, \nu}$ given in (18). \square

5 Appendix: Poincaré inequality and concentration phenomenon

A probability μ satisfies a Poincaré inequality if for any C^1 function $f : \mathbb{R} \rightarrow \mathbb{C}$ such that f and f' are in $L^2(\mu)$,

$$\mathbf{V}(f) \leq C_{PI} \int |f'|^2 d\mu,$$

with $\mathbf{V}(f) = \int |f - \int f d\mu|^2 d\mu$.

If the law of a random variable X satisfies the Poincaré inequality with constant C_{PI} then, for any fixed $\alpha \neq 0$, the law of αX satisfies the Poincaré inequality with constant $\alpha^2 C_{PI}$.

Assume that probability measures μ_1, \dots, μ_M on \mathbb{R} satisfy the Poincaré inequality with constant $C_{PI}(1), \dots, C_{PI}(M)$ respectively. Then the product measure $\mu_1 \otimes \dots \otimes \mu_M$ on \mathbb{R}^M satisfies the Poincaré inequality with constant $C_{PI}^* = \max_{i \in \{1, \dots, M\}} C_{PI}(i)$ in the sense that for any differentiable function f such that f and its gradient $\text{grad} f$ are in $L^2(\mu_1 \otimes \dots \otimes \mu_M)$,

$$\mathbf{V}(f) \leq C_{PI}^* \int \|\text{grad} f\|_2^2 d\mu_1 \otimes \dots \otimes \mu_M$$

with $\mathbf{V}(f) = \int |f - \int f d\mu_1 \otimes \dots \otimes \mu_M|^2 d\mu_1 \otimes \dots \otimes \mu_M$.

Lemma 17. *Lemma 4.4.3 and Exercise 4.4.5 in [1] or Chapter 3 in [27]. Let \mathbb{P} be a probability measure on \mathbb{R}^M which satisfies a Poincaré inequality with constant C_{PI} . Then there exists $K_1 > 0$ and $K_2 > 0$ such that, for any Lipschitz function F on \mathbb{R}^M with Lipschitz constant $|F|_{Lip}$,*

$$\forall \epsilon > 0, \mathbb{P}(|F - \mathbb{E}_{\mathbb{P}}(F)| > \epsilon) \leq K_1 \exp\left(-\frac{\epsilon}{K_2 \sqrt{C_{PI}} |F|_{Lip}}\right).$$

Acknowledgements

The authors want to thank an anonymous referee who encouraged them to establish more general results and to provide a better readability, which led to an overall improvement of the paper.

References

- [1] G.W. Anderson, A. Guionnet and O. Zeitouni. *An introduction to random matrices*, Cambridge University Press, Cambridge, 2010.

- [2] Z. D. Bai and J. W. Silverstein. No eigenvalues outside the support of the limiting spectral distribution of large-dimensional sample covariance matrices. *Ann. Probab.* 26, 316–345, 1998.
- [3] Z. Bai and J. Silverstein, *Spectral analysis of large dimensional random matrices* (Second edition. Springer Series in Statistics. Springer, New York, 2010).
- [4] Z. Bai and J. Yao. Central limit theorem for eigenvalues in a spiked population model. *Ann. Inst. H. Poincaré Probab. Statist.* Volume 44, Number 3, 447–474, 2008.
- [5] Z. D. Bai and Y. Q. Yin. Necessary and sufficient conditions for almost sure convergence of the largest eigenvalue of a Wigner matrix. *Ann. Probab.*, 16(4), 1729–1741, 1988.
- [6] J. Baik, G. Ben Arous and S. Péché. Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices. *Ann. Probab.* 33 no. 5, 1643–1697, 2005.
- [7] S. T. Belinschi and H. Bercovici, A new approach to subordination results in free probability, *J. Anal. Math.* 101, 357–365, 2007.
- [8] S. Belinschi, M. Capitaine. Spectral properties of polynomials in independent Wigner and deterministic matrices *J. Funct. Anal.*, Volume 273, Issue 12, 3901–3963, 2017.
- [9] F. Benaych-Georges and A. Knowles : Lectures on the local semicircle law for Wigner matrices. *Advanced Topics in Random Matrices*, Florent Benaych-Georges, Charles Bordenave, Mireille Capitaine, Catherine Donati-Martin, Antti Knowles (edited by F. Benaych-Georges, D. Chafai, S. Péché, B. de Tilière) Panoramas et synthèses 53, 2018.
- [10] P. Biane. On the free convolution with a semi-circular distribution. *Indiana Univ. Math. J.*, 46(3), 705–718, 1997.
- [11] P. Biane, Processes with free increments, *Math. Z.* 227, 143–174, 1998.
- [12] P. Billingsley. *Convergence of probability measures*. Second edition. Wiley Series in Probability and Statistics: Probability and Statistics. 1999.
- [13] A. Bloemendal, L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Isotropic local laws for sample covariance and generalized wigner matrices. *Electron. J. Probab.*, 19(33):1–53, 2014.

- [14] M. Capitaine. Additive/multiplicative free subordination property and limiting eigenvectors of spiked additive deformations of Wigner matrices and spiked sample covariance matrices *J. Theor. Probab.*, Volume 26 (3), 595–648, 2013.
- [15] M. Capitaine. Limiting eigenvectors of outliers for Spiked Information-Plus-Noise type matrices *Séminaire de Probabilités XLIX*, 119-164, 2018).
- [16] M. Capitaine. Nonuniversality of fluctuations of outliers for Hermitian polynomials in a complex Wigner matrix and a spiked diagonal matrix. *Random Matrices Theory Appl.*, <https://doi.org/10.1142/S2010326320500136>, 2019.
- [17] M. Capitaine, C. Donati-Martin. Spectrum of deformed random matrices and free probability. *Advanced Topics in Random Matrices*, Florent Benaych-Georges, Charles Bordenave, Mireille Capitaine, Catherine Donati-Martin, Antti Knowles (edited by F. Benaych-Georges, D. Chafai, S. Péché, B. de Tilière) Panoramas et synthèses 53, 2018.
- [18] M. Capitaine, C. Donati-Martin and D. Féral : The largest eigenvalues of finite rank deformation of large Wigner matrices: convergence and nonuniversality of the fluctuations. *Ann. Probab.* 37, no. 1, 1–47, 2009.
- [19] M. Capitaine, C. Donati-Martin and D. Féral. Central limit theorems for eigenvalues of deformations of Wigner matrices. *Ann. Inst. H. Poincaré Probab. Statist.* 48, 1, 107-133, 2012.
- [20] M. Capitaine, C. Donati-Martin, D. Féral and M. Février. Free convolution with a semicircular distribution and eigenvalues of spiked deformations of Wigner matrices. *Electron. J. Probab.* 16 no. 64, 1750–1792, 2011.
- [21] M. Capitaine, S. Péché. Fluctuations at the edges of the spectrum of the full rank deformed G.U.E. *Probab. Theory Relat. Fields*, Volume 165, Issue 1, 117–161, 2016.
- [22] L. Erdős, B. Schlein, B. and H.T Yau. Local semicircle law and complete delocalization for Wigner random matrices. *Comm. Math. Phys.* 287(2), 641–655, 2009.
- [23] L. Erdős, B. Schlein and H.-T. Yau. Semicircle law on short scales and delocalization of eigenvectors for Wigner random matrices. *Ann. Probab.*, 37(3):815–852, 2009.

- [24] Z. Füredi and J. Komlós. The eigenvalues of random symmetric matrices. *Combinatorica*, 1(3), 233–241, 1981.
- [25] D. Voiculescu, K. J. Dykema, and A. Nica. *Free random variables*, American Mathematical Society, Providence, RI, 1992.
- [26] A. Knowles and J. Yin. Eigenvector distribution of Wigner matrices. *Probab. Theory and Relat. Fields*. 155(3-4), 543–582, 2013.
- [27] M. Ledoux. *The Concentration of Measure Phenomenon*. Mathematical Surveys and Monographs, Volume 89, A.M.S, 2001.
- [28] J. O. Lee and J. Yin. A necessary and sufficient condition for edge universality of Wigner matrices. *Duke Math. J.* Volume 163, Number 1, 117–173, 2014.
- [29] MATHIAS R. (1993). The Hadamard operator norm of a circulant and applications, *SIAM J. Matrix Anal. Appl.* **14**, 1152–1167.
- [30] J.A. Mingo and R. Speicher. *Free probability and Random matrices*. Fields Institute Monographs Originally published by American Mathematical Society ISSN: 1069-5273. Springer.
- [31] J. Najim, J. Yao. Gaussian fluctuations for linear spectral statistics of large random covariance matrices. *The Annals of App. Prob.* 26, 1837–1887, 2016.
- [32] L. A. Pastur. On the spectrum of random matrices. *Theoretical and Mathematical Physics*, Volume 10, Issue 1, 67–74, 1972.
- [33] S. Péché. The largest eigenvalue of small rank perturbations of Hermitian random matrices. *Probab. Theory Relat. Fields* 134, no. 1, 127-173, 2006.
- [34] T. Shirai. Limit theorems for random analytic functions and their zeros. In *Functions in number theory and their probabilistic aspects*, RIMS Kôkyûroku Bessatsu, B34, 335–359. Res. Inst. Math. Sci. (RIMS), Kyoto, 2012.
- [35] A. Soshnikov. Universality at the edge of the spectrum in Wigner random matrices. *Comm. Math. Phys.* 207, 697–733, 1999.
- [36] T. Tao and V. Vu. Random matrices: universal properties of eigenvectors. *Random Matrices Theory Appl.* 1(1), 1150,001, 27, 2012.

- [37] C. Tracy, H. Widom. Level-spacing distributions and the Airy kernel. *Comm. Math. Phys.* 159, 151–174, 1994.
- [38] D. Voiculescu. The analogues of entropy and of Fisher’s information measure in free probability theory. I, *Comm. Math. Phys.* 155, 71–92, 1993.
- [39] E. P. Wigner. Characteristic vectors of bordered matrices with infinite dimensions. *Ann. of Math. (2)* 62, 548–564, 1955.
- [40] E. P. Wigner. On the distribution of the roots of certain symmetric matrices. *Ann. of Math. (2)* 67, 325–327, 1958.