

NONZERO-SUM STOCHASTIC DIFFERENTIAL GAMES WITH STOPPING TIMES AND FREE BOUNDARY PROBLEMS⁽¹⁾

BY

ALAIN BENSOUSSAN AND AVNER FRIEDMAN

ABSTRACT. One is given a diffusion process and two payoffs which depend on the process and on two stopping times τ_1, τ_2 . Two players are to choose their respective stopping times τ_1, τ_2 so as to achieve a Nash equilibrium point. The problem whether such times exist is reduced to finding a "regular" solution (u_1, u_2) of a quasi-variational inequality. Existence of a solution is established in the stationary case and, for one space dimension, in the nonstationary case; for the latter situation, the solution is shown to be regular if the game is of zero sum.

Introduction. We consider in this paper a nonzero-sum stochastic differential game with two players, where the decision variables of the players are stopping times. The system evolves according to a stochastic differential equation. The player who decides to stop first prevents the other from continuing and payoffs are computed for each of them. We are looking for a Nash equilibrium point. The model is a natural generalization of the two player zero-sum game studied in Friedman [14]. In Chapter 1 we give sufficient conditions for existence of a Nash equilibrium point. These conditions are stated in terms of functions $u_1(x, t)$, $u_2(x, t)$ which represent the payoffs for the Nash equilibrium point when the system starts at time t in position $x \in R^n$:

Suppose there exist functions u_1, u_2 satisfying the following system of differential inequalities:

- (0.1) $u_i(x, T) = h_i(x),$
- (0.2) $u_i \leq \phi_i,$
- (0.3) if $u_j = \phi_j$ for $j \neq i$, then $u_i = \psi_i,$
if $u_j < \phi_j$ for $j \neq i$, then

Received by the editors June 13, 1975.

AMS (MOS) subject classifications (1970). Primary 93E05, 49C10; Secondary 35J99, 35K60, 35K99, 60G40, 60H10, 93E20.

Key words and phrases. Stochastic differential games, stochastic differential equations, stopping time, payoff, Nash point, variational inequality, quasi-variational inequality, free boundary problem, nonzero-sum game, zero-sum game.

⁽¹⁾ This work was partially supported by National Science Foundation Grant GP-35347X.

© American Mathematical Society 1977

$$\begin{aligned}
 & -\frac{\partial u_i}{\partial t} + A(t)u_i \leq f_i, \\
 (0.4) \quad & (u_i - \phi_i) \left(-\frac{\partial u_i}{\partial t} + A(t)u_i - f_i \right) = 0,
 \end{aligned}$$

where $A(t)$ is a second order elliptic differential operator in x and ϕ_i, ψ_i, f_i, h_i are given functions (in terms of which the game is defined); suppose also that the u_i satisfy some regularity conditions. Then there exists a Nash equilibrium point for the initial problem.

In Chapter 2 we study the existence of solutions for (0.1)–(0.4) in the stationary case (i.e., the u_i do not depend on t). We formulate the problem as a quasi-variational inequality, and then prove an existence theorem based upon the study of fixed points of a monotone increasing mapping. The regularity as well as the uniqueness of the solution is left open, the latter being probably false.

In Chapter 3 we study the existence of a solution for (0.1)–(0.4) in the (nonstationary) case $n = 1$. Here again an existence theorem is established using a fixed point theorem for a monotone increasing mapping. Under suitable conditions it is proved that the free boundary consists of two monotone and continuous curves $x = s_i(t)$, $s_1(t) < s_2(t)$.

The fixed point theorem used in Chapters 2, 3 is due to Tartar [23]. It was also applied in [20] in order to solve a parabolic quasi-variational inequality arising from a Stefan problem of melting of ice with variable latent heat.

In Chapters 4, 5 we use the method of nonlinear Volterra integral equations in order to solve parabolic free boundary problems in one space dimension. In Chapter 4 we consider a Stefan type problem involving two “temperatures” θ_1 and θ_2 . By classical methods ([12], [13]) one establishes the existence of a unique regular solution as long as θ_{1x}, θ_{2x} are a priori bounded. When the initial “temperatures” are nonnegative, a priori bounds can be obtained without difficulty (see the proof of Theorem 4.1). When the initial “temperatures” take also negative values, it is harder to establish the necessary a priori estimates (see proofs of Theorems 4.3, 4.4). Our results in this case are extensions of the works of Van Moerbeke [24] and Friedman [16].

We also derive in Chapter 4 an estimate on $s_2(t) - s_1(t)$ as $t \rightarrow \infty$.

In Chapter 5 we apply the methods of Chapter 4 to a somewhat different system of Stefan type problem which corresponds to the zero-sum game (in the case $n = 1$); this system is in some sense a special case of the system studied in Chapter 4. We deduce that not only are the functions $u_1(x, t)$ and $u_2(x, t)$ regular (which is already known for general $n \geq 1$, by Friedman [14]) but also the free boundaries are regular curves.

CHAPTER 1. THE DIFFERENTIAL GAME WITH STOPPING TIMES AND
CHARACTERIZATION OF A NASH POINT BY A SYSTEM OF DIFFERENTIAL
INEQUALITIES

1.1. Assumptions; notation. Let (Ω, \mathcal{A}, P) be a probability space and \mathcal{F}_t , $t \geq 0$, be an increasing family of sub σ -algebras of \mathcal{A} . Let $b(t)$ be an n -dimensional (standard) Brownian motion with respect to the family \mathcal{F}_t . Let $g(x, t)$ be an n -vector defined on $R^n \times [0, \infty)$ and let $\sigma(x, t)$ be an $n \times n$ matrix defined on $R^n \times [0, \infty)$, such that

- (1.1) g is continuous and bounded,
- (1.2) $|g(x, t) - g(x', t)| \leq C|x - x'|$ for all x, x' (C constant),
- (1.3) σ is continuous and bounded,
- (1.4) $\partial\sigma(x, t)/\partial x$ is measurable and bounded,
- (1.5) σ^{-1} is continuous and bounded.

We denote by $y_{xt}(s)$, $s \geq t$, the solution of the stochastic differential equation

$$(1.6) \quad dy_{xt}(s) = g(y_{xt}(s), s) ds + \sigma(y_{xt}(s), s) db(s), \quad y_{xt}(t) = x.$$

In the following all the stopping times which we use are considered to be stopping times with respect to \mathcal{F}_t .

Let T be a positive number and let $f_i(x, t)$, $\phi_i(x, t)$, $\psi_i(x, t)$ ($i = 1, 2$) be functions such that

$$(1.7) \quad \begin{aligned} &f_i, \phi_i, \psi_i \text{ are continuous and bounded} \\ &\text{in } R^n \times [0, T], f_i \in L^2(R^n \times (0, T)), \\ &\psi_i \leq \phi_i \text{ for all } x, t \text{ in } R^n \times [0, T] \\ &(i = 1, 2). \end{aligned}$$

1.2. Definition of a Nash point in the nonstationary case. We consider two players such that each of them may decide to stop the process y_{xt} starting at time $t \leq T$. Given two functions $h_1(x)$, $h_2(x)$ satisfying

$$(1.8) \quad h_i \text{ are continuous and bounded,}$$

we define the payoff functions $J_{xt}^i(\tau_1, \tau_2)$, where τ_i are stopping times such that $t \leq \tau_i \leq T$, as follows:

$$(1.9) \quad \begin{aligned} J_{xt}^i(\tau_1, \tau_2) = E &\left[\int_t^{\tau_1 \wedge \tau_2} f_i(y_{xt}(s), s) ds + \chi_{\tau_1 < \tau_2} \phi_i(y_{xt}(\tau_1), \tau_1) \right. \\ &\left. + \chi_{\tau_1 > \tau_2} \psi_i(y_{xt}(\tau_2), \tau_2) + \chi_{\tau_1 = \tau_2 = T} h_i(y_{xt}(T)) \right] \\ &(j \neq i) \end{aligned}$$

where χ_A is the indicator function of A .

Our problem is to find a Nash point for the J_{xt}^i , i.e., to find $\hat{\tau}_1, \hat{\tau}_2$ such that

$$(1.10) \quad J_{xt}^1(\hat{\tau}_1, \hat{\tau}_2) \leq J_{xt}^1(\tau_1, \hat{\tau}_2), \quad J_{xt}^2(\hat{\tau}_1, \hat{\tau}_2) \leq J_{xt}^2(\hat{\tau}_1, \tau_2) \quad \text{for any } \tau_1, \tau_2.$$

1.3. System of differential inequalities. For t fixed, $t \in [0, T]$, we define the second order differential operator $A(t)$ by

$$(1.11) \quad A(t)w = - \sum_{i,j=1}^n \frac{1}{2} a_{ij}(x, t) \frac{\partial^2 w}{\partial x_i \partial x_j} - \sum_{j=1}^n g_j(x, t) \frac{\partial w}{\partial x_j}$$

where the $a_{ij}(x, t)$ are the components of the matrix $a(x, t) = \sigma \sigma^*(x, t)$, $\sigma^* = \text{transpose of } \sigma$, and $g_j(x, t)$ are the components of $g(x, t)$. Let $Q = R^n \times (0, T)$.

We introduce the following problem: Find two functions $u_1(x, t)$, $u_2(x, t)$ such that

$$(1.12) \quad u_i \text{ is continuous and bounded in } \bar{Q}, u_i \in L^2(0, T; H^1(R^n));$$

$$(1.13) \quad u_i(x, T) = h_i(x) \quad (x \in R^n);$$

$$(1.14) \quad u_i(x, t) \leq \phi_i(x, t) \text{ in } Q;$$

$$(1.15) \quad \text{if } u_j(x, t) = \phi_j(x, t) \text{ for } j \neq i \text{ and some } (x, t) \text{ in } Q, \\ \text{then } u_i(x, t) = \psi_i(x, t);$$

if $\Sigma_i = \{(x, t) \in Q; u_j(x, t) < \phi_j(x, t) \text{ for } j \neq i\}$, then

$$(1.16) \quad \frac{\partial u_i}{\partial t} - A(t)u_i \in L^2(\Sigma_i), \quad - \frac{\partial u_i}{\partial t} + A(t)u_i \leq f_i \quad \text{a.e. in } \Sigma_i, \\ (u_i - \phi_i) \left[- \frac{\partial u_i}{\partial t} + A(t)u_i - f_i \right] = 0 \quad \text{a.e. in } \Sigma_i.$$

We will refer to (1.12)–(1.16) as the system of differential inequalities (associated with the stochastic differential game).

To u_i we correspond a set C_i defined by

$$(1.17) \quad C_i = \{(x, t) \in Q; u_i(x, t) < \phi_i(x, t)\},$$

which by virtue of the continuity properties of the functions u_i and ϕ_i is an open subset of Q . (Notice that $C_1 = \Sigma_2$, $C_2 = \Sigma_1$.)

1.4. Sufficiency criterion for existence of a Nash point. We define the exit time of C_i by

$$(1.18) \quad \hat{\tau}_i = \inf\{s; t \leq T, y_{xt}(s) \notin C_i\} \quad \text{if such numbers } s \text{ exist,} \\ = T \quad \text{otherwise.}$$

THEOREM 1.1. *Let the assumptions (1.1)–(1.5), (1.7) and (1.8) hold. If there exist functions u_1, u_2 satisfying (1.12)–(1.16), then for any x, t , the $\hat{\tau}_i$ given by (1.18) form a Nash point for the payoff functions J_{xt}^i . Furthermore,*

$$(1.19) \quad u_i(x, t) = J_{xt}^i(\hat{\tau}_1, \hat{\tau}_2).$$

PROOF. We shall use the following slight extension of Ito's formula. Let

$$\Phi(x, t) \in L^2(0, T; H^1(R^n)), \quad \left[\frac{\partial \Phi}{\partial t} - A(t)\Phi \right] \in L^2(Q),$$

Φ continuous and bounded on \bar{Q} . Then for any stopping time θ , $t \leq \theta \leq T$, and for any $\varepsilon > 0$ we have

$$(1.20) \quad \begin{aligned} E\Phi(y_{xt}(\theta), \theta) &= E\Phi(y_{xt}(\theta \wedge (t + \varepsilon)), \theta \wedge (t + \varepsilon)) \\ &+ E \int_{\theta \wedge (t + \varepsilon)}^{\theta} \left(\frac{\partial \Phi}{\partial t} - A(t)\Phi \right)(y_{xt}(s), s) ds. \end{aligned}$$

For the proof of (1.20) see [3], [4]. Let us note that $\hat{\tau}_j$ ($j \neq i$) is the exit time of Σ_i . If we restrict ourselves to stopping times θ such that $t \leq \theta \leq \hat{\tau}_j$, then we may apply (1.20) with functions Φ such that

$$\Phi \in L^2(0, T; H^1(R^n)), \quad \left[\frac{\partial \Phi}{\partial t} - A(t)\Phi \right] \in L^2(\Sigma_i).$$

Indeed, for any $\lambda > 0$ let $\Sigma^\lambda = \{(x, t) \in \Sigma_i, \text{dist}((x, t), \partial \Sigma_i) > \lambda\}$, $\gamma_\lambda = \text{exit time from } \Sigma^\lambda$. Since we can modify Φ outside Σ^λ so that the modified function satisfies the regularity properties of (1.20), we conclude that (1.20) is valid with θ replaced by $\theta \wedge \gamma_\lambda$. Taking $\lambda \downarrow 0$, (1.20) follows.

Let $t \leq \tau_1, \tau_2 \leq T$ be stopping times. We shall prove that

$$(1.21) \quad u_1(x, t) \leq J_{xt}^1(\tau_1, \hat{\tau}_2), \quad u_2(x, t) \leq J_{xt}^2(\hat{\tau}_1, \tau_2).$$

Let $j = 1$ if $i = 2$ and $j = 2$ if $i = 1$. Applying (1.20) with $\Phi = u_i$ and $\theta = \tau_i \wedge \hat{\tau}_j$, we get

$$(1.22) \quad \begin{aligned} &Eu_i(y_{xt}(\tau_i \wedge \hat{\tau}_j), \tau_i \wedge \hat{\tau}_j) \\ &= Eu_i(y_{xt}(\tau_i \wedge \hat{\tau}_j \wedge (t + \varepsilon)), \tau_i \wedge \hat{\tau}_j \wedge (t + \varepsilon)) \\ &+ E \int_{\tau_i \wedge \hat{\tau}_j \wedge (t + \varepsilon)}^{\tau_i \wedge \hat{\tau}_j} \left(\frac{\partial u_i}{\partial t} + A(t)u_i \right)(y_{xt}(s), s) ds. \end{aligned}$$

From (1.16) it follows that

$$(1.23) \quad \begin{aligned} &E \int_{\tau_i \wedge \hat{\tau}_j \wedge (t + \varepsilon)}^{\tau_i \wedge \hat{\tau}_j} \left(-\frac{\partial u_i}{\partial t} + A(t)u_i \right)(y_{xt}(s), s) ds \\ &\leq E \int_{\tau_i \wedge \hat{\tau}_j \wedge (t + \varepsilon)}^{\tau_i \wedge \hat{\tau}_j} f_i(y_{xt}(s), s) ds. \end{aligned}$$

From (1.13) and (1.14) we also have

$$(1.24) \quad \begin{aligned} u_i(y_{xt}(\tau_i \wedge \hat{\tau}_j), \tau_i \wedge \hat{\tau}_j) &\leq \phi_i(y_{xt}(\tau_i), \tau_i) \chi_{\tau_i < \hat{\tau}_j} \\ &+ \chi_{\tau_i \geq \hat{\tau}_j, T > \hat{\tau}_j} \psi_i(y_{xt}(\hat{\tau}_j), \hat{\tau}_j) + \chi_{\tau_i = \hat{\tau}_j = T} h_i(y_{xt}(T)). \end{aligned}$$

Using (1.23), (1.24) in (1.22), letting $\varepsilon \rightarrow 0$ and noting that

$$Eu_i(y_{xt}(\tau_i \wedge \hat{\tau}_j \wedge (t + \varepsilon), \tau_i \wedge \hat{\tau}_j \wedge (t + \varepsilon))) \rightarrow u_i(x, t)$$

since u_i is continuous and bounded, we obtain (1.21).

It remains to prove that

$$(1.25) \quad u_i(x, t) = J_{xt}^i(\hat{\tau}_1, \hat{\tau}_2).$$

We use similar arguments to those above.

From (1.16) we get, if $j \neq i$,

$$(1.26) \quad \begin{aligned} E \int_{\hat{\tau}_i \wedge \hat{\tau}_j \wedge (t + \varepsilon)}^{\hat{\tau}_i \wedge \hat{\tau}_j} \left(-\frac{\partial u_i}{\partial t} + A(t)u_i \right)(y_{xt}(s), s) ds \\ = E \int_{\hat{\tau}_i \wedge \hat{\tau}_j}^{\hat{\tau}_i \wedge \hat{\tau}_j} f_i(y_{xt}(s), s) ds. \end{aligned}$$

From (1.13), (1.14) and the fact that if $\hat{\tau}_i < \hat{\tau}_j$ then

$$u_i(y_{xt}(\hat{\tau}_i), \hat{\tau}_i) = \phi_i(y_{xt}(\hat{\tau}_i), \hat{\tau}_i),$$

we get (1.24) with equality instead of inequality and $\hat{\tau}_i$ instead of τ_i . Taking $\tau_i = \hat{\tau}_i$ in (1.22), using (1.26) and (1.24) (with equality), and letting $\varepsilon \rightarrow 0$, we obtain (1.25), which completes the proof of the theorem.

1.5. Stationary case. We now assume that

$$(1.27) \quad g, \sigma, f_i, \phi_i, \psi_i \text{ do not depend on } t; T = +\infty.$$

We introduce a constant (discount factor) $\alpha > 0$ and define the payoff

$$(1.28) \quad \begin{aligned} J_x^i(\tau_1, \tau_2) = E \left[\int_0^{\tau_1 \wedge \tau_2} e^{-\alpha t} f_i(y_x(t)) dt + \psi_{\tau_i < \tau_j} \phi_i(y_x(\tau_i)) e^{-\alpha \tau_i} \right. \\ \left. + \chi_{\tau_i > \tau_j} \psi_i(y_x(\tau_j)) e^{-\alpha \tau_j} \right] \\ (j \neq i), \end{aligned}$$

where $y_x(t)$ denotes $y_{x0}(t)$. As in §1.3, we introduce functions $u_1(x)$, $u_2(x)$ (not depending on time) such that

$$(1.29) \quad u_i \text{ is continuous and bounded, } u_i \in H^1(R^n);$$

$$(1.30) \quad u_i \leq \phi_i \quad (x \in R^n);$$

$$(1.31) \quad \begin{aligned} \text{if } u_j(x) = \phi_j(x) \text{ for } j \neq i \text{ and some } x, \text{ then } u_i(x) = \psi_i(x); \\ \text{if } \Sigma_i = \{x \in R^n; u_j(x) < \phi_j(x)\} \ (j \neq i), \text{ then} \end{aligned}$$

$$(1.32) \quad \begin{aligned} Au_i \in L^2(\Sigma_i), \quad Au_i \leq f_i \quad \text{a.e. in } \Sigma_i, \\ (u_i - \phi_i)(Au_i - f_i) = 0 \quad \text{a.e. in } \Sigma_i. \end{aligned}$$

Here,

$$(1.33) \quad Aw = -\frac{1}{2} \sum_{i,j=1}^n a_{ij}(x) \frac{\partial^2 w}{\partial x_i \partial x_j} - \sum_{i=1}^n g_i(x) \frac{\partial w}{\partial x_i} + \alpha w$$

where (a_{ij}) is the matrix $\sigma\sigma^*(x)$ and $g_i(x)$ are the components of g . We define

$$(1.34) \quad C_i = \{x \in R^n; u_i(x) < \phi_i(x)\}$$

and state without proof a theorem analogous to Theorem 1.1:

THEOREM 1.2. *Under the assumptions of Theorem 1.1 and (1.27), if there exist functions $u_1(x)$, $u_2(x)$ satisfying (1.29)–(1.32), and if $\hat{\tau}_i = \inf\{s > 0, y_x(s) \notin C_i\}$, then the pair $(\hat{\tau}_1, \hat{\tau}_2)$ forms a Nash point for the payoff functions (1.28). Furthermore,*

$$(1.35) \quad u_i(x) = J_x^i(\hat{\tau}_1, \hat{\tau}_2).$$

It is assumed here that $\hat{\tau}_1, \hat{\tau}_2$ are well defined, i.e., that $y_x(s) \notin C_i$ for some time $s = s_i$ ($i = 1, 2$).

REMARK. The results of this chapter can be extended to N -person nonzero-sum game with payoffs

$$\begin{aligned} J_{xt}^i(\tau_1, \dots, \tau_N) = E \bigg[& \int_t^{\tau_1 \wedge \dots \wedge \tau_N} f_i(y_{xt}(s), s) ds + \chi_{\tau_i < \bigwedge_{j \neq i} \tau_j} \phi_i(y_{xt}(\tau_i), \tau_i) \\ & + \chi_{\tau_i > \bigwedge_{j \neq i} \tau_j, T > \bigwedge_{j \neq i} \tau_j} \psi_i \left(y_{xt} \left(\bigwedge_{j \neq i} \tau_j \right), \bigwedge_{j \neq i} \tau_j \right) \\ & + \chi_{\tau_1 = \dots = \tau_N = T} h_i(y_{xt}(T)) \bigg]. \end{aligned}$$

In the condition (1.15) one has to assume that $u_j(x, t) = \phi_j(x, t)$ for *some* $j \neq i$, and in the definition of Σ_i in (1.16) $u_j(x, t) < \phi_j(x, t)$ for *all* $j \neq i$. A Nash equilibrium point $(\hat{\tau}_1, \dots, \hat{\tau}_N)$ is defined by

$$J_i(\hat{\tau}_1, \dots, \hat{\tau}_N) \leq J_i(\hat{\tau}_1, \dots, \hat{\tau}_{i-1}, \tau_i, \hat{\tau}_{i+1}, \dots, \hat{\tau}_N) \quad (1 \leq i \leq N)$$

for any τ_1, \dots, τ_N . Theorems 1.1, 1.2 and their proofs readily extend to this case.

CHAPTER 2. STATIONARY QUASI-VARIATIONAL INEQUALITIES

In this Chapter we consider the stationary case for any dimension n .

2.1. Assumptions; notation. Consider a second order differential operator

$$(2.1) \quad Aw = -\frac{1}{2} \sum_{i,j=1}^n \frac{\partial}{\partial x_i} \left(a_{ij} \frac{\partial w}{\partial x_j} \right) + \sum_{j=1}^n a_j \frac{\partial w}{\partial x_j} + a_0 w,$$

where

$$(2.2) \quad \begin{cases} a_{ij}(x), a_j(x) \text{ are functions in } L^\infty(R^n), \\ a_0(x) \geq \alpha > 0; a_{ij} = a_{ji}. \end{cases}$$

When

$$(2.3) \quad a_j(x) = \frac{1}{2} \sum_i \frac{\partial a_{ij}}{\partial x_i} - g_j(x), \quad a_0(x) = \alpha,$$

we get the operator A of (1.33). We shall make the assumptions:

$$(2.4) \quad \sum_{i,j} a_{ij} \xi_i \xi_j \geq \gamma \sum_i \xi_i^2 \quad \text{for all real } \xi_i, \gamma \text{ positive constant,}$$

$$(2.5) \quad a_j \text{ is differentiable and } a_0(x) - \frac{1}{2} \sum_j \frac{\partial a_j}{\partial x_j} \geq \beta > 0 \quad (\beta \text{ constant}).$$

Let $V = H^1(R^n)$. We define on the Hilbert space V a bilinear continuous form $a(u, v)$:

$$(2.6) \quad a(u, v) = \frac{1}{2} \sum_{i,j} \int_{R^n} a_{ij} \frac{\partial u}{\partial x_j} \frac{\partial v}{\partial x_i} dx + \sum_j \int_{R^n} a_j \frac{\partial u}{\partial x_j} v dx + \int_{R^n} a_0 uv dx.$$

Noting that

$$(2.7) \quad \begin{aligned} a(u) = a(u, u) &= \frac{1}{2} \sum_{i,j} \int_{R^n} a_{ij} \frac{\partial u}{\partial x_j} \frac{\partial u}{\partial x_i} dx + \int_{R^n} \left(a_0 - \frac{1}{2} \sum_j \frac{\partial a_j}{\partial x_j} \right) u^2 dx \\ &\geq \beta \int_{R^n} u^2 dx, \end{aligned}$$

we conclude that

$$(2.8) \quad a(u) = 0 \text{ implies } u = 0.$$

We consider functions f_i, ϕ_i, ψ_i ($i = 1, 2$) satisfying (1.7) and (1.27). For u_1, u_2 in $L^2(R^n)$ define the sets

$$(2.9) \quad \begin{aligned} K_i(u_j) &= \{v \in V; v \leq \phi_i \text{ a.e. and a.e. in } x, \\ &\quad \text{if } u_j(x) \geq \phi_j(x) \text{ then } v(x) = \psi_i(x)\} \quad (j \neq i). \end{aligned}$$

Clearly $K_i(u_j)$ is a closed convex subset of V . It is not empty since $v = \psi_i \in K_i(u_j)$. We denote by $(,)$ the scalar product in $L^2(R^n)$ and by $((,))$ the scalar product in V .

2.2. System of Q.V.I. We consider the following problem: find u_1, u_2 in V such that

$$(2.10) \quad \begin{aligned} u_1 &\in K_1(u_2), & u_2 &\in K_2(u_1), \\ a(u_1, v - u_1) &\geq (f_1, v - u_1) & \text{for every } v \in K_1(u_2), \\ a(u_2, v - u_2) &\geq (f_2, v - u_2) & \text{for every } v \in K_2(u_1). \end{aligned}$$

We call (2.10) a system of *quasi-variational inequalities* (Q.V.I.). We can formulate (2.10) as one Q.V.I. in $V^2 = V \times V$, defining for $u = (u_1, u_2)$, $v = (v_1, v_2)$, the bilinear form

$$\tilde{a}(u, v) = \sum_{i=1}^2 a(u_i, v_i)$$

and setting $\tilde{f} = (f_1, f_2)$,

$$K(u) = \{v = (v_1, v_2); v_1 \in K_1(v_2), v_2 \in K_2(v_1)\}.$$

With this notation (2.10) is equivalent to

$$(2.11) \quad \tilde{a}(u, v - u) \geq (\tilde{f}, v - u) \quad \text{for every } v \in K(u), u \in K(u).$$

In (2.11) by $(\tilde{f}, v - u)$ we mean $\sum_{i=1}^2 (f_i, v_i - u_i)$.

Now (2.11) is a Q.V.I. as defined in [2], [4], [5], [6]. Systems of Q.V.I. have already been considered by Bensoussan-Lions [7]. However, the situation here differs from the one considered by these authors, because the convex set $K(u)$ is decreasing with u instead of increasing. Let us state this more precisely. We consider in V^2 the natural order relationship

$$(2.12) \quad v \geq v' \quad \text{if } v_i(x) \geq v'_i(x) \text{ a.e., for } i = 1, 2.$$

One easily checks that

$$(2.13) \quad K(v) \subset K(v') \quad \text{if } v \geq v'.$$

We say that the convex set *decreases*, and that the Q.V.I. (2.11) is a decreasing Q.V.I. For such Q.V.I. the general existence results of Bensoussan-Lions [7] and of Tartar [23] cannot be applied. Decreasing Q.V.I. have already been considered by Bensoussan-Lions [8], but with assumptions on the continuity of $K(v)$ with respect to v , which are not satisfied here.

Before giving existence results for (2.10), we shall show how (2.10) is related to (1.29)–(1.32). We have

THEOREM 2.1. *Let (2.3) hold. Suppose there exists a solution (u_1, u_2) of (2.10) such that the functions u_i are continuous and bounded. Define Σ_i as in (1.32) and suppose that Σ_i has a smooth boundary, that the complement of Σ_i is not of measure 0, and that $Au_i \in L^2(\Sigma_i)$. Then the u_i 's form a solution of (1.29)–(1.32).*

PROOF. Let S_i be the complement of Σ_i . Since $u_i \in K_i(u_j)$ ($j \neq i$), we have $u_i(x) = \psi_i(x)$ a.e. in S_i . By continuity we have $u_i = \psi_i$ everywhere in S_i , which proves (1.31).

Noting that if $v \in K_i(u_j)$, $j \neq i$, then $v = u_i$ on S_i and we get from (2.10)

$$(2.14) \quad \frac{1}{2} \int_{\Sigma_i} \left[\sum_{k,l=1}^n a_{kl} \frac{\partial u_i}{\partial x_l} \frac{\partial (v - u_i)}{\partial x_k} + \sum_{k=1}^n a_k \frac{\partial u_i}{\partial x_k} (v - u_i) + a_0 u_i (v - u_i) \right] dx \\ \geq \int_{\Sigma_i} f_i (v - u_i) dx.$$

Integrating by parts in the first integral and noting that $v = u_i$ on $\partial \Sigma_i$ we get, since $Au_i \in L^2(\Sigma_i)$,

$$(2.15) \quad \int_{\Sigma_i} (Au_i - f_i)(v - u_i) dx \geq 0 \quad \text{for any } v \in K_i(u_j) \ (j \neq i).$$

Let Θ be an open ball such that Θ is contained in Σ_i . Then we can find a family of smooth functions θ_ε , $0 \leq \theta_\varepsilon \leq 1$, such that $\theta_\varepsilon = 0$ outside Σ_i and $\theta_\varepsilon(x) \rightarrow 1$ in Θ , $\theta_\varepsilon \rightarrow 0$ outside Θ if $\varepsilon \rightarrow 0$. We may take in (2.15) $v = u_i - \theta_\varepsilon$, ε small, which is admissible. We then get

$$\int_{\Theta} \theta_\varepsilon (Au_i - f_i) dx + \int_{\Sigma_i - \Theta} \theta_\varepsilon (Au_i - f_i) dx \leq 0.$$

Letting $\varepsilon \downarrow 0$, we obtain $\int_{\Theta} (Au_i - f_i) dx \leq 0$, and since Θ is arbitrary,

$$(2.16) \quad Au_i - f_i \leq 0 \quad \text{a.e. in } \Sigma_i.$$

Now let Θ be as above and take in (2.15) the admissible function $v = \phi_i \theta_\varepsilon + (1 - \theta_\varepsilon)u_i$. Then

$$\begin{aligned} \int_{\Theta} (Au_i - f_i) [(\phi_i \theta_\varepsilon - u_i) + (1 - \theta_\varepsilon)u_i] dx \\ + \int_{\Sigma_i - \Theta} (Au_i - f_i) [\phi_i \theta_\varepsilon - \theta_\varepsilon u_i] dx \geq 0. \end{aligned}$$

Letting $\varepsilon \downarrow 0$, we get $\int_{\Theta} (Au_i - f_i)(\phi_i - u_i) dx \geq 0$. But from (2.16) and $u_i \leq \phi_i$, we also have the reverse inequality. Hence

$$(2.17) \quad \int_{\Theta} (Au_i - f_i)(\phi_i - u_i) dx = 0.$$

Since again Θ is arbitrary, the last relation in (1.32) follows. We have thereby completed the proof of (1.29)–(1.32).

REMARK. Theorem 2.1 extends to the case of N -person game. In the definition (2.9), $K_i(u_j)$ should be replaced by $K_i(u_1, \dots, u_{i-1}, u_{i+1}, \dots, u_N)$ and the assertion $v(x) = \psi_i(x)$ a.e. is required to follow from $u_j(x) \geq \phi_j(x)$ for some $j \neq i$.

2.3. Existence theorem. We shall now give an existence result for a system of Q.V.I. (2.10), making the following restrictive assumptions:

$$(2.18) \quad \begin{aligned} &f_i, \phi_i, \psi_i \text{ are continuous and bounded;} \\ &f \in L^2(R^n), \phi_i \in V, \psi_i \in V, \psi_i \leq \phi_i, \\ &\text{and } a(\psi_i, \chi) - (f_i, \chi) \leq 0 \text{ for any} \\ &\chi \in V, \chi \geq 0 \ (i = 1, 2). \end{aligned}$$

Notice that the last inequality holds if $\psi_i \in H^2(R^n)$ and $A\psi_i \leq f_i$.

THEOREM 2.2. *Let the assumptions (2.2)–(2.5), (2.18) hold. Then there exists a solution u_1, u_2 of the Q.V.I. (2.10) satisfying $u_i \geq \psi_i$ for $i = 1, 2$. Moreover there exist two pairs of solutions $(\underline{u}_1, \bar{u}_2)$ and $(\bar{u}_1, \underline{u}_2)$ such that if u_1, u_2 is a solution then*

$$(2.19) \quad \psi_1 \leq \underline{u}_1 \leq u_1 \leq \bar{u}_1 \leq \phi_1, \quad \psi_2 \leq \underline{u}_2 \leq u_2 \leq \bar{u}_2 \leq \phi_2.$$

PROOF. We shall define several mappings. Let $w_1, w_2 \in L^2(R^n)$. Recall that

$$(2.20) \quad \begin{aligned} K_2(w_1) &= \{v \in V; v \leq \phi_2 \text{ a.e. and a.e. if } w_1 \geq \phi_1 \text{ then } v = \psi_2\}, \\ K_1(w_2) &= \{v \in V; v \leq \phi_1 \text{ a.e. and a.e. if } w_2 \geq \phi_2 \text{ then } v = \psi_1\}. \end{aligned}$$

In $L^2(R^n)$ we define a mapping T_2 as follows: for a given $w_1 \in L^2(R^n)$, $u_2 = T_2 w_1$ is the solution of the variational inequality

$$(2.21) \quad a(u_2, v - u_2) \geq (f_2, v - u_2) \quad \text{for any } v \in K_2(w_1), u_2 \in K_2(w_1).$$

The solution of (2.21) exists and is unique (cf. [21]).

Next we define a mapping T_1 : for a given $w_2 \in L^2(R^n)$, $u_1 = T_1 w_2$ is the solution of the variational inequality

$$(2.22) \quad a(u_1, v - u_1) \geq (f_1, v - u_1) \quad \text{for all } v \in K_1(w_2), u_1 \in K_1(w_2).$$

We define a mapping S_1 in $L^2(R^n)$ by $S_1 = T_1 T_2$, i.e.,

$$(2.23) \quad u_1 = S_1 w_1 = T_1 T_2 w_1.$$

We also define $S_2 = T_2 T_1$. It is clear that if u_1 is a fixed point for S_1 , then $(u_1, T_2 u_1)$ is a solution of the Q.V.I.; further, $T_2 u_1$ is then a fixed point for S_2 (and, conversely, if u_2 is a fixed point for S_2 , then $T_1 u_2$ is a fixed point for S_1 , $(T_1 u_2, u_2)$ is a solution of the Q.V.I., and $T_1 u_2$ is a fixed point for S_1).

The crucial fact to be proved below is that

$$(2.24) \quad \begin{aligned} S_1 \text{ (resp. } S_2) &\text{ is increasing in the sense} \\ &\text{that if } w_1 \leq w'_1 \text{ a.e. (resp. } w_2 \leq w'_2 \text{ a.e.)} \\ &\text{then } S_1 w_1 \leq S_1 w'_1 \text{ a.e. (resp. } S_2 w_2 \leq \\ &S_2 w'_2 \text{ a.e.)}. \end{aligned}$$

We shall first verify that

$$(2.25) \quad u_2 = T_2 w_1 \geq \psi_2 \quad \text{for all } w_1, \quad u_1 = T_1 w_2 \geq \psi_1 \quad \text{for all } w_2.$$

We note that since $\psi_1 \in K_1(w_2)$, $\max(\psi_1, u_1) \in K_1(w_2)$. Using $v = \max(\psi_1, u_1)$ as a test function in (2.22) and noting that $\max(\psi_1, u_1) = u_1 + (u_1 - \psi_1)^-$, we get

$$a(u_1, (u_1 - \psi_1)^-) \geq (f_1, (u_1 - \psi_1)^-),$$

or

$$a((u_1 - \psi_1)^-) - a(\psi_1, (u_1 - \psi_1)^-) + (f_1, (u_1 - \psi_1)^-) \leq 0,$$

which with (2.18) implies $a((u_1 - \psi_1)^-) = 0$; hence by (2.8), $u_1 \geq \psi_1$. A similar argument shows that $u_2 \geq \psi_2$.

Now let $w_1 \leq w'_1$, and write $u_2 = T_2 w_1$ and $u_1 = T_1 T_2 w_1 = S_1 w_1$ (similarly define u'_2 and u'_1 for w'_1). We shall first prove that

$$(2.26) \quad u_2 \geq u'_2.$$

Since $w_1 \leq w'_1$, we have $K_2(w'_1) \subset K_2(w_1)$. Therefore $u_2, u'_2 \in K_2(w_1)$, and also $\max(u_2, u'_2) \in K_2(w_1)$. Now from $u'_2 \in K_2(w'_1)$ it follows that if $w'_1 \geq \phi_1$,

$\min(u_2, u'_2) \leq u'_2 = \psi_2$. But from (2.25) we also have $\min(u_2, u'_2) \geq \psi_2$; hence $\min(u_2, u'_2) = \psi_2$ if $w'_1 \geq \phi_1$. Since $\min(u_2, u'_2) \leq u'_2 \leq \phi_2$, it follows that $\min(u_2, u'_2) \in K_2(w'_1)$. We can thus take $\max(u_2, u'_2)$ as a test function in (2.21) and $\min(u_2, u'_2)$ as a test function in (2.21)' (i.e. (2.21) for w'_1). Noting that

$$\max(u_2, u'_2) = u_2 + (u'_2 - u_2)^+, \quad \min(u_2, u'_2) = u'_2 - (u'_2 - u_2)^+,$$

we get

$$\begin{aligned} a(u_2, (u'_2 - u_2)^+) &\geq (f_2, (u'_2 - u_2)^+), \\ -a(u'_2, (u'_2 - u_2)^+) &\geq -(f_2, (u'_2 - u_2)^+) \end{aligned}$$

and, by addition, $a((u'_2 - u_2)^+) \leq 0$ which with (2.8) implies $(u'_2 - u_2)^+ = 0$; hence (2.26). Now since $u_2 \geq u'_2$, we have $K_1(u_2) \subset K_1(u'_2)$. Therefore u_1 and $u'_1 \in K_1(u'_2)$; hence $\max(u_1, u'_1) \in K_1(u'_2)$. Taking (2.25) into account, we prove by a similar argument as above that $\min(u_1, u'_1) \in K_1(u_2)$. We use $\max(u_1, u'_1)$ (resp. $\min(u_1, u'_1)$) as a test function in the variational inequality for u'_1 (resp. u_1) and, by addition, we obtain $u_1 \leq u'_1$, which proves (2.24). From (2.25) it follows that

$$(2.27) \quad S_1 \psi_1 \geq \psi_1,$$

i.e., ψ_1 is a lower solution of the equation $S_1 u_1 = u_1$ in the terminology of Tartar [23]. By construction of S_1 and T_1 ,

$$(2.28) \quad S_1 \phi_1 \leq \phi_1,$$

i.e., ϕ_1 is an upper solution. Since $\psi_1 \leq \phi_1$, we can use a general theorem of Tartar [23], which asserts that S_1 has a fixed point between ψ_1 and ϕ_1 . Moreover there exists a maximal and a minimal solution. In other words, there exist \underline{u}_1 and \bar{u}_1 such that

$$(2.29) \quad \begin{aligned} \psi_1 &\leq \underline{u}_1 \leq \bar{u}_1 \leq \phi_1, \quad S_1 \underline{u}_1 = \underline{u}_1, \quad S_1 \bar{u}_1 = \\ &\bar{u}_1, \quad \text{and if } u_1 = S_1 u_1, \quad \psi_1 \leq u_1 \leq \phi_1 \\ &\text{then } \underline{u}_1 \leq u_1 \leq \bar{u}_1. \end{aligned}$$

Defining $\bar{u}_2 = T_2 \bar{u}_1$ and $\underline{u}_2 = T_2 \underline{u}_1$, we see that $(\underline{u}_1, \bar{u}_2)$ and $(\bar{u}_1, \underline{u}_2)$ are two pairs of solutions for the Q.V.I.

Let now u_1, u_2 be a solution. By (2.25), $\psi_1 \leq u_1 \leq \phi_1$ (and $\psi_2 \leq u_2 \leq \phi_2$). We thus have $u_1 = S_1 u_1$ and, according to (2.29), $\underline{u}_1 \leq u_1 \leq \bar{u}_1$. Therefore $T_2 \bar{u}_1 \leq u_2 \leq T_2 \underline{u}_1$ which proves (2.19) and completes the proof of the theorem.

CHAPTER 3. PARABOLIC QUASI-VARIATIONAL INEQUALITIES IN ONE SPACE DIMENSION

3.1. The problem. In this chapter we shall solve the following problem: Find functions $u_1(x, t)$, $u_2(x, t)$ and curves $s_1(t)$, $s_2(t)$ with

$$s_1(0) = -1, \quad s_2(0) = 1, \quad s_1(t) < s_2(t) \quad (0 < t \leq T)$$

such that

$$(3.1) \quad \begin{aligned} u_{1t} - u_{1xx} &\leq \tilde{f}_1 \text{ and } u_1 = \tilde{\phi}_1 \text{ if } x < s_1(t), 0 < t < T, \\ u_{1t} - u_{1xx} &= \tilde{f}_1 \text{ and } u_1 < \tilde{\phi}_1 \text{ if } s_1(t) < x < s_2(t), 0 < t < T, \\ u_1 &= \tilde{\psi}_1 \text{ if } x > s_2(t), 0 < t < T, \\ u_1(x, 0) &= h_1(x) \text{ if } -1 < x < 1; \end{aligned}$$

$$(3.2) \quad \begin{aligned} u_{2t} - u_{2xx} &\leq \tilde{f}_2 \text{ and } u_2 = \tilde{\phi}_2 \text{ if } x > s_2(t), 0 < t < T, \\ u_{2t} - u_{2xx} &= \tilde{f}_2 \text{ and } u_2 < \tilde{\phi}_2 \text{ if } s_1(t) < x < s_2(t), 0 < t < T, \\ u_2 &= \tilde{\psi}_2 \text{ if } x < s_1(t), 0 < t < T, \\ u_2(x, 0) &= h_2(x) \text{ if } -1 < x < 1. \end{aligned}$$

This system is a parabolic quasi-variational inequality in one space dimension. It can, in fact, be easily reformulated in the more standard Q.V.I. terminology, using convex cones $K_1(u_2(\cdot, t))$, $K_2(u_1(\cdot, t))$.

Assume now that

$$(3.3) \quad \begin{aligned} h_i(x) &> 0 \text{ if } -1 < x < 1 \ (i = 1, 2), \\ h_1(x) &= 0 \text{ if } x < -1, \\ h_2(x) &= 0 \text{ if } x > 1, \\ \sigma(x, t) &\equiv \sqrt{2}, \quad g(x, t) \equiv 0, \\ f_i(x, t) &= \tilde{f}_i(x, T - t), \quad \phi_i(x, t) = \tilde{\phi}_i(x, T - t), \\ \psi_i(x, t) &= \tilde{\psi}_i(x, T - t), \\ \psi_i(x, t) &< \phi_i(x, t) \quad (i = 1, 2). \end{aligned}$$

Suppose there exists a sufficiently "strong" solution of (3.1), (3.2) in the sense that

$$u_{it}, u_{ix}, u_{ixx} \text{ are in } L^2 \text{ and } u_i \text{ is continuous in the region } s_1(t) \leq x \leq s_2(t), 0 \leq t \leq T.$$

From Theorem 1.1 we then deduce that there exists a Nash equilibrium point (τ_1^*, τ_2^*) with τ_i^* being the exit time from the set

$$C_i = \{(x, t); u_i(x, t) < \tilde{\phi}_i(x, t)\}.$$

In this chapter we shall prove the existence of a "weak" solution of (3.1), (3.2). It will be shown, in fact, that there exists a "maximal" solution (\bar{u}_1, \bar{u}_2) and a "minimal" solution $(\underline{u}_1, \underline{u}_2)$, i.e., for any other solution (u_1, u_2) , $\underline{u}_1 \leq u_1 \leq \bar{u}_1$, $\underline{u}_2 \geq u_2 \geq \bar{u}_2$, where the functions are defined.

Set

$$\gamma_i = \tilde{\phi}_i - \tilde{\psi}_i, \quad k_i(x) = \tilde{\phi}_i(x, 0) - h_i(x).$$

Throughout this chapter it is always assumed that

$$(3.4) \quad \begin{aligned} &\gamma_{it}, \gamma_{ix}, \gamma_{ixx} \text{ are continuous and bounded for } (x, t) \in R^1 \times [0, T], \\ &k_{ix} \text{ are continuous for } x \in R^1, \\ &k_{1xx} \text{ is uniformly continuous for } -1 < x < \infty, \\ &k_{2xx} \text{ is uniformly continuous for } -\infty < x < 1, \\ &\gamma_i(x, t) > 0 \text{ for all } (x, t), \\ &k_i(x) > 0 \text{ if } -1 < x < 1, \\ &k_1(x) = 0 \text{ if } x < -1, \quad k_1(x) = \gamma_1(x, 0) \text{ if } x > 1, \\ &k_2(x) = \gamma_2(x, 0) \text{ if } x < -1, \quad k_2(x) = 0 \text{ if } x > 1, \end{aligned}$$

and that

$$(3.5) \quad \tilde{\phi}_{it}, \tilde{\phi}_{ix}, \tilde{\phi}_{ixx} \text{ are continuous,} \quad \tilde{\phi}_{it} - \tilde{\phi}_{ixx} - \tilde{f}_i \equiv -1.$$

The last condition implies, of course, that $u_{it} - u_{ixx} \leq \tilde{f}_i$ a.e. on the set $u_i = \tilde{\phi}_i$. Notice also that if $u_1 = \tilde{\psi}_1$ on the curve $x = s_2(t)$, then we can always define u_1 for $x > s_2(t)$ by $u_1 = \tilde{\psi}_1$. A similar remark applies to u_2 . Hence, setting

$$w_1 = \tilde{\phi}_1 - u_1, \quad w_2 = \tilde{\phi}_2 - u_2,$$

the system (3.1), (3.2) reduces to

$$(3.6) \quad \begin{aligned} w_{1t} - w_{1xx} &= -1 \text{ and } w_1 > 0 && \text{if } s_1(t) < x < s_2(t), 0 < t < T, \\ w_1 &= 0 && \text{if } x < s_1(t), 0 < t < T, \\ w_1 &= \gamma_1 && \text{if } x = s_2(t), 0 < t < T, \\ w_1(x, 0) &= k_1(x) && \text{if } -1 < x < 1; \end{aligned}$$

$$(3.7) \quad \begin{aligned} w_{2t} - w_{2xx} &= -1 \text{ and } w_2 > 0 && \text{if } s_1(t) < x < s_2(t), 0 < t < T, \\ w_2 &= 0 && \text{if } x > s_2(t), 0 < t < T, \\ w_2 &= \gamma_2 && \text{if } x = s_1(t), 0 < t < T, \\ w_2(x, 0) &= k_2(x) && \text{if } -1 < x < 1. \end{aligned}$$

We shall prove later on the existence of a “weak” solution of (3.6), (3.7) with $s_1(t)$, $s_2(t)$ (the *free boundary curves*) having the following properties:

$$(3.8) \quad \begin{aligned} &s_1(t) \text{ is continuous and strictly monotone decreasing,} \\ &s_2(t) \text{ is continuous and strictly monotone increasing.} \end{aligned}$$

We outline the idea of the proof: Given a monotone curve $x = s_2(t)$, we solve the variational inequality (3.6) and show that its free boundary is a curve $x = \sigma_1(t)$. Next we solve the variational inequality (3.7) with $s_1(t)$ replaced by $\sigma_1(t)$, and show that its free boundary is a curve $x = \bar{s}_2(t)$. We have thus constructed a mapping W , $\bar{s}_2 = Ws_2$. It will be shown (using

Tartar's fixed point theorem [23]) that W has fixed points; these points represent solutions of the Q.V.I. (3.6), (3.7). Some technical difficulties arise due to the fact that the monotone curves s_2 on which W is defined may not be continuous.

3.2. Auxiliary variational inequality. Let $s(t)$ be a monotone increasing function with Hölder continuous first derivative $\dot{s}(t)$, for $0 \leq t \leq T$, such that $s(0) = 1$. Set

$$G = \{(x, t); -\infty < x < s(t), 0 < t < T\}.$$

Consider the variational inequality: Find w such that

$$\begin{aligned} w &\geq 0 \quad \text{if } x < s(t), 0 < t < T, \\ w_t, w_x, w_{xx} &\text{ are in } L^2(G), \\ \int_{-\infty}^{s(t)} [w_t(x, t) - w_{xx}(x, t)] [v(x) - w(x, t)] dx \\ (3.9) \quad &\geq - \int_{-\infty}^{s(t)} [v(x) - w(x, t)] dx \\ &\text{for a.a. } t \in (0, T), \text{ for any } v \in L^\infty(R^1), v \geq 0 \text{ a.e.,} \\ w &= \gamma_1 \quad \text{if } x = s(t), 0 \leq t \leq T, \\ w(x, 0) &= k_1(x) \quad \text{if } x < 1. \end{aligned}$$

LEMMA 3.1. *There exists a unique solution w of (3.9) with compact support; further, w_t, w_x, w_{xx} belong to $L^p(G)$ for any $1 < p < \infty$.*

FIRST PROOF. Let $\beta_\varepsilon(t)$ be a family of C^∞ functions of t ($-\infty < t < \infty$, $0 < \varepsilon < 1$) such that

$$\begin{aligned} \beta_\varepsilon(t) &\leq 0, \quad \beta_\varepsilon(0) = -1, \quad \beta'_\varepsilon(t) \geq 0, \\ \beta_\varepsilon(t) &\rightarrow 0 \quad \text{if } t > 0, \varepsilon \rightarrow 0, \quad \beta_\varepsilon(t) \rightarrow -\infty \quad \text{if } t < 0, \varepsilon \rightarrow 0. \end{aligned}$$

Set $G_R = G \cap \{x; x > -R\}$ for any $R > 1$, and consider the parabolic problem

$$\begin{aligned} (3.10) \quad w_t - w_{xx} + \beta_\varepsilon(w) &= -1 \quad \text{in } G_R, \\ w(x, 0) &= k_1(x) \quad \text{if } -R < x < 1, \\ w(s(t), t) &= \gamma_1(s(t), t) \quad \text{if } 0 < t < T, \\ w(-R, t) &= 0 \quad \text{if } 0 < t < T. \end{aligned}$$

Denote the solution by $w = w_{\varepsilon, R}$. One can show (cf. [9], [17], [19]) that

$$(3.11) \quad -1 \leq \beta_\varepsilon(w) \leq 0.$$

Hence

$$(3.12) \quad |w_t - w_{xx}| \leq 1.$$

By the L^p estimates for parabolic equations (see, for instance, [22]) we then deduce that

$$(3.13) \quad \int \int_{G_R} [|w_t|^p + |w_x|^p + |w_{xx}|^p] dx dt \leq C \quad (1 < p < \infty)$$

where C is a constant independent of ε . Actually, in the L^p estimates one usually assumes that the domain G_R is cylindrical, i.e., $s(t) \equiv \text{const}$. Therefore in order to obtain (3.13) we first perform a transformation

$$(3.14) \quad y = \frac{x}{s(t)}, \quad \tilde{w}(y, t) = w(x, t)$$

and note that

$$(3.15) \quad w_t - w_{xx} = \tilde{w}_t - \frac{y \dot{s}(t)}{s(t)} \tilde{w}_y - \frac{1}{s^2(t)} \tilde{w}_{yy}.$$

We then apply the interior L^p estimates to $\tilde{w}(y, t)$ in some region $-\tilde{R} < y < 1$, $0 < t < T$. This yields the L^p estimates (3.13) in the region $G' = \{0 < x < s(t), 0 < t < T\}$, provided \tilde{R} was suitably chosen, depending on the function s . (We assume that $R > \tilde{R}$.) We also have, by interior L^p estimates, the estimate (3.13) in $G_R - G'$. Combining these estimates, (3.13) follows.

Recall that in (3.13), w stands for $w_{\varepsilon, R}$. Taking $\varepsilon \downarrow 0$ through such a sequence that $w_{\varepsilon, R}$ is weakly convergent in the norm $L^p(G_R)$ together with the derivatives $\partial/\partial t$, $\partial/\partial x$, $\partial^2/\partial x^2$, we find that the weak limit $w_R \equiv \lim w_{\varepsilon, R}$ in the unique solution of the variational inequality

$$(3.16) \quad \begin{aligned} & w \geq 0 \quad \text{in } G_R, \\ & w_t, w_x, w_{xx} \text{ are in } L^p(G_R), \quad 1 < p < \infty, \\ & \int_{-R}^{s(t)} (w_t - w_{xx})(v - w) dx \geq - \int_{-R}^{s(t)} (v - w) dx \text{ a.e. in } t \in [0, T], \\ & \text{for any } v \in L^\infty(R^1), v \geq 0 \text{ a.e.,} \\ & w = \gamma_1 \quad \text{if } x = s(t), 0 < t < T, \\ & w(x, 0) = k_1(x) \quad \text{if } -R < x < 1, \\ & w(-R, t) = 0 \quad \text{if } 0 < t < T. \end{aligned}$$

By [10], there exists a number R_0 sufficiently large such that if $R > R_0$ then

$$w \equiv w_R = 0 \quad \text{if } -R < x < -R_0, 0 < t < T.$$

Hence, by uniqueness,

$$w_{R'}(x, t) = w_R(x, t) \quad \text{if } -R < x < s(t), 0 < t < T, R' > R.$$

It follows that $w = \lim w_R$ is the asserted unique solution of (3.9).

SECOND PROOF. Performing the transformation (3.14), and using (3.15), the

variational inequality (3.9) reduces to the variational inequality:

$$\begin{aligned}
 & \tilde{w} \geq 0 \quad \text{in } \tilde{G}, \\
 & \tilde{w}_t, \tilde{w}_y, \tilde{w}_{yy} \text{ are in } L^2(\tilde{G}), \\
 & \int_{-\infty}^1 \left[\tilde{w}_t - \frac{1}{s^2(t)} \tilde{w}_{yy} - \frac{y\dot{s}(t)}{s(t)} \tilde{w}_y \right] (v - \tilde{w}) dy \geq - \int_{-\infty}^1 (v - \tilde{w}) dy \\
 & \text{for a.a. } t \in (0, T), \text{ for any } v \in L^\infty(R^1), v \geq 0 \text{ a.e.,} \\
 & \tilde{w}(1, t) = \gamma_1(s(t), t) \quad \text{if } 0 < t < T, \\
 & \tilde{w}(y, 0) = k_1(y) \quad \text{if } y < 1,
 \end{aligned}
 \tag{3.17}$$

where $\tilde{G} = \{(y, t); -\infty < y < 1, 0 < t < T\}$.

Let $\tilde{G}_R = \tilde{G} \cap \{y, y > -R\}$ and consider the parabolic problem:

$$\begin{aligned}
 & \tilde{w}_t - \frac{1}{s^2(t)} \tilde{w}_{yy} - \frac{y\dot{s}(t)}{s(t)} \tilde{w}_y + \beta_\varepsilon(w) = -1 \quad \text{in } \tilde{G}_R, \\
 & \tilde{w}(1, t) = \gamma_1(s(t), t) \quad \text{if } 0 < t < T, \\
 & \tilde{w}(y, 0) = k_1(y) \quad \text{if } -R < y < 1, \\
 & \tilde{w}(-R, t) = 0 \quad \text{if } 0 < t < T.
 \end{aligned}
 \tag{3.18}$$

Denote its solution by $\tilde{w} = \tilde{w}_{\varepsilon, R}$. Since the coefficients of the parabolic operator in (3.18) are Hölder continuous, the L^p parabolic estimates (cf. (3.13)) are valid. We can now proceed as in the first proof and show that when $\varepsilon \downarrow 0$ in a suitable way, the solutions $\tilde{w}_{\varepsilon, R}$ converge weakly in $L^p(\tilde{G}_R)$ to a function \tilde{w}_R (together with the derivatives $\partial/\partial t$, $\partial/\partial x$, $\partial^2/\partial x^2$), and \tilde{w}_R is a solution of the variational inequality obtained from (3.17) by replacing \tilde{G} by \tilde{G}_R , $\int_{-\infty}^1$ by \int_{-R}^1 and by replacing the last condition in (3.17) by the last two conditions of (3.18).

The techniques of [10] show that the support of \tilde{w}_R remains bounded as R increases to infinity. Hence $\tilde{w}_{R'} = \tilde{w}_R$ in \tilde{G}_R if $R' > R$, R sufficiently large. It follows that $\tilde{w} = \lim \tilde{w}_R$ is the desired solution.

REMARK 1. From the first proof, $w = \lim_R \lim_\varepsilon w_{\varepsilon, R}$. From the second proof, $w(x, t) = \tilde{w}(y, t)$ and $\tilde{w} = \lim_R \lim_\varepsilon \tilde{w}_{\varepsilon, R}$. Both relations will be needed in §3.3.

REMARK 2. From (3.12) and the maximum principle applied to $w = w_{\varepsilon, R}$ we deduce that

$$|w_{\varepsilon, R}| \leq A \tag{3.19}$$

where A is a constant independent of ε , R and of $s(t)$. Taking $\varepsilon \downarrow 0$, $R \uparrow \infty$ we deduce that

$$0 \leq w(x, t) \leq A \tag{3.20}$$

with the same A , which is independent of the curve $s(t)$.

REMARK 3. We can write, for $w = w_{\varepsilon, R}$,

$$(3.21) \quad (w - \gamma_1)_t - (w - \gamma_1)_{xx} = -1 - \beta_\varepsilon(w) - \gamma_{1t} + \gamma_{1xx} \equiv B$$

where, by (3.11), B is bounded by a constant A_0 independent of ε , R and the curve $s(t)$. Multiplying (3.21) by $w - \gamma_1$ and integrating over G_R , we obtain, after using (3.19) and taking $\varepsilon \downarrow 0$,

$$- \int \int_{G_R} (w - \gamma_1)(w - \gamma_1)_{xx} dx dt \leq A_1$$

where $w = w_R$ and A_1 is a constant independent of R and of the curve $s(t)$. If R is sufficiently large then $w = w_x = 0$ on $x = -R$. Hence, integrating by parts we find, after letting $R \rightarrow \infty$, that

$$(3.22) \quad \int \int_G w_x^2(x, t) dx dt \leq \tilde{A}$$

where \tilde{A} is a constant independent of the curve $s(t)$.

3.3. Further properties of w . We shall need the conditions:

$$(3.23) \quad \gamma_{1x} \geq 0, \gamma_{1t} \geq 0 \quad \text{if } x \geq 1, \quad k_{1x} \geq 0 \quad \text{if } -1 \leq x \leq 1.$$

LEMMA 3.2. *The function $w_x(x, t)$ is continuously differentiable in \bar{G} and $w_x \geq 0$.*

PROOF. The function $\tilde{w} = \tilde{w}_{\varepsilon, R}(y, t)$, occurring in the second proof of Lemma 3.1, satisfies

$$(3.24) \quad \tilde{w}_t - \frac{y\dot{s}(t)}{s(t)} \tilde{w}_y - \frac{1}{s^2(t)} \tilde{w}_{yy} + \beta_\varepsilon(\tilde{w}) = -1.$$

Differentiating this equation with respect to t , multiplying by $|\tilde{w}_t|^{p-2}\tilde{w}_t$ ($2 \leq p < \infty$) and integrating over \tilde{G}_R , we find (cf. [15]) that

$$(3.25) \quad \int_{-R}^1 \left| \frac{\partial}{\partial t} \tilde{w}_{\varepsilon, R} \right|^p dy < C \quad (0 < t < T)$$

where C is a constant independent of ε , R . We can now apply the elliptic L^p estimates to the elliptic operator $y\dot{s}\tilde{w}_y/s + \tilde{w}_{yy}/s^2$ and conclude that

$$(3.26) \quad \int_{-R}^1 \left[\left| \frac{\partial}{\partial y} \tilde{w}_{\varepsilon, R} \right|^p + \left| \frac{\partial^2}{\partial y^2} \tilde{w}_{\varepsilon, R} \right|^p \right] dy < C \quad (0 < t < T)$$

where C is a constant independent of ε , R .

Taking $\varepsilon \downarrow 0$, $R \uparrow \infty$, we deduce from (3.25), (3.26), after performing the transformation (3.14), that

$$(3.27) \quad \int_{R^1} [|w_t(x, t)|^p + |w_x(x, t)|^p + |w_{xx}(x, t)|^p] dx < C \quad (0 < t < T).$$

Since p is arbitrary, the Sobolev inequality implies that w is continuous in \bar{G} , and that the function $x \rightarrow w_x(x, t)$ is uniformly Hölder continuous in x , with exponent and coefficient that are independent of t . By the proof of Corollary 2.7 of [19] we then deduce that w_x is continuous in \bar{G} .

Set

$$\Omega = \{(x, t) \in G; w(x, t) > 0\}, \quad \Omega_\tau = \Omega \cap \{t < \tau\}.$$

The conditions on γ_1 in (3.23) together with the fact that $s(t)$ is increasing in t imply that

$$(3.28) \quad w(s(t), t) = \gamma_1(s(t), t) \text{ is increasing in } t.$$

Since $k_{1x} > 0$, $k_1(1) = \gamma_1(1, 0)$, we also have that

$$(3.29) \quad w(s(t), t) \geq k_1(x) \quad \text{if } x < 1.$$

We now apply the maximum principle to the function w in Ω_τ . Since $w_t - w_{xx} = -1 < 0$ in Ω and since $w = 0$ in that part of the parabolic boundary of Ω that lies in G , we deduce (using (3.28), (3.29)) that the maximum of w in $\bar{\Omega}_\tau$ is attained at $(s(\tau), \tau)$. Consequently,

$$w_x(s(\tau), \tau) \geq 0 \quad \text{if } 0 < \tau < t.$$

Next,

$$w_x(x, 0) = k_{1x} \geq 0 \quad \text{if } -1 < x < 1.$$

Finally, $w_x = 0$ on the free boundary. Since w_x is continuous in $\bar{\Omega}$ and

$$(w_x)_t - (w_x)_{xx} = 0 \quad \text{in } \Omega,$$

the maximum principle applied to w_x in Ω yields $w_x > 0$ in Ω . Since $w_x = 0$ in $G \setminus \Omega$, the proof of the lemma is complete.

We shall need the following additional assumptions:

$$(3.30) \quad k_{1xx}(x) - 1 \geq 0 \quad \text{if } -1 < x < 1;$$

there is a function $\tilde{\gamma}_1(x, t)$ with continuous derivatives

$\tilde{\gamma}_{1x}, \tilde{\gamma}_{1t}, \tilde{\gamma}_{1xx}$ such that

$$(3.31) \quad \begin{aligned} \tilde{\gamma}_1(x, t) &= \gamma_1(x, t) \quad \text{if } x > 1, \\ \tilde{\gamma}_1(x, 0) &\leq k_1(x) \quad \text{if } x < 1, \\ \lim_{x \rightarrow -\infty} \tilde{\gamma}_1(x, t) &\leq 0 \quad \text{uniformly in } t, 0 \leq t \leq T, \\ \tilde{\gamma}_{1xx} - \tilde{\gamma}_{1t} &\geq 1 \quad \text{if } x \in R^1, 0 < t < T. \end{aligned}$$

For example, we can take $\tilde{\gamma}_1(x, t) = k_1(x) + (x - 1)t$ if $\gamma_1(x, t) = k_1(x) + (x - 1)t$ when $x > 1$, provided $k_{1xx}(x) \geq x$ whenever $x > 1$.

LEMMA 3.3. *Under the additional assumptions (3.30), (3.31),*

$$(3.32) \quad w(x, t) > \gamma_1(x, t) \quad \text{if } 1 < x < s(t), 0 < t < T,$$

$$(3.33) \quad w_x - \gamma_{1x} \leq 0 \quad \text{if } x = s(t), 0 < t < T.$$

PROOF. Consider the function

$$u(x, t) = w_{\varepsilon, R}(x, t) - \tilde{\gamma}_1(x, t)$$

in the region G_R . From the last relation in (3.31) it follows that

$$u_t - u_{xx} \geq -\beta_\varepsilon(w_{\varepsilon, R}) \geq 0 \quad \text{in } G_R.$$

Also, for any $\delta > 0$,

$$u = 0 \quad \text{if } x = s(t), 0 < t < T,$$

$$u(x, 0) \geq 0 \quad \text{if } -R < x < 1,$$

$$u(-R, t) \geq -\delta \quad \text{if } 0 < t < T,$$

provided R is sufficiently large (depending on δ , but not on ε). By the maximum principle, $u \geq -\delta$ in G_R . Taking $\varepsilon \rightarrow 0$, $R \rightarrow \infty$ we get

$$w - \tilde{\gamma}_1 \geq -\delta \quad \text{in } G.$$

Since δ is arbitrary, $w \geq \tilde{\gamma}_1$ in G . By the strong maximum principle we actually then have $w > \tilde{\gamma}_1$ in Ω . This gives (3.32). The inequality (3.33) is an immediate consequence of (3.32).

LEMMA 3.4. *Under the additional assumptions (3.30), (3.31), $w_t \geq 0$ a.e.*

PROOF. Consider first the case where γ_1, k_1, s are in C^3 and

$$(3.34) \quad \gamma_{1x}(1, 0)s'(0) + \gamma_{1t}(1, 0) - k_{1xx}(1) = 0.$$

This is a consistency condition for the equation

$$(3.35) \quad \frac{\partial}{\partial t} w_{\varepsilon, R} - \frac{\partial^2}{\partial x^2} w_{\varepsilon, R} + \beta_\varepsilon(w_{\varepsilon, R}) = -1$$

at the point $(1, 0)$, since $\beta_\varepsilon(0) = -1$. At $(-R, 0)$ the consistency condition for (3.35) is also satisfied. We can now apply the Schauder estimates [13] to deduce that $\partial w_{\varepsilon, R}/\partial x$, $\partial^2 w_{\varepsilon, R}/\partial x^2$, $\partial w_{\varepsilon, R}/\partial t$ are continuous in \bar{G}_R .

Differentiating the relation

$$(3.36) \quad w_{\varepsilon, R}(s(t), t) - \gamma_1(s(t), t) = 0,$$

we get

$$\dot{s}(t) \frac{\partial}{\partial x} (w_{\varepsilon, R} - \gamma_1)(s(t), t) + \frac{\partial}{\partial t} w_{\varepsilon, R}(s(t), t) = \gamma_{1t}(s(t), t).$$

Since $w_{\varepsilon, R} \rightarrow w$ (weakly in $L^p(G_R)$) together with the first derivatives and, therefore, uniformly in \bar{G}_R , we can apply the Schauder boundary estimates (the consistency condition at $(1, 0)$ is used here) to deduce that

$$\frac{\partial}{\partial x} w_{\varepsilon, R}(s(t), t) \rightarrow \frac{\partial}{\partial x} w(s(t), t) \quad \text{as } \varepsilon \rightarrow 0, R \rightarrow 0,$$

uniformly in t , $0 \leq t \leq T$.

Using Lemma 3.3 we find that

$$\frac{\partial}{\partial x} (w_{\varepsilon, R} - \gamma_1)(s(t), t) < \frac{\delta}{A}$$

if $0 < \varepsilon < \varepsilon_0$, $R > R_0$ for some ε_0 , R_0 depending on δ/A ; here δ is an arbitrary positive number and $A = \max_{0 \leq t \leq T} \dot{s}(t)$.

Using this inequality in (3.36), and recalling that $\gamma_{1t} \geq 0$, we get

$$(3.37) \quad \frac{\partial}{\partial t} w_{\varepsilon, R}(s(t), t) > -\delta \quad (0 < \varepsilon < \varepsilon_0, R > R_0).$$

Setting $\xi = \partial w_{\varepsilon, R} / \partial t$ and differentiating (3.35) with respect to t , we get

$$(3.38) \quad \xi_t - \xi_{xx} + \beta'(w_{\varepsilon, R})\xi = 0 \quad \text{in } G_R.$$

Since $\beta_\varepsilon(0) = -1$ and since (3.30) holds,

$$\xi(x, 0) = -1 - \beta_\varepsilon(k_1(x)) + k_{1xx}(x) \geq 0 \quad \text{if } -R < x < 1.$$

On $x = -R$, $\xi = 0$. On $x = s(t)$, $\xi > -\delta$, by (3.37). Since ξ is continuous in \bar{G}_R , we can apply the maximum principle to conclude that $\xi \geq -\delta$ in \bar{G}_R . Taking $\varepsilon \rightarrow 0$, $R \rightarrow \infty$ the assertion of the lemma follows.

We have assumed in the above proof that γ_1 , k_1 , s are in C^3 and (3.34) holds. In the general case, we approximate γ_1 , k_1 , s by C^3 functions γ_1^m , k_1^m , s^m satisfying all the assumptions of Lemma 3.4 and (3.34). For the corresponding variational inequality, the solution w^m satisfies $\partial w^m / \partial t \geq 0$. By uniqueness, $w^m \rightarrow w$ (uniformly, say) as $m \rightarrow \infty$, and the assertion of the lemma readily follows.

We now introduce the free boundary curve $x = \sigma(t)$ of the variational inequality (3.9):

$$\sigma(t) = \inf\{x; x < s(t), w(x, t) > 0\}, \quad 0 \leq t \leq T.$$

LEMMA 3.5. *Let $s(t)$ be a monotone increasing function with Hölder continuous derivative, for $0 \leq t \leq T$, such that $s(0) = 1$. Let γ_1 , k_1 satisfy the conditions in (3.4), (3.23), (3.30), (3.31). Then there exists a unique solution w with compact support of (3.9); w and w_x are continuous in \bar{G} , $w_x \geq 0$, $w_t \geq 0$ a.e. in G . Furthermore, the function $\sigma(t)$ is continuous and monotone decreasing in t , $0 \leq t \leq T$.*

PROOF. All the assertions of the lemma, except those regarding $\sigma(t)$, follow from Lemmas 3.1, 3.2, 3.4. Since $w_x \geq 0$, $w_t \geq 0$ a.e. in G , the strong maximum principle implies that $w_x > 0$, $w_t > 0$ in Ω . It follows that $(x, t) \in \Omega$ if and only if $\sigma(t) < x < s(t)$ and $\sigma(t)$ is monotone decreasing.

Clearly $\sigma(t)$ is upper semicontinuous function. Hence $\sigma(t)$ is right continuous. It remains to prove that $\sigma(t)$ is left continuous. If this is not the case then there is a point $t_0 \in [0, T)$ such that $\sigma(t_0 + 0) < \sigma(t_0)$. Introduce the line segment $I = \{(x, t_0); \sigma(t_0 + 0) < x < \sigma(t_0)\}$. Then I is a part of the boundary of Ω . Since $D = \{(x, t); \sigma(t_0 + 0) < x < \sigma(t_0), t_0 < t < T\}$ is contained in Ω , $w_t - w_{xx} = -1$ in D , $w = 0$ on I ; it follows that w is smooth in $D \cup I$. But then $w_t = -1 + w_{xx} = -1 < 0$ on I . It follows that $w(x, t) < 0$ at the points (x, t) of Ω such that $\sigma(t_0 + 0) < x < \sigma(t_0)$ and such that $t > t_0$, $t - t_0$ is sufficiently small. This is impossible since $w > 0$ in Ω .

3.4. The case where $s(t)$ is not smooth.

LEMMA 3.6. *Let the conditions of Lemma 3.5 hold and let $s'(t)$ be a monotone increasing function with Hölder continuous derivative for $0 \leq t \leq T$ such that $s'(0) = 1$. Denote by $w'(x, t)$ the solution of the variational inequality (3.9) with $s(t)$ replaced by $s'(t)$, and denote by $x = \sigma'(t)$ the corresponding free boundary. If $s'(t) \geq s(t)$ for $0 \leq t \leq T$, then*

$$(3.39) \quad w'(x, t) \geq w(x, t) \quad \text{for all } 0 \leq x \leq s(t), 0 \leq t \leq T,$$

$$(3.40) \quad \sigma'(t) \leq \sigma(t) \quad \text{for all } 0 \leq t \leq T.$$

PROOF. We compare the solution $w_{\varepsilon, R}$ of (3.10) with the corresponding solution $w'_{\varepsilon, R}$ when s is replaced by s' . On $x = -R$ and on $t = 0$, $w'_{\varepsilon, R} - w_{\varepsilon, R} = 0$. Next, by the proof of Lemma 3.3, for any $\delta > 0$, $w'_{\varepsilon, R} > \gamma_1 - \delta$ on $x = s(t)$ provided R is sufficiently large, depending only on δ . Since $w_{\varepsilon, R} = \gamma_1$ on $x = s(t)$, we have $w'_{\varepsilon, R} - w_{\varepsilon, R} > -\delta$ on $x = s(t)$. Applying the maximum principle to $w'_{\varepsilon, R} - w_{\varepsilon, R}$, we find that $w'_{\varepsilon, R} - w_{\varepsilon, R} > -\delta$ in G_R . Taking $\varepsilon \rightarrow 0$, $R \rightarrow \infty$ we conclude that $w' - w \geq -\delta$ in G . Since δ is arbitrary the assertion (3.39) follows. Clearly, (3.40) is a consequence of (3.39).

We shall now consider the variational inequality (3.9) in case $s(t)$ is any monotone increasing function with $s(0) = 1$. We shall construct a "generalized" solution as a limit of solutions w_n of variational inequalities corresponding to smooth curves $s_n(t)$. In order to define $s_n(t)$, we define $s(t) = 1$ if $t < 0$,

$$\rho(t) = \begin{cases} c \exp[1/(|t-1|-1)] & \text{if } |t-1| < 1, \\ 0 & \text{if } |t-1| \geq 1, \end{cases}$$

$$\rho_n(t) = n\rho(nt) \quad \text{if } n = 1, 2, \dots,$$

where c is a positive constant such that $\int_{R^1} \rho(t) dt = 1$.

Let

$$(3.41) \quad s_n(t) = \int_{R^1} \rho_n(t-\tau)s(\tau) d\tau = \int_{t-2/n}^t \rho_n(t-\tau)s(\tau) d\tau.$$

We can write

$$(3.42) \quad s_n(t) = \int_{R^1} \rho_n(\tau) s(t - \tau) d\tau = \int_0^2 \rho(\tau) s(t - \tau/n) d\tau.$$

From (3.41) we see that $s_n(t)$ is a C^∞ function, and from (3.42) we see that $s_n(t)$ is a monotone increasing function in t , $s_n(0) = 1$, and

$$(3.43) \quad s_n(t) \leq s_{n+1}(t).$$

Denote by w_n the solution of the variational inequality (3.9) corresponding to $s = s_n$ and denote by $x = \sigma_n(t)$ the corresponding free boundary curve. By Lemma 3.6,

$$(3.44) \quad w_n \leq w_{n+1} \quad \text{if } -\infty < x < s_n(t), 0 \leq t \leq T,$$

$$(3.45) \quad \sigma_{n+1}(t) \leq \sigma_n(t) \quad \text{if } 0 \leq t \leq T.$$

Set

$$(3.46) \quad w(x, t) = \lim_{n \rightarrow \infty} w_n(x, t) \quad \text{if } -\infty < x \leq s(t), 0 \leq t \leq T,$$

$$(3.47) \quad \sigma(t) = \lim_{n \rightarrow \infty} \sigma_n(t) \quad \text{if } 0 \leq t \leq T.$$

Let $s^*(t)$ be any C^∞ function such that $s^*(t) \leq s(t) - \delta$ if $0 \leq t \leq T$, for some $\delta > 0$, and such that $\tilde{\gamma}_1(x, t) > 0$ in a \bar{G} -neighborhood of

$$\Gamma^* = \{(s^*(t), t); 0 \leq t \leq T\}.$$

By Remark 2 at the end of §3.2, $|w_n| \leq A$ in some \bar{G} -neighborhood N of Γ^* , where A and N are both independent of n . By the proof of Lemma 3.3, $w_n \geq \tilde{\gamma}_1 > 0$ in N and, consequently, $w_{nt} - w_{nxx} = -1$ in N . By standard results on parabolic equations we then deduce (if k_{1xx} is Hölder continuous) that w_{nx} , w_{nt} , w_{nxx} are continuous and uniformly bounded in some smaller \bar{G} -neighborhood of Γ^* .

We can now consider w_n as a solution of the variational inequality (3.9) with $s(t)$ replaced by $s^*(t)$. From the proof of Lemma 3.1 we get, for any $1 < p < \infty$,

$$(3.48) \quad \int_{-\infty}^{s^*(t)} [|w_{nx}|^p + |w_{nt}|^p + |w_{nxx}|^p] dx \leq C \quad \text{if } 0 < t < T,$$

where C is a constant independent of n ; by approximation, this inequality holds also when k_{1xx} is not assumed to be Hölder continuous.

We can now use the Sobolev inequality to deduce that

$$(3.49) \quad w_n \text{ is uniformly Hölder continuous in } (x, t),$$

uniformly with respect to n ,

$$(3.50) \quad w_{nx} \text{ is uniformly Hölder continuous in } x,$$

uniformly with respect to t, n .

In view of (3.46) we then have

$$(3.51) \quad w_n \rightarrow w \text{ uniformly in } \bar{G}^*,$$

$$(3.52) \quad w_x \text{ is Hölder continuous in } x, \text{ uniformly in } t, \text{ where} \\ G^* = \{(x, t); -\infty < x < s^*(t), 0 < t < T\}.$$

Notice also that

$$(3.53) \quad \int_{-\infty}^{s^*(t)} [|w_x|^p + |w_t|^p + |w_{xx}|^p] dx \leq C \quad \text{if } 0 < t < T.$$

From (3.52) and the proof of Corollary 2.7 in [19] we also deduce that

$$(3.54) \quad w_x \text{ is continuous in } \bar{G}^*.$$

DEFINITION. We shall call the function $w(x, t)$ a *generalized solution* of the variational inequality (3.9). The curve $x = \sigma(t)$ will be called the *free boundary*.

Notice that if w were continuous up to the curve $x = s(t)$ and if w_x, w_t, w_{xx} were in L^2 in a G -neighborhood of this curve, then w would be a solution of the variational inequality (3.9) in the usual sense.

LEMMA 3.7. *Let $s(t)$ be a monotone increasing function for $0 \leq t \leq T$, with $s(0) = 1$, and let γ_1, k_1 satisfy the conditions in (3.4), (3.23), (3.30), (3.31). Then*

$$(3.55) \quad \sigma(t) = \inf\{x; w(x, t) > 0\},$$

$w(x, t) = 0$ if $x < \sigma(t)$, and the function $\sigma(t)$ is monotone decreasing and continuous for $0 \leq t \leq T$.

PROOF. Let

$$\Omega = \{(x, t) \in G; w(x, t) > 0\},$$

$$\Omega_n = \{(x, t) \in G; w_n(x, t) > 0\},$$

$$\Omega_0 = \{(x, t) \in G; x > \sigma(t)\}.$$

In view of (3.44), (3.46), $\Omega_n \subset \Omega$. Hence

$$(3.56) \quad \Omega_0 = \bigcup_{n=1}^{\infty} \Omega_n \subset \Omega.$$

On the other hand, $w_n(\sigma_n(t), t) = 0$. Taking $n \rightarrow \infty$ we get $w(\sigma(t), t) = 0$. Observing that $w_x \geq 0$ a.e. (since $w_{nx} \geq 0$ a.e.) we deduce that $w(x, t) = 0$ if $x < \sigma(t)$. This, together with (3.56), completes the proof of (3.55).

It is clear that $\sigma(t)$ is monotone decreasing in t . The proof that $\sigma(t)$ is continuous is the same as in the proof of Lemma 3.5.

REMARK. On $x = s(t)$ the generalized solution w may not be continuous. From Remark 3 at the end of §3.2, when applied to w_n with $n \rightarrow \infty$, we see that

$$\int_0^T \int_{-\infty}^{s(t)} w_x^2 dx dt < \infty.$$

We conclude this section with a comparison lemma.

LEMMA 3.8. *Let the condition of Lemma 3.7 hold and let $s'(t)$ be a monotone increasing function for $0 \leq t \leq T$, with $s'(0) = 1$. Denote by $w'(x, t)$ the generalized solution of (3.9) corresponding to s' , and denote its free boundary curve by $x = \sigma'(t)$. If $s'(t) \geq s(t)$ for $0 \leq t \leq T$, then*

$$\begin{aligned} w'(x, t) &\geq w(x, t) \quad \text{if } -\infty < x < s(t), 0 \leq t \leq T, \\ \sigma'(t) &\leq \sigma(t) \quad \text{if } 0 \leq t \leq T. \end{aligned}$$

PROOF. Denote by s'_n the functions defined by (3.41) when s is replaced by s' . Then $s'_n(t) \geq s_n(t)$. Now apply Lemma 3.6 to the pair $s_n(t)$, $s'_n(t)$ and take $n \rightarrow \infty$.

3.5. Existence of solutions for the Q.V.I. Let $\sigma(t)$ be a monotone decreasing function for $0 \leq t \leq T$, with $\sigma(0) = -1$. Consider the variational inequality: Find $\bar{w}(x, t)$ such that

(3.57)

$$\begin{aligned} \bar{w} &\geq 0 \quad \text{if } x > \sigma(t), 0 < t < T, \\ \bar{w}_t, \bar{w}_x, \bar{w}_{xx} &\text{ are in } L^2(\hat{G}) \quad \text{where } \hat{G} = \{(x, t); \sigma(t) < x < \infty, 0 < t < T\}, \\ \int_{\sigma(t)}^{\infty} (\bar{w}_t - \bar{w}_{xx})(v - \bar{w}) dx &\geq - \int_{\sigma(t)}^{\infty} (v - \bar{w}) dx \quad \text{for a.a. } t \in (0, T), \\ &\text{for any } v = v(x) \in L^\infty(R^1), v \geq 0 \text{ a.e.,} \\ \bar{w} &= \gamma_2 \quad \text{if } x = \sigma(t), 0 < t < T, \\ \bar{w}(x, 0) &= k_2(x) \quad \text{if } x > -1. \end{aligned}$$

We shall assume that

$$(3.58) \quad \gamma_{2x} \leq 0, \gamma_{2t} \geq 0 \quad \text{if } x \leq -1, \quad k_{2x} \leq 0 \quad \text{if } -1 \leq x \leq 1.$$

Then we can prove an analog of Lemma 3.2, namely, if $\sigma(t)$ has Hölder continuous first derivatives then $w_x \leq 0$.

Next we assume

$$(3.59) \quad k_{2xx}(x) - 1 \geq 0 \quad \text{if } -1 < x < 1;$$

there is a function $\tilde{\gamma}_2(x, t)$ with continuous derivatives

$\tilde{\gamma}_{2x}, \tilde{\gamma}_{2t}, \tilde{\gamma}_{2xx}$ such that

$$\begin{aligned} (3.60) \quad \tilde{\gamma}_2(x, t) &= \gamma_2(x, t) \quad \text{if } x < -1, \\ \tilde{\gamma}_2(x, 0) &\leq k_2(x, 0) \quad \text{if } x > -1, \\ \overline{\lim}_{x \rightarrow \infty} \tilde{\gamma}_2(x, t) &\leq 0 \quad \text{uniformly in } t, 0 \leq t \leq T, \\ \tilde{\gamma}_{2xx} - \tilde{\gamma}_t &\geq 1 \quad \text{if } x \in R^1, 0 < t < T. \end{aligned}$$

Then we can establish, analogously to Lemmas 3.3, 3.4, that

$$\bar{w}(x, t) > \gamma_2(x, t) \quad \text{if } \sigma(t) < x < -1, 0 < t < T,$$

$$\bar{w}_x - \gamma_{2x} > 0 \quad \text{if } x = \sigma(t), 0 < t < T,$$

$$\bar{w}_t > 0 \quad \text{a.e.}$$

Next one can establish an analog of Lemma 3.5 for the variational inequality (3.57).

Consider now the case where $\sigma(t)$ is merely assumed to be monotone decreasing with $\sigma(0) = -1$. Define $\sigma(t) = -1$ if $t < 0$, and

$$\sigma_n(t) = \int_{R^1} \rho_n(t - \tau) \sigma(\tau) d\tau.$$

Denote by \bar{w}_n the solution of (3.57) when $\sigma(t)$ is replaced by $\sigma_n(t)$, and denote by $x = \bar{s}_n(t)$ the corresponding free boundary. Since $\sigma_{n+1}(t) \leq \sigma_n(t)$, we have, by an analog of Lemma 3.6,

$$\bar{w}_n \leq \bar{w}_{n+1}, \quad \bar{s}_n \leq \bar{s}_{n+1}.$$

Let

$$\bar{w}(x, t) = \lim_{n \rightarrow \infty} \bar{w}_n(x, t), \quad \bar{s}(t) = \lim_{n \rightarrow \infty} \bar{s}_n(t).$$

We call \bar{w} the *generalized solution* of the variational inequality (3.57), and we call $x = \bar{s}(t)$ the *free boundary curve* for (3.57).

An analog of Lemma 3.7 is valid. In particular,

$$\bar{w}(x, t) > 0 \quad \text{if } \sigma(t) < x < \bar{s}(t),$$

$$\bar{w}(x, t) = 0 \quad \text{if } x > \bar{s}(t),$$

$$\bar{s}(t) \text{ is monotone increasing and continuous for } 0 \leq t \leq T.$$

DEFINITION OF W . Let Σ denote the class of all monotone increasing functions $s(t)$, $0 \leq t \leq T$, with $s(0) = 1$. For any $s \in \Sigma$, denote by $x = \sigma(t)$ the free boundary of the generalized solution of the variational inequality (3.9). Denote by $x = \bar{s}(t)$ the free boundary of the generalized solution of the variational inequality (3.57). Then the mapping $s \rightarrow \bar{s}$ is denoted by W , i.e., $\bar{s} = Ws$.

Notice that every $s \in \Sigma$ determines a generalized solution w of (3.9) and a free boundary $x = \sigma(t)$ which, in turn, determines a generalized solution \bar{w} of (3.57).

DEFINITION. If for some $s \in \Sigma$, $Ws = s$ then the corresponding pair (w, \bar{w}) is said to form a *generalized solution* of the Q.V.I. (3.6), (3.7). The curves $x = s(t)$, $x = \sigma(t)$ are called the *free boundary curves*.

In view of Lemma 3.7 (and the analogous result for (3.57)), the free boundary curves are continuous.

The set Σ is partially ordered by the relation:

$s_1 < s_2$ if and only if $s_1(t) < s_2(t)$ for $0 \leq t \leq T$.

Notice that every subset of elements s_α in Σ has an upper bound s in Σ , namely, $s(t) = \sup_\alpha s_\alpha(t)$, and a lower bound \tilde{s} in Σ , namely, $\tilde{s}(t) = \inf_\alpha s_\alpha(t)$.

By Lemma 3.8 and its counterpart for the variational inequality (3.57) we see that if $s_1 < s_2$ then $Ws_1 < Ws_2$, i.e., W is a monotone increasing mapping.

Now, from the methods of [10] it follows that the support of the solution \bar{w} of (3.57) (for any smooth function σ) is bounded uniformly with respect to σ . Consequently there is a constant H such that $\bar{s}(t) \leq H$ whenever $\bar{s} = Ws$, $s \in \Sigma$. Defining

$$s_1(t) = \begin{cases} 0 & \text{if } t = 0, \\ H & \text{if } 0 < t \leq T, \end{cases}$$

we conclude that $s_1 > Ws_1$.

If $s_2(t) \equiv 1$ then clearly $s_2 < Ws_2$.

Thus, W satisfies the conditions in Lemma 2 of Tartar [23]. We deduce from this lemma that W has a maximal fixed point s^* and a minimal fixed point s_* in the interval (s_1, s_2) . Since every fixed point must lie in this interval, we conclude that

$$(3.61) \quad s^* > s_*, \quad Ws^* = s^*, \quad Ws_* = s_*, \quad \text{if } Ws = s \text{ then } s^* > s > s_*.$$

Denote by (w^*, \bar{w}^*) the generalized solution of (3.6), (3.7) corresponding to s^* , and denote by (w_*, \bar{w}_*) the generalized solution of (3.6), (3.7) corresponding to s_* . We then have:

THEOREM 3.1. *Let the conditions (3.4), (3.23), (3.30), (3.31) and (3.58)–(3.60) hold. Then:*

(i) *There exist generalized solutions (w^*, \bar{w}^*) , (w_*, \bar{w}_*) of (3.6), (3.7) with the corresponding free boundary functions (s^*, σ^*) and (s_*, σ_*) .*

(ii) *If (w, \bar{w}) is any generalized solution of (3.6), (3.7) with the corresponding free boundary functions (s, σ) , then*

$$(3.62) \quad s_* \leq s \leq s^*, \quad \sigma^* \leq \sigma \leq \sigma_*,$$

$$(3.63) \quad w_* \leq w \leq w^*, \quad \bar{w}^* \leq \bar{w} \leq \bar{w}_*,$$

each inequality in (3.63) is valid in the set where both sides are defined.

(iii) *For any solution (w, \bar{w}) the free boundary functions $s(t)$, $\sigma(t)$ are continuous functions for $0 \leq t \leq T$; $s(t)$ is monotone increasing and $\sigma(t)$ is monotone decreasing.*

CHAPTER 4. STEFAN TYPE FREE BOUNDARY PROBLEM FOR SYSTEMS

In Chapters 2, 3 we have solved Q.V.I. by a fixed point theorem for a monotone increasing operator. In Chapter 5 we shall use a method of integral

equations in order to find a unique regular solution with regular free boundaries of the parabolic Q.V.I. of Chapter 3 in the special case of zero-sum game. This method of integral equations is based on the fact that the Q.V.I. (for zero-sum game) can be reformulated as a Stefan problem of melting of ice; for the case of one player, this fact was first noted by Van Moerbeke [24].

In this chapter we study (by the method of integral equations) a Stefan type free boundary problem for a system with two "temperatures," θ_1 and θ_2 . The system in Chapter 5 is somewhat different (and, in some sense, it is a special case); it can be studied by the same methods as in this chapter.

4.1. Existence and uniqueness; the increasing case. Let $g_1(x)$, $g_2(x)$ be functions defined for $-1 \leq x \leq 1$ and let $\lambda_1(x, t)$, $\lambda_2(x, t)$ be functions defined for $-\infty < x < \infty$, $t \geq 0$, satisfying:

$$(4.1) \quad \begin{aligned} &g_i(x) \text{ are continuously differentiable and } \geq 0 \text{ for } -1 \leq x \leq 1; \\ &|g_{ix}| \leq C; g_1(-1) = 0, g_2(1) = 0; \end{aligned}$$

$$(4.2) \quad \begin{aligned} &\lambda_i(x, t) \text{ are continuously differentiable and } \geq 0; \\ &|\lambda_{ix}|, |\lambda_{it}| \leq C; \lambda_1(1, 0) = g_1(1), \lambda_2(-1, 0) = g_2(-1). \end{aligned}$$

We consider the following problem: find functions $\theta_1(x, t)$, $\theta_2(x, t)$, $s_1(t)$, $s_2(t)$ such that

$$(4.3) \quad \begin{aligned} &s_1(t) < s_2(t), \quad s_1(0) = -1, \quad s_2(0) = 1, \\ &\frac{\partial \theta_i}{\partial t} = \frac{\partial^2 \theta_i}{\partial x^2} \quad \text{if } s_1(t) < x < s_2(t), t > 0, \end{aligned}$$

$$(4.4) \quad \begin{aligned} &\theta_i(s_i(t), t) = 0 \quad \text{if } t \geq 0; \\ &\theta_1 = \lambda_1 \quad \text{if } x \geq s_2(t), t \geq 0, \\ &\theta_2 = \lambda_2 \quad \text{if } x \leq s_1(t), t \geq 0, \end{aligned}$$

$$(4.5) \quad \begin{aligned} &\frac{\partial}{\partial x} \theta_i(x, t)|_{x=s_i(t)} = -\dot{s}_i(t) \quad \text{if } t \geq 0; \\ &\theta_i(x, 0) = g_i(x) \quad \text{if } -1 \leq x \leq 1; \end{aligned}$$

$$(4.6) \quad \begin{aligned} &s_i(t) \text{ is continuously differentiable for } t \geq 0, \\ &\partial \theta_1 / \partial x \text{ is continuous for } t \geq 0, x \leq s_2(t), \\ &\partial \theta_2 / \partial x \text{ is continuous for } t \geq 0, x \geq s_1(t), \\ &\partial \theta_i / \partial t, \partial^2 \theta_i / \partial x^2 \text{ are continuous for } t > 0, x \in (s_1(t), s_2(t)). \end{aligned}$$

The main result of this section is the following

THEOREM 4.1. *Under the assumptions (4.1), (4.2) there exists a unique solution $(\theta_1, \theta_2, s_1, s_2)$ of (4.3)–(4.6); furthermore $s_1(t)$ decreases and $s_2(t)$ increases as t increases.*

Since $(-1)^i \dot{s}_i(t)$ is increasing, we are dealing here with (what we call) the increasing case. Later on we shall prove a more general theorem whereby $(-1)^i \dot{s}_i(t)$ is not necessarily increasing.

The proof of Theorem 4.1 is based on the method by Friedman [12], [13, Chapter 8]. Some details will be omitted.

LEMMA 4.1. Let $s_1(t), s_2(t)$ be two curves satisfying, for $0 \leq t \leq \sigma$,

$$\begin{aligned} s_1(t) &< s_2(t), s_i(t) \text{ is continuously differentiable,} \\ s_1(0) &= -1, s_2(0) = +1. \end{aligned}$$

Let $u(x, t)$ be a solution of

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} \quad \text{if } x \in (s_1(t), s_2(t)), 0 < t < \sigma,$$

and let $\partial u / \partial x$ be continuous for $x \in [s_1(t), s_2(t)], 0 \leq t \leq \sigma$. Then we have, for $x \in (s_1(t), s_2(t)), 0 < t \leq \sigma$, the integral representation of u ,

$$\begin{aligned} (4.7) \quad u(x, t) &= \int_{-1}^{+1} K(x, t; \xi, 0) u(\xi, 0) d\xi \\ &+ \int_0^t \dot{s}_2(\tau) K(x, t; s_2(\tau), \tau) u(s_2(\tau), \tau) d\tau \\ &- \int_0^t \dot{s}_1(\tau) K(x, t; s_1(\tau), \tau) u(s_1(\tau), \tau) d\tau \\ &+ \int_0^t \left[K(x, t; \xi, \tau) \frac{\partial u}{\partial \xi}(\xi, \tau) - u(\xi, \tau) \frac{\partial K}{\partial \xi}(x, t; \xi, \tau) \right]_{s_1(\tau)}^{s_2(\tau)} d\tau \end{aligned}$$

where

$$K(x, t; \xi, \tau) = \frac{1}{2\pi^{1/2}(t - \tau)^{1/2}} = \exp \left[-\frac{(x - \xi)^2}{4(t - \tau)} \right].$$

The proof follows by using Green's formula with u, K (see [12], [13]).

Suppose now that there is a solution of the problem (4.3)–(4.6). Let

$$(4.8) \quad v_1(t) = \dot{s}_1(t) = -\frac{\partial \theta_1}{\partial x}(s_1(t), t),$$

$$(4.9) \quad v_2(t) = -\dot{s}_2(t) = \frac{\partial \theta_2}{\partial x}(s_2(t), t),$$

$$(4.10) \quad w_1(t) = -\frac{\partial \theta_1}{\partial x}(s_2(t), t),$$

$$(4.11) \quad w_2(t) = \frac{\partial \theta_2}{\partial x}(s_2(t), t).$$

We apply (4.7) to $u = \theta_1$, differentiate with respect to x , let $x \rightarrow s_1(t)$ or $x \rightarrow s_2(t)$ and use a standard jump relation [12], [13, p. 217]. Making use of (4.3)–(4.5), we arrive at the formulas:

$$\begin{aligned}
 (4.12) \quad & -\frac{1}{2} v_1(t) = \int_{-1}^{+1} K(s_1(t), t; \xi, 0) \dot{g}_1(\xi) d\xi \\
 & + \int_0^t K(s_1(t), t; s_2(\tau), \tau) [-\lambda_{1x}(s_2(\tau), \tau) v_2(\tau) + \lambda_{1\tau}(s_2(\tau), \tau)] d\tau \\
 & + \int_0^t K_x(s_1(t), t; s_1(\tau), \tau) v_1(\tau) d\tau \\
 & - \int_0^t K_x(s_1(t), t; s_2(\tau), \tau) w_1(\tau) d\tau.
 \end{aligned}$$

$$\begin{aligned}
 (4.13) \quad & -\frac{1}{2} w_1(t) = \int_{-1}^{+1} K(s_2(t), t; \xi, 0) \dot{g}_1(\xi) d\xi \\
 & + \int_0^t K(s_2(t), t; s_2(\tau), \tau) [-\lambda_{1x}(s_2(\tau), \tau) v_2(\tau) + \lambda_{1\tau}(s_2(\tau), \tau)] d\tau \\
 & + \int_0^t K_x(s_2(t), t; s_1(\tau), \tau) v_1(\tau) d\tau \\
 & - \int_0^t K_x(s_2(t), t; s_2(\tau), \tau) w_1(\tau) d\tau.
 \end{aligned}$$

A similar argument applied to θ_2 leads to

$$\begin{aligned}
 (4.14) \quad & \frac{1}{2} v_2(t) = \int_{-1}^{+1} K(s_2(t), t; \xi, 0) \dot{g}_2(\xi) d\xi \\
 & - \int_{-1}^{+1} K(s_2(t), t; s_1(\tau), \tau) [\lambda_{2x}(s_1(\tau), \tau) v_1(\tau) + \lambda_{2\tau}(s_1(\tau), \tau)] d\tau \\
 & + \int_0^t K_x(s_2(t), t; s_2(\tau), \tau) v_2(\tau) d\tau \\
 & - \int_0^t K_x(s_2(t), t; s_1(\tau), \tau) w_2(\tau) d\tau,
 \end{aligned}$$

$$\begin{aligned}
 (4.15) \quad & \frac{1}{2} w_2(t) = \int_{-1}^{+1} K(s_1(t), t; \xi, 0) \dot{g}_2(\xi) d\xi \\
 & - \int_0^t K(s_1(t), t; s_1(\tau), \tau) [\lambda_{2x}(s_1(\tau), \tau) v_1(\tau) + \lambda_{2\tau}(s_1(\tau), \tau)] d\tau \\
 & + \int_0^t K_x(s_1(t), t; s_2(\tau), \tau) v_2(\tau) d\tau \\
 & - \int_0^t K_x(s_1(t), t; s_1(\tau), \tau) w_2(\tau) d\tau.
 \end{aligned}$$

The equations (4.12)–(4.15) form a system of nonlinear integral equations of Volterra type in v_1, v_2, w_1, w_2 , noting that s_1 and s_2 are given by

$$(4.16) \quad s_1(t) = \int_0^t v_1(\tau) d\tau - 1,$$

$$(4.17) \quad s_2(t) = - \int_0^t v_2(\tau) d\tau + 1.$$

LEMMA 4.2. *There exists a positive number σ depending only on C (cf. (4.1) and (4.2)) such that there exists a unique continuous solution $(v_1(t), v_2(t), w_1(t), w_2(t), s_1(t), s_2(t))$ of (4.12)–(4.17) for $0 \leq t \leq \sigma$.*

The proof is similar to the proof in the case of the Stefan problem [12].

Suppose now that we have a solution of (4.12)–(4.17). We wish to show that this solution yields a solution of the original free boundary problem (4.3)–(4.6) with

$$(4.18) \quad \begin{aligned} \theta_1(x, t) = & \int_{-1}^{+1} K(x, t; \xi, 0) g_1(\xi) d\xi - \int_0^t v_2(\tau) K(x, t; s_2(\tau), \tau) \lambda_1(s_2(\tau), \tau) d\tau \\ & + \int_0^t [-K(x, t; s_2(\tau), \tau) w_1(\tau) + K(x, t; s_1(\tau), \tau) v_1(\tau)] d\tau \\ & + \int_0^t K_x(x, t; s_2(\tau), \tau) \lambda_1(s_2(\tau), \tau) d\tau, \end{aligned}$$

$$(4.19) \quad \begin{aligned} \theta_2(x, t) = & \int_{-1}^{+1} K(x, t; \xi, 0) g_2(\xi) d\xi - \int_0^t v_1(\tau) K(x, t; s_1(\tau), \tau) \lambda_2(s_1(\tau), \tau) d\tau \\ & + \int_0^t [K(x, t; s_2(\tau), \tau) v_2(\tau) - K(x, t; s_1(\tau), \tau) w_2(\tau)] d\tau \\ & - \int_0^t K_x(x, t; s_1(\tau), \tau) \lambda_2(s_1(\tau), \tau) d\tau. \end{aligned}$$

LEMMA 4.3. *Let $(v_1, v_2, w_1, w_2, s_1, s_2)$ be a continuous solution of (4.12)–(4.17) for $0 \leq t \leq \sigma$ such that $s_2(t) - s_1(t) \geq \text{const} > 0$. Then θ_1, θ_2 defined by (4.18), (4.19) together with s_1, s_2 form a solution of the free boundary value problem (4.3)–(4.6) for $0 \leq t \leq \sigma$.*

The proof is an easy extension of the proof in [12] for the Stefan problem.

Using the maximum principle we can show that the solution of (4.3)–(4.6) asserted in Lemma 4.3 satisfies:

$$(4.20) \quad s_1(t) \downarrow \text{ and } s_2(t) \uparrow \text{ as } t \text{ increases.}$$

PROOF OF THEOREM 4.1. We shall prove the following a priori inequalities: For any solution of (4.3)–(4.6) in an interval $0 \leq t < t_0$

$$(4.21) \quad \dot{s}_1(t) \geq -M \quad (0 \leq t < t_0),$$

$$(4.22) \quad \dot{s}_2(t) \leq N \quad (0 < t < t_0),$$

where M, N are positive constants independent of t_0 .

Using Lemmas 4.2, 4.3 and (4.20) we can then proceed step-by-step to establish the existence of a unique solution of (4.3)–(4.6); the inequalities (4.21), (4.22) guarantee that the t -interval in each step is bounded from below by a positive constant; cf. [12].

We shall estimate $\dot{s}_1(t)$ at any point $t = \bar{t}$ near t_0 , $\bar{t} < t_0$. Let R be the rectangle given by $s_1(\bar{t}) < x < 1$, $0 < t < \bar{t}$.

Let w be the solution of the heat equation in R satisfying the boundary conditions:

$$w = 0 \quad \text{for } x = s_1(\bar{t}), 0 < t < \bar{t},$$

$$w = 0 \quad \text{for } t = 0 \text{ and } s_1(\bar{t}) \leq x \leq -1,$$

$$w = g_1 \quad \text{for } t = 0 \text{ and } -1 \leq x \leq 1,$$

$$w = k \quad \text{for } x = 1, 0 < t < \bar{t},$$

where k is a constant such that

$$k > \max_{-1 \leq x \leq 1} g_1 + \max_G \lambda_1$$

where the set G is defined by $s_1(\bar{t}) \leq x \leq s_2(\bar{t})$, $0 \leq t \leq \bar{t}$. By the maximum principle $w > \theta_1$ for $x = 1$, $0 \leq t < \bar{t}$. Also $w \geq 0$ in R ; therefore $w \geq \theta_1$ for $x = s_1(t)$, $0 \leq t \leq \bar{t}$. The maximum principle then shows that $w \geq \theta_1$ in the region $s_1(t) \leq x \leq 1$, $0 \leq t \leq \bar{t}$. Noting that $(w - \theta_1)(s_1(\bar{t}), \bar{t}) = 0$, we conclude that

$$\frac{\partial}{\partial x} (w - \theta_1) \geq 0 \quad \text{at } (s_1(\bar{t}), \bar{t}),$$

i.e.,

$$(4.23) \quad -\dot{s}_1(t) = \frac{\partial \theta_1}{\partial x} (s_1(\bar{t}), \bar{t}) \leq \frac{\partial w}{\partial x} (s_1(\bar{t}), \bar{t}).$$

We also have

$$(4.24) \quad \frac{\partial w}{\partial x} (s_1(\bar{t}), \bar{t}) \leq M$$

where M is a constant independent of \bar{t} and of the position of $s_1(\bar{t})$. This can be easily checked by using the explicit representation of w in terms of the Green function in the rectangle R (given, for instance, in [13]). The assertion (4.21) follows from (4.23), (4.24). The proof of (4.22) is similar.

4.2. Asymptotic estimates. In this section we give asymptotic estimates on the behavior of function $s_2(t) - s_1(t)$ as $t \rightarrow \infty$, namely, we shall prove

THEOREM 4.2. *Let the assumptions (4.1), (4.2) hold and let*

$$(4.25) \quad \overline{\lim}_{t \rightarrow \infty} \sup_x \lambda_i(x, t) < 1 - \alpha \quad (i = 1, 2; \alpha \text{ positive constant}),$$

$$(4.26) \quad \underline{\lim}_{t \rightarrow \infty} \left\{ \inf_x \lambda_1(x, t) + \inf_x \lambda_2(x, t) \right\} > \gamma > 0 \quad (\gamma \text{ constant}).$$

Then there exist positive constants γ_1, γ_2 such that

$$(4.27) \quad \gamma_1 \leq \frac{s_2(t) - s_1(t)}{\sqrt{t}} \leq \gamma_2 \quad \text{for all } t > 1.$$

To get the lower bound we actually do not need the condition (4.25); it suffices to assume that the λ_i are bounded.

PROOF. We shall first establish an identity, which is interesting by itself. Multiplying the equation $\theta_{1xx} = \theta_{1t}$ by $x - s_2(t)$ and integrating over the domain $s_1(t) < x < s_2(t)$, $0 < t < \sigma$, we get

$$(4.28) \quad \int_0^\sigma \int_{s_1(t)}^{s_2(t)} (x - s_2(t)) \theta_{1xx} dx dt = \int_0^\sigma \int_{s_1(t)}^{s_2(t)} (x - s_2(t)) \theta_{1t} dx dt.$$

By integration by parts,

$$(4.29) \quad \int_0^\sigma dt \int_{s_1(t)}^{s_2(t)} (x - s_2(t)) \theta_{1xx} dx \\ = \int_0^\sigma [(s_1(t) - s_2(t)) \dot{s}_1(t) - \lambda_1(s_2(t), t)] dt, \\ (4.30) \quad \int \int (x - s_2(t)) \theta_{1t}(x, t) dx dt = \int_{s_1(\sigma)}^{s_2(\sigma)} (x - s_2(\sigma)) \theta_1(x, \sigma) dx \\ - \int_{-1}^1 (x - 1) g_1(x) dx + \int \int \theta_1(x, t) \dot{s}_2(t) dx dt.$$

Next, we have

$$(4.31) \quad \int_0^\sigma \int_{s_1(t)}^{s_2(t)} (x - s_1(t)) \theta_{2xx} dx dt = \int_0^\sigma \int_{s_1(t)}^{s_2(t)} (x - s_1(t)) \theta_{2t} dx dt.$$

By integration by parts,

$$(4.32) \quad \int_0^\sigma dt \int_{s_1(t)}^{s_2(t)} (x - s_1(t)) \theta_{2xx} dx \\ = \int_0^\sigma [-(s_2(t) - s_1(t)) \dot{s}_2(t) + \lambda_2(s_1(t), t)] dt, \\ (4.33) \quad \int \int (x - s_1(t)) \theta_{2t}(x, t) dx dt = \int_{s_1(\sigma)}^{s_2(\sigma)} (x - s_1(\sigma)) \theta_2(x, \sigma) dx \\ - \int_{-1}^{+1} (x + 1) g_2(x) dx + \int \int \theta_2(x, t) \dot{s}_1(t) dx dt.$$

Since by (4.28) the right-hand sides of (4.29), (4.30) are equal and, by (4.31), the right-hand sides of (4.32), (4.33) are equal, we get, by adding,

$$\begin{aligned} \int_0^\sigma (s_1 - s_2)(\dot{s}_1 - \dot{s}_2) dt &= \int_0^\sigma [\lambda_1(s_2(t), t) + \lambda_2(s_1(t), t)] dt \\ &\quad + \int_{-1}^{+1} [(1-x)g_1(x) + (1+x)g_2(x)] dx \\ &\quad - \int_{s_1(\sigma)}^{s_2(\sigma)} [(s_2(\sigma) - x)\theta_1(x, \sigma) + (x - s_1(\sigma))\theta_2(x, \sigma)] dx \\ &\quad + \iint (\theta_1 \dot{s}_2 - \theta_2 \dot{s}_1) dx dt, \end{aligned}$$

or

$$\begin{aligned} \frac{1}{2} (s_2(\sigma) - s_1(\sigma))^2 &= 2 + \int_0^\sigma (\lambda_1 + \lambda_2) dt + \int_{-1}^{+1} [(1-x)g_1 + (1+x)g_2] dx \\ (4.34) \quad &- \int_{s_1(\sigma)}^{s_2(\sigma)} [(s_2(\sigma) - x)\theta_1(x, \sigma) + (x - s_1(\sigma))\theta_2(x, \sigma)] dx \\ &+ \int_0^\sigma \int_{s_1(t)}^{s_2(t)} (\theta_1 \dot{s}_2 - \theta_2 \dot{s}_1) dx dt. \end{aligned}$$

Let us now establish the lower bound in (4.27). We note that the double integral in (4.34) is ≥ 0 . Since $\theta_i \leq M$ (M constant), we get from (4.34)

$$\frac{1}{2} (s_2(\sigma) - s_1(\sigma))^2 + M (s_2(\sigma) - s_1(\sigma))^2 \geq \gamma \sigma - N$$

where γ, N are positive constants, from which follows the left-hand side of (4.27).

To prove the right-hand side, let t^* be such that

$$\lambda_i(x, t) \leq 1 - \alpha \quad \text{if } -\infty < x < \infty, t \geq t^*.$$

We claim that there exists a $t_0 > t^*$ such that

$$(4.35) \quad \theta_i(x, t) \leq 1 - \beta, \quad 0 < \beta < \alpha \text{ if } s_1(t) < x < s_2(t), t > t_0.$$

Indeed, let $M \geq g_i(x)$, $M \geq 1 - \alpha$. Consider the solution w of the heat equation in $-\infty < x < \infty, t > t^*$, with initial conditions

$$w(x, t) = \begin{cases} M & \text{if } s_1(t^*) < x < s_2(t^*), \\ 1 - \alpha & \text{if } x < s_1(t^*) \text{ or } x > s_2(t^*). \end{cases}$$

In the domain $s_1(t) < x < s_2(t), t > t^*$ we have, by the maximum principle, $w \geq \theta_i$. Now, with $s_i(t^*) = \alpha_i$,

$$\begin{aligned}
w(x, t) = & \int_{\alpha_1}^{\alpha_2} \frac{M}{(4\pi(t-t^*))^{1/2}} \exp\left[-\frac{(x-\xi)^2}{4(t-t^*)}\right] d\xi \\
& + \int_{\alpha_2}^{\infty} \frac{1-\alpha}{(4\pi(t-t^*))^{1/2}} \exp\left[-\frac{(x-\xi)^2}{4(t-t^*)}\right] d\xi \\
& + \int_{-\infty}^{\alpha_1} \frac{1-\alpha}{(4\pi(t-t^*))^{1/2}} \exp\left[-\frac{(x-\xi)^2}{4(t-t^*)}\right] d\xi.
\end{aligned}$$

Therefore

$$w(x, t) \leq 1 - \alpha + M(\alpha_2 - \alpha_1) / (4\pi(t - t^*))^{1/2}.$$

Taking $t \geq t_0$ where $t_0 - t^*$ is sufficiently large, we obtain the assertion (4.35).

We can now write an identity similar to (4.34), when $x \in (s_1(t), s_2(t))$, $t_0 \leq t \leq \sigma$ (instead of $0 \leq t \leq \sigma$):

$$\begin{aligned}
\frac{1}{2} (s_2(\sigma) - s_1(\sigma))^2 = & \frac{1}{2} (s_2(t_0) - s_1(t_0))^2 + \int_{t_0}^{\sigma} (\lambda_1 + \lambda_2) dt \\
& + \int_{s_1(t_0)}^{s_2(t_0)} [(s_2(t_0) - x)\theta_1(x, t_0) + (x - s_1(t_0))\theta_2(x, t_0)] dx \\
& - \int_{s_1(\sigma)}^{s_2(\sigma)} [(s_2(\sigma) - x)\theta_1(x, \sigma) + (x - s_1(\sigma))\theta_2(x, \sigma)] dx \\
& + \int \int (\theta_1 \dot{s}_2 - \theta_2 \dot{s}_1) dx dt.
\end{aligned}$$

Using (4.35) we get

$$\frac{1}{2} (s_2(\sigma) - s_1(\sigma))^2 \leq 2(1 - \alpha)\sigma + O(1) + (1 - \beta) \int_{t_0}^{\sigma} (\dot{s}_2 - \dot{s}_1)(s_2 - s_1) dt.$$

Hence

$$\beta (s_2(\sigma) - s_1(\sigma))^2 / 2 \leq 2(1 - \alpha)\sigma + O(1),$$

which completes the proof of (4.27).

4.3. The decreasing case. In this section we shall extend Theorem 4.1 to the case where the functions $g_1, g_2, \lambda_1, \lambda_2$ are nonpositive. Thus, we assume:

$$(4.36) \quad \begin{aligned} & g_i(x) \text{ are continuously differentiable and } \leq 0 \text{ for } -1 \leq x \leq 1; \\ & |g_{ix}| \leq C, g_1(-1) = 0, g_2(1) = 0; \end{aligned}$$

$$(4.37) \quad \begin{aligned} & \lambda_i(x, t) \text{ are continuously differentiable and } \leq 0; \\ & |\lambda_{ix}|, |\lambda_{it}| \leq C, \lambda_1(1, 0) = g_1(1), \lambda_2(-1, 0) = g_2(-1). \end{aligned}$$

We shall also assume:

$$(4.38) \quad \inf_{-1 \leq x \leq 1} g_1(x) > -1, \quad \inf_{-1 \leq x \leq 1} g_2(x) > -1, \\ \inf_{-1 \leq x \leq 1, t > 0} \lambda_1(x, t) > -1, \quad \inf_{-1 \leq x \leq 1, t > 0} \lambda_2(x, t) > -1.$$

Let $\lambda_i(t)$, $\tilde{\lambda}_i(t)$ be functions satisfying $\tilde{\lambda}_i(t) \leq \lambda_i(x, t) \leq \lambda_i(t) \leq 0$ and let

$$\begin{aligned} \Gamma_1 &= - \int_0^\infty \lambda_1(t) dt - \int_{-1}^1 (1-x) g_1(x) dx, \\ \Gamma_2 &= - \int_0^\infty \lambda_2(t) dt - \int_{-1}^1 (1+x) g_2(x) dx, \\ \tilde{\Gamma}_1 &= - \int_0^\infty \tilde{\lambda}_1(t) dt - \int_{-1}^1 (1-x) g_1(x) dx, \\ \tilde{\Gamma}_2 &= - \int_0^\infty \tilde{\lambda}_2(t) dt - \int_{-1}^1 (1+x) g_2(x) dx. \end{aligned}$$

Notice that $0 \leq \Gamma_i \leq \tilde{\Gamma}_i \leq \infty$ ($i = 1, 2$).

THEOREM 4.3. *Under the assumptions (4.36)–(4.38), there exists a unique solution $(\theta_1, \theta_2, s_1, s_2)$ of (4.3)–(4.6) for all $0 \leq t < T^*$ for some $T^* \in (0, \infty]$; further, $s_1(t)$ increases and $s_2(t)$ decreases as t increases. If*

$$(4.39) \quad \Gamma_1 + \Gamma_2 > 4$$

then $T^ < \infty$, and if*

$$(4.40) \quad \tilde{\Gamma}_1 + \tilde{\Gamma}_2 < 2$$

then $T^ = \infty$.*

PROOF. Let us first show that if there is a solution for $0 \leq t < t_0$, such that

$$(4.41) \quad s_2(t) - s_1(t) \geq \text{const} > 0 \quad \text{for } 0 \leq t < t_0,$$

then

$$(4.42) \quad \dot{s}_1(t) \leq M \quad \text{if } 0 \leq t < t_0,$$

$$(4.43) \quad -N \leq \dot{s}_2(t) \quad \text{if } 0 \leq t < t_0,$$

where M, N are positive constants.

To prove (4.42) it is sufficient to verify this inequality at any point $\bar{t} < t_0$ with the property that $\dot{s}_1(t) \leq \dot{s}_1(\bar{t})$ if $0 \leq t \leq \bar{t}$.

We shall need the following lemma.

LEMMA 4.4. *Let a, b, θ, σ, t be positive numbers such that $\theta = t - \sigma$, $a < b$. Let $\psi(\tau) = \alpha - a + b(\tau - \sigma)$ where α is a real number and τ varies in the interval $[\sigma, t]$. Let $W = \{(\xi, \tau); \psi(\tau) < \xi < \infty, \sigma < \tau < t\}$. Let w be the bounded solution of*

$$\frac{\partial w}{\partial \tau} = \frac{\partial^2 w}{\partial \xi^2} \quad \text{in } W,$$

$$w(\xi, \sigma) = 0 \quad \text{if } \alpha - a < \xi < \alpha,$$

$$w(\xi, \sigma) = k(\xi) \quad \text{if } \alpha < \xi < \infty \quad (-k_0 \leq k(\xi) \leq 0),$$

$$w(\psi(\tau), \tau) = 0 \quad \text{if } \sigma < \tau < t.$$

Then

$$\begin{aligned} (4.44) \quad 0 &\leq -\frac{\partial w}{\partial \xi}(\psi(t), t) \\ &\leq -b \int_{\sigma}^t \frac{b\tilde{k}(\tau)}{(4\pi(t-\tau))^{1/2}} \exp\left[-\frac{b^2(t-\tau)}{4}\right] d\tau + k_0 C(\theta) \end{aligned}$$

where

$$\tilde{k}(\tau) = \int_{\alpha}^{\infty} \frac{k(\xi)}{(4\pi(\tau-\sigma))^{1/2}} \exp\left[-\frac{(\psi(\tau)-\xi)^2}{4(\tau-\sigma)}\right] d\xi$$

and $C(\theta)$ is a constant independent of a, b, k .

The proof is given in the Appendix.

Notice that (4.44) implies that

$$0 \leq -\partial w(\psi(t), t)/\partial \xi \leq k_0 b + C(\theta).$$

We shall apply Lemma 4.4 with

$$\begin{aligned} t &= \bar{t}, \sigma = \bar{t} - \delta, b = \dot{s}_1(\bar{t}), \alpha = s_1(\bar{t} - \delta), s_1(\bar{t}) = \alpha - a + b\delta, \delta > 0, \\ k &= \min(\inf g_1, \inf \lambda_1) - \epsilon, \quad -1 < k < 0. \end{aligned}$$

Notice, by (4.38), that $\epsilon > 0$ can indeed be chosen so that $-1 < k < 0$. By the maximum principle, $w \leq 0$. Hence $w(s_1(\tau), \tau) \leq \theta_1(s_1(\tau), \tau)$ if $\bar{t} - \delta \leq \tau \leq \bar{t}$. Next

$$w(\xi, \bar{t} - \delta) < \theta_1(\xi, \bar{t} - \delta) \quad \text{if } s_1(\bar{t} - \delta) \leq \xi \leq s_2(\bar{t} - \delta).$$

Finally, if δ is sufficiently small (depending on ϵ) then

$$w(s_2(\tau), \tau) < \lambda_1(s_2(\tau), \tau) = \theta_1(s_2(\tau), \tau) \quad \text{for } \bar{t} - \delta < \tau < \bar{t}.$$

By the maximum principle we conclude that $w \leq \theta_1$ if $s_1(\tau) < \xi < s_2(\tau)$, $0 < \tau < \bar{t}$. Since $w(s_1(\bar{t}), \bar{t}) = \theta_1(s_1(\bar{t}), \bar{t})$, it follows that

$$\partial(\theta_1 - w)(s_1(\bar{t}), \bar{t})/\partial \xi \geq 0.$$

Consequently,

$$b = \dot{s}_1(\bar{t}) = -\frac{\partial}{\partial \xi} \theta_1(s_1(\bar{t}), \bar{t}) \leq -\frac{\partial}{\partial \xi} w(s_1(\bar{t}), \bar{t}) \leq -k(b + C(\theta))$$

by Lemma 4.4, i.e., $(1 + k)b \leq C(\theta)$. Since $1 + k > 0$, we get $b \leq \text{const}$. This completes the proof of (4.42). The proof of (4.43) is similar.

We can now proceed to complete the proof of existence and uniqueness for (4.3)–(4.6) as in the case of Theorem 4.1. As long as (4.41) holds for a constant arbitrarily small, the solution can be continued in a unique way. By letting the constant go to 0 we arrive at a maximal interval $0 \leq t < T^*$ where the solution of (4.3)–(4.6) exists and is unique. In general, $T^* \leq \infty$. If $T^* < \infty$ then we must have $s_1(T^* - 0) = s_2(T^* - 0)$, whereas if $T^* = \infty$ then $s_1(t) < s_2(t)$ for all $0 \leq t < \infty$.

We shall now assume that (4.39) holds and prove that $T^* < \infty$.

Let $0 < \sigma < T^*$. Multiplying the heat equation for θ_1 by $x - s_2(t)$ and integrating with respect to (x, t) , over the region $s_1(t) < x < s_2(t)$, $0 < t < \sigma$, we get

$$\int dt \int_{s_1(t)}^{s_2(t)} (x - s_2(t)) \frac{\partial^2 \theta_1}{\partial x^2} dx = \int_{-1}^1 dx \int_0^{l(x)} (x - s_2(t)) \frac{\partial \theta_1}{\partial t} dt$$

where

$$l(x) = \begin{cases} l_1(x) & \text{if } -1 < x < s_1(\sigma), \\ \sigma & \text{if } s_1(\sigma) < x < s_2(\sigma), \\ l_2(x) & \text{if } s_2(\sigma) < x < 1 \end{cases}$$

and l_i is the inverse function to s_i .

Integrating by parts, we obtain

$$\begin{aligned} \int_0^\sigma (s_1(t) - s_2(t)) \dot{s}_1(t) dt - \int_0^\sigma \lambda_1(s_2(t), t) dt \\ = \int_{-1}^1 (x - s_2(l(x))) \theta_1(x, l(x)) dx \\ - \int_{-1}^1 (x - 1) g_1(x) dx + \int_{-1}^1 \int_0^{l(x)} \dot{s}_2(t) \theta_1(x, t) dt dx, \end{aligned}$$

or,

$$\begin{aligned} - \int_0^\sigma \lambda_1(s_2(t), t) dt = \int_0^\sigma (s_2(t) - s_1(t)) \dot{s}_1(t) dt \\ + \int_{s_1(\sigma)}^{s_2(\sigma)} (x - s_2(\sigma)) \theta_1(x, \sigma) dx \\ + \int_{-1}^1 (1 - x) g_1(x) dx + \int_0^\sigma \left[\int_{s_1(t)}^{s_2(t)} \theta_1(x, t) dx \right] \dot{s}_2(t) dt. \end{aligned} \quad (4.45)$$

By the maximum principle, $-1 \leq \theta_1 \leq 0$. Using also the inequalities $\dot{s}_1 \geq 0$, $\dot{s}_2 \leq 0$, we find that the sum of the first and last integrals on the right-hand side of (4.45) is

$$\leq \int_0^\sigma (s_2(t) - s_1(t))(\dot{s}_1(t) - \dot{s}_2(t)) dt = 2 - \frac{1}{2} (s_2(\sigma) - s_1(\sigma))^2.$$

The second integral on the right is

$$\leq \int_{s_1(\sigma)}^{s_2(\sigma)} (s_2(\sigma) - x) dx = \frac{1}{2} (s_2(\sigma) - s_1(\sigma))^2.$$

Consequently

$$- \int_0^\sigma \lambda_1(s_2(t), t) dt \leq 2 + \int_{-1}^1 (1 - x) g_1(x) dx.$$

Similarly,

$$- \int_0^\sigma \lambda_2(s_1(t), t) dt \leq 2 + \int_{-1}^1 (1 + x) g_2(x) dx.$$

If $T^* = \infty$ then the last two inequalities hold also with $\sigma = \infty$. But then $\Gamma_1 + \Gamma_2 \leq 4$, which contradicts (4.39).

It remains to prove that if (4.40) holds then $T^* = \infty$. Suppose that $T^* < \infty$ and take $\sigma \uparrow T^*$ in (4.45). Then the second integral on the right converges to zero. Noting that the last integral on the right-hand side of (4.45) is nonnegative, we find that

$$(4.46) \quad - \int_0^{T^*} \tilde{\lambda}_1(t) dt - \int_{-1}^1 (1 - x) g_1(x) dx \geq \lim_{\sigma \uparrow T^*} \int_0^\sigma (s_2(t) - s_1(t)) \dot{s}_1(t) dt.$$

Similarly,

$$(4.47) \quad - \int_0^{T^*} \tilde{\lambda}_2(t) dt - \int_{-1}^1 (1 + x) g_2(x) dx \geq \lim_{\sigma \uparrow T^*} \int_0^\sigma (s_1(t) - s_2(t)) \dot{s}_2(t) dt.$$

Since

$$\int_0^\sigma (s_2 - s_1) \dot{s}_1 dt + \int_0^\sigma (s_1 - s_2) \dot{s}_2 dt = 2 - \frac{1}{2} (s_2(\sigma) - s_1(\sigma))^2 \rightarrow 2$$

as $\sigma \uparrow T^*$, we obtain, upon adding (4.46), (4.47), $\tilde{\Gamma}_1 + \tilde{\Gamma}_2 \geq 2$, which contradicts (4.40).

4.4. The general case. In this section we shall consider the general case where the functions g_i , λ_i need not have the same sign everywhere. More specifically we shall assume:

$$(4.48) \quad \begin{aligned} &g_{ix}(x) \text{ are continuously differentiable for } -1 \leq x \leq 1, \\ &|g_{ix}| \leq C, \text{ and } g_1(-1) = 0, g_2(1) = 0; \end{aligned}$$

$$(4.49) \quad \begin{aligned} &\lambda_i(x, t) \text{ are continuously differentiable and } |\lambda_{ix}|, |\lambda_{it}| \leq C; \\ &\lambda_1(1, 0) = g_1(1), \quad \lambda_2(-1, 0) = g_2(-1); \end{aligned}$$

$$(4.50) \quad \begin{aligned} &g_1(\alpha_i) = 0 \quad \text{for } \alpha_0 = -1 < \alpha_1 < \alpha_2 < \cdots < \alpha_k < 1, \\ &g_1(x) \neq 0 \quad \text{if } -1 < x < 1 \text{ and } x \neq \alpha_i \text{ for all } i, \\ &g_1(x) \text{ changes sign as } x \text{ crosses } \alpha_i \text{ (} 1 \leq i \leq k \text{),} \\ &g_2(\beta_i) = 0 \quad \text{for } \beta_0 = 1 > \beta_1 > \beta_2 > \cdots > \beta_l > -1, \\ &g_2(x) \neq 0 \quad \text{if } -1 < x < 1 \text{ and } x \neq \beta_i \text{ for all } i, \\ &g_2(x) \text{ changes sign as } x \text{ crosses } \beta_i \text{ (} 1 \leq i \leq l \text{).} \end{aligned}$$

THEOREM 4.4. *Under the assumptions (4.48)–(4.50) and (4.38), there exists a unique solution $(\theta_1, \theta_2, s_1, s_2)$ of (4.3)–(4.6) for all $0 \leq t < T$ for some $T \in (0, \infty)$; further $s_1(t)$ and $s_2(t)$ are both piecewise monotone functions, and $s_2(T - 0) = s_1(T - 0)$ in case $T < \infty$.*

From the proof it will follow that the direction of monotonicity of $s_1(t)$ ($s_2(t)$) changes at most k (l) times.

PROOF. One can prove the local existence and uniqueness of a solution in precisely the same way as in Theorem 4.1. Thus, all that remains to be shown is the piecewise monotonicity of $s_i(t)$ and the a priori bound on $\dot{s}_i(t)$.

Denote by γ_i the curve defined by $\theta_i = 0$, initiating at $(\alpha_i, 0)$; these curves are constructed in Friedman [16] and Van Moerbeke [24].

Suppose $g_1(x) > 0$ if $-1 < x < \alpha_1$. We claim that as long as γ_1 does not intersect $x = s_1(t)$ the function $s_1(t)$ is monotone decreasing. Indeed, by the maximum principle applied to θ_1 in the region bounded by $x = s_1(t)$, γ_1 and $t = 0$ we find that θ_1 takes its minimum on the boundary $x = s_1(t)$. Hence

$$\dot{s}_1(t) = -\theta_{1x}(s_1(t), t) \leq 0.$$

One can derive a priori bound on $\dot{s}_1(t)$ (as long as $x = s_1(t)$ does not intersect γ_1) by the method of §4.1.

Consider next the case where $g_1(x) < 0$ if $-1 < x < \alpha_1$. Using an argument similar to one given above we deduce that $\dot{s}_1(t) \geq 0$ as long as $x = s_1(t)$ does not intersect γ_1 . To find a priori bound on $\dot{s}_1(t)$ we use Lemma 4.4 (as in the proof of Theorem 4.3). However, here we may simply take $\sigma = 0$, i.e., $\delta = t$.

Denote by t_1 the first time γ_1 intersects $x = s_1(t)$. For $t > t_1$ we have to take into account the curve γ_2 . As long as γ_2 does not intersect $x = s_1(t)$, the function $s_1(t)$ is monotone (with the direction of monotonicity reversed to the direction of monotonicity in the interval $0 < t < t_1$). Furthermore, one can estimate $\dot{s}_1(t)$ as before. If γ_2 intersects $x = s_1(t)$ at time t_2 , then for $t > t_2$ we have to consider the curve γ_3 ; etc.

The monotone behavior of $s_2(t)$ and the a priori bounds on $\dot{s}_2(t)$ can be obtained in a similar manner.

Notice that as long as $s_2(t) - s_1(t)$ remains positive, we can continue with the construction of the solution step-by-step in t . After time $t = t_{k_0}$ when the last γ_{k_0} intersects $x = s_1(t)$ ($1 \leq k_0 \leq k$), the derivation of the a priori bound on $s_1(t)$ is slightly different than before, in the monotone decreasing case. Indeed, when we apply Lemma 4.4 we now have to take $t = \bar{t}$, $\sigma = \bar{t} - \delta$ with δ sufficiently small (cf. the proof of Theorem 4.3). The same discussion applies to the curve $x = s_2(t)$.

CHAPTER 5. ZERO-SUM STOCHASTIC GAME WITH STOPPING TIMES

5.1. Formal derivation of the Stefan problem. For zero-sum game,

$$f_2 = -f_1 = -f, \quad \phi_2 = -\psi_1 = -\phi, \quad \psi_2 = -\phi_1 = -\phi.$$

This case has been already studied by Friedman [14], as far as the existence and uniqueness are concerned. We want here to relate this problem to the methods of Chapter 4 and study the corresponding Stefan problem.

Let $Q = \{(x, t); -\infty < x < \infty, 0 < t < T\}$ and consider a function $u(x, t)$ satisfying

$$(5.1) \quad u \text{ is continuous in } \bar{Q}, (u_t + u_{xx}) \in L^2_{\text{loc}}(Q),$$

$$(5.2) \quad \psi \leq u \leq \phi \text{ in } Q,$$

$$(5.3) \quad \text{if } u > \psi \text{ then } -u_t - u_{xx} \leq f,$$

$$(5.4) \quad \text{if } u < \phi \text{ then } -u_t - u_{xx} \geq f,$$

$$(5.5) \quad u(x, T) = h(x) \text{ if } -\infty < x < \infty.$$

We assume throughout this section that ϕ, ψ are in $C^2(\bar{Q})$, f is in $C(\bar{Q})$, h, h' are continuous for all x , and h'' is continuous in $[-1, 1]$. It is easy to check that $u_1 = u$ and $u_2 = -u$ satisfy all the sufficient conditions of Theorem 1.1; consequently, a solution of (5.1)–(5.5) provides a Nash equilibrium point.

The existence and uniqueness of the solution of (5.1)–(5.5) follows from the general theory of parabolic variational inequalities of Lions-Stampacchia [21] (see also [14], [15]). The connection between parabolic variational inequalities and Stefan problems has been developed by Duvaut [11], Friedman [16] and Friedman and Kinderlehrer [19]; all these authors worked with one-sided inequality.

We shall first derive formally the Stefan problem corresponding to (5.1)–(5.5). For convenience, we shall change the time from t into $T - t$. If u satisfies:

$$(5.6) \quad u \text{ is continuous in } \bar{Q}, (u_t - u_{xx}) \in L^2_{\text{loc}}(Q),$$

$$(5.7) \quad \psi \leq u \leq \phi,$$

$$(5.8) \quad \text{if } u > \psi \text{ then } u_t - u_{xx} \leq f,$$

$$(5.9) \quad \text{if } u < \phi \text{ then } u_t - u_{xx} \geq f,$$

$$(5.10) \quad u(x, 0) = h(x),$$

then $u(x, T - t)$ satisfies (5.1)–(5.5) with $f(x, t)$, $\phi(x, t)$, $\psi(x, t)$ replaced by $f(x, T - t)$, $\phi(x, T - t)$, $\psi(x, T - t)$.

Let $s_1(t)$ and $s_2(t)$ be the two boundaries separating respectively $u < \phi$ from $u = \phi$ and $u > \psi$ from $u = \psi$ respectively. Let us assume that these functions satisfy

$$(5.11) \quad s_1(t) < s_2(t), s_1(0) = -1, s_2(0) = 1,$$

$$(5.12) \quad \dot{s}_1 \leq 0, \dot{s}_2 \geq 0.$$

Let $l_1(x)$, $l_2(x)$ be the inverse functions of $s_1(t)$, $s_2(t)$ respectively. We define $\theta(x, t)$ by the following relations:

$$(5.13) \quad \theta = 0 \quad \text{if } x \leq -1, t \leq l_1(x),$$

$$(5.14) \quad \phi - u = \int_{l_1(x)}^t \theta(x, \tau) d\tau \quad \text{if } x \leq 1, t \geq l_1(x),$$

$$(5.15) \quad \phi - u = \int_0^t \theta(x, \tau) d\tau - h(x) \quad \text{if } -1 \leq x \leq 1, t > 0,$$

$$(5.16) \quad \begin{aligned} \phi - u = \int_{l_2(x)}^t \theta(x, \tau) d\tau + \phi(x, l_2(x)) - \psi(x, l_2(x)) \\ \text{if } x \geq 1, t \geq l_2(x), \end{aligned}$$

$$(5.17) \quad \theta = \phi_t - \psi_t \quad \text{if } x \geq 1, t \leq l_2(x).$$

One easily checks that

$$(5.18) \quad \phi_t - u_t = \theta.$$

We shall assume:

$$(5.19) \quad h = 0 \quad \text{for } x \leq -1, \quad h \leq 0 \quad \text{if } x \geq -1;$$

$$h(x) \geq \psi(x, 0) \quad \text{for } x \in [-1, +1],$$

$$(5.20) \quad h(x) = \psi(x, 0) \quad \text{for } x \geq 1,$$

$$\phi(x, 0) \equiv 0 \quad \text{for } x \geq 1;$$

$$(5.21) \quad f - \phi_t + \phi_{xx} = 1.$$

From the regularity conditions (5.6) it follows that a.e. in t , u_x is continuous in x . Therefore on the free boundaries we have

$$(5.22) \quad u_x(s_1(t), t) = \phi_x(s_1(t), t),$$

$$(5.23) \quad u_x(s_2(t), t) = \psi_x(s_2(t), t).$$

Differentiating (5.14) in x and taking $t = l_1(x)$, we get, taking (5.22) into account,

$$(5.24) \quad \theta(s_1(t), t) = 0.$$

Differentiating (5.16) with respect to x , we get

$$(5.25) \quad \begin{aligned} \phi_x(x, t) - u_x(x, t) = & -\dot{l}_2(x)\theta(x, l_2(x)) + (\phi_x - \psi_x)(x, l_2(x)) \\ & + \dot{l}_2(x)(\phi_t - \psi_t)(x, l_2(x)) + \int_{l_2(x)}^t \theta_x(x, \tau) d\tau. \end{aligned}$$

Taking $t = l_2(x)$ and using (5.23), we get

$$(5.26) \quad \theta(s_2(t), t) = (\phi_t - \psi_t)(s_2(t), t).$$

In the region where (5.14) holds, we get by differentiating twice in x ,

$$(\phi_{xx} - u_{xx})(x, t) = -\dot{l}_1(x)\theta_x(x, l_1(x)) + \int_{l_1(x)}^t \theta_{xx}(x, \tau) d\tau.$$

Therefore, using the fact that $\theta(x, l_1(x)) = 0$ (i.e., (5.24)) we get, taking (5.21) into account,

$$1 = -\dot{l}_1(x)\theta_x(x, l_1(x)) - \int_{l_1(x)}^t (\theta_\tau - \theta_{xx})(x, \tau) d\tau$$

from which we deduce

$$(5.27) \quad \theta_x(s_1(t), t) = -\dot{s}_1(t),$$

$$(5.28) \quad \theta_1 - \theta_{xx} = 0 \quad \text{if } x < -1, t > l_1(x).$$

From (5.15) we deduce by similar calculations

$$1 + \int_0^t (\theta_\tau - \theta_{xx})(x, \tau) d\tau + \theta(x, 0) = -h''(x);$$

hence

$$(5.29) \quad \theta(x, 0) = -(1 + h''(x)) \quad \text{if } x \in (-1, +1)$$

$$(5.30) \quad \theta_t - \theta_{xx} = 0 \quad \text{if } x \in (-1, +1), t > 0.$$

Next, using (5.26) in (5.25), and differentiating (5.25) with respect to x , we get

$$\begin{aligned} (\phi_{xx} - u_{xx})(x, t) = & (\phi_{xx} - \psi_{xx})(x, l_2(x)) + \dot{l}_2(x)(\phi_{xt} - \psi_{xt})(x, l_2(x)) \\ & - \dot{l}_2(x)\theta_x(x, l_2(x)) + \int_{l_2(x)}^t \theta_{xx}(x, \tau) d\tau. \end{aligned}$$

After some rearrangement,

$$(5.31) \quad \begin{aligned} 1 + (\phi_t - \psi_t - (\phi_{xx} - \psi_{xx}))(x, l_2(x)) + \int_{l_2(x)}^t (\theta_\tau - \theta_{xx})(x, \tau) d\tau \\ = \dot{l}_2(x)((\phi_{xt} - \psi_{xt})(x, l_2(x)) - \theta_x(x, l_2(x))). \end{aligned}$$

Setting

$$(5.32) \quad \lambda = \phi_t - \psi_t, \quad -\mu = 1 + \phi_t - \psi_t - \phi_{xx} + \psi_{xx},$$

we get from (5.31)

$$(5.33) \quad \theta_x(s_2(t), t) - \lambda_x(s_2(t), t) = \dot{s}_2(t) \mu(s_2(t), t),$$

$$(5.34) \quad \theta_t - \theta_{xx} = 0 \quad \text{if } x > 1, t > l_2(x).$$

5.2. Verification theorem. We can now state our Stefan problem: find functions $\theta(x, t)$, $s_1(t)$, $s_2(t)$ such that

$$(5.35) \quad s_1(t) < s_2(t), \quad s_1(0) = -1, \quad s_2(0) = +1,$$

$$(5.36) \quad \partial\theta/\partial t - \partial^2\theta/\partial x^2 = 0 \quad \text{if } x \in (s_1(t), s_2(t)), t > 0,$$

$$(5.37) \quad \theta(s_1(t), t) = 0, \quad t \geq 0,$$

$$(5.38) \quad \theta(s_2(t), t) = \lambda(s_2(t), t), \quad t \geq 0,$$

$$(5.39) \quad \theta_x(s_1(t), t) = -\dot{s}_1(t), \quad t \geq 0,$$

$$(5.40) \quad \theta_x(s_2(t), t) = \lambda_x(s_2(t), t) + \mu(s_2(t), t)\dot{s}_2(t), \quad t \geq 0,$$

$$(5.41) \quad \theta(x, 0) = g(x), \quad x \in [-1, +1],$$

s_1, s_2 are continuously differentiable for $t \geq 0$,

$$(5.42) \quad \theta, \theta_x, \theta_t \text{ are continuous in } (x, t),$$

θ_{xx} is continuous for $x \in (s_1(t), s_2(t)), t > 0$.

We shall make the following assumptions:

$$(5.43) \quad g(x) = -(1 + h''(x)) \geq 0, \quad g(-1) = 0,$$

g is continuously differentiable in $[-1, +1]$;

$$(5.44) \quad \lambda \text{ is continuously differentiable, } |\lambda_x| \leq C, \quad |\lambda_t| \leq C, \quad \lambda \geq 0;$$

$$(5.45) \quad \lambda(x, 0) \geq g(x) \quad \text{if } -1 \leq x \leq 1, \quad \lambda(1, 0) = g(1);$$

$$(5.46) \quad 0 < \alpha \leq \mu \leq C_0, \mu_t \leq 0 \quad (\alpha, C_0 \text{ positive constants}).$$

Suppose now that there exists a solution of (5.35)–(5.42). Since $g \geq 0$, $\lambda \geq 0$, it follows from the maximum principle that $\theta \geq 0$. Since $\theta = 0$ on $x = s_1(t)$, we see that $\theta_x(s_1(t), t) \geq 0$; hence from (5.39) we get

$$(5.47) \quad \dot{s}_1(t) \leq 0.$$

Consider next the function $w = \theta - \lambda$. Clearly, $w = 0$ on $(s_2(t), t)$, $w = -\lambda \leq 0$ on $(s_1(t), t)$ and $w = g - \lambda(x, 0) \leq 0$ for $t = 0$. Since

$$w_t - w_{xx} = \theta_t - \theta_{xx} - \lambda_t + \lambda_{xx} = \mu_t \leq 0,$$

it follows from the maximum principle that $w \leq 0$. Since $w = 0$ on $(s_2(t), t)$, we have

$$(\theta_x - \lambda_x)(s_2(t), t) \geq 0.$$

Since $\mu > 0$, we get from (5.40) that

$$(5.48) \quad \dot{s}_2(t) \geq 0.$$

We next define $u(x, t)$ from $\theta(x, t)$ by using formulas (5.14)–(5.16) and

$$(5.49) \quad u = \phi \quad \text{for } x \leq -1, t \leq l_1(x),$$

$$(5.50) \quad u = \psi \quad \text{for } x \geq 1, t \leq l_2(x),$$

and check that it is indeed a solution of (5.6)–(5.10). Clearly u is continuous in \bar{Q} . We next have $\phi_t - u_t = \theta$ which proves that u_t is continuous. Next

$$u_x = \phi_x \quad \text{for } x \leq -1, t \leq l_1(x),$$

$$\phi_x - u_x = \int_{l_1(x)}^t \theta_x(x, \tau) d\tau \quad \text{for } x \leq -1, t > l_1(x),$$

$$\phi_x - u_x = \int_0^t \theta_x(x, \tau) d\tau - h'(x) \quad \text{for } -1 \leq x \leq 1, t > 0,$$

$$\phi_x - u_x = \int_{l_2(x)}^t \theta_x(x, \tau) d\tau + \phi_x(x, l_2(x)) - \psi_x(x, l_2(x))$$

$$\text{for } x \geq 1, t > l_2(x),$$

$$u_x = \psi_x \quad \text{for } x \geq 1, t \leq l_2(x);$$

it follows that u_x is continuous. Next,

$$(5.51) \quad u_{xx} = \phi_{xx} \quad \text{for } x < -1, t < l_1(x),$$

$$(5.52) \quad \phi_{xx} - u_{xx} = 1 + \int_{l_1(x)}^t \theta_{xx}(x, \tau) d\tau \quad \text{for } x < -1, t > l_1(x),$$

$$(5.53) \quad \phi_{xx} - u_{xx} = \int_0^t \theta_{xx}(x, \tau) d\tau - h''(x) \quad \text{for } -1 < x < 1, t > 0,$$

$$\phi_{xx} - u_{xx} = \int_{l_2(x)}^t \theta_{xx}(x, \tau) d\tau + 1 + (\phi_t - \psi_t)(x, l_2(x))$$

$$(5.54) \quad \text{for } x > 1, t > l_2(x),$$

$$(5.55) \quad u_{xx} = \psi_{xx} \quad \text{for } x > 1, t > l_2(x),$$

from which it easily follows that $u_{xx} \in L_{loc}^2$. Since $\theta \geq 0$ and $h \leq 0$, $\phi \geq \psi$, formulas (5.14)–(5.16) and (5.49), (5.50) imply that $u \leq \phi$.

Now from above we know that $w = \theta - \lambda \leq 0$; hence

$$(u - \psi_t) = \phi_t - \theta - \psi_t = \lambda - \theta \geq 0.$$

But $(u - \psi)(s_1(t), t) \geq 0$, $(u - \psi)(s_2(t), t) = 0$, $(u - \psi)(x, 0) = h - \psi(x, 0) \geq 0$ for $x \in [-1, 1]$. Therefore $u - \psi \geq 0$ for $s_1(t) \leq x \leq s_2(t)$, $t \geq 0$. From formulas (5.52), (5.53), (5.54) we get

$$(5.56) \quad u_t - u_{xx} = f \quad \text{if } s_1(t) < x < s_2(t), t > 0,$$

and

$$(5.57) \quad u_t - u_{xx} - f = \phi_t - \phi_{xx} - f = -1 \leq 0 \quad \text{if } x < -1, t < l_1(x),$$

$$u_t - u_{xx} - f = \psi_t - \psi_{xx} - f = \mu \geq 0 \quad \text{if } x > 1, t < l_2(x).$$

Hence (5.8), (5.9) are proved. Finally (5.10) follows from the definition of u and from the assumptions on h . We have thus proved

THEOREM 5.1. *Under the assumptions (5.19)–(5.21) and (5.43)–(5.46), a solution θ , s_1 , s_2 of (5.35)–(5.42) yields by formulas (5.14)–(5.16) and (5.49), (5.50) a solution of (5.6)–(5.10).*

5.3. Solution of the Stefan problem. Setting

$$v_1(t) = \dot{s}_1(t), \quad v_2(t) = -\dot{s}_2(t),$$

we can derive integral equations for v_1 , v_2 in a way similar to what we did in Chapter 4. Using the general formula (4.7) with $u = \theta$ we get, taking into account (5.37)–(5.40),

$$\begin{aligned} \theta(x, t) = & \int_{-1}^{+1} K(x, t; \xi, 0) g(\xi) d\xi - \int_0^t v_2(\tau) K(x, t; s_2(\tau), \tau) \lambda(s_2(\tau), \tau) d\tau \\ (5.58) \quad & + \int_0^t K(x, t; s_2(\tau), \tau) [\lambda_x(s_2(\tau), \tau) - v_2(\tau) \mu(s_2(\tau), \tau)] d\tau \\ & + \int_0^t K(x, t; s_1(\tau), \tau) v_1(\tau) d\tau - \int_0^t \lambda(s_2(\tau), \tau) \frac{\partial K}{\partial \xi}(x, t; s_2(\tau), \tau) d\tau. \end{aligned}$$

Differentiating in x and letting $x \rightarrow s_1(t)$, $s_2(t)$ respectively we get (cf. the derivation of (4.12))

$$\begin{aligned} -\frac{1}{2} v_1(t) = & \int_{-1}^{+1} K(s_1(t), t; \xi, 0) \dot{g}(\xi) d\xi \\ (5.59) \quad & + \int_0^t K_x(s_1(t), t; s_2(\tau), \tau) [\lambda_x(s_2(\tau), \tau) - \mu(s_2(\tau), \tau) v_2(\tau)] d\tau \\ & + \int_0^t K_x(s_1(t), t; s_1(\tau), \tau) v_1(\tau) d\tau \\ & + \int_0^t K(s_1(t), t; s_2(\tau), \tau) [-\lambda_x(s_2(\tau), \tau) v_2(\tau) + \lambda_t(s_2(\tau), \tau)] d\tau, \end{aligned}$$

$$\begin{aligned} & \frac{1}{2} [\lambda_x(s_2(t), t) - v_2(t) \mu(s_2(t), t)] \\ (5.60) \quad & = \int_{-1}^{+1} K(s_2(t), t; \xi, 0) \dot{g}(\xi) d\xi \\ & + \int_0^t K_x(s_2(t), t; s_2(\tau), \tau) [\lambda_x(s_2(\tau), \tau) - \mu(s_2(\tau), \tau) v_2(\tau)] d\tau \\ & + \int_0^t K_x(s_2(t), t; s_1(\tau), \tau) v_1(\tau) d\tau \\ & + \int_0^t K(s_2(t), t; s_2(\tau), \tau) [-\lambda_x(s_2(\tau), \tau) v_2(\tau) + \lambda_t(s_2(\tau), \tau)] d\tau. \end{aligned}$$

Since $\mu \geq \alpha > 0$, equations (5.59), (5.60) are two nonlinear integral equations of Volterra type, when one adds

$$(5.61) \quad s_1(t) = \int_0^t v_1(\tau) d\tau - 1,$$

$$(5.62) \quad s_2(t) = 1 - \int_0^t v_2(\tau) d\tau.$$

Existence uniqueness and monotonicity properties of the free boundaries can be established by the methods of §4.1. Hence

THEOREM 5.2. *Under the assumptions of Theorem 5.1, there exists a unique solution of (5.59)–(5.62). Defining θ by (5.58), the triple (θ, s_1, s_2) is a solution of the Stefan problem (5.35)–(5.42).*

REMARK 1. Theorem 4.2 can be extended to the present case. Thus, in particular, if

$$\mu \geq 1, \quad \lim_{t \rightarrow \infty} \sup_x \lambda(x, t) < 1,$$

then there is a positive constant γ such that

$$\frac{s_2(t) - s_1(t)}{\sqrt{t}} \leq \gamma \quad \text{if } t \geq 0.$$

The proof is obtained from an identity similar to (4.34).

REMARK 2. Theorems 4.3, 4.4 extend to the case where the function $g(x)$ takes also negative values. Here we need to apply the methods of §§4.3, 4.4. In case $\lambda \equiv \text{const}$, Lemma 4.4 can be applied and the extension of Theorems 4.3, 4.4 does not present any difficulties.

APPENDIX: PROOF OF LEMMA 4.4

By Green's formula

$$(A.1) \quad w(x, s) = \int_{\alpha}^{\infty} K(x, s; \xi, \sigma) k(\xi) d\xi + \int_{\sigma}^s K(x, s; \psi(\tau), \tau) v(\tau) d\tau$$

where $v(\tau) = -\partial w(\psi(\tau), \tau) / \partial \xi$. The condition $w(\psi(s), s) = 0$ can be expressed in the form

$$(A.2) \quad \int_{\sigma}^s K(\psi(s), s; \psi(\tau), \tau) v(\tau) d\tau = - \int_{\alpha}^{\infty} \frac{k(\xi)}{2\pi^{1/2}(s - \sigma)^{1/2}} \exp \left\{ - \frac{(\psi(s) - \xi)^2}{4(s - \sigma)} \right\} d\xi = \frac{f(s)}{2\pi^{1/2}},$$

i.e.,

$$(A.3) \quad \int_{\sigma}^s \frac{1}{(s-\tau)^{1/2}} \exp\left\{-\frac{b^2}{4}(s-\tau)\right\} v(\tau) d\tau = f(s).$$

Note that $f(\sigma) = 0$.

By Abel's method (see Van Moerbeke [24]) we can solve $v(\tau)$ in terms of $f(s)$ as follows: We multiply both sides of (A.3) by $(t-s)^{1/2} \exp(b^2 s/4)$ and integrate with respect to s , $\sigma < s < t$, and then differentiate both sides with respect to t . We find that

$$(A.4) \quad v(s) = \frac{1}{\pi} \int_{\sigma}^s \frac{1}{(s-\tau)^{1/2}} \exp\left(-\frac{b^2 s}{4}\right) \frac{d}{d\tau} \left\{ \exp\left(\frac{b^2 \tau}{4}\right) f(\tau) \right\} d\tau,$$

i.e.,

$$(A.5) \quad \begin{aligned} v(s) &= \frac{b^2}{4\pi} \int_{\sigma}^s \frac{f(\tau)}{(s-\tau)^{1/2}} \exp\left\{-\frac{b^2(s-\tau)}{4}\right\} d\tau \\ &+ \frac{1}{\pi} \int_{\sigma}^s \frac{f'(\tau)}{(s-\tau)^{1/2}} \exp\left\{-\frac{b^2(s-\tau)}{4}\right\} d\tau \\ &= I_1(s) + I_2(s). \end{aligned}$$

By direct computation we find that $I_1(t)$ is equal to the first term on the right-hand side of the second inequality of (4.44). Hence it remains to prove that

$$(A.6) \quad I_2(t) \leq k_0 C(\theta).$$

To prove (A.6) we compute

$$\begin{aligned} f'(\tau) &= - \int_{\alpha}^{\infty} k(\xi) \left\{ -\frac{1}{2} (\tau - \sigma)^{-3/2} + \frac{1}{4} (\tau - \sigma)^{-5/2} (\xi - \psi(\tau))^2 \right. \\ &\quad \left. + \frac{b}{2} (\tau - \sigma)^{-3/2} (\xi - \psi(\tau)) \right\} \\ &\quad \cdot \exp\left[-\frac{(\xi - \psi(\tau))^2}{4(\tau - \sigma)}\right] d\xi. \end{aligned}$$

In the last braces, the first term is negative and the other two terms are positive. Hence

$$\begin{aligned} f'(\tau) &\leq k_0 \int_{\alpha}^{\infty} \left\{ \frac{1}{4} (\tau - \sigma)^{-5/2} (\xi - \psi(\tau))^2 + \frac{b}{2} (\tau - \sigma)^{-3/2} (\xi - \psi(\tau)) \right\} \\ &\quad \cdot \exp\left[-\frac{(\xi - \psi(\tau))^2}{4(\tau - \sigma)}\right] d\xi. \end{aligned}$$

By integration by parts,

$$f'(\tau) \leq \frac{1}{2} k_0 \left\{ \frac{1}{2} (\tau - \sigma)^{-1} \int_{\alpha}^{\infty} \exp \left[-\frac{(\xi - \psi(\tau))^2}{4(\tau - \sigma)} \right] d\xi \right. \\ \left. + 2 \left[b(\tau - \sigma)^{1/2} + \frac{1}{2} (\tau - \sigma)^{-1} (\alpha - \psi(\tau)) \right] \right. \\ \left. \cdot \exp \left[-\frac{(\alpha - \psi(\tau))^2}{4(\tau - \sigma)} \right] \right\}.$$

Setting $\tau - \sigma = x$ we get

$$f'(\sigma + x) \\ \leq \frac{1}{2} k_0 \left\{ 2\pi^{1/2} x^{1/2} + 2 \left[bx^{-1/2} + \frac{1}{2} x^{-1} (a - bx) \right] \exp \left[-\frac{(a - bx)^2}{4x} \right] \right\}.$$

Substituting this into $I_2(t)$ we obtain

$$I_2(t) \leq \frac{k_0}{\pi^{1/2}} \int_0^{\theta} \frac{x^{1/2}}{(\theta - x)^{1/2}} \exp \left[-\frac{b^2(\theta - x)}{4} \right] dx \\ + \frac{k_0}{2\pi} \int_0^{\theta} \frac{a}{x(\theta - x)^{1/2}} \exp \left[-\frac{(a - bx)^2}{4x} \right] \exp \left[-\frac{b^2(\theta - x)}{4} \right] dx \\ (A.7) \quad + \frac{k_0 b}{\pi} \int_0^{\theta} \frac{1}{x^{1/2}(\theta - x)^{1/2}} \exp \left[-\frac{(a - bx)^2}{4x} \right] \exp \left[-\frac{b^2(\theta - x)}{4} \right] dx \\ \equiv J_1 + J_2 + J_3.$$

We have

$$(A.8) \quad J_1 \leq \frac{2k_0}{\pi^{1/2}}.$$

Next,

$$J_2 = \frac{k_0 a}{2\pi} \int_0^{\theta} \frac{1}{x(\theta - x)^{1/2}} \exp \left[-\frac{1}{4} \left(b^2 \theta - 2ab + \frac{a^2}{x} \right) \right] dx \\ = \frac{k_0 m}{2\pi} \int_0^{\theta} \frac{1}{x(\theta - x)^{1/2}} b \exp \left[-\frac{b^2}{4} \left(\theta - 2m + \frac{m^2}{x} \right) \right] dx,$$

where $m = a/b$. (Recall that $m < \theta$.)

Suppose first that $m \leq 3\theta/4$. Using the inequality

$$b \exp \left[-\frac{b^2}{4} \left(\theta - 2m + \frac{m^2}{x} \right) \right] \leq \frac{(2x)^{1/2}}{(m^2 + x(\theta - 2m))^{1/2}} \exp \left(-\frac{1}{2} \right),$$

we get

$$J_2 \leq \frac{k_0 m \exp[-(1/2)]}{\sqrt{2} \pi} \int_0^\theta \frac{dx}{x^{1/2}(\theta - x)^{1/2}(m^2 + x(\theta - 2m))^{1/2}}.$$

Since

$$m^2 + x(\theta - 2m) \geq \min(m^2, (\theta - m)^2) \geq \min\{m^2, (\theta/4)^2\},$$

$$(A.9) \quad J_2 \leq \frac{3}{\sqrt{2}} k_0 \exp \left(-\frac{1}{2} \right) \quad \text{if } m \leq \frac{3\theta}{4}.$$

If $m > 3\theta/4$, we first consider

$$\begin{aligned} J'_2 &\equiv \int_0^{\theta/2} \frac{1}{x(\theta - x)^{1/2}} b \exp \left[-\frac{b^2}{4} \left(\theta - 2m + \frac{m^2}{x} \right) \right] dx \\ (A.10) \quad &\leq \int_0^{\theta/2} \frac{1}{x(\theta - x)^{1/2}} b \exp \left(-\frac{b^2 m^2}{8x} \right) \exp \left[-\frac{b^2}{4\theta} (\theta - m)^2 \right] dx \\ &\leq \int_0^{\theta/2} \frac{1}{x(\theta - x)^{1/2}} b \exp \left(-\frac{b^2 \theta^2}{32x} \right) dx. \end{aligned}$$

Since

$$\begin{aligned} (A.11) \quad &b \exp \left(-\frac{b^2 \theta^2}{32x} \right) \leq \frac{4x^{1/2}}{\theta} \exp \left(-\frac{1}{2} \right), \\ J'_2 &\leq \frac{1}{(\theta/2)^{1/2}} \int_0^{\theta/2} \frac{dx}{\sqrt{x}} \cdot \frac{4}{\theta} \exp \left(-\frac{1}{2} \right) \leq \frac{8}{\theta} \exp \left(-\frac{1}{2} \right). \end{aligned}$$

Consider next

$$(A.12) \quad J''_2 \equiv \int_{\theta/2}^\theta \frac{1}{x(\theta - x)^{1/2}} b \exp \left[-\frac{b^2}{4} \left(\theta - 2m + \frac{m^2}{x} \right) \right] dx.$$

Since $m > 3\theta/4$,

$$(A.13) \quad \theta - 2m + \frac{m^2}{x} = \frac{(\theta - m)^2 + (2m - \theta)(\theta - x)}{x} > \frac{1}{2}(\theta - x).$$

Hence

$$(A.14) \quad J_2'' \leq \frac{2}{\theta} \int_{\theta/2}^{\theta} \frac{1}{(\theta-x)^{1/2}} b \exp\left[-\frac{b^2}{8}(\theta-x)\right] dx \leq \frac{2}{\theta} 2\sqrt{2\pi}.$$

From (A.10)–(A.12) and (A.14) it follows that

$$J_2 \leq 2k_0 \left[\frac{\sqrt{2}}{\sqrt{\pi}} + \frac{2}{\pi} \exp\left(-\frac{1}{2}\right) \right] \quad \text{if } m > \frac{3\theta}{4}.$$

Combining this with (A.9) we get

$$(A.15) \quad J_2 \leq 2k_0 \left[\frac{\sqrt{2}}{\sqrt{\pi}} + \frac{2}{\pi} \exp\left(-\frac{1}{2}\right) \right].$$

It remains to evaluate J_3 . Clearly,

$$J_3 = \frac{k_0}{\pi} \int_0^{\theta} \frac{1}{x^{1/2}(\theta-x)^{1/2}} b \exp\left[-\frac{b^2}{4}\left(\theta-2m+\frac{m^2}{x}\right)\right] dx.$$

Suppose first that $m \leq 3\theta/4$. Then

$$(A.16) \quad \begin{aligned} J_3 &\leq \frac{k_0}{\pi} \int_0^{\theta} \frac{1}{x^{1/2}(\theta-x)^{1/2}} b \exp\left[-\frac{b^2}{4}(\theta-m)^2\right] dx \\ &\leq \frac{k_0}{\pi} \int_0^{\theta} \frac{1}{x^{1/2}(\theta-x)^{1/2}} b \exp\left[-\frac{b^2\theta}{16}\right] dx \leq \frac{2k_0}{\sqrt{\theta}} \exp\left(-\frac{1}{2}\right). \end{aligned}$$

If $m > 3\theta/4$, we first consider

$$(A.17) \quad \begin{aligned} J_3' &\equiv \int_0^{\theta/2} \frac{1}{x^{1/2}(\theta-x)^{1/2}} b \exp\left[-\frac{b^2}{4}\left(\theta-2m+\frac{m^2}{x}\right)\right] dx \\ &\leq \int_0^{\theta/2} \frac{1}{x^{1/2}(\theta-x)^{1/2}} b \exp\left[-\frac{b^2\theta}{32x}\right] dx \\ &\leq \frac{4 \exp(-1/2)}{\theta} \int_0^{\theta/2} \frac{dx}{(\theta-x)^{1/2}} \leq \frac{8 \exp(-1/2)}{\theta^{1/2}}. \end{aligned}$$

If $m \leq 3\theta/4$ then we have, by (A.13),

$$(A.18) \quad \begin{aligned} J_3'' &\equiv \int_{\theta/2}^{\theta} \frac{1}{x^{1/2}(\theta-x)^{1/2}} b \exp\left[-\frac{b^2}{4}\left(\theta-2m+\frac{m^2}{4}\right)\right] dx \\ &\leq \left(\frac{2}{\theta}\right)^{1/2} \int_{\theta/2}^{\theta} \frac{1}{(\theta-x)^{1/2}} b \exp\left[-\frac{b^2}{8}(\theta-x)\right] dx \\ &\leq \frac{2\sqrt{2\pi}\sqrt{2}}{\theta^{1/2}} = \frac{4\sqrt{\pi}}{\theta^{1/2}}. \end{aligned}$$

Combining the estimates (A.16), (A.17) we find that

$$J_3 \leq \frac{4k_0}{\pi^{1/2}\theta^{1/2}} \left[1 + \frac{2}{\pi^{1/2}} \exp\left(-\frac{1}{2}\right) \right] \quad \text{if } m > \frac{3\theta}{4}.$$

In view of (A.16), the last inequality is satisfied also in case $m \leq 3\theta/4$. Combining this inequality with (A.15), (A.8), and using (A.7), the assertion (A.6) follows.

ADDED IN PROOF. (1) It was pointed out by Robert Jensen that the estimate (4.42) can be derived without recourse to Lemma 4.4. Instead of working with the function w introduced in Lemma 4.4, we work with

$$w(\xi, \tau) = k_0 \{ \exp[-b(\xi - \alpha + a) + b^2(\tau - \sigma)] - 1 \}.$$

(2) In a recent paper *Smoothness of the free boundary in the Stefan problem with supercooled water*, Robert Jensen proved the existence of a global solution for the Stefan problem corresponding to the variational inequality

$$u \geq 0, \quad (u_t - u_{xx})(v - u) \geq -(v - u) \text{ for every } v \geq 0, \\ u(x, 0) = h(x)$$

provided $h'(x)$ changes sign just once. His method can be applied to extend Theorem 5.2 (beyond the last remark of §5) to the case when no restrictions are made on the number of sign changes of the function $g(x)$ (defined in (5.43)).

REFERENCES

1. A. Bensoussan and A. Friedman, *Nonlinear variational inequalities and differential games with stopping times*, J. Functional Analysis **16** (1974), 305–352. MR **50** #6531.
2. A. Bensoussan, M. Goursat and J. L. Lions, *Contrôle impulsif et inéquations quasi-variationnelles stationnaires*, C. R. Acad. Sci. Paris Sér. A-B **276** (1973), A1279–A1284. MR **47** #5691.
3. A. Bensoussan and J. L. Lions, *Problèmes de temps d'arrêt optimal et inéquations variationnelles paraboliques*, Applicable Anal. **3** (1973), 267–295.
4. ———, *Nouvelles méthodes en contrôle impulsif*, J. Appl. Math. and Optimization **1** (1975), 289–312.
5. ———, *Nouvelles formulations de problèmes de contrôle impulsif et applications*, C. R. Acad. Sci. Paris Sér. A-B **276** (1973), A1189–A1192. MR **47** #5690.
6. ———, *Contrôle impulsif et inéquations quasi-variationnelles d'évolution*, C. R. Acad. Sci. Paris Sér. A-B **276** (1973), A1333–A1338. MR **47** #5692.
7. ———, *Contrôle impulsif et systèmes d'inéquations quasi-variationnelles*, C. R. Acad. Sci. Paris Sér. A **278** (1974), 747–751. MR **49** #5996.
8. ———, *Inéquations quasi-variationnelles décroissantes*, Congrès d'Analyse Convexe (St. Pierre de Chartreuse, 1974), Lecture Notes in Math., Springer-Verlag, Berlin and New York (to appear).
9. H. Brezis, *Problèmes unilatéraux*, J. Math. Pures Appl. **51** (1972), 1–168.
10. H. Brezis and A. Friedman, *Estimates on the support of solutions of parabolic variational inequalities*, Illinois J. Math. **20** (1976), 82–97.
11. G. Duvaut, *Résolution d'un problème de Stefan (fusion d'un bloc de glace à zéro degré)*, C. R. Acad. Sci. Paris Sér. A-B. **276** (1973), A1461–A1463. MR **48** #6688.

12. A. Friedman, *Free boundary problems for parabolic equations. I. Melting of solids*, J. Math. and Mech. **8** (1959), 499–517. MR **26** #1626.
13. ———, *Partial differential equations of parabolic type*, Prentice-Hall, Englewood Cliffs, N. J., 1964. MR **31** #6062.
14. ———, *Stochastic games and variational inequalities*, Arch. Rational Mech. Anal. **51** (1973), 321–346. MR **50** #4059.
15. ———, *Regularity theorem for variational inequalities in unbounded domains and applications to stopping time problems*, Arch. Rational Mech. Anal. **52** (1973), 134–160. MR **50** #5596.
16. ———, *Parabolic variational inequalities in one space dimension and smoothness of the free boundary*, J. Functional Analysis **18** (1975), 151–176.
17. ———, *The shape and smoothness of the free boundary for some elliptic variational inequalities*, Indiana Univ. Math. J. **25** (1976), 103–118.
18. ———, *A problem in hydraulics with non-monotone free boundary*, Indiana Univ. Math. J. **25** (1976), 577–592.
19. A. Friedman and D. Kinderlehrer, *A one phase Stefan problem*, Indiana Univ. Math. J. **24** (1975), 1005–1035.
20. ———, *A class of parabolic quasi-variational inequalities*, J. Differential Equations **21** (1976), 395–416.
21. J.-L. Lions and G. Stampacchia, *Variational inequalities*, Comm. Pure Appl. Math. **20** (1967), 493–519. MR **35** #7178.
22. V. A. Solonnikov, *A priori estimates for second-order parabolic equations*, Trudy Mat. Inst. Steklov. **70** (1964), 133–212; English transl., Amer. Math. Soc. Transl. (2) **65** (1967), 51–137. MR **28** #5267.
23. L. Tartar, *Inéquations quasi variationnelles abstraites*, C. R. Acad. Sci. Paris Sér. A **278** (1974), 1193–1196. MR **49** #9703.
24. P. Van Moerbeke, *An optional stopping time problem for linear reward*, Acta Math. **132** (1974), 1–41.

DEPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE PARIS IX, 75016 PARIS, FRANCE

INSTITUT DE RECHERCHE D'INFORMATIQUE ET D'AUTOMATIQUE (IRIA), 78150 LE CHESNAY, FRANCE

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, ILLINOIS 60201