

Samuel R. Chamberlain · Ulrich Müller ·  
Andrew D. Blackwell · Trevor W. Robbins ·  
Barbara J. Sahakian

## Noradrenergic modulation of working memory and emotional memory in humans

Received: 29 January 2006 / Accepted: 15 March 2006  
© Springer-Verlag 2006

**Abstract** *Rationale:* Noradrenaline (NA) is implicated in arousal. Working memory is dependent upon prefrontal cortex, and moderate levels of NA are thought to facilitate working memory whereas higher levels during extreme stress may impair working memory and engage more posterior cortical and sub-cortical circuitry. The NA system also influences emotional memory via modulation of the amygdalae and related mediotemporal structures. NA dysfunction and abnormalities in arousal-dependent memory functions are evident in a variety of neuropsychiatric illnesses. *Objectives:* The authors provide a concise overview of pharmacological studies that have investigated effects of selective NA manipulations on working memory and emotional memory functions in healthy human volunteers. *Materials and methods:* Selection of relevant peer-reviewed publications was based on a PubMed search. *Results:* Studies to date indicate that: (1) the beta-blocker propranolol impaired working and emotional memory, (2) clonidine frequently impaired working memory, and (3) reboxetine, a selective noradrenaline reuptake inhibitor, enhanced emotional memory for positive material. *Conclusions:* Improved understanding of coupling between NA, cortico-subcortical circuitry and human mnemonic functions will suggest

novel therapeutic directions for the treatment of neuropsychiatric conditions, such as attention deficit hyperactivity disorder and post-traumatic stress disorder. Future research directions are discussed in relation to neuroimaging techniques, functional central nervous system polymorphisms and study designs.

**Keywords** Noradrenaline · Norepinephrine · Memory · Cognition · Impulsivity · Emotion · Prefrontal

**Abbreviations** ADHD: Attention deficit hyperactivity disorder · fMRI: Functional magnetic resonance imaging · NA: Noradrenaline · PAL: Paired associates learning · PET: Positron emission tomography · PFC: Prefrontal cortex · PTSD: Post-traumatic stress disorder · SNRI: Selective noradrenaline reuptake inhibitor · SSRI: Selective serotonin reuptake inhibitor · SWM: Spatial working memory · WM: Working memory

### Introduction

Ascending noradrenergic (NA) projections from the locus coeruleus to higher cortical regions are implicated in arousal and responses to acute stress and are thought to modulate cognitive functions dependent upon prefrontal cortex and associated circuitry (Aston-Jones et al. 1999; Usher et al. 1999; Robbins 2000, 2005). Moderate levels of NA may act to enhance prefrontal cortex (PFC) control of behaviour including (short-term) working memory, but higher levels of NA during extreme stress may act as a chemical switch to take PFC off-line in favour of more posterior cortical and sub-cortical processes (Arnsten and Robbins 2002). In animal studies, moderate levels of NA enhance working memory whereas higher levels impair this function (Arnsten and Li 2005), and these findings appear attributable to dissociable effects at alpha-2a and alpha-1 adrenoceptors, respectively (Arnsten et al. 1988; Rama et al. 1996; Tanila et al. 1996; Franowicz et al. 2002; Arnsten and Li 2005). Emotional (long-term) memory, which is distinct from working memory, refers to the

S. R. Chamberlain (✉) · U. Müller (✉) · A. D. Blackwell ·  
B. J. Sahakian  
Department of Psychiatry,  
University of Cambridge School of Clinical Medicine,  
Addenbrooke's Hospital,  
Box 189, Cambridge CB2 2QQ, UK  
e-mail: src33@cam.ac.uk  
e-mail: um207@cam.ac.uk

U. Müller · T. W. Robbins  
Department of Experimental Psychology,  
University of Cambridge,  
Cambridge, UK

S. R. Chamberlain · U. Müller · A. D. Blackwell ·  
T. W. Robbins · B. J. Sahakian  
Behavioural and Clinical Neuroscience Institute,  
University of Cambridge,  
Cambridge, UK

formation, consolidation and retrieval of memories formed during times of heightened arousal or stress. Multiple studies implicate a pivotal role for the amygdalae in emotional memory, with mediotemporal structures being critical for conditioned fear learning and fearful responses in animals (LaBar and LeDoux 1996; Kilpatrick and Cahill 2003; McGaugh 2004). Augmentation of NA transmission (especially the beta-adrenergic system) in animals by systemic drug administration and local drug infusion into the amygdalae has been shown to enhance emotional memory, and such effects have been shown to be reversed by beta-blockade (McGaugh 2000, 2004; Roozendaal 2002; Phelps and LeDoux 2005).

Since Kraepelin's early pharmacological experiments in healthy volunteers using recreational and sedative drugs (Müller et al. 2006), increasingly sophisticated neuropsychological tasks and neurotransmitter-specific pharmacological agents have been developed for human use. Attempts to improve understanding of the role of NA in human working and emotional memory, and effects of NA agents, are of significant theoretical and clinical importance. NA drugs are used in the treatment of neuropsychiatric illnesses associated with PFC dysfunction and cognitive abnormalities, including attention deficit hyperactivity disorder (ADHD) (Arnsten and Li 2005) and depression (Harmer et al. 2003b; Chamberlain and Sahakian 2005). Noradrenergic (NA) transmission systems are also implicated in excessive emotional memory for traumatic events in post-traumatic stress disorder (PTSD) (Southwick et al. 1999, 2002; Nutt 2000; Giles 2005; Stone 2005). To characterise what is known of NA-mediated control of working and emotional memory in humans, we present a concise overview of available pharmacological studies conducted in healthy volunteers to date. We highlight limitations in the existing body of literature and promising new research directions.

## Materials and methods

To systematically search for studies in healthy volunteers using pharmacological manipulation with noradrenergic drugs and neuropsychological testing of memory functions, we (a) composed a list of selective noradrenergic drugs that have or can be used in humans (described in Table 1), (b) conducted a PubMed search using the string: "memory AND (noradrenaline OR atomoxetine OR dexmedetomidine OR clonidine OR epinephrine OR guanfacine OR idazoxan OR lofexidine OR norepinephrine OR prazosin OR propranolol OR reboxetine OR yohimbine)", (c) selected studies investigating emotional long-term memory or working (short-term) memory in healthy volunteers after inspection of titles and abstracts and (d) complemented the search by manually cross-checking the reference lists of selected papers.

## Results

### Effects of NA manipulations on working memory

The results from studies investigating the effects of NA drugs on working memory are presented in Table 2, where it can be seen that a variety of cognitive paradigms was employed. Examples include the spatial working memory (SWM) task from the Cambridge Neuropsychological Test Automated Battery (CANTAB), which assesses ability of volunteers to locate hidden tokens behind boxes using strategy (<http://www.camcog.com>), verbal tests of digit span and digit manipulation—see, e.g., (Müller et al. 2005a) and the Sternberg test (recognition probe for previously memorised digit or letter sequences) (Sternberg 1969).

**Table 1** Descriptions for key NA pharmacologic agents

Drug	Pharmacological profile	Approximate $t_{\max}/t_{1/2}$	Psychiatric dose range	Sample reference
Atomoxetine	Selective noradrenaline reuptake inhibitor (SNRI)	1–2/5 h	40–100 mg	(Bymaster et al. 2002)
Clonidine	Alpha2 noradrenaline receptor (adrenoceptor) agonist	3–5/6–24 h	0.1–1.8 mg	(Jakala et al. 1999a)
Dexmedetomidine	Alpha2 noradrenaline receptor (adrenoceptor) agonist	0.5 h/2 h	n/a	(Aho et al. 1993; Dutta et al. 2000)
Guanfacine	Alpha2 noradrenaline receptor (adrenoceptor) agonist (more selective than clonidine)	1–4/10–30 h	1–3 mg	(Jakala et al. 1999a)
Idazoxan	Alpha2 noradrenaline receptor (adrenoceptor) antagonist + 5-HT <sub>1A</sub> receptor agonist	1/6 h	n/a	(Middleton et al. 1999)
Lofexidine	Alpha2 noradrenaline receptor (adrenoceptor) agonist	3/11 h	0.2–2.4 mg	(Midgley et al. 1982)
Prazosin	Alpha1 noradrenaline receptor (adrenoceptor) antagonist	1.5/2 h	n/a	(Sawaki et al. 2003)
Propranolol	centrally active beta noradrenaline receptor antagonist (beta-blocker)	1–2/3–6 h	5–640 mg	(Müller et al. 2005b)
Reboxetine	Selective noradrenaline reuptake inhibitor (SNRI)	2/13 h	1–4(8) mg	(Harmer et al. 2003b)
Yohimbine	Alpha2 noradrenaline receptor (adrenoceptor) blocker	0.15/0.5 h	n/a	(Swann et al. 2005)

**Table 2** Effects of noradrenergic drugs on working memory in healthy volunteers

Author(s) and year	Drug(s) and dose(s) [mg]/design	N (females) [per drug condition]/Age [years]	Working memory task(s)	Results (and comment)
(Desai et al. 1983)	Oxprenolol 80, diazepam 5/ Parallel groups, PLC	44 (?) [5–12]/ 18–24	Running memory test	WM ↔ (after oxprenolol); WM ↓↑ (after diazepam; dependent on baseline anxiety)
(Frith et al. 1985)	Clonidine 0.2 (i.v.)/ Cross-over, PLC	8 (0) [8]/ 19–44	Digit span, PAL	WM ↓ (PAL); WM ↔ (other tasks)
(Freka and Lader 1988)	Propranolol 160, atenolol 50/ Long term (8 days)	12 (6) [12]/ 20–48	Digit symbol task, word list	WM ↓ (after propranolol)
(Coull et al. 1995a–c)	Clonidine 0.12/0.20 <sup>a</sup> , diazepam 5/10/ Mixed, PLC	88 (44) [12–16]/ ~22–25	PAL, RVIP, SWM (8), visual working memory	SWM ↓↑ (low-dose clonidine no effect; high dose improved SWM); visual WM ↓↑ (low dose impaired; high dose no effect) WM ↔ (after reboxetine)
(Kerr et al. 1996)	Reboxetine 0.5/1/4, amitriptyline 25/ Cross-over, PLC (alcohol)	10 (0) [10]/ 18–40	Sternberg	WM ↔; increase of thalamic rCBF after clonidine
(Coull et al. 1997)	Clonidine 0.1/0.12 <sup>a</sup> / Mixed, PLC, <sup>15</sup> O water PET	25 (0) [12–13]/ 18–47	PAL, RVIP	WM ↓↑ (after clonidine low and high doses impaired WM; medium dose no effect) WM ↑ (after guan-facine); small sample sizes
(Jakala et al. 1999a,b)	Clonidine 0.04/0.16/0.4 <sup>a</sup> , Guanfacine 0.56/2.32 <sup>a</sup> / Mixed, PLC	55 (15?) [6–12]/ 23–35	PAL, SWM (8)	WM ↓ (after combined idazoxan + clonidine) WM ↔ (after clonidine alone)
(Middleton et al. 1999)	Clonidine 0.12 <sup>a</sup> (i.v.), idazoxan 40, clonidine 0.12 <sup>a</sup> (i.v.) + Idazoxan 40/ Mixed, PLC	48 (23) [16]/ ~20–30	SWM (8), RVIP	
(Swartz et al. 2000)	Guanfacine/ Parallel groups, no placebo control, <sup>15</sup> O water PET	19 (7) [13]/ 20–59 (+ 24 epilepsy patients)	Delayed matching to sample	Increase of dorsal prefrontal rCBF after guanfacine; behavioural results not reported
(Rammsayer et al. 2001)	Reboxetine 2/4/ Cross-over, PLC	24 (0) [24]/ 20–38	Duration estimation	WM-related ↑ after lower dose; trend towards ↑ after higher dose
(Smith et al. 2003)	Clonidine 0.2, caffeine 120/ clonidine 0.2 + caffeine 120/ Cross-over, PLC	24 (0) [6]/ 18–35	Repeated digits, word list	WM ↓ (after clonidine, reversed by caffeine); single-blind design
(Veselis et al. 2004)	Dexmedetomidine propofol thiopental [infusion to steady state]	83 (32) [10 for dex/placebo]/ not specified	Auditory continuous recognition test	WM ↔ (all subjects after dexmedetomidine); small N
(Müller et al. 2005a)	Guanfacine 1/2/ Parallel groups, PLC	60 (0) [20]/ 20–39	SWM (12), digit span	WM ↔ (↓)
(Müller et al. 2005b)	Propranolol 25, atenolol 50/ Mixed, PLC	24 (12) [16]/ 19–27	Manipulation task	WM ↓ (after propranolol, in low-anxiety volunteers), WM ↔ (after atenolol)

**Table 2** (continued)

Author(s) and year	Drug(s) and dose(s) [mg]/design	N (females) [per drug condition]/Age [years]	Working memory task(s)	Results (and comment)
(Swann et al. 2005)	Yohimbine 20/30 or 40/ Cross-over, placebo first	9 (5) [9+8]/ ~23–37	Immediate and delayed memory task	WM ↔, more impulsive responses after yohimbine; small <i>n</i> + confounded by practice effects
(Tiplady et al. 2005)	Clonidine 0.15+0.3, temazepam 15+30/ Cross-over, PLC	15 (8) [15]/ 18–25	Sternberg, logical working memory, selective reminding	WM ↓ (after clonidine and temazepam, dose effects)

*PAL* Paired associates learning task, *PLC* placebo-controlled, *rCBF* regional cerebral blood flow, *Sternberg* Sternberg memory scanning task, *SWM* (*n*) *CANTAB* spatial working memory task (max. number of boxes), ↑/↓↔/↓↑ improved/impaired/unchanged/mixed effects (as compared to placebo)

<sup>a</sup>Individual doses, recalculated per 80 kg body weight

Propranolol impaired working memory across multiple studies (Frcka and Lader 1988; Müller et al. 2005b), but other beta-blockers including oxprenolol and atenolol generally did not (Desai et al. 1983; Frcka and Lader 1988; Müller et al. 2005b). In comparison to these other beta-blockers, propranolol is relatively lipophilic and, thus, able to penetrate the blood–brain barrier, suggesting that central blockade may be necessary for modulatory effects on working memory (though this does not rule out a possible role for peripheral system) (Müller et al. 2005b). Impairing effects of propranolol on working memory were found to be dependent upon baseline anxiety, with more significant effects in volunteers with low-state anxiety (Müller et al. 2005b). With regard to alpha-2-adrenoceptor agonists, all studies using clonidine found impairing effects on working memory at one or more doses, but evidence for dose-dependent effects was not very consistent. In one study, lower and higher doses of clonidine (0.04 and 0.4 mg) impaired working memory but an intermediate dose (0.16 mg) had no effect (Jakala et al. 1999a). In another, the deleterious effects of clonidine were more marked at a higher compared to a lower dose (0.30 vs 0.15 mg) (Tiplady et al. 2005). In another investigation, Coull et al. (1995b) found that 0.12 mg clonidine had no effect on spatial working memory but impaired visual working memory, whereas 0.20 mg improved spatial working memory with no effect on visual working memory. It is important to compare these findings to those of guanfacine, a more selective alpha-2-adrenoceptor agonist. Guanfacine improved working memory in one study (0.56- and 2.32-mg doses) (Jakala et al. 1999a). Yet, in a more recent study using similar cognitive tests (*CANTAB* *SWM*), no significant effects were noted after doses of 1 and 2 mg (and there was indeed a trend towards impairment on digit span, which was thought to be attributable to sedative effects of the drug) (Müller et al. 2005a). Another alpha-2-adrenoceptor agonist, dexmedetomidine, was found to have no significant effects on working memory overall, albeit this study used a small sample size (Veselis et al. 2004). The alpha-2-receptor blocker yohimbine appeared not to affect working memory per se but was found to increase a measure of impulsivity in that it led to increased numbers of inappropriate motor responses to non-target stimuli on a continuous performance test (Swann et al. 2005). For selective noradrenaline reuptake inhibitors (SNRIs), one study found no effects of reboxetine (0.5/1/4 mg) on a Sternberg test (Kerr et al. 1996), whereas another found that reboxetine (2 mg) improved temporal discrimination in the range of seconds, which is thought to reflect working memory and PFC function (Rammsayer et al. 2001).

In summary, propranolol (but not other beta-blockers with more limited central activity) was found to impair working memory across several studies, and clonidine impaired working memory in most studies. The single study using yohimbine found no evidence for general effects on working memory, and findings for guanfacine and SNRIs were inconsistent.

## Effects of NA manipulations on long-term emotional memory

Studies using NA drugs to manipulate emotional memory functions in healthy volunteers are summarised in Table 3. Subjects typically viewed a short slide show accompanied by a descriptive narrative, after receiving medication or placebo in a double-blind and parallel group design. Slides between conditions remained the same, and emotional arousal was manipulated by varying the narrative. For example, in the neutral condition, a typical narrative related to watching a practice disaster drill, with actors and fake badly scarred legs; in the arousal version, the narrative instead explained that this was an actual disaster, indicating that the badly scarred legs were real (van Stegeren et al. 1998). Arousing effects were assessed in terms of self-rating scales and physiological recordings. At a later date, the volunteers returned to undertake a surprise test of memory for the original story. Recall and recognition memory for the emotional and neutral segments of the original story were quantified. Problems with this paradigm include the limited number of critical items and the lack of parallel versions. As indicated in Table 3, some studies (e.g. van Stegeren et al. 2005) used other paradigms designed to overcome these limitations, including presentation of emotional and neutral pictures from the International Affective Picture System (IAPS) (Lang and Bradley 1997) or sets of emotionally valenced words.

In initial studies by Cahill et al. (1994), pre-encoding administration of propranolol (designed to maximise beta-adrenergic blockade at the time of initial viewing of slides) reduced recognition and recall for the emotional component of the story in the arousal condition after 1 week. These findings have been replicated in most subsequent studies (van Stegeren et al. 1998; Reist et al. 2001; Maheu et al. 2004; van Stegeren et al. 2005) but not all (O'Carroll et al. 1999a). It has been suggested that emotional arousal enhances memory for central story information in men and peripheral details in women, which has been attributed to differential hemispheric amygdala specialisation between the sexes (see Cahill and van Stegeren 2003, for discussion). van Stegeren et al. (2002) have also examined whether beta-blockade *after* the stage of encoding has any effect on later emotional recall. They found no evidence to support a role for NA after encoding or later at retrieval. No effects of peripheral beta-blockade or correlations between blood pressure decrease and emotional memory functions have been observed in studies with hydrophilic substances like atenolol or nadolol that do not cross the blood-brain barrier (van Stegeren et al. 1998; O'Carroll et al. 1999a). Few studies were identified that had examined effects of alpha adrenoceptor antagonists and agonists on emotional memory (Table 3). Administration of 20 mg yohimbine increased emotional memory in one study (O'Carroll et al. 1999b). Another study found no effect of (intravenous) yohimbine on emotional memory, but the dose was administered after slide presentation (Southwick et al. 2002). The single study using dexmedetomidine found no

effects on emotional memory albeit with a restricted sample size per group (Pryor et al. 2004).

Recent studies have used both positively and negatively valenced emotional material to explore the effects of NA drugs on emotional memory. Harmer et al. (2004) examined emotional memory for agreeable vs disagreeable personality characteristic words after a 7-day course of reboxetine (SNRI) (4 mg) (NB both encoding and retrieval were undertaken at end-of-study). Words were matched for frequency, meaningfulness and length. The treated group showed increased recall of positive personality characteristic words in particular compared to the placebo group, suggesting that NA manipulations can show differential effects depending upon the valence of stimuli.

Some studies were identified that had coupled NA manipulations with neuroimaging techniques. In one study using functional magnetic resonance imaging (fMRI), successful encoding of emotionally aversive nouns was found to engage the left amygdala, and this effect was abolished by administration of propranolol (Strange and Dolan 2004). Recognition of emotional noun words was found to engage the left hippocampus, but this effect was absent when beta-adrenergic blockade was used at the time of initial encoding. In another study, volunteers undertook fMRI while viewing affective pictures from the IAPS (van Stegeren et al. 2005). Viewing neutral pictures did not increase amygdala activation relative to baseline whereas viewing emotional pictures did. The increased activation after certain emotional stimuli was reduced in participants who had received 80 mg propranolol. These neuroimaging findings are consistent with an important role for NA-mediated modulation of amygdalae at the level of encoding, in particular, and possibly also during consolidation.

In summary, beta-blockade at the time of encoding of emotional stimuli with propranolol (but not other beta-blockers with more limited central actions) was found to reduce later recall (i.e. to reduce emotional memory) across the majority of studies. There was evidence from a single study that yohimbine increased emotional memory (O'Carroll et al. 1999b). Reboxetine (7-day treatment) increased emotional memory for positively valenced stimuli (Harmer et al. 2004). Neuroimaging evidence supported NA interactions with the amygdalae in emotional memory, especially at the encoding stage (Strange and Dolan 2004; van Stegeren et al. 2005).

---

## Discussion

Working memory refers to the short-term storage and manipulation of items in memory and is thought to be dependent upon the PFC (Baddeley 1986; Arnsten and Li 2005). Emotional memory refers to formation, consolidation, and long-term retrieval of memories formed during times of stress and is thought to be dependent upon the amygdalae and related mediotemporal structures (Cahill et al. 1994; van Stegeren et al. 1998, 2005; Cahill 2003; Strange and Dolan 2004). Both working and emotional memory are arousal-linked cognitive functions that are

**Table 3** Effects of noradrenergic drugs on emotional memory in healthy volunteers

Author(s) and year	Drug(s) and dose(s) [mg]/design	N (females) [per drug condition]/age [years]	Emotional memory task	Results and comment
(Cahill et al. 1994;	Propranolol 40, /	36 (19) [8–11]/	Emotional slide story	EM ↓; small sample sizes
Cahill and van Stegeren 2003)	Parallel groups, PLC	27.4±4.6		
(van Stegeren et al. 1998;	Propranolol 40, nadolol 40/	75 (52) [10–15]/	Emotional slide story	EM ↓ (after propranolol)
Cahill and van Stegeren 2003)	Parallel groups, PLC	22.6±0.8		EM ↔ (after nadolol)
(O'Carroll et al. 1999a)	Propranolol 40, nadolol 40/	36 (30) [12]/	Emotional slide story	EM ↔ (after both propranolol and nadolol)
	Parallel groups, PLC	21.4±2.5		
(O'Carroll et al. 1999b)	Yohimbine 20, metoprolol 50/	36 (18) [12]/	Emotional slide story	EM ↑ (after yohimbine), EM ↓ (after metoprolol)
	Parallel groups, PLC	~18–31		
(Reist et al. 2001)	Propranolol 40/	21 (0) [5–6]/	Emotional slide story	EM ↓, similar effect in controls and patients; small sample sizes
	Parallel groups, PLC	~35–65 (+ 17 PTSD patients)		
(Papps et al. 2002;	Reboxetine 4/8/	36 (10) [12]/	Emotional slide story	EM ↓ (dose-dependent); inverted U effect?
O'Carroll and Papps 2003)	Parallel groups, PLC	~18–25		
(Southwick et al. 2002)	Yohimbine 32* (i.v.)/	30 (9) [14–16]/	Emotional slide story	EM ↔, correlation with plasma MHPG levels; drug administration 5 min after slide presentation
	Parallel groups, PLC	32.4±10.9		EM ↔
(van Stegeren et al. 2002)	Propranolol 40/	60 (46) [15]/	Emotional slide story	EM ↔
	Parallel groups, PLC	~18–22		
(Cahill and Alkire 2003)	Epinephrine 9.6/19.2 (i.v.)/	42 (20) [?]/	Emotionally valenced slides	EM ↑ (only primary recall, slide 1–3); drug administration after slide presentation
	Parallel groups, PLC	21.9±0.7		EM ↑ (no negative bias)
(Harmer et al. 2003b)	Reboxetine 4/	24 (12) [12]/	Emotionally valenced word list	EM ↑ (after propranolol)
	Parallel groups, PLC	20–47		
(Strange et al. 2003)	Propranolol 40/	24 (12) [12]/	Emotionally valenced word list	EM ↓ (after propranolol)
	Parallel groups, PLC	19–32		
(Grillon et al. 2004)	Propranolol 40/	30 (?) [15]/	Cued feared conditioning	EM ↔, emotional arousal ↓
	Parallel groups, PLC	29±2.8		
(Harmer et al. 2004)	Reboxetine 8 (per day), citalopram 20 (per day)/	42 (21) [14]/	Emotionally valenced word list	EM ↑ (after both drugs, increased memory for positive stimuli)
	Parallel groups, PLC, long term (7 days)	25.0±4.2		EM ↓ (after high dose of propranolol, but not after low dose or metyrapone)
(Maheu et al. 2004, 2005)	Propranolol 40/80, metyrapone 2×750/	64 (0) [11–14]/	Emotional slide story	EM ↔ (dexmedetomidine) but small N for group
	Parallel groups, PLC	19–36		
(Pryor et al. 2004)	Dexmedetomidine thiopental propofol/	83 (32) [10 dex.; variable]/	Emotionally valenced slides	
	Parallel groups, PLC	18–50		

**Table 3** (continued)

Author(s) and year	Drug(s) and dose(s) [mg]/design	N (females) [per drug condition]/age [years]	Emotional memory task	Results and comment
(Strange and Dolan 2004)	Propranolol 40/ Parallel groups, PLC, fMRI	24 (12) [12]/ 20–39	Emotionally valenced word list	EM ↓ (after propranolol), reduced retrieval activation of amygdala/ hippocampus
(Schachinger et al. 2001; Moor et al. 2005)	Norepinephrine–nitroprusside sodium (i.v.), epinephrine–esmolol (i.v.) followed by placebo/ Cross-over	24 (0) [24]/ ~22–28	Emotionally valenced slides	EM ↑ (after norepinephrine); single blind design
(van Stegeren et al. 2005)	Propranolol 80/ Cross-over, PLC, fMRI	28/30 (15) [14/15]/ 18–28	Emotionally valenced slides	EM ↓, less amygdale activation

\*Approximation

important in everyday life and in understanding neuropsychiatric disorders. In this paper, we reviewed studies examining the effects of NA drugs on working and emotional memory in healthy human volunteers. The centrally and peripherally acting beta-blocker propranolol generally impaired working memory and reduced memory for emotional events/stimuli, and this was one of the most consistent findings in the literature. From a clinical perspective, these data suggest that the beta-adrenergic system represents a putative target for the treatment of PTSD (Southwick et al. 1999; Giles 2005; van Stegeren 2005). Neuroimaging evidence supports direct involvement of NA–amygdala coupling at the level of encoding, whereas the importance of this region during consolidation and retrieval remains disputed (Moor et al. 2005; van Stegeren et al. 2005). The alpha-2-adrenoceptor agonist clonidine generally impaired working memory, but findings with the more selective agent guanfacine were inconsistent, and the effects of these drugs on emotional memory could not be evaluated due to a paucity of studies. Effects of acute NA potentiation with reboxetine were inconsistent for working memory (Kerr et al. 1996; Rammsayer et al. 2001), while reboxetine was shown to increase emotional memory for positively valenced stimuli in one study (Harmer et al. 2004) and to enhance priming effects of emotional stimuli in another as a function of valence (Hurlemann et al. 2005). Thus, while NA is clearly implicated in emotional memory, differential effects have been reported as a function of stimulus valence and this remains an important area of investigation. In addition to effects on emotional long-term memory and priming, facilitating effects of reboxetine and atomoxetine on motor memory (learning of finger movement sequences) have been demonstrated that might also be arousal-dependent (Plewnia et al. 2004; Foster et al. 2006).

In reviewing the available studies, several methodological issues were identified. Many studies employed single drug doses, yet there is growing evidence that neurotransmitter systems may operate according to inverted-U functions, whereby too little or too much neurotransmitter can impair a given cognitive function in an optimal system (Cools and Robbins 2004). Therefore, it will be important in future works to clarify more precisely the effects of different drug doses. In this review, we focused on studies using selective NA agents rather than catecholaminergic drugs such as levodopa or methylphenidate that preferentially act on dopaminergic mechanisms. Nonetheless, NA drugs (Table 1) may exert secondary effects on other neurotransmitter systems, making it difficult to attribute behavioural effects explicitly to NA actions. One useful way to address this issue in future work would be to include drugs (or dietary amino acid manipulations) acting principally on other neurochemical systems as comparisons (Chamberlain et al. 2006b).

Another issue was that of possible ceiling effects on neuropsychological tasks, which can arise from reliance on high-functioning university students. Recruitment using community-orientated adverts or posters in places of work may be of greater utility. It is also possible to select

batteries of tests especially designed to avoid ceiling effects in high-functioning groups. With regard to sample sizes,  $N=15$  per group was commonplace (Tables 2 and 3, range 6–24). Restricted sample sizes limit power to detect effects of medication, making it difficult to interpret negative findings.

Many studies used a placebo-controlled double-blind approach, which is the best practice. The issue of within-subject (crossover) vs between-subject (parallel groups) pharmacological designs is difficult, with advantages and disadvantages to both. On the one hand, within-subject designs help to control for subject variability but can lead to critical problems with interpretation due to practice effects on neuropsychological tests; on the other hand, between-subject designs increase the possibility for variability between groups but eliminates the practice effect (as long as de novo volunteers with no prior exposure to the tasks are included). One method of addressing the potential issue of practice effects in within-subject designs is to train participants up to a certain level of baseline performance during a practice or screening session. However, the parallel groups design is especially recommended for studies with ‘single use’, highly strategy-dependent paradigms and designs with surprise stress or emotional challenge.

Studies frequently employed visual analogue scales to assess subjective effects, including sedative effects and emotional changes after drug administration. Inclusion of salivary cortisol measures (for example using the convenient Sarstedt Salivette TM collection system) or other measures of HPA axis function may be useful to exclude baseline group differences and to help characterise interplay between situational anxiety, stress, NA transmission and HPA axis function (Harmer et al. 2003a; Schule et al. 2004; Chamberlain and Sahakian 2005; Tse and Bond 2005). Another relevant recent development is that of salivary sampling for alpha-amylase, an indicator of sympathetic activity and/or NA activation (Rohleder et al. 2004; van Stegeren et al. 2006). Employing such a technique in future NA studies will be of great importance.

Several promising techniques are emerging in the field of neuropsychopharmacology, which will help to elucidate more precisely the role of NA in cognition and coupling with components of neural circuitry. Increasingly selective NA drugs are being developed, and it is becoming possible to use agonist/antagonist drug designs (O’Carroll et al. 1999a; Bullmore et al. 2003; Hurlmann et al. 2005). Positron emission tomography (PET) and functional magnetic resonance imaging have enabled visualisation of neural correlates of working and emotional memory processes in health and in the context of neuropsychiatric illnesses (Callicott and Weinberger 1999; Canli et al. 2000; Cahill 2003; Racine et al. 2005; van Stegeren et al. 2005). Pharmacological fMRI will facilitate detection of neuro-modulatory effects of NA agents on components of cortico-subcortical circuitry in a way that has not previously been possible (Gibbs and D’Esposito 2005a,b). Pure behavioural studies will continue to be important, as the behavioural sensitivity of cognitive tasks is frequently (and necessarily)

sacrificed when adapting tasks for the scanner (e.g. see Fletcher et al. 1998). With the advent of pharmacogenetics, future work should also investigate effects of functional polymorphisms in genes known to modulate NA actions (Diaz-Asper et al. 2006; Neumeister et al. 2006).

In all, well-designed human psychopharmacological studies are difficult to perform but are essential to our understanding of the role of NA in human cognition, especially when viewed alongside studies in experimental animals. Greater understanding of the precise control of working and emotional memory in humans has important implications in the context of everyday human cognition and in the manifestation and treatment of neuropsychiatric disorders. While NA is clearly implicated in working memory and emotional memory, examination of the role of NA on other cognitive processes, such as attention (Smith and Nutt 1996), response inhibition (Chamberlain et al. 2006a–c) and affective processing (Harmer et al. 2003b; Chamberlain and Sahakian 2004; Erickson et al. 2005), may help to clarify the interplay between these cognitive domains and memory and facilitate the development of increasingly sophisticated neurobiological models and candidate treatments for neuropsychiatric illnesses.

**Acknowledgements** This work was funded by the Wellcome Trust (Programme Grant 076274/Z/04/Z awarded to TWR, BJ Everitt, AC Roberts and BJS) and the Medical Research Council (Pathfinder Grant to UM, Priority Studentship to SRC). The Behavioural and Clinical Neuroscience Institute is funded by a joint award from the Medical Research Council and Wellcome Trust. ADB, TWR and BJS consult for Cambridge Cognition. We would like to thank the two anonymous reviewers for their very helpful suggestions and improvements on a draft version of the manuscript. Samuel R. Chamberlain and Ulrich Müller have contributed equally to this paper and are the corresponding authors.

---

## References

- Aho M, Erkola O, Kallio A, Scheinin H, Korttila K (1993) Comparison of dexmedetomidine and midazolam sedation and antagonism of dexmedetomidine with atipamezole. *J Clin Anesth* 5:194–203
- Arnsten A, Robbins TW (2002) Neurochemical modulation of prefrontal cortical function in humans and animals. In: Stuss DT, Knight R (eds) *Principles of frontal lobe function*. Oxford University Press, USA, pp 51–84
- Arnsten AF, Li BM (2005) Neurobiology of executive functions: catecholamine influences on prefrontal cortical functions. *Biol Psychiatry* 57:1377–1384
- Arnsten AF, Cai JX, Goldman-Rakic PS (1988) The alpha-2 adrenergic agonist guanfacine improves memory in aged monkeys without sedative or hypotensive side effects: evidence for alpha-2 receptor subtypes. *J Neurosci* 8:4287–4298
- Aston-Jones G, Rajkowski J, Cohen J (1999) Role of locus coeruleus in attention and behavioral flexibility. *Biol Psychiatry* 46:1309–1320
- Baddeley A (1986) *Working memory*. Oxford Science Publications, Clarendon
- Bullmore E, Suckling J, Zelaya F, Long C, Honey G, Reed L, Routledge C, Ng V, Fletcher P, Brown J, Williams SC (2003) Practice and difficulty evoke anatomically and pharmacologically dissociable brain activation dynamics. *Cereb Cortex* 13:144–154



- Bymaster FP, Katner JS, Nelson DL, Hemrick-Luecke SK, Threlkeld PG, Heiligenstein JH, Morin SM, Gehlert DR, Perry KW (2002) Atomoxetine increases extracellular levels of norepinephrine and dopamine in prefrontal cortex of rat: a potential mechanism for efficacy in attention deficit/hyperactivity disorder. *Neuropsychopharmacology* 27:699–711
- Cahill L (2003) Similar neural mechanisms for emotion-induced memory impairment and enhancement. *Proc Natl Acad Sci USA* 100:13123–13124
- Cahill L, Alkire MT (2003) Epinephrine enhancement of human memory consolidation: interaction with arousal at encoding. *Neurobiol Learn Mem* 79:194–198
- Cahill L, van Stegeren A (2003) Sex-related impairment of memory for emotional events with beta-adrenergic blockade. *Neurobiol Learn Mem* 79:81–88
- Cahill L, Prins B, Weber M, McGaugh JL (1994) Beta-adrenergic activation and memory for emotional events. *Nature* 371:702–704
- Callicott JH, Weinberger DR (1999) Neuropsychiatric dynamics: the study of mental illness using functional magnetic resonance imaging. *Eur J Radiol* 30:95–104
- Canli T, Zhao Z, Brewer J, Gabrieli JD, Cahill L (2000) Event-related activation in the human amygdala associates with later memory for individual emotional experience. *J Neurosci* 20:RC99
- Chamberlain SR, Sahakian BJ (2004) Cognition in mania and depression: psychological models and clinical implications. *Curr Psychiatry Rep* 6:451–458
- Chamberlain SR, Sahakian B (2005) Neuropsychological assessment of mood disorder. *Clin Neuropsychol Journ Treat Eval* 2
- Chamberlain SR, Blackwell AD, Fineberg N, Robbins T, Sahakian B (2006a) Motor inhibition and cognitive flexibility in OCD and trichotillomania. *Am J Psych* (In press)
- Chamberlain SR, Müller U, Blackwell AD, Clark L, Robbins TW, Sahakian BJ (2006b) Neurochemical modulation of response inhibition and probabilistic learning in humans. *Science* 311:861–863
- Chamberlain SR, Fineberg NA, Menzies LA, Blackwell AD, Bullmore ET, Robbins TW, Sahakian BJ (2006c) Impaired cognitive flexibility and motor inhibition in unaffected first-degree relatives of OCD patients: on the trail of endophenotypes. *Am J Psych* (In press)
- Cools R, Robbins TW (2004) Chemistry of the adaptive mind. *Philos Transact A Math Phys Eng Sci* 362:2871–2888
- Coull JT, Middleton HC, Robbins TW, Sahakian BJ (1995a) Clonidine and diazepam have differential effects on tests of attention and learning. *Psychopharmacology (Berl)* 120:322–332
- Coull JT, Middleton HC, Robbins TW, Sahakian BJ (1995b) Contrasting effects of clonidine and diazepam on tests of working memory and planning. *Psychopharmacology (Berl)* 120:311–321
- Coull JT, Sahakian BJ, Middleton HC, Young AH, Park SB, McShane RH, Cowen PJ, Robbins TW (1995c) Differential effects of clonidine, haloperidol, diazepam and tryptophan depletion on focused attention and attentional search. *Psychopharmacology (Berl)* 121:222–230
- Coull JT, Frith CD, Dolan RJ, Frackowiak RS, Grasby PM (1997) The neural correlates of the noradrenergic modulation of human attention, arousal and learning. *Eur J Neurosci* 9:589–598
- Desai N, Taylor-Davies A, Barnett DB (1983) The effects of diazepam and oxprenolol on short term memory in individuals of high and low state anxiety. *Br J Clin Pharmacol* 15:197–202
- Diaz-Asper CM, Weinberger DR, Goldberg TE (2006) Catechol-O-methyltransferase polymorphisms and some implications for cognitive therapeutics. *NeuroRx* 3:97–105
- Dutta S, Lal R, Karol MD, Cohen T, Ebert T (2000) Influence of cardiac output on dexmedetomidine pharmacokinetics. *J Pharm Sci* 89:519–527
- Erickson K, Drevets WC, Clark L, Cannon DM, Bain EE, Zarate CA Jr, Charney DS, Sahakian BJ (2005) Mood-congruent bias in affective go/no-go performance of unmedicated patients with major depressive disorder. *Am J Psychiatr* 162:2171–2173
- Fletcher PC, McKenna PJ, Frith CD, Grasby PM, Friston KJ, Dolan RJ (1998) Brain activations in schizophrenia during a graded memory task studied with functional neuroimaging. *Arch Gen Psychiatry* 55:1001–1008
- Foster DJ, Good DC, Fowlkes A, Sawaki L (2006) Atomoxetine enhances a short-term model of plasticity in humans. *Arch Phys Med Rehabil* 87:216–221
- Franowicz JS, Kessler LE, Borja CM, Kobilka BK, Limbird LE, Arnsten AF (2002) Mutation of the alpha2A-adrenoceptor impairs working memory performance and annuls cognitive enhancement by guanfacine. *J Neurosci* 22:8771–8777
- Frcka G, Lader M (1988) Psychotropic effects of repeated doses of enalapril, propranolol and atenolol in normal subjects. *Br J Clin Pharmacol* 25:67–73
- Frith CD, Dowdy J, Ferrier IN, Crow TJ (1985) Selective impairment of paired associate learning after administration of a centrally-acting adrenergic agonist (clonidine). *Psychopharmacology (Berl)* 87:490–493
- Gibbs SE, D'Esposito M (2005a) A functional MRI study of the effects of bromocriptine, a dopamine receptor agonist, on component processes of working memory. *Psychopharmacology (Berl)* 180:644–653
- Gibbs SE, D'Esposito M (2005b) Individual capacity differences predict working memory performance and prefrontal activity following dopamine receptor stimulation. *Cogn Affect Behav Neurosci* 5:212–221
- Giles J (2005) Beta-blockers tackle memories of horror. *Nature* 436:448–449
- Grillon C, Cordova J, Morgan CA, Charney DS, Davis M (2004) Effects of the beta-blocker propranolol on cued and contextual fear conditioning in humans. *Psychopharmacology (Berl)* 175:342–352
- Harmer CJ, Bhagwagar Z, Shelley N, Cowen PJ (2003a) Contrasting effects of citalopram and reboxetine on waking salivary cortisol. *Psychopharmacology (Berl)* 167:112–114
- Harmer CJ, Hill SA, Taylor MJ, Cowen PJ, Goodwin GM (2003b) Toward a neuropsychological theory of antidepressant drug action: increase in positive emotional bias after potentiation of norepinephrine activity. *Am J Psychiatr* 160:990–992
- Harmer CJ, Shelley NC, Cowen PJ, Goodwin GM (2004) Increased positive versus negative affective perception and memory in healthy volunteers following selective serotonin and norepinephrine reuptake inhibition. *Am J Psychiatr* 161:1256–1263
- Hurlemann R, Hawellek B, Matusch A, Kolsch H, Wollersen H, Madea B, Vogeley K, Maier W, Dolan RJ (2005) Noradrenergic modulation of emotion-induced forgetting and remembering. *J Neurosci* 25:6343–6349
- Jakala P, Riekkinen M, Sirvio J, Koivisto E, Kejonen K, Vanhanen M, Riekkinen P Jr (1999a) Guanfacine, but not clonidine, improves planning and working memory performance in humans. *Neuropsychopharmacology* 20:460–470
- Jakala P, Sirvio J, Riekkinen M, Koivisto E, Kejonen K, Vanhanen M, Riekkinen P Jr (1999b) Guanfacine and clonidine, alpha 2-agonists, improve paired associates learning, but not delayed matching to sample, in humans. *Neuropsychopharmacology* 20:119–130
- Kerr JS, Powell J, Hindmarch I (1996) The effects of reboxetine and amitriptyline, with and without alcohol on cognitive function and psychomotor performance. *Br J Clin Pharmacol* 42:239–241
- Kilpatrick L, Cahill L (2003) Amygdala modulation of parahippocampal and frontal regions during emotionally influenced memory storage. *Neuroimage* 20:2091–2099
- LaBar KS, LeDoux JE (1996) Partial disruption of fear conditioning in rats with unilateral amygdala damage: correspondence with unilateral temporal lobectomy in humans. *Behav Neurosci* 110:991–997

- Lang PJ, Bradley MM (1997) International affective picture system (IAPS) technical manual and affective ratings. NIMH center for the study of emotion and attention
- Maheu FS, Joobar R, Beaulieu S, Lupien SJ (2004) Differential effects of adrenergic and corticosteroid hormonal systems on human short- and long-term declarative memory for emotionally arousing material. *Behav Neurosci* 118:420–428
- Maheu FS, Joobar R, Lupien SJ (2005) Declarative memory after stress in humans: differential involvement of the beta-adrenergic and corticosteroid systems. *J Clin Endocrinol Metab* 90:1697–1704
- McGaugh JL (2000) Memory—a century of consolidation. *Science* 287:248–251
- McGaugh JL (2004) The amygdala modulates the consolidation of memories of emotionally arousing experiences. *Annu Rev Neurosci* 27:1–28
- Middleton HC, Sharma A, Agouzoul D, Sahakian BJ, Robbins TW (1999) Idazoxan potentiates rather than antagonizes some of the cognitive effects of clonidine. *Psychopharmacology (Berl)* 145:401–411
- Midgley I, Chasseaud LF, Taylor T, Darragh A (1982) The absorption and excretion of 14C-lofexidine hydrochloride in man. *Arzneimittelforschung* 32:972–975
- Moor T, Mundorff L, Bohringer A, Philippsen C, Langewitz W, Reino ST, Schachinger H (2005) Evidence that baroreflex feedback influences long-term incidental visual memory in men. *Neurobiol Learn Mem* 84:168–174
- Müller U, Clark L, Lam ML, Moore RM, Murphy CL, Richmond NK, Sandhu RS, Wilkins IA, Menon DK, Sahakian BJ, Robbins TW (2005a) Lack of effects of guanfacine on executive and memory functions in healthy male volunteers. *Psychopharmacology (Berl)* 1–9
- Müller U, Mottweiler E, Bublak P (2005b) Noradrenergic blockade and numeric working memory in humans. *J Psychopharmacol* 19:21–28
- Müller U, Fletcher PC, Steinberg H (2006) The origin of pharmacopsychology: Emil Kraepelin's experiments in Leipzig, Dorpat and Heidelberg (1882–1892). *Psychopharmacology (Berl)* 184:131–138
- Neumeister A, Drevets WC, Belfer I, Luckenbaugh DA, Henry S, Bonne O, Herscovitch P, Goldman D, Charney DS (2006) Effects of an alpha(2C)-adrenoreceptor gene polymorphism on neural responses to facial expressions in depression. *Neuropsychopharmacology (In press)*
- Nutt DJ (2000) The psychobiology of posttraumatic stress disorder. *J Clin Psychiatry* 61(Suppl 5):24–29 (Discussion 30–32)
- O'Carroll RE, Papps BP (2003) Decision making in humans: the effect of manipulating the central noradrenergic system. *J Neurol Neurosurg Psychiatry* 74:376–378
- O'Carroll RE, Drysdale E, Cahill L, Shajahan P, Ebmeier KP (1999a) Memory for emotional material: a comparison of central versus peripheral beta blockade. *J Psychopharmacol* 13:32–39
- O'Carroll RE, Drysdale E, Cahill L, Shajahan P, Ebmeier KP (1999b) Stimulation of the noradrenergic system enhances and blockade reduces memory for emotional material in man. *Psychol Med* 29:1083–1088
- Papps BP, Shajahan PM, Ebmeier KP, O'Carroll RE (2002) The effects of noradrenergic re-uptake inhibition on memory encoding in man. *Psychopharmacology (Berl)* 159:311–318
- Phelps EA, LeDoux JE (2005) Contributions of the amygdala to emotion processing: from animal models to human behavior. *Neuron* 48:175–187
- Plewnia C, Hoppe J, Cohen LG, Gerloff C (2004) Improved motor skill acquisition after selective stimulation of central norepinephrine. *Neurology* 62:2124–2126
- Pryor KO, Veselis RA, Reinsel RA, Feshchenko VA (2004) Enhanced visual memory effect for negative versus positive emotional content is potentiated at sub-anaesthetic concentrations of thiopental. *Br J Anaesth* 93:348–355
- Racine E, Bar-Ilan O, Illes J (2005) fMRI in the public eye. *Nat Rev Neurosci* 6:159–164
- Rama P, Linnankoski I, Tanila H, Pertovaara A, Carlson S (1996) Medetomidine, atipamezole, and guanfacine in delayed response performance of aged monkeys. *Pharmacol Biochem Behav* 55:415–422
- Rammesayer TH, Hennig J, Haag A, Lange N (2001) Effects of noradrenergic activity on temporal information processing in humans. *Q J Exp Psychol B* 54:247–258
- Reist C, Duffy JG, Fujimoto K, Cahill L (2001) beta-Adrenergic blockade and emotional memory in PTSD. *Int J Neuropsychopharmacol* 4:377–383
- Robbins TW (2000) Chemical neuromodulation of frontal-executive functions in humans and other animals. *Exp Brain Res* 133:130–138
- Robbins TW (2005) Chemistry of the mind: neurochemical modulation of prefrontal cortical function. *J Comp Neurol* 493:140–146
- Rohleder N, Nater UM, Wolf JM, Ehler U, Kirschbaum C (2004) Psychosocial stress-induced activation of salivary alpha-amylase: an indicator of sympathetic activity? *Ann N Y Acad Sci* 1032:258–263
- Roosendaal B (2002) Stress and memory: opposing effects of glucocorticoids on memory consolidation and memory retrieval. *Neurobiol Learn Mem* 78:578–595
- Sawaki L, Werhahn KJ, Barco R, Kopylev L, Cohen LG (2003) Effect of an alpha(1)-adrenergic blocker on plasticity elicited by motor training. *Exp Brain Res* 148:504–508
- Schachinger H, Weinbacher M, Kiss A, Ritz R, Langewitz W (2001) Cardiovascular indices of peripheral and central sympathetic activation. *Psychosom Med* 63:788–796
- Schule C, Baghai T, Schmidbauer S, Bidlingmaier M, Strasburger CJ, Laakmann G (2004) Reboxetine acutely stimulates cortisol, ACTH, growth hormone and prolactin secretion in healthy male subjects. *Psychoneuroendocrinology* 29:185–200
- Smith A, Nutt D (1996) Noradrenaline and attention lapses. *Nature* 380:291
- Smith A, Brice C, Nash J, Rich N, Nutt DJ (2003) Caffeine and central noradrenaline: effects on mood, cognitive performance, eye movements and cardiovascular function. *J Psychopharmacol* 17:283–292
- Southwick SM, Bremner JD, Rasmusson A, Morgan CA 3rd, Arnsten A, Charney DS (1999) Role of norepinephrine in the pathophysiology and treatment of posttraumatic stress disorder. *Biol Psychiatry* 46:1192–1204
- Southwick SM, Davis M, Horner B, Cahill L, Morgan CA 3rd, Gold PE, Bremner JD, Charney DC (2002) Relationship of enhanced norepinephrine activity during memory consolidation to enhanced long-term memory in humans. *Am J Psychiatr* 159:1420–1422
- Sternberg S (1969) Memory-scanning: mental processes revealed by reaction-time experiments. *Am Sci* 57:421–457
- Stone R (2005) Indian Ocean tsunami. In the wake: looking for keys to posttraumatic stress. *Science* 310:1605
- Strange BA, Dolan RJ (2004) Beta-adrenergic modulation of emotional memory-evoked human amygdala and hippocampal responses. *Proc Natl Acad Sci U S A* 101:11454–11458
- Strange BA, Hurlmann R, Dolan RJ (2003) An emotion-induced retrograde amnesia in humans is amygdala- and beta-adrenergic-dependent. *Proc Natl Acad Sci U S A* 100:13626–13631
- Swann AC, Birnbaum D, Jagar AA, Dougherty DM, Moeller FG (2005) Acute yohimbine increases laboratory-measured impulsivity in normal subjects. *Biol Psychiatry* 57:1209–1211
- Swartz BE, Kovalik E, Thomas K, Torgersen D, Mandelkern MA (2000) The effects of an alpha-2 adrenergic agonist, guanfacine, on rCBF in human cortex in normal controls and subjects with focal epilepsy. *Neuropsychopharmacology* 23:263–275
- Tanila H, Rama P, Carlson S (1996) The effects of prefrontal intracortical microinjections of an alpha-2 agonist, alpha-2 antagonist and lidocaine on the delayed alternation performance of aged rats. *Brain Res Bull* 40:117–119
- Tiplady B, Bowness E, Stien L, Drummond G (2005) Selective effects of clonidine and temazepam on attention and memory. *J Psychopharmacol* 19:259–265

- Tse WS, Bond AJ (2005) Sex differences in cortisol response to reboxetine. *J Psychopharmacol* 19:46–50
- Usher M, Cohen JD, Servan-Schreiber D, Rajkowski J, Aston-Jones G (1999) The role of locus coeruleus in the regulation of cognitive performance. *Science* 283:549–554
- van Stegeren A (2005) Beta-blockers in post-traumatic stress disorder: uses and controversies. *Expert Rev Neurother* 5:699–702
- van Stegeren AH, Everaerd W, Cahill L, McGaugh JL, Gooren LJ (1998) Memory for emotional events: differential effects of centrally versus peripherally acting beta-blocking agents. *Psychopharmacology (Berl)* 138:305–310
- van Stegeren AH, Everaerd W, Gooren LJ (2002) The effect of beta-adrenergic blockade after encoding on memory of an emotional event. *Psychopharmacology (Berl)* 163:202–212
- van Stegeren AH, Goekoop R, Everaerd W, Scheltens P, Barkhof F, Kuijjer JP, Rombouts SA (2005) Noradrenaline mediates amygdala activation in men and women during encoding of emotional material. *Neuroimage* 24:898–909
- van Stegeren A, Rohleder N, Everaerd W, Wolf OT (2006) Salivary alpha amylase as marker for adrenergic activity during stress: effect of betablockade. *Psychoneuroendocrinology* 31:137–141
- Veselis RA, Reinsel RA, Feshchenko VA, Johnson R Jr (2004) Information loss over time defines the memory defect of propofol: a comparative response with thiopental and dexmedetomidine. *Anesthesiology* 101:831–841