

RESEARCH

Open Access



Note on norm of an m -linear integral-type operator between weighted-type spaces

Stevo Stević^{1,2*}

*Correspondence: sstevic@ptt.rs

¹Mathematical Institute of the Serbian Academy of Sciences, Knez Mihailova 36/III, 11000 Beograd, Serbia

²Department of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan, Republic of China

Abstract

We find a necessary and sufficient condition for the boundedness of an m -linear integral-type operator between weighted-type spaces of functions, and calculate norm of the operator, complementing some results by L. Grafakos and his collaborators. We also present an inequality which explains a detail in the proof of the boundedness of the linear integral-type operator on $L^p(\mathbb{R}^n)$ space.

MSC: 47A30; 47B38

Keywords: Operator norm; m -linear integral-type operator; Weighted-type space; Locally compact group

1 Introduction

Throughout this note the set of natural numbers is denoted by \mathbb{N} , the set of reals by \mathbb{R} , the set of positive reals by \mathbb{R}_+ , the Euclidean n -dimensional space with the norm $|x| = (\sum_{j=1}^n x_j^2)^{1/2}$, $x = (x_1, \dots, x_n)$, by \mathbb{R}^n , the unit sphere in \mathbb{R}^n by \mathbb{S} , the $n - 1$ -dimensional surface measure by $d\sigma(\zeta)$, $\sigma_n = \sigma(\mathbb{S})$, the normalized surface measure $d\sigma(\zeta)/\sigma_n$ is denoted by $d\sigma_N(\zeta)$, the open unit ball in \mathbb{R}^n by \mathbb{B} , the open unit ball in \mathbb{R}^n centered at a and with radius r by $B(a, r)$, the Lebesgue volume measure on \mathbb{R}^n by $dV(x)$, $v_n = V(\mathbb{B})$, the normalized Lebesgue volume measure $dV(x)/v_n$ is denoted by $dV_N(x)$, whereas $L_\alpha^p(\Omega)$ denotes the weighted Lebesgue space on a domain $\Omega \subseteq \mathbb{R}^n$ with the weight $w(x) = |x|^\alpha$, that is,

$$L_\alpha^p(\Omega) = \left\{ f : \|f\|_{L_\alpha^p(\Omega)} := \left(\int_\Omega |f(x)|^p dV_\alpha(x) \right)^{1/p} < +\infty \right\},$$

where $1 \leq p < +\infty$, $\alpha > -n$, and $dV_\alpha(x) = |x|^\alpha dV(x)$ (for $\alpha = 0$, the space is reduced to the standard L^p space on the domain; see, e.g., [1]). If $k, l \in \mathbb{N}$ are such that $k \leq l$, then the notation $j = \overline{k, l}$ stands for the set of all $j \in \mathbb{N}$ such that $k \leq j \leq l$.

The weighted-type space $H_\alpha^\infty(\mathbb{R}^n)$, $\alpha > 0$, consists of all measurable functions f such that

$$\|f\|_{H_\alpha^\infty} := \operatorname{ess\,sup}_{x \in \mathbb{R}^n} |x|^\alpha |f(x)| < \infty.$$

The functional $\|\cdot\|_{H_\alpha^\infty}$ is a norm on the space, where, as usual, we identify functions which are dV almost everywhere equal. Weighted-type spaces on various domains frequently

© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

appear in the literature and are quite suitable for investigations (see, e.g., [2–7] and the references therein).

The following integral-type operator

$$L(f)(x) = \frac{1}{x} \int_0^x f(t) dt, \quad x \neq 0, \tag{1}$$

is a basic linear operator which has been studied on many spaces of functions. From the main result in [8] (see also [9]) we have that the operator is bounded on $L^p(\mathbb{R}_+)$ space when $p > 1$. This result was later improved in [10] by proving the following formula:

$$\|L\|_{L^p_\alpha(\mathbb{R}_+) \rightarrow L^p_\alpha(\mathbb{R}_+)} = \frac{p}{p - \alpha - 1} \tag{2}$$

for $p > \alpha + 1$ in nowadays terminology.

Although there are many linear operators whose norms can be calculated (see, e.g., [1, 4–28] and the related references therein), they are, in fact, quite rare since for many more other operators the norms can be only estimated by some quantities. Some of these operators are integral-type ones, a topic of a considerable recent interest (see, e.g., [11, 13–21, 23–25, 28–34] and the related references therein).

Operator (1) was generalized in [35] by introducing the following n -dimensional integral-type operator:

$$\mathcal{H}(f)(x) = \frac{1}{V(B(0, |x|))} \int_{B(0, |x|)} f(y) dV(y), \quad x \in \mathbb{R}^n \setminus \{0\}, \tag{3}$$

for nonnegative locally integrable functions on \mathbb{R}^n .

In [11] it was shown that the norm of the operator $\mathcal{H} : L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ can be calculated. Namely, the following formula holds:

$$\|\mathcal{H}\|_{L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)} = \frac{p}{p - 1}, \tag{4}$$

which matches the formula in (2) with $\alpha = 0$.

Note that $\mathcal{H}(f)(x\zeta) = \mathcal{H}(f)(x)$ for every $\zeta \in \mathbb{S}$, that is, the function $\mathcal{H}(f)(x)$ is radial for every f , and

$$\mathcal{H}(f)(x) = \frac{1}{V(B(0, |x|))} \int_{\mathbb{B}} f(|x|z) |x|^n dV(z) = \int_{\mathbb{B}} f(|x|z) dV_N(z),$$

where we have used the change of variables $y = |x|z$ and the fact that $V(B(0, |x|)) = |x|^n V(\mathbb{B})$.

In [13] the following m -linear extension of operator (3) was introduced:

$$\mathcal{H}^m(f_1, \dots, f_m)(x) = \frac{1}{v_{mn} |x|^{mn}} \int_{|(y_1, \dots, y_m)| < |x|} \prod_{j=1}^m f_j(y_j) dV(y_1) \cdots dV(y_m), \tag{5}$$

where $m \in \mathbb{N}$, $x \in \mathbb{R}^n \setminus \{0\}$, $y_1, \dots, y_m \in \mathbb{R}^n$, $y_j = (y_j^1, \dots, y_j^n)$, $j = \overline{1, m}$,

$$|(y_1, \dots, y_m)| = \left(\sum_{j=1}^m |y_j|^2 \right)^{1/2} = \left(\sum_{j=1}^m \sum_{l=1}^n (y_j^l)^2 \right)^{1/2},$$

$f_j, j = \overline{1, m}$, are nonnegative locally integrable functions on \mathbb{R}^n , and the norm of the m -linear operator was calculated from the product of weighted Lebesgue spaces $\prod_{j=1}^m L_{\alpha_j p_j/p}^{p_j}(\mathbb{R}^n)$ to $L_{\alpha}^p(\mathbb{R}^n)$, under some conditions posed on the parameters p, α, p_j , and $\alpha_j, j = \overline{1, m}$.

For the definition of norm of an m -linear operator and some basic examples, see, for example, [36, pp. 51–55].

Note also that $\mathcal{H}^m(f)(x\zeta) = \mathcal{H}^m(f)(x)$ for every $\zeta \in \mathbb{S}$, that is, the function $\mathcal{H}^m(f)(x)$ is radial for every f , and

$$\mathcal{H}^m(f_1, \dots, f_m)(x) = \frac{1}{v_{nm}} \int_{|(z_1, \dots, z_m)| < 1} \prod_{j=1}^m f_j(|x|z_j) dV(z_1) \cdots dV(z_m), \tag{6}$$

where we have used the change of variables $y_j = |x|z_j, j = \overline{1, m}$.

Our main aim here is to complement the results in [13] by calculating the norm of the operator $\mathcal{H}^m : \prod_{j=1}^m H_{\alpha_j}^{\infty}(\mathbb{R}^n) \rightarrow H_{\alpha}^{\infty}(\mathbb{R}^n)$ in the case $\alpha = \sum_{j=1}^m \alpha_j$. We also explain a detail appearing in the proof of the main result in [11].

The following known formula, which transforms integrals in the Descartes coordinates to the polar ones in \mathbb{R}^n , will be frequently used in the section that follows:

$$\int_{\mathbb{R}^n} f(x) dV(x) = \int_0^{\infty} \int_{\mathbb{S}} f(\rho\xi) d\sigma(\xi) \rho^{n-1} d\rho, \tag{7}$$

where f is a nonnegative measurable function (see, e.g., [1, pp. 149–150]).

2 Main results

This section presents and proves our main results in this note.

2.1 Boundedness of operator (5) between weighted-type spaces

First we consider operator (3). We consider it separately since its proof explains the first main step in the proof of the general case and does not request complex calculation.

Theorem 1 *Let $\alpha \in (0, n)$. Then the operator \mathcal{H} is bounded on $H_{\alpha}^{\infty}(\mathbb{R}^n)$. Moreover, the following formula holds:*

$$\|\mathcal{H}\|_{H_{\alpha}^{\infty}(\mathbb{R}^n) \rightarrow H_{\alpha}^{\infty}(\mathbb{R}^n)} = \frac{n}{n - \alpha}. \tag{8}$$

Proof Let

$$f_{\alpha}(x) = \begin{cases} 1/|x|^{\alpha}, & x \neq 0, \\ 0, & x = 0. \end{cases} \tag{9}$$

Then it is clear that

$$\|f_{\alpha}\|_{H_{\alpha}^{\infty}(\mathbb{R}^n)} = 1. \tag{10}$$

Further, we have

$$\begin{aligned} |x|^\alpha \left| \int_{\mathbb{B}} f_\alpha(|x|y) dV_N(y) \right| &= |x|^\alpha \left| \int_{\mathbb{B}} \frac{dV_N(y)}{(|x||y|)^\alpha} \right| \\ &= n \int_{\mathbb{S}} d\sigma_N(\zeta) \int_0^1 \rho^{n-1-\alpha} d\rho = \frac{n}{n-\alpha} \end{aligned}$$

for each $x \neq 0$, from which it follows that

$$\| \mathcal{H}(f_\alpha) \|_{H_\alpha^\infty(\mathbb{R}^n)} = \operatorname{ess\,sup}_{x \in \mathbb{R}^n} |x|^\alpha \left| \int_{\mathbb{B}} f_\alpha(|x|y) dV_N(y) \right| = \frac{n}{n-\alpha}. \tag{11}$$

Equalities (10) and (11) imply

$$\| \mathcal{H} \|_{H_\alpha^\infty(\mathbb{R}^n) \rightarrow H_\alpha^\infty(\mathbb{R}^n)} \geq \frac{n}{n-\alpha}. \tag{12}$$

On the other hand, we have

$$\begin{aligned} \| \mathcal{H}(f) \|_{H_\alpha^\infty(\mathbb{R}^n)} &= \operatorname{ess\,sup}_{x \in \mathbb{R}^n} |x|^\alpha \left| \int_{\mathbb{B}} f(|x|y) dV_N(y) \right| \\ &\leq \|f\|_{H_\alpha^\infty(\mathbb{R}^n)} \sup_{x \in \mathbb{R}^n \setminus \{0\}} |x|^\alpha \left| \int_{\mathbb{B}} \frac{dV_N(y)}{(|x||y|)^\alpha} \right| \\ &= \frac{n}{n-\alpha} \|f\|_{H_\alpha^\infty(\mathbb{R}^n)} \end{aligned}$$

for every $f \in H_\alpha^\infty(\mathbb{R}^n)$, from which it follows that

$$\| \mathcal{H} \|_{H_\alpha^\infty(\mathbb{R}^n) \rightarrow H_\alpha^\infty(\mathbb{R}^n)} \leq \frac{n}{n-\alpha}, \tag{13}$$

and consequently the boundedness of the operator when $\alpha \in (0, n)$. From (12) and (13) the equality in (8) follows. \square

Remark 1 Note that (10) holds for each $\alpha > 0$. However, if $\alpha \geq n$, then by using formula (7), we have

$$|x|^\alpha \left| \int_{\mathbb{B}} f_\alpha(|x|y) dV_N(y) \right| = n \int_0^1 \rho^{n-1-\alpha} d\rho = +\infty$$

for each $x \neq 0$, from which it follows that $\mathcal{H}(f_\alpha) \notin H_\alpha^\infty(\mathbb{R}^n)$. Hence, in this case the operator is not bounded on $H_\alpha^\infty(\mathbb{R}^n)$. From this and Theorem 1 we obtain the following corollary.

Corollary 1 *Let $\alpha > 0$. Then the operator \mathcal{H} is bounded on $H_\alpha^\infty(\mathbb{R}^n)$ if and only if $\alpha < n$. Moreover, if $\alpha \in (0, n)$, then formula (8) holds.*

The following theorem deals with the boundedness of the m -linear operator defined in (5).

Theorem 2 *Let $m \in \mathbb{N}$, $\alpha_j > 0$, $j = \overline{1, m}$, and*

$$\alpha = \sum_{j=1}^m \alpha_j. \tag{14}$$

Then the operator $\mathcal{H}^m : \prod_{j=1}^m H_{\alpha_j}^\infty(\mathbb{R}^n) \rightarrow H_\alpha^\infty(\mathbb{R}^n)$ is bounded if and only if

$$\int_{|\vec{y}|<1} \prod_{j=1}^m |y_j|^{-\alpha_j} \prod_{j=1}^m dV(y_j) < \infty, \tag{15}$$

where $\vec{y} = (y_1, y_2, \dots, y_m)$.

Moreover, if condition (15) is satisfied, then the following formula for the norm of the operator holds:

$$\|\mathcal{H}^m\|_{\prod_{j=1}^m H_{\alpha_j}^\infty(\mathbb{R}^n) \rightarrow H_\alpha^\infty(\mathbb{R}^n)} = \frac{1}{v_{nm}} \int_{|\vec{y}|<1} \prod_{j=1}^m |y_j|^{-\alpha_j} \prod_{j=1}^m dV(y_j), \tag{16}$$

where $\prod_{j=1}^m dV(y_j) := dV(y_1) \cdots dV(y_m)$.

Proof Let the family of functions f_α be defined in (9). Then clearly relation (10) holds. By using the condition in (14), after some simple calculation it follows that

$$\begin{aligned} |x|^\alpha \left| \int_{|\vec{y}|<1} \prod_{j=1}^m f_{\alpha_j}(|x|y_j) \prod_{j=1}^m dV(y_j) \right| &= |x|^\alpha \left| \int_{|\vec{y}|<1} \prod_{j=1}^m (|x|y_j)^{-\alpha_j} \prod_{j=1}^m dV(y_j) \right| \\ &= \int_{|\vec{y}|<1} \prod_{j=1}^m |y_j|^{-\alpha_j} \prod_{j=1}^m dV(y_j) \end{aligned} \tag{17}$$

for each $x \neq 0$.

From (6), (17) and by using the definition of norm on the space $H_\alpha^\infty(\mathbb{R}^n)$, it follows that

$$\begin{aligned} \|\mathcal{H}(f_{\alpha_1}, \dots, f_{\alpha_m})\|_{H_\alpha^\infty(\mathbb{R}^n)} &= \operatorname{ess\,sup}_{x \in \mathbb{R}^n} \frac{|x|^\alpha}{v_{nm}} \left| \int_{|\vec{y}|<1} \prod_{j=1}^m f_{\alpha_j}(|x|y_j) \prod_{j=1}^m dV(y_j) \right| \\ &= \frac{1}{v_{nm}} \int_{|\vec{y}|<1} \prod_{j=1}^m |y_j|^{-\alpha_j} \prod_{j=1}^m dV(y_j). \end{aligned} \tag{18}$$

Equality (10) implies

$$\prod_{j=1}^m \|f_{\alpha_j}\|_{H_{\alpha_j}^\infty} = 1,$$

from which along with (18) it follows that

$$\|\mathcal{H}^m\|_{\prod_{j=1}^m H_{\alpha_j}^\infty(\mathbb{R}^n) \rightarrow H_\alpha^\infty(\mathbb{R}^n)} \geq \frac{1}{v_{nm}} \int_{|\vec{y}|<1} \prod_{j=1}^m |y_j|^{-\alpha_j} \prod_{j=1}^m dV(y_j). \tag{19}$$

On the other hand, we have

$$\begin{aligned} \|\mathcal{H}^m(f_1, \dots, f_m)\|_{H_\alpha^\infty(\mathbb{R}^n)} &= \operatorname{ess\,sup}_{x \in \mathbb{R}^n} \frac{|x|^\alpha}{v_{mn}} \left| \int_{|\bar{y}| < 1} \prod_{j=1}^m f_j(|x|y_j) \prod_{j=1}^m dV(y_j) \right| \\ &\leq \prod_{j=1}^m \|f_j\|_{H_{\alpha_j}^\infty} \sup_{x \in \mathbb{R}^n \setminus \{0\}} \frac{|x|^\alpha}{v_{mn}} \left| \int_{|\bar{y}| < 1} \prod_{j=1}^m (|x||y_j|)^{-\alpha_j} \prod_{j=1}^m dV(y_j) \right| \\ &= \prod_{j=1}^m \|f_j\|_{H_{\alpha_j}^\infty} \frac{1}{v_{mn}} \int_{|\bar{y}| < 1} \prod_{j=1}^m |y_j|^{-\alpha_j} \prod_{j=1}^m dV(y_j) \end{aligned}$$

for every $(f_1, \dots, f_m) \in \prod_{j=1}^m H_{\alpha_j}^\infty(\mathbb{R}^n)$, from which by taking the supremum over the unit balls in $H_{\alpha_j}^\infty(\mathbb{R}^n)$, $j = \overline{1, m}$, it follows that

$$\|\mathcal{H}^m\|_{\prod_{j=1}^m H_{\alpha_j}^\infty(\mathbb{R}^n) \rightarrow H_\alpha^\infty(\mathbb{R}^n)} \leq \frac{1}{v_{mn}} \int_{|\bar{y}| < 1} \prod_{j=1}^m |y_j|^{-\alpha_j} \prod_{j=1}^m dV(y_j), \tag{20}$$

and consequently the boundedness of the operator. From (19) and (20) the equality in (16) follows. □

Let

$$I_m := \frac{1}{v_{mn}} \int_{|\bar{y}| < 1} \prod_{j=1}^m |y_j|^{-\alpha_j} \prod_{j=1}^m dV(y_j). \tag{21}$$

Employing the polar coordinates $y_j = \rho_j \zeta_j$, $j = \overline{1, m}$, and Fubini’s theorem, we obtain

$$\begin{aligned} I_m &= \frac{1}{v_{mn}} \underbrace{\int_{\mathbb{S}} \dots \int_{\mathbb{S}}}_{m \text{ times}} \int_{\sum_{j=1}^m \rho_j^2 < 1, \rho_j > 0, j = \overline{1, m}} \prod_{j=1}^m \rho_j^{-\alpha_j} \prod_{j=1}^m \rho_j^{n-1} d\rho_1 \dots d\rho_m d\sigma(\zeta_1) \dots d\sigma(\zeta_m) \\ &= \frac{\sigma_n^m}{v_{mn}} \int_{\sum_{j=1}^m \rho_j^2 < 1, \rho_j > 0, j = \overline{1, m}} \prod_{j=1}^m \rho_j^{n-1-\alpha_j} d\rho_1 \dots d\rho_m. \end{aligned} \tag{22}$$

By using the m -dimensional spherical coordinates

$$\begin{aligned} \rho_1 &= r \cos \varphi_1, \\ \rho_2 &= r \sin \varphi_1 \cos \varphi_2, \\ \rho_3 &= r \sin \varphi_1 \sin \varphi_2 \cos \varphi_3, \\ &\vdots \\ \rho_{m-1} &= r \sin \varphi_1 \sin \varphi_2 \dots \sin \varphi_{m-2} \cos \varphi_{m-1}, \\ \rho_m &= r \sin \varphi_1 \sin \varphi_2 \dots \sin \varphi_{m-2} \sin \varphi_{m-1}, \end{aligned}$$

where $r \geq 0$ is the radial coordinate and $\varphi_j, j = \overline{1, m-1}$, are angular coordinates, $\varphi_j \in [0, \pi]$, $j = \overline{1, m-2}$, $\varphi_{m-1} \in [0, 2\pi)$, and the known fact that the associated Jacobian is

$$|J_m| = r^{m-1} \sin^{m-2} \varphi_1 \sin^{m-3} \varphi_2 \cdots \sin \varphi_{m-2},$$

in (22), we have

$$\begin{aligned} I_m &= \frac{\sigma_n^m}{v_{mn}} \int_0^1 r^{mn-1-\sum_{j=1}^m \alpha_j} dr \\ &\quad \times \int_0^{\pi/2} \cdots \int_0^{\pi/2} \prod_{j=1}^{m-1} (\sin \varphi_j)^{n(m-j)-1-\sum_{i=j+1}^m \alpha_i} (\cos \varphi_j)^{n-1-\alpha_j} d\varphi_1 \cdots d\varphi_{m-1} \\ &= \frac{\sigma_n^m}{v_{mn}(mn-\alpha)} \prod_{j=1}^{m-1} \int_0^1 t_j^{n(m-j)-1-\sum_{i=j+1}^m \alpha_i} (1-t_j^2)^{\frac{n-2-\alpha_j}{2}} dt_j \\ &= \frac{\sigma_n^m 2^{1-m}}{v_{mn}(mn-\alpha)} \prod_{j=1}^{m-1} \int_0^1 s_j^{\frac{n(m-j)-\sum_{i=j+1}^m \alpha_i}{2}-1} (1-s_j)^{\frac{n-\alpha_j}{2}-1} ds_j \\ &= \frac{\sigma_n^m 2^{1-m}}{v_{mn}(mn-\alpha)} \prod_{j=1}^{m-1} B\left(\frac{n(m-j)-\sum_{i=j+1}^m \alpha_i}{2}, \frac{n-\alpha_j}{2}\right), \end{aligned} \tag{23}$$

where we have also used the fact that $\varphi_j \in (0, \pi/2)$, $j = \overline{1, m-1}$, which is a consequence of integrating over a set in the first orthant, the changes of variables $t_j = \sin \varphi_j, j = \overline{1, m-1}$, and $s_j = t_j^2, j = \overline{1, m-1}$, as well as the definition of the beta function (see, e.g., [36, p. 437]).

By using the following well-known relation between Euler’s beta and gamma functions:

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$$

(see, for example, [36]), after some simple calculations, we see that the following relations hold:

$$\begin{aligned} \prod_{j=1}^{m-1} B\left(\frac{n(m-j)-\sum_{i=j+1}^m \alpha_i}{2}, \frac{n-\alpha_j}{2}\right) &= \prod_{j=1}^{m-1} \frac{\Gamma\left(\frac{n(m-j)-\sum_{i=j+1}^m \alpha_i}{2}\right)\Gamma\left(\frac{n-\alpha_j}{2}\right)}{\Gamma\left(\frac{n(m-j-1)-\sum_{i=j}^m \alpha_i}{2}\right)} \\ &= \frac{\prod_{j=1}^m \Gamma\left(\frac{n-\alpha_j}{2}\right)}{\Gamma\left(\frac{nm-\alpha}{2}\right)}. \end{aligned} \tag{24}$$

Combining relations (16), (23), and (24), it follows that the following corollary holds.

Corollary 2 *Let $m \in \mathbb{N}$, $\alpha_j > 0, j = \overline{1, m}$, and $\alpha = \sum_{j=1}^m \alpha_j$. Then the operator $\mathcal{H}^m : \prod_{j=1}^m H_{\alpha_j}^\infty(\mathbb{R}^n) \rightarrow H_\alpha^\infty(\mathbb{R}^n)$ is bounded if and only if (15) holds. Moreover, if (15) holds, then the following formulas hold:*

$$\begin{aligned} \|\mathcal{H}^m\|_{\prod_{j=1}^m H_{\alpha_j}^\infty(\mathbb{R}^n) \rightarrow H_\alpha^\infty(\mathbb{R}^n)} &= \frac{\sigma_n^m 2^{1-m}}{v_{mn}(mn-\alpha)} \frac{\prod_{j=1}^m \Gamma\left(\frac{n-\alpha_j}{2}\right)}{\Gamma\left(\frac{nm-\alpha}{2}\right)} \\ &= \frac{\sigma_n^m 2^{1-m}}{v_{mn}(mn-\alpha)} \prod_{j=1}^{m-1} B\left(\frac{n(m-j)-\sum_{i=j+1}^m \alpha_i}{2}, \frac{n-\alpha_j}{2}\right). \end{aligned}$$

Remark 2 From the above consideration and Corollary 2 we see that when $\alpha_j > 0, j = \overline{1, m}$, condition (15) holds if and only if $\alpha_j \in (0, n), j = \overline{1, m}$.

2.2 A comment on the proof of the main result in [11]

As we have already mentioned, in [11] the norm of the operator $\mathcal{H} : L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ was calculated in a nice way by proving formula (4). In the proof of the result the authors applied the convolution inequality $\|g * L\|_{L^p} \leq \|g\|_{L^p} \|L\|_{L^1}$ in the group $(\mathbb{R}_+, \frac{dt}{t})$. However, the operator appearing there is not a convolution in the standard sense. On the other hand, the used inequality really holds. Hence, it needs some explanations related to the inequality, which might be useful in other similar situations. Namely, the following theorem holds and its proof is analogous to the one for the convolution operator. For some information on locally compact groups and related topics, see, e.g., [15, 37].

Theorem 3 *Let $p \in [1, \infty], f \in L^p(G), g \in L^1(G)$, where G is a locally compact group G , and μ be a right-invariant Haar measure on G . Then*

$$(f \odot g)(x) := \int_G f(xy)g(y) d\mu(y)$$

exists μ a.e. and

$$\|f \odot g\|_{L^p} \leq \|f\|_{L^p} \|g\|_{L^1}. \tag{25}$$

Proof If $p = 1$, then the result follows from Fubini’s theorem and the right-invariance of measure μ . If $p = \infty$, then we have

$$\begin{aligned} \|f \odot g\|_{L^\infty} &= \operatorname{ess\,sup}_{x \in G} \left| \int_G f(xy)g(y) d\mu(y) \right| \leq \operatorname{ess\,sup}_{x \in G} \int_G |f(xy)||g(y)| d\mu(y) \\ &\leq \int_G \|f\|_{L^\infty} |g(y)| d\mu(y) = \|f\|_{L^\infty} \|g\|_{L^1}. \end{aligned}$$

If $p \in (1, \infty)$ and if we use the notation $p' = p/(p - 1)$, then by using the Hölder inequality, Fubini’s theorem, a change of variables, and the right-invariance of measure μ , we have

$$\begin{aligned} \|f \odot g\|_{L^p}^p &= \int_G \left| \int_G f(xy)g(y) d\mu(y) \right|^p d\mu(x) \leq \int_G \left(\int_G |f(xy)||g(y)| d\mu(y) \right)^p d\mu(x) \\ &= \int_G \left(\int_G |f(xy)||g(y)|^{1/p} |g(y)|^{1/p'} d\mu(y) \right)^p d\mu(x) \\ &\leq \int_G \int_G |f(xy)|^p |g(y)| d\mu(y) \left(\int_G |g(y)| d\mu(y) \right)^{p/p'} d\mu(x) \\ &\leq \|g\|_{L^1}^{p/p'} \int_G |g(y)| \int_G |f(xy)|^p d\mu(x) d\mu(y) \\ &= \|g\|_{L^1}^{p-1} \int_G |g(y)| \int_G |f(z)|^p d\mu(z) d\mu(y) \\ &= \|f\|_{L^p}^p \|g\|_{L^1}^p, \end{aligned}$$

from which it follows that (25) holds, and since $f \odot g \in L^p(G)$ it follows that $(f \odot g)(x)$ exists μ a.e. □

Remark 3 Inequality (25) can be also obtained by using Minkowski's inequality. Indeed, by using the inequality and the right-invariance of measure μ , we have

$$\begin{aligned} \|f \odot g\|_{L^p} &= \left(\int_G \left| \int_G f(xy)g(y) d\mu(y) \right|^p d\mu(x) \right)^{1/p} \\ &\leq \left(\int_G \left(\int_G |f(xy)||g(y)| d\mu(y) \right)^p d\mu(x) \right)^{1/p} \\ &\leq \int_G \left(\int_G |f(xy)|^p d\mu(x) \right)^{1/p} |g(y)| d\mu(y) \\ &= \int_G \left(\int_G |f(z)|^p d\mu(z) \right)^{1/p} |g(y)| d\mu(y) = \|f\|_{L^p} \|g\|_{L^1}. \end{aligned}$$

Remark 4 By using inequality (25) instead of the corresponding one for the convolution operator, the proof of the main result in [11] is clear and complete.

Acknowledgements

Not applicable.

Funding

Not applicable.

Availability of data and materials

Not applicable.

Competing interests

The author declares that he has no competing interests.

Authors' contributions

The author has contributed solely to the writing of this paper. He read and approved the manuscript.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 28 October 2020 Accepted: 16 March 2021 Published online: 25 March 2021

References

- Rudin, W.: Real and Complex Analysis, 3rd edn. Higher Mathematics Series. McGraw-Hill, New York (1976)
- Avetisyan, K.: Hardy Bloch-type spaces and lacunary series on the polydisk. *Glasg. Math. J.* **49**, 345–356 (2007)
- Bierstedt, K.D., Summers, W.H.: Biduals of weighted Banach spaces of analytic functions. *J. Aust. Math. Soc. Ser. A* **54**(1), 70–79 (1993)
- Stević, S.: Norm of weighted composition operators from Bloch space to H_{μ}^{∞} on the unit ball. *Ars Comb.* **88**, 125–127 (2008)
- Stević, S.: Norm and essential norm of composition followed by differentiation from α -Bloch spaces to H_{μ}^{∞} . *Appl. Math. Comput.* **207**, 225–229 (2009)
- Stević, S.: Norm of weighted composition operators from α -Bloch spaces to weighted-type spaces. *Appl. Math. Comput.* **215**, 818–820 (2009)
- Stević, S.: Norm of some operators from logarithmic Bloch-type spaces to weighted-type spaces. *Appl. Math. Comput.* **218**, 11163–11170 (2012)
- Hardy, G.H.: Notes on some points in the integral calculus LX: an inequality between integrals. *Messenger Math.* **54**, 150–156 (1925)
- Hardy, G.H., Littlewood, J.E., Polya, G.: Inequalities. Cambridge University Press, Cambridge (1952)
- Hardy, G.H.: Notes on some points in the integral calculus, LXI: further inequalities between integrals. *Messenger Math.* **57**, 12–16 (1927)
- Christ, M., Grafakos, L.: Best constants for two nonconvolution inequalities. *Proc. Am. Math. Soc.* **123**, 1687–1693 (1995)
- Dunford, N., Schwartz, J.T.: Linear Operators, Part I: General Theory. Interscience, New York (1958)
- Fu, Z., Grafakos, L., Lu, S., Zhao, F.: Sharp bounds for m -linear Hardy and Hilbert operators. *Houst. J. Math.* **38**(1), 225–244 (2012)
- Grafakos, L.: Best bounds for the Hilbert transform on $L^p(\mathbb{R}^1)$. *Math. Res. Lett.* **4**(4), 469–471 (1997)
- Grafakos, L.: Classical Fourier Analysis, 3rd edn. Graduate Texts in Mathematics, vol. 249. Springer, New York (2014)
- Grafakos, L., Montgomery-Smith, S.: Best constants for uncentered maximal functions. *Bull. Lond. Math. Soc.* **29**(1), 60–64 (1997)

17. Grafakos, L., Montgomery-Smith, S., Motrunich, O.: A sharp estimate for the Hardy–Littlewood maximal function. *Stud. Math.* **134**(1), 57–67 (1999)
18. Grafakos, L., Savage, T.: Best bounds for the Hilbert transform on $L^p(\mathbb{R}^1)$, a corrigendum. *Math. Res. Lett.* **22**(5), 1333–1335 (2015)
19. Li, H., Li, S.: Norm of an integral operator on some analytic function spaces on the unit disk. *J. Inequal. Appl.* **2013**, Article ID 342 (2013)
20. Pelczynski, A.: Norms of classical operators in function spaces. *Astérisque* **131**, 137–162 (1985)
21. Stević, S.: Norms of some operators from Bergman spaces to weighted and Bloch-type space. *Util. Math.* **76**, 59–64 (2008)
22. Stević, S.: Norms of some operators on the Bergman and the Hardy space in the unit polydisk and the unit ball. *Appl. Math. Comput.* **215**(6), 2199–2205 (2009)
23. Stević, S.: On a new integral-type operator from the Bloch space to Bloch-type spaces on the unit ball. *J. Math. Anal. Appl.* **354**, 426–434 (2009)
24. Stević, S.: Norm and essential norm of an integral-type operator from the Dirichlet space to the Bloch-type space on the unit ball. *Abstr. Appl. Anal.* **2010**, Article ID 134969 (2010)
25. Stević, S.: Norm of an integral-type operator from Dirichlet to Bloch space on the unit disk. *Util. Math.* **83**, 301–303 (2010)
26. Stević, S.: Norms of multiplication operators on Hardy spaces and weighted composition operators from Hardy spaces to weighted-type spaces on bounded symmetric domains. *Appl. Math. Comput.* **217**, 2870–2876 (2010)
27. Stević, S.: Norms of some operators on bounded symmetric domains. *Appl. Math. Comput.* **216**, 187–191 (2010)
28. Stević, S.: Essential norm of some extensions of the generalized composition operators between k th weighted-type spaces. *J. Inequal. Appl.* **2017**, Article ID 220 (2017)
29. Benke, G., Chang, D.C.: A note on weighted Bergman spaces and the Cesáro operator. *Nagoya Math. J.* **159**, 25–43 (2000)
30. Li, S.: On an integral-type operator from the Bloch space into the $Q_k(p, q)$ space. *Filomat* **26**(2), 331–339 (2012)
31. Pan, C.: On an integral-type operator from $Q_k(p, q)$ spaces to α -Bloch spaces. *Filomat* **25**(3), 163–173 (2011)
32. Yang, W.: On an integral-type operator between Bloch-type spaces. *Appl. Math. Comput.* **215**(3), 954–960 (2009)
33. Zhu, X.: On an integral-type operator between H^2 space and weighted Bergman spaces. *Bull. Belg. Math. Soc. Simon Stevin* **18**(1), 63–71 (2011)
34. Zhu, X.: On an integral-type operator from Privalov spaces to Bloch-type spaces. *Ann. Pol. Math.* **101**(2), 139–148 (2011)
35. Faris, W.: Weak Lebesgue spaces and quantum mechanical binding. *Duke Math. J.* **43**, 365–373 (1976)
36. Zorich, V.A.: *Mathematical Analysis II*. Springer, Berlin (2004)
37. Rudin, W.: *Fourier Analysis on Groups*. Wiley, New York (1962)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)
