

Rheumatology 2000;39:1312–1315

Novel insights into the pathogenesis of osteoporosis: the role of intrauterine programming

Osteoporosis is a skeletal disorder characterized by low bone mass and microarchitectural deterioration of bony tissue with a consequent increase in the risk of fracture. These fractures typically occur at the hip, spine and distal forearm; the annual cost attributable to them in England and Wales is £942 million, with over 90% of this figure ascribed to hip fracture [1]. The bone mass of an individual in later adult life depends upon the peak attained during skeletal growth and the subsequent rate of bone loss. Preventive strategies against osteoporosis may be aimed at either increasing the peak bone mass attained or reducing the rates of bone loss. There is evidence to suggest that peak bone mass is inherited, but current genetic markers are able to explain only a small proportion of the variation in individual bone mass or fracture risk [2]. It is likely that environmental influences during early life interact with the genome in establishing the functional level of a variety of metabolic processes involved in skeletal growth. This editorial will address the role played by such environmental influences during intrauterine or very early postnatal life. It will review the conceptual basis of the fetal origins hypothesis by considering epidemiological studies pointing to the fetal programming of later osteoporosis risk and animal evidence relating to the programming of skeletal growth and metabolism.

Epidemiological studies of coronary heart disease performed over a decade ago demonstrated strong geographical associations between death rate from the disorder in 1968–1978 and infant mortality in 1901–1910 [3]. Subsequent research, based on individuals whose birth records had been preserved for seven decades, revealed that men and women who were undernourished during intrauterine life, and therefore had low birth-weight or were thin at birth, had an increased risk for coronary heart disease, hypertension, non-insulin-dependent diabetes and hypercholesterolaemia [4]. These associations are explained by a phenomenon known as programming [5]; this term describes persisting changes in structure and function caused by environmental stimuli acting at critical periods during early development. During embryonic life, the basic form of the human baby is laid down in miniature. However, the body does not increase greatly in size until

the fetal period; a rapid growth phase then commences, and continues until after birth [6]. The main feature of fetal growth is cell division. Different tissues of the body grow during periods of rapid cell division, the so-called critical periods [7]. Their timing differs for different tissues; for example, the kidney has one in the weeks immediately before birth, while the long bones accelerate their rate of growth during the second trimester of gestation. The main adaptive response to a lack of nutrients and oxygen during this period of growth is to slow the rate of cell division, especially in tissues which are undergoing critical periods at the time. This reduction in cell division is either direct or mediated through altered concentrations of growth factors or hormones (in particular insulin, growth hormone and cortisol).

It is not in question that the human skeleton can be programmed by undernutrition. Rickets has served as a long-standing example of undernutrition at a critical stage of early life, leading to persisting changes in structure. What is new is the realization that some of the body's 'memories' of early undernutrition become translated into pathology and thereby determine disease in later life. Evidence has now accumulated that such intrauterine programming contributes to the risk of osteoporosis in later life.

Evidence that the risk of osteoporosis might be modified by environmental influences during early life stems from four groups of studies: (i) bone mineral measurements undertaken in cohorts of adults whose detailed birth and/or childhood records have been preserved; (ii) detailed physiological studies exploring the relationship between candidate endocrine systems which might be programmed [growth hormone (GH)/insulin-like growth factor (IGF)-1]; hypothalamic–pituitary–adrenal, gonadal steroid) and age-related bone loss; (iii) studies characterizing the nutrition, body build and lifestyle of pregnant women and relating these to the bone mass of their newborn offspring; and (iv) studies relating childhood growth rates to the later risk of hip fracture.

The first epidemiological evidence that osteoporosis risk might be programmed came from a study of 153 women born in Bath during 1968–69 who were traced

and studied at age 21 yr [8]. Data on childhood growth were obtained from linked birth and school health records. There were statistically significant ($P < 0.05$) associations between weight at 1 yr and bone mineral content (BMC), but not bone density, at the lumbar spine and femoral neck; these relationships were independent of adult weight and body mass index. The data suggested a discordance between the processes which govern skeletal growth and those which influence mineralization. They also provide direct evidence that the trajectory of bone growth might be programmed, an assertion previously supported only by inference from measurements of body height. The association between weight in infancy and adult bone mass was replicated in a second cohort study of 238 men and 201 women aged 60–75 yr, who were born and still lived in Hertfordshire [9]. In this study, there were highly significant relationships between weight at 1 yr and adult bone area at the spine and hip ($P < 0.005$); the relationships with BMC at these two sites were weaker but remained statistically significant ($P < 0.02$). They also remained after adjustment for known genetic markers of osteoporosis risk, such as polymorphisms in the genes for the vitamin D receptor [10] and for collagen IA1, and after adjustment for lifestyle characteristics in adulthood which might have influenced bone mass (physical activity, dietary calcium intake, cigarette smoking and alcohol consumption).

To explore further the potential role of hypothalamic–pituitary function and its relevance to the pathogenesis of osteoporosis, profiles of circulating GH and cortisol were compared with bone density among groups of men and women whose birth records had been preserved. These studies revealed that birthweight and weight in infancy were predictors of basal levels of GH and cortisol during late adult life [11–13]. The levels of these two skeletally active hormones were also found to be determinants of prospectively determined bone loss rate. The data are compatible with the hypothesis that environmental stressors during intrauterine or early postnatal life alter the sensitivity of the growth plate to GH and cortisol. The consequence of such endocrine programming would be to reduce peak skeletal size, perhaps also to reduce mineralization, and to predispose to an accelerated rate of bone loss during later life [11–13].

The third piece of epidemiological evidence that osteoporosis might be programmed stems from investigation of a series of mothers through pregnancy; anthropometric and lifestyle maternal characteristics were related to the bone mineral of their newborn offspring [14, 15]. After adjusting for sex and gestational age, neonatal bone mass was strongly, positively associated with birthweight, birth length and placental weight. Other determinants included maternal and paternal birthweight, and maternal triceps skinfold thickness at 28 weeks. Maternal smoking and maternal energy intake at 18 weeks' gestation were negatively

associated with neonatal BMC in both the spine and the whole body. The independent effects of maternal and paternal birthweight on fetal skeletal development support the notion that paternal influences, for example through the imprinting of growth-promoting genes such as IGF-2, contribute strongly to the establishment of the early skeletal growth trajectory, while maternal nutrition and body build modify fetal nutrient supply and subsequent bone accretion, predominantly through influences on placentation.

Most evidence relating the intrauterine environment to later osteoporosis stems from studies using non-invasive assessment of bone mineral. The clinically important consequence of reduced bone mass is fracture, and data are now available which directly link growth rates in childhood with the risk of later hip fracture [16]. Studies of a unique Finnish cohort in which birth and childhood growth data were linked to later hospital discharge records for hip fracture have permitted the follow-up of around 7000 men and women who were born in Helsinki University Central Hospital during 1924–1933. Body size at birth was recorded and an average of 10 measurements of height and weight were obtained throughout childhood. Hip fracture incidence was assessed in this cohort using the Finnish hospital discharge registration system. After adjustment for age and sex, there were two major determinants of hip fracture risk: tall maternal height ($P < 0.001$) and low rate of childhood growth (height, $P = 0.006$; weight, $P = 0.01$). The effects of maternal height and childhood growth rate were statistically independent of each other, and remained after adjusting for socioeconomic status. The data are compatible with endocrine programming influencing the risk of hip fracture. In addition, the observation that fracture subjects were shorter at birth but of average height by age 7 yr suggests that hip fracture risk might be particularly high among children in whom growth of the skeletal envelope is forced ahead of the capacity to mineralize, a phenomenon which is accelerated during pubertal growth.

Numerous animal experiments have shown that hormones, undernutrition and other influences that affect development during sensitive periods of early life permanently programme the structure and physiology of the body's tissues and systems. A remarkable example is the effect of temperature on the sex of reptiles [17]. If the eggs of an American alligator are incubated at 30°C, all the offspring are female. If the eggs are incubated at 33°C, all the offspring are male. At temperatures between 30 and 33°C, there are varying proportions of females and males. It is believed that the fundamental sex is female, and a transcription factor is required to divert growth along a male pathway. Instead of the transcription factor being controlled genetically by a sex chromosome, it depends on the environment, specifically temperature.

Organ systems in the body are most susceptible to programming during periods when they are growing rapidly. During the first 2 months of life, the embryonic

period, there is extensive differentiation of progenitor cells without rapid cell replication. Thereafter, in the fetal period, the highest growth rates are observed. Growth slows in late gestation and continues to slow in childhood. The high growth rate of the fetus compared with the child is mostly the result of cell replication; the proportion of cells which are dividing becomes progressively less as the fetus becomes older. Slowing of growth is a major adaptation to undernutrition. Experiments on rats, mice, sheep and pigs have demonstrated that protein or calorie restriction of the mother during pregnancy and lactation is associated with smaller offspring [17–20]. In general, the earlier in life that undernutrition occurs, the more likely it is to have permanent effects on body size [21]. Early in embryonic life, growth is regulated by the supply of nutrients and oxygen. At some point shortly after birth, growth begins to track. In humans, tracking is demonstrated by the way in which infants grow along centile curves. Once tracking is established, it is no longer possible to make animals grow faster by offering them unlimited food. The rate of growth has become set, homeostatically controlled by feedback systems. After a period of undernutrition, they will regain their expected size. This contrasts with the effects of undernutrition during intrauterine life, in which skeletal development is slowed and the peak skeletal proportions attained following the completion of linear growth are reduced.

There are three cellular mechanisms for the induction of programming. First, the nutrient environment may permanently alter gene expression; one example of this is permanent change in the activity of metabolic enzymes, such as HMG-CoA (hydroxymethylglutaryl-coenzyme A) reductase [22]. Secondly, early nutrition may permanently reduce cell numbers. The small but normally proportioned rat produced by undernutrition before weaning has been shown to have fewer cells in its organs and tissues [23]. Growth-retarded human babies have reduced numbers of cells in their pancreas, which may limit insulin secretion [24], and reduced size of their airways, which may limit respiratory function [25]. Thirdly, certain clones of cells may be altered by environmental adversity during development; for example, an altered balance of the Th-1 and Th-2 lymphocyte subtypes might predispose to atopic disease in later life [26]. Evidence from the human studies outlined above suggests that skeletal development might be programmed as a consequence of the first two of these mechanisms.

Undernutrition and other adverse influences arising in fetal life or immediately after birth have a permanent effect on body structure, physiology and metabolism. The specific effects of undernutrition depend on the time and development at which it occurs; rapidly growing fetuses and neonates are more vulnerable. Its effects include altered gene expression, reduced cell numbers, imbalance between cell types, altered organ structure, and changes in the pattern of hormonal release and

tissue sensitivity to these hormones. Evidence is now accumulating from human studies that programming of bone growth might be an important contributor to the later risk of osteoporotic fracture. Body weight in infancy is a determinant of adult BMC as well as of the basal levels of activity of the GH/IGF-1 and hypothalamic–pituitary–adrenal axes. Epidemiological studies have suggested that maternal smoking and nutrition during pregnancy influence intrauterine skeletal mineralization. Finally, childhood growth rates have been directly linked to the risk of hip fracture many decades later. Further studies of this phenomenon are required in order that effective preventive strategies against osteoporosis throughout the life course may be delineated and more effectively applied.

C. COOPER, K. WALKER-BONE, N. ARDEN
and E. DENNISON

MRC Environmental Epidemiology Unit, University of Southampton, Southampton General Hospital, Southampton SO16 6YD, UK

Correspondence to: C. Cooper.

References

1. Royal College of Physicians. Osteoporosis: clinical guidelines for prevention and treatment. London: Royal College of Physicians of London, 1999.
2. Ralston S. Do genetic markers aid in risk assessment? *Osteoporosis Int* 1998;8(Suppl. 1):S37–42.
3. Barker DJP. Fetal origins of coronary heart disease. *Br Med J* 1995;311:171–4.
4. Barker DJP. The fetal origins of adult disease. *Proc R Soc Lond B Biol Sci* 1995;262:37–43.
5. Lucas A. Programming by early nutrition in man. In: Bock GR, Whelan J, eds. *The childhood environment and adult disease*. New York: John Wiley, 1991;38–55.
6. Walpert L. *The triumph of the embryo*. Oxford: Oxford University Press, 1991.
7. Widdowson EM, McCance RA. The determinants of growth and form. *Proc R Soc Lond B Biol Sci* 1974; 185:1–17.
8. Cooper C, Cawley MID, Bhalla A, Egger P, Ring F, Morton L *et al*. Childhood growth, physical activity and peak bone mass in women. *J Bone Miner Res* 1995;10: 940–7.
9. Cooper C, Fall C, Egger P, Hobbs R, Eastell R, Barker D. Growth in infancy and bone mass in later life. *Ann Rheum Dis* 1997;56:17–21.
10. Keen R, Egger P, Fall C, Major P, Lanchbury J, Spector TD *et al*. Polymorphisms of the vitamin D receptor, infant growth and adult bone mass. *Calcif Tiss Int* 1977; 60:233–5.
11. Fall C, Hindmarsh P, Dennison E, Kellingray S, Barker D, Cooper C. Programming of growth hormone secretion and bone mineral density in elderly men: an hypothesis. *J Clin Endocrinol Metab* 1998;83:135–9.
12. Dennison E, Hindmarsh P, Fall C, Kellingray S, Barker D, Phillips D *et al*. Profiles of endogenous circulating cortisol and bone mineral density in healthy elderly men. *J Clin Endocrinol Metab* 1999;84:3058–63.
13. Phillips DIW, Barker DJP, Fall CHD *et al*. Elevated plasma cortisol concentrations: a link between low

- birthweight and the insulin resistance syndrome? *J Clin Endocrinol Metab* 1998;83:757–60.
14. Dennison E, Godfrey K, Wheeler T, Taylor S, Kellingray S, Cooper C. Determinants of neonatal bone mass [abstract]. *Bone* 1997;20:26S.
 15. Godfrey KM, Breier BH, Cooper C. Constraint of the materno-placental supply of nutrients: causes and consequences. In: O'Brien PMS, Wheeler T, Barker DJP, eds. *Fetal programming: influences on development and disease in later life*. London: Royal College of Gynaecology Press, 1999;283–98.
 16. Cooper C, Eriksson JF, Foresen T, Osmond C, Tuomilehto J, Barker DJP. Childhood growth trajectory as a strong determinant of later risk of hip fracture: a prospective study. *Calcif Tiss Int* 1999;64(Suppl. 1):S38.
 17. Barker DJP. Programming the baby. In: Barker DJP, ed. *Mothers, babies and health in later life*. London: Churchill Livingstone, 1998;13–41.
 18. McCance RA, Widdowson EM. The determinants of growth and form. *Proc R Soc Lond B Biol Sci* 1974; 185:1–17.
 19. Smart JL, Massey RF, Nash SC, Tonkiss J. Effects of early life undernutrition in artificially reared rats: subsequent body and organ growth. *Br J Nutr* 1987;58:245–55.
 20. Chow BF, Lee CJ. Effect of dietary restriction of pregnant rats on body weight gain of the offspring. *J Nutr* 1964;84:10–8.
 21. McCance RA, Widdowson EM. Nutrition and growth. *Proc R Soc Lond B Biol Sci* 1962;156:326–37.
 22. Brown SA, Rogers LK, Dunn JK, Gotto AM, Patsch W. Development of cholesterol homeostatic memory in the rat is influenced by maternal diets. *Metabolism* 1990;39: 468–73.
 23. McLeod KI, Goldrick RB, Whyte HM. The effect of maternal malnutrition on the progeny in the rat: studies on growth, body composition and organ cellularity in first and second generation progeny. *Aust J Exp Biol Med Sci* 1972;50:435–46.
 24. Snoeck A, Remacle C, Reusens B, Hoet JJ. Effect of a low protein diet during pregnancy on the fetal rat endocrine pancreas. *Biol Neonate* 1990;57:107–18.
 25. Matsui R, Thurlbeck WM, Fujita Y, Yu SY, Kida K. Connective tissue, mechanical, and morphometric changes in the lungs of weanling rats fed a low protein diet. *Pediatr Pulmonol* 1989;7:159–66.
 26. Godfrey KM, Barker DJP, Osmond C. Disproportionate fetal growth and raised IGE concentration in adult life. *Clin Exp Allergy* 1994;24:641–8.