arXiv:2208.13300v2 [nucl-th] 5 Oct 2022

Nuclear fragmentation reactions as a probe of neutron skins in nuclei

E.A. Teixeira,^{1,2} T. Aumann,^{3,4,5,*} C.A. Bertulani,^{1,3,5,†} and B.V. Carlson^{2,‡}

¹Department of Physics and Astronomy, Texas A&M University-Commerce, TX 75429-3011, USA

²Departamento de Física, Instituto Tecnológico de Aeronáutica-CTA, 12228-900, São José dos Campos, Brazil[§]

³Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

⁴GSI Helmholtzzentrum für Schwerionenforschung GmbH, D-64291 Darmstadt, Germany

⁵Helmholtz Research Academy Hesse for FAIR, D-64289 Darmstadt, Germany

We investigate the contributions of various reaction channels to the interaction, reaction, chargechanging and neutron-changing cross sections. The goal is to investigate the relation between microscopic interactions and the symmetry energy component of the equation of state (EoS) of interest for the structure of neutron stars. We have made a comparison of the neutron skins extracted from diverse experimental techniques with those obtained with Hartree-Fock-Bogoliubov calculations with 23 Skyrme and with 8 density-dependent interactions used in the relativistic mean field method. We have shown that no particular conclusion can be drawn on the best EoS in view of the wide range of uncertainty in the experimental data. We have further investigated the prospects of using neutron-changing reactions to assess the isospin dependence of the neutron-skin in neutron-rich nuclei.

INTRODUCTION

A neutron star (NS) is a very dense baryonic system, with about 20 times more neutrons than protons and a central density reaching 5 to 6 times the saturation density of the matter in nuclei on earth, $\rho_0 \simeq 0.16 \text{ fm}^{-3}$. NS are very complex systems where all four fundamental forces have an essential role in defining their structure. One cannot rule out the possibility that hyperons and quarks exist in their dense cores. Several astronomical observables such as NS masses, radii and tidal deformabilities, together with experiments with nuclei on earth, have provided an insight on the equation of state (EoS) of symmetric and asymmetric nuclear matter. The EoS is the main ingredient for the determination of the basic NS properties [1].

The EoS for asymmetric nuclear matter is a relation between the energy per nucleon \mathcal{E} , the nucleon density ρ , and the asymmetry parameter $\delta = (\rho_n - \rho_p)/\rho$, where ρ_n (ρ_p) is the neutron (proton) density. It can be written as

$$\mathcal{E}(\rho,\delta) = \mathcal{E}(\rho) + S(\rho)\delta^2 + \mathcal{O}(\delta^4), \tag{1}$$

where the first term depends only on the total nucleon density $\rho = \rho_n + \rho_p$. The second term accounts for the deviation from symmetric nuclear matter and higher order corrections on δ are included in $\mathcal{O}(\delta^4)$. The so-called symmetry energy term $S(\rho)$ of the EoS is the most uncertain one, especially at high nuclear matter densities. To gain further insight on the symmetry energy term, it is common to expand it in a Taylor series around the nuclear matter (NM) saturation density, so that

$$S(\rho) = J + \frac{L}{3}\alpha + \frac{1}{18}K_{sym}\alpha^2 + \mathcal{O}(\alpha^3), \qquad (2)$$

where $\alpha = (\rho - \rho_0)/\rho_0$ is the expansion parameter, $J = S(\rho_0)$ is the symmetry energy at saturation density ρ_0 , L

is the slope parameter and K_{sym} is the curvature parameter. Here will will only discuss the roles of J and L. The relevance of the K_{sym} term has been addressed extensively in the literature (see, e.g., Refs. [2–4]). Whereas the value of $J \simeq 30$ MeV is compatible with numerous theoretical predictions, $L = (3\rho\partial S/\partial\rho)_{\rho=\rho_0}$ is still poorly constrained [5, 6]. This poses a problem when extrapolating from the knowledge of symmetric $\rho_n \simeq \rho_p$ to asymmetric nuclear matter (ANM) with $\rho_p \simeq 0$. The pressure in homogeneous nuclear matter is extracted from the EoS by using the relation $p(\rho, \delta) = \rho^2 d\mathcal{E}(\rho, \delta)/d\rho$. For asymmetric nuclear matter at saturation density one obtains to leading order an additional pressure from the symmetry energy term, given by $p = L\rho_0/3$. Therefore, the slope parameter L needs to be well determined for a good description of a neutron star.

Theoretically, several models have been developed to obtain the EoS of nuclear matter, all of them being constrained by comparison with experiment nuclear observables, such as nuclear masses, charge radii, excitation energies of giant monopole and dipole resonances, particles produced in central nuclear collisions at intermediate energies (\sim a few 100 MeV/nucleon). Standard microscopic models like the non-relativistic Skyrme-Hartree-Fock (SHF), or the relativistic mean-field (RMF), have been used quite successfully to describe nuclear properties [7–29].

Each of these microscopic models with different interactions yields a distinct energy density functional or EoS for nuclear matter. However, a universal energy density functional able to describe most nuclear properties and leading to a consistent EoS, has not yet been found. In other words, different NM quantities predicted by these models, such as incompressibility and symmetry energy, have a broad range of values as a function of the NM density, even when many other nuclear properties are well described for a limited set of nuclei. For example, in Ref. [30] it has been shown that while the parameter J is approximately constant, around 32 MeV, for about 23 popular Skyrme interactions, the slope parameter L can vary from about 30 MeV to 130 MeV. This result has been known for decades and highlights the fact that the 10 or more parameters used to describe Skyrme interactions can lead to very different EoS, albeit being able to describe very well a limited set of nuclear properties [31].

Numerous experiments using complementary techniques have been designed to constrain the symmetry energy term of the EoS. Astronomical observations have also helped to assess this property of the EoS. Here we will discuss one class of laboratory experiments linking the symmetry energy term to the neutron skin in nuclei. Experiments assessing the size of the neutron skin in nuclei also use numerous techniques, although an accurate result is still lacking. The neutron skin in a nucleus is often defined in terms of the difference between its neutron and proton root-mean-squared (rms) radius, $\Delta r_{np} = \langle r_n^2 \rangle^{1/2} - \langle r_p^2 \rangle^{1/2}$. It has been widely advertised that neutron skins in nuclei are directly correlated with the slope parameter L of NM (see, e.g., Refs. [2, 32]). A recent experiment [33] performed at the Jefferson Laboratory, USA, has looked at the analyzing power for polarized electron scattering due to its parity violating (PV) term in the interactions with a ²⁰⁸Pb target and found an unexpectedly large neutron skin[34] value of $\Delta r_{np} = 0.283 \pm 0.071$ fm. Intriguingly, this result implies a slope parameter $L = 106 \pm 37$ MeV, larger than expected from most microscopic calculations and also from most other experimental results. For example, Ref. [35] reported a value of $\Delta r_{np} = 0.156^{+0.025}_{-0.021}$ fm. Recent astronomical observations from heavy pulsar masses, NICER telescope and LIGO/Virgo laboratory seem to clearly point to much smaller values for Δr_{np} and L.

Experimental efforts have also been made to deduce the neutron skin from fragmentation reactions of relativistic nuclei incident on light nuclear targets. Fragmentation reactions can be thought as occurring in two steps: First nucleons are stripped off the projectile and energy is deposited in the primary fragments. After that, the primary fragments decay by evaporation of nucleons, alpha particles, and even by fission, leading to secondary fragments which are actually the ones reaching the detectors. While the first step can be rather accurately determined by theory, the second one involves numerous assumptions about excitation energies, level densities and optical potentials that are not as easily handled by theory. А way to avoid most of the complications due to the calculation of the evaporation stage is the measurement of total (a) charge- or (b) neutron-changing cross sections, i.e, all the fragments with (a) charge, or (b) isotones, different than the projectile [36-41]. The idea is that in both cases the cross sections for secondary fragments is nearly the same as those for primary fragments. Assuming that the main evaporation probability is for decay by emission of neutrons, the detection of all possible (a) isotopes, or (b) isotones, yields the same cross sections as for the total primary (a) isotopes, or (b) isotones, because all decay channels have been accounted for. The method fails when a primary fragment with neutrons (protons) removed decays by proton, or alpha, emission leading to charge changing and neutron changing cross sections that are not the same for primary and secondary fragments.

With the radioactive beams now provided in major nuclear facilities and with the increasing accuracy of betterbuilt detectors, experiments with neutron-rich nuclei are now possible, increasing the constraints on the cross section measurements and their viability to study neutron skins in several nuclei, a clear advantage over fixed target experiments. Total neutron changing cross section measurements with good accuracy are now possible with inverse kinematics, allowing the study of the fragmentation of projectiles along an isotopic chain [41].

Any experimental method using a reaction to extract an observable sensitive to EoS is model dependent, since ab initio calculations are not available. The determination of this model uncertainty, i.e., the systematic uncertainties of the reaction theory have to be evaluated and should be included in the error budget of quantities extracted from the measured cross sections.

The Glauber scattering theory, based on a description of the nucleus-nucleus reaction by means of individual nucleon-nucleon scattering is believed to be a good approximation at high beam energies above few hundred MeV [42–46]. Since we discuss total reaction probabilities, multiple reactions are not important and we start with the Glauber theory in its simplest form in first order without free parameters. The ingredients are pointneutron and -proton densities (whenever possible derived from experimental data), and the NN interaction is implemented by computing the eikonal wave function for an optical potential using the densities and measured pp and pn cross sections.

In this work, we make an assessment of uncertainties related to the first-order eikonal approximation and the fact that the interaction is introduced as a free nuclear NN interaction. Deviations due to Coulomb recoil, internal Fermi motion of nucleons, as well as due to Pauli blocking are estimated. Calculations are extended up to the 3rd eikonal order.

Even more important is the assessment of deviations from the Glauber approximation due to interactions beyond individual NN scattering. These are nuclear excitations of collective modes, such as giant resonances, in the scattering process. For heavier nuclei, these processes yield sizable contributions to the cross sections. The cross sections for nuclear- and electromagneticallyinduced excitations of collective states are considered here. Such high-lying continuum excitations contribute (for heavy nuclei) to neutron emission and add thus to the total reaction cross section and to the total-neutron removal cross sections. These calculations have systematic uncertainties but can be used as a basis to estimate for the limitations of the Glauber approach. Alternatively, these cross sections can be measured separately in the experiment, as proposed in [41], to avoid this theoretical model dependence.

CROSS SECTIONS OF PRIMARY FRAGMENTS

Glauber model for fragment production

Normalizing the projectile P and target T proton (p) and neutron (n) densities to one, the abrasion-ablation model described in Refs. [42, 44–47], yields the following expression for the probability that a proton from the projectile *survives* the collision with the target (N_T, Z_T) for a nucleus-nucleus collision with an impact parameter b:

$$P_{p}(\mathbf{b}) = \int d^{2}s \, dz \, \rho_{p}^{P}(z, \mathbf{s} - \mathbf{b}) \left[1 - \sigma_{pp} \int dz' \, \rho_{p}^{T}(z', \mathbf{s}) \right]^{Z_{T}} \\ \times \left[1 - \sigma_{pn} \int dz' \, \rho_{n}^{T}(z', \mathbf{s}) \right]^{N_{T}}, \qquad (3)$$

where, σ_{pp} and σ_{pn} are the total (minus Coulomb) proton-proton and proton-neutron scattering cross sections, respectively. An analogous expression holds for the neutron survival probability by swapping the role of the neutron and proton in the expression above. Notice that the factors $1 - \sigma \int dz(\cdots)$ are small for any nuclear system, and we can safely use the approximation $1 - x \sim \exp(-x)$ in the equations above [44–46].

In the abrasion-ablation model, the cross section for the *primary production* of a fragment with N_F neutrons and Z_F protons is

$$\sigma(N_F, Z_F) = \binom{N_P}{N_F} \binom{Z_P}{Z_F} \int d^2 b \ [P_p(\mathbf{b})]^{Z_F} [P_n(\mathbf{b})]^{N_F} \times [1 - P_n(\mathbf{b})]^{N_P - N_F} [1 - P_p(\mathbf{b})]^{Z_P - Z_F} .$$
(4)

Interaction, neutron- and charge-changing cross sections

The *interaction cross section* is the sum of all channels for which at least one nucleon is removed, that is

$$\sigma_I = \left[\sum_{N_F=0}^{N_P} \sum_{Z_F=0}^{Z_P} \sigma(N_F, Z_F)\right] - \sigma(N_P, Z_P), \quad (5)$$

where the second term removes the undisturbed projectile from the sum. Using the binomial sum, one obtains

$$\sigma_I = \int d^2 b \left[1 - [P_p(\mathbf{b})]^{Z_P} [P_n(\mathbf{b})]^{N_P} \right].$$
 (6)

Thus, the interaction cross section is the integral of one minus the probability that all protons and neutrons simultaneously survive the collision.

The neutron-changing cross section $\sigma_{\Delta N}$ is the cross section to produce all fragments with the same charge as the projectile by removing at least one of its neutrons. It is obtained by replacing $Z_F = Z_P$ in Eq. (4) and summing from $N_F = 0$ up to $N_P - 1$. The sum over the binomial coefficients yields

$$\sigma_{\Delta N} = \sigma_{\text{all } n}^{Z_P \text{ survives}} = \sum_{N_F=0}^{N_P-1} \sigma(N_F, Z_P)$$
$$= \int d^2 b \left[P_p(b) \right]^{Z_P} \left\{ \left[1 - \left[P_n(b) \right]^{N_P} \right\}.$$
(7)

This means that the probability that Z_P protons survive while all possible numbers of neutrons are removed is equal to the probability that all protons survive (irrespective to what happens to any neutron) minus the probability that all protons and neutrons survive, simultaneously (i.e, that the projectile remains intact). The charge-changing cross section $\sigma_{\Delta Z}$ is obtained by adding all fragments in which at least one proton is removed. This means that $\sigma_{\Delta Z} = \sigma_I - \sigma_{\Delta N}$.

The optical potential, eikonal phase and S-matrices

The eikonal S-matrices for a projectile proton scattering off a target nucleus is given by $S_p(\mathbf{b}) = \exp[i\chi_p(\mathbf{b})]$, where [46] $\chi_p(\mathbf{b}) = -(\hbar v)^{-1} \int dz \ U_p(\mathbf{r})$ is the eikonal phase, v is the projectile velocity, assumed undisturbed in high energy collisions, and U_p is the optical potential for proton scattering.

For nucleon removal reactions, only the imaginary part of the optical potential is of relevance and it can be related to the nucleon-nucleon cross section and nuclear densities as [46]

$$U_{p}(\mathbf{r}) = -i\frac{\hbar v}{2} \int d^{2}s \ \rho_{p}^{P} \left(z, \mathbf{s} - \mathbf{b}\right)$$
$$\times \int dz' \left[Z_{T} \sigma_{pp} \ \rho_{p}^{T} \left(z', \mathbf{s}\right) + N_{T} \sigma_{np} \ \rho_{n}^{T} \left(z', \mathbf{s}\right) \right]. (8)$$

The eikonal phase-shift is therefore

$$\chi_{p}(\mathbf{b}) = i\frac{1}{2} \int d^{2}s dz \ \rho_{p}^{P}(z, \mathbf{s} - \mathbf{b})$$
$$\times \int dz' \left[Z_{T} \sigma_{pp} \ \rho_{p}^{T}(z', \mathbf{s}) + N_{T} \sigma_{np} \ \rho_{n}^{T}(z', \mathbf{s}) \right], \quad (9)$$

and the S-matrix is

$$S_{p}(\mathbf{b}) = \exp\left\{-\frac{1}{2}\int d^{2}sdz \ \rho_{p}^{P}\left(z,\mathbf{s}-\mathbf{b}\right) \\ \times \int dz' \left[Z_{T}\sigma_{pp} \ \rho_{p}^{T}\left(z',\mathbf{s}\right) + N_{T}\sigma_{np} \ \rho_{n}^{T}\left(z',\mathbf{s}\right)\right]\right\}. (10)$$

The probability in Eq. (3) is given by $P_p(\mathbf{b}) = |S_p(\mathbf{b})|^2$. Analogous expressions are obtained for neutron removal with the roles of neutrons and protons inverted. Notice that Eq. (10) can be deduced on probability grounds without using the optical potential, Eq. (8). But later we will need the concept of an optical potential to study the higher-order corrections in the eikonal wavefunctions.

The optical limit (OL) of the Glauber cross sections is obtained when the second term inside brackets in Eq. (3) is very small so that $1 - x \sim \exp(-x)$. Then it is straightforward to show that $\sigma_I \to \sigma_{OL}$ where

$$\sigma_{OL} = \int d^2 b \left[1 - \exp\left\{ -Z_P \int d^2 s dz \; \rho_p^P \left(z, \mathbf{s} - \mathbf{b}\right) \right. \\ \left. \times \int dz' \left[Z_T \sigma_{pp} \; \rho_p^T \left(z', \mathbf{s}\right) + N_T \sigma_{np} \; \rho_n^T \left(z', \mathbf{s}\right) \right] \right. \\ \left. - N_P \int d^2 s dz \; \rho_n^P \left(z, \mathbf{s} - \mathbf{b}\right) \right. \\ \left. \times \int dz' \left[Z_T \sigma_{np} \; \rho_p^T \left(z', \mathbf{s}\right) + N_T \sigma_{pp} \; \rho_n^T \left(z', \mathbf{s}\right) \right] \right\} \right]. (11)$$

The equations developed above have been used extensively in the literature to describe interaction, reaction, charge- and neutron-changing cross sections. But, in order to achieve the accuracy needed to extract values of neutron skins and constraints on the EoS of nuclear matter, corrections to these expressions must be considered. In the next sections we study most of the relevant corrections known and investigate the sensitivity of the cross sections on each correction.

CORRECTIONS OF THE FRAGMENTATION CROSS SECTIONS

Medium corrections of NN cross sections

Here we discuss the nucleon-nucleon cross sections used as input in Eqs. (3-11). To account for medium effects, we use the Pauli-blocking corrections for nucleusnucleus collisions extensively discussed in Refs. [43, 48– 50]. The reduction of the nucleon-nucleon cross section in the medium due to Pauli-blocking is considered as given by

$$\sigma_{NN}^{Pauli}(E,\rho_1,\rho_2) = \sigma_{NN}^{free}(E) \frac{1}{1 + 1.892 \left(\frac{|\rho_1 - \rho_2|}{\tilde{\rho}\rho_0}\right)^{2.75}} \\ \times \begin{cases} 1 - \frac{37.02\tilde{\rho}^{2/3}}{E}, & \text{if } E > 46.27\tilde{\rho}^{2/3}, \\ \frac{E}{231.38\tilde{\rho}^{2/3}}, & \text{if } E \le 46.27\tilde{\rho}^{2/3}, \end{cases}$$

where E is the laboratory energy in MeV, $\tilde{\rho} = (\rho_1 + \rho_2)/\rho_0$, with $\rho_0 = 0.16 \text{ fm}^{-3}$, and $\rho_i(r)$ is the local den-

sity at position r within nucleus i. The free nucleonnucleon cross sections were adopted from Ref. [50] where a parametrization was developed to fit the experimental data as a function of the energy.

For proton-nucleus collisions, the Pauli blocking correction reads [51]

$$\sigma_{NN}^{Pauli}(E,\rho_2) = \sigma_{NN}^{free}(E) \left(1 - \frac{8}{5} \frac{E_F}{E}\right), \text{ for } E \ge \frac{8}{5} E_F,$$
(13)

where $E_F = \hbar^2 [3\pi^2 \rho_2(r)/2]^{2/3}/2m_N$, ρ_2 is the total target nucleon density, with $\rho_1(\mathbf{r}) = \delta(\mathbf{r})$. Equation (13) is derived under the assumption that the nucleon-nucleon cross section has an 1/E dependence which is a good approximation for E < 100 MeV. As we show later, for larger energies where $\sigma_{NN}^{free} \sim const.$, the Pauli-blocking contribution to proton-nucleus collisions is small and the exact form of Eq. (13) becomes irrelevant.

The average NN cross section at the distance of closest approach between the projectile and the target is obtained using

$$\langle \sigma_{NN}(E,b) \rangle_{Pauli} = \frac{\int d^3 r \rho_1(\mathbf{r}) \rho_2(\mathbf{r} + \mathbf{b}) \ \sigma_{NN}^{Pauli}(E,\rho_1,\rho_2)}{\int d^3 r \rho_1(\mathbf{r}) \rho_2(\mathbf{r} + \mathbf{b})}$$
(14)

where \mathbf{b} is the impact parameter vector, perpendicular to the beam axis.

To assess the relevance of Pauli-blocking in collisions at low energies, we consider first the p + ²⁰⁸Pb interaction cross section. For ²⁰⁸Pb we generate the density using a Skyrme SLy5 interaction in the Hartree-Fock-Bogoliubov (HFB) approximation [52]. The Skyrme force is a function of numerous terms with contact interactions accounting for coordinate, spin, and isospin dependence. Numerical methods have been developed to calculate nuclear binding energies and other nuclear properties such as the energy density functional $E[\rho]$. To each Skyrme interaction, we added a mixed pairing interaction of the form

$$v(\mathbf{r}, \mathbf{r}') = v_0 \left(1 - \frac{\rho}{2\rho_0}\right) \delta(\mathbf{r} - \mathbf{r}') \tag{15}$$

where $\rho(r) = \rho_n(r) + \rho_p(r)$ is the isoscalar local density. The pairing strength adopted is the same for neutrons and protons, $v_0 = -131.6$ MeV, and the saturation density is fixed at $\rho_0 = 0.16$ fm⁻³.

The energy dependence of the cross section is shown in Figure 1 with data collected from Refs. [53–55]. The solid curve, denoted σ_{Pauli} , includes the effect of Pauli blocking, while the dashed curve, denoted σ_I , does not. The effect of Pauli blocking in σ_{Pauli} is to reduce the nucleon-nucleon cross sections in the medium and consequently the probability for nucleon removal. The interaction cross section is therefore also reduced. It is noticeable from Figure 1 that the inclusion of Pauli-blocking



FIG. 1. Interaction cross sections for $p + {}^{208}Pb$ scattering. Data are from Refs. [53–55]. The solid curve includes the effect of Pauli blocking while the dashed curve does not.



FIG. 2. Interaction cross section for ${}^{12}C + {}^{12}C$ collisions as a function of the bombarding energy in MeV/nucleon. The solid (dashed) curve describes the cross sections (Eq. (6)) calculated with (without) Pauli blocking corrections.

becomes important at proton bombarding energies below 50 MeV, leading to a better agreement with the experimental data. It is also visible that the effect is much reduced at large proton bombarding energies beyond a few hundreds MeV. At the very low energies, below $E_p \sim 30$ MeV the calculations taking into account the Pauli blocking effect run into problems, because the equations erroneously imply that the nucleus becomes fully transparent at the large central densities. This is an overestimation of the effect of Pauli blocking. In this limit, an appropriate Brueckner g-matrix method (see, e.g., Ref. [56]) would be more appropriate than the approach adopted here. However, the simpler method adopted here yields similar results to those using the g-matrix method reported in Ref. [56].

For nucleus-nucleus collisions, we consider the interaction cross section for ${}^{12}C + {}^{12}C$ reactions as a function of the bombarding energy in MeV/nucleon. This is shown in Figure 2 with data collected from Ref. [57, 58]. The calculations were made with charge densities extracted from electron scattering, as also used in Ref. [57]. The same profile density was assumed for neutrons and for protons. Calculations were performed with Eq. (6) which yields the same results as the optical limit cross sections, given by Eq. (11).

Intriguingly, and contrary to expectations, inclusion of Pauli blocking does not appear improve the comparison with the experimental data at low bombarding energies, as seen in Figure 2. The data presented in this figure also seem to be in apparent contradiction with the protonnucleus collision data presented in Figure 1, because the inclusion of Pauli-blocking at low energies seems to play an important role in the proton-nucleus case. The experimental data presented in Figure 2 are also at odds with the studies reported in Refs. [43] for ${}^{12}C + {}^{12}C$ collisions. We have not found a good explanation for this inconsistency. Notice that the lowest data point in Figure 2 is at E = 33 MeV/nucleon. Looking at Figure 1 we see that this is not much below the point of inflection of the solid curve which includes the medium effect. Thus it might seem natural to think that Pauli blocking effects should not play an important role at the energies of the data of Figure 2. However, as reported in Refs. [43, 48–50], the inclusion of two occupied Fermi spheres in momentum space also leads to an expected larger reduction of nucleon-nucleon cross sections in nucleus-nucleus collisions at this energy.

Fermi motion

Now we discuss the effect of Fermi motion on the nucleon-nucleon cross sections. This effect has been studied previously using different methods than the one we describe here. A few examples are given in Refs. [57, 59–62]. For a given impact parameter, the Fermi motion of nucleons inside the projectile and target modify the collision momentum between the nucleons, leading to an effective momentum $p_{eff} = p_0 + \Delta p$, with the unmodified collision energy $E = p_0^2/2m_N$, and Δp denoting the additional momentum due to Fermi motion at the point of collision in the overlap region between the nuclei. We define the Fermi motion averaged NN cross section as

$$\left\langle \sigma_{NN}(b) \right\rangle_{Fermi} = \frac{\int d^3 r \rho_1(\mathbf{r}) \rho_2(\mathbf{r} + \mathbf{b}) \sigma_{NN}^{Fermi}(p_0, \mathbf{r})}{\int d^3 r \rho_1(\mathbf{r}) \rho_2(\mathbf{r} + \mathbf{b})},\tag{16}$$

where the local momentum averaged cross section is

$$\sigma_{NN}^{Fermi}(p_0, \mathbf{r}) = \frac{1}{2\Delta p} \int_{p_0 - \Delta p}^{p_0 + \Delta p} dp \, \sigma_{NN}^{free}(p), \qquad (17)$$

and $\Delta p = p_{F_1} + p_{F_2}$, with the local Fermi momenta $p_{F_i} = \hbar [3\pi^2 \rho_i(r)/2]^{1/3}$, where $\rho_i(r)$ are the total projectile (i = 1) and target (i = 2) densities at position **r**. The Fermi

motion averaged interaction cross section $\langle \sigma_{NN}(b) \rangle_{Fermi}$ is then provided as input in Eqs. (3-11).

We have found that the Fermi motion increases the interaction cross section for ${}^{12}C + {}^{12}C$ collisions, albeit the change is very small. At 30 MeV/nucleon the Fermi motion modified cross section, σ_{FM} , is about 3% larger than the interaction cross section without inclusion of Fermi motion. However, this decreases dramatically as the energy increases, becoming less that 0.1% at energies larger than 200 MeV/nucleon. This is a physically reasonable result, as the net effect of Fermi motion is to increase the interaction cross section at low energies. The reason for the slight increase at low energies is that the NN cross section decreases as 1/E, thus favoring the lower relative energies created by the Fermi motion. The same trend was observed in Ref. [57] where Fermi motion was included using an average based on the Goldhaber model for the internal momentum distribution. We find a smaller correction due to the Fermi motion than reported by those authors. The reason for this difference probably lies in the different prescriptions used to tackle the Fermi motion problem. Notice that our method inherently washes out the effect of the Fermi motion, due to nucleons colliding symmetrically around the bombarding momentum p_0 along the beam axis, as clearly displayed in Eq. (17). It is also clear that the slight increase of the cross sections due to Fermi motion is not sufficient to offset the decrease induced by the Pauli blocking effect.

Higher-order eikonal corrections

The first-order eikonal approximation neglects higher derivatives of the optical potential. In Refs. [63, 64], Wallace has investigated analytical corrections arising from higher-order derivatives in the eikonal S-matrices. To increasing order it was found that the total S-matrix can be cast as a sum of the form

$$S(b) = S^{(0)}(b) + S^{(\text{LO})}(b) + S^{(\text{NLO})}(b) + S^{(\text{NNLO})}(b) + \dots,$$
(18)

with the solutions

$$S^{(0)}(b) = \exp[i\chi_0(b)]$$

$$S^{(LO)}(b) = S^{(0)}(b) \exp[i\chi_1(b)]$$

$$S^{(NLO)}(b) = S^{(LO)}(b) \exp[i\chi_2(b) - \psi_2(b)]$$

$$S^{(NNLO)}(b) = S^{(NLO)}(b) \exp\{i[\chi_3(b) + \lambda_3(b)] + \psi_3(b)\},$$

(19)

where $S^{(0)}(b)$ is the usual eikonal S-matrix, as in Eq. (10), and we use a notation in which the corrections are to leading-order (LO), next-to-leading order (NLO) and next-to-next-to-leading-order (NNLO). The analytical expressions for the phase shifts entering the modified

S-matrices are [63, 64]

$$\chi_{0}(b) = -2k\epsilon \int_{0}^{\infty} dz \, u(r)$$

$$\chi_{1}(b) = -k\epsilon^{2}(1+\beta_{1}) \int_{0}^{\infty} dz \, u^{2}(r)$$

$$\chi_{2}(b) = -k\epsilon^{3} \left(1 + \frac{5}{3}\beta_{1} + \frac{1}{3}\beta_{2}\right) \int_{0}^{\infty} dz \, u^{3}(r)$$

$$- \frac{b}{24k^{2}} \left[\chi_{0}'(b)\right]^{3},$$
(20)

and

$$\begin{split} \psi_{2}(b) &= \frac{b}{8k^{2}}\chi_{0}'(b)\nabla^{2}\chi_{0}(b)\\ \chi_{3}(b) &= -k\epsilon^{4}\left(\frac{5}{4} + \frac{11}{4}\beta_{1} + \beta_{2} + \frac{1}{12}\beta_{3}\right)\int_{0}^{\infty}dz\,u^{4}(r)\\ &- \frac{b}{8k^{2}}\chi_{1}'(b)\left[\chi_{0}'(b)\right]^{2}\\ \lambda_{3}(b) &= -k\epsilon^{2}\left(1 + \frac{5}{3}\beta_{1} + \frac{1}{3}\beta_{2}\right)\int_{0}^{\infty}dz\left[\frac{1}{2k}\frac{\partial u(r)}{\partial r}\right]^{2}\\ \psi_{3}(b) &= \frac{1}{8k^{2}}\left[b\chi_{0}'(b)\nabla^{2}\chi_{1}(b) + b\chi_{1}'(b)\nabla^{2}\chi_{0}(b)\right]. \end{split}$$
(21)

where $\beta_n = b^n \partial^n / \partial b^n$, is a dimensionless derivative of order *n* in the transverse direction, u(r) = U(r)/U(0), where *U* is the optical potential, and $\epsilon = U(0)/2E$ can be identified as an expansion parameter. For large bombarding energies, *E*, the higher order corrections should become irrelevant. Typical depths of the optical potential are of the order of 100 MeV, leading to a prediction that these corrections should be small for energies above 100 MeV for proton-nucleus collisions and at smaller energies for nucleus-nucleus collisions.

The corrections of the eikonal S-matrix as proposed by Wallace have been incorporated in recent studies of one-nucleon knockout reactions in Ref. [65] and for elastic scattering and breakup reactions involving halo nuclei in Ref. [66]. In both cases, it has been shown that higher-order eikonal corrections are only relevant below 50 MeV/nucleon. Here we investigate if these corrections could modify the fragmentation cross sections appreciably. Higher-order eikonal corrections should apply to protons and neutron removal probabilities, adding to Eqs. (9) and (10). Due to the derivatives connected to β_n the corrections are expected to be larger at the nuclear surface. The usual eikonal approximation is thus valid if $ka \gg 1$, where a is the nuclear diffuseness. The higher order corrections may become important for the nuclear potential, but are irrelevant and cancel out for the Coulomb potential which decreases with 1/r [63, 64].

A few remarks are in order before we proceed. Since the "fragmentation optical potential" represented by Eq. (8), and the corresponding one for its neutron U_n counterpart, are purely imaginary, u(r) in the equations above is real and χ_0 and ϵ are imaginary. Moreover, since the nucleon removal probabilities are obtained from the product SS^* , the additional phases introduced in $S^{(\text{LO})}$ or $S^{(\text{NNLO})}$ do not change the removal probabilities. Additionally, $\psi_2(b)$ does not contribute to the fragment production cross sections. These claims can be checked by inspection of Eqs. (19) and (21). In summary, for the purposes of calculating the isotope production and interaction cross sections, the only higher-order correction needed for the eikonal phase is the addition of the phase χ_2 in equation (20) to the standard eikonal phase χ_0 . Thus, it is naturally expected that higher-order eikonal corrections will be smaller in the case studied here than those reported in Refs. [65, 66]

In Table I we show in the second and third columns the interaction cross section for $^{12}C + ^{12}C$ collisions at selected bombarding energies. In the third column we show the interaction cross section including higher-order eikonal corrections, σ_{eik} . These corrections are small, except for energies below 100 MeV/nucleon, and even in this case they do not comprise more than a 1% correction.

TABLE I. Interaction cross sections σ_I for ${}^{12}\text{C} + {}^{12}\text{C}$ collisions at selected bombarding energies. σ_{heik} denotes interaction cross sections including higher-order eikonal corrections, while σ_{Coul} denotes interaction cross sections including Coulomb repulsion.

E (MeV/nucl)	σ_I	σ_{heik}	σ_{Coul}
50	1139	1133	1026
70	980	977	969
100	909.3	908	902
200	801.9	802.0	796
500	806.6	806.6	804

Coulomb repulsion correction

At low collision energies one must add a correction due to the Coulomb deflection of the nuclear trajectory. This correction amounts to replacing the impact parameter variable within the integral with b' due to repulsion as nuclei pass by at closest distance. That is, [46]

$$b' = a_0 + \sqrt{a_0^2 + b^2},\tag{22}$$

where $a_0 = Z_P Z_T e^2 / (\gamma \mu v^2)$ is half the distance of closest approach in a head-on collision of point charged particles. $\gamma = (1 - v^2/c^2)^{-1/2}$ is the Lorentz contraction factor and μ is the reduced mass of the system. This correction leads to an improvement of the eikonal amplitudes for the scattering of heavy systems in collisions at low energies (see Ref. [46] for more details).

The fourth column in Table I shows the cross section corrected by Coulomb recoil. Both the higher-order eikonal correction and the Coulomb repulsion correction are small. But the Coulomb recoil affects the cross section more, specially at lower energies. The Coulomb recoil increases the closest distance between the projectile and the target and the corresponding nucleon removal probability also decreases.

Relativistic corrections

It is evident that relativistic corrections are very important in nuclear reactions at high energies as low as 100 MeV/nucleon where the nucleon mass has already increased by about 10%. This is very important for inelastic processes, such as the excitation of giant resonances, discussed in the next section. The corrections are not only related to kinematics, but also in the reaction dynamics. Except for a few studies [67], this important issue has been largely ignored in the literature. Notably, the basic concept of optical potential is not a relativistic one, as one needs a four-potential to comply with proper Lorentz transformations. Besides, simultaneity and retardation are a major theory hurdle when dealing with relativistic many-body systems [67, 68]. Therefore, the concept of an optical potential, as displayed by Eq. (8)lacks the proper relativistic treatment. In this work we use the optical potential concept to relate it to the calculation of the eikonal phase, Eq. (9), and the eikonal S-matrix, Eq. (10), although they can also be deduced on probabilistic grounds. The optical potential of Eq. (8) is also a necessary condition to calculate the higher order eikonal corrections based on the adopted formalism from Refs. [63, 64].

Glauber models depend on the transverse coordinates of the colliding systems, usually identified as impact parameters. These coordinates are Lorentz covariant, the probabilities and cross sections being thus unchanged by Lorentz transformations. However, in Eq. (3) and the following ones contain integrations along the longitudinal direction. Longitudinal directions are not Lorentz invariant, as they contract along the direction of motion. This amounts to change $dz \rightarrow dz/\gamma$ and $\rho(\mathbf{b}, z) \rightarrow \gamma \rho(\mathbf{b}, z)$ and the Lorentz factor cancels out. This comes at no surprise because our adopted Glauber procedure amounts to calculate the probability of a binary collision along a tube of thickness d^2b in the direction of motion. The number of nucleons within the tube is unchanged by Lorentz contraction.

Reaction versus interaction cross section

The excitation of nuclear states leading to decay by nucleon emission can also contribute to the fragmentation cross sections. Adding these contributions to the interaction cross section yields the so-called reaction cross



FIG. 3. Corrections of the neutron removal cross sections for 124 Sn and 132 Sn incident on carbon and hydrogen targets at 900 MeV/nucleon. Corrections include Pauli blocking (1), Coulomb recoil (2), higher order eikonal corrections (3), Fermi motion (4), and Coulomb and nuclear excitation of giant resonances (5). The lines are guides to the eye.

section. The major contribution comes from the excitation of giant resonances (GR), a highly collective excitation mode of the nucleus. Denoting this contribution by σ_{GR} , the reaction cross section is $\sigma_R = \sigma_I + \sigma_{GR}$. The excitation of giant resonances through the nuclear and Coulomb interaction between the nuclei has been extensively discussed in past publications. In Ref. [30] it was studied in the same context as the one discussed in this work, namely its contribution to fragmentation reactions through the excitation and decay of giant resonances, mainly by neutron emission. Here we adopt the same method as in Ref. [30] to calculate the nuclear and Coulomb excitation cross sections with relativistic corrections included.

Since the motivation of this work is to study the effect of the neutron skin along an isotopic chain via the analysis of experiments with neutron rich nuclei at high energies, we consider here the fragmentation cross sections of heavier nuclei incident on proton and carbon targets. In Figure 3 we show the contribution of the several corrections of the neutron removal cross sections for ¹²⁴Sn and ¹³²Sn incident on carbon and hydrogen targets at 900 MeV/nucleon. Here we use the SLY4 [15] interaction to obtain the ¹²⁴Sn and ¹³²Sn proton and neutron densities. Corrections include Fermi motion (1), Coulomb recoil (2), higher order eikonal corrections (3), Pauli blocking correction (4), and Coulomb and nuclear excitation of giant resonances (5). The lines are guides to the eye. The parameters used for the excitation of giant resonances are the same as reported in Ref. [30] and we assume that they decay by neutron emission only. It is evident that all corrections, except for the last two (Pauli blocking and excitation of giant resonances) are very small at this bombarding energy. For proton targets, the Pauli correction is less than 2% and the contribution of nuclear and Coulomb excitation cross section is below 0.3%. However, for carbon targets, the nuclear excitation cross sections are of the order of 60-80 mb, making an important contribution to the neutron removal reaction cross sections.

Secondary binary collisions

As the mass of the fragment nucleus increases, secondary collisions of abraded nucleons with others in the fragment become an increasingly important source of additional abraded nucleons. This is a simple consequence of the increase in the thickness of the matter that an abraded nucleon must traverse to escape the fragment. These secondary collisions occurring after the abrasion stage are not taken into account in the Glauber formalism adopted here. However, due to the smaller energy of the secondary nucleons and the higher Coulomb barrier of a heavier fragment, the secondary emitted particles are predominantly neutrons. They thus tend to broaden the mass distribution of each charge state of the fragment without changing the charge distribution. The total cross section for each charge state, in particular, the total neutron removal cross section, is therefore relatively immune to this process. We do not consider such secondary binary collisions in this work.

TESTING NEUTRON DENSITIES FROM MEAN-FIELD MODELS

Comparison to fragmentation reactions

We compare our calculations of reaction cross sections, σ_R , charge changing cross sections, $\sigma_{\Delta Z}$, and neutron removal cross sections, $\sigma_{\Delta N}$, to several experiments using nuclear densities generated using the nonrelativistic Hartree-Fock-Bogoliubov (HFB) approximation with 23 different Skyrme interactions and the relativistic mean field (RMF) approximation using 6 nonlinear and 2 density-dependent interactions. The Skyrme interactions used are the SIII [7], SKA and SKB [8], SKM* [9], SKP [10], UNE0 and UNE1 [11], SKMP [12], SKI2, SKI3, SKI4 and SKI5 [13], SLY230A [14], SLY4, SLY5, SLY6, and SLY7 [15], SKX [16], SKO [17], SK255 and SK272 [18], HFB9 [19] and SKXS20 [20]. The calculations for the ENE0 and UNE1 interactions performed done with the modified version of the code HFBTO code [11]. To all Skyrme interactions a pairing force was added with the form described in Eq. (15). The RMF calculations used the six nonlinear interactions BSR6 and BSR14 [21, 22], FSUZG00 [23, 69], NL3* [24], NLRA1 [25], NL3-II [26], and the two density-dependent interactions DDME2 [27] and PKDD [28]. The RMF calculations used the effective pairing force described in Ref. [29].

The large number of interactions covered in this work encompasses a wide range of nuclear matter properties. A sample of the variation of the parameters of the symmetric matter EOS is shown for few of these interactions in Table II. It is worthwhile noticing that the slope parameter L is the most uncertain of all quantities.

TABLE II. For a few interactions used in this work, we show a sample of the nuclear matter properties at saturation density. All quantities are in MeV units.

Skyrme	K_0	J	L	Skyrme	K_0	J	L
SIII	355.	28.2	9.91	DDME2	251.	32.3	51.
SKP	201.	30.0	19.7	FSUG00	240.	31.43	62.19
SKX	271.	31.1	33.2	SKXS20	202.	35.5	67.1
HFB9	231.	30.0	39.9	SKO	223.	31.9	79.1
SLY5	230.	32.0	48.2	SKI5	255.	36.6	129.

We consider reactions in the energy range from 100 MeV/nucleon to 1 GeV/nucleon, including corrections due to Pauli blocking, Fermi motion, eikonal higher-order corrections, and Coulomb recoil. These are small corrections, at the level of 2% or less, and we can easily quantify them with the theory discussed in the previous sections. The most relevant additions to the interaction cross section are due to the excitation of giant resonances, discussed in the last section. In Figure 4 we compare the results of our calculations with total reaction cross sections for $p+{}^{12}C$ (upper panel) and $p+{}^{208}Pb$ (lower panel) collisions compared to experimental data from Refs. [55, 70]. The shaded bands represent calculations using densities obtained with non-relativistic and relativistic mean-field calculations with the 31 different interactions. We arbitrarily select the five best interactions based on the smallest chi-square fit to the data. For the data presented in the upper panel of Fig. 4 they are the DDME2, SKO, SKX, SLY230A and SKXS20, yielding an average slope parameter of $\langle L \rangle = (54.9 \pm 23)$ MeV, where the error reflects the range of values obtained from the interactions. The nucleus ¹²C is not expected to have a sizable neutron skin and the neutron and proton densities should be well described by the electron scattering data reported in Ref. [57]. The large variation of the reaction cross sections suggests that the density calculations using mean field methods is not appropriate for such a light nucleus. In fact, the lower panel of Fig. 4 shows that for a large nucleus the calculated mean field densities are more uncertain allowing for a bet-



FIG. 4. Calculated total reaction cross sections for $p+{}^{12}C$ (upper panel) and $p+{}^{208}Pb$ (lower panel) collisions compared to experimental data from Refs. [55, 70]. The shaded band represents the spread of calculations using densities obtained with non-relativistic and relativistic mean-field calculations using a total of 31 different interactions.

ter inference of the neutron skin by a comparison with experimental data. In this case, best χ^2 comparisons with the data are obtained with the BSR14, SLY4, SKA, SKI4, and UNE0. They yield an average slope parameter $\langle L \rangle = (59.5 \pm 14.1)$ MeV and an average neutron skin of $\langle \Delta r_{np} \rangle = (0.188 \pm 0.021)$ fm for the ²⁰⁸Pb nucleus. This is in rather good agreement with $\Delta r_{np} = (0.156 \pm 0.023)$ fm reported in the experimental analysis of Ref. [35] and substantially smaller than the value $\Delta r_{np} = 0.283 \pm 0.071$ fm from the average of PREX1 and PREX2 experiments implying $\langle L \rangle = (106 \pm 37)$ MeV [33].

It must be pointed out that the extraction of the neutron skin by comparison with the reaction cross section data presented in the lower panel of Fig. 4 is rather deceiving. It is evident from the figure that the accuracy of the data is not good enough to constrain the best interactions reproducing the data. The limited information we can extract from this comparison is that the energy dependence of the cross section data can be rather well described with the densities from microscopic calculations, except for the highest energy points. Besides, the neutron skin extracted from such a comparison is more a convergence of the mean field calculations toward a narrow range of neutron skins than a true constraint placed by the experimental data.

Total charge changing reactions, $\sigma_{\Delta Z}$ [36–41], namely



FIG. 5. Calculated total charge changing cross section, $\sigma_{\Delta Z}$, for the reaction ²⁸Si + ¹²C compared to experimental data from Refs. [71–77]. The shaded band represents densities for ²⁸Si obtained with non-relativistic and relativistic mean-field calculations with a total of 31 different interactions.

fragmentation reactions in which all isotopes are measured, can be a good probe of neutron skins because, if the measurements are accurate, they can be compared to a simple Glauber calculation of primary fragments, similar to Eq. (7). A comparison of our calculations to the experimental data from Refs. [71–77] for the reaction 28 Si + 12 C is made in Fig. 5. The shaded band represents the spread of all calculations done with the densities for ²⁸Si obtained from non-relativistic and relativistic mean-field calculations using a total of 31 different interactions. The ¹²C density was taken from electron scattering experiments [57]. The comparison is again not good enough to constrain the best microscopic interactions and the corresponding EoS. It is also apparent that the calculations fail to reproduce the experimental data beyond 700 MeV/nucleon. This is an unexpected result because the experimental data seem to indicate that the nuclear transparency increases at large energies. In this case, the 5 smallest χ^2 comparison with the data correspond to the SLY7, SKI4, BSR14, SKP and SKB interactions. They yield an average slope parameter $\langle L \rangle = (43.6 \pm 20.0)$ MeV and an average neutron skin of $\langle \Delta r_{np} \rangle = (0.0348 \pm 0.0081)$ fm for the ²⁸Si nucleus. There are currently no other experiments dedicated to study the neutron skin of ²⁸Si than the ones based on reaction cross sections and charge-changing cross sections. It is clear from the experimental data accumulated so far that more experimental efforts are needed.

In Figure 6, upper panel, we shown the calculated interaction cross section, σ_I , for the reaction Ca + 12 C and different calcium isotopes compared to experimental data from Ref. [78]. The shaded band represents the spread of calculations using densities obtained with nonrelativistic and relativistic mean-field calculations with a total of 31 different interactions. The lower panel dis-



FIG. 6. User panel: Calculated interaction cross section σ_I at 280 MeV/nucleon for the reaction Ca + ¹²C and different calcium isotopes compared to experimental data from Ref. [78]. The shaded band represents the spread of calculations using calcium densities obtained with non-relativistic and relativistic mean-field calculations with a total of 31 different interactions. Lower panel: The neutron skin for calcium isotopes calculated with the different mean-field models. The data from the CREX experiment for ⁴⁸Ca [80] is also shown.

plays the neutron skin for calcium isotopes calculated with the different mean-field models. The data from the CREX experiment for ${}^{48}Ca$ [80] is also shown.In Figure 7 we show the calculated charge changing cross section $\sigma_{\Delta Z}$ at 280 MeV/nucleon for the reaction Fe + ¹²C and different iron isotopes compared to experimental data from Ref. [79]. The shaded band represents the spread of calculations using densities obtained with nonrelativistic and relativistic mean-field calculations with a total of 31 different interactions. In both cases, it is clear that the measurements for interaction cross sections and for charge changing cross sections are still not accurate enough to extract accurate information on neutron skins by comparison with theoretical predictions for the cross sections. In particular, for the data presented in Figure 7, the agreement with theory is very poor. The reason for these differences are not well understood. In the case of $Ca + {}^{12}C$ collisions presented in Figure 6, one is tempted to extract numbers for the ⁴⁸Ca nucleus, where a reasonable agreement with experimental data is found,



FIG. 7. Calculated charge changing cross section $\sigma_{\Delta Z}$ at 280 MeV/nucleon for the reaction Fe + ¹²C and different iron isotopes compared to experimental data from Ref. [79]. The shaded band represents the spread of calculations using densities obtained with non-relativistic and relativistic mean-field calculations with a total of 31 different interactions.

to compare with the findings of the CREX experiment [80]. The HFB9 interaction yields the closest value for the interaction cross section, predicting a neutron skin of $\Delta r_{np} = 0.156$ fm and a slope parameter L = 39.8 MeV. The second best result is obtained with the SKX interaction, yielding $\Delta r_{np} = 0.167$ fm and L = 33.2 MeV. These results for the neutron skin are about 20% higher than the value $\Delta r_{np} = 0.121 \pm 0.026$ fm reported by the CREX experiment [80] and, in contrast, the slope parameter L derived from these two interactions are also much smaller than the value $L \simeq 119$ MeV obtained with the NL3 interaction [81] and favored by the CREX experiment [80].

Comparison to past and presently proposed methods

It is clear from our limited analysis that the present status of the experimental data on fragmentation reactions is not yet at the level of accuracy to allow a good constraint on the best possible microscopic interactions and in the process to obtain the best possible EoS for neutron stars. Other experimental methods have been devised in the past but also lack the necessary accuracy needed to constrain the best possible EoS.

Recently, a new method was introduced, with the neutron skin extracted from an analysis of the parity violation effect on electron scattering off lead and calcium nuclei [82–84]. This method would in principle allow a measurement independent of the complications of the strong force. As we discussed before, a recent experiment based on this technique were performed at the Jefferson Laboratory [33] with a ²⁰⁸Pb target and found a large neutron skin of $\Delta r_{np} = 0.283 \pm 0.071$ fm, implying a slope parameter of $L = 106\pm37$ MeV, larger than expected from most microscopic calculations and also from other experiments

with the same nucleus. For example, Ref. [35] reported a value of $\Delta r_{np} = 0.156^{+0.025}_{-0.021}$ fm. Moreover, astronomical observations from NICER and LIGO/Virgo collaborations are compatible with much smaller values for Δr_{np} and *L*. Although the JLab experiment was successful, statistical significance was not achieved [33]. It has been advertised by the JLab [80] that the CREX experiment, a variant of the PREX experiment but this time using ⁴⁸Ca as target, has obtained a much smaller value of the neutron skin, $\Delta r_{np} = 0.121 \pm 0.026$ fm. Hence, the PREX and the CREX experiments seem to provide incompatible results for the size of the neutron skin.

In Figure 8 we show the neutron skin Δr_{np} of several nuclei calculated with the SLY4 interaction as a function of the isospin asymmetry parameter $\delta = (N-Z)/A$. The experimental values were extracted from antiproton annihilation data [85, 86] and with data from the CREX experiment for 48 Ca [80] from the PREX experiment for 208 Pb (two data points) [33, 87]. It is evident that, as expected, both the experimental data and theory display a trend of increasing neutron skin with increasing isospin asymmetry parameter $\delta = (N - Z)/A$ for all nuclei studied so far. It is also clear from the plot that the popular SLY4 interaction, as many others, yield a pretty reasonable description of the average behavior of the neutron skin of most studied nuclei with the antiproton and electron PV scattering techniques. The PREXI [33] and PREXII [87] data are hard to explain and are in tension with the microscopic calculations. It is worthwhile mentioning that the extraction of neutron skins using antiprotons is more of a direct technique for this purpose. The antiproton-nucleon interaction is very strong and antiprotons interacting with nuclei are absorbed and annihilate already at the nuclear periphery, where the nucleon density is significantly smaller than the central nuclear density. One uses radiochemical methods [88], supplemented with radiochemical data [89] by means of in-beam antiprotonic x-ray studies [85, 86] thus determining strong-interaction level widths and shifts in isotopically enriched targets. The major hurdle in analyzing the antiproton data is the theoretical description of antiproton-nucleus interactions (optical potential) [90]. A renewed interest in theoretical derivations of the antiproton-nucleus optical potential has recently emerged to support future experiments [91]. In contrast, both electron PV scattering experiments and nuclear fragmentation data analysis are based on a comparison with theory input for matter density obtained with microscopic calculations either through neutron density form factors, or Glauber nucleon knockout profile functions. Therefore, the three techniques described here are very different, complementary, but strongly dependent on theory input. It is also clear that one needs to increase the number, accuracy and methods used in the experiments to narrow down the proper EoS of nuclear matter.

A promising method to extract the nuclear EoS is still



FIG. 8. Neutron skin thickness as a function of the isospin asymmetry parameter $\delta = (N-Z)/A$ compared to experimental data extracted from antiproton annihilation data [85, 86] and with data from the CREX experiment for ⁴⁸Ca [80] and for the PREX experiment for ²⁰⁸Pb (two data points) [33, 87]. The small triangles guided by a dashed line correspond to the prediction of the SLY4 interaction in the HFB formalism.

based on the effect on the nucleus-nucleus collisions of the growth of the neutron skin along an isotopic chain. This method can be tested in different experiments using hadronic probes and a plethora of techniques such as elastic and inelastic scattering, photo absorption, fragmentation, etc. In particular, radioactive beam facilities can be very helpful, as they provide isotopic beams with different neutron/proton ratios. In Figure 9 we present the neutron skin Δr_{np} of Sn isotopes calculated with the DDME2 microscopic interaction. The experimental data for Sn nuclei were measured with (p, p) reactions (triangles) [92], antiproton atoms (stars) [85], giant dipole resonance method (squares) [93] and spin dipole resonance method (circles) [94, 95]. The small triangles and dashed line displays the numerical calculations for the neutron skin, Δr_{np} , calculated with the DDME2 relativistic mean field interaction. This interaction predicts a slope parameter L = 51 MeV. It is noticeable that the distinct probes of the neutron skin seem to yield inconsistent results, requiring a further assessment of the experiments systematics. The small error bars and apparent well behaved trend of the experimental data based on polarized proton scattering [85, 96] seem to provide a promising way to move forward with radioactive beams and indirect techniques. The same sort of comparison of more accurate data along an isotopic chain with theory prediction as in Fig. 9 will provide another useful constraint on the symmetry energy content of the ANM EOS.

The experiments using fragmentation reactions can and will certainly help in determining which additional information that can be gleaned from the experiments. In particular, fragmentation reactions can be performed



FIG. 9. Neutron skin Δr_{np} of Sn isotopes calculated with the DDME2 interaction in the RMF formalism. The experimental data Sn were measured with (p, p) reactions (triangles) [92], antiproton atoms (stars) [85], giant dipole resonance method (squares) [93] and spin dipole resonance method (circles) [94, 95] To avoid superposition of the data, we displace them slightly and label each group according to their masses A.

using combinations of numerous projectiles and targets [41]. Using proton and carbon targets have the advantage of testing the skin with a deeply penetrating probe (proton) and a more surface oriented one (carbon). Light targets are preferable to avoid large contributions from fragmentation due to Coulomb excitation. Fragmentation reactions can also use projectiles with a varying combination of neutron to proton ratio, in particular with radioactive beam facilities. Tin, for example, has a long isotopic chain and is one of the candidates for testing an isospin dependence of the neutron skin Δr_{np} . In Table III we summarize the calculations for the total neutron removal cross sections using ¹²C and proton targets for a selected number of tin isotopes and energies compatible with the GSI facility in Germany. Here we use the SLY4 interaction in the HFB formalism. For this interaction, the slope parameter is L = 45.9 MeV the neutron skins and are: (a) 124 Sn, $\Delta r_{np} = 0.171$ fm, (b) 128 Sn, $\Delta r_{np} = 0.198$ fm, (c) 132 Sn, $\Delta r_{np} = 0.222$ fm, and (d) 134 Sn, $\Delta r_{np} = 0.250$ fm. The increase of the skin as a function of the neutron-proton asymmetry attests the value of studying the neutron skin evolution along an isotopic chain to better constrain the best interaction to fit the data.

From Table III one sees that the most challenging correction to the rather clean and commonly used theoretical picture presented by the Glauber-like method described by Eqs. (3-7) is the excitation of collective modes

TABLE III. Total neutron removal cross sections for Sn projectiles incident on carbon and hydrogen targets using the SLY4 interaction in the HFB formalism. The corrections due to Pauli blocking, Coulomb repulsion (Coul), and excitation of collective (GDR+GQR) states are added in sequence.

	E_{beam}	Target	$\sigma_{\Delta N}$	(+ Pauli $)$	$(+ \operatorname{Coul})$	(+ GR)
	$\mathrm{MeV}/\mathrm{nucl}$		$^{\mathrm{mb}}$	$^{\mathrm{mb}}$	${\rm mb}$	$^{\mathrm{mb}}$
124 Sn	400	С	419	429	427	493
		Η	254	263	263	271
	900	\mathbf{C}	420	425	424	469
		Η	268	273	272	279
$^{128}\mathrm{Sn}$	900	\mathbf{C}	449	455	454	516
		Η	288	291	290	297
$^{132}\mathrm{Sn}$	600	\mathbf{C}	472	481	480	541
		Η	365	370	367	374
	900	\mathbf{C}	476	481	480	542
		Η	306	309	308	315
$^{134}\mathrm{Sn}$	600	\mathbf{C}	496	505	504	563
		Η	378	383	381	387
	900	\mathbf{C}	501	506	505	565
		Η	321	323	322	328

leading to neutron emission. This has been rarely discussed in the literature in this context, except in a few publications [30]. But using both proton and carbons targets will help to disentangle the effects of this correction. Performing experiments at different energies will also help to constrain the dependence of the cross sections on the neutron and proton nuclear densities. On the other hand, in inclusive fragmentation experiments as proposed in Ref. [97], the total neutron removal cross section could be determined with a precision of around 1%. In order to avoid the theoretical uncertainty of calculation the contribution due to collective excitations as mentioned above, a recently proposed experiment will directly determine in addition this part of the cross section, which contributes to about 10% to the total cross section. If neutron removal is caused only by neutron evaporation after excitation, the neutrons will be detectable at small angles in the laboratory around the beam axis (due to the high beam velocity). The total neutron-changing cross section minus this part can be directly compared to Glauber calculations using different theoretical densities (corresponding to different L values and neutron skins). By this comparison, the L value and $\langle r^2 \rangle_n$ will be constrained. A detailed sensitivity study for this approach is discussed in Ref. [41]. The uncertainty of the extracted value will contain besides the experimental error the uncertainty of the Glauber model used.

CONCLUSIONS

In this work we have made an analysis of the contributions of numerous physics effects on the interaction, reaction, charge-changing and neutron-changing cross sections. We have studied the contribution of (a) Pauliblocking, (b) Fermi motion, (c) eikonal corrections, (d) Coulomb recoil, (d) Coulomb and (e) nuclear excitation of giant resonances. Clear discrepancies with published experimental data have been found for interaction, reaction, charge-changing and neutron-changing cross sections. Notably, the data reported in the literature for charge-changing reactions seem to favor a transparency scenario for the cross sections at the largest energies. This is not easy to understand, as the nucleon-nucleon cross sections are rather energy independent at these energies.

It is also clear that the data for total neutron removal, or neutron-changing, cross sections are not yet accurate enough to provide a tight constraint on the mean-field calculations and the corresponding EoS. In particular, we have observed in our calculations that old interac-⁼ tions such as the SKIII, SKA and SKB interactions [7, 8] fare better in reproducing some of the published data than more modern and celebrated interactions such as the SLY, DDME2, or UNEDF interactions [11, 15, 27]. No single interaction stands out as a better one to reproduce the experimental data that we have discussed in this work. At this stage, a global fit or machine learning approach to constrain the parameters of the best theory does not seem to be very useful in view of the scarcity and inconsistency of the data reported in the literature. Also, the celebrated PREX experiment has not been conclusive, yielding an unexpectedly large neutron skin for lead. The CREX experiment has obtained a much smaller neutron skin than the expected by an extrapolation of the PREX experiment.

The recently proposed experimental campaign to determine neutron-changing cross sections in inverse kinematics for projectiles with a large range of isospin asymmetry [41] will help to increase the available data to assess the dependence of the neutron skin on the isospin asymmetry parameter $\delta = (N - Z)/A$. These experiments will have a direct impact on the determination of the symmetry term of the EoS.

Acknowledgments. E.A.T. acknowledges partial support by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brazil (CAPES). T.A. acknowledges support by the German Federal Ministry of Education and Research (BMBF, project 05P2015RDFN1). C.A.B. acknowledges support by the U.S. DOE grant DE-FG02-08ER41533 and the Helmholtz Research Academy Hesse for FAIR. B.V.C. acknowledges support from grant 2017/05660-0 of the São Paulo Research Foundation (FAPESP), grant 303131/2021-7 of the CNPq and the

- * thomas.aumann@tu-darmstadt.de
- [†] carlos.bertulani@tamuc.edu
- [‡] brettvc@gmail.com
- [§] soberania@gmail.com
- Gordon Baym, Tetsuo Hatsuda, Toru Kojo, Philip D Powell, Yifan Song, and Tatsuyuki Takatsuka. From hadrons to quarks in neutron stars: a review. *Reports* on Progress in Physics, 81(5):056902, 2018.
- [2] Jeremy W. Holt and Yeunhwan Lim. Universal correlations in the nuclear symmetry energy, slope parameter, and curvature. *Physics Letters B*, 784:77–81, 2018.
- [3] J.M. Lattimer. Neutron stars and the nuclear matter equation of state. Annual Review of Nuclear and Particle Science, 71(1):433-464, 2021.
- [4] Nguyen Hoang Dang Khoa, Ngo Hai Tan, and Dao T. Khoa. Spin symmetry energy and equation of state of spin-polarized neutron star matter. *Phys. Rev. C*, 105:065802, 2022.
- [5] X. Roca-Maza, M. Centelles, X. Viñas, and M. Warda. Neutron skin of ²⁰⁸Pb, nuclear symmetry energy, and the parity radius experiment. *Phys. Rev. Lett.*, 106:252501, 2011.
- [6] X. Roca-Maza, X. Viñas, M. Centelles, B. K. Agrawal, G. Colò, N. Paar, J. Piekarewicz, and D. Vretenar. Neutron skin thickness from the measured electric dipole polarizability in ⁶⁸Ni, ¹²⁰Sn, and ²⁰⁸Pb. *Phys. Rev. C*, 92:064304, 2015.
- [7] M. Beiner, H. Flocard, Nguyen Van Giai, and P. Quentin. Nuclear ground-state properties and self-consistent calculations with the skyrme interaction: (i). spherical description. *Nuclear Physics A*, 238(1):29–69, 1975.
- [8] H.S. Köhler. Skyrme force and the mass formula. Nuclear Physics A, 258(2):301–316, 1976.
- [9] J. Bartel, P. Quentin, M. Brack, C. Guet, and H.-B. Håkansson. Towards a better parametrisation of Skyrmelike effective forces: A critical study of the SkM force. *Nuclear Physics A*, 386(1):79–100, 1982.
- [10] J Dobaczewski, H Flocard, and J Treiner. Hartree-Fock-Bogolyubov description of nuclei near the neutron-drip line. Nuclear Physics A, 422(1):103–139, 1984.
- [11] M.V. Stoitsov, N. Schunck, M. Kortelainen, N. Michel, H. Nam, E. Olsen, J. Sarich, and S. Wild. Axially deformed solution of the Skyrme-Hartree-Fock-Bogoliubov equations using the transformed harmonic oscillator basis (ii) hfbtho v2.00d: A new version of the program. *Computer Physics Communications*, 184(6):1592–1604, 2013.
- [12] L. Bennour, P-H. Heenen, P. Bonche, J. Dobaczewski, and H. Flocard. Charge distributions of ²⁰⁸Pb, ²⁰⁶Pb, and ²⁰⁵Tl and the mean-field approximation. *Phys. Rev. C*, 40:2834–2839, 1989.
- [13] P.-G. Reinhard and H. Flocard. Nuclear effective forces and isotope shifts. *Nuclear Physics A*, 584(3):467–488, 1995.
- [14] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and R. Schaeffer. A Skyrme parametrization from subnuclear to neutron star densities. *Nuclear Physics A*, 627(4):710– 746, 1997.
- [15] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and

R. Schaeffer. A skyrme parametrization from subnuclear to neutron star densities part ii. nuclei far from stabilities. *Nuclear Physics A*, 635(1):231–256, 1998.

- [16] B. Alex Brown. New Skyrme interaction for normal and exotic nuclei. *Phys. Rev. C*, 58:220–231, 1998.
- [17] P.-G. Reinhard, D. J. Dean, W. Nazarewicz, J. Dobaczewski, J. A. Maruhn, and M. R. Strayer. Shape coexistence and the effective nucleon-nucleon interaction. *Phys. Rev. C*, 60:014316, 1999.
- [18] B. K. Agrawal, S. Shlomo, and V. Kim Au. Nuclear matter incompressibility coefficient in relativistic and nonrelativistic microscopic models. *Phys. Rev. C*, 68:031304, 2003.
- [19] S. Goriely, M. Samyn, J.M. Pearson, and M. Onsi. Further explorations of Skyrme-Hartree-Fock-Bogoliubov mass formulas. iv: Neutron-matter constraint. *Nuclear Physics A*, 750(2):425–443, 2005.
- [20] M. Dutra, O. Lourenço, J. S. Sá Martins, A. Delfino, J. R. Stone, and P. D. Stevenson. Skyrme interaction and nuclear matter constraints. *Phys. Rev. C*, 85:035201, 2012.
- [21] Shashi K. Dhiman, Raj Kumar, and B. K. Agrawal. Nonrotating and rotating neutron stars in the extended field theoretical model. *Phys. Rev. C*, 76:045801, 2007.
- [22] B. K. Agrawal. Asymmetric nuclear matter and neutron skin in an extended relativistic mean-field model. *Phys. Rev. C*, 81:034323, 2010.
- [23] Raj Kumar, B. K. Agrawal, and Shashi K. Dhiman. Effects of ω meson self-coupling on the properties of finite nuclei and neutron stars. *Phys. Rev. C*, 74:034323, 2006.
- [24] G.A. Lalazissis, S. Karatzikos, R. Fossion, D. Pena Arteaga, A.V. Afanasjev, and P. Ring. The effective force nl3 revisited. *Physics Letters B*, 671(1):36–41, 2009.
- [25] M. Rashdan. Structure of exotic nuclei and superheavy elements in a relativistic shell model. *Phys. Rev. C*, 63:044303, 2001.
- [26] G. A. Lalazissis, J. König, and P. Ring. New parametrization for the lagrangian density of relativistic mean field theory. *Phys. Rev. C*, 55:540–543, 1997.
- [27] G. A. Lalazissis, T. Nikšić, D. Vretenar, and P. Ring. New relativistic mean-field interaction with density-dependent meson-nucleon couplings. *Phys. Rev. C*, 71:024312, 2005.
- [28] Wenhui Long, Jie Meng, Nguyen Van Giai, and Shan-Gui Zhou. New effective interactions in relativistic mean field theory with nonlinear terms and density-dependent meson-nucleon coupling. *Phys. Rev. C*, 69:034319, 2004.
- [29] B. V. Carlson and D. Hirata. Dirac-Hartree-Bogoliubov approximation for finite nuclei. *Phys. Rev. C*, 62:054310, 2000.
- [30] C. A. Bertulani and J. Valencia. Neutron skins as laboratory constraints on properties of neutron stars and on what we can learn from heavy ion fragmentation reactions. *Phys. Rev. C*, 100:015802, 2019.
- [31] G.F. Burgio, H.-J. Schulze, I. Vidana, and J.-B. Wei. Neutron stars and the nuclear equation of state. *Progress in Particle and Nuclear Physics*, 120:103879, 2021.
- [32] X. Viñas, M. Centelles, X. Roca-Maza, and M. Warda. Density dependence of the symmetry energy from neutron skin thickness in finite nuclei. *The European Physi*cal Journal A, 50(2):27, 2014.
- [33] D. Adhikari et al. Accurate determination of the neutron skin thickness of ²⁰⁸Pb through parity-violation in electron scattering. *Phys. Rev. Lett.*, 126:172502, 2021.
- [34] The value and error bar reported in Ref. [33] is a combina-

tion of two separate measurements, denoted as PREX-1 and PREX-2.

- [35] A. Tamii et al. Complete electric dipole response and the neutron skin in Pb-208. *Phys. Rev. Lett.*, 107(6), 2011.
- [36] B. Blank et al. Charge-changing cross sections of the neutron-rich isotopes ^{8,9,11}Li. Zeitschrift für Physik A Hadrons and Nuclei, 343(4):375–379, 1992.
- [37] T. Yamaguchi, I. Hachiuma, A. Kitagawa, K. Namihira, S. Sato, T. Suzuki, I. Tanihata, and M. Fukuda. Scaling of charge-changing interaction cross sections and pointproton radii of neutron-rich carbon isotopes. *Phys. Rev. Lett.*, 107:032502, 2011.
- [38] S. Terashima et al. Proton radius of ¹⁴Be from measurement of charge-changing cross sections. Progress of Theoretical and Experimental Physics, 2014(10), 2014. 101D02.
- [39] A. Estradé et al. Proton radii of ¹²⁻¹⁷B define a thick neutron surface in ¹⁷B. *Phys. Rev. Lett.*, 113:132501, 2014.
- [40] D. T. Tran et al. Charge-changing cross-section measurements of ¹²⁻¹⁶C at around 45A MeV and development of a glauber model for incident energies 10 – 2100A MeV. *Phys. Rev. C*, 94:064604, 2016.
- [41] T. Aumann, C. A. Bertulani, F. Schindler, and S. Typel. Peeling off neutron skins from neutron-rich nuclei: Constraints on the symmetry energy from neutron-removal cross sections. *Phys. Rev. Lett.*, 119:262501, 2017.
- [42] J. Hüfner, K. Schäfer, and B. Schürmann. Abrasionablation in reactions between relativistic heavy ions. *Phys. Rev. C*, 12:1888–1898, 1975.
- [43] M.S. Hussein, R.A. Rego, and C.A. Bertulani. Microscopic theory of the total cross-section and application to stable and exotic nuclei. *Physics Reports*, 201(5):279– 334, 1991.
- [44] B. V. Carlson, R. C. Mastroleo, and M. S. Hussein. Fragment production in heavy-ion reactions. *Phys. Rev. C*, 46:R30–R33, 1992.
- [45] B. V. Carlson. Microscopic abrasion-ablation approximation to projectile fragmentation. *Phys. Rev. C*, 51:252– 268, 1995.
- [46] C. A. Bertulani and P. Danielewicz. Introduction to Nuclear Reactions. CRC Press, London, 2021 - second edition.
- [47] W. J. Swiatecki J. D. Bowman and C. F. Tsang. LBL Report LBL-2908 (unpublished), 1973.
- [48] C. A. Bertulani. Heavy ion reaction cross-sections. Braz. J. Phys., 16:380–412, 1986.
- [49] C.A. Bertulani. Density dependence of in-medium nucleon-nucleon cross sections. J. Phys. G, 27(8):L67– L71, 2001.
- [50] C. A. Bertulani and C. De Conti. Pauli blocking and medium effects in nucleon knockout reactions. *Phys. Rev.* C, 81(6), 2010.
- [51] E. Clementel and C. Villi. The nucleon effective mass and the statistical model of the nucleus. Il Nuovo Cimento (1955-1965), 9(6):950–989, 1958.
- [52] Peter Ring and Peter Schuck. The Nuclear Many-Body Problem. Texts and Monographs in Physics. Springer-Verlage, Heidelberg, 2004.
- [53] R. F. Carlson et al. Proton total reaction cross sections for the doubly magic nuclei ¹⁶O, ⁴⁰Ca, and ²⁰⁸Pb in the energy range 20-50 MeV. *Phys. Rev. C*, 12:1167–1175, 1975.
- [54] A. Ingemarsson et al. Reaction cross sections for 65 MeV

protons on targets from 9Be to 208Pb. Nuclear Physics A, 653(4):341-354, 1999.

- [55] A. Auce et al. Reaction cross sections for protons on ${}^{12}C$, ${}^{40}Ca$, ${}^{90}Zr$, and ${}^{208}Pb$ at energies between 80 and 180 MeV. *Phys. Rev. C*, 71:064606, 2005.
- [56] Shingo Tagami, Tomotsugu Wakasa, Jun Matsui, Masanobu Yahiro, and Maya Takechi. Neutron skin thickness of ²⁰⁸Pb determined from the reaction cross section for proton scattering. *Phys. Rev. C*, 104:024606, 2021.
- [57] M. Takechi et al. Reaction cross sections at intermediate energies and fermi-motion effect. *Phys. Rev. C*, 79:061601, 2009.
- [58] A. Ozawa, T. Suzuki, and I. Tanihata. Nuclear size and related topics. *Nuclear Physics A*, 693(1):32–62, 2001. Radioactive Nuclear Beams.
- [59] A.S. Goldhaber. Statistical models of fragmentation processes. *Physics Letters B*, 53(4):306–308, 1974.
- [60] L. Bertocchi and A. Tékou. Fermi-motion effects in neutron stripping cross-section from relativistic deuteronnucleus collisions. Il Nuovo Cimento A (1965-1970), 21(2):223-235, 1974.
- [61] W G Lynch. Nuclear fragmentation in proton- and heavyion-induced reactions. Annual Review of Nuclear and Particle Science, 37(1):493–535, 1987.
- [62] Guang wei Fan, W. Y. Kong, T. F. Han, X. Li, J. B. Ma, Zong qiang Sheng, G. Z. Shi, F. Tian, J. Wang, and C Zhang. Fermi-motion effect on the intermediate energy nucleus-nucleus collision. *Modern Physics Letters* A, 31:1650216, 2016.
- [63] S. J. Wallace. Eikonal expansion. Phys. Rev. Lett., 27:622–625, 1971.
- [64] Stephen J Wallace. Eikonal expansion. Annals of Physics, 78(1):190–257, 1973.
- [65] Micah Buuck and Gerald A. Miller. Corrections to the eikonal approximation for nuclear scattering at medium energies. *Phys. Rev. C*, 90:024606, 2014.
- [66] C. Hebborn and P. Capel. Analysis of corrections to the eikonal approximation. *Phys. Rev. C*, 96:054607, 2017.
- [67] C. A. Bertulani. Relativistic continuum-continuum coupling in the dissociation of halo nuclei. *Phys. Rev. Lett.*, 94:072701, 2005.
- [68] Wen Hui Long and C. A. Bertulani. Nucleus-nucleus interaction between boosted nuclei. *Phys. Rev. C*, 83:024907, 2011.
- [69] B. G. Todd-Rutel and J. Piekarewicz. Neutron-rich nuclei and neutron stars: A new accurately calibrated interaction for the study of neutron-rich matter. *Phys. Rev. Lett.*, 95:122501, 2005.
- [70] W. Bauhoff. Tables of reaction and total cross sections for proton-nucleus scattering below 1 GeV. Atomic Data and Nuclear Data Tables, 35(3):429–447, 1986.
- [71] W. R. Webber, J. C. Kish, and D. A. Schrier. Individual charge changing fragmentation cross sections of relativistic nuclei in hydrogen, helium, and carbon targets. *Phys. Rev. C*, 41:533–546, 1990.
- [72] F Flesch, G Iancu, W Heinrich, and H Yasuda. Projectile fragmentation of silicon ions at 490 A Mev. *Radiation Measurements*, 34(1):237–240, 2001. Proceedings of the 20th International Conference on Nuclear Tracks in Solids.
- [73] C. Zeitlin, A. Fukumura, S.B. Guetersloh, L.H. Heilbronn, Y. Iwata, J. Miller, and T. Murakami. Fragmentation cross sections of 28Si at beam energies from 290A

to 1200A MeV. Nuclear Physics A, 784(1):341-367, 2007.

- [74] S. Cecchini et al. Fragmentation cross sections of Fe26, Si14 and C6 ions of 0.3-10 A GeV on polyethylene, CR39 and aluminum targets. *Nuclear Physics A*, 807(3):206– 213, 2008.
- [75] T. Yamaguchi et al. Energy-dependent charge-changing cross sections and proton distribution of ²⁸Si. *Phys. Rev.* C, 82:014609, 2010.
- [76] Jun-Sheng Li, Dong-Hai Zhang, Jin-Xia Cheng, S. Kodaira, and N. Yasuda. Fragmentation cross sections of 788A MeV 28Si on carbon and polyethylene targets. *Chi*nese Journal of Physics, 54(2):314–317, 2016.
- [77] Jun-Sheng Li, Ying-Hua Dang, Dong-Hai Zhang, Jin-Xia Cheng, S. Kodaira, and N. Yasuda. Charge-changing cross sections of 736A MeV ²⁸Si on carbon targets. *Chinese Physics Letters*, 34(10):102501, 2017.
- [78] M. Tanaka et al. Swelling of doubly magic 48 Ca core in ca isotopes beyond n = 28. *Phys. Rev. Lett.*, 124:102501, 2020.
- [79] S. Yamaki et al. Systematic study of individual chargechanging cross sections of intermediate-energy secondary beams. Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms, 317:774–778, 2013. XVIth International Conference on ElectroMagnetic Isotope Separators and Techniques Related to their Applications, December 2-7, 2012 at Matsue, Japan.
- [80] D. Adhikari et al. Precision determination of the neutral weak form factor of ⁴⁸Ca. *Phys. Rev. Lett.*, 129:042501, 2022.
- [81] G. A. Lalazissis, J. König, and P. Ring. New parametrization for the lagrangian density of relativistic mean field theory. *Phys. Rev. C*, 55:540–543, 1997.
- [82] T.W. Donnelly, J. Dubach, and Ingo Sick. Isospin dependences in parity-violating electron scattering. *Nuclear Physics A*, 503(3):589–631, 1989.
- [83] C. J. Horowitz, S. J. Pollock, P. A. Souder, and R. Michaels. Parity violating measurements of neutron densities. *Phys. Rev. C*, 63:025501, 2001.
- [84] M Thiel, C Sfienti, J Piekarewicz, C J Horowitz, and M Vanderhaeghen. Neutron skins of atomic nuclei: per aspera ad astra. *Journal of Physics G: Nuclear and Particle Physics*, 46(9):093003, 2019.
- [85] A. Trzcińska, J. Jastrzębski, P. Lubiński, F. J. Hartmann, R. Schmidt, T. von Egidy, and B. Kłos. Neutron density distributions deduced from antiprotonic atoms. *Phys.*

Rev. Lett., 87:082501, 2001.

- [86] J. Jastrzebski, A. Trzcinska, P. Lubinski, B. Klos, F. J. Hartmann, T. von Egidy, and S. Wycech. Neutron density distributions from antiprotonic atmos compared with hadron scattering data. *International Journal of Modern Physics E*, 13(01):343–351, 2004.
- [87] S. Abrahamyan et al. Measurement of the neutron radius of ²⁰⁸Pb through parity violation in electron scattering. *Phys. Rev. Lett.*, 108:112502, 2012.
- [88] J. Jastrzebski, H. Daniel, T. von Egidy, A. Grabowska, Y.S. Kim, W. Kurcewicz, P. Lubiński, G. Riepe, W. Schmid, A. Stolarz, and S. Wycech. Signature of a neutron halo in 232Th from antiproton absorption. *Nuclear Physics A*, 558:405–414, 1993.
- [89] R. Schmidt, F. J. Hartmann, T. von Egidy, T. Czosnyka, J. Iwanicki, J. Jastrzębski, M. Kisieliński, P. Lubiński, P. Napiorkowski, L. Pieńkowski, A. Trzcińska, J. Kulpa, R. Smolańczuk, S. Wycech, B. Kłos, K. Gulda, W. Kurcewicz, and E. Widmann. Nucleon density of ¹⁷²Yb and ¹⁷⁶Yb at the nuclear periphery determined with antiprotonic x rays. *Phys. Rev. C*, 58:3195–3204, 1998.
- [90] E. Friedman, A. Gal, and J. Mares. Antiproton-nucleus potentials from global fits to antiprotonic x-rays and radiochemical data. *Nuclear Physics A*, 761(3):283–295, 2005.
- [91] R. Lazauskas and A. Obertelli. personal communication.
- [92] L. Ray. Neutron isotopic density differences deduced from 0.8 GeV polarized proton elastic scattering. *Phys. Rev.* C, 19:1855–1872, 1979.
- [93] M. Csatlos et al. Measurement of neutron-skin thickness in 208Pb by excitation of the GDR via inelastic α scattering. *Nuclear Physics A*, 719:C304–C307, 2003.
- [94] A. Krasznahorkay et al. Excitation of isovector spindipole resonances and neutron skin of nuclei. *Phys. Rev. Lett.*, 82:3216–3219, 1999.
- [95] A. Krasznahorkay et al. Neutron-skin thickness in neutron-rich isotopes. Nuclear Physics A, 731:224–234, 2004.
- [96] G. W. Hoffmann et al. Elastic scattering of 500 MeV polarized protons from ^{40,48}Ca, ⁹⁰Zr, and ²⁰⁸Pb, and breakdown of the impulse approximation at small momentum transfer. *Phys. Rev. Lett.*, 47:1436–1440, 1981.
- [97] Thomas Aumann and Andrea Horvat. Constraining energy-density functionals and the density-dependence of the symmetry energy by measurements of accurate cross sections with large acceptance at R3B. *GSI-PAC*, 2020.