



## NRC Publications Archive Archives des publications du CNRC

### **Nuclear quadrupole resonance measurement of the anisotropic magnetic shielding and quadrupole coupling constants of $^{151}\text{Eu}^{3+}$ and $^{153}\text{Eu}^{3+}$ dilute in $\text{YAIO}_3$ single crystal**

Erickson, L. E.; Sharma, K. K.

This publication could be one of several versions: author's original, accepted manuscript or the publisher's version. / La version de cette publication peut être l'une des suivantes : la version prépublication de l'auteur, la version acceptée du manuscrit ou la version de l'éditeur.

For the publisher's version, please access the DOI link below. / Pour consulter la version de l'éditeur, utilisez le lien DOI ci-dessous.

#### **Publisher's version / Version de l'éditeur:**

<https://doi.org/10.1103/PhysRevB.24.3697>

*Physical Review B*, 24, 7, pp. 3697-3700, 1981-10

#### **NRC Publications Record / Notice d'Archives des publications de CNRC:**

<https://nrc-publications.canada.ca/eng/view/object/?id=29c4e713-0967-4397-99b6-ec9a8a501d45>

<https://publications-cnrc.canada.ca/fra/voir/objet/?id=29c4e713-0967-4397-99b6-ec9a8a501d45>

Access and use of this website and the material on it are subject to the Terms and Conditions set forth at

<https://nrc-publications.canada.ca/eng/copyright>

READ THESE TERMS AND CONDITIONS CAREFULLY BEFORE USING THIS WEBSITE.

L'accès à ce site Web et l'utilisation de son contenu sont assujettis aux conditions présentées dans le site

<https://publications-cnrc.canada.ca/fra/droits>

LISEZ CES CONDITIONS ATTENTIVEMENT AVANT D'UTILISER CE SITE WEB.

#### **Questions?** Contact the NRC Publications Archive team at

PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca. If you wish to email the authors directly, please see the first page of the publication for their contact information.

**Vous avez des questions?** Nous pouvons vous aider. Pour communiquer directement avec un auteur, consultez la première page de la revue dans laquelle son article a été publié afin de trouver ses coordonnées. Si vous n'arrivez pas à les repérer, communiquez avec nous à PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca.



**Nuclear quadrupole resonance measurement of the anisotropic magnetic shielding  
and quadrupole coupling constants of  $^{151}\text{Eu}^{3+}$  and  $^{153}\text{Eu}^{3+}$   
dilute in  $\text{YAlO}_3$  single crystal**

L. E. Erickson and K. K. Sharma\*

*National Research Council of Canada, Ottawa, Canada K1A 0R8*

(Received 16 April 1981)

The nuclear quadrupole resonance spectra of  $\text{Eu}^{3+}$  in  $\text{YAlO}_3$  single crystal at  $\sim 10$  K are reported. From the ratio of the quadrupole interaction parameters of  $^{151}\text{Eu}$  and  $^{153}\text{Eu}$ , the ratio of the quadrupole moments corrected for the pseudoquadrupole interaction is obtained  $Q^{153}/Q^{151} = 2.5812 \pm 0.0010$ . With low-magnetic-field studies the anisotropic magnetic shielding factors are measured. They are  $\alpha_x = 0.58$ ,  $\alpha_y = 0.80$ ,  $\alpha_z = 0.47$  compared to an estimate of  $\alpha_z = 0.89$  by Elliott. A measured value  $\langle r^{-3} \rangle = 39 \text{ \AA}^{-3}$  is obtained using Elliott's formulation and the experimental values of the anisotropic magnetic shielding factors.

### I. INTRODUCTION

It is commonly believed that nuclear quadrupolar resonance measurements (NQR) of the europium nucleus in a solid have not been successful because of a large diamagnetic shielding of the nucleus from external static and rf magnetic fields by the crystal-field split  $f^6$  electron shell. Elliott<sup>1</sup> suggested that second-order interactions which are cross terms of the electronic Zeeman and the magnetic hyperfine interactions between the singlet  $^7F_0$  ground state and other states  $^7F_J (J \neq 0)$  of the ground manifold were responsible for this shielding. Judd<sup>2</sup> has found that other similar interactions were responsible for the anomalously large quadrupole splitting of  $^{152}\text{Eu}$  and  $^{154}\text{Eu}$  observed via angular correlation studies of radioactive decay products. Edmonds<sup>3</sup> in a room-temperature NMR study of  $\text{La}^{3+}$  in several different host crystals established that the anomalously large quadrupole splitting parameter  $P$  for the La nucleus (and also the lattice contribution to  $P$  for other rare-earth nuclei) is due to an unusually large *antishielding* of the nucleus from the field gradients of the host lattice by the closed-shell electrons of the rare-earth ion. Macfarlane *et al.*<sup>4</sup> has demonstrated that magnetic dipole transitions in Eu (in  $\text{EuP}_3\text{O}_{14}$ ) could be observed optically if a large pulsed (8 G, 25 msec) resonant rf magnetic field is applied to the sample. However, to our knowledge, no measurements have been made of the anisotropic diamagnetic shielding of  $\text{Eu}^{3+}$  in a solid.

In this paper, we report cw nuclear quadrupole resonance measurements of the two naturally occurring isotopes of  $^{151,153}\text{Eu}^{3+}$  which are substituted for 0.25% of the Y in a  $\text{YAlO}_3$  single crystal, in both zero field and in a low static magnetic field. A detailed magnetic field study of one transition ( $I_z = \frac{1}{2}$ ) - ( $I_z = \frac{3}{2}$ )

of one isotope, yielded all three anisotropic shielding factors for the Eu nucleus in the host crystal.

The ground term of  $\text{Eu}^{3+}$  ( $f^6$ ),  $^7F$  is split into seven multiplets by the spin-orbit interaction. In a solid host, each of the multiplets is further split by the action of the crystalline electric field. In  $\text{YAlO}_3$ , the ground term  $J=0$  is separated from the  $^7F_1$  triplet by only  $378 \text{ cm}^{-1}$ .<sup>5</sup> The usual Hamiltonian for a rare-earth ion in a solid is written

$$\mathcal{H} = A \vec{J} \cdot \vec{I} + P \left\{ [I_z^2 - \frac{1}{3} I(I+1)] + \eta(I_x^2 - I_y^2) \right\} + \beta \vec{H} \cdot (\vec{J} + \vec{S}) + \hbar \gamma \vec{H} \cdot \vec{I} \quad (1)$$

where the terms are the magnetic hyperfine, nuclear quadrupole, and finally the electronic and nuclear Zeeman interactions. If one considers only the  $J=0$  state, it appears as if only the nuclear quadrupole and nuclear Zeeman terms remain, leaving a simple spectrum with an isotropic nuclear Zeeman effect. However, Elliott<sup>1</sup> has pointed out that second-order interactions which couple the Zeeman and magnetic hyperfine terms via  $^7F_1$  and higher levels produce a nuclear Zeeman-like term which results in an anisotropic  $\gamma$ . The simplified Hamiltonian for this ion may be written

$$\mathcal{H} = P' \left\{ [I_z^2 - \frac{1}{3} I(I+1)] + \eta(I_x^2 - I_y^2) \right\} + \hbar (\gamma_x H_x I_x + \gamma_y H_y I_y + \gamma_z H_z I_z) \quad (2)$$

where  $P'$  is the effective quadrupole interaction parameter,  $\gamma_i = \gamma(1 - \alpha_i)$  ( $i = x, y, z$ ) is the effective gyromagnetic ratio and  $\alpha_i$  are Elliott's<sup>1</sup> shielding parameters. This reduction in  $\gamma$  from the bare nuclear value reduces the sensitivity of NQR experiments by the  $\gamma_i/\gamma$  ratio squared. The nuclear quadrupole interaction  $P$  is dominated not by the crystal-

field gradient directly but by an electronic distortion which is proportional to the nuclear quadrupole moment.<sup>1</sup> The pseudoquadrupole effect which is predominant in  $\text{Pr}^{3+}$  ( $J=4$ ) is estimated to be much smaller than  $P$  for  $\text{Eu}^{3+}$ . In Sec. III, we calculate the very small pseudoquadrupole corrections and obtain a ratio of the quadrupole moments for the two isotopes.

## II. EXPERIMENT

The NQR measurements were made using an optical rf double-resonance technique previously described.<sup>6</sup> A single crystal of  $\text{YAlO}_3:\text{Eu}^{3+}$  (0.25 at.%) at  $\sim 10$  K was illuminated by a 5815-Å single-mode, frequency-stabilized laser beam to selectively excite hyperfine levels of the  ${}^7F_0 \rightarrow {}^5D_0$  transition. The optical excitation produced a substantial redistribution of population in the ground-state hyperfine levels which was probed by an  $\sim 0.5$ -G rf magnetic field. This rf field was derived from a frequency synthesizer, 20-W amplifier, and a 2-turn coil surrounding the sample. The rf-induced magnetic dipole transitions tend to equalize the populations and this results in a large change in the optical absorption and therefore the crystal fluorescence. The  ${}^5D_0 \rightarrow {}^7F_1$  fluorescence was monitored. In Fig. 1, a NQR of  ${}^{153}\text{Eu}$  at 23.007 MHz is shown. This measurement consists of the




FIG. 1. Optically detected zero-field ground state NQR of the ( $I_z = \pm \frac{1}{2} \rightarrow I_z = \pm \frac{3}{2}$ ) transition of  ${}^{151}\text{Eu}^{3+}$  (0.25 at.%) in  $\text{YAlO}_3$  at 10 K. The linewidth is 28 kHz (FWHM). The solid line is a smooth curve drawn through the experimental points. The slight asymmetry which is present is due to rapid sweeping of the transition. Each point represents two 0.5-sec stops on a staircase of 2-kHz steps. The center frequency is dependent upon the position of the exciting laser in the inhomogeneously broadened  ${}^7F_0 \rightarrow {}^5D_0$  transition which is 1 GHz above line center.

sum of two 2-min sweeps of the peak in the absence of an external magnetic field. Each sweep consists of 250 frequency steps. The effective time constant is 1 sec. Four such resonances were observed at  $22.993 \pm 0.003$ ,  $45.970 \pm 0.006$ ,  $59.594 \pm 0.006$ ,  $119.150 \pm 0.012$  MHz, when the laser was exciting the inhomogeneously broadened  ${}^5D_0 \leftarrow {}^7F_0$  transition  $\sim 0.5$  GHz below line center. The linewidths full width at half maximum (FWHM) are 28, 56, 56, 110 kHz, respectively. The 23.0-MHz line position was observed to vary at  $+10$  kHz/GHz when the laser frequency was varied from  $-5$  to  $+5$  GHz in relation to line center. Only two isotopes,  ${}^{151}\text{Eu}$  ( $I = \frac{5}{2}$ ) and  ${}^{153}\text{Eu}$  ( $I = \frac{5}{2}$ ) occur naturally. The ratio of natural abundances is 0.48/0.52. We interpret the two lowest observed transitions as those of  ${}^{151}\text{Eu}$ , ( $I_z = \frac{1}{2}, -\frac{3}{2}$ ) and  $I_z = (\frac{3}{2}, -\frac{5}{2})$ . Since the ratio of zero field frequencies is almost exactly 2, the quadrupole splitting parameters are very nearly axially symmetric as was observed for the ground state of  $\text{Pr}^{3+}$  in this host. The two higher frequency transitions are then  ${}^{153}\text{Eu}^{3+}$ . The observed ratio of these two higher frequencies is 1.9994, differing slightly from the 1.9993 ratio between the two lower frequencies. ( ${}^{153}\eta = 0.0157 \pm 0.0030$ ,  ${}^{151}\eta = 0.0164 \pm 0.0030$ .) The ratio of the quadrupole parameters  $P$  for the two isotopes (at line center) is equal to

$${}^{153}P'/{}^{151}P' = 2.5919 \pm 0.0005$$

This compares to the ratio of quadrupole moments measured by Baker and Williams<sup>7</sup> in an ENDOR study of  $\text{Eu}^{2+}$ ,  $Q^{153}/Q^{151} = 2.5835 \pm 0.0191$  and by Sandars and Woodgate<sup>8</sup> in an atomic beam experiment  $Q^{153}/Q^{151} = 2.5445 \pm 0.0194$ . If the estimates of Elliott are correct, then the measured ratio  ${}^{153}P'/{}^{151}P'$  quadrupole should be nearly equal to the measured ratio of the nuclear quadrupole moment. The difference is due to the contribution of the pseudoquadrupole moment to  $P'$  which he estimates as  $\frac{1}{1500}$  of the total. The zero-field linewidths are approximately proportional to the resonance frequency. This would indicate that these linewidths are due primarily to an inhomogeneous crystal-field broadening rather than being dominated by magnetic dipolar broadening as in the case of  $\text{Pr}^{3+}$ . This is also consistent with the large variation in resonant frequency depending on which part of the inhomogeneous line is excited. The measured ratio of quadrupole parameters is independent of line position. It is obvious then that this site-selective technique enables one to obtain a more accurate determination of the ratio of quadrupole parameters than is possible if the entire line is observed as in conventional NQR. A corollary is that for nonsite selective techniques the resonances would be significantly broader than observed here. A crystal with a different residual strain would likely give a

different set of zero-field frequencies, but the ratios of those parameters should be the same as that measured in this work within the accuracy of the experiments.

Detailed magnetic resonance studies were made in a static magnetic field of 50.0 G on the 23-MHz transition ( $^{151}\text{Eu}$ ) as a function of the angle between the crystal axes and the field. This isotope was studied because its magnetic moment is a factor of 2 larger than the other and also because the linewidth of the lowest-frequency transition is 2 times smaller. Indeed, at 50 G no splitting was resolved for the 59.6-MHz transition. The 23-MHz transition was studied because the  $\frac{1}{2}-\frac{3}{2}$  transition is sensitive to all three magnetic splitting factors in first order unlike the  $\frac{3}{2}-\frac{5}{2}$  where only  $\gamma_z$  is effective. Figure 2 shows a typical NQR result for an external static field of 50 G at an angle of  $75^\circ$  to the  $z$  axis of the  $\text{Eu}^{3+}$  site. The small magnetic splitting factors do not give clearly resolved spectra in this low magnetic field. This limits the accuracy of the experiment somewhat. Figure 3 shows the resonance frequencies versus the field direction where the field lies in the  $xz$  plane of the  $\text{Eu}$  site. Figure 4 gives similar data for the field in the  $xy$  plane of the site. Only one  $\text{Eu}^{3+}$  site is observed. The Hamiltonian (2) was fitted to the data. The solid curves are those calculated by diagonalizing a complex  $6 \times 6$  matrix of  $\mathcal{H}$  assuming  $\eta=0$  using our best-fit parameters.  $P = 11.500 \pm 0.002$  MHz,  $\gamma_x/2\pi = 0.44 \pm 0.02$ ,  $\gamma_y/2\pi = 0.21 \pm 0.02$ ,  $\gamma_z/2\pi = 0.56 \pm 0.02$  kHz/G. ( $\alpha_x = 0.58 \pm 0.02$ ,  $\alpha_y = 0.80 \pm 0.02$ ,  $\alpha_z = 0.47 \pm 0.02$ .) This clearly demonstrates the anisotropy of the magnetic shielding parameters as predicted by Elliott. They differ significantly in mag-




FIG. 2. Optically detected NQR line in a field of 50 G. The field lies in the  $xz$  plane of the  $\text{Eu}^{3+}$  site at an angle  $75^\circ$  to the  $z$  axis.




FIG. 3.  $^{151}\text{Eu}^{3+}$  nuclear quadrupole resonance frequencies ( $I_z = \frac{1}{2} - I_z = \frac{3}{2}$ ) in an external static magnetic field of 50.0 G. The field is always perpendicular to the  $y$  axis of the  $\text{Eu}^{3+}$  site. As the field approaches the  $z$  axis, the outer transitions become very weak. The solid lines are calculated using the best-fit data assuming the asymmetry parameter  $\eta=0$ .

nitude from his estimated  $\alpha_z = 0.89$ . Judd<sup>2</sup> has pointed out that the  $\langle r^{-3} \rangle$  used by Elliott is too large. The accuracy of these measurements is limited somewhat by the low field in which the measurements were made. The resonance linewidths are about  $\frac{1}{10}$  of the maximum splitting. This leads to some scatter in Figs. 3 and 4.




FIG. 4.  $^{151}\text{Eu}^{3+}$  nuclear quadrupole resonance frequencies ( $I_z = \frac{1}{2} - I_z = \frac{3}{2}$ ) in an external magnetic field of 50.0 G. The field is always perpendicular to the  $z$  axis of the  $\text{Eu}^{3+}$  site. The solid lines are calculated using the best-fit data.

In order to confirm the assignment of isotopes (beyond the quadrupole ratios), a single measurement of magnetic splitting in a field of 182 G was made for the 23-MHz and the 59.6-MHz transitions. The splitting of the 23-MHz transition was  $2.16 \pm 0.22$  times that of the 59-MHz transition. The measured ratio of their respective magnetic moments is equal to  $2.2632 \pm 0.0026$ .<sup>7</sup>

### III. DISCUSSION

Elliott<sup>1</sup> has given an expression for the anisotropic diamagnetic shielding which can be used to derive an experimental value for  $\langle r^{-3} \rangle$  using measured values of  $\alpha_i$

$$\alpha_i = \frac{40\beta^2 \langle r^{-3} \rangle}{3\Delta_i} \quad (3)$$

where  $\Delta_i$  is an energy level of  ${}^7F_1$  and  $i$  indicates  $x$ ,  $y$ , or  $z$  depending on which level is chosen. The  $J_z = 0$  level at  $500 \text{ cm}^{-1}$  must be used to obtain  $\alpha_z$ . From this we calculate

$$\langle r^{-3} \rangle_z = 40.8 \text{ \AA}^{-3}$$

The shielding factors for  $x$ ,  $y$  yield

$$\langle r^{-3} \rangle_x = 37.1 \text{ \AA}^{-3}, \quad \langle r^{-3} \rangle_y = 36.8 \text{ \AA}^{-3}$$

This is to be compared with Judd and Lindgren's value<sup>9</sup> of  $45.4 \text{ \AA}^{-3}$ .

The ratio of nuclear quadrupole moments for the two isotopes differs from the measured ratio of quadrupole interaction constants  ${}^{151}P'/{}^{153}P'$  due to the pseudoquadrupole interaction which is proportional to the nuclear magnetic moment of each isotope rather than the nuclear electric quadrupole moment. Elliott gives an equation for the pseudoquadrupole correction in second order,

$$P'' = \frac{5}{2}(\alpha_x + \alpha_y - 2\alpha_z) \langle r^{-3} \rangle \beta_N^2 \mu_N^2 / 6I^2 \quad (4)$$

If the mean of the experimental  $\langle r^{-3} \rangle = 39 \text{ \AA}^{-3}$  and the Pichanick, Sandars, and Woodgate<sup>10</sup> values of  $\mu_N$  for these two isotopes is used, one obtains

TABLE I. Europium quadrupole moment ratios  $Q^{153}/Q^{151}$ .

This work	$2.5812 \pm 0.0010$
Sandars and Woodgate <sup>a</sup>	$2.5445 \pm 0.0194$
Baker and Williams <sup>b</sup>	$2.5835 \pm 0.0191$

<sup>a</sup> Eu atomic beam study. See Ref. 8.

<sup>b</sup> Eu<sup>2+</sup> ENDOR study. See Ref. 7.

${}^{151}P'' = 0.0515$  and  ${}^{153}P'' = 0.0100$  MHz. This gives a quadrupole moment ratio

$$Q^{153}/Q^{151} = 2.5812 \pm 0.0010$$

The error is due almost equally to the uncertainty in the measured ratio  ${}^{153}P'/{}^{151}P'$  and an estimated 10% accuracy of the pseudoquadrupole correction.<sup>11</sup>

This result is compared with previously published values of the quadrupole moment ratio in Table I. It lies within the experimental precision of Baker and Williams, and just outside that of Sandars and Woodgate.

### IV. CONCLUSION

NQR measurements of Eu<sup>3+</sup> in solids are reported. The anisotropic diamagnetic shielding predicted by Elliott for a rhombic host are observed and agree very well with his second-order formalism. A "measured" value of  $\langle r^{-3} \rangle = 39 \text{ \AA}^{-3}$  for Eu<sup>3+</sup> is obtained which is approximately 15% less than the calculated value of Judd and Lindgren. Finally, with corrections to the measured values of the quadrupole interaction constant, a new value of the quadrupole moment ratio  $Q^{153}/Q^{151} = 2.5812 \pm 0.0010$  is obtained.

### ACKNOWLEDGMENT

We thank M. Weber for the loan of the YAlO<sub>3</sub>:Eu<sup>3+</sup> crystal used in these experiments.

\*Permanent address: Indian Institute of Technology, Kanpur, India.

<sup>1</sup>R. J. Elliott, Proc. Phys. Soc. London Sect. B **70**, 119 (1957).

<sup>2</sup>B. R. Judd, C. A. Lovejoy, and D. A. Shirley, Phys. Rev. **128**, 1733 (1962).

<sup>3</sup>D. T. Edmonds, Phys. Rev. Lett. **10**, 129 (1963).

<sup>4</sup>R. M. Macfarlane, R. M. Shelby, A. Z. Genack, and D. A. Weitz, Opt. Lett. **5**, 462 (1980).

<sup>5</sup>Fluorescence from  ${}^5D_0$  was observed at 5906, 5942.5, and 5989. This corresponds to energy levels of  ${}^7F_1$  at 265, 369, and  $500 \text{ cm}^{-1}$ . C. A. Morrison, N. Karayianis and D. E. Wortman, in Harry Diamond Lab Report No. HDL-TR-1788, have calculated the three levels to be at 287, 371, and  $473 \text{ cm}^{-1}$  using crystal-field parameters extrapo-

lated from best-fit values of other rare-earth ions in YAlO<sub>3</sub>.

<sup>6</sup>L. E. Erickson, Phys. Rev. B **19**, 4412 (1979).

<sup>7</sup>J. M. Baker and F. I. B. Williams, Proc. R. Soc. London Ser. A **267**, 283 (1962).

<sup>8</sup>P. G. H. Sandars and G. K. Woodgate, Proc. R. Soc. London Ser. A **257**, 269 (1960).

<sup>9</sup>B. R. Judd and I. P. K. Lindgren, Phys. Rev. **122**, 1802 (1961).

<sup>10</sup>F. M. Pichanick, P. G. H. Sandars, and G. K. Woodgate, Proc. R. Soc. London Ser. A **257**, 277 (1960).

<sup>11</sup>Following Judd (Ref. 2) and Edmonds (Ref. 3), a negative value of  $P$  is chosen. The pseudoquadrupole correction is positive.