

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/53149>

Please be advised that this information was generated on 2022-08-25 and may be subject to change.

Nucleoplasmic LAP2 α –lamin A complexes are required to maintain a proliferative state in human fibroblasts

Vanja Pekovic,¹ Jens Harborth,² Jos L.V. Broers,³ Frans C.S. Ramaekers,³ Baziel van Engelen,⁴ Martin Lammens,⁴ Thomas von Zglinicki,⁵ Roland Foisner,⁶ Chris Hutchison,¹ and Ewa Markiewicz¹

¹School of Biological and Biomedical Sciences, University of Durham, Durham DH1 3LE, England, UK

²Alnylam Pharmaceuticals, Inc., Bioanalytics and Preclinical, Cambridge, MA 02142

³Department of Molecular Cell Biology, Cardiovascular Research Institute Maastricht, and Research Institute Growth and Development, University of Maastricht, Maastricht, Netherlands

⁴Neuromuscular Centre Nijmegen, Institute of Neurology, University Medical Centre, Nijmegen, Netherlands

⁵Henry Wellcome Laboratory for Biogerontology Research, University of Newcastle, Newcastle General Hospital, Newcastle-upon-Tyne NE4 6BE, England, UK

⁶Max F. Perutz Laboratories, Medical University Vienna, 1030 Vienna, Austria

In human diploid fibroblasts (HDFs), expression of lamina-associated polypeptide 2 α (LAP2 α) upon entry and exit from G₀ is tightly correlated with phosphorylation and subnuclear localization of retinoblastoma protein (Rb). Phosphoisoforms of Rb and LAP2 α are down-regulated in G₀. Although RbS780 phosphoform and LAP2 α are up-regulated upon reentry into G₁ and colocalize in the nucleoplasm, RbS795 migrates between nucleoplasmic and speckle compartments. In HDFs, which are null for lamins A/C, LAP2 α is mislocalized within nuclear aggregates,

and this is correlated with cell cycle arrest and accumulation of Rb within speckles. Nuclear retention of nucleoplasmic Rb during G₁ phase but not of speckle-associated Rb depends on lamin A/C. siRNA knock down of LAP2 α or lamin A/C in HDFs leads to accumulation of Rb in speckles and G₁ arrest, probably because of activation of a cell cycle checkpoint. Our results suggest that LAP2 α and lamin A/C are involved in controlling Rb localization and phosphorylation, and a lack or mislocalization of either protein leads to cell cycle arrest in HDFs.

Introduction

The nuclear envelope (NE) is composed of inner and outer nuclear membranes that are perforated by nuclear pore complexes and supported by the nuclear lamina (Hutchison, 2002). Proteins of the inner nuclear membrane have been implicated in organization of nuclear architecture and cell cycle control. One such family of inner nuclear membrane proteins is lamina-associated polypeptide 2 (LAP2), generated by alternative splicing from a single gene (Foisner and Gerace 1993; Harris et al., 1994; Berger et al., 1996). LAPs have only been found in vertebrates; up to six isoforms exist in humans and mice (α , β , γ , δ , ϵ , and ζ). Most LAP2 isoforms have a closely related N-terminal nucleoplasmic domain of variable length with a nuclear localization signal sequence, a single-membrane spanning region, and a

short luminal domain at their C terminus. LAP2 α is structurally and functionally unique. It shares the first N-terminal 187 residues with all other LAPs but contains a unique C-terminal domain of 506 residues, which lacks a membrane spanning domain. As a result, LAP2 α is distributed diffusely throughout the interphase nucleus except for nucleoli (Dechat et al., 1998). At their N terminus, LAP2 proteins share a LEM (LAP2, emerin, MAN1) domain (aa 111–152; Lin et al., 2000), which binds to the DNA bridging protein barrier-to-auto-integration factor (Furukawa, 1999; Shumaker et al., 2001), and a LEM-like domain (aa 1–85), which binds to DNA and chromosomes (Cai et al., 2001). In addition, LAP2 α interacts with chromosomes via its α -specific C-terminal domain in a phosphorylation-dependent manner (Vlcek et al., 1999; Dechat et al., 2004). Overexpression of C-terminal fragments of LAP2 α dominantly inhibits assembly of endogenous LAP2 α , nuclear membranes, and A-type lamins in *in vitro* nuclear assembly assays and causes a cell cycle arrest in interphase, indicating a role for LAP2 α in cell cycle progression (Vlcek et al., 2002).

Correspondence to Chris Hutchison: c.j.hutchison@durham.ac.uk

Abbreviations used in this paper: HDF, human diploid fibroblast; LAP2, lamina-associated polypeptide 2; MEF, mouse embryonic fibroblast; NCS, newborn calf serum; NE, nuclear envelope; Rb, retinoblastoma protein.

The online version of this article contains supplemental material.

Supplemental Material can be found at:
<http://jcb.rupress.org/content/suppl/2007/01/16/jcb.200606139.DC1.html>

LAP2 α is found in stable complexes with the type V intermediate filament proteins lamins A/C in the nucleoplasm (Dechat et al., 2000). The LAP2 α -specific C terminus associates with the C terminus of lamins A/C (aa 319–566) *in vivo* and *in vitro*. Dominant-negative lamin mutants, which cause aggregation of lamins A/C, also cause LAP2 α to redistribute to the same aggregates, indicating a functional association between LAP2 α and lamin A/C (Dechat et al., 2000). Mutations in the gene encoding lamins A/C (*LMNA*) cause a spectra of human diseases (termed laminopathies), including muscular dystrophies, lipodystrophies, cardiomyopathies, neuropathies, dermopathies, and premature aging syndromes (Hutchison and Worman, 2004). It has been suggested that altered lamin A/C–LAP2 α associations might occur in laminopathies (Goldman et al., 2004) because LAP2 α binds to a region of lamin A/C where many different mutations have been found. In support of this hypothesis, a mutation in the LAP2 gene causing cardiomyopathy and affecting the C-terminal domain of LAP2 α alters LAP2 α interaction with lamins A/C (Taylor et al., 2005).

The retinoblastoma protein (Rb) has functions in muscle and fat cell differentiation and in coordinating proliferation and differentiation during muscle regeneration (Smith and Kudlow, 2005). The C-terminal domain of Rb is involved in its nuclear tethering, which is essential for its growth-suppressing activity (Mittnacht, 1998). In early G₁, hypophosphorylated Rb is tethered in the nucleus and is capable of binding transcription factor E2F-1, which prevents the transcriptional activation of S phase-specific genes and traverse to S phase. Hyperphosphorylation of Rb during late G₁ phase releases it from nuclear tethers, which in turn releases and derepresses E2F and leads to passage through S phase. A-type lamins (Mancini et al., 1994; Ozaki et al., 1994) and LAP2 α (Markiewicz et al., 2002; Dorner et al., 2006) both interact with the C-terminal nuclear anchorage domain of Rb. In line with an Rb tethering function of LAP2 α –lamins A/C complexes, LAP2 α has recently been shown to be involved in an Rb-mediated negative cell cycle control (Dorner et al., 2006).

One model for laminopathies proposes that mutations in *LMNA* impair the control of cell proliferation, particularly, Rb-mediated regulatory mechanisms controlling the exit and reentry into the cell cycle (Gotzmann and Foisner, 2006). In support of this model, embryonic fibroblasts from a *Lmna*^{-/-} mouse show cell cycle defects because of reduced levels of Rb (Johnson et al., 2004). Interestingly, despite the proposed growth-suppressive function of LAP2 α (Dorner et al., 2006), its expression is dramatically down-regulated upon cell cycle exit in human diploid fibroblasts (HDFs) and conversely up-regulated during reentry into the cell cycle (Markiewicz et al., 2002). Therefore, we tested the hypothesis that in HDFs, LAP2 α expression is required for cell cycle progression via direct regulation of Rb. Here, we report that expression and lamin A-dependent organization of LAP2 α is indeed required for maintaining HDFs in a proliferative state by promoting uniform nucleoplasmic localization of Rb. Our findings support the “cell cycle–proliferating–aging model” for laminopathies and provide a novel biological function for LAP2 α in cell cycle regulation of human nontransformed adult cells.

Results

Changes in expression, solubility, and distribution of lamins, LAPs, and pRb isoforms in quiescent human fibroblasts

We previously showed that the expression of LAP2 α is down-regulated as HDFs progress from a proliferating to a G₀ state after serum starvation (Markiewicz et al., 2002). To further investigate LAP2 α responsiveness to factors that induce quiescence in HDFs, we grew cultures to confluence and used immunoblotting to investigate the expression of proteins of interest. The level of expression of LAP2 α declined dramatically as cultures reached confluence and the protein was undetectable in postconfluent cultures. In contrast, no changes in the level of expression of LAP2 β , lamin A/C, or lamin B2 were detected in confluent or postconfluent cultures (Fig. S1 a, available at <http://www.jcb.org/cgi/content/full/jcb.200606139/DC1>). The level of expression of hyperphosphorylated Rb and, in particular, the RbS780 isoform also declined progressively in confluent and postconfluent cultures and correlated precisely with changes in the expression of LAP2 α (Fig. S1, a and b). These data confirm that loss of expression of LAP2 α occurs as a direct consequence of entry into G₀ and is closely correlated with dephosphorylation of pRb at serine 780.

To further investigate the correlation between changes in expression of LAP2 α and phosphorylation of pRb, we performed double immunofluorescence microscopy to investigate both the expression and distribution of total Rb and forms phosphorylated at serines 780 and 795. In cultures induced to enter G₀ through confluence, LAP2 α as well as total Rb (Ab2) and RbS780 were barely detected in a majority of cells (Fig. 1, a–c). Cells still expressing higher levels of LAP2 α also expressed relatively high levels of RbS780 and RbS795 (Fig. 1, a and c). In contrast to Rb780, RbS795 was still detectable in cells with low levels of LAP2 α and distributed in a small number of nucleoplasmic foci (Fig. 1 c, arrows).

To see how the distribution and phosphorylation of Rb is correlated with expression of LAP2 α , G₀ cultures were restimulated to reenter the cell cycle. 12 h after serum restimulation, LAP2 α was detected in the nuclei of all cells and was distributed uniformly within the nucleoplasm, excluding nucleoli. Total and RbS780 were also expressed in nearly all cells and had a distribution that was very similar to that of LAP2 α (Fig. 1, a and b). In contrast, RbS795 was located in a restricted number of nucleoplasmic foci (Fig. 1 c). 18 h after serum restimulation, the intensity of LAP2 α , Rb, and RbS780 had increased but remained uniformly distributed throughout the nucleoplasm, whereas RbS795 remained within nucleoplasmic foci. Only at 24 h after serum restimulation, when the staining intensity of LAP2 α , Rb, RbS780, and RbS795 was maximal, were all four proteins distributed uniformly throughout the nucleoplasm.

Next, we investigated the solubility properties of lamins, LAPs, and Rb in HDFs using a nuclear matrix extraction procedure. Using immunoblotting, we found that lamins A/C and lamin B2 were relatively more soluble in proliferating HDFs than in quiescent HDFs. In contrast, both LAP2 α and LAP2 β displayed increased solubility properties in quiescent HDFs




Figure 1. LAP2 α expression is correlated with phosphorylation of Rb at serine 780 and a nucleoplasmic distribution of Rb. Confluent (C) or serum-starved HDF cultures, which were restimulated to enter the cell cycle for 12, 18, or 24 h, were prepared for immunofluorescence either before (a–c) or after extraction with detergents. Cultures were costained with antibodies against LAP2 α and Ab2 (a and d), LAP2 α and RbS780 (b and e), or LAP2 α and RbS795 (c and f). Micrographs are presented as single black-and-white images. Arrows in c denote cells with RbS795 in speckles. Bars, 10 μ m.

compared with proliferating HDFs. Similarly, Rb was more soluble in quiescent HDFs than in proliferating HDFs (Fig. S1 c).

To investigate the solubility properties of the total Rb and the two Rb phosphoisoforms more closely, G₀ or serum-restimulated HDFs were extracted in situ. In most HDFs induced to enter G₀, total Rb, RbS780, and RbS795 were barely detectable (Fig. 1, d–f). Between 12 and 18 h after serum restimulation, all three forms of Rb were detectable and resistant to extraction. However, although Rb and RbS780 showed a more uniform staining throughout the nucleoplasm (Fig. 1, d and e), RbS795 was located in brightly stained nucleoplasmic foci (Fig. 1 f). 24 h after serum restimulation, when cells enter into S phase, all isoforms were relatively more soluble after extraction with Triton X-100 (Fig. 1, e and f).

Collectively, our data suggest that progression of HDFs from G₀ to a proliferating state (and vice versa) leads to a substantial remodeling of lamina proteins correlated with a more complex pattern of Rb nuclear anchorage than had previously been thought. In HDFs entering or exiting a quiescent state, Rb appears to be present in at least two different compartments depending on its phosphorylation status. Only when cells enter S phase (24 h after serum restimulation) are all


forms of Rb detected in an apparently uniform distribution and largely soluble.

RbS795 associates with splicing speckle compartments during cell cycle exit and entry

The nature of the foci detected by antibodies against RbS795 was examined by colocalization with a range of antibodies previously reported to detect nuclear foci. There was no substantial colocalization between RbS795 and dense chromatin compartments (detected by DAPI; Fig. S2 a, available at <http://www.jcb.org/cgi/content/full/jcb.200606139/DC1>), phosphohistone H2A.X, or PML bodies (Fig. S2, c and d). In contrast, there was almost complete colocalization of RbS795 foci with splicing factor SC-35, which resides within splicing speckle compartments (Fig. S2 b), indicating that in cells entering G₀, this form of Rb is localized in splicing speckle compartments.

To further investigate the distribution and nuclear anchorage of RbS795, we performed double immunofluorescence and confocal microscopy on HDFs that had been restimulated from G₀, before and after nuclear matrix extraction. Between 6 and 12 h after serum restimulation, RbS795 colocalized with

Figure 2. RbS795 associates with splicing speckles during early G₁ phase of the cell cycle. HDFs were restimulated to divide from a quiescent state for 6, 12, 18, or 24 h and prepared for immunofluorescence before (a) or after extraction with detergents, nucleases, and salt (b). Cultures were costained with antibodies against splicing factor SC-35 to detect splicing speckles and RbS795 and investigated by confocal microscopy. Micrographs are projected as single black-and-white or two-color merged images in which SC-35 fluorescence is in red and RbS795 fluorescence is in green. Bars, 10 μ m.



splicing speckle compartments and remained insoluble within this compartment. However, after 18 h, colocalization between RbS795 and SC-35 was lost, but RbS795 was still resistant to extraction. Only upon entry into S phase (24 h) did RbS795 become completely soluble (Fig. 2). Thus, associations of Rb with the splicing speckles are both growth dependent and related to the expression of LAP2 α . HDFs just entering the cell cycle and expressing low levels of LAP2 α have RbS795 associated with splicing speckles. As LAP2 α levels increase, RbS795 is no longer associated with speckle compartments but codistributes with LAP2 α and other Rb forms within the nucleoplasm.

Absence of lamin A/C in HDFs correlates with a mislocalization of LAP2 α , cell cycle arrest, and accumulation of pRb in speckles

To investigate whether lamins A/C directly influence the distribution of Rb within speckles, we performed double immunofluorescence on HDFs from a patient with lethal fetal akinesia, which harbors a homozygous mutation Y259X in the LMNA gene and is null for lamins A/C (Muchir et al., 2003). When grown for 2–4 d, an increasing number of Y259X HDFs (20–40%) displayed an abnormal accumulation of LAP2 α into aggregates typically at one pole of the nucleus (Fig. 3). These cells stained negatively with antibodies against the proliferation marker Ki67 (Fig. 3 a). In addition, cells that were either negative for LAP2 α or in which LAP2 α was entirely located within aggregates had greatly reduced levels of total Rb and RbS780 (Fig. 3, b and c). In contrast, nearly all cells containing LAP2 α aggregates expressed RbS795, which was mainly located in nuclear speckles (Fig. 3 d) colocalizing with anti-SC35–positive splicing speckles (Fig. S3, available at <http://www.jcb.org/cgi/content/full/jcb.200606139/DC1>). Therefore, the absence of lamins A/C is correlated with aggregation of LAP2 α and accumulation of RbS795 in speckle compartments.

To investigate the influence of lamins A/C on nuclear anchorage of the different Rb isoforms, control or Y259X HDFs were subjected to extraction with detergents before staining with different Rb antibodies. In exponentially dividing HDFs, the nuclei of most cells stained positively for total Rb and RbS780 before extraction. After extraction, \sim 40% cells had greatly reduced or absent staining for total Rb and RbS780, whereas \sim 60% remained strongly positive (Fig. 4, a and b), reflecting the different cell cycle stages. RbS795 was uniformly distributed in the nucleus of most cells before extraction, whereas it was found in speckles in \sim 60% of cells and greatly reduced or absent in \sim 40% of cells after extraction (Fig. 4 c). In contrast to control cells, the majority of Y259X HDFs were negative or only weakly positive for total Rb and RbS780 before extraction, and nearly 100% of cells were negative for Rb and RbS780 after extraction (Fig. 4, d and e), whereas \sim 90% of Y259X HDFs still contained RbS795 restricted entirely to speckles after extraction (Fig. 4 f).

These data suggest that nuclear anchorage of RbS780 was lamin A/C dependent. To test this hypothesis, Y259X HDFs were transfected with either GFP–lamin A or GFP–lamin C and prepared for immunofluorescence before or after detergent extraction. RbS780 was strongly retained in most cells that expressed GFP–lamin A after extraction (Fig. 4 g). Surprisingly, RbS780 was not retained in cells expressing GFP–lamin C (Fig. 4 h). Our data suggest that anchorage of nucleoplasmic forms of Rb is dependent on expression of lamin A, whereas anchorage of forms of Rb located in speckles is independent of lamins A and C.

siRNA knock down of lamin A/C and LAP2 α leads to cell cycle arrest and accumulation of Rb in speckles

In Y259X HDFs, aggregation of LAP2 α and cell cycle arrest occurs in only 40% of cells, suggesting that a compensatory mechanism might be abrogating the effects of loss of lamin A/C

in culture. To investigate the consequences of loss of lamin A/C or LAP2 α more directly, we used siRNA to knock down each protein independently. HDFs were transfected with siRNA targeted to LAP2 α or lamins A/C or as a control, scrambled siRNA. We observed a $\sim 70\%$ reduction in the level of LAP2 α or lamins A/C expression 48 h after transfection with specific siRNA as compared with control siRNA, whereas expression of LAP2 β was unaffected. Surprisingly, lamin A expression decreased by $\sim 30\%$ in cells transfected with LAP2 α RNAi, and LAP2 α decreased similarly in lamin A/C RNAi-treated cells. The total amount of Rb and, more dramatically (by $\sim 70\%$), the levels of RbS780 were reduced in both LAP2 α and lamins A/C RNAi cultures (Fig. 5 a).

To investigate the effects of siRNA knock down of LAP2 α or lamins A/C on cell proliferation, we tested the expression of Ki67. After transfection with scrambled siRNA, only 10% of cells were negative for Ki67. In contrast, ~ 70 and $\sim 60\%$ of cells transfected with LAP2 α -specific siRNA or lamin A/C-specific siRNA, respectively, were negative for Ki67 (Fig. 5 b). We concluded that knock down of LAP2 α or lamins A/C in HDFs leads to cell cycle arrest.

To confirm these findings, we performed DNA flow cytometry on HDFs. Cells were arrested in G₀ by serum starvation, transfected with siRNA, restimulated by addition of complete serum medium, and harvested 0, 24, 48, or 72 h after transfection (Fig. 5 c). After 24 h, a large proportion of cells had entered S phase, and there was no difference in the cell cycle profiles between any of the transfected cultures. 48 h after transfection (the time at which knock down of LAP2 α and lamins A/C were detected), control RNAi cultures proceeded normally through G₂/M and displayed a cell cycle profile similar to asynchronously dividing fibroblasts. In contrast, cultures in which LAP2 α or lamin A/C had been knocked down had a greater proportion of cells in S phase and an abnormally high proportion of cells in G₂/M phase. In view of previous data, showing a negative effect of LAP2 α on the G₁-S phase transition (Dorner et al., 2006), our result may indicate that loss of LAP2 α or lamins A/C promotes more rapid progression through G₁ and premature entry into S phase, leading to a checkpoint arrest in G₂. However, 72 h after transfection, the LAP2 α RNAi-treated cultures appeared to accumulate in G₁ phase of the cell cycle. At the same time point, many cells in lamin A/C knockdown cultures had accumulated in G₁, but some remained arrested in G₂. Control cultures were dividing asynchronously at this time point (Fig. 5 c). These data are consistent with a cell cycle arrest of LAP2 α - and lamin A/C-deficient cells in the G₁ phase, presumably by activating a checkpoint to overcome defects due to a premature S phase entry in the previous cell cycle round.

We previously reported that down-regulation of LAP2 α after cell cycle arrest during myoblast differentiation is correlated with relocation of lamins A/C from the nucleoplasm to the NE (Markiewicz et al., 2005). To investigate the effects of siRNA knock down of LAP2 α on lamins A/C distribution, fibroblasts were costained with antibodies against LAP2 α and either lamin A or lamin C. In cells in which LAP2 α was reduced or absent, both lamin A (Fig. 6 a) and lamin C (not depicted) concentrated at the NE. In contrast, in control





Figure 3. Organization of LAP2 α and expression of Rb isoforms in lamin A/C-null human fibroblasts. (a) HDFs with Y259X LMNA mutation were grown in culture for 2 or 4 d and costained with antibodies against LAP2 α and Ki67 (a), LAP2 α and Ab2 (b), LAP2 α and RbS780 (c), or LAP2 α and RbS795 (d). All cultures were stained with DAPI (blue fluorescence) and investigated by confocal microscopy. Micrographs are projected as individual black-and-white images or three-color merged images with LAP2 α fluorescence in red and Ki67 or Rb fluorescence in green. Bar, 10 μ m.

fibroblasts or in those fibroblasts in knockdown cultures that still expressed LAP2 α , both lamins were found in the nucleoplasm and at the NE.

To investigate the influence of LAP2 α on the nuclear distribution of Rb, control and LAP2 α siRNA-transfected cells were costained with antibodies against LAP2 α and different forms of Rb. Although most LAP2 α -positive control cells had Rb, RbS780, and RbS795 distributed uniformly throughout the nucleoplasm (Fig. 6, b, c, and d), in LAP2 α knockdown cultures, Rb was greatly reduced and had a granular appearance (Fig. 6 b) and RbS780 was largely absent (Fig. 6 c). In contrast, RbS795 was present in all LAP2 α -deficient cells but was

Figure 4. Nuclear anchorage of Rb isoforms in lamin A/C-null human fibroblasts. (a–f) Control or Y259X HDFs were prepared for immunofluorescence either before or after extraction with detergents. Cultures were stained with antibodies against Ab2 (a and d), RbS780 (b and e), or RbS795 (c and f), counterstained with DAPI, and viewed by confocal microscopy. (g and h) Y259X fibroblasts were transfected with either GFP–lamin A (g) or GFP–lamin C (h). After transfection, cultures were prepared for immunofluorescence either before or after extraction with detergents. Cultures were stained with antibodies against RbS780, counterstained with DAPI, and viewed by confocal microscopy. Micrographs are presented as single black-and-white images. Bars, 10 μ M.



restricted to nuclear speckles (Fig. 6 d), whereas it had more uniform distribution in LAP2 α -expressing cells. Our data suggest that knock down of LAP2 α or lamin A/C expression in HDFs leads to cell cycle arrest, dephosphorylation, and migration of Rb into nuclear speckles.

Discussion

LAP2 α function is required for cell proliferation in human fibroblasts

Here, we show that entry of HDFs into G₀ is correlated with loss of expression of LAP2 α . Furthermore, in HDFs that are null for lamins A/C, LAP2 α accumulates in aggregates, which is correlated with cell cycle arrest. Finally, siRNA knock down of LAP2 α or lamins A/C in HDFs results in cell cycle arrest, and this is correlated with accumulation of Rb into speckles. Our data suggests that the expression and normal distribution of LAP2 α allows proper regulation of Rb and thus maintains a proliferative state in HDFs. Our findings are consistent with other reports that show that HDFs lacking lamins A/C grow very slowly in culture (Muchir et al., 2003) and that HDFs harboring mutant lamins enter a senescent state prematurely (Goldman et al., 2004). Other papers, however, appear to partly contradict these findings. Embryonic fibroblasts from a *Lmna*^{-/-} mouse (-/- mouse embryonic fibroblasts [MEFs]) display a rapid growth phenotype with characteristics of Rb-null fibro-

blasts resulting from proteosomal degradation of Rb (Johnson et al., 2004). A second paper found that *Lmna*^{-/-} MEFs displayed a rapid growth phenotype but suggested that this resulted from inhibition of TGF β signaling, leading to increased phosphorylation of Rb (Van Berlo et al., 2005). We have shown that in mouse fibroblasts and adipocytes, LAP2 α represses E2F activity via an Rb-dependent pathway and that knock down of LAP2 α in HeLa cells leads to rapid growth (Dorner et al., 2006). Although there are discrepancies between each paper as to the precise mechanism involved, all three agree that a loss of lamin A/C or LAP2 α function leads to rapid proliferation through pathways involving Rb.

The apparent discrepancies between the findings reported here and those reported previously (Johnson et al., 2004; Van Berlo et al., 2005; Dorner et al., 2006) probably reflect fundamental differences between the model systems used. Our study exclusively made use of HDFs, whereas the previous papers used either MEFs or transformed human cell lines. HDFs are able to respond to stimuli, such as genotoxic stresses, by inducing a checkpoint arrest transiently in G₂ and eventual G₁ arrest. This checkpoint is nonfunctional in MEFs as well as in transformed human cell lines with inactivated pocket proteins, which continue to divide or enter apoptosis after exposure to genotoxic agents (Baus et al., 2003; Jackson et al., 2005). Interestingly, FACS analysis of LAP2 α and lamin A/C knockdown HDFs reveals that, initially, cells accumulate in G₂, before entering a G₁




Figure 5. siRNA knock down of LAP2 α induces delayed G₂/M progression and eventual cell cycle arrest in G₁ phase. HDFs were transfected with either control LAP2 α or lamin A/C RNAi. (a) After 48 h, transfected cultures were harvested for immunoblotting and probed with antibodies against LAP2 α (75 kD), LAP2 β (55 kD), lamins A/C (70/65 kD), total pRb (110 kD), RbS780 (115 kD), or β -actin (43 kD). (b) After 72 h in culture, transfected cells on triplicate coverslips were stained with antibodies against LAP2 α or lamin A/C and a proliferation marker Ki67. 300 cells on each coverslip were scored for Ki67. The chart shows the fraction of nuclei (expressed as a mean percentage) that scored negative for Ki67 and standard error in two independent experiments. (c) Synchronized human fibroblasts were transfected with either control LAP2 α or lamin A/C RNAi and harvested after 0, 24, 48, or 72 h. Transfected cells were prepared for cell cycle analysis, and cell cycle phase distribution (percentage of G₁, S, and G₂/M cells) was determined by the Dean/Jett/Fox model of collected DNA histograms.

arrest. Therefore, it appears that HDFs respond to a loss of LAP2 α or lamin A/C function as if they were treated with a genotoxic stress. Alternatively, the loss of Rb repressor activity upon LAP2 α or lamin A/C down-regulation may prematurely drive the cells into S phase, causing activation of an incomplete S phase cell cycle arrest initially in G₂ and eventually in G₁. Therefore, the reason that previous studies have not found that a loss of either lamins A/C or LAP2 α leads to cell cycle arrest is most likely because these studies used cell lines in which this checkpoint pathway was abrogated.

LAP2 α maintains pRb within a nucleoplasmic compartment


It has been shown that Rb binds to lamin A in vitro and associates with intermediate filament-like structures in the nucleus (Mancini et al., 1994; Ozaki et al., 1994). As Rb is no longer anchored in the nucleus of *Lmna*^{-/-} MEFs (Johnson et al., 2004), these findings are widely interpreted as evidence that anchorage of Rb within the nucleoplasm is dependent on lamins A/C. Our current data suggest that anchorage of Rb within the nucleoplasm depends on both lamin A and LAP2 α . We show that nucleoplasmic forms of Rb are entirely absent from

cells that contain either no LAP2 α or in which LAP2 α is entirely restricted to aggregates. These findings are consistent with our previous observations that Rb is anchored within the nucleoplasm via its C-terminal pocket C domain, which also binds to LAP2 α in vitro (Markiewicz et al., 2002). Therefore, our results suggest that functional complexes of lamin A and LAP2 α are required for anchorage of Rb within the nucleoplasm of HDFs.

We also show that absence of LAP2 α or its accumulation in aggregates is correlated with the preferential association of Rb with speckle compartments. Previous studies have suggested that Rb can bind speckle-associated protein p84 via sequences within its N-terminal domain (Durfee et al., 1994). These data are entirely consistent with our current findings because the N-terminal domain of Rb does not bind to lamins A/C or LAP2 α (Markiewicz et al., 2002), and we show that Rb anchorage in speckles is independent of lamins A/C or LAP2 α .

We propose that Rb is distributed predominantly within a nucleoplasmic compartment, through an association of pocket C with LAP2 α and lamin A. However, Rb can also associate with speckle compartments via its N-terminal domain. Although in principle these two modes of association are noncompetitive,

Figure 6. siRNA knock down of LAP2 α causes changes in subnuclear distribution of lamins A/C and pRb expression patterns. HDFs were transfected with either control or LAP2 α RNAi. (a) After 72 h in culture, transfected cells were prepared for immunofluorescence and costained with antibodies against lamin A and LAP2 α (a), LAP2 α and Ab2 (b), LAP2 α and RbS780 (c), or LAP2 α and RbS795 (d). All cultures were counterstained with DAPI and viewed by confocal microscopy. Micrographs are presented as individual black-and-white images. Bar, 10 μ m.



we propose that in practice Rb only associates with speckles when LAP2 α expression is down-regulated.

How do lamin A and LAP2 α influence cell proliferation?

The cell cycle arrest caused by down-regulation of LAP2 α and lamin A/C in HDFs is correlated with rapid dephosphorylation of pRb. In a related study, we have shown that the introduction of dominant-negative lamin A mutants into C2C12 myoblasts causes loss of expression of LAP2 α , which is also correlated with the absence of RbS780 (Markiewicz et al., 2005). Dephosphorylated forms of Rb bind to and inhibit the transcription factor E2F, thereby suppressing growth in both normally dividing cells and during differentiation of several mesenchymal cell types (for review see Mittnacht, 1998). Therefore, it seems likely that the growth arrest caused by knock down of LAP2 α results from dephosphorylation of Rb. Whether association of Rb into speckle domains is also a prerequisite for growth arrest is unclear. Certainly, the N-terminal region of Rb, through which Rb can associate with speckle-associated protein p84, is crucial for both terminal differentiation and growth suppression (Riley et al., 1997). We propose that functional LAP2 α –lamin A nucleoplasmic complexes might be required to anchor Rb in a nucleoplasmic compartment. This may allow correct regulation of Rb by cyclin-dependent kinases and protein phosphatases, which in turn makes cells responsive to environmental stresses, such as genotoxic agents. In conclusion, the corollary of this hypothesis is that a loss of function of lamin A or LAP2 α , in cells with functional checkpoint pathways, might lead to irreversible cell cycle arrest and possibly cellular senescence because Rb can no longer be maintained in a phosphorylated state within the nucleoplasm and instead enters speckle compartments by default. This could in turn explain why mutations in lamins A/C and LAP2 α both cause diseases associated with premature aging (Hutchison and Worman, 2004).

Materials and methods

Cell culture and media

HDFs from a needle biopsy of the forearm were cultured in DME (Invitrogen) supplemented with 10% newborn calf serum (NCS) and 10 U/ml penicillin plus 50 μ g/ml streptomycin, at 37°C in humidified incubators containing 5% CO₂. HDFs from a patient with a homozygous Y259X LMNA mutation were obtained as an autopsy sample after an informed consent. Cultures were grown to 70–80% confluence and subcultured thereafter at a seeding density of 3×10^5 cells per 75 cm² flasks. In this study, control and Y259X HDFs were used between passages p8 and 12.

Serum starvation, serum restimulation, and contact inhibition

To induce quiescence by serum starvation, control HDFs were grown for 3 d in complete medium and maintained in starvation medium (0.5% NCS) for 5 d. To induce cell cycle reentry, quiescent HDFs were serum restimulated (10% NCS) for 6, 12, 18, or 24 h and prepared for immunofluorescence microscopy. To induce quiescence by contact inhibition, HDFs were grown in complete medium (10% serum) for 7 d (confluent stage) or for 10 d (postconfluent stage).

Immunofluorescence and confocal microscopy

Immunofluorescence was performed according to established laboratory procedures (Markiewicz et al., 2002). The primary antibodies used and their dilutions are described in Table I. Secondary antibodies were donkey anti-mouse and anti-rabbit IgG conjugated to rhodamine (TRITC) or fluorescein (FITC; Strata-Tech) and, for viewing, DNA in cell coverslips were mounted in DAPI. For imaging cells, a confocal microscope imaging system (Radiance 2000; Bio-Rad Laboratories) with LaserSharp software (Bio-Rad Laboratories) or confocal microscope imaging system (LSM 510 META; Carl Zeiss MicroImaging, Inc.) with LSM510 image browser software (Carl Zeiss MicroImaging, Inc.) were used at ambient temperature, equipped with 40 \times /1.3 and 63 \times /1.4 oil-immersion lens and nonimaging photodetection device (photomultiplier tube; Carl Zeiss MicroImaging, Inc.). The imaging medium used was immersion oil (Immersol 518; Carl Zeiss MicroImaging, Inc.). A dynamic range adjustment was used to optimize the signal for the fluorophores, and images were collected in sequential mode (Bio-Rad Laboratories) or multitrack mode (Carl Zeiss MicroImaging, Inc.). Any brightness and contrast adjustments were performed in Photoshop (Adobe).

In situ nuclear matrix extraction

In situ nuclear matrix extractions using sequential treatment with detergents, nucleases, and salt were performed as described by Dyer et al. (1997). After extraction, cells were prepared for immunofluorescence microscopy.

Preparation of whole cell extracts

HDF cell pellets were washed with ice-cold PBS and lysed in 0.1 ml of ice-cold hypotonic buffer per 10^6 cells (10 mM Tris, pH 7.4, 10 mM KCl,

Table 1. Primary antibodies used in this study

Antibody	Target	Antibody type	Dilution		Source
			IF	IB	
JOL2	Lamin A/C	Mouse/m		1:200	Dyer et al., 1997
133A2	Lamin A	Mouse/m	1:200		Novus Biologicals
LN43	Lamin B2	Mouse/m		1:100	Dyer et al., 1997
LAP15	LAP2 α	Mouse/m	1:10	1:100	Dechat et al., 1998
245-2	LAP2 α	Rabbit/p	1:250		Vlcek et al., 2002
LAP17	LAP2 β	Mouse/m		1:100	Dechat et al., 1998
IF8	Pocket A of Rb	Mouse/m		1:10	D. Lane, University of Dundee, Dundee, UK
Rb Ser780	Phosphoserine 780	Rabbit/p	1:100	1:1,000	Cell Signalling
Rb Ser795	Phosphoserine 795	Rabbit/p	1:100		Cell Signalling
H2.A.X (Ser139)	Phosphoserine 139; Histone 2.A.X	Mouse/m	1:1,000		Upstate Biotechnology
Ab2	Pocket C of Rb	Rabbit/p	1:50		Santa Cruz Biotechnology, Inc.
Ki67	Ki67	Rabbit/p	1:150		DakoCytomation
SC-35	SC-35	Mouse/m	1:2,000		Sigma-Aldrich
PML	PML bodies	Mouse/m	1:100		A. Vaughan, University of St. Andrews, St. Andrews, UK
AC-40	β -actin	Mouse/m		1:2,000	Sigma-Aldrich

IF, immunofluorescence; IB, immunoblotting; m, monoclonal; p, polyclonal.

3 mM MgCl₂, and 0.1% Triton X-100), containing protease inhibitor cocktail and 100 U/ml of RNase-free DNase I (Sigma-Aldrich) for 10 min on ice. Cell lysates were analyzed by SDS-PAGE. Alternatively, before the above, cell pellets were subjected to sequential extraction according to the protocol of Dyer et al. (1997) using ice-cold buffers and freshly added protease inhibitor cocktail.

Gel electrophoresis and immunoblotting

1D SDS-PAGE was performed according to Laemmli (1970). For immunoblotting, proteins separated on gels were electrophoretically transferred onto nitrocellulose membranes (Schleicher & Schuell) using the Mini Trans-Blot system (Bio-Rad Laboratories) and processed according to standard protocols (Markiewicz et al., 2002). Secondary antibodies were donkey anti-mouse or donkey anti-rabbit IgG conjugated to HRP (Jackson Immuno-Research Laboratories). For the immunological detection of proteins, membranes were incubated in ECL reagents (GE Healthcare) and visualized using either LAS-1000 intelligent dark box (FujiFilm) or autoradiography. Densitometry of signals obtained for the protein bands was performed using Image Gauge analysis software (FujiFilm).

Selection of siRNA sequences, transfection of siRNAs, and determination of transfection efficiency

LAP2 α - and lamin A/C-specific siRNA duplexes were obtained from Ambion. The sequences were selected from the open reading frames to obtain 21-nt sense and 21-nt antisense strand with symmetric 2-nt 3' overhangs of identical sequence. The sequences of each strand of siRNA oligos were as follows: LAP2 sense, 5'-GCUAAGAAAGUACAUACUUtt-3'; LAP2 anti-sense, 5'-AAGUAUGUACUUUCUUAGCtt-3'; lamin A/C sense, 5'-CUGGACUUCAGAGAACAAtt-3'; and lamin A/C anti-sense, 5'-UGUUCUUCUGGAAGUCCAGtt-3'. RNAi transfection procedure was modified from Harborth et al. (2001). On the day of transfection, cells were seeded at 5 × 10⁴ cells/well in 6-well plates in the presence of 10% NCS and no antibiotics and transfected in tandem with specific or control (scrambled) siRNAs using Oligofectamine reagent (Life Technologies). 24 h after transfection, medium was replaced by fresh medium (10% NCS) without antibiotics. Cells were assayed 48–72 h after transfection. Transfection efficiency was determined by immunofluorescence microscopy and immunoblotting. Specific silencing of LAP2 α or lamin A/C was confirmed by four independent experiments.

Flow cytometry

HDFs were synchronized by serum starvation and restimulation. Cultures were trypsinized, counted, and transfected with LAP2 α , lamin A/C, or control siRNA. Transfected cells were harvested by trypsinization after 0, 24, 48, or 72 h and resuspended in PBS and methanol prechilled

at –20°C (1:9 ratio). Subsequently, washed cell pellets were incubated in PBS containing 100 μ g/ml RNase and 25 μ g/ml propidium iodide, washed in PBS, centrifuged, and diluted in PBS for cell cycle analysis on a FACSCaliber flow cytometer (Becton Dickinson). Data were collected as DNA histograms from 5,000 single-cell events, and cell cycle phase distribution (percentage of G₁, S, and G₂/M cells) was determined by the Dean/Jett/Fox model using FlowJo software.

Online supplemental material

Fig. S1 shows expression and solubility properties of confluent HDFs. Fig. S2 shows colocalization of RbS795 foci with splicing speckles in confluent HDFs. Fig. S3 shows colocalization of RbS795 foci with splicing speckles in Y259X HDFs. Online supplemental material is available at <http://www.jcb.org/cgi/content/full/jcb.200606139/DC1>.

This work was supported by grants from the Association for International Cancer Research (to C. Hutchison and R. Foissner), the Muscular Dystrophy Campaign (to C. Hutchison), the Austrian Science Research Fund (FWF P17871; to R. Foissner), an EU FP6 Programme, Eurolaminopathies (to R. Foissner and C. Hutchison), and a Medical Research Council Career Development award to E. Markiewicz.

We wish to thank Ian Dimmick (Institute of Human Genetics, Bioscience Centre, Newcastle-upon-Tyne, UK) for his help with FACS analyses.

Submitted: 29 June 2006

Accepted: 15 December 2006

References

- Baus, F., V. Gire, D. Fisher, J. Piette, and V. Dulic. 2003. Permanent cell cycle exit in G₂ phase after DNA damage in normal human fibroblasts. *EMBO J.* 22:3992–4002.
- Berger, R., L. Theodor, J. Shoham, E. Gokkel, F. Brok-Simoni, K.B. Avraham, N.G. Copeland, N.A. Jenkins, G. Rechavi, and A.J. Simon. 1996. The characterization and localization of the mouse thymopoietin/lamina-associated polypeptide 2 gene and its alternatively spliced products. *Genome Res.* 6:361–370.
- Cai, M., Y. Huang, R. Ghirlando, K.L. Wilson, R. Craigie, and G.M. Clore. 2001. Solution structure of the constant region of nuclear envelope protein LAP2 reveals two LEM-domain structures: one binds BAF and the other binds DNA. *EMBO J.* 20:4399–4407.
- Dechat, T., J. Gotzmann, A. Stockinger, C.A. Harris, M.A. Talle, J.J. Siekierka, and R. Foissner. 1998. Detergent-salt resistance of LAP2alpha in interphase nuclei and phosphorylation-dependent association with

- chromosomes early in nuclear assembly implies functions in nuclear structure dynamics. *EMBO J.* 17:4887–4902.
- Dechat, T., B. Korbei, O.A. Vaughan, S. Vlcek, C.J. Hutchison, and R. Foisner. 2000. Lamina-associated polypeptide 2alpha binds intranuclear A-type lamins. *J. Cell Sci.* 113:3473–3484.
- Dechat, T., A. Gajewski, B. Korbei, D. Gerlich, N. Daigle, T. Haraguchi, K. Furukawa, J. Ellenberg, and R. Foisner. 2004. LAP2alpha and BAF transiently localize to telomeres and specific regions on chromatin during nuclear assembly. *J. Cell Sci.* 117:6117–6128.
- Dorner, D., S. Vlcek, N. Foeger, A. Gajewski, C. Makolm, J. Gotzmann, C.J. Hutchison, and R. Foisner. 2006. Lamina-associated polypeptide 2alpha regulates cell cycle progression and differentiation via the retinoblastoma-E2F pathway. *J. Cell Biol.* 173:83–93.
- Durfee, T., M.A. Mancini, D. Jones, S.J. Elledge, and W.H. Lee. 1994. The amino-terminal region of the retinoblastoma gene product binds a novel nuclear matrix protein that co-localizes to centers for RNA processing. *J. Cell Biol.* 127:609–622.
- Dyer, J.A., I.R. Kill, G. Pugh, R.A. Quinlan, E.B. Lane, and C.J. Hutchison. 1997. Cell cycle changes in A-type lamin associations detected in human dermal fibroblasts using monoclonal antibodies. *Chromosome Res.* 5:383–394.
- Foisner, R., and L. Gerace. 1993. Integral membrane proteins of the nuclear envelope interact with lamins and chromosomes, and binding is modulated by mitotic phosphorylation. *Cell.* 73:1267–1279.
- Furukawa, K. 1999. LAP2 binding protein 1 (L2BP1/BAF) is a candidate mediator of LAP2-chromatin interaction. *J. Cell Sci.* 112:2485–2492.
- Goldman, R.D., D.K. Shumaker, M.R. Erdos, M. Eriksson, A.E. Goldman, L.B. Gordon, Y. Gruenbaum, S. Khuon, M. Mendez, R. Varga, and F.S. Collins. 2004. Accumulation of mutant lamin A causes progressive changes in nuclear architecture in Hutchinson-Gilford progeria syndrome. *Proc. Natl. Acad. Sci. USA.* 101:8963–8968.
- Gotzmann, J., and R. Foisner. 2006. A-type lamin complexes and regenerative potential: a step towards understanding laminopathic diseases? *Histochem. Cell Biol.* 125:33–41.
- Harborth, J., S.M. Elbashir, K. Bechert, T. Tuschl, and K. Weber. 2001. Identification of essential genes in cultured mammalian cells using small interfering RNAs. *J. Cell Sci.* 114:4557–4565.
- Harris, C.A., P.J. Andryuk, S. Cline, H.K. Chan, A. Natarajan, J.J. Siekierka, and G. Goldstein. 1994. Three distinct human thymopoietins are derived from alternatively spliced mRNAs. *Proc. Natl. Acad. Sci. USA.* 91:6283–6287.
- Hutchison, C.J. 2002. Lamins: building blocks or regulators of gene expression? *Nat. Rev. Mol. Cell Biol.* 3:848–858.
- Hutchison, C.J., and H.J. Worman. 2004. A-type lamins: guardians of the soma? *Nat. Cell Biol.* 6:1062–1067.
- Jackson, M.W., M.K. Agarwal, J. Yang, P. Bruss, T. Uchiyumi, M.L. Agarwal, G.R. Stark, and W.R. Taylor. 2005. p130/p107/p105Rb-dependent transcriptional repression during DNA-damage-induced cell-cycle exit at G2. *J. Cell Sci.* 118:1821–1832.
- Johnson, B.R., R.T. Nitta, R.L. Frock, L. Mounkes, D.A. Barbie, C.L. Stewart, E. Harlow, and B.K. Kennedy. 2004. A-type lamins regulate retinoblastoma protein function by promoting subnuclear localization and preventing proteasomal degradation. *Proc. Natl. Acad. Sci. USA.* 101:9677–9682.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 227:680–685.
- Lin, F., D.L. Blake, I. Callebaut, I.S. Skerjanc, L. Holmer, M.W. McBurney, M. Paulin-Levasseur, and H.J. Worman. 2000. MAN1, an inner nuclear membrane protein that shares the LEM domain with lamina-associated polypeptide 2 and emerin. *J. Biol. Chem.* 275:4840–4847.
- Mancini, M.A., B. Shan, J.A. Nickerson, S. Penman, and W.H. Lee. 1994. The retinoblastoma gene product is a cell cycle-dependent, nuclear matrix-associated protein. *Proc. Natl. Acad. Sci. USA.* 91:418–422.
- Markiewicz, E., T. Dechat, R. Foisner, R.A. Quinlan, and C.J. Hutchison. 2002. Lamin A/C binding protein LAP2alpha is required for nuclear anchorage of retinoblastoma protein. *Mol. Biol. Cell.* 13:4401–4413.
- Markiewicz, E., M. Ledran, and C.J. Hutchison. 2005. Remodelling of the nuclear lamina and nucleoskeleton is required for skeletal muscle differentiation in vitro. *J. Cell Sci.* 118:409–420.
- Mittnacht, S. 1998. Control of pRB phosphorylation. *Curr. Opin. Genet. Dev.* 8:21–27.
- Muchir, A., B.G. van Engelen, M. Lammens, J.M. Mislow, E. McNally, K. Schwartz, and G. Bonne. 2003. Nuclear envelope alterations in fibroblasts from LGMD1B patients carrying nonsense Y259X heterozygous or homozygous mutation in lamin A/C gene. *Exp. Cell Res.* 291:352–362.
- Ozaki, T., M. Saijo, K. Murakami, H. Enomoto, Y. Taya, and S. Sakiyama. 1994. Complex formation between lamin A and the retinoblastoma gene product: identification of the domain on lamin A required for its interaction. *Oncogene.* 9:2649–2653.
- Riley, D.J., C.Y. Liu, and W.H. Lee. 1997. Mutations of N-terminal regions render the retinoblastoma protein insufficient for functions in development and tumour suppression. *Mol. Cell. Biol.* 17:7342–7352.
- Shumaker, D.K., K.K. Lee, Y.C. Tanheco, R. Craigie, and K.L. Wilson. 2001. LAP2 binds to BAF/DNA complexes: requirement for the LEM domain and modulation by variable regions. *EMBO J.* 20:1754–1764.
- Smith, E.D., and B.A. Kudlow. 2005. A-type nuclear lamins, progerias and other degenerative disorders. *Mech. Ageing Dev.* 126:447–460.
- Taylor, M.R., D. Slavov, A. Gajewski, S. Vlcek, L. Ku, P.R. Fain, E. Carniel, A. Di Lenarda, G. Sinagra, M.M. Boucek, et al. 2005. Thymopoietin (lamina-associated polypeptide 2) gene mutation associated with dilated cardiomyopathy. *Hum. Mutat.* 26:566–574.
- Van Berlo, J.H., J.W. Voncken, N. Kubben, J.L. Broers, R. Duisters, R.E. van Leeuwen, H.J. Crijns, F.C. Ramaekers, C.J. Hutchison, and Y.M. Pinto. 2005. A-type lamins are essential for TGF-beta1 induced PP2A to dephosphorylate transcription factors. *Hum. Mol. Genet.* 14:2839–2849.
- Vlcek, S., H. Just, T. Dechat, and R. Foisner. 1999. Functional diversity of LAP2alpha and LAP2beta in postmitotic chromosome association is caused by an alpha-specific nuclear targeting domain. *EMBO J.* 18:6370–6384.
- Vlcek, S., B. Korbei, and R. Foisner. 2002. Distinct functions of the unique C terminus of LAP2alpha in cell proliferation and nuclear assembly. *J. Biol. Chem.* 277:18898–18907.